

VISIT ONLINE  
OXFORD  
QUICK  
REFERENCE



OFFERS AUTHORITATIVE AND EXPANSIVE  
COVERAGE OF THE EARTH SCIENCES

# Oxford



DICTIONARY OF  
*Geology &*  
*Earth Sciences*



MICHAEL ALLABY

## **How to search for terms in *A Dictionary of Geology and Earth Sciences***

To find an entry in this e-book you can:

- Browse the [Alphabetical List of Entries](#) and select the entry you would like to view
- or
- Use your Search function to be taken to a complete list of references to your search term in the *Dictionary*
  - If your search term has its own entry, it will usually be listed at the top of your results
  - In cases where your search term appears in more than one entry heading, the results will be listed alphabetically

### **A note on special characters**

While most e-readers can display special characters (such as é and â), many cannot search for words containing them, unless the special characters themselves are typed into the search box. If you are unable to type these characters, please browse for your term using the [Alphabetical List of Entries](#).

## Alphabetical List of Entries

A B C D E F G H I J K L M N O P Q R  
S T U V W X Y Z

aa

AABW

AAC

Aalenian

Aalto-1

AAV

Ab

abandoned channel

abandonment facies association

abapical

abaptation

Abbé refractometer

abiogenesis

abiotic

ablation

ablation till

ablation zone

aboral

abrasion

abrasion ramp

absarokite

absolute age

absolute humidity

absolute plate motion

absolute pollen frequency

absolute porosity

absolute temperature

absolute vorticity

absolute zero

absorptance  
absorptance band  
absorption  
abstraction  
Abukama-type metamorphism  
abundance zone  
ABW  
abyssal hills  
abyssal plain  
abyssal storm  
abyssal zone  
Acadian orogeny  
Acado-Baltic Province  
acanthite  
acanthodians  
Acanthodii  
Acanthograptidae  
Acanthostega  
acceleration  
acceleration, gravitational  
accelerograph  
accelerometer  
accessory, lithic  
accessory cloud  
accessory mineral  
accessory plate  
accidental lithic  
accommodation space  
accommodation zone  
accordion fold  
accretion  
accretionary heating  
accretionary basin  
accretionary lapilli

accretionary levée  
accretionary prism  
accretionary wedge  
accumulated temperature  
accumulation zone  
ACD  
ACF  
ACF diagram  
achnelith  
achondrite  
achromatic line  
acicular  
acid  
acidophile  
acid rain  
acid rock  
acid soil  
acclinic line  
acme zone  
acoustic basement  
acoustic impedance  
acquired characteristics  
ACRIMSAT  
acrisols  
acritarchs  
Acrothoracica  
acrozone  
actinides  
actinium series  
actinoids  
actinolite  
Actinopterygii  
activation analysis  
activation energy

Active Cavity Radiometer Irradiance Monitor Satellite  
active geophysical methods  
active layer  
Active Magnetosphere and Planetary Electrodynamics Response  
Experiment  
active margin  
active methods  
active pool  
active remote sensing  
activity  
activity coefficient  
Actonian  
actual evapotranspiration  
actualism  
acuity  
ACV  
Adam  
adamantine  
adamellite  
Adams–Williamson equation  
adapical  
adaptation  
adaptive radiation  
adaptive zone  
addition rule  
additive primary colours  
adductor muscles  
Adelaidean  
Adelaidean orogeny  
ADEOS  
adhesion ripples  
adhesion warts  
adiabat  
adiabatic

adit  
admission  
adobe  
adoral  
Adrastea  
adsorption  
adsorption complex  
adularia  
Advanced Earth Observing Satellite  
Advanced Land Observing Satellite  
Advanced Satellite with New System Architecture for Observation  
advection  
adventive cone  
adventurine  
AE  
aedificinia  
aegirine  
Aegyptopithecus zeuxis  
AEM-2  
Aeneas  
aeolian abrasion  
aeolianite  
aeolian processes  
aeolian ripple  
Aeolis Quadrangle  
Aeolus  
aerial photograph  
aerial photography  
aerobic  
aerodynamic roughness  
aerological diagram  
aeromagnetic survey  
Aeronian  
Aeronomy of Ice in the Mesosphere

aerosol  
Aëtosauria  
AFC  
AF demagnetization  
affine  
AFM diagram  
AFMAG EM system  
African Plate  
Afternoon Constellation  
aftershock  
afterslip  
Aftonian  
Agassiz, Jean Louis Rodolphe  
agate  
age  
ageostrophic wind  
agglomerate  
agglutinate  
aggradation  
aggregate  
aggregate abrasion value  
aggregate crushing value  
aggregate impact value  
aggregate tests  
aggregation  
Aglaophyton major  
Agnatha  
Agnostida  
agonic line  
agric horizon  
agrichnion  
Agricola, Georgius  
agrometeorology  
AGU

Agulhas current  
ahermatypic  
AIM  
AIPG  
airborne dust analysis  
airborne gravity survey  
airgun  
air-lift pump  
air mass  
air wave  
Airy, George Biddell  
Airy model  
Airy phase  
AISat  
Aitken nuclei counter  
Aitken nucleus  
AIV  
AIW  
åkermanite  
A'KF diagram  
aklé  
Aksayan  
aktuopalaeontology  
alabaster  
alar  
alas  
Alaska current  
A-layer  
albedo  
albedo feature  
albeluvisols  
Alberta low  
Albertan  
Albian

albic  
albite  
albite–epidote–amphibolite facies  
albite twin  
albitization  
alcove  
alcrete  
Aldingan  
alete  
Aleutian current  
Aleutian low  
Aleutian Trench  
Alexandrian  
alexandrite  
ALEXIS  
alfisols  
Alfvén waves  
alga  
algal bloom  
algal limestone  
algal mat  
Algeria Satellite  
alginite  
Algoma-type granular iron formation  
Algonkian  
aliquot  
aliasing  
alisols  
alkali–aggregate reaction  
alkali basalt  
alkalic  
alkali-calcic series  
alkalic series  
alkali feldspar

alkaline  
alkaline rock  
alkaline soil  
alkaliphile  
alkenones  
allanite  
Alleghanian orogeny  
allele  
allelomorph  
Allen's rule  
Allerød  
allochem  
allochemical  
allochthon  
allochthonous  
allochthonous terrane  
aloclast  
allocyclic mechanisms  
allogenic  
allogenic stream  
allogroup  
allomember  
allometry  
allopatric speciation  
allopatry  
allophane  
allostratigraphic units  
allostratigraphy  
allotriomorphic  
allotrope  
allotropy  
allowable bearing pressure

alluvial  
alluvial cone  
alluvial fan  
alluvium  
almandine  
alnöite  
ALOS  
alpha decay  
alpha diversity  
alpha-mesohaline water  
alpha-proton-X-ray spectrometer  
alpine glow  
Alpine-Himalayan orogeny  
Alportian  
alteration  
alteration halo  
alternating current  
alternating-magnetic-field demagnetization  
altiplanation  
altocumulus  
Altonian  
altostratus  
alumstone  
alunite  
alveolus  
ALVIN  
A/m  
Amalthea  
Amarassian  
Amasia  
Amazonian  
amazonite  
amazonstone  
amb

amber  
ambient pressure  
ambient temperature  
ambitus  
ambulacral  
ambulacral groove  
ambulacrum  
Ambulocetus natans  
amensalism  
American Geophysical Union  
American Institute of Professional Geologists  
American Province  
Amersfoort  
amesite  
amethyst  
Amgan  
amino acid  
amino group  
ammonites  
Ammonoidea  
ammonoids  
amnion  
amniotic  
Amontons' laws of friction  
amorphous clay  
amorphous cloud  
amosite  
AMPERE  
amperes per metre  
Amferer subduction  
Amphibia  
amphibians  
amphiboles  
amphibolite

amphibolite facies  
amphicoelous  
Amphicyonidae  
amphidetic  
amphidromic point  
Amphineura  
amphitheatre  
amplitude  
ampulla  
amygdale  
amygdaloidal  
amygdule  
An  
anabatic wind  
anabranching channel  
anacline  
anaerobic  
anaerobic ammonium oxidation  
anafrost  
anagenesis  
analcime  
analcite  
analcite-basanite  
analcitite  
analog data  
analogous structures  
analogue image  
analyser  
anammox  
anamniotic  
Ananke  
Anapsida  
Anaspida  
anastomosing channel

anatase  
anatexis  
Anatolepis heintzi  
anchialine  
anchimetamorphism  
anchor  
ancient biomolecule  
Ancient Cratered Terrain  
andalusite  
ANDE  
Andean orogenic belt  
andesine  
andesite  
andhis  
andic horizon  
Andino-type margin  
andisols  
andosols  
andradite  
anelastic  
anemometer  
anemometry  
aneroid barometer  
Angara  
Angaraland  
angiophyte  
angiosperm  
Angiospermae  
angle of draw  
angle of incidence  
angle of internal friction  
angle of reflection  
angle of refraction  
angle of repose

angle of shearing resistance  
anglesite  
Anglian  
ångstrom  
angularity number  
angular momentum  
angular shear strain  
angular unconformity  
angular velocity  
anhedral  
anhydrite  
anhysteretic magnetization  
Animalia  
Animikian  
anion  
Anisian  
Anisograptidae  
anisometric growth  
anisomyarian  
anisotropic  
anisotropic meter  
anisotropy  
ankerite  
Ankylosaurus  
ankylosaurs  
Anna University Microsatellite  
annealing  
Annelida  
Anning, Mary  
annual snow-line  
annulus  
anode  
anomalous lead  
anomaly

anomphalous  
anorogenic  
anorthite  
anorthoclase  
anorthosite  
anoxic  
Antarctic air  
Antarctic bottom water  
Antarctic Circumpolar Current  
Antarctic Coastal Current  
Antarctic convergence  
Antarctic front  
Antarctic intermediate water  
Antarctic meteorites  
Antarctic Ocean  
Antarctic Plate  
Antarctic polar current  
Antarctic polar front  
ante-  
antecedent drainage  
anterior  
Anthocyathea  
anthophyllite  
Anthophyta  
Anthozoa  
anthracite  
Anthracosauria  
anthropogenic  
anthropogeomorphology  
Anthropoidea  
anthrosols  
anti-  
Antian  
anticlinal trap

anticlinal valley  
anticline  
anticlinorium  
anticoincidence circuit  
anticyclogenesis  
anticyclolysis  
anticyclone  
anticyclonic gloom  
antidune  
antiferromagnetic  
antiform  
antigorite  
antimonite  
antimony, native  
antimony glance  
antiperthite  
antipode  
antithetic fault  
antitrade  
ANUSat  
anvil  
apatite  
Apatosaurus  
ape  
aperture  
apex  
APF  
aphanitic  
Aphebian  
aphelion  
aphyric  
API gravity  
aplite  
apodeme

apogee  
Apollo  
apomorph  
apomorphic  
apophysis  
Appalachian orogenic belt  
apparent age  
apparent cohesion  
apparent conductivity  
apparent dip  
apparent polar wander  
apparent resistivity  
apparent velocity  
apparent wavelength  
appinites  
AprizeSat-3 and -4  
apsacine  
aptation  
Aptian  
aptychus  
APW  
APXS  
Aqua  
aquamarine  
Aquarius  
aquiclude  
aquic moisture regime  
aquifer  
aquifer test  
aquifuge  
Aquitania  
aquitard  
Arabian Plate  
arachnid

Arachnida  
arachnid structure  
arachnoids  
aragonite  
aragonite compensation depth  
aragonite mud  
Aratauran  
arborescent  
arc  
archae-  
Archaea  
Archaean  
archaeobacteria  
Archaeocalamites radiatus  
Archaeoceti  
Archaeocyatha  
Archaeogastropoda  
archaeological geology  
archaeomagnetism  
Archaeopteris  
Archaeopteryx lithographica  
archaeopyle  
Archaeosperma arnoldii  
Archaeosphaeroides  
Archaeozoic  
archaic sapiens  
arche-  
archetype  
Archie's law  
arching  
archipelago  
architecture of sandbodies  
archosaur  
Archosauria

## Arcsecond Space Telescope Enabling Research in Astrophysics

arctic air

arctic bottom water

arctic front

Arctic Ocean

arctic sea smoke

arc-trench gap

arcus

Ardipithecus ramidus

Arduino, Giovanni

areal erosion

arenaceous

Arenicolites

Arenig

arenite

arenosols

areology

arête

arfvedsonite

argentite

argic horizon

argillaceous

argillaceous limestone

argillans

argillic horizon

argillite

Argo

argon–argon dating

argon-40

Argos DCS

aridic moisture regime

aridisols

aridity index

Ariel

aristogenesis  
Aristotle's lantern  
arkose  
arkosic arenite  
arkosic wacke  
arls  
Arnsbergian  
Arowhanan  
array  
Array of Low-Energy X-Ray Imaging Sensors  
arrow worms  
arroyo  
arsenopyrite  
artesian water  
artesian well  
Arthrodira  
Arthropoda  
arthropods  
Articulata  
artificial freezing  
artificial rain  
artificial recharge  
Artinskian  
Artiodactyla  
Arundian  
Asaphida  
asbestos  
Asbian  
aseismic  
aseismic margin  
aseismic ridge  
ash  
ash-cloud surge  
ash cone

ash-flow  
Ashgill  
asiderite  
ASIM  
ASNARO  
asperitas  
asphalt  
asphaltite  
assay  
Asselian  
assemblage zone  
assimilation  
assimilation-fractional crystallization  
astatic magnetometer  
ASTERIA  
Asteriacites  
asteroid  
Asterosoma  
Asteroxylon  
Asterozoa  
asthenosphere  
astogenetic heterochrony  
astra  
astraeoid  
astragalus  
astrobleme  
astrogeology  
astronomical unit  
astrum  
A-subduction  
asymmetrical fold  
asymmetric valley  
Atdabanian  
Athabasca

Atlantic  
Atlantic conveyor  
Atlantic Ocean  
Atlantic Province  
Atlantic-type coast  
Atlantic-type margin  
Atlantis Massif  
Atlas  
atlas vertebra  
atmometer  
atmophile  
atmosphere  
Atmosphere-Space Interactions Monitor  
Atmospheric Neutral Density Experiment  
atmospheric pollution  
atmospheric pressure  
atmospheric shimmer  
atmospheric structure  
atmospheric 'window'  
Atokan  
atoll  
atollon  
atomic absorption spectrometry  
atomic number  
A-train  
attapulgate  
attenuation  
Atterberg limits  
attitude  
atritus  
AU  
aubrite  
augen-gneiss  
auger

augite  
augite-minette  
Aulacocerida  
aulacogen  
aulodont  
Aura  
aureole  
aurora  
Australian faunal realm  
australopithecines  
Australopithecus  
autapomorphy  
authigenic  
autobrecciated lava  
autochthonous  
autoclast  
autocorrelation  
autocyclic mechanisms  
automatic point counter  
automatic weather station  
autometamorphism  
autometasomatism  
AutoNaut  
autosuspension  
autotheca  
Autunian  
autunite  
auxiliary reference section  
available nutrients  
available relief  
available water  
avalanche  
avalanche wind  
Avalonian orogeny

aventurescence  
aventurine  
average  
average velocity  
Aves  
Avicenna  
Avogadro constant  
avulsion  
Awamoan  
axial modulus  
axial plane  
axial plane cleavage  
axial ratio  
axial rift  
axial surface  
axial tilt  
axial trace  
axial trough  
axinite  
axiolitic structure  
axis of rotation  
axis of symmetry  
axis vertebra  
Ayusokkanian  
azimuth  
azimuthal distribution  
azimuth resolution  
Azoic  
Azores high  
azurite  
b  
B  
Bacillariophyceae  
back-arc basin

back-arc spreading  
backing  
backreef  
backscatter  
backshore  
backswamp  
back thrust  
backwash  
Bacteria  
bacterial chemosynthesis  
badlands  
Badr-B  
bafflestone  
baguio  
bahada  
Bairnsdalian  
Bai-u season  
bajada  
Bajocian  
balanced sections  
Balanidae  
Balcombian  
Balfour  
ball and pillow structure  
ball clay  
balloon sounding  
Baltica  
Baltoscandia  
band  
banded iron formation  
band filter  
band-pass filter  
band-reject filter  
band silicate

bank calving  
bankfull flow  
bankfull stage  
banner cloud  
bar  
Baragwanathia longifolia  
barat  
barchan  
barchanoid  
barite  
barkevikite  
barnacles  
baroclinic  
baroduric  
barograph  
barometer  
barothermograph  
barotropic  
barred basin  
barrel  
Barrell, Joseph  
barrel trend  
Barremian  
barren interzone  
barren intrazone  
barrier  
barrier bar  
barrier beach  
barrier island  
barrier reef  
Barrovian-type metamorphism  
Barrow's zones  
Bartonian  
barycentric reference frame

baryte  
basal conglomerate  
basal sliding  
basalt  
basal thrust  
basaltic meteorites  
basanite  
base  
basecourse  
baseflow  
baselap  
base level  
basement  
base saturation  
base station  
base surge  
Bashkirian  
basic  
basic rock  
basic soil  
Basilosaurus  
basin  
basin-and-range crustal type  
basin-and-range province  
basin-and-swell sedimentation  
basin modelling  
Basleoan  
bastnäsité  
Batesfordian  
batholith  
Bathonian  
bathy-  
bathyal zone  
bathymetry

bathymograph  
bats  
Batyrbayan  
Bauplan  
bauxite  
Bavarian  
bay bar  
Bayesian  
bayhead barrier  
bayhead beach  
baymouth barrier  
bayou  
Bazin's average velocity equation  
bbl  
beach  
beach cusp  
beach drift  
beach rock  
beaded lightning  
beak  
BEAMS  
beardworms  
bearing capacity  
Beaufort scale  
Beche, Henry Thomas de la  
Becke, Friedrich Johann Karl  
Becke line test  
bed  
bedding  
bedding foliation  
bedding plane  
bedform  
bed load  
bed roughness

Beekmantownian  
Beestonian  
belemnites  
Belemnitida  
Belinda  
bellerophontiform  
bell pit  
bell trend  
bench  
Bendigonian  
beneficiate  
Benguela current  
Benioff, Hugo  
Benioff zone  
benmoreite  
Bennettiales  
Benson's flood-peak formula  
benthic storm  
benthos  
bentonite  
BepiColombo  
Bergen School  
Bergeron theory  
Bergman, Torbern Olof  
Bergmann's rule  
bergschrund  
berg wind  
Beringia  
Bering land bridge  
berm  
Berman balance  
Bermuda high  
Bernoulli, Daniel  
Bernoulli equation

Berriasian  
bertrandite  
Bertrand lens  
beryl  
beta decay  
beta diagram  
beta-mesohaline water  
bevelled cliff  
Bianca  
biaxial interference figure  
Bibymalagasia  
biconical  
BIF  
bifurcation  
bifurcation ratio  
big bang theory  
bilateral symmetry  
Bilateria  
billow clouds  
bilophodonty  
bimodal distribution  
bin  
binary system  
bindstone  
Bingham fluid  
binomial distribution  
bio-  
biochemical oxygen demand  
biochron  
biochronology  
bioclast  
biocoenosis  
biofacies  
biogenesis

biogenetic law  
biogenic  
biogenic deposit  
biogeochemical cycle  
biogeochemical exploration  
biogeochemistry  
biogeocoenosis  
biogeography  
biogeomorphology  
bioglyph  
bioherm  
biohorizon  
bioimmuration  
biointermediate elements  
biolimiting elements  
biolithite  
biological marker  
biological oxygen demand  
biomagnetism  
biomarker  
biome  
biomicrite  
biomineralization  
biophile  
biosparite  
biosphere  
biostratigraphic interval zone  
biostratigraphic unit  
biostratigraphic zone  
biostratigraphy  
biostratinomy  
biostrome  
biota  
biotic

biotic index  
biotite  
bioturbation  
biounlimiting elements  
biozone  
bipolar distribution  
bipolar seesaw  
bipyramid  
biramous  
bird  
birdseye fabric  
bireflectance  
birefringence  
birefringence chart  
BIRPS  
Birimian orogeny  
bischofite  
bise  
biserial  
bismuth, native  
bismuthinite  
bistatic radar  
bisulcate  
bit  
Bitauanian  
bitheca  
bitter lake  
bittern salts  
bitumen  
bituminous coal  
bivalves  
Bivalvia  
Bjerknes, Vilhelm Frimann Koren  
black body

black earth  
Blackett, Patrick Maynard Stuart  
black ice  
black jack  
Blackriverian  
black shale  
BlackSky Constellation  
'black smoker'  
blade  
bladed  
Blake  
blanket bog  
-blast  
blasto-  
blastoporphyratic  
Blastozoa  
blast ratio  
B-layer  
bleb  
bleicherde  
blind hole  
blind pores  
blind thrust  
blind valley  
blind zone  
blizzard  
block-and-ash deposit  
blockfield  
block glide  
blocking  
blocking anticyclone  
blocking high  
blocking temperature  
blocking volume

block volume  
blocky lava  
bloedite  
blood rain  
blow-hole  
blow-out  
blue  
blueberries  
blue-green algae  
blue jet  
Blue John  
blue Moon  
blueschist  
blueschist facies  
blue starter  
blue Sun  
bocca  
BOD  
Bode, Johann Elert  
body chamber  
body plan  
body wave  
boehmite  
Bogen structure  
boghead coal  
bog iron ore  
bolide  
Bolindian  
Bølling-Allerød interstadial  
bolson  
Boltwood, Bertram Borden  
Boltzmann constant  
bomb  
bomb sag

bone  
bony fish  
Boomer  
Boomerangian  
bootstrapping  
bora  
borax  
bord and pillar  
border fault  
bore  
boreal  
Boreal  
boreal climate  
Boreal realm  
borehole  
borehole effect  
borehole logging  
borehole sonde  
boring  
bornhardt  
bornite  
bort  
Bortonian  
bortz  
boss  
Bothriocidaroida  
Botomian  
botryoidal  
bottleneck  
bottomset beds  
bottom water  
boudin  
boudinage  
Bouguer, Pierre

Bouguer anomaly  
Bouguer correction  
Bouguer gravity map  
boulder  
boulder clay  
Bouma sequence  
bounce mark  
boundary current  
boundary layer  
boundary stratotype  
boundary wave  
boundary zonal  
boundstone  
bourn  
bourne  
bournonite  
bow-and-arrow rule  
Bowen, Norman Levi  
Bowen's ratio  
Bowen's reaction principle  
Bowen's reaction series  
bow shock  
bow-tie reflection  
bow trend  
boxcar trend  
box classification  
box fold  
BP  
brachia  
brachial plates  
brachidium  
brachiole  
Brachiopoda  
Brachiosaurus

brachium  
brachydont  
bradyseism  
bradytelic evolution  
Bragg, William Lawrence  
Bragg equation  
Bragg's law  
braided stream  
branch  
branchial arch  
branchial basket  
branching decay  
Brandenburg  
Brandon  
braunerde  
braunite  
Bravais lattice  
Bravais law  
Bravais rule  
Brazil current  
Brazilian CubeSat Project-1  
bread-crust bomb  
break  
break-back thrust  
breaker  
breccia  
breccio-conglomerate  
brecciola  
breeze  
breviconic  
brick  
brickearth  
Bridgerian  
Bridgman, Percy Williams

bridgmanite  
bridgmanite-enriched mantle structure  
Brigantian  
bright spot  
Brillouin spectroscopy  
brine  
Brioverian  
bristlecone pine  
British classification  
British Geological Survey  
brittle  
brittle behaviour  
Brøgger, Waldemar Christofer  
Brongniart, Alexandre  
Brontosaurus  
bronzite  
brookite  
Brørup  
brown clay  
brown coal  
brown earth  
brown forest soil  
Brownian motion  
brown podzolic soil  
brown soil  
brucite  
Brückner cycle  
Brunhes  
Bruun rule  
Bryophyta  
Bryozoa  
B-subduction  
bubble pulse  
bubnoff unit

Buchan metamorphic zones  
Buchan spells  
Buckland, William  
buckle folding  
Buffon, Georges Louis Leclerc, Comte de  
Buganda-Toro-Kibalian orogeny  
bulb of pressure  
Bulitian  
bulk composition of Earth  
bulk density  
bulking  
bulk minerals  
bulk modulus  
Bullard, Edward Crisp  
bumpiness  
Buntsandstein  
buoyancy  
buran  
Burdigalian  
Burgers model  
Burgess Shale  
burial metamorphism  
buried soil  
buried topography  
burn  
burner reactor  
Burnet, Thomas  
burrow  
'burst of monsoon'  
Burzyan  
butte  
Buys Ballot's law  
Byerlee's rule  
by-product

byssate  
byssus  
byte  
bytowntite  
b-zone  
14C  
cable drilling  
cadicone  
CADRE  
Caerfai epoch  
Cainozoic  
cairngorm  
cake  
Calabrian  
calamine  
Calamites cistiiformes  
calc-alkaline  
Calcarea  
calcarenite  
calcareous ooze  
calcareous soil  
calceolid  
Calcichordata  
calcichordates  
calcic horizon  
calcic series  
calcification  
calcilutite  
calcirudite  
calcisiltite  
calcisols  
calcispheres  
Calcispongea  
calcite

calcium feldspar  
calcrete  
calcrete uranium  
calcsilicate  
calc-sinter  
caldera  
Caledonian orogeny  
Caliban  
calibration graph  
caliche  
calichnia  
California bearing ratio  
California current  
caliper log  
CALIPSO  
Callisto  
Callovia  
calm  
Calorian  
calving  
calvus  
calyce  
Calymmian  
Calypso  
Calyptomatida  
calyx  
calyx drilling  
camara  
camaral  
cambering  
cambic horizon  
cambisols  
Cambrian  
camera

cameral  
cameral fluid  
Camerata  
camouflage  
Campanian  
Campbell–Stokes sunshine recorder  
camptonite  
Canada balsam resin  
Canadian  
'canali'  
canalizing selection  
Canaries current  
cancrinite  
Canidae  
Caniformia  
cannel coal  
cannel shale  
cannonball bomb  
canyon  
cap  
capacity  
capillarity  
capillary action  
capillary fringe  
capillary moisture  
capillary water  
capillary wave  
capillary zone  
capillatus  
Capitanian  
cap rock  
Captorhinomorpha  
capture  
capuliform

Caradoc  
carbon  
carbon-14  
carbonaceous chondrite  
carbonate compensation depth  
carbonate lump  
carbonate platform  
carbonate ramp  
carbonates  
carbonation  
carbonatite  
carbon 'burning'  
carbon cycle  
carbon dating  
Carbon Dioxide Monitoring Mission  
Carboniferous  
carbon isotopes  
carbonization  
cardinalia  
cardinal septum  
cardinal tooth  
Caribbean current  
Caribbean Plate  
carina  
Carlsbad twin  
Carlsberg Ridge  
Carne  
carnallite  
carnassial  
Carnian  
Carnivora  
carnosaur  
carnotite  
carpal

Carpentarian  
Carpenter, William Benjamin  
carpoids  
carrier element  
Cartesian projection  
cartilage  
cartilaginous fish  
CartoSat  
Cascade SmallSat and Ionospheric Polar Explorer  
cascading system  
casing  
Cassadagian  
Cassini  
Cassini Division  
CASSIOPE  
cassiterite  
cast  
Castalia  
castellanus  
Castlecliffian  
castle koppie  
Castlemainian  
CAT  
cataclasis  
cataclasite  
catagenesis  
cataracta  
cataractagenitus  
catarrhine  
catastrophic evolution  
catastrophism  
catazone  
catchment  
catena

cateniform  
cathode  
cathodoluminescence  
cation  
cation exchange  
cation-exchange capacity  
cation ordering  
CATS  
cat's eye  
cauda  
caudal  
caudal vertebra  
cauldron-subsidence  
Cautleyan  
cavate  
cavern porosity  
cavitation  
cavum  
cavus  
cay  
Cayugan  
Cazenovian  
CBERS  
CBR  
CCD  
CCL  
CCN  
14C dating  
CDP  
CDP stack  
CEC  
Celastrophyllum circinerve  
celerity  
celestial reference frame

celestite  
cement  
cementation  
cemented  
Cenomanian  
Cenozoic  
centipedes  
Centrales  
Central European Sea  
central limit theorem  
central vent volcano  
centre of curvature  
centre of symmetry  
centric diatoms  
centrifugal pump  
centripetal drainage pattern  
Centroceratida  
centrum  
Cephalaspida  
Cephalaspis  
cephalic  
cephalic spine  
cephalic suture  
Cephalochordata  
cephalon  
Cephalopoda  
ceratites  
ceratoid  
Ceratopsia  
Ceres  
cerioid  
cerussite  
cervical vertebra  
CFESat

CFOSAT  
c.g.s. system  
Chadian  
Chaetognatha  
chain-former  
chain lightning  
chain-modifier  
chain silicate  
chalcedony  
chalcocite  
chalcophile  
chalcopyrite  
chalk  
Challenger expedition  
Challenging Minisatellite Payload  
chalybdite  
chalybeate  
Chamberlin, Thomas Chrowder  
Chambers, Robert  
chamosite  
Chamovnicheskian  
CHAMP  
Champlainian  
Chandler wobble  
Changhsingian  
Changlangpuan  
channel  
channel and vug porosity  
Channel Deposits  
channel fill  
channel wave  
chaos  
char  
character

character states  
chara marl  
charcoal  
charge  
chargeability  
charge-coupled device  
charged-body potential method  
Charnian  
Charniodiscus  
charnockite  
charnockitic gneiss  
Charon  
Charophyceae  
Charophyta  
Charpentier, Jean de  
chart datum  
chasma  
chatoyance  
chatoyancy  
chattermark  
Chattian  
Chautauquan  
Chazyan  
Chebotarev sequence  
Cheirolepis trailli  
chelation  
Chelford  
cheliceræ  
Chelicerata  
Cheltenhamian  
chemical demagnetization  
chemical fossil  
chemical garden  
chemical layering

chemical oxygen demand  
chemical potential  
chemical remanent magnetization  
chemical weathering  
chemocline  
chemofossil  
chemostratigraphy  
chemosymbiosis  
chemosynthesis  
chenier  
chenier plain  
Cheollian-1  
Cheremshanskian  
chernozem  
chert  
Chesterian  
chevron fold  
chevron marks  
Chewtonian  
Chezy's formula  
chiastolite  
Chibis-M  
chickenwire structure  
Chicxulub  
Chile Rise  
Chile saltpetre  
chilidial plates  
chilidium  
chilled edge  
chilled margin  
Chilopoda  
China–Brazil Earth Resources Satellite  
china clay  
China Seismo-Electromagnetic Satellite

chinastone  
chine  
Chinese–French Oceanography Satellite  
chinook  
chip sampling  
Chiron  
Chiroptera  
chi-squared test  
chitin  
Chitinodendron franconianum  
chloralgal  
chlorinity  
chlorite  
chloritoid  
Chlorophyta  
chlorozoan  
Choanichthyes  
Chokerian  
Chokierian  
Chondrichthyes  
chondrite  
chondrite model  
Chondrites  
chondritic Earth model  
chondritic unfractionated reservoir  
chondrocranium  
chondrodite  
chondrophore  
chondrosteans  
Chondrostei  
chondrule  
Choquette and Pray classification  
chord  
Chordata

chroma  
chromatid  
chromatography  
chrome diopside  
chromite  
chromosome  
chron  
chronohorizon  
chronomere  
chronometry  
chronosequence  
chronosome  
chronospecies  
chronostratic scale  
chronostratigraphic correlation chart  
chronostratigraphic horizon  
chronostratigraphic scale  
chronostratigraphic unit  
chronostratigraphy  
chronozone  
chrysoberyl  
chrysocolla  
Chrysophyceae  
chrysotile  
ChubuSat-1  
CHUR  
Churchillian orogeny  
Cibola Flight Experiment Satellite  
CICERO  
cilia  
cilium  
Cincinnatian  
cinder cone  
CINEMA

cingulum  
cinnabar  
CIPW norm calculation  
CIRAS  
circalittoral zone  
circularity index  
circularity ratio  
circular polarization  
circulation index  
circum-oral canal  
cirque  
cirque glacier  
cirri  
Cirripedia  
cirrocumulus  
cirrostratus  
cirrus  
citrine  
clade  
cladism  
cladistic analysis  
cladistics  
cladogenesis  
cladogram  
Cladoselache  
Cladoselachiformes  
Clapeyron–Clausius equation  
Clapeyron equation  
clarain  
Clarence  
Clarke, Frank Wigglesworth  
Clarke, William Branwhite  
Clarke orbit  
Clarkfordian

classification  
clast  
clastic  
clastic dyke  
clastic rock  
clastogenic flow  
clathrate  
Clausius–Clapeyron equation  
Clavatipollenites  
clay  
clay dune  
C-layer  
clay films  
clay minerals  
clay pan  
clayskins  
claystone  
cleaning, magnetic  
cleaning-up trend  
clear-air turbulence  
clear ice  
cleat  
cleavage  
cleavage refraction  
Clementine  
Cliffdenian  
CLIMAP  
CLIMAPP  
climate classification  
Climate–Leaf Analysis Multivariate Program  
Climate/Long-ranged Investigation Mapping and Predictions Project  
climatic geomorphology  
climatic optimum  
climatic station

climatic zone  
Climatiiformes  
climatostratigraphy  
climax trace fossil  
climbing dune  
climbing-ripple cross-lamination  
climosequence  
cline  
clino-  
clinoclone  
clinoform  
clinohumite  
clinometer  
clinopyroxene  
clinosequence  
clinothem  
clinozoisite  
clint  
Clinton ironstone  
clipped trace  
clitter  
clod  
closed form  
closed system  
close fold  
closure age  
closure temperature  
Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations  
Cloud-Aerosol Transport System  
cloud amount  
cloud base  
cloudburst  
cloud classification  
cloud condensation nuclei

cloud discharge  
cloud droplet  
CloudSat  
cloud seeding  
cloud street  
Cluster  
cluster analysis  
CMP  
Cnidaria  
CNMMN  
CNMNC  
C/NOFS  
co-  
co-adaptation  
Coahulian  
coal  
coalescence  
coalification  
coal lithotype  
coal maceral  
coal-maceral group  
coal measures  
coal series  
coarsening-upward succession  
coastal onlap  
coastal processes  
coastal toplap  
coated stone  
co-axial correlation  
cobalt glance  
cobaltite  
cobble  
Coble creep  
coccolithophorids

coccoliths  
Cochiti  
COCORP  
Cocos Plate  
COD  
coefficient of compressibility  
coefficient of consolidation  
coefficient of nivity  
coefficient of sliding friction  
coefficient of variation  
coelacanth  
Coelacanthiformes  
Coelenterata  
coelom  
coelomate  
Coelophysis  
Coelurosauria  
coelurosaurs  
Coenopteridales  
coenosteum  
coenozone  
coercivity, magnetic  
coesite  
co-evolution  
cognate lithic  
cogwheel ore  
coherence  
cohesion  
cohort  
coign  
co-ignimbrite breccia  
coiling  
coke  
col

colatitude  
cold front  
cold-front clearance  
cold glacier  
cold low  
cold pole  
cold pool  
cold sector  
cold seep  
cold wave  
cold working  
colemanite  
coliform bacteria  
coliform count  
collagen  
colles  
collinite  
collision theory  
collision zone  
colloform banding  
colloid  
collophane  
colluvial  
colonization window  
colonnade  
Colorado Student Space Weather Experiment  
colorimetric analysis  
colour  
colour index  
Columbia  
columbite  
columella  
columnal  
columnar

columnar joint  
columnar section  
com-  
COM  
coma  
Comanchean  
comber  
combe rock  
combination trap  
combined plate margin  
Comely epoch  
comet  
Comet Nucleus Tour  
comfort zone  
comminution  
Commission on Classification of Minerals  
Commission on New Minerals and Mineral Names  
Commission on New Minerals, Nomenclature, and Classification  
Commission on Ore Mineralogy  
commissure  
common canal  
common depth point  
common-depth-point stack  
common lead  
common mid-point  
common strontium  
Communication/Navigation Outage Forecast System  
Communication, Ocean and Meteorological Satellite  
community  
Community Initiative for Continuous Earth Remote Observation  
compaction  
compaction test  
COMPASS-1  
COMPASS-2

compass clinometer  
compensation level  
competence  
competent rock  
complex  
complexation  
Complex Orbital Magneto-Plasma Autonomous Small Satellite-2  
complex twins  
component-stratotype  
composite fault-line scarp  
composite intrusion  
composite-stratotype  
composite volcano  
compositional immaturity  
compositional maturity  
composition plane  
composition surface  
compound corals  
compound twins  
compressed air  
compressibility  
compressional wave  
compressive stress  
COMS  
con-  
concealed coalfield  
concentration  
concentration factor  
concentration-Lagerstätte  
concentric fold  
conchoidal fracture  
concordant  
concordant age  
Concordia diagram

concrete  
concrete dam  
concrete minimum temperature  
concretion  
concretionary  
concurrent range zone  
condensation level  
condensation nucleus  
condensation trail  
condensed bed  
condenser  
conditional instability  
Condobolinian  
conductance  
Condylarthra  
condylarths  
cone-in-cone structure  
cone of depression  
cone penetrometer  
cone sheet  
Conewangan  
confidence interval  
confidence level  
confined aquifer  
confining pressure  
confluence  
conformable  
congelifraction  
congeliturbation  
congestus  
conglomerate  
congruent dissolution  
congruent solution  
Coniacian

Coniferales  
Coniferophyta  
Coniferopsida  
conifers  
conjugate fault set  
conjugate fold  
conjunct  
connate water  
Conodontophora  
conodonts  
conoscopic  
Conrad discontinuity  
consequent stream  
conservation-Lagerstätte  
conservative margin  
consistence  
consistency  
consistency index  
consolidation  
constant head permeameter  
constant offset  
constant-separation traversing  
constant-velocity gather  
constant-velocity stack  
Constellation Observing System for Meteorology, Ionosphere and Climate  
constructive boundary  
constructive margin  
constructive wave  
contact  
contact aureole  
contact goniometer  
contact metamorphism  
contact resistance  
contact twin

contained fragments  
contessa del vento  
continental air  
continental crust  
continental drift  
continental freeboard  
continentality  
continental margin  
continental rise  
continental shelf  
continental-shelf waves  
continental shield crustal type  
continental slope  
continuation  
continuous distribution  
continuous profiling  
continuous reaction series  
continuous velocity logging  
CONTOUR  
contour current  
contour diagram  
contourites  
contracting Earth hypothesis  
contractional kink band  
contraction limit  
contrail  
contrast  
contrast stretching  
control system  
convection  
convection current  
convective cell  
convective condensation level  
convective instability

convergence  
convergent evolution  
convergent margin  
converted wave  
convex slope  
convolute  
convolute lamination  
convolution  
Conybeare, William Daniel  
Cook, James  
Cooksonia hemispherica  
cooling age  
cooling joint  
coombe rock  
coordinated stasis  
coordination number  
COP  
Copernican System  
Copernicus  
Cope's rule  
copper, native  
copper glance  
copper pyrites  
coppice dune  
co-product  
coprolite  
coquina  
coral  
coral growth lines  
coralline limestone  
corallite  
coralloid  
corallum  
Cordaitales

Cordelia  
cordierite  
cordillera  
core  
core barrel  
core-logging  
core recovery  
core slicer  
corestone  
core wall  
Cor F  
Coriolis effect  
Coriolis parameter  
Coriolis/WindSat  
corona  
corrasion  
correlated progression  
correlation  
correlation diagram  
correlogram  
corridor dispersal route  
corrie  
corundum  
Corynexochida  
coseismic period  
coset  
COSMIC  
cosmic abundance of elements  
cosmic dust  
cosmic radiation  
cosmic-ray track  
cosmine  
cosmochemistry  
cosmoid scale

cosmology  
cosmopolitan distribution  
Cosmorhaphé  
COSMO-SkyMed  
costa  
Costonian  
cotectic curve  
cotectic surface  
coterminous  
cotidal line  
Cotylosauria  
coulée flow  
Coulomb failure criterion  
Coulomb–Terzaghi shear strength equation  
coultter counter  
counterpoised beam balance  
country rock  
couple  
coupled substitution  
covalent bond  
covalent compound  
covalent radius  
covariance  
covellite  
cow-dung bomb  
coxa  
cpx  
crabs  
crachin  
cracking  
crag  
crag and tail  
Craniata  
cranidium

cranium  
crater  
crater counting  
crater density studies  
Cratered Plains  
Cratered units  
crateriform  
craton  
creep  
creep mechanisms  
creep strength  
Crenarchaeota  
crenulation cleavage  
Creodonta  
creodont-like teeth  
crepuscular rays  
crescent-and-mushroom  
Cressida  
crest  
crestal plane  
crest line  
Cretaceous  
crevasse  
crevasse deposit  
crevasse splay  
Crinoidea  
crinoids  
cristobalite  
critical angle  
critical damping  
critical erosion velocity  
critical flow  
critical reflection  
critical temperature

critical velocity  
CRM  
crocidolite  
Crocodilia  
Croixian  
Cromerian  
Crommelin  
Cromwell current  
cronstedite  
cross-bedding  
cross-correlation  
crosscut  
cross-cutting relationships  
cross-dating  
crossed nicols  
crossed polars  
cross-hairs  
cross-lamination  
Crossopterygii  
cross-over distance  
cross set  
cross-stratification  
cross-well seismic  
cross-wires  
crotovina  
crown group  
crude oil  
Crudinian  
crumb structure  
crura  
crus  
crushing  
crust  
Crustacea

crustaceans  
crustal abundance of elements  
Cruziana  
cryergic  
cryic  
cryoconite  
cryoconite hole  
Cryogenian  
cryogenic  
cryogeysers  
cryolite  
cryonival  
cryopediment  
cryoplanation  
Cryosat-2  
cryosols  
cryosphere  
cryoturbation  
cryovolcanism  
Cryptic  
cryptocrystalline  
cryptodome  
Cryptodonta  
cryptoperthite  
cryptotephra  
Cryptozoic  
crystal  
crystal class  
crystal face  
crystal-field theory  
crystal group  
crystalline  
crystalline carbonate  
crystalline limestone

crystalline remanent magnetization  
crystallite  
crystalloblastic  
crystallographic axes  
crystallography  
crystal symmetry  
crystal system  
crystal twinning  
crystal zoning  
CSES  
CSSWE  
CST  
CTD  
cube  
CubeRRT  
CubeSat  
CubeSat for Ions, Neutrals, Electrons & Magnetic fields  
CubeSat Infrared Atmospheric Sounder  
CubeSat – Launch 1  
CubeSat Radiometer Radio Frequency Interference Technology  
cubic  
cubichnia  
cuesta  
culmination  
cumec  
cumingtonite  
cumulate  
cumulative percentage curve  
cumuliform  
cumulonimbus  
cumulus  
Cunninghamian  
Cupid  
cupola

cuprite  
Curie point  
Curie symmetry principle  
Curie temperature  
Curie–Weiss law  
current electrode  
current meter  
Curvolithus  
cusate foreland  
cuspidate  
cutan  
cut and fill  
cuticle  
cutin  
cutinite  
cut-off  
cut-off grade  
cut-off high  
cut-off low  
cut-off trench  
cutting bar  
cutting boom  
Cuvier, Chrétien Frédéric Dagobert ('Georges'), Baron  
CVG  
CVS  
cwm  
cyanobacteria  
Cyanophyta  
cyanophyte  
cycad  
Cycadaceae  
Cycadales  
Cycadopsida  
cycle of erosion

cyclic sedimentation  
cycling pool  
cyclogenesis  
cyclolysis  
cyclone  
Cyclone Global Navigation Satellite System  
cyclopean concrete  
cyclopel  
cyclopsam  
cyclosilicate  
cyclostratigraphy  
cyclothem  
CYGNSS  
cylindrical  
cylindrical trend  
cylindroidal fold  
cymatogeny  
cyrenoid  
cyrtoconic  
Cystoidea  
cystoids  
Cytherean  
dacite  
Dactyl  
Dactylodites ottoii  
Daedalus  
dagalas  
Dalmatian-type coast  
Dalradian  
Dalslandian  
Dalslandian orogeny  
Daly, Reginald Aldworth  
DAMPE  
damping

Dana, James Dwight  
Danian  
Dansgaard–Oeschger event  
darcy  
Darcy's law  
Dark Matter Particle Explorer  
d'Arrest  
Darriwilian  
Darwin, Charles Robert  
darwin  
Dasycladales  
data  
dating errors  
dating methods  
Datsonian  
datum  
datum level  
daughter  
daughter minerals  
Davis, William Morris  
Davisian cycle  
day degrees  
day length  
dB  
D days  
dead end  
death assemblage  
débâcle  
debris fall  
debris flow  
debris slide  
debrite  
decalcification  
decay constant

decay curve  
decay index  
decay series  
Deccan Traps  
decibel  
declination  
declined  
décollement  
décollement plane  
decompression melting  
deconvolution  
decorrelation stretching  
decussate  
dedolomite  
deepening  
deep scattering layer  
Deep Sea Drilling Programme  
deep-sea fan  
deep-sea geomorphology  
deep-sea trench  
Deep Space Climate Observatory  
Deerparkian  
Defense Meteorological Satellite Program  
deflation  
deflation hollow  
deflexed  
deformation lamellae  
deformation twinning  
de Geer moraine  
degree days  
degrees of freedom  
dehydration  
dehydration curve  
Deimos

delamination  
Delaware effect  
delayed flow  
delay time  
Dellingr  
Delmontian  
delta  
delta front  
Deltatheridium  
 $\Delta T$  method  
delthyrium  
deltidial plate  
DEM  
demagnetization  
demagnetizer  
de Maillet, Benoît  
Demospongea  
Demospongiae  
demultiplexing  
dendritic  
dendritic drainage  
dendrochronology  
dendroclimatology  
dendrogeomorphology  
dendrogram  
Dendrograptidae  
dendrohydrology  
dendroid  
Dendroidea  
dendroid graptolites  
Denekamp  
Dense-medium separation  
density  
density current

density–depth profile  
density determination  
density log  
denticle  
dentine  
dentition  
denudation  
denudation chronology  
depleted mantle  
depocentre  
deposit feeder  
deposit gauge  
deposition  
depositional remanent magnetization  
depositional sequence  
depositional sequence model  
depositional system  
depositional systems tract  
depression  
depression angle  
depth point  
deranged drainage  
derived  
dermal bone  
dermal denticle  
dermatocranium  
Derryan  
desalination  
Desdemona  
desert  
desertification  
desert pavement  
desert rose  
desert varnish

desiccation  
desiccation cracks  
Desmarest, Nicolas  
desmodont  
Desmoinesian  
Despina  
desquamation  
destructive margin  
destructive wave  
detachment  
detachment control  
detachment fault  
detachment horizon  
detrital  
detrital remanent magnetization  
detritivore  
detritus  
detritus feeder  
deuteric alteration  
deuteric reaction  
Devensian  
deviatoric stress  
devitrification  
Devonian  
dewatering  
dew-point  
dew-point hygrometer  
dextral coiling  
dextral fault  
D:H ratio  
diabase  
diachronous  
diad  
diadochy

diagenesis  
diagnostic horizon  
diallage  
diamagnetism  
diamict  
diamictite  
diamicton  
diamond  
diamond drilling  
diaphragm wall  
diaphoresis  
diapir  
diapirism  
diapsid  
Diapsida  
Diarthrognathus broomi  
diaspore  
diastem  
diastrophism  
diatom  
diatomaceous earth  
diatomite  
diatom ooze  
diatrema  
DICE  
dichograptid  
Dichograptina  
dichotomize  
dichroic  
dichroism  
Dickinsonia  
dickite  
Dictyonema flabelliforme  
diductor muscle

dielectric  
dielectric constant  
dielectric loss factor  
dielectric permittivity  
Dienerian  
Dietz, Robert Sinclair  
differential settlement  
differential stress  
differentiation  
differentiation index  
diffraction  
diffuse reflection  
diffusion  
diffusion coefficient  
diffusion-controlled growth  
diffusion creep  
digenite  
digital elevation map  
digital image  
digital number  
digitize  
dihedron  
dike  
dilatancy  
dilatation  
dilatational wave  
dilation  
diluvialism  
dimension stone  
Dimetrodon angelensis  
dimorphism  
dim spot  
dimyarian  
Dinantian

dinocyst  
dinoflagellates  
Dinophyceae  
Dinophysiales  
dinosaurs  
diogenites  
Dione  
diopside  
dioptase  
diorite  
dip  
dip circle magnetometer  
dip fault  
dip-isogon method  
diphycercal tail  
Diplichnites  
Diplocraterion  
diplograptids  
Diplopleurozoa  
Diplopoda  
Diploporita  
dipmeter log  
dip moveout  
Dipneusti  
Dipnoi  
dipole field  
dip pole  
dip shooting  
dip-slip fault  
dip slope  
dipyramid  
Dirac function  
direct circulation  
direct current

directional fabric  
directional filter  
direct problem  
direct wave  
dirtying-up trend  
Disaster Monitoring Constellation for International Imaging  
discharge  
discharge hydrograph  
discoid  
disconformity  
discontinuity  
discontinuous reaction series  
discordant  
discordant age  
discordant drainage  
discordant intrusion  
Discovery Program  
disharmonic fold  
dish-pan experiment  
dish structure  
disjunct  
disjunct shell  
dislocation  
dislocation creep  
dismicrite  
dispersion  
displaced terrane  
displacement  
displacive transformation  
diss  
disseminated deposit  
dissepiment  
dissepimentarium  
dissipation trail

dissolved load  
dissolved-oxygen level  
distal  
distributary channel  
distribution coefficient  
disturbance  
disturbed days  
Ditomopyge  
Dittmar's law  
diurnal temperature variation  
diurnal variation  
divariant assemblage  
divariant equilibrium  
divaricator muscle  
divergence  
divergent margin  
diversification  
diversity  
divide  
divining  
Diwata-1  
Dix formula  
D-layer  
DMCII  
DMO  
DMSP  
Dnepr-Samarovo  
doctor, the  
documentation map  
dodecahedron  
DO event  
Dogger  
dog-tooth spar  
Dokuchayev, Vasily Vasilyevich

doldrums  
dolerite  
Dolgellian  
doline  
Dollo, Louis Antoine Marie Joseph  
Dollo's law  
dololithite  
Dolomieu, Déodat de Gratet de  
dolomite  
dolomitization  
domain  
domal uplift  
dome  
dome and basin  
dome–crescent–mushroom  
domichnia  
Dominian Reef  
Donau  
Donau/Günz Interglacial  
Doppler Orbitography and Radiopositioning Integrated by Satellite  
Doppler radar  
Doppler shift  
DORIS  
Dorogomilovskian  
dorsa  
dorsal  
dorsum  
dot chart  
Dott classification  
double core barrel  
double couple  
double planation  
double refraction  
double salt

double sulphides  
double zig-zag  
doublure  
down-hole hammer drilling  
downlap  
downthrow  
Downtonian  
downward continuation  
downwelling  
dowsing  
draa  
drag  
drainage  
drainage basin morphometry  
drainage basin relief ratio  
drainage basin shape index  
drainage density  
drainage morphometry  
drainage network  
drainage-network analysis  
drainage pattern  
drainage-sediment survey  
drainage wind  
drained test  
dravite  
drawdown  
dreikanter  
Dresbachian  
dribble cone  
drift  
drifter  
drift map  
drill hole  
drilling

drilling bit  
drilling mud  
drill string  
dripstone  
drizzle  
DRM  
dromaeosaurid  
drop ball  
dropstone  
drought  
drought cycle  
drumlin  
drumlin field  
drumlin swarm  
druse  
drusy  
dry adiabatic lapse rate  
dry air  
Dryas  
dry-bulb thermometer  
dry continental morphoclimatic zone  
dry ice  
dry melt  
dry season  
dry valley  
dry-weather flow  
DSCOVER  
DSDP  
DSL  
d-spacing  
dual decay  
DubaiSat  
ductile behaviour  
dug well

dump structure  
dune  
dune bedform  
dune slack  
Dunham classification  
dunite  
Duntroonian  
duplex  
duplicatus  
durain  
duric horizon  
duricrust  
duripan  
durisols  
durophagic  
dust  
dust-bowl  
dust detection instrument  
dust devil  
dust storm  
Dutch cone  
Du Toit, James Alexander Logie  
Dutton, Clarence Edward  
dwarf planet  
dyke  
dyke set  
dyke swarm  
dynamic correction  
dynamic correlation  
dynamic equilibrium  
Dynamic Ionosphere CubeSat Experiment  
dynamic metamorphism  
dynamic viscosity  
dysaerobic

dysodont  
E1P-2  
Early Cambrian epoch  
EARS  
Earth  
EarthCARE  
Earth Clouds, Aerosols and Radiation Explorer  
Earth compositional layering  
Earth Explorer Opportunity Mission-2  
earth fall  
earthflow  
Earth Observing System  
earth pillar  
earthquake  
earthquake accelerometer  
earthquake energy  
earthquake intensity  
earthquake magnitude  
earthquake mechanisms  
earthquake prediction  
Earth Remote Observation System  
Earth rheological layering  
Earth rotation  
Earth Science Enterprise  
Earth tides  
earthy  
East Australian current  
easterly wave  
Eastonian  
East Pacific Rise  
East Wind Drift  
ebb tide  
ebullient  
ebullition

ebullitive  
Eburonian  
Ecardines  
ecdysis  
Echinodermata  
Echinoidea  
Echmatocrinus  
echo dune  
echo-sounding  
eclipse  
ecliptic  
eclogite  
eclogite facies  
ecologic reef  
ecology  
economic basement  
economic geology  
ecophenotype  
ecophenotypic effects  
ecophenotypy  
ecosphere  
ecostratigraphy  
ECOSTRESS  
ecosystem  
ECOSystem Spaceborne Thermal Radiometer Experiment on Space Station  
ecotone  
Ectasian  
ectocochlear  
Ectoprocta  
ectotherm  
edaphic  
eddy  
eddy currents  
eddy viscosity

Edenian  
edenite  
edentulous  
edge  
edge dislocation  
edge enhancement  
Ediacaran  
Ediacaran fossils  
Eemian  
effective porosity  
effective precipitation  
effective stress  
effective temperature  
EGS  
EGT  
Egyptian jasper  
EH  
Eifelian  
Eildonian  
einkanter  
ejecta blanket  
Ekman depth  
Ekman spiral  
Ekman transport  
Elara  
Elasmobranchii  
elastic constants  
elastic deformation  
elastic limit  
elastic rebound theory  
elastic wave  
elastoviscous behaviour  
E-layer  
elbaite

elbow of capture  
electrical charge  
electrical conductivity  
electrical conductivity (in Earth)  
electrical drilling  
electrical sondes  
electrical sounding  
electrical tomography  
electrode array  
electrode configuration  
electrode potential  
electrolyte  
electrolytic conduction  
electrolytic polarization  
electromagnetic methods  
electromagnetic radiation  
electromagnetic spectrum  
electromagnetic wave  
electrometer  
electron  
electron capture  
electronegativity  
Electron Losses and Fields Investigation  
electron microscope  
electron-probe microanalyser  
electro-osmosis  
electrophoresis  
electropositive element  
Elektro  
Elektro-L  
eleutherozoan  
elevation correction  
elevation head  
elevation potential energy

elf  
ELFIN  
Élie de Beaumont, Léonce  
elite  
elliptical polarization  
ellipticity  
El Niño  
elongation index  
elongation ratio  
ELR  
Elsasser, Walter Maurice  
Elsonian orogeny  
Elsterian  
elutriation  
eluvial deposit  
eluviation  
elvan  
embankment dam  
Embolomeri  
Embry and Clovan classification  
embryophytes  
emerald  
emery  
Emilian  
emission spectrum  
emissivity  
EMR  
Emsian  
emu  
En  
en–  
enamel  
enantiomorphy  
enargite

enation theory  
Enceladus  
Encke  
Endeavour hydrothermal vents  
endemism  
endichnia  
end-member  
end-member textural classification  
end moraine  
endobiont  
endobyssate  
endochondral bone  
endocochlear  
endocone  
endogenetic processes  
endogenous dome  
endolith  
endopunctate  
endorheic lake  
endoskeleton  
endothem  
endrumpf  
en échelon  
energetic particles detector  
energy budget  
energy of activation  
engineering geophysics  
englacial  
Enhanced Fujita Scale  
enhancement seismograph  
enigmatic taxon  
EnMAP  
enriched uranium  
ensialic belt

ENSO event  
enstatite  
entablature  
entelechy  
enterolithic structure  
enteron  
enthalpy (H)  
entisols  
Entoprocta  
entrainment  
entrenched meander  
entropy  
environmental geology  
environmental lapse rate  
Environmental Monitoring and Analysis Program  
EnviSat  
Environmental Satellite  
Eoarchaeon  
Eobactrites sandbergeri  
Eocambrian  
Eocene  
Eocrinoidea  
eocrinoids  
Eodelphis  
Eoembryophytic  
'Eohippus'  
eolian  
eon  
eonothem  
EOS  
Eosimias  
Eosphaera  
Eotracheophytic  
Eötvös effect

EPD

epeiric sea

epeirogenesis

ephemeral stream

ephemerides

ephemeris

epi-

epibenthos

epibole

epibyssate

epicentral angle ( $\Delta$ )

epicentre

epichnia

epiclast

epicontinental sea

epicratonic

epidote

epifaunal

epigene

epigenesis

epigenetic drainage

epigenetic ore

epilimnion

Epimetheus

Epiphyton

episodic evolution

epitaxy

epitheca

epithermal

epoch

e-process

epsilon cross-bedding

epsomite

Epsom salts

equal-area net  
equant  
Equatorial countercurrent  
Equatorial current  
equatorial orbit  
equatorial plane  
equatorial trough  
Equatorial undercurrent  
Equidae  
equifinality  
equilibrium  
equilibrium flow  
equilibrium line  
equilibrium process  
equinoctial gale  
equipotential  
Equisetites hemingwayi  
Equisetum  
equivalence  
equivalent aperture width  
Equus  
era  
erathem  
Eratosthenian system  
e-ray  
ERG  
erg  
ergodic hypothesis  
Erian  
Eris  
EROS  
Eros  
erosion  
erosion rate

erosion surface  
erratic  
errors  
eruption  
eruptive centres  
ESA  
escape tectonics  
escape velocity  
escarpment  
escutcheon  
ESE  
esker  
Eskola, Pentti Elias  
essential mineral  
estuary  
Etalian  
Etalon  
Etched Plains  
etch figures  
etch marks  
etchplain  
etesian winds  
Ethiopian faunal realm  
ethotype  
eu-  
Eubacteria  
Eucaryota  
eucrite  
Euechinoidea  
eugeocline  
eugeosyncline  
euhaline water  
euhedral  
Eukarya

Eukaryota  
eukaryote  
Euler angles  
Eulerian current measurement  
Euler pole  
Euler's rotation theorem  
eulite  
eulysite  
EUMETSAT Advanced Retransmission Service  
Euparkeria  
euphotic zone  
Euramerica  
Eurasian Plate  
Europa  
European Federation of Geologists  
European Geosciences Union  
European GeoTraverse  
European Province  
European Space Agency  
europium anomaly  
Euryapsida  
Euryarchaeota  
euryhaline  
eurypterid  
eurythermal  
eurytopic  
eustasy  
eutaxitic structure  
eutectic point  
eutectic system  
Eutelsat 172B  
Eutheria  
Eutracheophytic  
eutrophic

euxinia  
euxinic  
EV  
evaporation  
evaporation pan  
evaporimeter  
evaporite  
evapotranspiration  
evapotron  
event deposit  
event stratigraphy  
event stratinomy  
evolute  
evolution  
evolutionary lineage  
evolutionary rate  
evolutionary species  
evolutionary trend  
evolutionary zone  
evolved magma  
Ewing, Maurice  
ex-  
ExactView  
exaerobic  
exaptation  
excavation  
exchangeable ions  
exchange capacity  
exchange pool  
exfoliation  
exhumed topography  
exichnia  
exine  
exinite

exitance  
exobiology  
ExoCube  
exocuticle  
exogenetic processes  
exogenous dome  
exorheic lake  
exoskeleton  
exosphere  
exothecal  
exotherm  
exotic  
expanding Earth  
expanding spread  
Experimental Geodetic Satellite  
Exploration of energization and Radiation in Geospace  
Explorer-1 PRIME-2  
Explorer 59  
Explorer Ridge  
explosive charge  
exposure age  
exsiccation  
exsolution  
extension  
external mould  
extinction  
extraclast  
extraction  
extraformational  
extraordinary ray  
extra-orogen basin  
extremophile  
extrusion  
extrusive

'eye' of storm  
eyepiece  
eyot  
fabric  
fabric analysis  
Fabrosaurus australis  
face pole  
facial suture  
facies  
facies association  
facies fossils  
facies sequence  
facing direction  
facula  
faculae  
faecal pellet  
fahlband  
failed arm  
failed rift  
failure  
failure strength  
failure stress envelope  
fairweather wave-base  
falling dune  
falling head permeameter  
falling stage systems tract  
falls  
fall-stripes  
false body  
false cirrus  
false colour  
famatinite  
Famennian  
family

FAMOUS project  
fan cleavage  
fanglomerate  
fan shooting  
FAO  
Farallon Plate  
far-field barrier  
farra  
farrum  
fasciculate  
fasciole  
fast breeder reactor  
fast Fourier transform  
Fast On-Orbit Recording of Transient Events  
fathom  
fatty acid  
fault  
fault block  
fault-block mountains  
fault line  
fault-line scarp  
fault outcrop  
fault plane  
fault-plane solution  
fault scarp  
fault slice  
fault trace  
fault trap  
fault zone  
fauna  
faunal province  
faunal realm  
faunal succession  
faunizone

fayalite  
Faye correction  
FDSN  
Feammox  
feather angle  
feather ore  
Federov stereographic net  
feldspars  
feldspathic greywacke  
feldspathic wacke  
feldspathoid  
Felidae  
Feliformia  
felsenmeer  
felsic  
felsite  
felsitic  
femic  
femtoplankton  
femtosatellite  
femur  
fence diagram  
fenestrae  
fenestral fabric  
fenestral porosity  
fenestrated  
FengYun  
fenite  
fenitization  
Fennoscandian Border Zone  
Fennoscandian uplift  
Ferdinand  
ferns  
ferralitization

ferralization  
ferralsols  
Ferrel cell  
ferric horizon  
ferricrete  
ferrimagnetism  
ferro-  
ferroactinolite  
ferroan  
ferroaugite  
ferroelectricity  
ferrohastingsite  
ferromagnesian minerals  
ferromagnetic  
ferromagnetism  
ferropericlase  
ferrosilite  
fertility, soil  
Ferungulata  
Festiniogian  
fetch  
Ffestiniogian  
FFT  
fiamme  
fiard  
fibratus  
fibril  
fibrolite  
fibrous  
fibula  
Fick's laws of diffusion  
fiducial point  
field capacity  
field reversal

Figtree  
Filicopsida  
filiform  
filling  
film water  
filter  
filter route  
filtrate  
filtration  
fin  
finds  
fines  
fines 10% test  
finger lake  
Fingerlakian  
fining-upward succession  
finite resource  
finite rotation  
finite strain  
fiord  
fire-ball  
fire damp  
Firefly  
fire-fountain  
firn  
firn limit  
firn line  
firn wind  
first arrival  
first break  
firth  
fish  
Fisher, Osmond  
fish-hook beach

fish-tail bit  
fissility  
fission  
fission hypothesis  
fission-track dating  
fissure volcano  
fixation  
fixed-source method  
fixigena  
fjord  
fjord  
f-k space  
Fladbury  
flagstone  
flake tectonics  
flakiness index  
flame photometry  
flame spectrometry  
flame structure  
flamma  
flammagenitus  
Flandrian  
flank eruption  
flaser bedding  
flaser gneiss  
flaser rock  
flash flood  
flat  
flat bed  
flat-iron  
flat spot  
flattening  
F-layer  
FLEX

flexural basin  
flexural isostasy  
flexural rigidity  
flexural slip  
flexure  
flexus  
flight  
Flinn diagram  
flint  
floating point  
floatstone  
flocculation  
floccus  
Flock 1  
flood basalt  
flood forecasting  
flood-peak formulae  
floodplain  
flood prediction  
flood tide  
flood zone  
floor thrust  
flora  
Florian  
Florida current  
floristics  
flotation separation  
flow  
flow cleavage  
flower structure  
flow folding  
flowmeter  
flow slide  
flowstone

flow till  
fluctus  
fluid mud  
fluid inclusion  
fluidization  
flume  
flumen  
flumina  
fluorescence  
Fluorescence Explorer  
fluorite  
fluorometer  
fluorspar  
flushed zone  
flute cast  
fluted moraine  
flute mark  
fluvial  
fluvial processes  
fluviate  
fluvic horizon  
fluviokarst  
fluvisols  
fluxgate magnetometer  
fluxoturbidite  
flyer  
Flying Laptop  
flysch  
focal mechanism, earthquake  
focus  
fodinichnia  
foehn wall  
foehn wind  
fog

föhn wall  
föhn wind  
foid  
foiolite  
fold  
fold-and-thrust belt  
fold angle  
fold axis  
fold belt  
folding frequency  
fold limb  
fold test  
foliation  
folivorous  
Folk limestone classification  
fondathem  
fondoform  
fool's gold  
footwall  
foram  
foramen  
foramen magnum  
foramina  
foraminifera  
Foraminiferida  
foramol  
forams  
forced convection  
forced regression  
forced regressive systems tract  
fore-arc  
fore-arc basin  
forebulge  
foredeep

foredune  
foreland  
foreland basin  
foreland thrust belt  
fore reef  
foreset  
foreshock  
foreshore  
foreshortening  
forked lightning  
form  
formal  
formation  
formation age  
formation evaluation  
formation velocity  
form factor  
form-genus  
FormoSat-3  
form roughness  
form set  
forsterite  
FORTE  
Fortin barometer  
Förtsch discontinuity  
forward problem  
fossa  
fossil  
fossil fuel  
fossiliferous micrite  
fossilization  
fossil-Lagerstätte  
fossula  
foundation

founder effect  
founder lineage  
fourchite  
Fourier analysis  
Fourier synthesis  
Fourier transform  
fracking  
fractal  
fractional crystallization  
fractionation  
fracto-  
fractography  
fracture  
fracture cleavage  
fracture porosity  
fracture spacing index  
fracture zone  
fractus  
fragipan  
fragmental  
framestone  
framework grains  
framework porosity  
Francisco  
Franconian  
franklinite  
Frasien  
Frasnian  
frazil ice  
free-air anomaly  
free-air correction  
free atmosphere  
freeboard  
free dune

free face  
free oscillations  
freestone  
freezing nuclei  
freibergite  
frequency  
frequency domain  
fresh water  
freta  
Fretted and Chaotic Hummocky Terrain  
fretum  
friable  
friction  
frictional angle  
frictional resistance, angle of  
frigid  
frigid zone  
fringing reef  
Fronian  
front  
frontal arc  
frontal fog  
frontal wave  
frontal zone  
frontogenesis  
frontolysis  
frost  
frost boil  
frost heave  
frost heave test  
frost heaving  
frost hollow  
frost pull and frost push  
frost-shattering

frost smoke  
frost table  
frost wedging  
Froude number  
frustule  
Fs  
fugacity  
fugichnia  
Fujita Tornado Intensity Scale  
fulgurite  
fuller's earth  
fulvic acid  
fumarole  
fumarolic stage  
functional morphology  
fundamental form  
fundamental strength  
Fungi  
funnel cloud  
funnelling  
furious fifties  
fusain  
fusiform  
fusinite  
fusion  
fusuline  
Fuvelian  
FY  
G  
G  
g  
gabbro  
gaging station  
gahnite

Gaian hypothesis  
gaining stream  
gal  
galactic cosmic rays  
Galápagos Rift  
Galápagos Rise  
Galassia  
Galatea  
galaxite  
gale  
galena  
Galilean satellites  
Galileo  
Gallic  
gamma  
gamma–gamma sonde  
gamma-ray log  
gamma rays  
gamma-ray sonde  
gamma-ray spectrometer  
gamma-ray spectrometry  
gangué  
ganister  
gannister  
ganoid scale  
Ganymede  
Gaofen  
GaoJing  
gap  
Gardar rifting  
garnet  
GARP  
gas  
gas chimney

gas chromatography  
Gasconadian  
gas-liquid chromatography  
Gaspra  
gas-retention age  
gas-solid chromatography  
gastrolith  
Gastropoda  
gauging station  
Gault Clay  
Gauss, Karl Friedrich  
Gauss  
gauss  
Gaussian distribution  
Gaussian process regression  
Gazzi-Dickinson method  
GCM  
GCOM  
GCOM-C1  
GEDI  
Gedinnian  
gedrite  
gehlenite  
Geiger counter  
Geiger-Müller counter  
Geikie, Archibald  
gel  
Gelasian  
gel-filtration  
gelifluction  
gelifraction  
geliturbate  
geliturbation  
gelivation

gemstone  
genal angle  
genal spine  
gendarme  
gene  
gene flow  
gene pool  
genera  
general adaptation  
general circulation  
general circulation model  
generalized reciprocal method  
generating curve  
Genesis  
Genesis Inflatable Space Complex  
genetic drift  
genetic stratigraphic sequence model  
genetic stratigraphic unit  
geniculate twin  
genitus  
genotype  
genus  
genus-zone  
GEO  
geo-  
geo  
geobarometer  
geobotanical anomaly  
geobotanical exploration  
geochemical affinity  
geochemical anomaly  
geochemical cycle  
geochemical differentiation  
geochemical soil survey

geochemistry  
geochronologic unit  
geochronology  
geochronometric scale  
geochronometry  
geode  
geodesy  
geodetic latitude  
geodetic measurement  
geodynamics  
geoecology  
geo-electric section  
GeoEye-1  
geognosy  
geographic information system  
Geographos  
geoid  
GEO-KOMPSAT-1  
geological barrier  
Geological Long Range Inclined Asdic  
Geological Society of America  
Geological Society of London  
geologic cross-section  
geologic map  
geologic map symbols  
geologic timescale  
geologic-time unit  
geology  
geomagnetic dipole field  
geomagnetic equator  
geomagnetic field  
geomagnetic polarity interval  
geomagnetic pole  
geomagnetic reversal timescale

geomatics  
geometric distribution  
geometric factor  
geomicrobiology  
geomorphic sequence  
geomorphology  
geopetal structure  
geophone  
geophysics  
geophysiology  
Georgian epoch  
geosphere  
geostatic stress  
Geostationary Meteorological Satellite  
Geostationary Operational Environmental Satellite  
Geostationary Operational Meteorological Satellite-2  
geostationary orbit  
geostrophic current  
geostrophic wind  
geosynchronous orbit  
geosyncline  
GEOTAIL  
geotechnical map  
geotherm  
geothermal brine  
geothermal field  
geothermal gradient  
geothermic survey  
geothermometer  
germanate system  
geyser  
GF  
GHGSat-D  
GHOST

ghost  
ghost stratigraphy  
Ghyben–Herzberg relationship  
Giacobini–Zinner  
gibber  
gibber plain  
Gibbs free energy  
Gibbs function  
gibbsite  
giga-  
gigantic jet  
Gigantoproductus giganteus  
Gilbert  
Gilbert, Grove Karl  
Gilbert, William  
Gilbert-type delta  
gilgai  
gills  
gilsonite  
Ginkgoales  
Giotto  
gipfflur  
GIS  
Gisbornian  
GISP  
Givetian  
glabella  
glaci-  
glacial breach  
glacial diffluence  
glacial diversion  
glacial drainage channel  
glacial horn  
glacial lake outburst flood

glacial limit  
glacial morphoclimatic zone  
glacial period  
glacial plucking  
glacial stairway  
glacial theory  
glacial transfluence  
glacial trough  
glaciaquatic  
glaciated rock knob  
glaciation  
glacier  
glacier creep  
glacier ice  
glacier power  
glacier surge  
glacier wind  
glacifluvial  
glacilacustrine  
glacio-  
glacioeustasy  
glacioisostasy  
glaciology  
glaciomarine  
glaciomarine sediment  
glaciotectonics  
glaebules  
glance  
glass  
glass-plate reflector  
glass shards  
glass sponges  
glassy  
glaucinite

glaucony  
glaucophane  
glaucophane-schist facies  
G-layer  
glaze  
Gleedonian  
Glen's power flow law  
gley  
gleying  
gley soil  
gleysols  
gleyzation  
gliding tectonics  
gliding twins  
glimmerite  
Glinka, Konstantin Dimitrievich  
Global Atmospheric Research Programme  
Global Change Observation Mission-Climate 1  
Global Change Observation Mission-Water  
Global Ecosystem Dynamics Investigation  
Global Heritage Stone Resource  
Global Horizontal Sounding Technique  
Global Positioning System  
Global Precipitation Measurement  
Global-scale Observations of the Limb and Disk  
Global Seismographic Network  
global tectonics  
global warming potential  
global water budget  
Globigerina ooze  
Gloger's rule  
glomeroporphyritic  
GLORIA  
Glory

Glossifungites  
Glossopteris flora  
gloup  
glow curve  
GMES  
GMS  
gnamma  
Gnathostomata  
gneiss  
gneissose banding  
gneissosity  
Gnetales  
gnome  
gnomon  
goaf  
gobi  
GOCE  
GOES  
goethite  
Göktürk  
GOLD  
gold  
golden algae  
golden-brown algae  
golden spike  
Goldschmidt, Victor Moritz  
Goldschmidt's rules  
GomeSpace Express  
Gomphotheriidae  
GOMS-2  
GOMX  
gonatoparian suture  
Gondwana  
Gondwanaland

goniatites  
goniometer  
goniometry  
Gorda Plate  
Gore  
Gorstian  
GOSAT  
gossan  
Gothenburg  
Gothian  
Gothian orogeny  
gouge  
GovSat-1  
GPM  
GPS  
Grabau, Amadeus William  
graben  
grab sampling  
GRACE  
grade  
graded bedding  
graded reach  
graded sediment  
graded slope  
gradient  
gradient wind  
grading curve  
gradiometer  
gradualism  
grain  
grain boundary sliding  
grain flow  
grain roughness  
grain shape

grain size, igneous rocks  
grainstone  
grain-support  
granite  
granite minimum  
granitic layer  
granitization  
granitoid  
granoblastic  
granodiorite  
granofels  
granophyre  
granophyric  
granular  
granular iron formation  
granulation  
granule  
granulestone  
granulite  
granulite facies  
granulometry  
grapestone  
graphic  
graphic log  
graphite  
graphoglyptid  
graptolite  
Graptolithina  
Graptoloidea  
grass minimum temperature  
graupel  
gravel  
gravimeter  
gravimetric analysis

gravimetry  
gravitational acceleration  
gravitational constant  
gravitational equipotential  
gravitational field  
gravitational potential energy  
gravitational signature  
gravitational water  
gravity anomaly  
gravity assist  
gravity corer  
Gravity field and steady-state Ocean Circulation Explorer  
gravity gliding  
gravity meter  
Gravity Recovery and Climate Experiment  
gravity separation  
gravity settling  
gravity sliding  
gravity survey  
gravity tectonics  
gravity unit  
graywacke  
grazer  
greasy  
great circle  
Great Interglacial  
Great Oxidation Event  
Great Red Spot  
green algae  
greenalite  
greenhouse effect  
greenhouse gas  
Greenhouse Gases Observing Satellite  
Greenhouse Gas Satellite – Demonstrator

greenhouse period  
Greenlandian  
Greenland Ice Core Project  
Greenland Ice Sheet Project  
greensand  
greenschist  
greenschist facies  
greenstone  
greenstone belt  
greisen  
Grenvillian orogeny  
Grenz horizon  
grey-brown podzolic  
grey level  
greywacke  
grèze litée  
grid reference  
Griesbachian  
Griffith cracks  
Griffith failure criterion  
Griffith–Murrell failure criterion  
Grigg–Skjellerup  
grike  
GRIP  
GRM  
groin  
Grooved Terrain  
groove mark  
grossular  
ground anchor  
ground-control point  
ground data  
ground frost  
ground ice

ground information  
groundmass  
ground moraine  
ground range  
ground roll  
ground surge  
ground truth  
groundwater  
groundwater facies  
groundwater flow  
group  
group interval  
group speed (of wave)  
group velocity  
grouting  
growan  
growth band  
growth curve  
growth fault  
growth-fibre analysis  
growth fibres  
growth line  
growth twinning  
groyne  
Grüneisen ratios  
grunerite  
grus  
GSA  
GSL  
GSN  
Guadalupian  
guano  
guard  
Guettard, Jean Étienne

guild  
Gulfian  
Gulf Stream  
gull  
gulling  
gully  
Gunflint Chert  
Günz  
Günz/Mindel Interglacial  
gust  
Gutenberg, Beno  
Gutenberg discontinuity  
gutter cast  
guyot  
G-wave  
GWP  
Gymnodiniales  
gymnosperm  
Gymnospermae  
gypcrete  
gypsic  
gypsisols  
gypsum  
gyre  
gyrocompass  
gyroconic  
gyrogonite  
gyroremanent magnetization  
gyttja  
Gzelian  
Gzhelian  
H  
H 0  
haar

habit  
haboob  
hackly fracture  
HAD  
hadal zone  
hade  
Hadean  
Hadley cell  
Hadrosauridae  
hadrosaurs  
Hadrynian  
haematite  
haemocyanin  
Hagedoorn method  
hail  
hair hygrometer  
hairpin  
Haiyang-1B  
haldane  
Hale–Bopp  
Hale cycle  
half-field prism  
half-graben  
half-life  
half-shadow test  
half-space  
half-spreading rate  
half-width  
halides  
Halimede  
halinity  
Haliomma vetustum  
halite  
Halley

Hall, James  
Hall, James  
Hall effect  
Hallian  
halloysite  
halmyrolysis  
halo  
haloclasty  
halocline  
halokinesis  
halophile  
halotectonism  
hamada  
hammada  
hammer chart  
hammer source  
hand lens  
hanging valley  
hanging wall  
hardground  
hardness  
hardpan  
HARI  
Harker diagram  
harmattan wind  
harmonic  
harmonic fold  
harmonic tremor  
Harnagian  
HARP  
Hastarian  
hastingsite  
Haumurian  
Hauterivian

Haüy, Abbé René-Just  
häüyne  
Hawaiian-Emperor chain  
Hawaiian eruption  
hawaiite  
Hawera  
Hawker  
Hayabusa 2  
haze  
HDR  
head  
head cut  
heading blasting  
head shield  
headwall, glacial  
head wave  
heat capacity  
heat flow  
heat-flow anomaly  
heat-flow unit  
heat flux  
heat of formation  
heave  
heavy liquids  
heavy-medium separation  
hedenbergite  
Heezen, Bruce Charles  
Heim, Albert  
Heinrich events  
Heinrich layers  
Helderbergian  
Helene  
helicitic structure  
helictite

Helikian  
Heliopora  
helium abundance interferometer  
helium 'burning'  
helium clock  
Hellenic Arc  
Hellenic Plate  
Hellenic Trench  
Helminthoida  
helm wind  
Helvetian  
hematite  
hemera  
hemi-  
Hemichordata  
hemimorphism  
hemimorphite  
hemipelagic sediment  
hemipelagite  
Hengelo  
Hennig, Willi  
Hennig's dilemma  
Herangi  
Herbert Smith refractometer  
Hercynian orogeny  
hercynite  
Heretaungan  
hermatypic  
herringbone cross-bedding  
Hervyan  
Hesperian  
Hess, Harry Hammond  
hessenbergite  
Heterian

hetero-  
heterocercal tail  
heterochrony  
heterocoelus  
Heterocorallia  
heterodont  
Heterodonta  
Heterodontosauridae  
heterogeneity  
heterogeneous accretion  
heterogeneous simple shear  
heterolithic bedding  
heterolithic unconformity  
heteropygous  
heterosphere  
heterospory  
Heterostraci  
heterotopy  
Hettangian  
Hexacorallia  
Hexactinellida  
hexad  
hexagonal  
Hexapoda  
HFS  
HFU  
hiatus  
hidden layer  
hierarchical method  
high-alumina basalt  
high-aspect-ratio ignimbrite  
high-field-strength elements  
high-level waste  
high-pass filter

high-plateau crustal type  
high-potassium basalt  
highstand  
highstand systems tract  
high symmetry  
Hiller borer  
Hiller peat-borer  
hill fog  
Himalayan orogenic belt  
Himalia  
Himawari-8 and -9  
hinge  
hinge line  
hinterland  
Hippocamp  
Hirnantian  
Hiscock Radiation Belt Explorer  
histic epipedon  
histogram  
histosols  
HJ-1  
Hjulström effect  
hlaup  
HMS  
hoar frost  
Hoboken  
hogback  
hog's back  
Hohmann transfer orbit  
Holkerian  
Holmes, Arthur  
holo-  
Holocene  
holochroal

holocrystalline  
holohyaline  
holokarst  
hololeucocratic  
holometabolous  
holomictic  
holophyletic  
Holostei  
holostomatous  
holostratotype  
holosymmetric  
holothuroid  
Holothuroidea  
holotype  
Holsteinian  
Homalozoa  
homeomorph  
homeostasis  
homeotherm  
Homerian  
Hominidae  
Hominoidea  
homo-  
homocercal tail  
homodont  
homoeostasis  
homogeneous accretion  
homogeneous non-rotational strain  
homogeneous nucleation  
homogeneous rotational strain  
homogeneous strain  
homogenitus  
homogenization temperature  
homoiotherm

homologous  
homomutatus  
homoplasy  
homopycnal flow  
homosphere  
homospory  
homotaxis  
Honda–Mrkos–Pajdusakova  
hoodoo  
Hooke, Robert  
Hooke’s law  
hopeful monster  
‘hopper’ crystals  
horizon  
horizontal drilling  
horizontal hole  
horizontal stack  
horn  
hornblende  
hornblende–hornfels facies  
hornblendite  
hornfels  
hornitos  
horotely  
horse  
horse latitudes  
horsetails  
horst  
hortic horizon  
Horton analysis  
Hortonian flow  
hot dry rock  
hot spot  
hot spring

hot working  
Howard, Luke  
Hoxnian  
HRBE CubeSat Mission  
HREE  
HSR  
Huan Jing-1  
Hubble constant  
Hubble parameter ( $H_0$ )  
Hubble's law  
Hubble Space Telescope  
Hudsonian orogeny  
hue  
Hugoniot  
Humberian orogeny  
Humboldt, Friedrich Heinrich Alexander von  
Humboldt current  
humerus  
humic acid  
humic coal  
humidity  
humid mid-latitude morphoclimatic zone  
humid tropical morphoclimatic zone  
humification  
humilis  
humite  
hummocky cross-bedding  
hummocky moraine  
humus  
Huronian  
hurricane  
Hutchinsonian  
Hutton, James  
Huygenian eyepiece

Huygens  
Huygens' principle  
HY-1B  
Hyaenidae  
Hyakutake  
hyaline  
hyaloclastite  
hyalopilitic  
Hyalospongea  
hydration  
hydraulic boundary  
hydraulic conductivity  
hydraulic corer  
hydraulic equivalent  
hydraulic fracture  
hydraulic fracturing  
hydraulic geometry  
hydraulic gradient  
hydraulic head  
hydraulic radius  
hydric  
hydrocarbon  
hydrochemistry  
hydroclast  
hydrocollapsibility  
hydrofracturing  
hydrogen 'burning'  
hydrogeologic map  
hydrogeology  
hydrograph  
hydroids  
hydro-isostasy  
hydrologic cycle  
hydrologic modelling

hydrologic network  
hydrologic regions  
hydrologic simulation  
hydrology  
Hydrology and Water Resources Programme  
hydrolysate  
hydrolysis  
hydromuscovite  
hydrophone  
hydrosphere  
hydrospire  
hydrostatic stress  
hydrostratigraphic units  
hydrothermal activity  
hydrothermal metamorphism  
hydrothermal mineral  
hydrothermal vent  
hydrovolcanic processes  
hydroxides  
hydroxyapatite  
Hydrozoa  
hyetograph  
hygromagnetophile elements  
hygrometer  
hygroscopic nucleus  
hygroscopic water  
hygrothermograph  
Hylonomus lyelli  
hyoid  
Hyalithida  
hyp-  
hypabyssal  
hyper-  
Hyper-Angular Rainbow Polarimeter

Hyperion  
hypermorphosis  
hyperpycnal flow  
hypersolvus granite  
hypersolvus syenite  
Hyperspectral Precursor and Application Mission  
hypersthene  
hyperthermic  
hyperthermophile  
hypichnia  
hypidiomorphic fabric  
hypidiotopic fabric  
hypo-  
hypocentre  
hypocrystalline  
hypogene  
hypolimnion  
hyponome  
hyponomic sinus  
hypopycnal flow  
hypostomal suture  
hypostratotype  
hypothea  
hypothermal  
hypothesis  
hypothesis testing  
Hypsilophodontidae  
Hypsithermal  
hypsographic curve  
hypsometric curve  
Hyracotherium  
hysteresis loop  
hystricospheres  
IAEG

-ian  
Iapetus  
Iapetus Ocean  
Iberomesornis  
IBEX  
Ibexian  
Icarus  
ICE  
ice  
ice ages  
iceberg calving  
ice blink  
ice cap  
Ice, Cloud, and land Elevation Satellite  
ice core  
ice crystal  
IceCube  
ice dome  
ice field  
icehouse period  
Iceland low  
ice mound  
ice nucleus  
ice-rafted detritus  
ICESat  
ice sheet  
ice shelf  
ice stream  
ice wedge  
ice-wedge polygon  
ichnoclast  
ichnocoenosis  
ichnofabric  
ichnofacies

ichnofossil  
ichnogenus  
ichnoguild  
ichnology  
ichnospecies  
ichnotaxobase  
ichnotaxonomy  
Ichthyornis  
Ichthyosauria  
Ichthyostega  
Ichthyostegopsis  
ICP  
-id  
Ida  
Idamean  
-ide  
-ides  
idioblastic  
idiomorphic  
idiomorphic fabric  
idiotopic fabric  
idocrase  
Idwian  
I f  
IfSAR  
IGC  
igen  
igneous  
igneous rock classification  
ignimbrite  
IGRF  
IGTS  
Iguanodontidae  
ijolite

ilium  
Illinoian  
illite  
illuminator  
illuviation  
ilmenite  
IMA  
IMAGE  
image intensifying  
imaginary component  
Imager for Magnetopause-to-Aurora Exploration  
imaging  
imaging radar  
imaging spectrometer  
imbibition  
Imbrian  
imbricate  
imbricate structure  
immersion objective method  
immiscible  
immobilization  
IMP-8  
impact cratering  
impactite  
impactogen  
impeded dune  
imperforate  
impervious rock  
implosion  
impulse response function  
impunctate  
Inarticulata  
Inarticulate brachiopods  
inceptisols

incident angle  
incipient hardground  
incised meander  
inclination  
inclined extinction  
inclined fold  
inclinometer, geomagnetic  
included fragments  
incompatible elements  
incompetent  
incompressibility modulus  
incongruent dissolution  
incongruent melting  
inconsequent drainage  
incumbent replacement  
incus  
INDEX  
index ellipsoid  
index fossil  
index mineral  
index species  
Indian jade  
Indian National Satellite-3  
Indian Ocean  
Indian Remote Sensing  
Indian summer  
indicated reserve  
indicator  
indicatrix  
Indo-Australian Plate  
Induan  
induced polarization  
induced potential  
induced pulsed transient

induction  
induction log  
induction sonde  
inductively coupled plasma emission spectrometry  
induration  
industrial mineral  
inertial reference frame  
inertinite  
infaunal  
inferred reserve  
inferred tree  
infiltration  
infiltration capacity  
infinitesimal strain  
inflexion point  
inflow band  
influent stream  
influx  
informal  
infrared radiation  
infrared remote sensing  
ingrown meander  
inhomogeneity  
inhomogeneous strain  
initial levée  
initial strontium ratio  
inland sea  
inlier  
inner planet  
Innovative Technology Demonstration Experiment  
inosilicate  
in-phase component  
INPUT  
InSAR

INSAT-3  
Insecta  
inselberg  
insequent drainage  
InSight  
insolation  
insolation weathering  
instability  
instantaneous field of view  
instantaneous rotation  
instantaneous strength  
Institute of Space and Astronautical Science  
insula  
intact rock strength  
Integrated Ocean Drilling Programme  
intensification  
intensity  
intensity  
intensity-hue-saturation processing  
inter-  
interambulacral  
interambulacrum  
inter-arc basin  
inter-arc trough  
interarea  
interbiohorizon zone  
interception  
intercept ratio  
intercept time  
intercrystalline boundary  
intercrystalline porosity  
interdigitating  
interface  
interface-controlled growth

interfacial angle  
interfacial polarization  
interference  
interference colour chart  
interference colours  
interference figure  
interference pattern  
interferogram  
interferometer  
interferometric synthetic aperture radar  
interfingering  
interflow  
interfluve  
interglacial  
intergrade  
intergranular  
intergranular displacement  
intergranular pores  
Interior Exploration using Seismic Investigations, Geodesy and Heat Transport  
inter-limb angle  
interlocking  
intermediate rock  
intermittent stream  
intermontane  
internal angle of friction  
internal mould  
internal node  
internal reflection  
internal standard  
internal wave  
International Association for Engineering Geology and the Environment  
International Cometary Explorer  
International Federation of Digital Seismic Networks

International Geomagnetic Reference Field  
International Gravity Formula  
International Gravity Standardization Network  
International Mineralogical Association  
International Monitoring Platform  
International Polar Year  
International Programme of Ocean Drilling  
International Research Institutions for Seismology  
International Sun–Earth Explorer  
International Union of Geological Sciences  
interparticle porosity  
interpenetrant twin  
inter-record gap  
intersection cleavage  
intersection lineation  
interseismic period  
intersertal  
interstade  
interstadial  
Interstellar Boundary Explorer  
interstellar clouds  
interstellar medium  
interstitial  
intertidal zone  
intertropical confluence  
intertropical convergence zone  
intertropical front  
interval  
intervallum  
interval time  
interval velocity  
interval zone  
intortus  
intra-

intraclast  
intrafolial fold  
intraformational  
intramicrite  
intra-orogen basin  
intraparticle porosity  
intrasparite  
intrinsic permeability  
intrusion  
intrusive  
intrusive phonolite  
invaded zone  
Inverian  
inverse problem  
inversion  
inversion axis  
inversion tectonics  
invertebrate  
inverted metamorphism  
inverted relief  
inviscid  
involute  
involution  
Io  
ion  
ion drive  
ion engine  
ion exchange  
ionic bond  
ionic charge  
ionic potential  
ionic radius  
ionic substitution  
ionization potential

ionosphere  
ion pair  
ion thrust  
IPOD  
Ipswichian  
IPY  
IR  
IRAZÚ  
iridescence  
iridescent clouds  
iridium anomaly  
Iridium NEXT  
IRIS  
Irish elk  
IRM  
iron, native  
iron formation  
iron glance  
iron hypothesis  
iron meteorite  
iron pan  
ironstone  
irradiance  
irragic horizon  
irregular echinoids  
Irregulares  
irrigation  
irrotational wave  
Is  
IRS  
ISAS  
ischium  
ISEE  
island arc

island-arc crustal type  
iso-  
isobar  
isobaric gas-tight sampler  
isobaric surface  
isobase  
isobath  
isochore  
isochron  
isochron map  
isochronous  
isoclinal fold  
isoclinic chart  
isoconductivity map  
isodynamic line  
isofrigid  
isogal  
isogon  
isogonic  
isograd  
isogyre  
isohel  
isohyet  
isohyperthermic  
isomagnetic chart  
isomesic  
isometric  
isomorphous  
isomyarian  
isoneph  
isopach  
isopach map  
isopycnal  
isopygous

isoresistivity map  
isoseismal map  
isospore  
isostasy  
isostatic anomaly  
isostatic compensation  
isostructural  
isotherm  
isothermal remanence  
isothermic  
isotope  
isotope dilution  
isotope fractionation  
isotope geochemistry  
isotope hydrology  
isotope tracer  
isotopic dating  
isotrophic  
isotropic  
isotropic indicatrix  
isotropy  
isotypic  
isovelocity plot  
isovol  
isp  
ISSC  
Isuan  
itabirite  
ITCZ  
iterative evolution  
Itokawa  
IUGS  
IVD  
Ivorian

I-wave  
IX  
Jaccard's index  
jacobsite  
Jacob's staff  
jade  
jadeite  
Jameson, Robert  
jamesonite  
Janjukian  
Janus  
Japan Aerospace Exploration Agency  
Japanese Meteorological Agency seismic intensity scale  
Japan Trench  
Japan-type margin  
Jaramillo  
jarosite  
Jason  
Jason-1  
Jason-2  
jasper  
jaspillite  
Java Trench  
jawless fish  
JAXA  
Jeffreys, Sir Harold  
Jeffreys–Bullen curves  
jet  
jet stream  
JMA scale  
Johannian  
JOIDES  
joint  
Joint Oceanographic Institutions for Deep Earth Sampling

Joint Polar Satellite System-1

joint set

joint system

jöklar

jökulhlaup

jökull

Jolly balance

Joly, John

Jotnian

Jotnian orogeny

jovian

jovian planet

jovian satellites

JPSS-1

Juan de Fuca Plate

Juan de Fuca Ridge

jug

Juliet

Juno

Jupiter

Jurassic

Jura-type relief

Juvavic

juvenile

juvenile water

J-wave

Kaena

kaersutite

KAGUYA

Kaiatan

Kaihikuan

kainite

Kainozoic

Kalb light line

Kalimnan  
kaliophilite  
Kalpana-1  
kalsilite  
Kama  
kame  
kame delta  
kamenitza  
kame terrace  
kamikaze grains  
kandite  
Kanopus-V1  
Kanopus-V-1K 1  
Kanopus-Vulkan-Infra-Krasny-1  
Kanopus-Vulkan N1  
Kansan I and II  
kaolin  
kaolinite  
kaolinitization  
kaolinization  
Kapitean  
Karatau  
Karatavian  
Karelian  
Karelian orogeny  
Karginisky  
von Karman–Prandtl equation  
K–Ar method  
Karnian  
karren  
karst  
karstic aquifer  
Kashirskian  
Kasimovian

kastanozems  
katabatic wind  
katafront  
katatectic layer  
katazone  
katophorite  
Kawhia  
Kazanian  
KazEOSat-1  
kb  
K-band  
keel  
Keewatinian  
KEI  
Keiloran  
kelly  
kelp  
Kelvin, Lord  
kelvin  
Kelvin scale  
Kelvin–Voigt model  
kelyphitic rim  
Kenoran orogeny  
Kenorland  
Kent Ridge 1  
kenyte  
Kepler's laws of planetary motion  
keratophyre  
kernal function  
kerogen  
kersantite  
Ketilidian orogeny  
kettle hole  
kettle lake

Kew barometer  
Keweenawan  
key bed  
key evolutionary innovation  
khamsin  
Kibalian orogeny  
kidney ore  
kieselguhr  
kill curve  
kilobar  
Kimberella quadrata  
kimberlite  
Kimmeridgian  
Kinderhookian  
Kinderscoutian  
K-index  
kinematics  
kinematic viscosity  
kinematic wave  
king crab  
kink band  
kipuka  
Kirkfield  
Kirkwood Gaps  
Klazminskian  
klippe  
knee twin  
knick point  
knob and kettle  
Knobby Terrain  
knock and lochan  
knot  
Knott, Cargill Gilston  
Knott's equations

Kohoutek  
Kolmogorov Wiener prediction  
komatiite  
KOMPSAT  
kona storm  
Königsberger ratio  
kopje  
Köppen, Wladimir Peter  
Köppen climate classification  
koppie  
Korangan  
Korea Multi-Purpose Satellite  
kosava  
Kotlassia prima  
KR1  
KREEP volcanism  
Krevyakinskian  
kriging  
krotovina  
K-selection  
K/T boundary event  
Kuehneosaurus latus  
Kuenen, Philip Henry  
Kuiper, Gerard Peter  
Kuiper belt  
Kuiperian  
Kula Plate  
Kungurian  
kunkar  
Kuril Trench  
Kuroshio current  
kurtosis  
Kutorginida  
K-wave

kyanite  
Kyushu Satellite for Earth Observation System Demonstration  
labes  
Labrador current  
labradorite  
Labyrinthodontia  
labyrinthus  
laccolith  
lacuna  
lacunosus  
lacus  
lacustrine  
LAD  
Ladinian  
lag  
lag breccia  
lag deposit  
LAGEOS  
Lagerstätte  
lagoon  
Lagrangian current measurement  
Lagrangian points  
lahar  
Lake Mungo  
Lamarckism  
Lamarck, Jean Baptiste Pierre Antoine de Monet, Chevalier de  
Lamb, Hubert Horace  
Lambeophyllum profundum  
Lambertian reflector  
Lamb's dust-veil index  
Lamé constant  
lamellar  
Lamellibranchia  
lamina

laminar flow  
lamine  
lamination  
laminite  
laminoid fenestrae  
lamprophyre  
lamp shells  
Lancefieldian  
land and sea breezes  
land bridge  
landfill  
Landsat  
landscape ecology  
landslide  
Langhian  
La Niña  
lanthanide  
Laomedea  
LAPAN-A2  
LAPAN-A3  
lapiés  
lapilli  
lapilli-tuff  
lapis lazuli  
Laplace, Pierre Simon, Marquis de  
Laplace's equation  
lapout  
lappets  
lapse rate  
lapse time  
Lapworth, Charles  
Laramide-Columbian orogeny  
Laramide Province  
LARES

large-aperture seismic array  
large-ion lithophile  
large low shear velocity province  
LARI  
Larissa  
larvikite  
LASA  
Laschamp  
laser  
laser altimeter  
Laser Geodynamics Satellite  
laser ranging  
LAser RElativity Satellite  
last-appearance datum  
Late-Devensian Interstadial  
late-glacial  
late heavy bombardment  
latent heat of transition  
lateral  
lateral accretion deposit  
lateral dispersion  
lateral moraine  
laterite  
laterolog sonde  
latite  
latite-andesite  
latitude correction  
latosol  
lattice  
lattice energy  
lattice gliding  
laumontite  
Laurasia  
Laurentia

Laurentian  
Laurentian Shield  
Laurentide ice sheet  
lava  
lava blister  
lava channel  
lava haze  
lava lake  
lava levée  
lava tube  
law of constancy of interfacial angles  
law of constant proportions  
law of correlation of facies  
law of cross-cutting relationships  
law of faunal succession  
law of Haüy  
law of original horizontality  
law of rational indices  
law of rational ratios of intercepts  
law of superposition of strata  
law of the wall  
laze  
Laxfordian  
Laxfordian orogeny  
layer cloud  
Layered Deposits  
layered silicate  
layer-parallel shortening  
layover  
Lazarus taxon  
lazurite  
L-band  
leachate  
leaching

lead–lead dating  
lead loss  
Leaf Margin Analysis  
leaky transform fault  
least-work principle  
least-work profile  
lebensspuren  
lechatelierite  
lectostratotype  
lectotype  
Leda  
lee depression  
lee dune  
lee waves  
left lateral fault  
Lehmann, Inge  
Lehmann, Johann Gottlob  
Leighton–Pendexter classification  
Leitz–Jelley refractometer  
Lemur-2  
Lenian  
lenticula  
lenticulae  
lenticular bedding  
lenticularis  
Lenz's law  
LEO  
Leonardian  
Le Pichon, Xavier  
lepidoblastic  
lepidocrocite  
Lepidodendron selaginoides  
lepidolite  
lepidomelane

Lepidophloios kilpatrickense  
Lepidosauria  
lepisphere  
Lepospondyli  
leptograptid  
leptokurtic  
leptosols  
lessivage  
leste  
leucite  
leucite–basanite  
leucitite  
leuco-  
leucocratic  
leucosome  
levanter  
leveche  
levée  
level of compensation  
Lewisian  
LHB  
lherzolite  
Lias  
Libby, Willard Frank  
libeccio  
libration  
librigena  
lichenometry  
Lichida  
life assemblage  
lifting condensation level  
ligand  
light absorption  
Lightning Imaging Sensor

lignite  
Likhvin  
LIL  
Lillburnian  
lime  
lime boundstone  
lime grainstone  
lime mud  
lime mudstone  
lime packstone  
limestone  
limestone pavement  
lime wackestone  
limit-equilibrium analysis  
limnic  
limnic eruption  
limnology  
limonite  
limpet  
Limulus  
Lindgren, Waldemar  
linea  
lineae  
lineage zone  
lineament  
linear regression  
linear sand ridge  
lineation  
line scanner  
line spectrum  
line squall  
Lingulata  
Lingulella viridis  
Lingulida

linguoid  
liptinite  
liquation  
liquefaction  
liquid limit  
liquidus  
LIS  
listric fault  
lithic arenite  
lithic fragment  
lithic greywacke  
lithic wacke  
lithification  
litho-  
lithoclast  
lithofacies  
lithofacies map  
lithologic symbol  
lithologic trap  
lithology  
lithology classification  
lithomarge  
lithomorphic soil  
lithophile  
lithophysae  
lithosequence  
lithosome  
lithosphere  
lithospheric plate  
lithostatic stress  
lithostratigraphic unit  
lithostratigraphy  
Lithothamnion  
litter

Little Ice Age  
littoral drift  
littoral zone  
living fossil  
lixisols  
lizardite  
Llandeilo  
Llandovery  
Llanvirn  
L-layer  
LLSVP  
LMA  
load  
load cast  
loadstone  
loam  
lobe  
lobe fins  
lobsters  
local range zone  
local wind  
lochan  
Lochkovian  
Loch Lomond stadial  
Lockportian  
lode  
lodestone  
lodgement till  
loess  
logging  
log-normal distribution  
logs  
Longfordian  
longiconic

Longisquama insignis  
longitudinal conductance  
longitudinal dispersion  
longitudinal-type coast  
Longmyndian  
long-range forecasting  
longshore bar  
longshore current  
longshore drift  
Longtanian  
Longvillian  
Longwangmioan  
long wave  
Lonsdale, William  
look angle  
looping  
lophodont  
lophophore  
lopolith  
Lopstedt  
Los Angeles abrasion test  
losing stream  
Lost City Hydrothermal Field  
Love, Augustus Edward Hugh  
Love wave  
low  
low-angle fault  
low-aspect-ratio ignimbrite  
low Earth orbit  
low-level waste  
low-potassium tholeiite  
lowstand  
lowstand systems tract  
low symmetry

low-velocity layer  
low-velocity zone  
Lq  
LQ-wave  
LREE  
LST  
L-S-tectonite  
L-tectonite  
lucinoid  
Ludfordian  
Ludhamian  
Ludlow  
Luisian  
lumbar vertebra  
luminance  
luminous night clouds  
lumps, carbonate  
Luna  
lunar  
Lunar-A  
Lunar Highlands  
lunar magnetic variation  
Lunar Orbiter  
lunar timescale  
lunate  
lunette  
lunule  
lustre  
lustre mottling  
Lutetian  
lutite  
luvisols  
luxullianite  
LVL

LVZ  
L-wave  
lycopods  
Lycopsida  
Lyell, Charles  
lysimeter  
Lysithea  
lysocline  
lystric fault  
m-  
M-  
M  
Ma  
maar  
Maastrichtian  
Mab  
McAuliffe  
mackerel sky  
Mackereth corer  
macroclimate  
macroevolution  
macropygous  
macrotidal  
macula  
maculae  
maculose  
Madagascar  
made ground  
Madelung constant  
Madreporaria  
madreporite  
Maentwrogian  
maerl  
Maestrichtian

maestro  
mafic  
Magellan  
Magic Mountain  
magma  
magma chamber  
magmaphile elements  
magmatic arc  
magmatic differentiation  
magmatic fractionation  
magmatic-segregation deposit  
magnesiocromite  
magnesioferrite  
magnesiowüstite  
magnesite  
magnesium number  
magnesium-sulphate soundness test  
magnetic age  
magnetic anomaly  
magnetic anomaly pattern  
magnetic dating  
magnetic domain  
magnetic equator  
magnetic fabric determination  
magnetic field  
magnetic flux  
magnetic gradiometry  
magnetic induction  
magnetic moment  
magnetic orientation  
magnetic permeability  
magnetic profile  
magnetic quiet zone  
magnetic sampling

magnetic separator  
magnetic signature  
magnetic storm  
magnetic survey  
magnetic susceptibility  
magnetic variations  
magnetite  
magnetochemistry  
magnetogram  
magnetograph  
magneto hydrodynamics  
magnetometer  
magnetopause  
magnetosheath  
magnetosphere  
magnetostratigraphic timescale  
magnetostratigraphy  
magnetotelluric sounding  
magnetozone  
magnitude, earthquake  
Magnolia  
magnon  
main-stage sequence  
majanna  
malachite  
Malacostraca  
Malagasy armadillo  
Malakonian  
malleolus  
Mallet, Robert  
malleus  
Malm  
Malvernian  
mamma

Mammalia  
mammal-like reptiles  
mammatous  
mamillary  
mamillated  
mamillated topography  
Mammoth  
mammoth  
Mammuthus  
Mammutidae  
mandible  
manganese nodule  
manganite  
Mangaorapan  
Mangaotanean  
Mangapanian  
mangrove swamp  
man-made soil  
Manning equation  
manometer  
Mansurian  
Mantell, Gideon Algernon  
mantle  
mantle array  
mantle bedding  
mantle cavity  
mantle convection  
mantle creep  
mantled gneiss dome  
mantle lobe  
mantle plume  
manto  
Maozhangian  
map projection

marble  
marcasite  
mare  
mare ridge  
mares' tails  
Margaret  
margarite  
marginal basin  
marginal sea  
marginal suture  
marialite  
Marianas Trench  
marine platform  
Mariner  
maritime air  
maritime climate  
marker bed  
Markov chain  
marl  
marlstone  
Mars  
Mars 96  
Marsdenian  
Mars Global Surveyor  
Marshbrookian  
Mars Observer  
Mars Pathfinder  
marsquake  
Mars Surveyor  
Marsupialia  
marsupials  
martian  
martian canals  
martian terrain units

mascon  
maser  
mass balance  
mass extinction  
mass flow  
massif  
massive  
massive sulphide deposit  
mass movement  
mass number  
mass spectrometry  
mass-wasting  
master curve  
mastodon  
Mata  
Mathilde  
matrix  
matrix-support  
maturation  
Matuyama, Motonori  
Matuyama  
Maury, Matthew Fontaine  
maximum-likelihood classification  
maximum-likelihood tree  
maximum-parsimony tree  
maximum thermometer  
Maxwell mantle  
Maxwell model  
Mayan  
Mayvillian  
mean  
meander  
meander belt  
meander core

meander migration  
meander neck  
meander scroll  
meander wavelength  
mean sea level  
mean square  
measure  
mechanical layering  
mechanical weathering  
medial moraine  
median  
median filter  
median network  
median suture  
median valley  
Medinian  
mediocris  
Mediterranean climate  
Mediterranean-type margin  
Mediterranean water  
medium Earth orbit  
Medusina mawsoni  
Medusinites  
mega-  
megabreccia  
megacryst  
Meganeura  
megaphyll  
megaregolith  
megaripple  
megasclere  
megasequence  
megasplay fault  
megasporangia

megaspore  
megathermal  
megathermal climate  
Meghalayan  
Megha-Tropiques  
meionite  
Meishuchuan  
mela-  
mélange  
melanocratic  
melanosome  
Melbournian  
Melekesskian  
melilite  
melilitite  
melt  
melteigite  
meltemi  
meltout till  
meltwater channel  
MEM  
member  
membrane stress  
Menapian  
mendip  
Menevian  
mensa  
MEO  
Meramecian  
Mercalli, Giuseppe  
Mercalli scale  
Mercurial geologic timescale  
Mercury  
Mercury Orbiter

meridional circulation  
meridional overturning circulation  
Merioneth  
Merionsian  
MERLIN  
merocrystalline  
meromictic  
Merostomata  
mesa  
mesentery  
mesic  
Mesoarchaeon  
mesoclimate  
mesocratic  
mesocyclone  
mesohaline water  
mesokurtic  
mesopause  
Mesoproterozoic  
Mesosauria  
mesosphere  
mesothem  
mesothermal  
mesothermal climate  
mesotidal  
mesotype  
Mesozoic  
Messenger probe  
Messinian  
meta-  
metabasalt  
metacarpal  
metacryst  
metagenesis

metal factor  
metalimnion  
metallic  
metallic bond  
metallogenesis  
metallogenic province  
metaluminous  
metameric segmentation  
metamorphic aureole  
metamorphic differentiation  
metamorphic facies  
metamorphic grade  
metamorphic rock  
metamorphic rock classification  
metamorphic zone  
metamorphism  
metapelite  
Metaphyta  
metaquartzite  
metasediment  
metaseptum  
metasilicate  
metasomatism  
metaspecies  
metastable  
metatarsal  
Metazoa  
metazoan  
meteor  
Meteor  
Meteor-3M  
meteoric diagenesis  
meteoritic abundance of elements  
meteoric water

meteorites  
meteoroid  
Meteorological LEO Observations in the Intertropical Zone  
Meteorological Operational Satellite Program of Europe  
Meteorological Satellite-1  
Meteosat  
methane hydrates  
Methane Remote Sensing Lidar Mission  
methanogen  
Metis  
metocean data  
MetOp  
MetOp-SG  
MetSat-1  
MF  
M-fold  
Mg#  
Miacidae  
miarolitic  
micas  
Michel–Lévy chart  
Michell, John  
Micraster  
micrinite  
micrite  
-micrite  
micrite envelope  
micritic limestone  
micritization  
micro-  
Microaccelerometric Measurements of Satellite Accelerations  
microbialite  
MicroCarb  
microclimate

microcline  
microcontinent  
microcraters  
microcrystalline  
microdiorite  
microearthquake  
microerosion meter  
microevolution  
microfabric  
microfossil  
microfracture  
microgabbro  
microgranite  
microgranodiorite  
micrographic  
microkrystite  
microlite  
microlog  
MicroMAS-2  
micropalaeontology  
microperthite  
microphenocryst  
microphyll  
micropiracy  
microplate  
microporphyrific  
micropulsations  
micropygous  
microsatellite  
Micro-Satellite à trainée pour l'Observation du Principe d'Equivalence  
microsclore  
MicroSCOPE  
microseism  
Micro-sized Microwave Atmospheric Satellite-2

microspar  
microsparite  
microsporangium  
microspore  
microstylolite  
microsyenite  
microtektite  
microthermal  
microthermal climate  
microthermometry  
microtidal  
microwave  
microwave demagnetization  
microwave sounding unit  
Mid-Atlantic Ridge  
mid-continent crustal type  
mid-infrared  
mid-oceanic crustal type  
mid-ocean ridge  
mid-ocean-ridge basalt  
mid-Pleistocene transition  
midstand systems tract  
Mie scattering  
migmatite  
migmatization  
migration  
migration route  
Mikulino  
Milankovitch, Milutin  
Milankovitch cycles  
Milazzian  
milioline winding  
Milleporina  
Miller, William Hallowes

Miller–Bravais indices  
Miller indexes  
Miller indices  
millerite  
milli-  
millibar  
milligal  
millipedes  
Millstone Grit series  
Milne, John  
Mimas  
mimetic twins  
mimetite  
Mimistrobell  
MIMOSA  
Mindel  
Mindel/Riss Interglacial  
Mindyallan  
mineral  
mineralization  
mineral layering  
mineralogy  
mineraloid  
mineral saturation index  
mineral soil  
minette  
minette ironstone  
minimum-distance-to-means classification  
minimum melting curve  
minimum temperature  
minimum thermometer  
minisatellite  
minor fold  
mio-

Miocene  
miogeocline  
miogeosyncline  
mirage  
Miranda  
mirror plane  
miscibility  
miscible  
mise-à-la-masse method  
misfit stream  
mispickel  
Mississippian  
Missourian  
mist  
mistral  
Mitchellian  
mitchellite  
mites  
mitochondrial-DNA  
mitochondrial Eve  
mixed cloud  
mixed pixel  
mixing condensation level  
mixing depth  
mixing ratio  
mixolimnion  
MMS  
moberg  
mobile belt  
mocha stone  
modal analysis  
mode  
model  
Modified units

Moershoofd  
mofette  
moffette  
Mogi doughnut hypothesis  
mogote  
Mohawkian  
Mohnian  
Moho  
Mohorovic'ić, Andrija  
Mohorovic'ić discontinuity  
Mohr stress diagram  
Mohs, Friedrich  
Mohs's scale of hardness  
Moinian  
moisture balance  
moisture budget  
moisture index  
Mokoiwian  
Mokolian  
mol  
molarity  
molasse  
Molasse Basin  
moldic porosity  
mole  
molecular sieve action  
mole drain  
mollic horizon  
mollisols  
Mollusca  
molluscs  
molybdenite  
moment magnitude scale  
moment of inertia

monadnock  
monazite  
monchiquite  
Monera  
Monian  
monimolimnion  
Monitor-E  
mono-  
monochromator  
monocline  
monoclinic  
Monocraterion  
Monocyathea  
monodactyl  
monograptid  
Monograptus  
monolete  
monomictic  
monomineralic  
monomyarian  
monophyletic  
Monoplacophora  
monopodial  
Monorhina  
monostatic radar  
monosulcate  
Monotremata  
Monotrematum sudamericanum  
monotremes  
monotypic  
mons  
monsoon  
monster curve  
montes

month degrees  
Montian  
montmorillonite  
monzodiorite  
monzogabbro  
monzonite  
moom  
Moon  
moonquake  
moonstone  
mor  
moraine  
Morarian orogeny  
MORB  
Morisawa flood-peak formula  
morphoclimatic zone  
morphogen  
morphogenesis  
morphogenetic zone  
morphological mapping  
morphological system  
morphology  
morphometric analysis  
morphometrics  
morphometry  
morphospace  
morphospecies  
morphotectonics  
morphotype  
Morrowan  
mortar texture  
Mortensnes  
mosaic evolution  
mosaic heterochrony

mosaic texture  
Mosasauridae  
mosasaurs  
Moscovian  
Moskva Drift  
moss agate  
Mössbauer spectroscopy  
mossy  
mother cloud  
mother-of-coal  
mother-of-pearl cloud  
mottling  
motu  
Motuan  
mould  
moulin  
mouldic porosity  
Mount Agung  
mountain breeze  
mountain wind  
mouthbar  
moveout  
MPT  
M sat  
M-shape  
MSU  
MTSAT  
mud, drilling  
mud cake  
mudcracks  
mud drape  
mud filtrate  
mudflat  
mudflow

mud log  
mudlump  
mud mound  
mudrock  
mud roll  
mud/shale diapir  
mudstone  
mud-support  
mud volcano  
mugearite  
mull  
mullion structure  
multichannel seismic reflection  
Multifunction Transport Satellite  
multifurcation  
multilocular  
multiple  
multiple common-depth-point coverage  
multiple twinning  
multiplexing  
multi-ring basin  
multispectral imaging  
multispectral scanner  
multispectral scanning systems  
multistate character  
multistorey sandbody  
Multituberculata  
Multi-User System for Earth Sensing  
multivariate analysis  
Munsell colour  
Murchison, Roderick Impey  
Murray, John  
murus  
Muschelkalk

muscle scar  
muscovite  
MUSES  
Muses-C  
mushroom rock  
Mustelidae  
mutation  
mutation rate  
mutatus  
muxing  
Myachkovskian  
mylonite  
myophore  
Myriapoda  
Mytiloidea  
N  
n  
NA  
Nabarro–Herring creep  
nabkha  
nacreous cloud  
nacrite  
nadir  
NADW  
Nafe–Drake relationship  
Nagssugtoqidian orogeny  
Naiad  
Namibian  
Nammalian  
Namurian  
nanofossil  
nano-  
nanofossil  
nanoplankton

NanoRacks-QB50  
NanoSatC-Br1  
nanosatellite  
nappe  
Naraoiidae  
Narizian  
NASA  
naticiform  
native element  
natric horizon  
natrolite  
natron lake  
'natural break'  
natural cast  
natural gas  
natural gas hydrate  
natural remanent magnetism  
natural selection  
natural theology  
nautical mile  
nautilicone  
nautiliconic  
Nautiloidea  
Nazca Plate  
neap tide  
NEAR  
Nearctic faunal realm  
Near Earth Asteroid Rendezvous  
near-field barrier  
near-infrared  
near-infrared mapping spectrometer  
near-infrared mapping spectrophotometer  
near-shore current system  
nebkha

Nebraskan  
nebulosus  
necrology  
Nectarian  
Nectarian system  
needle ice  
Néel, Louis Eugène Félix  
Néel temperature  
negative inversion  
nekton  
nema  
Nemakit-Daldynian  
Nematoda  
Nemertini  
Neoarchaeon  
Neobothriocidaris  
Neocomian  
Neogene  
Neohelikian  
neoichnology  
neomineralization  
neomorphism  
Neoproterozoic  
neostratotype  
neotectonics  
neoteny  
Neotethys  
Neotropical faunal realm  
neotype  
NEP  
nephanalysis  
nepheline  
nepheline-basanite  
nepheline-monzonite

nepheline-syenite  
nephelinite  
nepheloid layer  
nephelometer  
nephrite  
Neptune  
Neptune XIV  
neptunian dyke  
neptunian satellites  
neptunism  
Nereid  
Nereites  
Nereus  
neritic province  
neritic zone  
Neso  
nesosilicates  
net flux radiometer  
net slip  
network-former  
network-modifier  
neural arch  
neural spine  
neutral fold  
neutrally buoyant float  
neutral soil  
neutron-activation analysis  
neutron-gamma sonde  
neutron log  
neutron moisture meter  
neutron-neutron sonde  
neutron soil-moisture probe  
névé  
Newer Drift

New Horizons  
New Millennium Deep Space-1  
New Millennium Deep Space-2  
New Millennium Program  
New Red Sandstone  
newton  
Newtonian behaviour  
Newtonian fluid  
Newton's law of gravitation  
NEXRAD  
Next Generation Weather Radar  
Ngaterian  
Niagaran  
niccolite  
Nicol, William  
nicol prism  
NigeriaSat-2  
NigeriaSat-X  
Niggli, Paul  
Niggli method  
nimbostratus  
NIMS  
niobite  
nitisols  
nitratine  
nitre  
nitric horizon  
nitrogen fixation  
nival  
nivation  
NMO  
NMP  
NOAA-20  
NOAA-20/JPSS-1

NOAA-class satellite  
Noachian  
NOAA POES Series  
noctilucent clouds  
nocturnal radiation  
node  
nodes  
nodular  
nodule  
Noginskian  
Noguerornis  
noise  
nomogenesis  
non-conformity  
non-dipole field  
non-frontal depression  
non-metallic  
non-polarizable electrode  
non-recoverable strain  
non-renewable resource  
non-rotational strain  
non-selective scattering  
non-sequence  
non-spectral hue  
non-steady flow  
non-strophic  
non-umbilicate  
non-uniform flow  
norbergite  
Nor'easter  
Norian  
norite  
norm  
normal distribution

normal fault  
normal field  
normal incidence  
normalized vegetation index  
normally consolidated clay  
normal moveout  
normal problem  
normal stress  
normal travel time  
normal twins  
normal zoning  
normative constituents  
normative mineral  
NorSat-1 and -2  
norte  
North Alpine foreland basin  
North American Plate  
North Atlantic deep water  
North Atlantic Drift  
norther  
'northern lights'  
Northgrippian  
North Pacific current  
nor'wester  
nosean  
Nothosauria  
nothosaurs  
notochord  
notothyrium  
nova  
novaculite  
NovaSAR-S  
NRM  
nSight-1

nuclear–magnetic log  
nuclear-precession magnetometer  
nuclear waste  
nucleation  
nucleic acids  
nucleosynthesis  
nucleus  
nucleus number  
nuclide  
nuée ardente  
Nukumaruan  
Nullaginian  
null hypothesis  
null point  
numerical aperture  
numerical taxonomy  
Nummulites  
Nuna  
nunatak  
Nunivak  
Nusselt number  
nutation  
nutrient cycle  
Nyquist frequency  
oasis  
obduction  
Obdurodon  
Oberon  
Obik Sea  
objective  
oblate  
oblate uniaxial strain  
oblique extinction  
oblique-slip fault

oblique-slip margin  
obliquity of the ecliptic  
obrusion  
obrution  
obrution deposit  
obsequent  
observation well  
obsidian  
Occam's razor  
occipital  
occipital condyle  
occluded front  
occlusion  
ocean  
Ocean-1B  
ocean-basin crustal type  
ocean-basin floor  
ocean current  
ocean gyre  
oceanic crust  
oceanicity  
oceanic plateau  
oceanic trench  
oceanic-trench crustal type  
ocean-island basalt  
OceanSat-2  
oceanus  
ocean wave  
Ochoan  
ochre  
ochric horizon  
Ockham's razor  
OCO-2  
octahedrite

octahedron  
Octocorallia  
octocorals  
ocular  
OD  
Odderade  
Oddo–Harkins rule  
Odin  
Odintsovo  
ODP  
Oe  
oedometer  
oersted  
offlap  
offset  
offshore bar  
offshore zone  
ogive  
Ohauan  
Ohm's law  
OIB  
oikocryst  
oil  
oil immersion  
oil shale  
Oka/Demyanka  
okta  
Older Drift  
Older Dryas  
Oldest Dryas  
Oldham, Richard Dixon  
Old Red Sandstone Continent  
Olduvai  
Olenekian

Olenelloidea  
oligo-  
Oligocene  
Oligochaeta  
oligoclase  
oligohaline water  
oligomictic  
oligotaxic times  
oligotrophic  
olisthostrome  
olistolith  
olistostrome  
olivine  
olivine dolerite  
omphacite  
oncoid  
oncolite  
oncolith  
one-circle reflecting goniometer  
Onesquethawian  
one-way travel time  
onion weathering  
onlap  
Onnian  
Onondagan  
Ontarian  
ontogenetic heterochrony  
ontogeny  
Onverwacht  
onyx  
oo-  
oobiosparite  
ooid  
oolite

oolith  
oolitic  
oolitic limestone  
oomicrite  
oomoldic porosity  
oomouldic porosity  
18O:16O ratio  
Oort cloud  
oosparite  
ooze  
opacus  
opal  
opalescence  
opaque mineral  
open-cast mining  
open fold  
open form  
open hole  
open-pit mining  
open system  
operculum  
Ophelia  
ophicalcite  
Ophiocistioidea  
ophiocistoids  
ophiolite  
Ophiomorpha  
ophitic texture  
Ophiuroidea  
Opisthobranchia  
opisthodetic  
opisthogyral  
opisthogyrate  
opisthoparian suture

opisthosoma  
Opoitian  
Oppel, Albert  
Oppel zone  
optical continuity  
optical emission spectrum  
optical goniometry  
optical indicatrix  
Optical Satellite-3000  
optic axis  
OptSat-3000  
opx  
Or  
o-ray  
orbicular  
orbicular texture  
orbit  
orbital forcing  
orbit period  
Orbiting Carbon Observatory-2  
OrbView-5  
order  
Ordian  
ordinary ray  
Ordnance Survey  
Ordovician  
Ordovician V  
Ordovician VI  
ore  
orebody  
ore genesis  
ore grade  
ore microscope  
ore microscopy

ore mineral  
Oretian  
organic soil  
organic weathering  
Oriental faunal realm  
orientation survey  
original horizontality  
origination  
Oriskanyan  
Ornithischia  
ornithischian dinosaur  
ornithomimid  
Ornithopoda  
orogen  
orogenesis  
orogenic belt  
orogenic cycle  
orogeny  
orographic  
Orosirian  
orpiment  
Orthida  
orthids  
orthite  
orthoamphiboles  
orthochemical  
orthoclase  
orthoconglomerate  
orthoconic  
orthoferrosilite  
orthogenesis  
orthogonal thickness  
orthomagmatic  
orthophotograph

orthopyroxene  
orthoquartzite  
orthorhombic  
orthorhombic amphiboles  
orthoscopic  
orthoselection  
orthosilicates  
ortstein  
oryctognosy  
Osagean  
Osborn, Henry Fairfield  
oscillation ripple  
oscillatory wave  
oscillatory zoning  
osculum  
osmosis  
osmotic potential  
ossicles  
Osteichthyes  
osteostracans  
Osteostraci  
ostia  
Ostracoda  
Ostracodermi  
ostracoderms  
ostracum  
ostrich dinosaur  
Ostwald's step rule  
Otaian  
Otamitan  
Otapirian  
Otariidae  
Oteke  
ottrelite

outburst flood  
outcrop  
outerarc  
outer planet  
outgassing  
outgroup  
outlet glacier  
outlier  
out-of-phase component  
outwash  
outwash plain  
outwelling  
overbank deposit  
overbreak  
overburden  
overburden ratio  
over-consolidated clay  
overflow channel  
overflowing well  
overflow levée  
overfold  
overland flow  
overlap  
overpressure  
overspecialization  
overstep  
overthrust  
overturned fold  
overvoltage  
Owen, Richard  
oxbow  
Oxfordian  
oxic horizon  
oxidation

oxidation potential  
oxidation–reduction  
oxides  
oxisol  
oxycone  
oxygen ‘burning’  
oxygen-isotope analysis  
oxygen-isotope curve  
oxygen-isotope ratio  
oxygen isotopes  
oxygen-isotope stage  
oxyhornblende  
oxyluminescence  
Oyashio current  
ozone layer  
ozonesonde  
Pa  
Pachycephalosauridae  
pachydont  
Pacific- and Indian-Ocean common water  
Pacific-Antarctic Ridge  
Pacific Ocean  
Pacific Plate  
Pacific Province  
Pacific Ring of Fire  
Pacific-type coast  
Pacific-type margin  
packed biomicrite  
packer test  
packstone  
paedomorphosis  
pahoehoe  
Paibian  
paired metamorphic belts

pakakota  
Palaeartic faunal realm  
palaeo-  
Palaeoarchaean  
palaeoautecology  
palaeobiogeography  
palaeobiology  
palaeobotany  
Palaeocene  
Palaeocene-Eocene thermal maximum  
palaeoclimatic indicator  
palaeoclimatology  
palaeocurrent analysis  
Palaeodictyon  
palaeoecology  
palaeoflow  
palaeofluminology  
Palaeogene  
palaeogeography  
palaeoguild  
Palaeohelikian  
palaeohydraulics  
palaeoichnology  
palaeolatitude  
palaeomagnetic pole  
palaeomagnetism  
palaeoniscids  
palaeontology  
palaeoplacer  
Palaeoproterozoic  
Palaeopterygii  
palaeoseismology  
palaeosol  
palaeosome

palaeospecies  
palaeosynecology  
palaeotaxodont  
Palaeotaxus rediviva  
Palaeotethys  
Palaeozoic  
palagonite  
palate  
paleo-  
Paleocene  
palimpsest  
palingenesis  
palinspastic map  
Pallas  
pallasite  
pallial line  
pallial sinus  
pallium  
Palmer method  
palsa  
paludal  
palus  
palustrine  
palynology  
palynomorph  
pampero  
Pan  
pan  
Panama Isthmus  
Pancrustacea  
pandemic distribution  
Pandora  
Pangaea  
panhole

panidiomorphic  
panning  
Pannotia  
pannus  
panplain  
panplane  
Panthalassa  
Pantotheria  
papagayo  
paper shale  
para-  
Parablastoidea  
parabolic dune  
paraclade  
paraconglomerate  
Paracrinoidea  
paracycle  
paradigm  
paragenesis  
paragenetic sequence  
paragonite  
paralic  
parallax  
parallel evolution  
parallel extinction  
parallel fold  
parallel twins  
paramagnetism  
parameter  
parametral form  
parametral plane  
paramorph  
parapatric  
paraphyletic

Parapithecidae  
parasequence  
parasequence set  
parasitic cone  
parasitic fold  
parasitic magnetization  
PARASOL  
parastratotype  
parataxon  
Paratethys  
paratype  
parcel of air  
Pareiasauridae  
parent  
parent material  
pargasite  
parivincular  
parna  
parsimony  
partial annealing  
partial melting  
partial pressure  
partial range zone  
particle density  
particle shape  
particle size  
particle velocity  
parting lineation  
partition coefficient  
pascal  
pascichnia  
Pasiphae  
pass band  
passive margin

passive margin basin  
passive margin-distal orogen basin  
passive microwave  
passive remote sensing  
Pasteur effect  
Pasteur point  
Pastonian  
patch  
patch reef  
'patella' beach  
patellate  
Patella vulgata  
patera  
paternoster lake  
patterned ground  
Pauling's rules  
Pauropoda  
pavement  
Payntonian  
paystreak  
P-band  
PBRs  
PcP-wave  
PE  
peacock ore  
peak shear strength  
peak zone  
pearl-necklace lightning  
pearlspar  
pearly  
peat  
peat-borer  
peat podzol  
peat soil

pebble  
pectolite  
pectoral girdle  
ped  
pedestal rock  
pedicel  
pedicle  
pedicle foramen  
pediment  
pedion  
pediplain  
pedofacies  
pedogenesis  
pedogeomorphology  
pedology  
pedon  
pedoturbation  
peel technique  
pegmatite  
pegmatitic  
pel-  
pelagic  
pelagic ooze  
pelagic sediment  
pelagite  
Peléean eruption  
Pelecypoda  
Pelé's hair  
Pelé's tears  
pelite  
pelitic  
pellet limestone  
pellets  
pellicular water

pelmatozoan  
pelmicrite  
peloid  
pelosols  
pelsparite  
pelvic girdle  
pelvis  
Pelycosauria  
Penck, Albrecht  
Penck, Walther  
pendent  
Pendleian  
pene-  
peneplain  
peneplane  
penesaline  
penetration test  
penetration twin  
penitentes  
Pennaes  
pennate diatoms  
Pennatulacea  
penninite  
Pennsylvanian  
pentadactyl  
pentagonal dodecahedron  
pentameral symmetry  
Pentamerida  
pentamerids  
pentectic point  
Pentevrian  
pentlandite  
Penutian  
peperite

peralkaline  
peraluminous  
peramorphosis  
perched aquifer  
perched block  
percolation  
percussion boring  
Perdita  
perennial stream  
pergelic  
pergelisol  
peri-  
periclaste  
periclinal fold  
pericline  
pericline twinning  
pericontinental sea  
periderm  
Peridinales  
peridotite  
peridotite model  
perigee  
periglacial  
periglacial morphoclimatic zone  
perignathic girdle  
perihelion  
period  
periodic bedrock ridges  
periostracum  
peripatric speciation  
peripheral foreland basin  
periproct  
Perissodactyla  
peristome

peritectic point  
peritidal  
perlite  
perlitic  
perlucidus  
permafrost  
permafrost table  
permanent strain  
permanent wilting percentage  
permanent wilting point  
permeability  
permeameter  
Permian  
permineralization  
permitted intrusion  
permittivity  
perovskite  
perovskite model  
persistence  
perthite  
Peru–Chile Trench  
Peru current  
PeruSat-1  
petals  
petit-spot volcano  
petrification  
petrocalcic horizon  
petrofabric  
petrogenesis  
petrogenetic grid  
petrographic microscope  
petrography  
petrogypsic horizon  
petroleum

petroleum geology  
petrological microscope  
petrology  
petrophysics  
p-form  
PGF  
pH  
phaceloid  
phacoidal  
phacolith  
Phacopida  
Phaeophyceae  
phaeozems  
phaneritic  
phanerocrystalline  
Phanerozoic  
phase  
phase angle  
phase boundary  
phase diagram  
phase layering  
phase rule  
phase transitions in the mantle  
phase velocity  
phena  
phenoclast  
phenocryst  
phenogram  
phenon  
phenotype  
Philippine Plate  
Philippine Scientific Earth Observation Microsatellite-1  
phi scale  
PHL-Microsat-1

phlogopite  
Phobos  
Phocidae  
Phoebe  
Phoenix Plate  
Pholas  
phonolite  
phosphates  
phosphorescence  
phosphorite  
photo-  
photochemical smog  
photodisintegration  
photodissociation  
photogeology  
photogrammetry  
photographic infrared  
photohydrometer  
photometer  
photometry  
photon log  
photopolarimeter-radiometer  
photosphere  
photosymbiosis  
photosynthesis  
phragma  
Phragmites cliffwoodensis  
phragmocone  
phreatic activity  
phreatic zone  
phreatomagmatic activity  
phreatoplinian deposit  
Phycosiphon  
phyletic

phyletic evolution  
phyletic gradualism  
phyllite alteration  
phyllite  
phyllonite  
phyllosilicate  
phylogenetics  
phylogenetic tree  
phylogenetic zone  
phylogeny  
phylozone  
phylum  
-phyre  
physico-theology  
phytogeography  
phytophagous  
Piacenzian  
pick  
pickup  
pico-  
picoplankton  
picosatellite  
Picosatellite for Remote-sensing and Innovative Space Missions  
picrite  
pi diagram  
piedmont  
piedmont glacier  
piercing fold  
piezoelectricity  
piezometer  
piezometric surface  
piezoremanent magnetization  
pigeonite  
piggyback basin

piggyback thrust sequence

Pikaia

pile

pileus

pillar and stall

pillar structure

pillow lava

pilotaxitic

pinacoid

pinch-and-swell

pinger

pingo

pinite

pinnacle reef

pinnate fractures

pinnular plates

pinnules

*Pinus longaeva*

PIOCW

Pioneer

pipe

pipette analysis

piprake

Piripauan

pisoid

pisolith

pisolithic

pistacite

piston corer

piston sampler

*Pistosaurus grandaevus*

pitch

pitchblende

pitchstone

pitting  
pixel  
pixel colour  
pixie  
PKP-wave  
placental mammals  
placer deposit  
place value  
placic horizon  
Placodermi  
Placodontidae  
placoid scale  
plaggen  
plagioclase feldspar  
plagioclase series  
plagiogranite  
plana  
planar cross-stratification  
planar slide  
planar slump  
planation surface  
plane bed  
plane of projection  
plane of symmetry  
plane-polarized light  
plane strain  
planetary boundary layer  
planetary geodesy  
planetary geology  
planetary geomorphology  
planetary geoscience  
Planet-B  
planetismal  
planetismal hypothesis

plane wave  
planèze  
planispiral  
planitia  
plankton  
planktonic geochronology  
Planolites  
planosols  
Plantae  
planum  
plasma  
plasma engine  
plasma instrument  
plasma wave sub-system  
plastic deformation  
plasticity index  
plastic limit  
plastron  
plate  
plateau basalt  
plate bearing test  
plate boundary  
plate kinematics  
plate margin  
plate motions  
plate stratigraphy  
plate tectonics  
platform  
platform conodonts  
platy  
platykurtic  
play  
playa  
Playfair, John

Plectronoceras cambria  
Pleiades-HR  
Pleistocene  
Pleistocene refugium  
Pleistogene  
pleochroic halo  
pleochroism  
pleonaste  
plesiomorph  
plesiomorphic  
plesion  
Plesiorycteropus  
Plesiosauroidea  
plesiosaurs  
pleura  
pleural  
pleuron  
plicate  
Pliensbachian  
Plinian eruption  
plinthic horizon  
plinthite  
plinthosols  
Pliocene  
plough mark  
plug  
plunge  
plunging breaker  
plus–minus method  
Pluto  
pluton  
plutonic  
plutonism  
pluvial period

pneumatolysis  
poached soil  
Podolskian  
Podopteryx mirabilis  
podsol  
podsolization  
podzol  
podzolic soils  
podzolization  
podzols  
Pogonophora  
poikil-  
poikilitic  
poikiloaerobic  
poikiloblast  
poikilotherm  
poikilotopic  
point bar  
point counting  
point group  
point load index  
point load tester  
point source  
Poisson distribution  
Poisson's ratio  
polar air  
polar-air depression  
polar climate  
polar-desert soil  
polar front  
polar-front jet stream  
polar glacier  
polarimetry  
polarity

polarity chron  
polarity chronozone  
polarity epoch  
polarity event  
polarity excursion  
polarity interval  
polarity reversal, geomagnetic  
polarity subchron  
polarity subchronozone  
polarity subzone  
polarity superchron  
polarity superchronozone  
polarity superzone  
polarity timescale  
polarity transition period  
polarity zone  
polarization, electrode  
polarization colours  
polarized radiation  
polarizer  
polarizing microscope  
Polaroid  
polar orbit  
Polar Orbiting Passive Atmospheric Calibration Sphere  
polar stratospheric cloud  
Polar units  
polar wander path  
polder  
pole of a face  
pole of rotation  
'Polflucht'  
polished section  
polished stone value  
polishing relief

polje  
pollen  
pollen analysis  
pollen-assemblage zone  
pollen diagram  
pollen zone  
poloidal field  
poly-  
polyanions  
Polychaeta  
polycyclic landscape  
polygenetic  
polygonization  
polyhaline water  
polyhalite  
polymetallic sulphide  
polymetamorphism  
polymictic  
polymineralic  
polymorph  
polymorphic  
polymorphic minerals  
polymorphic transformation  
polymorphism  
polypedon  
polyphase landscape  
polyphyletism  
Polyplacophora  
polysynthetic twin  
polytaxic times  
polytetrahedron  
Pomeranian  
ponente  
Pongola

ponor  
pool-and-riffle  
poorly washed biosparite  
POPACS  
Poperinge  
pop-shooting  
pop-up  
Porangan  
porcelain jasper  
porcellanite  
pore  
pore fluid pressure  
pore space  
pore-water pressure  
Porifera  
porosity  
porosity and permeability determination  
porphyrin  
porphyritic  
porphyroblast  
porphyroclast  
porphyry  
porphyry copper  
porphyry deposit  
porphyry gold  
porphyry molybdenum  
Portia  
Portlandian  
positive inversion  
post-  
post-depositional remanent magnetization  
post-deuteric alteration  
postdisplacement  
posterior

post-glacial  
post-perovskite  
postseismic slip  
post-tectonic  
postzygapophyses  
potassium–argon dating  
potassium–calcium dating  
potassium feldspar  
potential electrode  
potential energy  
potential evapotranspiration  
potential instability  
potential reserve  
potential temperature  
potentiometric surface  
pot-hole  
Potsdam gravity  
Potter’s flood-peak formula  
Poundian  
powder photograph  
powder technology  
powellite  
power-law creep  
pozzolan  
p-parameter  
ppb  
PPL  
ppm  
PPR  
p-process  
praecipitatio  
praedichnia  
Praghian  
Prandtl number

Pratt, John Henry  
Pratt model  
pre-  
pre-adaptation  
Preboreal  
Precambrian  
precession  
precession of the equinoxes  
precipitable water  
precipitation  
precipitation-efficiency index  
precipitation fog  
precision  
predator  
predisplacement  
pre-ferns  
preferred orientation  
pre-Hadean  
prehnite  
prehnite-pumpellyite facies  
Pre-Imbrian  
Pre-Nectarian system  
pre-splitting  
pressure-depth profile  
pressure dissolution  
pressure fringe  
pressure-gradient force  
pressure head  
pressure melting  
pressure ridge  
pressure shadow  
pressure solution  
pressure-tube anemometer  
pressure wave

pressure welding  
pre-tectonic  
pre-Tolstojan  
prevailing wind  
prey  
prezygapophyses  
Priabonian  
Priapulida  
Pridoli  
primary creep  
primary crushing  
primary geochemical differentiation  
primary geochemical dispersion  
primary magma  
primary melt  
primary migration  
primary mineral  
primary porosity  
primary productivity  
primary sedimentary structure  
primary wave  
primitive  
primitive circle  
primordial  
principal component analysis  
principal point  
principal shock  
principal strain axes  
principal strain ratio  
principle stress axes  
principle of contained fragments  
principle of included fragments  
principle of superposition  
principle of uniformitarianism

Priscoan  
PRISM  
prism  
PRISMA  
prismatic  
protactinium dating  
protactinium/thorium isotopes  
probability density function  
probability distribution  
PROBA-V  
'Problematica'  
Proboscidea  
proboscidean  
Procaryotae  
Procellarum system  
processes, geomorphological  
process-response system  
prochoanitic  
Procolophonia  
Procyonidae  
pro-delta  
prod mark  
production log  
production well  
Productus giganteus  
Proetida  
profiling  
pro-foreland basin  
Proganochelys quenstedii  
progenesis  
proglacial  
progradation  
prograde metamorphism  
progressive deformation

progressive evolution  
progressive metamorphism  
progressive wave  
Progymnospermopsida  
progymnosperms  
Project for On-Board Autonomy – Vegetation  
prokaryote  
prokaryotic  
prolate  
prolate uniaxial strain  
proloculus  
Prometheus  
promontorium  
proparian  
proper motion  
propylitization  
proseptum  
prosobranch gastropods  
prosogyral  
prosogyrate  
prosoma  
Prospero  
protalus rampart  
protaspis  
Proterosuchus  
Proterozoic  
Proteus  
protist  
Protista  
proto-  
proto-Atlantic  
Protoaulopora ramosa  
Protoceratops andrewsi  
protoconch

Protoctista  
protolith  
proton-adding process  
proton magnetometer  
protoparian suture  
protoplanet  
Protopteridales  
protore  
protostar  
protozoa  
protrusive  
proved reserve  
provenance  
province  
provinciality  
proximal  
proximity log  
proxy  
Psamathe  
psammite  
PSC  
psephite  
pseudobreccia  
pseudoextinction  
pseudofossil  
pseudo-gravitational field  
pseudo-magnetic field  
pseudomorph  
pseudonodule  
pseudopunctate  
pseudosection  
pseudospar  
Pseudosycidium  
pseudotachylite

psilomelane  
Psilonichnus  
Psilophytales  
psilophytes  
Psilopsida  
PSV  
psychrometer  
psychrophile  
Pteraspida  
Pteraspis  
Pteridophyta  
pteridophytes  
Pteridospermales  
Pterobranchia  
pteropod ooze  
Pteropsida  
Pterosauria  
Ptilograptidae  
Ptychopariida  
ptygmatic fold  
Puaroan  
pubis  
Puck  
puddingstone  
puddled soil  
Puercan  
pull-apart basin  
pulsations, geomagnetic  
pulse length  
pumice  
pump  
pumpellyite  
pumping test  
punctae

punctate  
punctuated equilibrium  
pupaeiform  
Purbeckian  
pure shear  
purga  
pushbroom system  
push moraine  
push-pull wave  
Pusgillian  
puy  
P-wave  
PWS  
pycnocline  
pycnometer  
pygidium  
pyknometer  
pyralspite  
pyramid  
pyramidal peak  
pyrite  
pyritohedron  
pyro-  
pyrochlore  
pyroclastic  
pyroclastic flow  
pyroelectricity  
pyrogenetic minerals  
pyrolite model  
pyrolusite  
pyrolysis  
pyrometasomatic deposit  
pyromorphite  
pyrope

pyrophyllite  
pyroxene  
pyroxene gneiss  
pyroxene hornfels facies  
pyroxenite  
pyroxenoid  
pyrrhotite  
p-zone  
Q  
QAPF classification  
QAP triangle  
QBITO  
Q days  
Q-factor  
QFL diagram  
QFL triangle  
Qn  
QSat-EOS  
quadrature  
quaquaversal  
quarrying  
quartz  
quartz arenite  
quartz dolerite  
quartzite  
quartz overgrowth  
quartz porphyry  
quartz sandstone  
quartz wacke  
quartz wedge  
quasi-equilibrium  
quasi-section  
Quaternary  
quaternary system

quenching  
Quetzalcoatlus northropi  
quick clay  
quickflow  
quiet days  
quiet zone  
QuikSCAT  
Qungzusian  
Q-wave  
Ra  
radar  
radar altimetry  
radar cross-section  
radar imaging  
Radar Imaging Satellite  
Radar in a CubeSat  
RADARSAT  
radar scattering coefficient  
radar scatterometer  
radial drainage  
radial dykes  
radial fault  
radial fibrous  
radial relief displacement  
radial symmetry  
radiance  
radiant flux density  
Radiata  
radiating  
radiation  
Radiation Belt Storm Probes  
radiation budget  
radiation densimeter  
radiation fog

radiation inversion  
radiation night  
radiation tracks  
radiatus  
radial  
radioactive decay  
radioactive logging  
radioactive survey  
radioactive waste  
Radio Aurora eXplorer  
radiocarbon dating  
radiogenic  
radiogenic heating  
radioimmunoassay  
Radiolaria  
radiolarian earth  
radiolarian ooze  
radiolarite  
radiometer  
Radiometer Assessment using Vertically Aligned Nanotubes  
radiometric dating  
radiometry  
radio occultation  
radiosonde  
Radiozoa  
radius  
radius ratio  
raft foundation  
rainbow  
RainCube  
raindrop  
rain-gauge  
rain-making  
rain print

rain-shadow  
rain-splash  
rain-wash  
raised beach  
raised bog  
rake  
Ramapithecus  
rammer  
ramose  
ramp  
rampart craters  
Ramsden eyepiece  
Randian  
random sampling  
range  
Ranger  
range zone  
rank  
rapakivi texture  
RapidEye Earth Observation Constellation  
rapid flow  
rapid-neutron process  
Rapitan-type granular iron formation  
rare-earth element  
RASAT  
raster  
ratio  
Raukumara  
RAVAN  
ravinement surface  
ravine wind  
rawinsonde  
Rawtheyan  
RAX

Rayleigh, Lord  
Rayleigh criterion  
Rayleigh number  
Rayleigh scattering  
Rayleigh wave  
rays  
RBSP  
RE  
reaction time  
reactivation surface  
real-aperture radar  
real component  
realgar  
realistic reaction  
recapitulation of phylogeny  
Recent  
recessional moraine  
recharge  
recharge area  
reclined  
reclined fold  
reconstructive transformation  
recovery factor  
recrystallization  
rectangular drainage  
rectilinear slope  
rectimarginate  
recumbent fold  
red algae  
red-bed copper  
red beds  
red clay  
red copper ore  
red edge

red iron ore  
Redlichiida  
redox potential  
redox reaction  
red podzolic soil  
Red Sea  
red sprite  
reduction  
reduction potential  
reduction to pole  
REE  
reef  
reef flat  
reef front  
reef trap  
re-entrant  
reference ellipsoid  
reference section  
refertilization  
reflectance  
reflectance spectrometry  
reflected infrared  
reflected-light microscopy  
reflection  
reflection coefficient  
reflection pleochroism  
reflectivity  
reflector  
reflexed  
reflux theory  
refolded fold  
refraction  
refraction survey  
refractive index

refractometer  
refractory mineral  
refugia  
Refugian  
reg  
regelation  
regio  
regional field  
Regionally Important Geological/Geomorphological Sites  
regional metamorphism  
regional stratigraphic scale  
regolith  
regosols  
regression  
regressive systems tract  
regular echinoids  
Regulares  
Reid, Harry Fielding  
REIMI  
rejuvenation  
relative age  
relative humidity  
relative permittivity  
relative plate motion  
relative pollen frequency  
relative timescale  
relative vorticity  
relaxation  
relaxation frequency  
relaxation time  
releasing bend  
relict  
relict land-form  
relict sediment

relict structure  
relief  
Relizian  
remanent magnetization  
remanié  
remanié beds  
remnant arc  
remote sensing  
remoulding  
removal time  
rendzina  
renewable resource  
reniform  
repeated twinning  
Repettian  
repichnia  
replacement  
reptation  
Reptilia  
reptiles  
Reptilomorpha  
resequent  
resequent fault-line scarp  
reserve  
reservoir  
reservoir pool  
reservoir rock  
reshabar  
residence time  
residual deposit  
residual gravity map  
residual shear strength  
resinite  
resinous

resistate mineral  
resistivity logging  
resistivity methods  
resonance  
resorption  
ResourceSat-2 and -2A  
restraining bend  
resurgence  
resurgent caldera  
Resurs-DK1  
reticula  
reticulated  
reticulite  
reticulum  
retro-arc basin  
retro-arc foreland basin  
retrochoanitic  
retrograde  
retrograde metamorphism  
retrogression  
retrogressive metamorphism  
retrosiphonate  
retrusive  
return flow  
return period  
Réunion  
reverberation  
reversal  
reversal timescale  
reversed field  
reversed kink band  
reverse fault  
reverse zoning  
reversing dune

revolving storm  
reworked  
Reykjanes Ridge  
Reynolds number  
rhabdosome  
Rhaetian  
Rhea  
rheid flow  
rheidity  
rhenium–osmium dating  
rheology  
Rhine graben  
Rhipidistia  
Rhizocorallium  
rhodochrosite  
rhodonite  
Rhodophyceae  
rhombododecahedron  
rhombic dodecahedron  
rhombochasm  
rhombohedral  
rhombohedral  
rhomboid  
Rhuddanian  
Rhyacian  
Rhynchocephalia  
rhyncholite  
Rhynchonellida  
rhynchonellids  
Rhynia  
Rhyniophytina  
rhyodacite  
rhyolite  
rhythmic sedimentation

rhythmite  
ria  
ribbon bomb  
ribbon jasper  
ribbon lakes  
ribbons  
Richmondian  
Richter, Charles Francis  
Richter denudation slope  
Richter scale  
Ricker pulse  
ridge  
ridge-and-ravine topography  
ridge and runnel  
ridge crest  
ridge-push  
riebeckite  
Riedel shear structure  
riegel  
riffle  
rift  
rift valley  
right lateral fault  
rigidity modulus  
RIGS  
rill  
rille  
rill-wash  
rima  
rime  
rimmed shelves  
rim syncline  
ring canal  
ring-dyke

ringed basin  
ring fracture  
ringing  
Ring of Fire  
ringshine  
ring silicate  
ringwall  
Ringwood's rule  
rip current  
Riphean  
rippability  
ripple  
ripple-drift cross-lamination  
ripple index  
ripple mark  
ripple-symmetry index  
ripple train  
rip-rap  
RISAT  
Rising-2  
Riss  
Riss/Würm Interglacial  
river capture  
river grade  
river profile  
river-sediment analysis  
river terrace  
river water, major constituents  
RMQ  
road base  
Roadian  
roadstone  
roadway construction  
roaring forties

Roche limit  
roche moutonnée  
rock  
rock bench  
rock bolt  
rock burst  
rock crystal  
rock drumlin  
rock fall  
rock fill  
rock flour  
rock glacier  
rock head  
Rocklandian  
rock mass  
rock-mass quality  
rock-mass strength  
rock mechanics  
rock pavement  
rock-quality designation  
rock salt  
rock slide  
rock:soil ratio  
rock-stratigraphic unit  
rock-structure rating  
rock unit  
Rodda's flood-peak formula  
rodding structure  
Rodebaek  
rodingite  
Rodinia  
rogen moraine  
roll along  
rollover

rollover anticline  
roll-type uranium ore  
Romer, Alfred Sherwood  
roof pendant  
roof thrust  
room and pillar  
rooted tree  
rootlet  
root-mean-square velocity  
root zone  
rope bomb  
Rosalind  
rose diagram  
Rose Garden  
rose quartz  
Rosetta  
Rossby waves  
Rossi–Forel scale  
rostral suture  
rostrum  
rotary drilling  
rotating-cups anemometer  
rotational remanent magnetism  
rotational shear  
rotational slip  
rotational slump  
rotation of the Earth  
Rotliegende  
Rotliegendes  
rotor cloud  
roughness  
rounded biosparite  
roundness index  
Rowe cell

RPF  
r-process  
RQD  
RRM  
RRR junction  
R-selection  
RSR  
RSS  
RST  
rubble levée  
rubefaction  
rubidium–strontium dating  
ruby  
rudaceous rock  
rudist bivalves  
rudite  
rudstone  
Rugosa  
rugose  
rule of Vs  
Rumford, Count  
Runangan  
Runcorn, Stanley Keith  
runnel  
Rupelian  
rupes  
Rusophycus  
Russian borer  
Rutherford, Ernest  
rutile  
ruware  
Ryukyu Trench  
Ryzanian  
S

Saalian  
sabkha  
saccate  
Saccominopsis  
saccus  
saddle  
safe bearing capacity  
Saffir–Simpson Hurricane Wind Scale  
sag and swell topography  
sagduction  
SAGE-III  
sag pond  
St David's  
Sakian  
Sakigake  
Sakmarian  
salcrete  
salic  
salic horizon  
salina  
salination  
saline giant  
saline-sodic soil  
saline soil  
salinity  
salinization  
SALR  
salt  
saltation  
salt dome  
salt-dome trap  
salt fingering  
salt flat  
salt lake

salt marsh  
salton  
salt pan  
saltpetre  
salt weathering  
salt wedge  
samarium–neodymium dating  
sampling frequency  
sampling interval  
sampling methods  
sand  
sandbody  
Sander's symmetry principle  
sand line  
sand ribbon  
sand sheet  
sandstone  
sandstone dyke  
sandstorm  
sandur  
sand volcano  
sand wave  
Sangamonian  
sanidine  
Santerian  
Santonian  
Sao  
sapphire  
saprofite  
saprofite  
saprofite coal  
saprofite  
SAR  
SARAL

SAR-Lupe  
Sarcopterygii  
sardonix  
Sargasso Sea  
sarl  
saros unit  
sarsen stone  
Sartan  
sastrugi  
Satélite de Coleta de Dados  
satellite  
satellite photography  
satellite sounding  
Satellite with ARGos and ALtiKa  
satin spar  
saturated  
saturated adiabatic lapse rate  
saturated air  
saturated flow  
saturation  
saturation deficit  
saturation magnetization and moment  
saturation moisture content  
saturation vapour pressure  
Saturn  
saturnian satellites  
Saucesian  
saurian  
Saurischia  
saurischian dinosaur  
Sauropoda  
Sauropterygia  
Saussure, Horace Bénédict de  
saussuritization

Saxonian  
S-band  
scalenohedron  
scandent  
Scandinavian ice sheet  
scanning electron microscope  
scapolite  
scar  
scarp  
scarp, lobate  
scarp-and-vale topography  
scarp-foot knick  
scarp retreat  
scarp slope  
SCATSat-1  
scatter diagram  
scattering  
Scatterometer Satellite-1  
scavenger  
scavenging  
SCD  
scheelite  
Scheinwoodian  
Scheuchzer, Johann Jacob  
schiller  
Schindewolf, Otto H.  
schist  
schistosity  
schizochroal  
schizodont  
Schizomycophyta  
schlieren  
Schlumberger array  
Schmidt hammer

Schmidt hammer test  
Schmidt–Lambert net  
schorl  
Schroeder Van Der Kolk method  
schuppen structure  
Schwassmann–Wachmann 3  
Science Satellite/Atmospheric Chemistry Experiment  
scintillation counter  
Scintillation Observations and Response of the Ionosphere to  
Electrodynamics  
Scintillation Prediction Observations Research Task  
scintillometer  
scintillometer survey  
sirocco  
SciSat-1/ACE  
Scleractinia  
sclerotinite  
SCLM  
scolecodont  
scolecoid  
Scolicia  
scopuli  
scopulus  
scoria  
scoriaceous  
scoria cone  
scorpions  
Scotia Plate  
scour and fill  
Scourian  
Scourian orogeny  
scour lag  
scree  
screw dislocation

scroll bar  
ScS-wave  
scud  
scute  
Scyphozoa  
Scythian  
sea  
sea-anemones  
sea breeze  
sea-floor spreading  
sea fret  
sea ice  
seamount  
sea pens  
seatearth  
sea water, major constituents  
SeaWiFS  
Secchi disc  
second arrival  
secondary blasting  
secondary creep  
secondary crushing  
secondary depressions  
secondary enrichment  
secondary front  
secondary geochemical differentiation  
secondary geochemical dispersion  
secondary matrix  
secondary migration  
secondary mineral  
secondary porosity  
secondary quartz  
secondary recovery methods  
secondary sedimentary structure

secondary wave  
second derivative  
secular variation  
SEDEX  
Sedgwick, Adam  
sedigraph  
sediment  
sedimentary basin  
sedimentary cycle  
sedimentary environments  
sedimentary exhalative processes  
sedimentary mélange  
sedimentary rock  
sedimentary rock classification  
sedimentary structure  
sedimentation, rate of  
sedimentation coefficient  
sedimentation tube  
sedimentology  
seed plants  
seep  
seepage  
seepage velocity  
seiche  
seif dune  
seism-  
seismic anisotropy  
seismic blind zone  
seismic gap  
seismicity  
seismic margin  
seismic moment  
seismic record  
seismic reflection

seismic-reflection profiling  
seismic refraction  
seismic section  
seismic stratigraphy  
seismic survey  
seismic tomography  
seismic velocity  
seismic wave  
seismic-wave modes  
seismic zone  
seismogram  
seismograph  
seismology  
seismometer  
seismostratigraphy  
seismotectonics  
Selandian  
SELENE  
selenite  
selenizone  
selenology  
self diffusion  
self-exciting dynamo  
self-potential method  
self-potential sonde  
self-reversal  
SEM  
Senecan  
Senonian  
Sensitive High Resolution Ion MicroProbe  
sensitive tint  
sensitivity  
Sentinel  
SEOSat/Ingenio

septa  
septarian nodule  
septomaxilla  
septum  
sequence stratigraphy  
serac  
SERB  
serein  
sericite  
series  
serir  
serpens  
serpentes  
serpenticone  
serpentine  
serpentine barrens  
serpentinite  
serpentinization  
Serpukhovian  
Serravallian  
sesquioxides  
sessile  
seston  
seta  
Setebos  
settlement  
settling lag  
sexual dimorphism  
Seymouria  
sferics  
S-fold  
SG  
SGCS  
shade temperature

shadow test  
shadow zone  
shaft well  
shale  
shale line  
shallowing-upward carbonate cycle  
shamal  
shape fabric  
sharp sand  
shatter cones  
shear box  
shear direction  
shear modulus  
shear plane  
shear strain  
shear strength  
shear stress  
shear wave  
shear zone  
sheath fold  
sheep-walk  
sheet flood  
sheet lightning  
sheet sand  
sheet silicate  
sheetwash  
Sheinwoodian  
shelf  
shell beak  
shell bed  
shell structure  
shelly limestone  
shelter porosity  
Shenzhou

shergottite/nakhlite/chassignite meteorites

Shermanian

shield

shield volcano

shingle

Shipka

shoal

shoaling

shoal retreat massif

shock metamorphism

shock-remanent magnetization

Shoemaker, Eugene Merle

shoestring sand

shonkinite

shooting flow

shoreface

shore platform

shortening

short wavelength infrared

shot

shot bounce

shotcrete

shot depth

shotpoint gap

shower

shrieking sixties

SHRIMP

shrimps

shrinkage

shrinkage cracks

shrinkage joint

shrub-coppice dune

shutter ridge

SH-wave

SI

Siberian high

sichelwannen

Sicilian

sicula

side-looking airborne radar

sidereal day

sidereal month

Siderian

siderite

siderolite

siderophile

side-scan sonar

sidewall corer

Sidufjall

Siegennian

siemens

sieve

sieve deposit

sieving

sigma-t density

signature

significant wave height

silcrete

silent quake

Silesian

silex

silica

silica-oversaturated rock

silica saturation

silicates

silica-undersaturated rock

siliceous ooze

siliceous sinter

siliciclastic  
silicification  
silicon 'burning'  
silky  
sill  
sillar  
sillimanite  
silt  
siltstone  
Silurian  
silva  
silvagenitus  
silver, native  
silver glance  
silver iodide  
silver spike  
silver thaw  
Simiiformes  
similar fold  
simoom  
simple shear  
Simpson, George Gaylord  
Sinemurian  
single couple  
single-stage lead  
singularity  
Sinian  
sinistral coiling  
sinistral fault  
sink  
sink-hole  
sinking  
Sinope  
Sinornis

sinter  
sinus  
siphon  
siphonal canal  
siphonate  
siphonostomatous  
siphuncle  
Sistema Satelital para la Observación de la Terra  
sister groups  
sister taxa  
site investigation  
Sivapithecus  
skarn  
skeletal limestone  
skeletal material  
skeletal micritic limestone  
skewness  
skin depth  
skip mark  
Skolithos  
SKS-wave  
skull  
Skylab  
SkySat  
Skythian  
sky-view factor  
slab-pull  
slake-durability test  
slaking  
slant range  
slant-range resolution  
SLAR  
slate  
slaty cleavage

sleet  
sleeve exploder  
slick  
slickenfibres  
slickenside  
slide  
slide-rock  
slingram method  
sling psychrometer  
slip  
slipface  
slip-off slope  
sloc  
slope angle  
slope processes  
slope profile  
slope stability  
slope stabilization  
slow earthquake  
slow-neutron process  
slow slip  
sluff  
slump  
slump structure  
Smålfjord  
small circle  
small satellite  
SMAP  
smarl  
SMC  
smectite  
Smith, William  
Smithian  
smithsonite

Smith's rule  
smog  
smoker  
SMOS  
SMOW  
smudging  
SNC  
Snell's law  
snout  
snowball Earth  
snowblitz theory  
snowflake  
snow-gauge  
snow grain  
snow line  
soapstone  
soda lake  
sodalite  
soda nitre  
sodication  
sodic soil  
sodium-adsorption ratio  
sodium feldspar  
sodium-sulphate soundness test  
SOFAR channel  
soil  
soil air  
soil anchor  
soil association  
soil-atmosphere survey  
soil borrow  
soil complex  
soil conservation  
soil formation

soil geomorphology  
soil grading curve  
soil horizon  
soil individual  
soil line  
soil lithology  
soil management  
soil mechanics  
Soil Moisture Active/Passive mission  
Soil Moisture and Ocean Salinity mission  
soil-moisture content  
soil-moisture deficit  
soil-moisture index  
soil-moisture regime  
soil profile  
soil separates  
soil series  
soil structure  
soil survey  
soil taxonomy  
soil variant  
soil-wash  
soil-water zone  
Sojourner  
sol  
sola  
solar abundance of elements  
solar constant  
solar cosmic rays  
solar flare  
solarimeter  
Solar Irradiance and Earth Radiation Budget  
solar magnetic variation  
solar nebula

solar pond  
solar radiation  
Solar Radiation and Climate Experiment  
solar system  
solar wind  
sole mark  
Solenopora  
sole structure  
sole thrust  
solfataric activity  
solid  
solid-melt equilibrium  
solid solution  
solid-state imaging camera  
solidus  
solifluction  
solifluxion  
solitary corals  
solodic soil  
solonchaks  
solonetz  
solstice  
solubility product  
solum  
solution  
solution channel  
solution cleavage  
solution pan  
solution pipe  
Solvan  
solvus  
solvus temperature  
Somali Plate  
sombric horizon

Somoholoan  
sonar  
sonde  
sonde self-potential  
sonic log  
sonic sonde  
sonobuoy  
sonograph  
Sorby, Henry Clifton  
SORCE  
Sordes pilosus  
sorosilicate  
sorption  
sorted biosparite  
SORTIE  
sorting  
Soudleyan  
sound channel  
sunder  
sound speed  
source region  
source rock  
South American Plate  
South-east Pacific Plate  
southerly burster  
Southern Ocean  
southern oscillation  
sövite  
SP  
spaced cleavage  
space lattice  
space–time substitution  
spallation  
spandrels of San Marco

spar  
sparite  
-sparite  
sparker  
sparse biomicrite  
spastolith  
Spathian  
spathose iron  
spatial frequency  
spatial-frequency filter  
spatter  
spatter cone  
spatter-fed flow  
specialization  
species  
species longevity  
species selection  
species zone  
specific gravity  
specific-gravity determinations  
specific humidity  
specific retention  
specific yield  
spectra  
spectral hue  
spectral radiance  
spectrochemical analysis  
spectrograph  
spectrometer  
spectrophotometer  
spectroradiometer  
spectroscope  
spectroscopic binary  
spectrum

specular  
specularite  
specular reflection  
spelean  
speleothem  
Spermatophyta  
spessartine  
spessartite  
sp. gr.  
sphaericone  
sphalerite  
sphene  
sphenoid  
Sphenopsida  
sphericity  
spheroid  
spheroidal oscillation  
spheroidal weathering  
spherule  
spherulite  
spicular chert  
spicule  
spiculite  
spider diagram  
spiders  
spike  
spilite  
spilling breaker  
spillway  
spinal column  
spindle bomb  
spinel  
spinifex texture  
spiny sharks

spire  
Spiriferida  
spiriferids  
spissatus  
spit  
splanchnocranium  
splay fault  
splendent  
SP method  
spodic horizon  
spodosols  
spodumene  
spondylium  
sponges  
Spongiaria  
spontaneous potential  
spontaneous-potential method  
sporangium  
spore  
SporeSat  
sporinite  
SPORT  
SPOT  
spotting  
spread  
spreading rate  
spreiten  
Spriggina  
spring  
spring balance  
Springerian  
spring sapping  
spring tide  
sprite halo

s-process  
SPS  
spur  
squall  
squall line  
square array  
squeeze-up  
squeezing ground  
Sq variation  
SSI  
SSOT  
SSS  
stability  
stability field  
stable isotope  
stable-isotope studies  
stack  
stacking  
stacking fault  
stacking velocity  
stade  
stadial  
staff gauge  
stage  
stage hydrograph  
stagnosols  
stains and staining techniques  
stalactite  
stalagmite  
Stampian  
standard deviation  
Standard Global Chronostratigraphic Scale  
standard mean open water  
Standard Stratigraphic Scale

standing wave  
stand of the tide  
stannite  
stapes  
star dune  
Stardust  
Starlette and Stella  
star pair  
star phylogeny  
star twinkling  
stasigenesis  
stasis  
Statherian  
static correction  
statics  
stationary front  
station frequency  
station interval  
stauroilite  
steady flow  
steam fog  
steatite  
S-tectonite  
Stefan–Boltzmann law  
Stegosauridae  
steinkern  
Steinmann trinity  
Stella  
stem group  
stem reptiles  
Stenian  
Steno, Nicolaus  
stenothermal  
stenotopic

Stensen, Niels  
step faulting  
Stephanian  
Stephano  
stepout  
steptoe  
stereogram  
stereographic net  
stereographic projection  
stereom  
stereonet  
stereophotography  
stereoptic vision  
stereoscope  
Steropodon  
Stettin  
stibnite  
stick-slip  
Stigmaria  
Stille, Wilhelm Hans  
stillstand  
stilpnomelane  
stipe  
stishovite  
stock  
stockwork  
Stokes's law  
stolon  
stolotheca  
stomodeum  
-stone  
stone canal  
stone circle  
stone garland

stone net  
stone polygon  
stone steps  
stone stripes  
stony-iron meteorite  
stony meteorite  
stoop and room  
stopping  
storage coefficient  
storativity  
storm  
storm beach  
storm bed  
storm deposit  
storm surge  
storm wave-base  
stoss  
stoss and lee  
stoss-and-lee topography  
Strahler climate classification  
straight extinction  
strain  
strain ellipse  
strain ellipsoid  
strain gauge  
strain marker  
strain parallelepiped  
strain rate  
strain–slip cleavage  
strain–time diagram  
strandflat  
strandline  
strandplain  
strata

strath terrace  
stratified sampling  
stratiform deposit  
stratiformis  
stratigraphic  
stratigraphic column  
stratigraphic correlation  
stratigraphic cross-section  
stratigraphic nomenclature  
stratigraphic reef  
stratigraphic scale  
stratigraphic trap  
stratigraphic unit  
stratigraphy  
stratocumulus  
stratomere  
stratopause  
stratophenetic classification  
stratophenetics  
stratosphere  
Stratospheric Aerosol and Gas Experiment-III  
stratotype  
stratovolcano  
stratum  
stratus  
streak  
streak lightning  
streak plate  
streamer  
stream flood  
stream grade  
streamline  
stream order  
stream power

stream-sediment analysis  
stream terrace  
stress  
stress axial cross  
stress difference  
stress field  
stress meter  
stress–strain diagram  
stress trajectory  
stretch  
strewnfield  
striation  
strike  
strike fault  
strike ridge  
strike-slip fault  
strike stream  
strike valley  
string  
strip mining  
stromatactis  
Stromatocystites walcotti  
stromatolite  
Stromatoporoidea  
Strombolian eruption  
strontianite  
strophic  
Strophomenida  
strophomenids  
structural contour map  
structural geology  
structural trap  
structure grumeleuse  
Strutt, John William

Sturtian  
sturzstrom  
Stylasterina  
stylolite  
stylolitization  
sub-  
Sub-arctic current  
subarkose  
Sub-Atlantic  
subbase  
Sub-Boreal  
sub-continental lithospheric mantle  
subcritical reflection  
subduction  
subduction earthquake cycle  
subduction zone  
subglacial  
subgrade  
subgroup  
subhedral  
sublimate  
sublimation  
sublitharenite  
sublittoral zone  
submarine canyon  
submersible  
sub-metallic  
sub-Plinian eruption  
subpolar glacier  
subsequent stream  
subsidence  
subsoiling  
subsolvus granite  
subsolvus syenite

substage  
subsurface flow  
subtidal  
subtractive primary colours  
subtropical high  
subtropical jet stream  
subzone  
sucrosic limestone  
Suess, Eduard  
Suess wriggles  
suevite  
suffusion  
Suisei  
sulci  
sulcus  
sulphates  
sulphides  
sulpho-salts  
sulphur, native  
sumatra  
Sun  
suncracks  
Sundaland  
Sundance Sea  
sunshine recorder  
Sun-synchronous orbit  
sun-tan age  
Suomi NPP  
super-  
super-adiabatic lapse rate  
supercell  
supercontinent  
supercooled cloud  
supercooling

supercritical fluid  
supercritical reflection  
supergene enrichment  
supergroup  
superimposed drainage  
superinterval  
Superior-type granular iron formation  
superposition  
supersaturation  
Supersaurus  
SuperView Earth Observation Constellation  
supplementary forms  
suppressed layer  
supra-  
supraglacial  
supralittoral zone  
supratidal  
surf  
surface inversion  
surface runoff  
surface tension  
Surface Water Ocean Topography mission  
surface wave  
surface wind  
surf wave  
surge  
Surtseyan eruption  
Surveyor  
survivorship curve  
susceptibility, magnetic  
susceptibility meter  
suspect terrane  
suspended load  
suspension feeder

sutural angle  
suture  
Svecofennian orogeny  
Sveconorwegian orogeny  
Svedberg unit  
swale  
swaley cross-bedding  
Swallow buoy  
swallow hole  
swallowtail twinning  
Swarm  
swash  
S-wave  
Swazian  
sweepstakes dispersal route  
swell  
swelling coefficient  
swirl  
SwissCube  
SWOT  
Sycorax  
syenite  
syenodiorite  
syenogabbro  
syenoids  
sylvite  
symbiosis  
symmetrical extinction  
symmetrical fold  
symmetrical trend  
Symmetrodonata  
symmetry plane  
sympatric evolution  
sympatry

Symphyla  
symplectic texture  
symplectite  
symplesiomorphy  
syn-  
synaeresis  
synaeresis cracks  
synapomorphy  
Synapsida  
synclinal ridge  
syncline  
synclinorium  
syneresis cracks  
synform  
syngenetic ore  
synkinematic  
synneusis  
synodic month  
synoptic meteorology  
synorogenic  
synrhabdosome  
synroc  
synrock  
symsacrum  
symsedimentary fault  
symsedimentary fold  
syntaxial growth  
syntectonic  
syntexis  
synthem  
synthetic-aperture radar  
synthetic fault  
synthetic seismogram  
synthetic thrust

syntype  
Syringopora fischeri  
system  
systematic errors  
systematic sampling  
Système Probatoire d'Observation de la Terre  
systems tract  
SZ  
T  
T  
Tabianian  
tabula  
tabulae  
tabular  
tabular cross-stratification  
Tabulata  
tabulation  
tachylite  
tachylyte  
tachytely  
Taconic orogeny  
taconite  
tadpole plot  
Tae Weian  
tafoni  
Taghanician  
tagma  
tagmosis  
tailings  
tailings dam  
Taitai  
talc  
talc schist  
taleolae

talik  
tail cloud  
talus  
taluvium  
Tame Valley  
Tandem-L  
tangential longitudinal strain  
TanSat  
tantalite  
taphichnia  
taphofacies  
taphonomic facies  
taphonomic grade  
taphonomy  
taphrogenesis  
TARANIS  
tarsal bone  
tar sand  
tarsometatarsus  
Taupo  
taxa  
taxodont  
taxon  
taxonomy  
taxon range zone  
Taylor, Frank Bursley  
Taylor number  
TCR  
t-d  
T-d curve  
T e  
tear fault  
technosols  
tectofacies

tectonic  
tectonism  
tectonite  
tectosilicates  
teeth  
tegeminal plates  
tegmen  
Teichichnus  
teilchron  
teilzone  
tektites  
teleconnections  
TeLEOS-1  
Teleostei  
teleseism  
Telesto  
telethermal  
Television and Infrared Observation Satellite  
telinite  
telluric anomaly  
telluric current  
telome theory  
telson  
Telychian  
TEM  
Temaikan  
Temispack  
Tempel–Tuttle  
temperate climate  
temperate glacier  
temperate zone  
temperature–composition diagram  
temperature distribution with depth  
temperature inversion

temperature log  
temperature range  
TEMPEST-D  
tempestite  
Templetonian  
Temple–Tuttle  
Temporal Experiment for Storms and Tropical Systems Technology –  
Demonstration  
tenacity  
tennantite  
tensile strength  
tensile stress  
tensiometer  
tension crack  
tension fracture  
tension gash  
tent rocks  
tepee  
tephigram  
tephra  
tephrite  
tephrochronology  
Teratan  
Terebratulida  
terebratulids  
terminal node  
terminator  
ternary system  
Terra  
terra  
terrace  
terracette  
terrae  
terrain

terrain component  
terrain correction  
terrain evaluation  
terrain pattern  
terrain unit  
terrane  
terra rossa  
terrestrial planet  
terrestrial radiation  
terric horizon  
terrigenous  
Tertiary  
tertiary creep  
teschenite  
tesla  
tessera  
tesserae  
test  
testate  
Tethyan realm  
Tethys  
Tethys Sea  
Tetracorallia  
tetracorals  
tetrad  
tetragonal  
tetrahedrite  
tetrahedron  
Tetrapoda  
Teurian  
textural maturity  
texture  
Thailand Earth Observation System  
Thalassa

Thalassinoidea  
thalweg  
thanatocoenosis  
Thanetian  
thaw  
Thebe  
theca  
Thecodontia  
thecodonts  
thematic map  
thematic mapper  
THEMIS  
theoretical morphology  
THEOS  
Therapsida  
thermal  
thermal cleaning  
thermal conductivity  
thermal emission  
thermal equator  
thermal inertia  
thermal infrared  
thermal low  
thermal metamorphism  
thermal resistivity  
thermal wind  
thermic  
thermistor  
thermochronology  
thermoclastic  
thermocline  
thermograph  
thermohaline circulation  
thermo-hygrograph

thermokarst  
thermoluminescence  
thermoluminescence dating  
thermonuclear reactions  
thermophile  
thermopile  
thermoremanent magnetization  
thermosphere  
Thermosphere Ionosphere Mesosphere Energetics and Dynamics  
Theropoda  
thickness  
thin section  
thixotropic mud  
thixotropy  
tholeiite  
tholi  
tholins  
tholoid  
tholus  
Thompson, Benjamin  
Thomson, Charles Wyville  
Thomson, William  
Thoracica  
thoracic vertebra  
thorax  
thorium–lead dating  
Thornthwaite, Charles Warren  
Thornthwaite climate classification  
three-cell model  
three-dimensional seismology  
threshold  
threshold-slope concept  
thrombolites  
throughfall

throughflow  
throw  
thrust  
thrust-sheet-top basin  
thufur  
thulite  
thundercloud  
thunderstorm  
Thuringian  
thuringite  
Thurnian  
Thvera  
Tibetan Plate  
Tibetan Plateau  
tibia  
tibiotarsus  
tidal barrage  
tidal bundle  
tidal correction  
tidal current  
tidal flat  
tidal friction  
tidal heating  
tidal inlet  
tidalite  
tidal range  
tidal rhythmite  
tidal stream  
tidal theory  
tide  
tieline  
tie point  
Tiffanian  
'tiger's eye'

tight gas  
tight sand  
Tiglian  
tile drain  
till  
till fabric analysis  
tillite  
tilloid  
till plain  
tilt-block tectonics  
tiltmeter  
time-averaged velocity  
TIMED  
time–distance curve  
time domain  
Time History of Events and Macroscale Interactions during Substorms  
time plane  
time-rock unit  
timescale  
time-stratigraphic unit  
tinguaite  
tinstone  
Tioughniogan  
TIROS  
Titan  
titanaugite  
Titania  
titanite  
titanohaematites  
titanohematites  
titanomagnetite  
Tithonian  
Titius–Bode law  
titrimetric analysis

tjaele  
TLE  
toad's eye tin  
Toarcian  
toeset  
Tolstojan  
tombolo  
Tommotian  
tomography  
tonalite  
Tonawandan  
Tonga–Kermadec Trench  
Tongaporutuan  
tonguing  
tonhäutchens  
Tonian  
tonstein  
Tool for the Analysis of RAdiations from lightNings and Sprites  
tool mark  
topaz  
TOPEX/Poseidon  
toplap  
topographic correction  
topology  
toposequence  
topotype  
topozone  
topple  
topset beds  
topsoil  
tor  
torbanite  
torbernite  
tornado

Tornquist Line  
toroidal field  
Torrejonian  
Torridonian  
torrid zone  
torsion  
torsion balance  
torta  
Tortonian  
torus  
toscanite  
total core recovery  
total intensity  
total internal reflection  
total range zone  
total stress  
tourmaline  
tourmalinization  
Tournaisian  
tourquoise  
Toutatis  
tower karst  
Toyonian  
T-peg  
trace  
trace element  
trace-element fractionation  
trace fossil  
tracer  
Tracheophyta  
trachyandesite  
trachybasalt  
trachyte  
trachytic texture

trachytoidal  
track  
traction carpet  
traction load  
trade-wind inversion  
trade winds  
Traditional Stratigraphic Scale  
trail  
trailing edge  
training area  
tramontana  
tranquil flow  
transcurrent fault  
transfer fault  
transfluence  
transformation twinning  
transform fault  
transgression  
transgressive systems tract  
transient creep  
transient electromagnetic method  
transient flow  
transient luminous event  
transient variation  
transit time  
translation  
translational slide  
translational slump  
translation gliding  
translocation  
translucent  
translucidus  
transmission coefficient  
transmission electron microscope

transmissivity  
transmittance  
trans-orogen basin  
transparency  
transparent  
transpiration  
transpolar drift  
transpression  
trans-Saharan seaway  
transtension  
transverse dune  
transverse-type coast  
transverse wave  
trap  
trap-door caldera  
trapezohedron  
travel time  
travel-time curve  
traverse  
travertine  
tree-ring analysis  
trellis drainage pattern  
trema  
Tremadocian  
tremata  
tremolite  
Trempealeauan  
trench  
trend  
Treptichnus pedum  
trevorite  
triad  
Triana  
triangle zone

triangular facet  
Triassic  
triaxial cell  
triaxial compression test  
triaxial ellipsoid  
tributary  
triclinic  
tricolpate sulci  
Triconodonta  
tridymite  
trigonal  
Trilete  
Trilobita  
trilobite  
trilobite eye  
Trimerophytina  
trim line  
trimorphism  
Trinculo  
triple core barrel  
triple junction  
triple-junction method  
triple point  
triserial  
tritium clock  
Triton  
TRMM  
trochiform  
trochoid  
trochospiral  
troctolite  
troilite  
troll  
trona

trondhemite  
tropical air  
tropical arid morphoclimatic zone  
tropical cyclone  
Tropical Cyclone Programme  
Tropical Rainfall Measuring Mission  
tropical semi-arid morphoclimatic zone  
tropical wet–dry morphoclimatic zone  
tropopause  
troposphere  
trough  
trough cross-stratification  
trowal  
true age  
true dip  
true thickness  
truncated spur  
Ts  
tschermakite  
TSS  
TST  
tsunami  
tsunamites  
t-test  
tuba  
tube feet  
tubercle  
tubular fenestrae  
tufa  
tuff  
tumulus  
tundra  
Tundra Soil  
tungstates

tunicates  
tunnel trend  
tunnel valley  
TURAM method  
turbidite  
turbidity current  
turbidity flow  
turbinate  
turbulence  
turbulent flow  
turnover rate  
turnover time  
Turonian  
turquoise  
turreted  
turriculate  
tuya  
Twenhofel, William Henry  
twilight  
twin axis  
twin gliding  
twinkling  
twin law  
twin plane  
TWINS  
two-way travel time  
Two Wide-angle Imaging Neutral-atom Spectrometers  
t-x curve  
t<sup>2</sup>-x<sup>2</sup> graph  
Tycho  
tympanic bone  
Type I earthquake source  
Type II earthquake source  
type area

type locality  
type section  
type series  
type specimen  
typhoon  
typological method  
Tyrannosaurus rex  
Tyrrhenian  
Ubendian orogeny  
Udden–Wentworth scale  
Udocanian  
Ufimian  
Uivakian orogeny  
UK-DMC-2  
Ulatisian  
Ulcanian  
ulexite  
ulna  
Ulsterian  
ultimate bearing capacity  
ultimate strength  
ultisols  
ultrabasic rock  
ultramafic  
ultramylonite  
ultraplinian eruption  
ultraviolet radiation  
ultraviolet spectrometer  
ultraviolet spectrophotometer  
ultraviolet–visual spectrophotometry  
Ulysses  
umber  
umbilicus  
umbo

umbones  
umbric epipedon  
umbric horizon  
Umbriel  
umbrisols  
unavailable water  
uncinus  
unconfined aquifer  
unconfined compressive strength  
unconformity  
unconformity trap  
unconformity-type uranium ore  
unconsolidated  
undae  
undaform  
undathem  
undepleted mantle  
undercliff  
undercooling  
underfit stream  
underflow  
underplating  
undersaturated  
underthrusting  
undertow  
Undillian  
undrained test  
undulatus  
undulose extinction  
ungulate  
uniaxial compression test  
uniaxial compressive strength  
uniaxial interference figure  
Unibothriocidaris

unicarinate  
Unified Stratigraphic Timescale  
UNIFORM-1  
uniform flow  
uniformitarianism  
unilocular  
uniramous  
uniserial  
unit cell  
United Kingdom – Disaster Monitoring Constellation-2  
United States Department of Agriculture  
United States Geological Survey  
unit form  
unit hydrograph  
unit-stratotype  
unit stress  
univariant assemblage  
universal soil loss equation  
University International Formation Mission-1  
unloading joint  
unsaturated zone  
unsaturation index  
unsorted biosparite  
unsteady flow  
uphole survey  
uphole time  
upright fold  
upslope fog  
upthrow  
upthrust  
Upton Warren  
upward continuation  
upward lightning  
upwelling

Ur  
Uralian orogeny  
uralite  
uralitization  
Ural Sea  
uranian satellites  
uraninite  
uranium deposit  
uranium–lead dating  
uranium–protactinium dating  
uranium series  
Uranus  
urban climate  
urban dome  
urban heat island  
Urey, Harold Clayton  
Ureconian  
Urochordata  
Ursidae  
urtite  
Ururoan  
Urutawan  
USDA  
UTS  
UV  
uvala  
uvarovite  
UVS  
UV–Vis spectrophotometry  
[UVW]  
V  
Vaalbara  
Vaalian  
vadose zone

Valanginian  
Valdayan/Zyryanka  
Valdonnian  
valency  
valley bog  
valley bulging  
valley glacier  
valley-side bench  
valley train  
valley wind  
vallis  
valve  
vanadinite  
Van Allen belts  
Van Allen Probes  
van der Waals force  
vane  
vane test  
van't Hoff, Jacobus Henricus  
vapour-phase crystallization  
vapour pressure  
vapour-pressure curve  
VAR  
Varanger  
Varangian  
vardar  
variable-area display  
variance  
variolitic  
varve  
varve analysis  
varve chronology  
varve count  
vascular plants

vastitas  
vastitates  
veering  
Vega  
Vegetation and Environment monitoring on a New MicroSatellite  
vegetation index  
vein  
vein deposit  
velocity-depth distribution  
velocity inversion  
velocity log  
velocity profile  
velocity survey  
VELOX-CI  
VELOX-1  
velum  
vendavale  
Vendian  
Vendobionta  
Venera  
Venezuelan Remote Sensing Satellite-1  
Venice system  
Vening-Meinesz, Felix Andries  
VEN $\mu$ S  
vent breccia  
vent conglomerate  
venter  
Ventersdorp  
ventifact  
ventral  
Venturian  
Venus  
venusian  
Venus snow

veranillo  
verano  
Vereiskian  
vergence  
vermiculite  
Verrucano  
vertebra  
vertebral column  
Vertebrata  
vertebrates  
vertebratus  
vertical component  
vertical electrical sounding  
vertical fold  
vertical seismic profile  
vertical stacking  
vertic horizon  
vertisols  
very-long-baseline interferometry  
very-low-frequency method  
very-near infrared  
VES  
vesicle  
vesicular basalt  
Vesta  
Vesuvian eruption  
vesuvianite  
VGP  
VHN  
vibration, ground  
vibration direction  
Vibroseis  
vicariance  
Vickers hardness number

Victoriapithecus  
vidicon  
Vietnam Natural Resources, Environment and Disaster-monitoring  
Satellite-1A  
Viking  
Villafranchian  
virga  
virgella  
Virgellina  
Virgilian  
virtual geomagnetic pole  
viscoelastic behaviour  
viscosity  
viscous remanent magnetism  
Visean  
vishnevite  
visible radiation  
visual binary  
vitrage  
vitrain  
vitreous  
vitric horizon  
vitrinite  
vitroclastic  
vitrophyric  
Viverridae  
vivianite  
VLBI  
VLF method  
VNREDSat-1  
vog  
vogesite  
Vogt, Johan Hermann Lie  
void

void ratio  
Voigt model  
volatile  
volcanic-arc zone  
volcanic bomb  
volcanic centres  
volcanic conduit  
volcanic cone  
Volcanic Constructs  
volcanic dome  
volcanic dust  
volcanic-exhalative processes  
Volcanic Explosivity Index  
volcanic haze  
volcanicity  
volcaniclastic  
volcanism  
volcanic neck  
volcanic pile  
Volcanic Plains  
volcanic plug  
Volcanic units  
volcano  
volcano-tectonic depression  
Volgian  
volume diameter  
volume diffusion  
volumenometer  
volume scattering  
von Karman–Prandtl equation  
vorticity  
Voyager  
VRM  
VRSS-1

VSP  
vug  
vuggy porosity  
vugh  
Vulcanian eruption  
vulcanicity  
vulcanism  
Waalian  
wacke  
wackestone  
wad  
Wadati-Benioff zone  
wadi  
Waiauan  
Waiaun  
Waipawan  
Waipipian  
Waitakian  
walkaway vertical seismic profile  
Walker's steelyard  
Wallace's line  
wall cloud  
wall, law of the  
wall rock  
wall-rock alteration  
Walther's law  
Waltonian  
Wangerripian  
waning slope  
Warendian  
Warepan  
warm front  
warm glacier  
warm rain

warm sector  
Wasatchian  
washboard moraine  
washover delta  
washover fan  
washplain  
water  
water-absorption test  
water balance  
water budget  
water budget, global  
water-drive reservoir  
water gun  
water inventory  
water potential  
watershed  
waterspout  
water storage  
water-table  
water vapour  
water vascular system  
water velocity  
water-witching  
Waucoban  
wave  
wave base  
wave-built terrace  
wave clouds  
wave-cut bench  
wave-cut platform  
wave depression  
wave diffraction  
wave equation  
wave-front

wavelength  
wavellite  
wavenumber  
wave period  
wave refraction  
wave-ripple cross-lamination  
wave ripple mark  
wave spectrum  
wave velocity  
wavy bedding  
waxing slope  
waxy  
weakening  
Wealden  
Wealdien  
wearing course  
weathering  
weathering correction  
weathering front  
weathering index  
weathering layer  
weathering microindices  
weathering pit  
weathering profile  
weathering series  
weathering zone  
Weathernews Inc. Satellite-1  
weather report  
weather satellite  
wedge  
wedge-edge trap  
wedge tornado  
Wegener, Alfred  
Weichselian

weight drop  
weighted average  
weir  
Weissliegende  
Weiss zone law  
welded ignimbrite  
welded tuff  
well  
well injection method  
well logging  
well-point drainage  
well screen  
well shooting  
Weltian  
Wenlock  
Wenner electrode array  
Wentworth scale  
Werner, Abraham Gottlob  
Werrickooian  
West Australia current  
westerlies  
Western Boundary Undercurrent  
western intensification  
Western Pacific Satellite  
West Greenland current  
West-Kohoutek-Ikemura  
WESTPAC  
Westphalian  
westward drift  
West Wind Drift  
wet-bulb depression  
wet-bulb thermometer  
'wet chemistry'  
wet mass

wet melt  
wetted perimeter  
W-fold  
Whaingaroan  
Wheelerian  
Whewell, William  
whinstone  
whirling psychrometer  
whirlwind  
white-out  
Whiterockian  
'white smoker'  
whittings  
Whitwellian  
whole Earth composition  
whole-mantle convection  
whole-rock dating  
whorl  
Widmanstätten structure  
Wiechert, Emil  
Wiener filter  
Wien's displacement law  
wiggle trace  
wigwams  
Wild 2  
wildflysch  
willemite  
Wilson, John Tuzo  
Wilson cycle  
wilting coefficient  
wilting point  
WIND  
Windermere Interstadial  
wind noise

window  
wind rose  
windrow  
wind shear  
windward  
wireline  
wireline logging  
Wirtanen  
Wisconsinian  
witherite  
within-plate basalt  
Witwatersrand  
WMO  
WNISAT-1  
Wolfcampian  
wolframite  
wollastonite  
Wolstonian  
Wonokan  
wood tin  
Woodward, John  
woolsack  
Wordian  
work hardening  
World Climate Programme  
World Meteorological Organization  
World Reference Base for Soil Resources  
WorldView  
World Weather Watch  
WorldWide Standard Seismograph Network  
WPB  
WRB  
wrench fault  
wrinkle ridge

Wuchiapingian  
wulfenite  
Wulff stereographic net  
Würm  
WWSSN  
WWW  
xanthophyllite  
Xenian  
xeno-  
xenoblastic  
xenocryst  
xenolith  
xenotime  
xenotopic fabric  
XPL  
xpols  
X-ray diffraction crystallography  
X-ray fluorescence  
X-ray fluorescence spectrometry  
X-ray photography  
X-ray powder photograph  
X-ray spectrometer  
XRF  
yardang  
Yarmouthian  
Yatalan  
yazoo stream  
Yeadonian  
yedoma permafrost  
yellowcake  
yield–depression curve  
yield point  
yield stress  
Ynezian

Younger Dryas  
younging  
young mountain belt crustal type  
Young's modulus  
Yovian  
Ypeenian  
Ypresian  
Yurmatian  
Z  
Zanclean  
'zap pits'  
zastrugi  
Zechstein  
Zechstein Sea  
Zedian  
Zelzate  
Zemorrian  
zeolite facies  
zeolites  
zephyr  
zero-length spring  
zeta-form beach  
zeugen  
Z-fold  
Zhangheng 1  
zibar  
zig-zag fold  
zinc blende  
zincite  
Zingg diagram  
zinnwaldite  
zircon  
ZiYuan  
Zoantharia

zodiacal light  
Zoeppritz's equations  
zoisite  
zonal  
zonal flow  
zonal index  
zonal scheme  
zonation  
Zond  
zone  
zone axis  
zone fossil  
zone of aeration  
zone of saturation  
zone refining  
zone symbol  
zonule  
zoogeographical region  
zoogeomorphology  
zoophycus  
Zosterophyllum  
Zosterophyllophytina  
ZTR index  
Zungxian  
Zuzhuangian  
zygapophyses

# A Dictionary of Geology and Earth Sciences

**Michael Allaby** has written many books on environmental science and especially on climatology and meteorology. These include *Encyclopedia of Weather and Climate*; *The Facts On File Weather and Climate Handbook*; and the *DK Guide to Weather*. He is the General Editor of the Oxford Dictionaries of *Ecology, Zoology, and Plant Sciences*.



<http://www.oxfordreference.com/page/geol>

For recommended web links for this title, visit  
[www.oxfordreference.com/page/geol](http://www.oxfordreference.com/page/geol) when you see this sign.

A Dictionary of  
**Geology and Earth Sciences**

FIFTH EDITION

*Edited by* MICHAEL ALLABY

**OXFORD**  
UNIVERSITY PRESS

# OXFORD

UNIVERSITY PRESS

Great Clarendon Street, Oxford, OX2 6DP, United Kingdom

Oxford University Press is a department of the University of Oxford. It furthers the University's objective of excellence in research, scholarship, and education by publishing worldwide. Oxford is a registered trade mark of Oxford University Press in the UK and in certain other countries

Text © Oxford University Press 1990, 1999, 2003, 2008, 2013, 2020 © in the compilation and editorial matter Michael Allaby 1990, 1999, 2003, 2008, 2013, 2020

The moral rights of the author have been asserted

First published as *A Dictionary of Earth Sciences* 1990

First issued as an Oxford University Press paperback 1991

Second edition 1999

Reissued with new covers and corrections 2003

Third edition 2008

Fourth edition published as *A Dictionary of Geology and Earth Sciences* 2013

Fifth edition 2020

Impression: 1

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, without the prior permission in writing of Oxford University Press, or as expressly permitted by law, by licence or under terms agreed with the appropriate reprographics rights organization. Enquiries concerning reproduction outside the scope of the above should be sent to the Rights Department, Oxford University Press, at the address above

You must not circulate this work in any other form and you must impose this same condition on any acquirer

Published in the United States of America by Oxford University Press  
198 Madison Avenue, New York, NY 10016, United States of America

British Library Cataloguing in Publication Data

Data available

Library of Congress Control Number: 2019952629

ISBN 978-0-19-883903-3

ebook ISBN 978-0-19-257570-8

Printed and bound in Great Britain by Clays Ltd, Elcograf S.p.A.

Links to third party websites are provided by Oxford in good faith and for information only. Oxford disclaims any responsibility for the materials contained in any third party website referenced in this work.

---

## Preface to the Fourth Edition

In preparing this new edition of our *Dictionary of Geology and Earth Sciences* I have changed the emphasis very slightly by increasing the number of entries for terms that are directly geological. The new title reflects this minor revision. At the same time, however, I have found it necessary to add more entries describing satellite missions and planetary discoveries. I have also added a new Appendix, listing web addresses for all the satellite missions described in dictionary entries.

Inevitably, the dictionary has grown a little longer. That is the way of dictionaries. New terms emerge, but old ones take a long time to die and disappear from the vocabulary.

As with the third edition, I have worked alone. I have scrutinized every entry and made many small changes where I thought I could improve the clarity of definitions. If, inadvertently, I have introduced any errors, or failed to spot and remove earlier errors, I have no one to blame but myself. If you find a mistake I will be grateful if you draw it to my attention. You can contact me through my website.

If you are meeting the dictionary for the first time, I hope you find it helpful.

MICHAEL ALLABY

TIGHNABRUAICH, ARGYLL



[www.michaelallaby.com](http://www.michaelallaby.com)

---

## Preface to the Fifth Edition

Over the decades since the first edition of our dictionary was published, satellite observations of the Earth's surface and atmosphere have grown increasingly important and increasingly valuable to geologists and Earth scientists. There are now so many satellites in orbit above us that it would be impossible to describe them all in a volume of this size. Nevertheless, I have done my best by adding a substantial number of new satellite entries, and in many cases links to web sites, describing those that seem to me the most relevant and interesting.

I have also updated many of the entries for stages. Again, it would be impractical to include all the stage names from every part of the world, but I have added quite a number.

I hope you find the dictionary useful and its entries clear.

MICHAEL ALLABY

TIGNABRUAICH, ARGYLL



[www.michaelallaby.com](http://www.michaelallaby.com)

---

# Contents

How to search for terms

Contributors and Advisers

Alphabetical List of Entries

## **A Dictionary of Geology and Earth Sciences**

A B C D E F G H I J K L M N O P Q R S T U  
V W X Y Z

### Appendices

Appendix A: Stratigraphic Units

Appendix B: Timescales

Appendix C: Wind Strength

Appendix D: SI Units, Conversions, and Multiples

Appendix E: Volcanic Explosivity Index (VEI)

Appendix F: Torino Impact Hazard Scale

Appendix G: Avalanche Classes

Appendix H: List of Useful Websites

Appendix I: Satellite Missions

---

## Contributors and Advisers

**Ailsa Allaby**

**Michael Allaby**

**Robin Allaby**, University of Manchester Institute of Science and Technology

**Dr Keith Atkinson**, Camborne School of Mines

**Dr R. L. Atkinson**, Camborne School of Mines

**Dr T. C. Atkinson**, University of East Anglia

**Dr A. V. Bromley**, Camborne School of Mines

**Denise Crook**

**J. G. Cruickshank**, Department of Agriculture for Northern Ireland, Belfast

**Dr P. Francis**, Open University; Lunar and Planetary Institute, Houston

**Professor K. J. Gregory**, Goldsmith's College, University of London

**Dr C. D. Gribble**, University of Glasgow

**Dr Colin Groves**, \*Australian National University

**Dr W. J. R. Harries**, University of Plymouth

**Professor M. Hart**, University of Plymouth

**Professor Emeritus H. H. Lamb**, \*University of East Anglia

**John Macadam**

**Dr R. J. T. Moody**, Kingston University

**Dr J. Penn**, Kingston University

**Dr John M. Reynolds**, Reynolds Geo-Science Ltd.

**Dr D. Rolls**, Kingston University

**Dr I. Roxburgh**

**Dr N. A. Rupke**, Wolfson College, Oxford

**Dr Stuart Scott**, University of Plymouth

**Dr B. W. Sellwood**, University of Reading

**Dr P. J. C. Sutcliffe**, Kingston University

**Professor D. H. Tarling**, University of Plymouth

**Joan Taylor**

**Professor S. R. Taylor**, Australian National University

**Dr R. J. Towse**, \*Kingston University

**Dr I. Tunbridge**, University of Plymouth

**Dr C. E. Vincent**, University of East Anglia

**Professor Brian F. Windley**, University of Leicester

**Andrew Yelland**, Birkbeck College, London

\* Contributor to *The Oxford Dictionary of Natural History* whose earlier entries have been transferred to this book.

# A

**aa** See LAVA.

**AABW** See ANTARCTIC BOTTOM WATER.

**AAC** See ANTARCTIC CONVERGENCE.

**Aalenian** A \*stage in the European Middle \*Jurassic (175.6–171.6 Ma ago, Int. Commission on Stratigraphy, 2004). See also DOGGER.

**Aalto-1** A student project at the University of Aalto, Finland, in which students built an Earth-observation \*nanosatellite based on \*CubeSat standards, launched on 23 June 2017, from the Satish Dhawan Space Centre, India.

**AAV** See AGGREGATE TESTS.

**Ab** See ALKALI FELDSPAR.

**abandoned channel** A former stream channel through which water no longer flows (e.g. a \*cut-off).

**abandonment facies association** A \*facies association formed under conditions of rising sea level, when \*clastic deposition has ceased and sediment is deposited very slowly.

**abapical** A directional term meaning away from the shell \*apex.

**abaptation** The process by which an organism is fitted to its environment as a consequence of the characters it inherits, which have been filtered by \*natural selection in previous environments. Because present environments seldom differ greatly from recent past environments, adaptive fitness can resemble \*adaptation. In this sense, however, adaptation appears to imply advance planning, or design, which is misleading.

**Abbé refractometer** See REFRACTOMETER.

**abiogenesis** Development of living organisms from non-living matter; as in the supposed origin of life on Earth, or in the concept of spontaneous generation, which was once held to account for the origin of life but which modern understanding of evolutionary processes (see EVOLUTION) has rendered outdated.

**abiotic** Non-living; devoid of life. Compare BIOTIC.

**ablation** 1. Removal of snow and ice by melting and by direct alteration from the solid to the gaseous phase (sublimation). The rate of loss is controlled chiefly by air temperature, wind velocity, \*humidity, rainfall, and \*solar radiation. Ablation on snowfields is also influenced by aspect, depth of snow, and the nature of the underlying surface. Ablation \*till is the glacial debris that may be released. The ablation zone of a glacier is that area in which losses, including \*calving, exceed additions. 2. Removal of \*rock material, especially by wind action.

**ablation till** See ABLATION 1; TILL.

**ablation zone** See ABLATION.

**aboral** Away from the mouth; on the opposite side of the body from the mouth.

**abrasion (corrasion)** The erosive (see EROSION) action that occurs when \*rock particles of varying size are dragged over or hurled against a surface. Some common agents of abrasion are the \*bed load of streams, rock debris embedded in the bases of \*glaciers, and \*sand and \*shingle transported by wind or waves.

**abrasion ramp** A gentle, seaward slope, with a gradient of approximately 1°, in an intertidal \*shore platform that is caused by wave \*abrasion. The removal of material by wave action leaves the base of the cliff exposed, leading to further cliff retreat. Abrasion ramps usually terminate on the seaward side where the sea depth reaches about 10 m.

**absarokite** An \*alkaline \*basalt that contains \*olivine, \*augite, \*labradorite, and sanidine (see ALKALI FELDSPARS) in approximately equal proportions, together with \*accessory minerals \*biotite, \*apatite, and

opaque oxides (see [OPAQUE MINERALS](#)). Absarokite is abundant in the Absaroka Range of mountains in Yellowstone National Park, USA.

**absolute age (true age)** The age of a geologic phenomenon measured in present Earth years, rather than its age relative to other geologic phenomena (compare [RELATIVE AGE](#)). The term 'absolute age' has been considered rather misleading, as the means for measuring ages (*\*radiometric dating*, *\*dendrochronology*, *\*varve analysis*) are subject to experimental error and the dates obtained are not precise. The alternative term 'apparent age' has been suggested. See also [DATING METHODS](#); [GEOCHRONOLOGY](#).

**absolute humidity** See [HUMIDITY](#).

**absolute plate motion** The motion of a lithospheric *\*plate* (see [LITHOSPHERE](#)) with respect to a fixed frame of reference. Various frames of reference have been used, including those defined by *\*hot spots*, no net torque of all the plates, and palaeomagnetic (see [PALAEOMAGNETISM](#)) Euler poles (see [POLE OF ROTATION](#)).

**absolute pollen frequency (APF)** *\*Pollen* data from sediments, expressed in terms of the absolute numbers for each *\*species*, *\*genus*, or *\*family*, per unit volume of sediment and, where deposition rates are known, per unit time. In certain circumstances this approach gives clearer information than does the traditional way of expressing pollen data as *\*relative pollen frequencies* (RPFs). APFs are particularly useful in site comparisons in which one or more high pollen producers vary. For example, when trees first appear in the regional pollen rain their prolific pollen may, in an RPF method, give the impression of declining herbaceous species, whereas examination by an APF method will show constant values for herb species.

**absolute porosity** See [POROSITY](#).

**absolute temperature** Temperature measured using the *\*Kelvin scale*.

**absolute vorticity** See [VORTICITY](#).

**absolute zero** See [KELVIN SCALE](#).

**absorptance** The ability of a material to absorb *\*electromagnetic radiation* of a specified wavelength. See also [ABSORPTANCE BAND](#).

**absorptance band** The range of wavelengths of *\*electromagnetic radiation* which are absorbed by a material. *See also* ABSORPTANCE.

**absorption** 1. A chemical or physical process in which an amount of one substance enters a solid or liquid bulk phase of another; the substance absorbed is taken up by the volume of the absorbing substance. *Compare* ADSORPTION; *see also* SORPTION. 2. The amount of seismic energy lost during transmission, by conversion to heat. The absorption coefficient is the fractional loss of energy over a distance of one *\*wavelength*; hence higher-*\*frequency* signals are attenuated more readily than those of lower frequencies over the same path. Typical values for *\*rocks* range from 0.25 to 0.75 dB per wavelength.

**abstraction (extraction)** The artificial removal of water from a well, *\*reservoir*, or river.

**Abukama-type metamorphism** The *\*recrystallization* of *\*rocks* under a high *\*geothermal gradient* so that at any given temperature the pressure is relatively low. The term originally referred to a belt of *\*metamorphic rocks* stretching south-westwards from the Abukama Plateau in Japan, and characterized by the development of *\*andalusite* and *\*sillimanite* in rocks that were originally *\*shales* (*\*pelites*). This belt lies parallel to, and on the continental side of, a high-pressure metamorphic belt.

**abundance zone** *See* ACME ZONE.

**ABW** *See* ARCTIC BOTTOM WATER.

**abyssal hills** Relatively small topographic features of a dominantly flat, deep-ocean floor, commonly 50–250 m in height and a few kilometres in width. They are most typical of the *\*Pacific Ocean* floor at depths of 3000–6000 m.

**abyssal plain** Smooth, almost level area of the deep-ocean floor in which the gradient is likely to be as low as 1:10 000. The covering sediments are usually thin deposits of a *\*pelagic ooze* or *\*distal \*turbidite*.

**abyssal storm (benthic storm)** A large pulse of energy, possibly transferred from the surface, that accelerates *\*contour currents* on the ocean floor to about 40 cm/s, raising large amounts of fine sediment.

**abyssal zone** Zone of greatest ocean depth, i.e. below a depth of 2000 m. This zone lies seaward of, and deeper than, the **\*bathyal zone**, and covers approximately 75% of the total ocean floor. It is the most extensive Earth environment, cold, dark, with slow-moving currents (less than a few centimetres per second), supporting **\*fauna** that typically are black or grey, delicately structured, and not streamlined.

**Acadian orogeny** A phase of mountain building affecting an area from the northern Appalachians in what is now New York State to the Bay of Fundy in maritime Canada (the name refers to the colony of Acadie in that region of French Canada). It occurred in the **\*Devonian** about 390 Ma ago, although the precise date and duration are uncertain, and was most intense east of the Taconic area (see **TACONIC OROGENY**). It was caused by the westward movement of the Avalon **\*terrane**. See **APPALACHIAN OROGENIC BELT**.

**Acado-Baltic Province** See **ATLANTIC PROVINCE**.

**acanthite** An **\*opaque mineral**,  $\text{Ag}_2\text{S}$ ; density 7.2–7.4 g/cm<sup>3</sup>; **\*hardness** 2–2.5; **\*monoclinic**; lead grey or black; lead grey **\*streak**; metallic **\*lustre**; crystals pseudo-cubic, pseudo-octahedral, or monoclinic; no **\*cleavage**; fracture sub-conchoidal; widely distributed in silver deposits and zones of **\*secondary enrichment**. It is the low-temperature modification of silver sulphide, and all natural silver sulphide at room temperature is acanthite.

**acanthodians** See **ACANTHODII**.

**Acanthodii (acanthodians, spiny sharks)** Class of primitive, fossil fish, characterized by the presence of a true bony skeleton (see **BONE**), a **\*heterocercal tail fin**, a persistent **\*notochord**, **\*ganoid scales**, and stout spines in front of the fins. The acanthodians lived from the **\*Silurian** to the **\*Permian** period and may be related to ancestors of the more modern bony fish.

**Acanthograptidae** See **DENDROIDEA**.

**Acanthostega** See **ICHTHYOSTEGA**.

**acceleration** 1. An increase in speed or velocity. 2. **\*Evolution** that occurs by increasing the rate of ontogenetic (see **ONTOGENY**) development, so that

further stages can be added before growth is completed. This form of **\*heterochrony** was proposed by E. H. Haeckel as one of the principal modes of evolution.

**acceleration, gravitational** See GRAVITATIONAL ACCELERATION.

**accelerograph (earthquake seismometer)** An instrument used to measure **\*earthquake** movements that are too strong for more sensitive **\*seismometers** to register accurately. An accelerograph contains three **\*accelerometer** heads aligned to measure movement in three directions. The accelerograph is often connected directly to the Internet.

**accelerometer** A device whose output is directly proportional to acceleration. Accelerometers are used in the measurement of the motion of a ship, helicopter, or aircraft during **\*gravity surveys**. A **\*seismometer** or moving-coil **\*geophone** can also function as an accelerometer.

**accessory, lithic** See LITHIC FRAGMENT.

**accessory cloud** A small cloud that is seen to be associated with a much larger cloud belonging to one of the ten cloud genera (see CLOUD CLASSIFICATION). **\*Pileus**, **\*tuba**, and **\*velum** are accessory clouds.

**accessory mineral** A **\*mineral** **\*phase** within a rock whose presence does not affect the root name of the rock. For instance, the root name 'granite' is defined by the presence of **\*quartz**, **\*alkali feldspar**, and **\*mica**. These are the '**\*essential minerals**'. The presence of the mineral **\*sphene** does not affect the root name and hence would be an example of an accessory mineral. **\*Apatite** and **\*zircon** are also common accessory minerals.

**accessory plate (sensitive tint)** In optical microscopy, a plate used to determine the optical properties of **\*minerals**. **\*Quartz**, **\*mica**, and **\*gypsum** are the common minerals used to determine the slow and fast **\*vibration** directions that relate to the two **\*refractive** indices of an **\*anisotropic** mineral. The terms 'length-fast' and 'length-slow' may then be assigned to a given mineral for identification purposes. A wedge of quartz (quartz wedge) is used to determine the order of **\*interference colour** exhibited by a mineral.

**accidental lithic** See LITHIC FRAGMENT.

**accommodation space** The space in which sediment may accumulate.

**accommodation zone** A region of intermeshed *\*normal faults*, with very complex geometry, that lies between the boundary faults of a series of interlinked half-grabens (see *GRABEN*) in an area of crustal extension.

**accordion fold** See *CHEVRON FOLD*.

**accretion** **1.** Process by which an inorganic body grows in size by the addition of new particles to its exterior. It is the mechanism by which primitive planetary bodies are believed to form as a result of the accumulation of minute, cold, homogeneous particles (homogeneous accretion). An alternative hypothesis is that iron-rich cores accumulated first and were later surrounded by silicate material (heterogeneous accretion). Homogeneous accretion yields a planet that initially has the same composition from centre to surface; heterogeneous accretion yields a planet that has a layered structure from the start. **2.** The accumulation of sediments from any cause, representing an excess of deposition over *\*erosion*. **3.** The addition of continental material to a pre-existing continent, usually at its edge. The use of ‘accretion’ in this sense has evolved from theories of *\*nucleation* to newer theories of the horizontal addition of *\*allochthonous \*terrane*s of initially coherent bodies of continental *\*rock*, usually more than 100 km<sup>2</sup> in area, which can collide, rotate, and fragment as they become sutured to a continent.

**accretionary heating** The heating of bodies orbiting a star due to bombardment by smaller objects, the kinetic energy of the impacting body ( $\frac{1}{2}mv^2$ , where  $m$  is mass and  $v$  velocity) being released mainly as heat.


**accretionary basin** A small basin, much smaller than a *\*fore-arc basin*, that develops on the top of an *\*accretionary wedge* and fills with *\*debris-flow* material, *\*turbidites*, and volcanic rocks.

**accretionary lapilli** Pellets of *\*ash*, ranging in size from 2 mm to 64 mm, which commonly exhibit a concentric (‘onion skin’) internal structure. The *\*lapilli* are formed by the accretion of very fine ash around condensing water droplets or solid particles, particularly in steam-rich eruptive columns (see *ERUPTION*). Once formed they can be transported and deposited by *\*pyroclastic* fall, *\*surge*, or flow processes.

**accretionary levée** See LAVA LEVÉE.

**accretionary prism** See ACCRETIONARY WEDGE.

**accretionary wedge (accretionary prism)** A tectonically thickened wedge of *\*sediment* found on the landward side of some *\*trenches*. The accretionary wedge consists of oceanic sediment scraped off the subducting *\*plate* (see SUBDUCTION), plus sediment derived from landward and deposited in the trench. Slices of sediment are added to the wedge by *\*underthrusting* and the trench migrates seaward, the continuation of this process producing an *\*inversion*.



### Accretionary wedge

**accumulated temperature** Surplus or deficit of temperature with respect to a defined mean value and expressed as an accumulation over a given period, e.g. a month, season, or year. For example, a datum value of 6 °C is used as a critical temperature for sustained vegetation growth, against which accumulated surpluses or deficits may be measured.

**accumulation zone** That part of a *\*glacier* where the mean annual gain of *\*ice*, *\*firn*, and snow is greater than the mean annual loss. The zone consists of stratified firn and snow together with ice from frozen meltwater. Its lower boundary is the *\*equilibrium line*.

**ACD** See ARAGONITE COMPENSATION DEPTH.

**ACF** See ACF DIAGRAM; AUTOCORRELATION.

**ACF diagram** A three-component, triangular graph used to show how metamorphic *\*mineral* assemblages vary as a function of *\*rock* composition within one *\*metamorphic facies*. Besides SiO<sub>2</sub>, the five most abundant oxides found in *\*metamorphic rocks* are Al<sub>2</sub>O<sub>3</sub>, CaO, FeO, MgO, and K<sub>2</sub>O. The three components plotted on ACF diagrams are A (Al<sub>2</sub>O<sub>3</sub>), C (CaO), and F (FeO + MgO), making the diagrams particularly useful for showing assemblage variations in metamorphosed, *\*basic*, *\*igneous* rocks, and impure *\*limestones*. However, each of these components has to be modified slightly to account for the presence of other, minor components in the rock. Such modification leads to: A (Al<sub>2</sub>O<sub>3</sub> - Na<sub>2</sub>O - K<sub>2</sub>O); C (CaO - [(10/3)P<sub>2</sub>O<sub>5</sub>] - CO<sub>2</sub>); and F (FeO + MgO - Fe<sub>2</sub>O<sub>3</sub> - TiO<sub>2</sub>). The minerals *\*quartz* and *\*albite* are assumed to be present in the rocks and are not shown on the diagram. *\*Tielines* connect minerals which coexist in equilibrium and can thus define triangular areas in which three minerals are in equilibrium in the rock, lines on which two minerals are in equilibrium in the rock, and points at which one mineral is in equilibrium in the rock (in addition to the ubiquitous quartz and albite). See *AFM DIAGRAM*. See also *A'KF DIAGRAM*.

**achnelith** See *PELÉ'S HAIR*.

**achondrite** Rare stony *\*meteorite* lacking *\*chondrules* and with low nickel-iron content. It is more coarsely crystalline than a *\*chondrite*. Basaltic achondrites resemble terrestrial *\*lavas*.

**achromatic line** In the three-dimensional graph which plots quantities of the three *\*additive* primary colours contributing to *\*pixels* against each other, the line which runs at 45° to the axes. Pixels which plot close to this line will not be strongly coloured and may be subject to *\*decorrelation stretching*.

**acicular** Pointed or needle-shaped.

**acid** According to the Brønsted–Lowry theory, a substance that in solution liberates hydrogen *\*ions* or protons. The Lewis theory states that an acid is a substance that acts as an electron-pair acceptor. An acid reacts with a *\*base* to give a salt and water (neutralization), and has a *\*pH* of less than 7.0. The theory was proposed in 1923 by the Danish physical chemist Johannes Nicolaus Brønsted (1879–1947) and the British chemist Thomas

Martin Lowry (1874–1936), and independently by the American theoretical chemist Gilbert Newton Lewis (1875–1946).

**acidophile** An *\*extremophile* (domain *\*Archaea*) that thrives in environments where the *\*pH* is below 5.0.

**acid rain** Precipitation with a *\*pH* of less than about 5.0, which is the value produced when naturally occurring carbon dioxide, sulphate, nitrogen oxides, and formic acid dissolve into cloud droplets. The effects of increased acidity on surface waters, soils, and vegetation are complex.

**acid rock** *\*Igneous* rock containing more than about 60% *\*silica* (SiO<sub>2</sub>) by weight, most of the silica being in the form of silicate minerals, but with the excess of about 10% as free *\*quartz*. Typical acid rocks are *\*granites*, *\*granodiorites*, and *\*rhyolites*. Compare BASIC ROCK; INTERMEDIATE ROCK. See also ALKALINE ROCK.

**acid soil** *\*Soil* having a *\*pH* less than 7.0. Degrees of soil acidity are recognized. Soil is regarded as 'very acid' when the reaction is less than pH 5.0. The *\*USDA* lists five standard ranges of soil acidity (less than pH 4.5, extremely acid; 4.5–5.0, very strongly acid; 5.1–5.5, strongly acid; 5.6–6.0, medium acid; and 6.1–6.5, slightly acid). Surface *\*soil horizons* of acid *\*brown earths* have a reaction of pH 5.0 or less.

**aclinic line** See MAGNETIC EQUATOR.

**acme zone** (**peak zone, flood zone, epibole, abundance zone**) An *\*informal* term for a body of *\*strata* containing the maximum abundance of a particular *\*taxon* occurring within the stratigraphic range of that taxon, and after which the *\*zone* is named.

**acoustic basement** The region of the Earth lying below the limit that can be imaged by *\*seismic surveys*.

**acoustic impedance** (**Z**) The product of density ( $\rho$ ) and the acoustic velocity ( $v$ ) for a given rock mass;  $Z = \rho v$ . The *\*reflection coefficient* for an interface is governed by the contrast in the acoustic impedances of the two adjacent *\*rock masses*.

**acquired characteristics** Characteristics that are acquired in the lifetime of an organism. The evolutionary theorist *\*Lamarck* suggested that traits

acquired in one generation in response to environmental stimuli would be inherited by the next generation. Thus over several generations a particular type of organism would become better adapted (see [ADAPTATION](#)) to its environment. The kinds of acquisition envisaged by Lamarck and their heritability are now discredited, although there has been a recent revival of some aspects of Lamarckism in modified form.

**ACRIMSAT** See [ACTIVE CAVITY RADIOMETER IRRADIANCE MONITOR SATELLITE](#).

**acrisols** A reference soil group in the [\\*World Reference Base for Soil Resources](#) classification scheme used by the [\\*FAO](#). Acrisols are [\\*acid soils](#) with an argic B horizon (see [ARGIC HORIZON](#)) having a [\\*cation-exchange](#) capacity of less than 24 cmol<sub>c</sub>/kg.

**acritarchs** Hollow fossil structures, 5–240 μm in diameter, inside which dinoflagellates and single-celled algae survived dry periods. They range from [\\*Precambrian](#) to [\\*Recent](#) times. They are found in marine strata, although some non-marine examples are reported from Recent beds. Acritarchs are used in [\\*correlation](#) and to distinguish onshore from offshore [\\*sediments](#).

**Acrothoracica** See [CIRRIPEDIA](#).

**acrozone** See [RANGE ZONE](#).

**actinides (actinoids)** The 15 metallic elements with [\\*atomic numbers](#) 89–103. Actinium (<sup>89</sup>Ac) is the first and gives its name to the group. The others are: thorium (<sup>90</sup>Th), proactinium (<sup>91</sup>Pa), uranium (<sup>92</sup>U), neptunium (<sup>93</sup>Np), plutonium (<sup>94</sup>Pu), americium (<sup>95</sup>Am), curium (<sup>96</sup>Cm), berkelium (<sup>97</sup>Bk), californium (<sup>98</sup>Cf), einsteinium (<sup>99</sup>Es), fermium (<sup>100</sup>Fm), mendelevium (<sup>101</sup>Md), nobelium (<sup>102</sup>No), and lawrencium (<sup>103</sup>Lr). All the actinides are radioactive, highly electropositive, tarnish when exposed to air, and react with water, releasing hydrogen.

**actinium series** See [DECAY SERIES](#).

**actinoids** See [ACTINIDES](#).

**actinolite** A member of the *\*amphiboles*,  $\text{Ca}_2(\text{Mg,Fe})_5(\text{Si}_4\text{O}_{11})_2(\text{OH,F})_2$ , with the ratio  $\text{Fe}/(\text{Fe} + \text{Mg}) = 0.9$  to  $0.5$ , belonging to the *\*tremolite–ferroactinolite* series of Ca-rich amphiboles; sp. gr. 3.0–3.4; *\*hardness* 5–6; *\*monoclinic*; light greenish-grey to dark green; white *\*streak*; *\*vitreous lustre*; habit *\*acicular*, often fibrous and felted; *\*cleavage prismatic*, good {110}; occurs widely in low- to medium-grade *\*schists* and some *\*igneous* rocks. The asbestiform variety is called *\*nephrite* and such felted forms were used in the past for insulation and fire-resistant materials, but the development of asbestosis in workers has severely restricted their use.

**Actinopterygii (ray-finned fish)** A class of the *\*Osteichthyes* (bony fish, see **BONE**), comprising the ray-finned fish, which include the majority of living bony fish of sea and fresh water. The *\*fins* are composed of a membranous web of skin supported by a varying number of spines and soft rays. They appeared first during the *\*Devonian*.

**activation analysis** See **NEUTRON-ACTIVATION ANALYSIS**.

**activation energy (energy of activation)** The energy that must be delivered to a system in order to increase the incidence within it of reactive molecules, thus initiating a reaction.

**Active Cavity Radiometer Irradiance Monitor Satellite (ACRIMSAT)** A *\*NASA* satellite instrument that was launched on 20 December 1999 and became operational on 5 April 2000. ACRIMSAT measures total solar irradiance. It is the third ACRIM experiment and extends the database that began with ACRIM 1 in 1980.



<http://www.acrim.com/>

- A NASA mission launched in 1978 to monitor total solar irradiance.

**active geophysical methods** Geophysical exploration methods which require an artificial signal to be generated. For example, exploration seismology, some *\*electromagnetic* techniques, electrical resistivity, *\*remote sensing*, and *\*induced* polarization are said to be active geophysical methods. The term is contrasted with *\*passive* geophysical methods.

**active layer** A seasonally thawed surface layer between a few centimetres and about 3 m thick, lying above the permanently frozen ground in a periglacial environment. It may be subject to considerable expansion on freezing, especially if silt-sized particles dominate, with important engineering implications. *See also* MOLLISOLS; PERMAFROST.

**Active Magnetosphere and Planetary Electrodynamics Response Experiment (AMPERE)** A US Earth-observing programme that since 2010 has used data from sixty-six Iridium Communications commercial satellites in polar orbits to provide magnetic field measurements in near-real time to assist in forecasting weather in space and to enable 24-hour tracking of Earth's response to blasts of plasma ejected at supersonic speed by the Sun.

**active margin (seismic margin)** The margin of a continent that is also a **\*plate margin**. The alternative term, 'Pacific-type margin', indicates the range of features (e.g. **\*earthquakes**, andesitic (*see* ANDESITE) volcanic chains, offshore oceanic **\*trenches**, and young fold mountains) which may be associated with active margins. Some authors distinguish an 'Andino-type margin', involving an oceanic and a continental plate, from a 'Japan-type margin', involving an oceanic plate and an **\*island arc**. The term 'Mediterranean-type margin' is also in use, although to a lesser extent, to signify the coincidence of continental edges and plate margins in a **\*collision zone**.

**active methods** *See* ACTIVE GEOPHYSICAL METHODS.

**active pool** The part of a **\*biogeochemical cycle** in which the nutrient element under consideration exchanges rapidly between the biotic and abiotic components. Usually the active pool is smaller than the **\*reservoir pool**, and it is sometimes referred to as the 'exchange' or 'cycling' pool.

**active remote sensing** **\*Remote** sensing which is based on the illumination of a scene by use of artificial radiation. An example is **\*radar**. *Compare* PASSIVE REMOTE SENSING.

**activity** A broadly used term which refers to the rate or extent of a change associated with some substance or system. For example, it may be the tendency of a metal high in the electromotive series to replace another metal lower in the series, e.g. magnesium displacing copper from most of

its compounds. It may also be used to describe the rate of decay of atoms by radioactivity.

**activity coefficient ( $\gamma$ )** The ratio of chemical activity (i.e. the effective concentration ( $a$ ) of a component in a solution, to the actual mole fraction ( $X$ ) present in solution: ( $\gamma = a/X$ ). Values for activities are determined experimentally in a number of ways, including measuring the ratio of the **\*vapour pressure ( $p$ )** of a known concentration of the substance in solution to the vapour pressure ( $p^*$ ) of the pure substance:  $a = p/p^*$ . In an ideal solution the activity coefficient = 1, and the activity of the component is equal to its mole fraction. In general, the greater the amount of dissolved material, the lower the activity coefficients of each of the species present.

**Actonian** A **\*stage** of the **\*Ordovician** (454–453 Ma ago) in the Upper **\*Caradoc**, underlain by the **\*Marshbrookian** and overlain by the **\*Onnian**.

**actual evapotranspiration (AE)** The amount of water that evaporates from the surface and is transpired by plants if the total amount of water is limited. Compare **POTENTIAL EVAPOTRANSPIRATION**.

**actualism** The theory that present-day processes provide a sufficient explanation for past geomorphological phenomena, although the rate of activity of these processes may have varied. The theory was first clearly expressed in 1749 by G. L. L. **\*Buffon** (1707–88), and was the essential principle of **\*uniformitarianism** as presented in 1830 by C. **\*Lyell** (1797–1875).

**acuity** The ability of a human to discern spatial variation in a scene.

**ACV** See **AGGREGATE TESTS**.

**Adam** The postulated male ancestor for all modern humans, who lived in Africa between about 100 000 and 200 000 years ago. ‘Adam’ is based on a change in the human Y chromosome that occurred at that time in one descendant of Adam and is now present in all human males, except for some Africans. See also **MITOCHONDRIAL EVE**.

**adamantine** Of mineral **\*lustre**, brilliant, like a polished diamond.

**adamellite** A rock of granitic composition (see **GRANITE**) characterized by the presence of **\*quartz**, **\*plagioclase** feldspar, and potassic feldspar (see

ALKALI FELDSPAR) accompanied by \*biotite and/or \*hornblende. The two feldspar types occur in approximately equal proportions, the plagioclase composition lying within the oligoclase range. The name is derived from the type locality of Adamello in the Tyrol, Austria, where granites of this type were originally defined. In Britain the best-known example occurs at Shap Fell in Cumbria.

**Adams–Williamson equation** Equation describing a fundamental relationship between seismic velocities ( $v_p$  and  $v_w$ ), the \*gravitational acceleration ( $g$ ), and the adiabatic change in density ( $d\rho$ ) within the \*Earth (assuming only hydrostatic pressure) as a function of radius ( $dr$ ):

$$d\rho = g\rho / (drv_p^2 - (4/3)v_w^2)$$

This equation is directly applicable to the lower \*mantle and outer \*core, but is invalid where the composition is variable, the pressure is not hydrostatic, or the increase in pressure is not adiabatic.

**adapical** A directional term meaning towards the shell \*apex.

**adaptation** 1. Generally, the adjustments that occur in animals in respect of their environments. The adjustments may occur by \*natural selection, as individuals with favourable genetic traits breed more prolifically than those lacking these traits (genotypic adaptation), or they may involve non-genetic changes in individuals, such as physiological modification (e.g. acclimatization) or behavioural changes (phenotypic adaptation). *Compare* ABAPTATION. 2. In an evolutionary sense, that which fits an organism both generally and specifically to exploit a given environmental zone.

**adaptive radiation** 1. A burst of evolution, with rapid divergence from a single ancestral form, resulting in the exploitation of an array of habitats. The term is applied at many \*taxonomic levels, e.g. the radiation of the mammals at the base of the \*Cenozoic refers to \*orders, whereas the radiation of ‘Darwin’s finches’ in the Galápagos Islands resulted in a proliferation of \*species. 2. Term used synonymously with ‘\*cladogenesis’ by some authors.

**adaptive zone** The adaptive specialization(s) that fit the \*taxon to its environment, e.g. feeding habits.

**addition rule (Weiss zone law)** With reference to crystallographic notation, the rule stating that the indices (see [MILLER INDEXES](#)) of two [\\*crystal](#) faces in the same [\\*zone](#) always add up to the indices of a face bevelling the edge lying between them. The rule may be used to index faces on a [\\*stereogram](#), or faces at the intersection of two zones.

**additive primary colours** The spectral colours red, green, and blue, which, when mixed together by projection through filters, can be used to produce all other colours. None of the primary colours can be produced by combinations of the other two. *See also* [SUBTRACTIVE PRIMARY COLOURS](#).

**adductor muscles** *See* [MUSCLE SCAR](#).

**Adelaidean** A [\\*stage](#) (542–1300 Ma ago) of the Upper [\\*Proterozoic](#) of south-eastern Australia, underlain by the [\\*Carpentarian](#) and overlain by the Early [\\*Cambrian](#).

**Adelaidean orogeny** A late [\\*Proterozoic](#) and [\\*Ordovician](#) phase of mountain building, affecting what is now southern Australia, in which [\\*sedimentary rocks](#) of the Adelaidean system were raised by severe thrusting and overfolding, first in the south and later along the northern margin of the system.

**ADEOS** *See* [ADVANCED EARTH OBSERVING SATELLITE](#).

**adhesion ripples** *See* [ADHESION WARTS](#).

**adhesion warts (adhesion ripples)** A [\\*sedimentary structure](#) consisting of an irregular, wart-like or blistered, [\\*sand](#) surface, formed by the wind blowing dry sand over a moist surface. The warts tend to be slightly asymmetrical, with steeper sides in the up-wind direction.

**adiabat** The rate at which a [\\*parcel](#) of air cools as it rises and warms as it descends, as indicated by two lines (dry adiabat and wet adiabat) on a [\\*tephigram](#).

**adiabatic** Applied to the changes in temperature, pressure, and volume in a [\\*parcel](#) of air or liquid that occur as a consequence of the vertical movement of the fluid, and without any exchange of energy with the surrounding fluid. *See also* [DRY ADIABATIC LAPSE RATE](#); [SATURATED ADIABATIC LAPSE RATE](#).

**adit** Horizontal or nearly horizontal tunnel from the surface into a mine, for entry, drainage, or exploration.

**admission** The substitution of a *\*trace element* for a major element with a similar *\*ionic radius* but a higher *\*valency* during the crystallization of a *\*magma*, e.g. the substitution of  $\text{Li}^+$  for  $\text{Mg}^{2+}$  in the *\*pyroxenes*, *\*amphiboles*, and *\*micas*.

**adobe** A silty *\*clay*, often calcareous, found in dry, desert-lake basins. This fine-grained *\*sediment* is usually deposited by desert floods which have eroded wind-blown *\*loess* deposits. The term is of Spanish origin ('adobe' means sun-dried brick).

**adoral** On the same side of the body as the mouth.

**Adrastea (Jupiter XV)** A jovian satellite (a *\*moom*) that orbits within the main ring of Jupiter; it and *\*Metis* may be the source of the material comprising the ring. Both are considered too small to suffer tidal disruption, but eventually their orbits will decay. Adrastea is one of the smallest satellites in the solar system. It was discovered in 1979 by David Jewitt. Its diameter is 20 km ( $\pm 20$ ) ( $23 \times 20 \times 15$  km); mass  $1.91 \times 10^6$  kg; mean distance from Jupiter 129 000 km.

**adsorption** The attachment of an ion, molecule, or compound to the charged surface of a particle, usually of *\*clay* or *\*humus*, from where it may be subsequently replaced or exchanged. Ions carrying positive charges (e.g. those of calcium, magnesium, sodium, and potassium) become attached to, or adsorbed by, negatively charged surfaces (e.g. those of clay or humus). *Compare* ABSORPTION. *See also* SORPTION.

**adsorption complex** Various materials of the soil, mainly *\*clay* and *\*humus* and to a lesser degree other particles, capable of adsorbing ions and molecules.

**adularia** *See* ALKALI FELDSPAR.

**Advanced Earth Observing Satellite (ADEOS)** A satellite mission led by the *\*Japan Aerospace Exploration Agency* with international collaborators, sometimes known in Japan as *Midori* meaning green, dedicated to environmental research, observation of geophysical parameters, observation of atmospheric processes, as well as ocean colour and sea-surface

temperature. ADEOS was launched on 17 August 1996, from Tanegashima Space Center, Japan. It ceased to function on 30 June 1997, apparently due to structural damage to its solar array. ADEOS-2 was launched on 14 December 2002 but suffered a catastrophic power failure on 24 October 2003.



[http://www.jaxa.jp/projects/sat/adeos2/index\\_e.html](http://www.jaxa.jp/projects/sat/adeos2/index_e.html)

- A joint CNES, JAXA, NASA, and NASDA mission to study the Earth's environment.

**Advanced Land Observing Satellite (ALOS)** An Earth-observation satellite programme developed by the *\*Japan Aerospace Exploration Agency* to provide high-resolution data for use in cartography and environmental and hazard monitoring. The first satellite was launched on 24 January 2006 and retired in May 2011; ALOS-2 was launched on 24 May 2014. ALOS-3 is to be launched in 2019 or 2020.

**Advanced Satellite with New System Architecture for Observation (ASNARO)** A Japanese optical high-resolution mission that aims to develop a next-generation high-performance *\*minisatellite* bus system that will reduce the cost and development period by using advanced electronic technologies. The spacecraft was launched on 6 November 2014, from Dombarovsky, Russia.

**advection** The horizontal transfer of heat by means of a moving gas (usually air).

**adventive cone** See PARASITIC CONE.

**adventurine** See AVENTURINE.

**AE** See ACTUAL EVAPOTRANSPIRATION.

**aedifichnia** A category of *\*trace fossils* that comprises structures in full relief that were constructed by organisms from raw materials, e.g. mud nests of wasps, caddis fly cases, spiders' larders consisting of concentrations of insects, insect remains, and spiders.

**aegirine** \*Pyroxene mineral,  $\text{NaFe}^{3+}\text{Si}_2\text{O}_6$ ; sp. gr. 3.5; \*hardness 6; \*monoclinic; greenish-black or brown; occurs as fairly short, \*prismatic crystals in \*igneous and \*metamorphic rocks. A variety intermediate in composition between aegirine and augite is called 'aegirine–augite'. See also AUGITE; CLINOPYROXENE.

**Aegyptopithecus zeuxis** A genus and species of early \*catarrhine primates, known from abundant remains, including several nearly complete skulls, from the early \*Oligocene of the Jebelal-Qatrani Formation, Fayum, Egypt. The size of a small, living monkey, it had a long tail and could jump from branch to branch. It possessed the dental and some of the cranial characteristics of living catarrhines, but lacked many of the other cranial and most of the postcranial diagnostic features, and so represents a time when catarrhines had separated from other primates, but remained more primitive than living hominoids (\*Hominoidea) or Old World monkeys and it could have been ancestral to living catarrhines.

**AEM-2** A NASA \*minisatellite mission to monitor stratospheric \*aerosol and ozone that was launched on 18 February 1979 but experienced power problems in May 1979 and its battery failed on 7 January 1982. The satellite decayed in the atmosphere on 11 April 1989.

**Aeneas** A nanosatellite, comprising a 3-unit \*CubeSat, developed by teams of students at the University of Southern California Engineering Research Center, that was launched into an elliptical orbit on 13 September 2012. Its purpose is to track the location of cargo containers.

**aeolian abrasion** The erosion of a surface that is caused by bombardment with loose particles carried by the wind.

**aeolianite** General term for the sedimentary products of wind (aeolian) deposition.

**aeolian processes (eolian processes)** The erosion, transport, and deposition of material due to the action of the wind at or near the Earth's surface. Aeolian processes are at their most effective when the vegetation cover is discontinuous or absent.

**aeolian ripple (eolian ripple)** A ripple on the surface of a sedimentary rock that is caused by saltating grains. Aeolian ripples have a wavelength

approximately equal to the [\\*saltation](#) path of the grains, and they usually have no internal [\\*cross-lamination](#).

**Aeolis Quadrangle** A region of Mars formed in the Late [\\*Noachian](#) or Early [\\*Hesperian](#) epoch, containing both extensional and compressional land-forms and valleys (see [VALLIS](#)), some of which may be outflow channels, but some of which may be tectonic rift features.

**Aeolus** An Earth Explorer Core Mission space satellite, operated by the [\\*European Space Agency](#), that was launched on 22 August 2018, from Kourou, French Guiana, into a Sun-synchronous orbit at a mean altitude of 320 km, to measure wind profiles with an accuracy acceptable to the [\\*World Meteorological Organization](#).

**aerial photograph** A photograph taken from an aircraft. In hydrology, false-colour infrared photographs are used to determine the wetness and temperature of soils and to detect [\\*springs](#).

**aerial photography** The taking of aerial photographs of rock exposures and of the ground surface for purposes of geologic interpretation. The photographs may be taken vertically, or at a high-oblique or low-oblique angle, and may be assembled like a mosaic to provide a picture of a large area. Stereoscopic cameras (two cameras within a single body) may be used to produce pairs of pictures that provide three-dimensional pictures when observed through a stereoscopic viewer. See [PHOTOGEOLOGY](#).

**aerobic** **1.** Of an environment: one in which air (oxygen) is present. In the case of a depositional environment, one with more than 1 ml of dissolved oxygen per litre of water. Compare [ANAEROBIC](#); [DYSAEROBIC](#). **2.** Of an organism: one requiring the presence of oxygen for growth, i.e. an aerobe. **3.** Of a process: one that occurs only in the presence of oxygen.

**aerodynamic roughness** Surface irregularities that impede the passage of air, causing uneven flow. The surface over which the flow takes place may be solid, liquid (e.g. large ocean waves), or of air of different density.

**aerological diagram** Diagram to demonstrate variations with height of the physical characteristics of the atmosphere, particularly its temperature, pressure, and [\\*humidity](#).

**aeromagnetic survey** Survey of the Earth's magnetic field, based on data from **\*magnetometers** towed behind aircraft or suspended below helicopters. These instruments measure the total intensity of the **\*geomagnetic field** or, occasionally, components of this field. The resulting measurements can then be compared with theoretical models for the value of the field and the differences (**\*magnetic anomalies**) can be interpreted in terms of changes in the magnetic properties of the rocks below the survey line or grid. The magnetometers are usually flown with other instrumentation, e.g. **\*radiometric** and electromagnetic, at the lowest practicable constant height above the ground. Usually the magnetometer is housed in a 'bird' towed behind the aircraft, or in a wing-tip pod, or in a 'stinger' in the tail. In cases where the magnetometer is on board, in-board coil systems compensate for the aircraft's own magnetic field.

**Aeronian** A **\*stage** (439–436 Ma ago) of the Lower **\*Silurian** (**\*Llandovery** period) underlain by the **\*Rhuddanian** and overlain by the **\*Telychian**.

**Aeronomy of Ice in the Mesosphere (AIM)** A NASA **\*minisatellite** that was launched on 25 April 2007, from California, and that provides data relevant to research into the cause of polar mesospheric clouds, which form every summer 50–90 km above polar regions.

**aerosol** Colloidal substance, either natural or man-made, that is suspended in the air because the small size (0.01–10  $\mu\text{m}$ ) of its particles makes them fall slowly. Aerosols in the **\*troposphere** are usually removed by **\*precipitation** and their **\*residence** time is measured in days or weeks. Aerosols that are carried into the **\*stratosphere** usually remain there much longer. Tropospheric aerosols may act as **\*Aitken** nuclei but the general effect of aerosols is to absorb, reflect, or scatter radiation. Stratospheric aerosols, mainly sulphate particles resulting from volcanic **\*eruptions**, may reduce **\*insolation** significantly. About 30% of tropospheric dust particles are the result of human activities. See **ATMOSPHERIC STRUCTURE**; **MIE SCATTERING**; **RAYLEIGH SCATTERING**; **VOLCANIC DUST**.

**Aëtosauria** Mainly **\*Triassic** group of primitive thecodontian ('tooth-in-socket') reptiles (see **THECODONTIA**). They resembled heavily armoured crocodiles, and appear to have been specialized herbivores or possibly omnivores. They grew up to 3 m long, and their armour plating comprised rows of bony **\*plates**.



<http://www.ucmp.berkeley.edu/diapsids/pseudosuchia/aetosauria.html>

- Introduction to the Aëtosauria.

**AFC** See ASSIMILATION-FRACTIONAL CRYSTALLIZATION.

**AF demagnetization** See ALTERNATING-MAGNETIC-FIELD DEMAGNETIZATION.

**affine** In geometry, a transformation that preserves straight lines and the ratios of distances between points lying on straight lines. Parallel lines remain parallel. Angles and lengths of lines are not necessarily preserved.



<http://homepages.inf.ed.ac.uk/rbf/HIPR2/affine.htm>

- Affine transformation.

**AFM diagram** A three-component, triangular graph used to show how metamorphic *\*mineral* assemblages vary as a function of *\*rock* composition within one *\*metamorphic facies*. Besides SiO<sub>2</sub>, the five most abundant oxides found in *\*metamorphic rocks* are Al<sub>2</sub>O<sub>3</sub>, CaO, FeO, MgO, and K<sub>2</sub>O. The three components plotted on AFM diagrams are derived from a tetragonal diagram, with species Al<sub>2</sub>O<sub>3</sub>, K<sub>2</sub>O, FeO, and MgO, and are ideal for showing mineral assemblage variations as a function of the composition of *\*pelites*. Mineral and rock compositions plotting within this diagram are projected on to the Al<sub>2</sub>O<sub>3</sub>-FeO-MgO face from either the *\*muscovite* or K-feldspar point on the Al<sub>2</sub>O<sub>3</sub>-FeO edge. The components of the diagram are thus A (Al<sub>2</sub>O<sub>3</sub>), F (FeO), and M (MgO), with the projection geometry being accommodated on specially scaled axes. Each of these components has to be modified slightly to account for the presence of other, minor components in the rock, leading to: A (Al<sub>2</sub>O<sub>3</sub> - 3K<sub>2</sub>O); F (FeO - TiO<sub>2</sub> - Fe<sub>2</sub>O<sub>3</sub>); and M (MgO). The minerals *\*quartz* and *\*albite* are assumed to be present in the rocks and are not shown on the diagram. As in *\*ACF* diagrams, *\*tielines* connect minerals which coexist in equilibrium.

**AFMAG EM system** Audio-Frequency Magnetic ElectroMagnetic method, which uses natural electromagnetic (EM) fields (*\*sferics*) in the audio-

frequency range (1–1000 Hz) generated by thunderstorms to investigate lateral changes in the **\*resistivity methods** of the Earth's surface.

**African Plate** One of the present-day major lithospheric **\*plates**, consisting of the continental mass of Africa surrounded, except to the north, by **\*oceanic** crust and oceanic **\*ridges**. To the north, a complex picture of collision and **\*subduction zones** and **\*transform faults** has been postulated for the boundary with the **\*Eurasian Plate** and various minor plates, e.g. the Aegean Plate. The northern part of the African Plate also contains remnants of the oceanic crust of **\*Tethys**. To the north-east the Red Sea is interpreted as an actively forming ocean, at the young stage of the **\*Wilson** cycle, while the E. African **\*rifts**, partially defining what is called by some the 'Somali Plate' to the east, may be at the embryonic stage of ocean development, or possibly a stillborn ocean.

**Afternoon Constellation** See **A-TRAIN**.

**aftershock** A seismic event that occurs after an **\*earthquake**, usually within days or weeks. Although often of small **\*magnitude**, aftershocks can be more destructive as buildings and structures have already been weakened.

**afterslip (post-seismic slip)** Movement that continues for a year or longer in the aftermath of a major **\*earthquake**. Afterslip occurs in discrete pulses of up to 10 mm, these becoming less frequent over time.

**Aftonian** The earliest (2.6–0.117 Ma ago) of four **\*interglacial \*stages** in N. America, following the **\*Nebraskan** glacial episode, and approximately equivalent to the **\*Donau/Günz interglacial** of Alpine terminology. Climatically it was marked by mild summers and winters warmer than those in present-day N. America.

**Agassiz, Jean Louis Rodolphe** (1807–73) A Swiss geologist who worked initially on fossil fish, Agassiz is better known for his **\*glacial theory** (1837). He met **\*Buckland** in 1840, and persuaded him that **\*drift** deposits in Britain were evidence of a glacial epoch. In 1846 he moved to the USA to become professor of zoology and geology at Harvard, where he founded the Museum of Comparative Zoology (1859).

**agate (mocha stone)** Variety of chalcedonic silica (SiO<sub>2</sub>) that is **\*cryptocrystalline**. It is similar to **\*chalcedony** except that impurities of iron and manganese may give it a distinct colour banding which is frequently

precipitated in concentric zones. Moss agate contains delicate, fern-like, dendritic patterns. Agates may be cut and polished as decorative stones.

**age** **1.** The interval of geologic time equivalent to the *\*chronostratigraphic unit* '*\*stage*'. Ages are subdivisions of *\*epochs* and may themselves be subdivided into *\*chrons*. An age takes its name from the corresponding stage, so like the stage name it carries the suffix '-ian' (or sometimes '-an'); the term 'age' is sometimes capitalized when used in this formal sense, e.g. '*\*Oxfordian Age*'. **2.** An *\*informal* term to denote a time span marked by some specific feature, e.g. '*\*Villafranchian mammalian age*'.

**ageostrophic wind** The vector difference between the *\*geostrophic* and the actual winds.

**agglomerate** Coarse-grained volcanic rock with rounded to subangular fragments. These fragments are mainly larger than 2 cm in size, but the mixture of fragments is typically ill sorted and the *\*matrix* may be fine grained. An agglomerate may be the product of a volcanic explosion and therefore a *\*pyroclastic* rock, but often the term 'agglomerate' is applied to brecciated volcanic rocks of uncertain origin. Those deposits may range from vent *\*breccias* to debris from mudflow or *\*lahar* deposits.

**agglutinate** A constituent of lunar soils comprising glass-bonded *\*aggregates*, which consist of *\*glasses* and rock and mineral fragments welded together by glass. These aggregates form during the impact of micrometeorites into lunar soils. Their abundance in a lunar soil is an index of exposure to micrometeorite bombardment, and hence to soil maturity. The average size of agglutinates in mature soils varies, but tends toward a mean of 60  $\mu\text{m}$ .

**aggradation** The general accumulation of unconsolidated sediments on a surface, which thereby raise its level. A large range of mechanisms may be involved, including *\*fluvial*, *\*aeolian*, marine, and *\*slope processes*.

**aggregate** **1.** In the building and construction industry, a mixture of mineral substances (bulk *\*minerals*), e.g. sand, gravel, crushed *\*rock*, stone, slag, and other materials (e.g. colliery spoil, pulverized fuel ash) which, when cemented, forms *\*concrete*, mastic, mortar, plaster, etc. Uncemented, it can be used as road-making material, railway ballasts, filter beds, and in some manufacturing processes as flux. In road-making, aggregate mixed with

**\*bitumen** is called ‘coated stone’, and different physical characteristics are required for the different layers comprising the road **\*pavement**. Fine aggregate is less than 6.35 mm in diameter, coarse aggregate greater than 6.35 mm. See **AGGREGATE TESTS**; **PAVEMENT**. **2.** Group of soil particles adhering together in a cluster; the smallest structural unit, or ped, of soil. Aggregates join together to make up the major structural soil units.

**aggregate abrasion value** See **AGGREGATE TESTS**.

**aggregate crushing value** See **AGGREGATE TESTS**.

**aggregate impact value** See **AGGREGATE TESTS**.

**aggregate tests** Specific tests used to determine the suitability of **\*aggregates** for special purposes. There are tests for: (a) shape and texture (the angularity number), to determine whether particles have a large angle of friction with good bonding properties; (b) size and grading, to determine whether particles will pack well; (c) moisture content, to discover whether materials absorb so much water that freeze–thaw action might cause the break-up of structures; (d) rock density, which may affect the economics of an operation; (e) strength, determined by subjecting the rock to hammering in a standard test and measuring the percentage of fine material produced (the aggregate impact value, or AIV); (f) resistance to crushing (the aggregate crushing value, or ACV), measured in a similar manner; (g) resistance to abrasion, measured by standard equipment to give the aggregate abrasion value (AAV)—the lower the AAV, the more resistant the rock; and (h) resistance to polishing, measured in the laboratory to give the polished stone value (PSV)—the higher the PSV, the greater the resistance to polishing and therefore skidding, and the more valuable the material.

**aggregation** **1.** Process in which soil particles coalesce and adhere to form soil aggregates. The process is encouraged by the presence of bonding agents such as organic substances, **\*clay**, iron oxides, and ions (e.g. calcium and magnesium). **2.** Progressive attachment of particles (e.g. ice or snow) or droplets around a nucleus, thereby causing its growth.

**Aglaophyton major** See **RHYNIA**.

**Agnatha** (phylum **\*Chordata**, subphylum **\*Vertebrata**) Superclass of jawless, fish-like vertebrates, with sucker-like mouths, including the extant

lampreys, slime-eels and hagfish, and some of the earliest primitive vertebrates, with heavily armoured forms, e.g. *\*Cephalaspis* (see also *OSTEOSTRACI*), *Pteraspis* (see *HETEROSTRACI*), and *Jamoytius* (see *ANASPIDA*). They appeared first during the *\*Ordovician*.

**Agnostida** An order of *\*Trilobita* that lived from the Lower *\*Cambrian* to Upper *\*Ordovician*. Most were blind, lacking sutures, and typically are found rolled up. They had a subequal *\*cephalon* and *\*pygidium*. There were two suborders. They are important stratigraphic markers.

**agonic line** A line joining points where the magnetic *\*declination* is zero.

**agric horizon** Mineral-soil diagnostic horizon formed from an accumulation of *\*clay*, *\*silt*, and *\*humus*, which has moved down from an overlying, cultivated soil layer. It is a *\*soil horizon* created by agricultural management, and is identified by its near-surface position, and by *\*colloids* accumulated in the pores of the soil.

**agricrion** (*pl. agrichnia*) A *\*trace fossil* comprising a burrow that formed the permanent dwelling of an organism and was used to trap or culture smaller organisms for food or use them in chemosymbiosis.

**Agricola, Georgius (Georg Bauer Agricola)** (1494–1555) The author of works on ‘geology’ and mineral classification, and of the first comprehensive record of mining, *De Re Metallica* (1556). Using Roman sources and contemporary German knowledge, his books became basic reference material for two centuries.

**agrometeorology** The study of the relationship between conditions in the surface layers of the atmosphere and those in the surface of the Earth, as this affects agriculture.

**AGU** See *AMERICAN GEOPHYSICAL UNION*.

**Agulhas current** Part of the large-scale circulation of the southern Indian Ocean. It is a surface-water current that flows off the east coast of southern Africa between latitudes 25° S and 40° S in a south-westerly direction. Flow velocity varies seasonally between 0.2 and 0.6 m/s.

**ahermatypic** Applied to corals that lack zooxanthellae (symbiotic unicellular *\*algae*) and that are not *\*reef*-forming.

**AIM** See AERONOMY OF ICE IN THE MESOSPHERE.

**AIPG** See AMERICAN INSTITUTE OF PROFESSIONAL GEOLOGISTS.

**airborne dust analysis** Sampling and determination of airborne particles. This technique requires size segregation of the particles and a device for collection during updrafts in order to obtain only local particles. Modern equipment sucks dust directly off vegetation for analysis.

**airborne gravity survey** A regional \*gravity survey undertaken from the air. Such surveys are now rapid and precise because of the development of \*gravimeters capable of being compensated for changes in the motion and flight path of an aircraft, particularly a helicopter.

**airgun** A seismic source which discharges a bubble of highly compressed air into water. Airguns are most commonly used in marine seismic exploration, but can also be used as a down-hole seismic source.

**air-lift pump** A device composed of two pipes, one inside the other. Air is blown down the inner pipe, which is slightly shorter than the outer pipe. The result of this is to push an air–water mixture up the gap between the two pipes. This is a useful pump for obtaining samples from very small diameter boreholes.


**air mass (airmass)** Large body of air (sometimes of oceanic or continental proportions) identified primarily by an approximately constant wet-bulb-potential temperature (i.e. the lowest temperature to which the air can be cooled by the evaporation of water into it). The temperature and \*humidity characteristics of an air mass, which are roughly the same within the one air mass at a particular latitude and height, are modified by and modify the atmospheric environment through which the air mass passes.

**air wave** A sound wave which travels through the air from a seismic shot. The speed of such a wave is approximately 330 m/s.

**Airy, George Biddell** (1801–92) A Cambridge astronomer and mathematician, Airy became Astronomer Royal in 1835. He investigated planetary motion and tides, and studied the Earth and its density, using gravity measurements. His name is used to describe one version of the theory of \*isostasy. His wide-ranging advice to the government on

scientific issues created, for the first time, the role of a professional scientific civil servant.

**Airy model** A model to account for *\*isostasy* which in the *\*lithosphere* assumes a constant density ( $\rho_c = 2\,670\text{ kg/m}^3$ ), but in which topographic elevations ( $h$ ) are compensated by the presence of ‘roots’ replacing high-density *\*mantle* rocks ( $\rho_m = 3\,300\text{ kg/m}^3$ ) by lower-density lithospheric rocks. The depth of the root ( $d$ ) is equal to  $h\rho_c/(\rho - \rho_c)$ . See also [PRATT MODEL](#).



**Airy model**

**Airy phase** When a high-frequency seismic wave is superimposed on a low-frequency ground wave, the two frequencies gradually approach one another until they merge, at which point they form a single wave with a relatively large amplitude, called the ‘Airy phase’.

**AlSat** See [ALGERIA SATELLITE](#).

**Aitken nuclei counter** Device for the estimation of the concentration of particles with radii of more than  $0.001\ \mu\text{m}$  in a sample of air. Air is made to expand in a chamber: this causes it to cool. Water vapour in it condenses on to particles, forming a mist whose opacity allows estimation of the number of particles present. See also [AITKEN NUCLEUS](#).

**Aitken nucleus** Suspended, atmospheric, solid particle with a radius of less than 0.2  $\mu\text{m}$ . On average, their concentration varies from less than 1000/cm<sup>3</sup> over oceans to 150 000/cm<sup>3</sup> in urban areas. See [AITKEN NUCLEI COUNTER](#); [NUCLEUS](#).

**AIV** See [AGGREGATE TESTS](#).

**AIW** See [ANTARCTIC INTERMEDIATE WATER](#).

**åkermanite** See [MELILITE](#).


**A'KF diagram** A three-component triangular graph that is used to show how metamorphic *\*mineral* assemblages vary as a function of rock composition within one *\*metamorphic facies*. The three components are: A' = (Al<sub>2</sub>O<sub>3</sub> + Fe<sub>2</sub>O<sub>3</sub>) - (Na<sub>2</sub> + K<sub>2</sub>O + CaO); K = (K<sub>2</sub>O); F = (FeO + MgO + MnO). See also [ACF DIAGRAM](#).



[http://www.tulane.edu/~sanelson/eens212/triangular\\_plots\\_metamorphic\\_petrology.htm](http://www.tulane.edu/~sanelson/eens212/triangular_plots_metamorphic_petrology.htm)

- Describes triangular plots in metamorphic petrology.

**aklé** French term for a network of sand *\*dunes* found especially in the western Sahara. The basic unit of the network is a sinuous ridge, at right angles to the wind, made up of crescent-shaped sections which alternately face the wind (linguoid) and back to the wind (barchanoid). Aklé patterns require winds from one direction, and a large quantity of sand.



**Aklé dune**

**Aksayan** A Russian-Kazakhstanian *\*stage* (493–491.5 Ma ago) of the Late *\*Cambrian \*epoch*, preceded by the *\*Sakian* and followed by the *\*Batyrbayan*.

**aktuopalaeontology** A branch of *\*taphonomy* in which experimenters observe the decay of recently living organisms under natural conditions.

**alabaster** See GYPSUM.

**alar** The first lateral protosepta (see SEPTUM) on either side of the *\*cardinal* septum. The term is used in descriptions of the septal development of the rugose corals (*\*Rugosa*), and may also be applied to *\*fossulae* which occur in a similar position.

**alas** A *\*thermokarst* depression with relatively steep sides and a flat floor, which may be occupied by a lake. Alases are well developed in Siberia (the word ‘alas’ is of Yakutian origin) where they can occupy 40–50% of the land surface.

**Alaska current** Oceanic water boundary current produced by the deflection of the *\*N. Pacific current* by the N. American continent. It flows in a north-westward direction along the south-eastern margin of Alaska. The Alaska current is also called the *\*Aleutian* current in some texts.

**A-layer** The seismic layer corresponding to the *\*crust* of the Earth. It varies in thickness from a few kilometres to 70–90 km. The base lies on the *\*mantle* and this boundary is the *\*Mohorovičić* discontinuity.

**albedo** The proportion of *\*insolation* that is reflected back from the Earth, from the tops of the clouds, and from the atmosphere, without heating the receiving surface. It averages about 30%, but varies widely according to the substance and texture of the surface, and the angle and wavelength of the incident radiation. The value for green grass and forest is 8–27% (over 30% for yellowing deciduous forest in autumn); for cities and rock surfaces, 12–18% (over 40% for chalk and light-coloured rock and buildings); for sand up to 40%; for fresh, flat snow up to 90%; and for calm water only 2% in the case of vertically incident radiation but up to 78% where there is a low angle of incidence. The albedo for cloud surfaces averages 55%, but can be up to 80% for thick *\*stratocumulus*.

**albedo feature** An area on the surface of a planet or other extraterrestrial body that is markedly lighter or darker (i.e. shows a contrast in **\*albedo**) than its surroundings.

**albeluvisols** A reference soil group in the **\*World Reference Base for Soil Resources**. Albeluvisols have an argic B horizon (see **ARGIC HORIZON**) with an irregular upper boundary (a condition sometimes known as ‘tonguing’).

**Alberta low** Storms common in Alberta, Canada, and associated with heavy rain and snow. The storms form as a result of **\*cyclone** regeneration after passage over the Canadian Rockies: as they move eastwards they bring very cold conditions, with blizzards.

**Albertan** A **\*series** (501–513 Ma ago) of the Middle **\*Cambrian** of N. America, equivalent to the **\*St David’s**.

**Albian \*Stage** (99.6–112 Ma ago) in the **\*Cretaceous**, underlain by the **\*Aptian**, and overlain by the **\*Cenomanian**. It is known to contain a great variety of **\*molluscs**, with the **\*gastropods** in particular being useful **\*zonal** indicators between continents. The Gault and Speeton Clays of England are Albian.

**albic** Applied to an almost white soil in which there is little **\*clay** or oxides coating the sandy or silty particles. The albic **\*horizon** lies at or below the surface.

**albite** See **ALKALI FELDSPAR; PLAGIOCLASE FELDSPAR**.

**albite–epidote–amphibolite facies** A set of metamorphic **\*mineral** assemblages that is produced by the **\*metamorphism** of a wide range of initial rock types under the same metamorphic conditions, and is typically characterized by the development of the mineral assemblage **\*albite–\*epidote–\*hornblende** in rocks of **\*basic \*igneous** composition such as **\*basalts**. Other rocks of contrasting composition, e.g. **\*shales** or **\*limestones**, would each develop their own specific mineral assemblage, even though they are all being metamorphosed under the same conditions. The variation of mineral assemblage with starting rock composition reflects a particular range of pressure, temperature, and  $P(\text{H}_2\text{O})$ . Experimental studies of mineral  $P$ – $T$  stability fields indicate that the **\*facies** represents a range of low-pressure, moderate-temperature conditions. See **AMPHIBOLITE**.

**albite twin** The *\*plagioclase feldspars*, particularly *\*albite* ( $\text{NaAlSi}_3\text{O}_8$ ), are frequently twinned on the albite law (see **TWIN LAW**) where the *\*twin plane* and composition plane is (010). This twinning is often repeated to give a series of fine lamellae, seen in the hand specimens as striations (particularly on the basal plane); such twinning is usually called 'polysynthetic' or '*\*lamellar*' twinning.

**albitization** The partial or complete replacement of pre-existing *\*plagioclase* or *\*alkali feldspar* by albite. There are a number of ways in which this can be achieved. A common process involves the residual water-rich vapour released during the final stages of crystallization of a *\*granite* body. This vapour, which can carry high concentrations of  $\text{Na}^+$  in solution, rises through the granite body and reacts with the feldspars present in the granite, converting them to albite which is stable under the lower temperature vapour-rich conditions. A typical reaction that partially or completely replaces plagioclase would be:  $\text{CaAl}_2\text{Si}_2\text{O}_8 + 4\text{SiO}_2 + 2\text{Na}^+ \rightarrow 2\text{NaAlSi}_3\text{O}_8 + \text{Ca}^{2+}$ ; anorthite + quartz + sodium (in aqueous solution)  $\rightarrow$  albite + calcium (in aqueous solution). This type of reaction, where a rock simmers in its own juices, is termed a '*\*deuteric* reaction'. Another way in which albitization can be achieved is by the reaction of ocean-floor *\*basalts* with sea water in thermal circulation cells within the basalt layer of the *\*oceanic crust*.

**alcove** A steep-sided hollow eroded by a stream from an exposed rock face.

**alcrete** See **DURICRUST**.

**Aldingan** A *\*stage* (33–36 Ma ago) in the Lower *\*Tertiary* of south-eastern Australia, underlain by the *\*Johannian*, overlain by the *\*Janjukian*, and roughly contemporaneous with the *\*Bartonian* and *\*Priabonian* stages.

**alete** See **SPORE**.

**Aleutian current (Sub-arctic current)** The oceanic current that flows westwards south of the Aleutian Islands and parallel to, but north of, the *\*N. Pacific current*. The water mass is a mixture of water from the *\*Kuroshio* and *\*Oyashio* currents. See also **ALASKA CURRENT**.

**Aleutian low** Region of the N. Pacific, near the Aleutian Islands, where the average value of atmospheric pressure is low, owing to the frequency of

low-pressure systems (cyclones) moving into and occupying the region. Any one of these systems, when present on an individual day, may be called 'an Aleutian low'. Some of them are intense, others much less so. The term is the Pacific equivalent of '\*Iceland low', used in the Atlantic.

**Aleutian Trench** The oceanic \*trench which marks the boundary between the \*N. American Plate and the \*Pacific Plate. The \*subduction of the Pacific Plate changes from normal to oblique from west to east along the trench, with the boundary becoming a \*transform fault before subduction continues in the \*Kuril Trench. Towards the eastern end of the Aleutian Trench there is an increasingly wide \*accretionary wedge, and an absence of andesitic \*volcanoes.

**Alexandrian** A \*series of the Lower \*Silurian of N. America equivalent to the Lower–Middle \*Llandovery.

**alexandrite** See CHRYSOBERYL.

**ALEXIS** See ARRAY OF LOW-ENERGY X-RAY IMAGING SENSORS.

**alfisols (grey-brown podzolics)** An order of mineral soils that have \*clay-enriched or \*argillic B \*horizons; are alkaline to intermediate in reaction, with the \*base saturation in the B horizon more than 35%; are usually derived from base-rich parent materials; and are drier than –15 bars moisture potential for at least three months when plants could grow.

**Alfvén waves** Magnetohydrodynamic waves that are produced by coupling forces between the \*geomagnetic field and highly conductive fluids. Alfvén waves travel along magnetic field lines when jets of highly conductive fluid or charged particles flow across the field lines. The waves were discovered by the Swedish astrophysicist Hannes Olof Gösta Alfvén (1908–95).

**alga** (*pl.* **algae**) Common (non-\*taxonomic) name for a relatively simple type of plant which is never differentiated into root, stem, and leaves; which contains chlorophyll *a* as the primary photosynthetic pigment; which has no true vascular (water-conducting) system; and in which there is no sterile layer of cells surrounding the reproductive organs. The algae range in form from single cells (\*Protista) to plants many metres in length; algae can be found in most habitats on Earth, although the majority occur in freshwater or marine environments. See BACILLARIOPHYCEAE; CHAROPHYCEAE;

CHLOROPHYTA; CHRYSOPHYCEAE; DINOPHYCEAE; PHAEOPHYCEAE;  
RHODOPHYCEAE.

**algal bloom** Sudden growth of algae in an aquatic ecosystem. It can occur naturally in spring or early summer when primary production exceeds consumption by aquatic herbivores (see PRIMARY PRODUCTIVITY). Algal blooms may also be induced by nutrient enrichment of waters due to pollution.

**algal limestone** See LEIGHTON-PENDEXTER CLASSIFICATION.

**algal mat** A sheet-like accumulation of \*Cyanobacteria (blue-green algae) developed in shallow marine \*subtidal to \*supratidal environments, as well as in lakes and swamps. The algae cover the \*sediment surface, and will in turn trap sediment to produce a laminated alternation of dark, organic-rich algal layers and organic-poor sediment layers. See also STROMATOLITE.

**Algeria Satellite (AlSat)** An optical Earth-observation project of the Algerian National Space Technology Centre (CNTS). AlSat-1, launched on 28 November 2002, is a five-satellite constellation; AlSat-2 comprises two satellites (A and B), launched in 2010 and 2014. The project aims to supply high-quality surface images.

**alginite** See COAL MACERAL.

**Algoma-type granular iron formation** A type of \*granular iron formation that comprises lenticular deposits up to 100 m thick, derived from volcanic rocks and \*turbidites. They are associated with \*back-arc basins and rift zones within \*cratons. See also RAPITAN-TYPE GRANULAR IRON FORMATION; SUPERIOR-TYPE GRANULAR IRON FORMATION.

**Algonkian** A \*Precambrian \*system (Van Eysinga, 1975) of equivalent time period to the \*Proterozoic.

**aliquot** 1. A portion of the total amount of a \*solution. 2. A section of land divided from a larger area by measuring distances along its edges, rather than its area.

**aliasing** A distortion in the frequency of sampled data produced by insufficient sampling per wavelength, which can result in spurious frequencies. When the sampling rate is too low to represent the wave-form

accurately, then aliasing will occur. To avoid aliasing, the sampling frequency should be at least twice that of the highest-frequency component contained within the sampled wave-form. Alternatively, an anti-alias filter can be applied, which removes frequency components above the [\\*Nyquist frequency](#).

**alisols** A reference soil group in the [\\*World Reference Base](#) for Soil Resources used by the [\\*FAO](#). Alisols have an argic B horizon (see [ARGIC HORIZON](#)) with a [\\*cation-exchange](#) capacity of more than 24 cmol<sub>c</sub>/kg clay and a [\\*base saturation](#) of less than 50% within 100 cm of the soil surface. Alisols have a high concentration of aluminium.

**alkali–aggregate reaction** A chemical reaction that can lead to damage in [\\*concrete](#) structures. Free lime (CaO) in [\\*cement](#) reacts with CO<sub>2</sub> in the atmosphere to precipitate CaCO<sub>3</sub> around the cement grains. This protects them from [\\*weathering](#) and also gives an alkalinity level ([\\*pH](#) higher than 7.0) which helps to protect steel from corrosion. If the aggregate contains soluble [\\*silica](#), however, new minerals may precipitate by reaction between the aggregate and the cement. These may absorb water, causing the concrete to swell and eventually crack. Water entering these cracks may cause rusting of reinforcement bars and repeated wetting and drying may eventually destroy a structure.

**alkali basalt** A fine-grained, dark-coloured, volcanic rock characterized by [\\*phenocrysts](#) of [\\*olivine](#), titanium-rich [\\*augite](#), [\\*plagioclase](#), and iron oxides. For similar SiO<sub>2</sub> concentrations, alkali basalts have a higher content of Na<sub>2</sub>O and K<sub>2</sub>O than other [\\*basalt](#) types such as [\\*tholeiites](#). They are also characterized by the development of [\\*modal](#) [\\*nepheline](#) in their [\\*groundmass](#) (only seen with the highest powered lens on a petrological microscope) and normative nepheline (Ne) in their [\\*CIPW](#) norms. Alkali basalts are typically found on updomed and rifted [\\*continental crust](#), and on oceanic islands such as Hawaii and Ascension Island.

**alkalic** See [ALKALINE](#).

**alkali-calcic series** See [CALC-ALKALINE](#).

**alkalic series** See [CALC-ALKALINE](#).

**alkali feldspar** A group of *\*silicate* minerals that contain the alkali metal elements potassium and sodium. The normal feldspar minerals (including the calcium-bearing varieties) can be plotted on a chemical basis into a triangle which has  $\text{KAlSi}_3\text{O}_8$  (potassium feldspar, sanidine, orthoclase (Or), or microcline),  $\text{NaAlSi}_3\text{O}_8$  (sodium feldspar, albite, or Ab), and  $\text{CaAl}_2\text{Si}_2\text{O}_8$  (calcium feldspar, anorthite, or An) at the three apices. The alkali feldspars are represented by the edge of the triangle joining  $\text{KAlSi}_3\text{O}_8$  and  $\text{NaAlSi}_3\text{O}_8$  and these minerals may also contain up to 10% by weight of the third phase ( $\text{CaAl}_2\text{Si}_2\text{O}_8$ ). At high temperatures the alkali feldspars show complete *\*solid solution* between the potassium and sodium *\*end-members*, but as the temperature drops unmixing occurs and potassium feldspar and sodium feldspar separate out to produce a perthitic texture. Depending upon the final temperature, a range of perthites may result, from coarse (*\*perthite*), representing perthites formed during a large drop in temperature, to fine (*\*micropertthite*), and finally to very fine (*\*cryptopertthite*), representing perthites invisible to the naked eye and often invisible under the microscope, but observed by *\*X-ray diffraction (XRD)* techniques. If the amount of potassium exceeds that of sodium, then potassium feldspar is the host and sodium feldspar occurs within the host mineral as *\*blebs*, irregular patches, etc. In the alkali feldspars, perthitic textures occur in the compositional range  $\text{Or}_{85}\text{Ab}_{15}$  to  $\text{Or}_{15}\text{Ab}_{85}$  (or  $\text{Or}_{85}$  to  $\text{Or}_{15}$ ). K-feldspar ( $\text{KAlSi}_3\text{O}_8$ ) is the general name for the *\*monoclinic*, potassium-rich end-member: sp. gr. 2.6; *\*hardness* 6; white, sometimes with a reddish tint; *\*vitreous* *\*lustre*; crystalline, *\*prismatic*, with simple twins (see *CRYSTAL TWINNING*). It is an *\*essential* constituent of *\*acid* *\*igneous* rocks and *\*arkoses* and is used in the manufacture of glazes, porcelain, and pottery. Microcline has the same physical properties and composition as orthoclase, but is *\*triclinic* and is characterized by 'cross-hatched' twinning. It is greyish-white, but bright green in the variety known as 'amazonstone' ('amazonite'). Anorthoclase is very similar to microcline, but the amount of sodium exceeds that of potassium. Crystal twinning is common particularly along the *\*pericline* and albite laws. Sanidine is the high-temperature variety of orthoclase and the inversion temperature is at 900°C. It occurs in quickly cooled lavas. Adularia is a variety of microcline, but with up to 10% sodium substituting for potassium. It may show an opalescent play of colours to give a variety known as 'moonstone'. Albite

( $\text{NaAlSi}_3\text{O}_8$ ) is the sodium-rich end-member of both the alkali feldspars and the *\*plagioclase* feldspars. The semi-precious moonstone, with its characteristic bluish sheen or *\*schiller*, is an example of a perthitic alkali feldspar.

**alkaline (alkalic)** 1. Having a *\*pH* greater than 7.0. 2. See *ALKALINE ROCK*.

**alkaline rock** *\*Igneous* rock containing a relatively high concentration of the alkali (lithium, sodium, potassium, rubidium, caesium, and francium) and alkaline earth metals (magnesium, calcium, strontium, barium, and radium). Both silica-saturated and silica-undersaturated varieties exist, expressed in the presence of *\*alkali* feldspars and *\*feldspathoids* respectively. Alkali *\*ferromagnesian* minerals are usually present, and their identity depends on the composition of the rock. Igneous rocks of the alkaline suite span the composition range from *\*basic* to *\*acid*, and may be *\*intrusive* or *\*extrusive*.

**alkaline soil** Soil with a *\*pH* greater than 7.0. Degrees of soil alkalinity are recognized. The *\*USDA* lists soils with pH 7.4–7.8 as mildly alkaline; 7.9–8.4 as moderately alkaline; 8.5–9.0 as strongly alkaline; and more than 9.0 as very strongly alkaline. Soil is not regarded as highly alkaline unless the reaction is between 8.0 and 10.0. The full range of the pH scale (0–14) is not used in soils, as the reaction of most soils is between pH 3.5 and pH 10.0. A *\*base saturation* of 100% indicates a pH of about 7.0 or higher.

**alkaliphile** An *\*extremophile* (domain *\*Archaea*) that thrives in environments where the *\*pH* is above 9.0.

**alkenones** A group of long-chain unsaturated methyl and ethyl *n*-ketones with 35–41 carbon atoms and 1–4 double bonds that are produced by a few species of haptophyte phytoplankton (e.g. *\*coccolithophorids*). These organisms require sunlight and occur mainly in the photic zone. Alkenones are found in marine sediment cores of *\*Eocene* to modern age, in some *\*lacustrine* environments, and in exposed outcrops of uplifted marine sediment. Species producing alkenones alter the relative proportions of their alkenones by increasing the proportion of saturated alkenones as the water temperature rises. Thus, the degree of unsaturation allows the past water temperature to be estimated. See *UNSATURATION INDEX*.

**allanite (orthite)** \*Mineral, with the formula  $(Ca,Ce,Y,La,Th)_2(AlFe)_3Si_3O_{12}(OH)$ ; sp. gr. 3.4–4.2; \*hardness 5.0–6.5; \*monoclinic; light brown to black; pitchy to \*sub-metallic \*lustre; faintly radioactive; \*crystals normally \*prismatic, often \*tabular, sometimes \*massive; \*cleavage imperfect {001}; often occurs as an \*accessory mineral in granitic rocks, \*syenites, \*gneisses, and \*skarns.

**Alleghanian orogeny** A phase of mountain building, that began in the Early \*Carboniferous and was completed by the end of the \*Permian, caused by the collision between N. America and Africa. It formed part of the general WSW to ENE \*Hercynian belt. The orogeny affected the Lower \*Palaeozoic \*basement and Lower Permian strata along the western margin of the southern and central parts of the Appalachian Mountains extending from what is now Pennsylvania to Alabama, with effects as far north as New Brunswick and Newfoundland. See APPALACHIAN OROGENIC BELT.

**allele** Common shortening of the term ‘allelomorph’. One of two or more forms of a \*gene arising by mutation and occupying the same relative position (locus) on homologous \*chromosomes.

**allelomorph** Term that is commonly shortened to ‘\*allele’.

**Allen’s rule** A corollary to \*Bergmann’s rule and \*Gloger’s rule, holding that a race of warm-blooded species in a cold climate typically has shorter protruding body parts (nose, ears, tail, and legs) relative to body size than another race of the same species in a warm climate. This is because long protruding parts emit more body heat, and so are disadvantageous in a cool environment, but advantageous in a warm environment. The idea is disputed, critics pointing to many other adaptations for heat conservation which probably are more important, notably fat layers, feathers, fur, and behavioural adaptations to avoid extreme temperatures.

**Allerød** See BØLLING-ALLERØD INTERSTADIAL.

**allochem** The collective term for particles (grains) which form the framework in mechanically deposited \*limestones. In the limestone classification of \*Folk, allochems are often found together with a \*carbonate mud \*matrix (\*micrite) and may subsequently have \*pore spaces filled by sparry \*calcite \*cement (\*sparite). Common allochems include skeletal fragments (\*bioclasts), \*ooids, \*peloids, and \*intraclasts.

**allochemical** A *\*limestone* defined by the *\*Folk* classification as comprising *\*allochems* with either a sparry *\*calcite* *\*cement* (*\*sparite*), or a *\*microcrystalline* *\*calcite* (*\*micrite*) *\*matrix*. Limestones lacking allochems are defined by Folk's classification of limestones as *\*orthochemical* limestones or *\*autochthonous* *\*reef* rocks.

**allochthon** A body of rock that has been transported to its present position, usually over a considerable distance. See ALLOCHTHONOUS.

**allochthonous** Not indigenous; acquired. In the Earth sciences the term is applied to geologic units that originated at a distance from their present position. Such displacement may be due to lateral thrusting and overfolding, or to gravity gliding. Compare AUTOCHTHONOUS.

**allochthonous terrane** See TERRANE.

**alloclast** A *\*clast* produced by subterranean, igneous processes that break up pre-existing volcanic rocks. Compare AUTOCLAST; EPICLAST; HYDROCLAST.

**allocyclic mechanisms** Events responsible for the accumulation of sediments that are external to the sedimentary system itself (e.g. sea-level changes, tectonic activity, or climate). Compare AUTOCYCLIC MECHANISMS.

**allogenic** Applied to materials deposited by turbidity (see TURBIDITY CURRENT) or *\*mass flow*, particularly used in relation to *\*limestones* deposited by mass flow.

**alloformation** See ALLOSTRATIGRAPHIC UNITS.

**allogenic** Applied to minerals, or other components of a rock, that have been derived from pre-existing rocks and transported some distance to form part of the present unit, e.g. *\*quartz* grains in a *\*sandstone*. Compare AUTHIGENIC.

**allogenic stream** Stream originating outside a particular area and whose continuation is inconsistent with its new surroundings. Type examples are the Nile and Indus, whose discharges are sufficient to carry them through arid regions, and the European Neretva, which is large enough to pass over permeable limestone.

**allogroup** See ALLOSTRATIGRAPHIC UNITS.

**allomember** See ALLOSTRATIGRAPHIC UNITS.

**allometry** Differential rate of growth such that the size of one part (or more) of the body changes in proportion to another part, or to the whole body, but at a constant exponential rate. For example, the antlers of the extinct Irish elk (*Megaloceros giganteus*), the largest of all deer, grew 2.5 times faster than the rest of its body to reach an adult span of up to 3.5 m in the largest individuals. Allometry may in other cases be negative, leading to comparatively smaller parts.

**allopatric speciation** Formation of new \*species from the ancestral species as a result of the geographic separation or fragmentation of the breeding population. Genetic divergence in the newly isolated daughter populations ultimately leads to new species; divergence may be gradual or, according to punctuationist models, very rapid. See also PUNCTUATED EQUILIBRIUM.

**allopatry** The occurrence of \*species in different geographic regions. When closely related species are separated, differences between them that minimized their competition for food, shelter, or other resources usually decrease (i.e. the characteristics converge). The process is called character displacement and may be morphological or ecological.

**allophane (kandite)** \*Clay mineral of the \*kaolinite group,  $\text{Al}_2\text{Si}_2\text{O}_5(\text{OH})_2$ ; whitish; amorphous, non-crystalline; occurs along \*faults or \*joint planes in a variety of rocks.

**allostratigraphic units** Allogroups, alloformations, and allomembers; these are subdivisions of sedimentary structures that are the subject of \*allostratigraphy.

**allostratigraphy** The study of sedimentary strata that can be defined and identified from the discontinuities bounding them, and that can be mapped.

**allotriomorphic** See ANHEDRAL.

**allotrope** See ALLOTROPY.

**allotropy** The property of some chemical elements existing in two or more forms, called allotropes. For example, dioxygen ( $\text{O}_2$ ) and ozone ( $\text{O}_3$ ) are allotropes of oxygen.

**allowable bearing pressure** The *\*bearing* capacity that takes account of both the weight of the built structure and the strength of the geological structure beneath the foundation to ensure that geological deformation and the settlement of the building remain within limits the building can tolerate. The allowable bearing capacity ( $q_a$ ) is always equal to or less than the *\*safe* bearing capacity ( $q_s$ ).

**alluvial** Applied to the environments, action, and products of rivers or streams. Alluvial deposits (alluvium) are *\*clastic*, *\*detrital* materials transported by a stream or river and deposited as the river floodplain. The term is also applied to surface flow, as in *\*alluvial fans*, *\*bajadas*, etc.

**alluvial cone** See ALLUVIAL FAN.

**alluvial fan (alluvial cone)** Mass of sediment deposited at some point along a stream course at which there is a sharp decrease in gradient, e.g. between a mountain range and a plain. Essentially, a fan is the terrestrial equivalent of a river-delta formation.

**alluvium** An *\*alluvial* deposit.

**almandine** Member of the *\*garnet* group of *\*minerals*,  $\text{Fe}_3\text{Al}_2(\text{SiO}_4)_3$ ; sp. gr. 4.25; *\*hardness* 6.5–7.5; *\*cubic*; red, brown-red, or black; greasy to vitreous *\*lustre*; most common *\*crystals* are dodecahedra, and many are irregular grains; widely distributed in *\*metamorphic* and *\*igneous* rocks, and in beach *\*sands* and *\*placers*. Transparent crystals are used as *\*gemstones*, and the mineral is useful in general as an abrasive.

**alnöite** An *\*intrusive*, *\*basic*, *\*igneous*, *\*carbonatite* rock, distinctive in possessing primary *\*calcite*, and consisting of *\*melilite* ( $1/3$ ); *\*biotite* ( $1/3$ ); and *\*pyroxene*, calcite, and *\*olivine* ( $1/3$ ). *\*Feldspar* is not present in the rock, its place being taken by the mineral melilite which has the general formula:  $\text{X}_2\text{YZ}_2\text{O}_7$ ; where X = Ca, Na; Y = Mg, Al; Z = Si, Al. The type location for this rock is Alnö island off the coast of Sweden.

**ALOS** See ADVANCED LAND OBSERVING SATELLITE.

**alpha decay** Certain radionuclides (radioactive *\*nuclides*) decay by the spontaneous emission of alpha particles from their nuclei. The alpha particle is composed of two protons and two neutrons and has a charge of

+2. It also has an appreciable mass and its ejection from the nuclide creates a certain amount of recoil energy in the nucleus. The total energy ( $E_x$ ) created by alpha decay is, therefore, the sum of the kinetic energy of the particle, the recoil energy given to the new *\*nucleus* and the total energy of any emitted *\*gamma rays*. See also RADIOACTIVE DECAY.

**alpha diversity** Diversity among members of a species within a single population.

**alpha-mesohaline water** See HALINITY.

**alpha-proton-X-ray spectrometer (APXS)** A set of instruments carried on Russian *\*Vega* and *\*Phobos* missions and by *Sojourner*, the rover vehicle carried on the 1997 *\*Mars Pathfinder* mission, that measures the elemental chemistry of surface materials. The sensor head of the instrument contains curium, as a source of alpha particles, an alpha particle detector, a proton detector, and an X-ray detector. The head is placed in contact with a sample and remains there for 10 hours. Alpha particles of known energy bombard the sample. Scattered alpha particles, protons from alpha-proton reactions, and X-rays produced by excitation by the alpha particles of the atomic structure of the sample are measured by the detectors. The energy spectrum of detections by all three instruments is then recorded and transmitted to Earth.

**alpine glow** At sunset, beginning as the Sun nears the horizon, mountains exposed to direct sunlight in the east, particularly if snow-covered, assume a series of colours changing from yellow-orange to a rosy pink, which finally becomes purplish. The same series of colours in reverse order is seen on mountains in the west at sunrise.

**Alpine-Himalayan orogeny** Period of mountain building that affected both northern and southern margins of the ancient *\*Tethyan* ocean. It began in the *\*Triassic*, but reached its high point during the Late *\*Oligocene* and *\*Miocene*. The Alps are an obvious testament to this orogeny, while the gentle folds of northern France, and the Weald and London Basin in England, reflect its outer effects.

**Alportian** The final stage (324.5–318.1 Ma ago) of the *\*Serpukhovian* epoch, underlain by the *\*Chokierian*.

**alteration** A change produced in a rock by chemical or physical action.

**alteration halo** A border of minerals produced by *\*hydrothermal* *\*alteration* in the rock surrounding a *\*vein*.

**alternating current** The current output, with a sinusoidal wave-form, from an alternator or dynamo.

**alternating-magnetic-field demagnetization (AF demagnetization, thermal cleaning)** A common method for demagnetizing (see DEMAGNETIZATION) rock samples that is widely used in *\*palaeomagnetism* and *\*archaeomagnetism* because of its simplicity and because it produces no chemical change in the samples. It can cause problems associated with *\*anhysteretic* and *\*rotational* remanences, and is only fully suitable for *\*magnetite*-bearing rock samples.

**altiplanation** Process of relief reduction or planation (i.e. the smoothing of the surface) under periglacial conditions. Two mechanisms are involved: destruction of upstanding relief features by *\*gelifraction* or *\*nivation*, and accumulation of debris in depressions or as terraces. In many areas only partial altiplanation has been achieved, with the emergence of altiplanation terraces, such as those of Cox Tor on Dartmoor, England.

**altocumulus** From the Latin *altum* (height) and *cumulus* (heap). A genus of cloud composed largely of water droplets, and consisting of grey-white sheets, or banded layers and rolls, which may also be broken up into cells. Sometimes it has a banded appearance, occasionally giving a mackerel-sky effect; this is probably associated with strong vertical wind shear in middle altitudes. See also CLOUD CLASSIFICATION.

**Altonian** A *\*stage* (17.5–16.5 Ma ago) in the Upper *\*Tertiary* of New Zealand, underlain by the *\*Otaian*, overlain by the *\*Cliffdenian*, and roughly contemporaneous with the upper *\*Burdigalian* stage.

**altostratus** From the Latin *altum* (height) and *stratus* (spread out). A genus of cloud consisting of greyish sheets or layers; the cloud may be striated, fibrous, or uniform. It may be composed of ice crystals as well as water droplets. See also CLOUD CLASSIFICATION.

**alumstone** See ALUNITE.

**alunite (alumstone)** *\*Mineral*,  $\text{KAl}_3(\text{SO}_4)_2(\text{OH})_6$ ; sp. gr. 2.6–2.8; *\*hardness* 3.5–4.0; *\*trigonal*; white, sometimes grey to reddish; white

**\*streak**; **\*vitreous lustre**; **\*crystals** rare, **\*habit** **\*massive**; **\*cleavage** basal {0001}, distinct; **\*fracture** uneven, **\*conchoidal**; slightly astringent taste. Occurs as a **\*secondary** mineral in volcanic rocks containing potassic **\*feldspars** altered by sulphuric-acid solutions. It is difficult to distinguish from **\*dolomite**, **\*anhydrite**, and **\*magnesite**.

**alveolus** See BELEMNITIDA.

**ALVIN** A submersible that enables two scientists to collect data and observations at an ocean depth of up to 4500 m for dives lasting up to 10 hours. It was commissioned in 1964 and has been updated several times; it is operated by the Woods Hole Oceanographic Institution. It is named after the WHOI engineer Allyn Vine.

**A/m** See AMPERES PER METRE.

**Amalthea (Jupiter V)** The jovian satellite with the closest orbit to Jupiter. Its surface colour is reddish, apparently because of sulphur emitted from **\*Io**. Its diameter is 189 km (262 × 146 × 134 km), the irregular shape suggesting a rigid body. Its mass is  $7.17 \times 10^{18}$  kg; mean distance from Jupiter 181 000 km. It radiates more heat than it receives from the Sun.

**Amarassian** See CHANGHSINGIAN; KAZANIAN.

**Amasia** The name of a proposed **\*supercontinent** that is predicted to form approximately 100 million years from today through the joining of Eurasia to N. America and Arabia, Arabia to Africa, and S. America to N. America, with the disappearance of the Arctic Ocean and Caribbean Sea.

**Amazonian** A division of **\*areological** time, lasting from 1.80 Gy to the present in the Hartmann–Tanaka Model and 3.55 Gy to the present in the Neukum–Wise Model, and divided into three epochs: Lower Amazonian (1.80–0.70 or 3.55–2.50 Gy); Middle Amazonian (0.70–0.25 or 2.50–0.70 Gy); and Upper Amazonian (0.25–0.00 or 0.70–0.00 Gy).

**amazonite** See ALKALI FELDSPAR.

**amazonstone** See ALKALI FELDSPAR.

**amb** See AMBULACRUM.

**amber** Fossil conifer resin which is brittle and hard, translucent to transparent, and yellow to brown in colour. It is found in **\*sediments** or on the shore and takes a fine polish.

**ambient pressure** Atmospheric pressure in the surrounding air.


**ambient temperature** The dry-bulb temperature prevailing in the surrounding air.

**ambitus** The outline or edge of an echinoid (**\*Echinoidea**) when seen from above or below. Usually it is the place where the **\*plates** of the **\*test** are at their widest.

**ambulacral** Applied to those areas of the body of an **\*echinoderm** that bear.

**ambulacral groove** See **AMBULACRUM**.

**ambulacrum (amb)** In **\*Echinodermata**, an area of the body surface (covered in most classes by calcitic (see **CALCITE**) **\*plates**), that overlies one of the radial canals of the internal water vascular system, and bears the **\*tube** feet. In some echinoderms, e.g. Asteroidea, Blastoidea, and **\*Crinoidea**, the ambulacrum is marked by a deep linear depression, the ambulacral groove. Typically, echinoderms have five ambulacral areas, or a multiple of five. See **ECHINOIDEA**.



### Ambulacrum

***Ambulocetus natans*** The most completely known early cetacean, described in 1994 by J. G. M. Thewissen, S. T. Hussain, and M. Arif, from Lower to Middle *Eocene* beds in Pakistan. It is known by parts from most of the skeleton, showing that it had a long neck, relatively long hind limbs, and five separate (hoofed) digits on each limb. It was the size of a sea lion.

**amensalism** An interaction of species populations, in which one population is inhibited while the other (the amensal) is unaffected.

**American Geophysical Union (AGU)** A society with approximately 35 000 members that was founded in 1919 and is based in Washington, DC. It is the leading academic body for geophysicists.

**American Institute of Professional Geologists (AIPG)** The body that certifies professional geologists in the United States, based on their competence, integrity, and ethical standards. The AIPG presents testimony and position papers to legislators and agencies at federal and state level on matters pertaining to the interests and employment opportunities of geologists. Through its contacts with equivalent bodies in other countries, the AIPG provides American geologists with access to professional registration in those countries.

**American Province** See PACIFIC PROVINCE.

**Amersfoort** An *\*interstadial* in the last *\*glaciation* of the Netherlands during the early *\*Devensian* (somewhere between 70 000 and 60 000 Ma ago). The July temperature (based on floral evidence) was perhaps 15–20 °C.

**amesite** See CHAMOSITE.

**amethyst** See QUARTZ.

**Amgan** A Russian-Kazakhstanian *\*stage* (513–502 Ma ago) of the Middle *\*Cambrian* *\*epoch*, preceded by the *\*Toyonian* and followed by the *\*Mayan*.

**amino acid** Organic compound containing an acidic carboxyl (COOH) group and a basic amino (NH<sub>2</sub>) group. They constitute the fundamental building blocks of peptides and proteins and are classified either (*a*) as neutral, basic, or acidic (according to their *\*pH*), or (*b*) as non-polar, polar, or charged (according to their electrical configuration).

**amino group** The chemical group –NH<sub>2</sub>.

**ammonites** Family of *\*Ammonoidea*.

**Ammonoidea (ammonoids)** (phylum *\*Mollusca*, class *\*Cephalopoda*) Subclass of cephalopods which generally have *\*planispirally* coiled, septate shells (see SEPTUM). Characteristically the shells are tightly coiled and planispiral, although some are coiled loosely or spirally; the *\*protoconch* is globular; the shells may be either *\*involute* or *\*evolute*. Some forms have marked *\*ventral* *\*keels*; ribs and nodes may also be present. The *\*siphuncle* is variable but mainly ventral in position. *\*Sutures* are often very complex. *\*Camaral* deposits are absent. The Ammonoidea were probably tetrabranchiate (four-gilled) cephalopods. They constitute the largest cephalopod subclass, with 163 families including the ammonites, in which the suture lines form very complex patterns; the ceratites, in which part of the suture line is frilled; and the goniatites, with relatively simple suture lines. They range in age from *\*Devonian* to Upper *\*Cretaceous*. All members are now extinct. See also APERTURE; APTYCHUS; FORAMEN; PHRAGMOCONE; VENTER.

**ammonoids** See AMMONOIDEA.

**amnion** See AMNIOTIC.

**amniotic** Applied to a type of development typical of higher *\*vertebrates* (*\*reptiles*, *\*birds*, and *\*mammals*), in which the amnion (a protective membrane) surrounds the embryo in a bag of (amniotic) fluid. Evolutionarily, the amnion is *\*primitively* associated with a shell and is capable of gaseous exchange; its development thus enabled eggs to be laid on dry land for the first time in vertebrate evolution. Compare ANAMNIOTIC.

**Amontons' laws of friction** Leonardo da Vinci (1452–1519) investigated the phenomenon of *\*friction* and the conclusions he reached were eventually discovered and formalized by the French physicist and inventor Guillaume Amontons (1663–1705) as two laws: **1.** The force of friction is directly proportional to the applied load; **2.** The force of friction is independent of the apparent area of contact (i.e. if two bodies of equal mass and made from similar material are both in contact with the same third body, the same amount of force will be required to move them across the surface of the third body regardless of differences in the area of contact).

**amorphous clay** *\*Clay* that is not composed of *\*crystals*. It occurs most commonly in volcanic ash and forms during the early stages of *\*weathering* in soils derived from volcanic material.

**amorphous cloud** Continuous cover of low, featureless cloud (e.g. *\*nimbostratus*), often producing rain.

**amosite** See ANTHOPHYLLITE; ASBESTOS.

**AMPERE** See ACTIVE MAGNETOSPHERE AND PLANETARY ELECTRODYNAMICS RESPONSE EXPERIMENT.

**amperes per metre (A/m)** The SI unit of magnetic moment per unit volume.  $1 \text{ A/m} = 10^3 \text{ gauss}$ .

**Ampferer subduction** See A-SUBDUCTION.

**Amphibia (amphibians)** Class that appeared first in the *\*Devonian*, having evolved from rhipidistian (lobe-finned) fish (see RHIPIDISTIA). They flourished in the *\*Carboniferous* and *\*Permian*. During the *\*Triassic* some

forms, e.g. Mastodontosaurus, grew to 6 m long, and the first modern types were established. Today the amphibians are represented by just three groups, of which the Urodela (salamanders) and Anura (frogs and toads) are the best known (the third group, the caecilians (Apoda), are worm-like and burrowing). Most amphibians are found in damp environments and they occur on all continents except Antarctica.

**amphibians** See AMPHIBIA.

**amphiboles** A group of \*minerals possessing double chains of silicon-oxygen  $[\text{SiO}_4]$  tetrahedra with a composition of  $[\text{Si}_4\text{O}_{11}]_n$  running parallel to the \*crystallographic axis; i.e. parallel to the \*prism zone of a crystal. The double chains are held together by monovalent, divalent, or trivalent cations, of which  $\text{Na}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Fe}^{2+}$ ,  $\text{Al}^{3+}$ , and  $\text{Fe}^{3+}$  are the most important; hydroxyl ions also occur. There are three main groups of amphibole minerals: (a) calcium-poor amphiboles with the general formula  $\text{X}_2\text{Y}_5[\text{Z}_4\text{O}_{11}]_2(\text{OH},\text{F})_2$ , where  $\text{X} = \text{Mg}$  or  $\text{Fe}^{2+}$ ,  $\text{Y} = \text{Mg}$ ,  $\text{Fe}^{2+}$ ,  $\text{Fe}^{3+}$ ,  $\text{Al}^{3+}$ , etc., and  $\text{Z} = \text{Si}$  or  $\text{Al}$ ; (b) the calcium-rich amphiboles with the general formula  $\text{AX}_2\text{Y}_5[\text{Z}_4\text{O}_{11}]_2(\text{OH},\text{F})_2$ , where  $\text{A} = \text{Na}$ ,  $\text{X} = \text{Ca}$ ,  $\text{Y} = \text{Mg}$ ,  $\text{Fe}^{2+}$ ,  $\text{Fe}^{3+}$ ,  $\text{Al}$ , etc., and  $\text{Z} = \text{Si}$  or  $\text{Al}$ ; and (c) the alkali amphiboles in which  $\text{Na} > \text{Ca}$  and with the general formula  $\text{AX}_2\text{Y}_5[\text{Z}_4\text{O}_{11}]_2(\text{OH},\text{F})_2$ , where  $\text{A} = \text{Na}$  or  $\text{K}$ ,  $\text{X} = \text{Na}$  (or  $\text{Na}$  and  $\text{Ca}$ ),  $\text{Y} = \text{Mg}$ ,  $\text{Fe}^{2+}$ ,  $\text{Fe}^{3+}$ ,  $\text{Al}$ , etc., and  $\text{Z} = \text{Si}$  or  $\text{Al}$ . Calcium-poor amphiboles include the \*orthorhombic amphiboles (called the orthoamphiboles) and include \*anthophyllite and gedrite, but the other two groups are \*monoclinic and include the common \*hornblendes \*tremolite and \*actinolite, as well as the sodium-rich varieties such as \*glaucophane and \*riebeckite. Amphiboles are common rock-forming silicate minerals that occur in \*intermediate and \*alkaline \*igneous rocks and also in many regional \*metamorphic rock types.

**amphibolite** A medium-grained, dark-coloured, regional \*metamorphic rock composed of \*hornblende and \*plagioclase with minor \*epidote, \*sphene, \*biotite, and \*quartz. The rock may show a well-developed planar or linear alignment of elongate hornblende crystals as a result of suffering deformation at the same time as \*regional metamorphism. These alignments define \*fabrics within the rock known as \*schistosity and \*lineation respectively. Amphibolites are formed by medium-grade

\*metamorphism of \*basic \*igneous rocks such as \*basalts, both \*extrusive and \*intrusive types.

**amphibolite facies** A set of metamorphic \*mineral assemblages produced by the \*metamorphism of a wide range of starting rock types under the same metamorphic conditions and typically characterized by the development of the mineral assemblage andesine (\*plagioclase)–\*hornblende in rocks of \*basic \*igneous composition. Other rocks of contrasting composition, e.g. \*shales or \*limestones, would each develop their own specific mineral assemblage, even though they are all being metamorphosed under the same conditions. The variation of mineral assemblage with starting rock composition reflects a particular range of pressure, temperature, and  $P(H_2O)$  conditions. Experimental studies of mineral  $P$ – $T$  stability fields indicate that the \*facies represents a range of moderate-pressure, moderate-to high-temperature conditions.

**amphicoelous** Applied to the condition in which the central part of each \*vertebra (the centrum) is concave on both anterior and posterior faces. *Compare* HETEROCOELUS.

**Amphicyonidae** See CARNIVORA.

**amphidetic** Applied to a bivalve (\*Bivalvia) ligament located around the \*umbones in both anterior and posterior areas. *Compare* OPISTHODETIC.

**amphidromic point** The centre of an amphidromic or tidal system; a no-tide or nodal point around which the crest of a standing wave or a high-water level rotates once in each tidal cycle. The tidal range increases progressively with increasing distance from the central point. The high water rotates anticlockwise around the central point in the northern hemisphere and clockwise in the southern hemisphere.

**Amphineura** (phylum \*Mollusca) Class of elongate, bilaterally symmetrical, marine molluscs, first appearing in the Upper \*Cambrian, in which the shell, if present, consists of seven or eight overlapping, calcareous plates on the dorsal surface. The class comprises the \*Polyplacophora (chitons), and the Aplacophora which have no fossil record.

**amphitheatre** Flat-topped, steep-walled depression, shaped like a horseshoe and resembling an ancient Greek theatre. It may be the result of glacial erosion, forming a **\*cirque**, or the collapse of a volcano. In the case of Mount St Helens, the catastrophic collapse of the summit and northern slopes of the cone and the subsequent **\*pyroclastic flow** eruption on 18 May 1980 created a north-facing amphitheatre enclosing a small active vent, later occupied by a small **\*dacite** dome.

**amplitude** 1. Of a **\*fold**, the height of the maximum displacement of a folded layer measured from a median trace which passes through the **\*inflexion** points of adjacent limbs. 2. Of a wave, the distance between the highest point of the wave and the position of zero displacement.

**ampulla** An inflatable sac at the end of each **\*tube foot** in the echinoids (**\*Echinoidea**). It controls water pressure in the tube foot when the muscles in its walls are contracted.

**amygdale** (**amygdule**; *adj.* **amygdaloidal**) Spheroidal, ellipsoidal, or (literally) ‘almond-shaped’ cavity or **\*vesicle** within a **\*lava**, filled with secondary minerals, e.g. **\*calcite**, **\*quartz**, or **\*zeolites**.

**amygdaloidal** See **AMYGDALE**.

**amygdule** See **AMYGDALE**.

**An** See **ALKALI FELDSPAR**.

**anabatic wind** A wind that blows up a slope, often gently, and usually when the sloping ground surface has been warmed by the Sun. Compare **KATABATIC WIND**.

**anabranching channel** A type of **\*distributary** river channel that separates from its trunk stream and may flow parallel to it for several kilometres before rejoining it. The anabranching channel remains undivided, and so differs from an anastomosing channel which has major distributaries that branch and then rejoin it.

**anacline** A common condition of the **\*interarea** in a strophic brachiopod (**\*Brachiopoda**) shell. It describes the situation where the interarea is at an angle of less than 90° to the plane of **\*commisure**. Compare **APSACLINE**.

**anaerobic** **1.** Of an environment: one in which air (oxygen) is absent. In the case of a depositional environment, one with 0.0–0.1 ml of dissolved oxygen per litre of water. *Compare* AEROBIC; DYSAEROBIC. **2.** Of an organism: one able to grow only in the absence of oxygen, i.e. an anaerobe. **3.** Of a process: one that can occur only in the absence of oxygen.

**anaerobic ammonium oxidation** *See* ANAMMOX.

**anafront** Front (warm or cold) at which there is upward movement of the warm-sector air, commonly producing clouds and precipitation. *Compare* KATAFRONT.

**anagenesis** In the original sense, refers to evolutionary advance. Now it is often applied more widely, to virtually all sorts of evolutionary change, along a single, unbranching lineage.

**analcime (analcite)** \*Mineral,  $\text{Na}(\text{AlSi}_2\text{O}_6) \text{H}_2\text{O}$ ; sp. gr. 2.2–2.3; \*hardness 5.0–5.5; \*cubic; can be colourless or white, with grey, red, and greenish tints, sometimes pigmented with iron oxides; \*vitreous \*lustre; \*crystals polyhedra and tetragonal trioctahedra, also found in granular aggregates; \*cleavage poor, cubic; occurs in rocks of low-temperature \*igneous origin that are rich in soda and water (e.g. \*nepheline-syenites), and also volcanic \*tuffs.

**analcite** *See* ANALCIME.

**analcite-basanite** *See* BASANITE.

**analcitite** An \*igneous rock in which the abundant \*plagioclases have been replaced by the \*mineral \*analcime during late-stage deuteric activity (*see* DEUTERIC REACTION). \*Magma-derived, late-stage aqueous fluids penetrate along \*cleavages and \*fractures in primary plagioclase within the crystalline rock body and react with it, converting the plagioclase to analcime which is stable in the presence of a late-stage fluid.

**analog data (analogue data)** Data which are recorded continuously, as opposed to digital data which are recorded by discrete (digital) sampling.


**analogous structures** Features with similar functions which have developed independently in unrelated taxonomic groups, in response to a

similar way of life, or similar method of locomotion, or similar food source, etc. Thus the wings of birds and insects are analogous.

**analogue image** In *\*remote* sensing, an image in which continuous variation in the scene being sensed is represented by continuous variation in image tone, such as the image produced from photosensitive chemicals in a photographic film. *Compare* **DIGITAL IMAGE**.

**analyser** A piece of *\*Polaroid* in a transmitted- or *\*reflected-light microscope* which may be inserted into the light path above the *\*mineral* section. When the analyser is out of the path of light through the microscope observations are made in *\*plane-polarized light (PPL)*; when the analyser is in the path of light, observations are made in crossed polarized light, or *\*crossed* polars (crossed nicols, xpols, or XPL). *See* **NICOL PRISM**.

**anammox (anaerobic ammonium oxidation)** A process in which anaerobic bacteria oxidize nitrite (NO<sub>2</sub>) and ammonium (NH<sub>4</sub>) to dinitrogen (N<sub>2</sub>):




This occurs in wetland soils but mainly in freshwater and marine ecosystems, where it accounts for up to 67% of N<sub>2</sub> production and is therefore an important component of the nitrogen cycle. *Compare* **FEAMMOX**.

**anamniotic** Applied to a type of development typical of lower *\*vertebrates* (*\*fish* and *\*amphibians*). The egg lacks a shell and protective embryonic membranes: consequently it must be laid in water or in a suitably damp environment. *Compare* **AMNIOTIC**.

**Ananke (Jupiter XII)** One of the lesser satellites of *\*Jupiter*, with a diameter of 20 km; its orbit is *\*retrograde*.

**Anapsida** *\*Clade* of *\*reptiles* characterized by a skull that lacks apertures in the temple regions behind the eyes. This was the condition in the earliest reptiles, the *\*Cotylosauria* or 'stem reptiles'. Four groups flourished in the late *\*Palaeozoic*: the *\*Captorhinomorpha*; the *\*Mesosauria*; the

\*Pareiasauridae; and the \*Procolophonia, which survived into the \*Triassic. The only living representatives are the turtles (Chelonia).



### Anapsida

**Anaspida (anaspids)** \ (superclass \*Agnatha) Extinct order of fish-like \*vertebrates. The body, including the head region, was covered in dermal scales (see **PLACOID SCALE**), and was less flattened than in the \*Osteostraci. The tail was hypocercal (tilting downwards). Anaspids were small (up to 15 cm in length). Genera included Jaymoytius, Pharyngolepis, and Pterygolepis. Anaspids ranged from the Late \*Silurian to the Late \*Devonian.

**anastomosing channel** See ANABRANCHING CHANNEL.

**anatase (octahedrite)** \*Mineral,  $\text{TiO}_2$ ; sp. gr. 3.9, \*hardness 5.5–6.0; \*tetragonal; yellow, or shades of brown, blue, or black; white \*streak; \*vitreous \*lustre; \*crystals normally \*bipyramidal octahedra, but sometimes \*tabular; \*cleavage perfect, basal, and \*pyramidal; occurs as an \*accessory mineral in \*metamorphic and \*igneous rocks, also in \*hydrothermal veins and \*granite \*pegmatites.

**anatexis** The partial or incomplete melting of a rock in response to an increase in temperature (at constant pressure), or a drop in pressure (at constant temperature). Melting takes place along grain boundaries, and the \*melt can either be extracted from the partially molten rock system, or can remain within the system. Typical examples of anatexis would be the

generation of granitic melts (see [GRANITE](#)) by partially melting aluminous crustal rocks, and the generation of [\\*basalts](#) by partially melting [\\*mantle](#) [\\*peridotite](#).

**[Anatolepis heintzi](#)** Possibly one of the earliest jawless fish, known from a number of associated scales and fragments discovered in Lower [\\*Ordovician](#) ([\\*Arenig](#)) strata at Spitzbergen.

**[anchialine](#)** Applied to flooded caves lying deep below the ground surface. They form in limestone environments and from volcanic tubes.

**[anchimetamorphism](#)** A [\\*metamorphic grade](#) in sedimentary rocks where changes due to [\\*diagenesis](#) are overtaken by the very earliest phases of [\\*metamorphism](#).

**[anchor](#)** 1. Rock anchors are long bolts or cables with one end [\\*grouted](#) into a [\\*drill](#) hole and with a plate and nut on the exposed end. These can carry considerable [\\*loads](#), although slow [\\*failure](#) of the rock will lessen the support. 2. Soil anchors may be used in sediments where the material is strong enough to provide sufficient reaction to the load. Holes must be drilled and the anchor installed and grouted quickly, as soil around the hole may crumble and reduce the strength of the bond.

**[ancient biomolecule](#)** Any molecule produced by an organism during its lifetime which persists post mortem. Identification and characterization of an ancient biomolecule may allow inferences to be drawn about the organism in which it was produced, i.e. the type of organism or evolutionary relationships. The types of molecule studied include nucleic acids, lipids, proteins, and complex polysaccharides such as lignin.

**[Ancient Cratered Terrain](#)** See [MARTIAN TERRAIN UNITS](#).

**[andalusite](#) ([chiastolite](#))** A member of the  $\text{Al}_2\text{SiO}_5$  (or  $\text{Al}_2\text{O}_3 \cdot \text{SiO}_2$ ) group of [\\*polymorphic](#) minerals; the other members of this group are [\\*kyanite](#) and [\\*sillimanite](#). The mineral mullite ( $3\text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2$ ) is related to this group of minerals; sp. gr. 3.2; [\\*hardness](#) 6.5–7.5; [\\*orthorhombic](#); pink or red, sometimes grey-brown and green, transparent varieties used as [\\*gemstones](#); [\\*vitreous](#) [\\*lustre](#); [\\*crystals](#) [\\*prismatic](#) with a pseudo-tetragonal form which has a characteristic cross-section, can also be [\\*massive](#); [\\*cleavage](#) at right angles, distinct {110}; occurs principally in

thermally *\*metamorphosed* rocks that show alteration to aggregates of white *\*mica*, in low-grade, regionally metamorphosed rocks, and also in *\*granites* and *\*pegmatites* together with *\*corundum*, *\*tourmaline*, *\*topaz*, and other minerals. The name is derived from Andalusia in Spain.

**ANDE** See *ATMOSPHERIC NEUTRAL DENSITY EXPERIMENT*.

**Andean orogenic belt** Part of the active Cordillera Mountain Belt on the eastern margin of the Pacific Ocean, the Andes extend approximately 10 000 km from the Caribbean Sea to the Scotia Sea. They developed during the *\*Mesozoic* and *\*Cenozoic* eras, largely as a result of the *\*subduction* of normal oceanic *\*lithosphere*. Unlike the N. American Cordillera, very few *\*allochthonous* *\*terrane*s have been recognized, and there have been no collisions of major continental masses as in the Himalayas. These differences have led to the concept of a distinctive 'Andean-type orogeny'.

**andesine** See *PLAGIOCLASE FELDSPAR*.

**andesite** Fine-grained volcanic rock (named after the Andes mountains) characterized by the presence of *\*plagioclase* feldspars of the *\*oligoclase*–andesine end of the range, and of some combination of *\*augite*, *\*orthopyroxene*, and *\*hornblende*. In chemical and mineralogical terms andesites are similar to *\*diorites*. The feldspars frequently occur as strongly zoned *\*phenocrysts*. Glassy andesites are rare. An increase in the *\*quartz* percentage marks the gradation of andesites into *\*dacites*. An increase in the percentages of *\*alkali* feldspars marks the change from andesites into trachyandesites, some of which (e.g. mugearite and hawaiite) exhibit basaltic features, including a higher *\*olivine* content. See *BASALT*.

**andhis** Local term, used in the north-west of the Indian subcontinent, for dust storms accompanying violent squalls, caused by strong convection.

**andic horizon** A *\*soil horizon* containing weathered *\*pyroclastic* deposits.

**Andino-type margin** See *ACTIVE MARGIN*.

**andisols** In the US *\*soil taxonomy*, an order of soils that are derived from volcanic ash; they contain glass and *\*amorphous clay*.

**andosols** A reference soil group in the *\*World Reference Base for Soil Resources*, comprising volcanic soils with a *\*vitric* horizon at a depth of

more than 30 cm or an **\*andic** horizon within 25 cm of the surface.

**andradite** Member of the **\*garnet** group of **\*minerals**,  $\text{Ca}_3\text{Fe}_2(\text{SiO}_4)_3$ ; sp. gr. 3.75; **\*hardness** 6.5–7.5; **\*cubic**; yellow, greenish, or brownish-red, to black; greasy to **\*vitreous** **\*lustre**; most **\*crystals** dodecahedral; occurs widely in **\*metamorphic rock** and **\*igneous** rocks often as a result of **\*metasomatism**, also in beach **\*sands** and **\*alluvial** **\*placers**. Transparent crystals are used as **\*gemstones**, and the mineral can be used as an abrasive.

**anelastic** Applied to a material in which there is a temporal shift between the application of **\*stress** and the resulting **\*strain**. If the stress is periodic, there will be a phase shift between stress and strain in an anelastic material.

**anemometer** An instrument for measuring the speed of flow in a fluid. Meteorological anemometers measure wind speed. Most do so either from the rotation of a wind-driven turbine ('rotating cups anemometer') or from the wind pressure through a tube aligned by a vane to point into the wind ('pressure-tube anemometer').

**anemometry** The measurement of the speed of flow in a fluid.

**aneroid barometer** Instrument for measuring atmospheric pressure as represented by the height (or thickness) of a sealed, flexible, metal, concertina-shaped drum from which air has been partially evacuated. Variations in the height of the drum are transmitted to a pressure scale either mechanically, through a series of levers and a pointer, or electronically.

**Angara** A continental mass of Asia, China, and the Far East that existed during the **\*Palaeozoic**. In the latest Palaeozoic it became joined to **\*Euramerica** along the line of the present Ural Mountains.

**Angaraland** Name given by the Austrian geologist **Eduard \*Suess** (1831–1914) to a small **\*shield** in north-central Siberia, where **\*Precambrian** rocks are exposed, and which was considered to be the nucleus for subsequently developed structural features in Asia.

**angiophyte** A plant belonging to the evolutionary line leading to the **\*angiosperms**. Angiophytes may have begun to diversify in the Late **\*Triassic**, become rarer during the **\*Jurassic**, but radiated vigorously in the Early **\*Cretaceous**.

**angiosperm** A flowering seed plant (*\*Spermatophyta*), distinguished by producing seeds that are enclosed fully by fruits. Angiosperms are the most highly evolved of plants, and appear first in rocks of Lower *\*Cretaceous* age.

**Angiospermae** In earlier classification, a division of the *\*Spermophyta* (seed plants) containing all the flowering plants (*\*angiosperms*). Modern classifications regard the angiosperms as a *\*paraphyletic* group.

**angle of draw** The angle between the end of an underground working and the point on the ground surface to which *\*subsidence*, due to that working, may extend. Draw usually proceeds at an angle of 65–75° to the horizontal.

**angle of incidence (incident angle)** The angle between an incident ray or wave and the normal to a reflecting or refracting surface. The angle of incidence is related to the *\*angle of refraction* by *\*Snell's law*. See also CRITICAL ANGLE.

**angle of internal friction (friction angle)** A measure of the ability of a unit of rock or soil to withstand a *\*shear* stress. It is the angle ( $\phi$ ), measured between the normal force ( $N$ ) and resultant force ( $R$ ), that is attained when failure just occurs in response to a shearing stress ( $S$ ). Its tangent ( $S/N$ ) is the coefficient of sliding friction. Its value is determined experimentally.

**angle of reflection** The angle between a reflected ray or wave and the normal to a reflecting surface.

**angle of refraction** The angle between a refracted ray or wave and the normal to a refracting surface. The angle of refraction is related to the *\*angle of incidence* by *\*Snell's law*. A ray or wave entering a medium in which its speed is higher than in the medium it is leaving will deviate away from the normal, whereas when it enters a medium of lower speed it will deviate towards the normal.

**angle of repose** Maximum angle at which unconsolidated material can stand. Various factors determine this angle, including particle size and angularity, the degree of interlocking between particles, and *\*pore-water pressure*. A typical angle of repose for coarse scree is 32–36°.

**angle of shearing resistance (internal angle of friction, angle of frictional resistance;  $\phi$ )** Approximate angle of repose for clean *\*sand*; it

reduces with moisture content and is zero for a sheared, saturated **\*clay**. For rocks, a rough determination of the angle may be made by placing two small blocks of material together and inclining them until the top one slides.

**anglesite** **\*Secondary** lead mineral,  $\text{PbSO}_4$ ; sp. gr. 6.3–6.4; **\*hardness** 3; **\*orthorhombic**; normally colourless to white, but sometimes yellow, grey, or coloured by a bluish tinge; white **\*streak**; **\*adamantine** **\*lustre**; **\*crystals** **\*tabular**, **\*prismatic**, or pyramidal, or occurring as **\*massive**, granular, or compact aggregations; normally occurs in the oxidized zone of lead deposits, and often surrounds a core of **\*galena**.

**Anglian** 1. A glacial **\*stage** of the middle **\*Pleistocene** in Britain. 2. A middle-**\*Pleistocene**, cold-climate series of deposits. There were a number of glacial advances in East Anglia (England), which are difficult to understand. Near Lowestoft there are two **\*till** sheets separated by stratified sands. The lower, North Sea Drift, contains Scandinavian **\*erratics**, and the upper, thicker till, contains erratics of **\*Jurassic** or **\*Cretaceous** material. The highest terraces of the Thames may belong to this period and perhaps could also be correlated with the **\*Elsterian** deposits of Europe.

**ångstrom** (**Å**) Unit of length equal to  $10^{-10}$  m, used in measuring **\*electromagnetic** radiation including visible light and X-rays. Replaced in SI units by the nanometre ( $10 \text{ Å} = 1 \text{ nm}$ ).

**angularity number** See AGGREGATE TESTS.

**angular momentum** The momentum of a body rotating in a plane around a point. It is formally the product of the mass of the body, the radius of the orbit, and the square of the angular velocity ( $mrv^2$ ). Because of its rotation about its axis, the Earth has rotational angular momentum, and orbital angular momentum on account of its annual revolution around the Sun. The angular momentum of the Earth–Moon system ( $3.45 \times 10^{34} \text{ rad/kg/m}^2/\text{s}$ ) is the sum of the rotational angular momentum of the Earth and the Moon's orbital angular momentum, and is high compared to that of the other terrestrial planets.

**angular shear strain** A measure of the angular rotation from two mutually perpendicular reference axes following simple **\*shear**. The shear strain is given as the tangent of angular shear  $\psi$ , and may be positive or negative

with respective clockwise or anticlockwise rotations in relation to the reference perpendicular.

**angular unconformity** A *\*discordant* surface of contact between the deposits of two episodes of sedimentation in which the older, underlying *\*strata* have undergone folding, uplift, and *\*erosion* before the deposition of the younger *\*sediments*, so that the younger strata truncate the older. *See also UNCONFORMITY.*

**angular velocity ( $\Omega$ )** The speed of a body that is moving along a curved path, usually expressed in radians per second;  $\Omega = 2\pi/T$ , where  $T$  is the time taken to complete one revolution.

**anhedral (allotriomorphic)** A morphological term referring to grains in *\*igneous* rocks which have no regular crystalline shape. Anhedral forms are developed when a *\*crystal's* free growth in a *\*melt* is inhibited by the presence of surrounding crystals. The shape of the growing crystal is thus controlled by the arrangement and orientation of the surrounding pre-existing crystals. *Compare EUHEDRAL.*

**anhydrite** Mineral sulphate,  $\text{CaSO}_4$ , found in *\*evaporite* deposits; sp. gr. 2.9–3.0; *\*hardness* 3.0–3.5; *\*orthorhombic*; normally white to colourless, sometimes with a bluish tinge, but occasionally grey and reddish; white *\*streak*; *\*vitreous* to pearly *\*lustre*; *\*crystals* rare, the mineral being usually *\*massive*, granular, and fibrous; three *\*cleavages* at right angles, perfect {010}, good {100}, {001}; occurs with *\*gypsum* and *\*halite*, deposited directly from sea water at temperatures in excess of 42 °C, can form from the dehydration of gypsum; may occur as a *\*cap rock* above *\*salt \*domes*, and as a minor *\*gangue* mineral in *\*hydrothermal* veins. It is used as a raw material for cements.

**anhysteretic magnetization** The magnetization acquired when a direct magnetic field is applied to a specimen which is simultaneously held in an alternating magnetic field. *See ALTERNATING-MAGNETIC-FIELD DEMAGNETIZATION.*

**Animalia (Metazoa)** Multicellular organisms that develop from embryos; one of the three kingdoms of multicellular organisms (the other two being *\*Fungi* and *\*Plantae*). The kingdom includes all animals other than protozoons (some of which are colonial); *\*Porifera* (sponges) are

sometimes excluded because their structure differs markedly from that of other animals. Animals first appeared in the **\*Precambrian**, the Porifera from one kind of **\*protist** forebear, and all other animals from another (or possibly more than one other) protist. The oldest fossils are burrows of a **\*coelomate** in rocks rather less than 700 Ma old.

**Animikian** A **\*system** of the **\*Palaeoproterozoic** dated at about 2225–2050 Ma ago (Int. Commission on Stratigraphy, 2004).

**anion** A negative ion, i.e. an atom, or complex of atoms, that has gained one or more electrons and thereby carries a negative electric charge, e.g.  $\text{Cl}^-$ ,  $\text{OH}^-$ , and  $\text{SO}_4^{2-}$ . So-called because when an electric current is passed through a conducting solution the negative ions present in the solution are attracted to the anode (the positive electrode). *Compare* **CATION**.

**Anisian 1.** A Middle **\*Triassic \*age**, preceded by the Early **\*Triassic** epoch, followed by the **\*Ladinian \*stage**, and dated at 245–237 Ma ago (Int. Commission on Stratigraphy, 2004). **2.** The name of the corresponding European **\*stage**, which is roughly contemporaneous with the Guan Ling (China) and **\*Etalian** (New Zealand).

**Anisograptidae** *See* **GRAPTOLOIDEA**.

**anisometric growth** A change in the ratio between the sizes of two parts of an organism during its **\*ontogeny**, such that if an organism grows anisometrically its shape will change.

**anisomyarian** *See* **MUSCLE SCAR**.

**anisotropic 1.** Applied to substances whose optical or other physical properties vary according to the direction from which they are observed. In optical mineralogy the term is applied to **\*minerals** other than **\*cubic** that split the light passing through them into two **\*vibration** directions, each with a different velocity (provided the light is not travelling along an **\*optic axis**), as a result of the light being doubly refracted (*see* **DOUBLE REFRACTION**). In addition, each beam is refracted differently for different colours of light. **2.** In engineering geology, anisotropy refers to a rock whose engineering properties vary with direction. For example, **\*schist**, a highly anisotropic rock, has a compressive strength that varies depending

upon the orientation of the **\*foliation** to the applied load (see **COMPRESSIVE STRESS**).

**anisotropic meter** Any instrument that measures differences of particular parameters in different directions. Specific instruments measure **\*electrical conductivity**, **\*seismic waves**, and low- and high-field magnetic **\*susceptibility**.

**anisotropy** The variations in the physical properties of a medium that depend on the direction in which they are measured. *Compare* **INHOMOGENEITY**; **ISOTROPY**.

**ankerite** **\*Mineral**,  $\text{Ca}(\text{Mg,Fe})(\text{CO}_3)_2$ ; sp. gr. 2.9–3.2; **\*hardness** 3.5–4.0; **\*trigonal**; yellowish-brown, sometimes white, yellow, or grey; white **\*streak**; **\*vitreous lustre**; **\*crystals** can be rhombohedral, but ankerite also occurs **\*massive** and granular; **\*cleavage** rhombohedral, perfect {1011}, faint {0221}; occurs as a **\*gangue** mineral with iron ores, and as fillings associated with coal seams, and in an environment similar to that of **\*dolomite**. It is a form of **\*ferroan** dolomite where ferrous iron substitutes for Mg in a **\*solid solution** series from dolomite ( $\text{CaMg}(\text{CO}_3)_2$ ) to ankerite ( $(\text{CaMg}_{0.75}\text{Fe}_{0.25})(\text{CO}_3)_2$ ). Ankerite often forms as a **\*cement** by diagenetic reaction (see **DIAGENESIS**) from ferroan **\*calcite** at burial depths of about 2.5 km. It is named after the Austrian mineralogist M. J. Anker. *See also* **CARBONATITE**.

**Ankylosaurus (ankylosaurs)** Genus of **\*Cretaceous dinosaurs**, heavily armoured with bony plates, rather in the manner of armadillos. They had small heads with insignificant teeth, and some forms were toothless.

**ankylosaurs** *See* **ANKYLOSAURUS**.

**Anna University Microsatellite (ANUSat)** A cooperative **\*microsatellite** mission by Anna University, Chennai, India, and the Indian Space Research Organization (ISRO) to foster interest in Earth observation and space technology among students. The satellite was launched on 20 April 2009 and ended in April 2012.

**annealing** *See* **HOT WORKING**.

**Annelida** Phylum of *\*coelomate* worms which possess a definite head and good *\*metameric* segmentation. The body is elongate, and each segment has bristles. Their fossils are found in rocks dating from the *\*Cambrian*, and possible fossil annelid worms are known from *\*Precambrian* sediments in S. Australia. They are represented today by the earthworms, sandworms, and leeches.

**Anning, Mary** (1799–1847) The daughter of a cabinet maker who sold ‘natural curiosities’ in Lyme Regis, on the Dorset coast in the south of England, Mary Anning learned from her father how to collect *\*fossils* from the *\*Jurassic* rocks exposed in the cliffs along the shore. After her father died in 1810, Mary began collecting fossils on her own behalf. In 1811 she saw bones protruding from the cliff and hired men to dig out the rock in which they were embedded. When the material was finally exposed it proved to be the skeleton of an animal 5 m long, and eventually identified as *Ichthyosaurus platyodon* (see *ICHTHYOSAURIA*). Mary Anning later found fossils of cephalopods (see *CEPHALOPODA*), Plesiosaurus (see *PLESIOSAUROIDEA*), *Dimorphodon* (*\*Pterosauria*), and fish. Her discoveries attracted many distinguished palaeontologists to Lyme Regis and made her famous. She is commemorated by a stained glass window in the parish church and in the tongue-twister ‘She sells sea shells on the sea shore’.

**annual snow-line** See *FIRN LINE*.

**annulus (planetary)** A ring-like structure used, for example, to describe a discrete ring of ejecta around a martian *\*rampart crater*.

**anode** A positive electrode. See *ANION*.

**anomalous lead** *\*Lead–lead dating* is based on the assumption that the decay of uranium and thorium add the *\*isotopes*  $^{206}\text{Pb}$ ,  $^{207}\text{Pb}$ , and  $^{208}\text{Pb}$  at a constant rate to the lead being sampled. Certain processes (e.g. *\*pneumatolysis*) can redistribute lead isotopes, however, altering the proportions in lead that is then isotopically anomalous. Anomalous lead can give dates that are older than the rock in which the lead is embedded, so it appears that nearby *\*common lead* will not form until some time in the distant future.

**anomaly** See GEOBOTANICAL ANOMALY; GEOCHEMICAL ANOMALY; GRAVITY ANOMALY; HEAT-FLOW ANOMALY; MAGNETIC ANOMALY.

**anomphalous** See IMPERFORATE.

**anorogenic** Applied to 1. a feature that was not formed as a result of tectonic disturbance, i.e. in an interval between orogenic events, and 2. such an interval.

**anorthite** See ALKALI FELDSPAR.

**anorthoclase** See ALKALI FELDSPAR.

**anorthosite** \*Plutonic rock composed almost entirely of \*plagioclase feldspar (90% or more) which may be oligoclase, andesine, labradorite, or bytownite. These rocks occur most frequently in \*Precambrian \*shield areas as non-stratiform intrusions or in layered complexes. Anorthosite also constitutes the \*Lunar Highlands where it is largely composed of the calcium \*end-member anorthite.

**anoxic** The condition of oxygen deficiency or absence of oxygen. Anoxic \*sediments and anoxic bottom waters are commonly produced where there is a deficiency of oxygen due to very high organic productivity and a lack of oxygen replenishment to the water or sediment, as in the case of stagnation or stratification of the water body.

**Antarctic air** Very cold, continental polar \*air mass, which originates over the frozen surface of Antarctica and the surrounding pack-ice, and over the coldest waters of the \*Antarctic Ocean. As this air moves northward over warmer waters it becomes convectively unstable.

**Antarctic bottom water (AABW)** A dense bottom-water mass, formed in the Weddell and Ross Seas, which moves in an easterly direction around Antarctica under the influence of the deep-reaching, wind-driven surface \*Antarctic Circumpolar Current (West Wind Drift). It is typified by a \*salinity of 34.66 parts per thousand and a low temperature ( $-2^{\circ}\text{C}$  to  $-0.4^{\circ}\text{C}$ ). The high salinity and dense nature is caused by the removal of pure water as sea ice.

**Antarctic Circumpolar Current (West Wind Drift)** The largest and most important ocean current in the southern hemisphere. It flows in an eastward

direction around Antarctica, and occupies a wide tract of water in the S. \*Pacific Oceans, S. \*Atlantic Oceans, and \*Indian Oceans. There is very little separation between surface- and \*bottom-water circulation within this area. The current is remarkably constant and is characterized by low \*salinity (less than 34.7 parts per thousand), and cold waters (−1 to 5 °C). It is the only current which flows right around the world.

**Antarctic Coastal Current (East Wind Drift)** A narrow ocean current that flows in a westerly direction parallel to the coast of Antarctica at about 65° S. Its average speed is 0.1 m/s. The current is not continuous, being absorbed into the \*gyres circulating in the Weddell and Ross Seas.

**Antarctic convergence (AAC, Antarctic Polar Front)** A convergence line in the seas that circle Antarctica between latitudes 50° S and 60° S. It is where the cold waters from the Antarctic region meet and sink beneath the warm waters from the middle latitudes, so forming the \*Antarctic intermediate water.

**Antarctic front** Frontal boundary between cold Antarctic air and warmer air, usually lying to the north of it.

**Antarctic intermediate water (AIW)** A water mass formed at the surface near to the \*Antarctic convergence, at about 50° S. It is typified by a low \*salinity (33.8 parts per thousand) and low temperature (2.2 °C). As it spreads northwards it sinks to depths of 900 m and can be traced even in the North Atlantic at 25° N.

**Antarctic meteorites** \*Meteorite finds from \*ablation zones on Antarctic glaciers, which represent about 25% of all the meteorites so far discovered. They vary in composition but are usually quite fresh and may have fallen from one million years ago to the present day. They are therefore of great importance in the study of meteorites. Antarctic meteorites include the first meteorites known to have been derived from the \*Moon and \*Mars.

**Antarctic Ocean (Southern Ocean)** The ocean surrounding Antarctica. It extends northwards to about 40° S latitude, the limit of the northward drift of ice from the Antarctic region, where there is a marked change in water temperature and \*salinity. The Antarctic Ocean is typified by low water temperatures (−1.8 °C to 10 °C).

**Antarctic Plate** A present-day major lithospheric plate that extends beyond the Antarctic continent to the surrounding **\*constructive** plate margins.

**Antarctic polar current** A surface current which flows in a westward direction around Antarctica under the influence of easterly winds blowing off the ice cap.

**Antarctic polar front** See ANTARCTIC CONVERGENCE.

**ante-** From the Latin *ante*, meaning 'before', a prefix meaning 'preceding' or 'previous'.

**antecedent drainage** The hypothesis that a stream crossing a geologic structure may be older than that structure. Such a stream is believed to maintain its course across a developing **\*fold** or **\*fault**. This is a favoured explanation for streams crossing structures in geologically active areas, as do the Arun in the Himalayas, and the Grand Canyon section of the Colorado in the Rockies.

**anterior (biological)** In animals, applied to the front part of the body when the animal is moving forward. In plants, that part of the inflorescence that is farthest from the main stem.

**Anthocyathea** See IRREGULARES.

**anthophyllite** **\*End-member**  $(\text{Mg,Fe})_2(\text{Mg,Fe})_5[\text{Si}_4\text{O}_{11}]_2(\text{OH,F})_2$ , along with gedrite, of the calcium-poor, orthorhombic **\*amphiboles**. In anthophyllite  $\text{Mg} > \text{Fe}$ , whereas in gedrite  $\text{Mg} < \text{Fe}$ . It is found exclusively in **\*metamorphic rocks**, where it may occur with **\*cordierite**. Amosite is the asbestiform variety of anthophyllite.

**Anthophyta** In earlier classifications, a phylum or division that contained all plants bearing flower-like structures, including the **\*angiosperms**, **\*Bennettitales**, and **\*Gnetales**. The group is now known to be **\*paraphyletic**, and the term is no longer used formally.

**Anthozoa (sea-anemones, corals, sea pens)** (phylum **\*Cnidaria**) A class of exclusively polyploid, marine cnidarians. They probably first appeared in the **\*Ordovician** although there are possible records for some groups in the **\*Cambrian**. They are solitary or colonial and usually sedentary. The oral end is expanded as an oral disc with a central mouth that has one or more rings of hollow tentacles. Anthozoans have a well-developed stomodeum

(gullet) leading from the mouth to the enteron (gastrovascular cavity). The interior of the enteron is divided by **\*mesenteries** (infoldings of the gut wall). Those members of the class that secrete hard, calcareous skeletons are important in the geologic record from the **\*Palaeozoic** onwards, and at some levels form true coral **\*reefs**. In the Palaeozoic they are often associated with other organisms, e.g. stromatoporoids (**\*Stromatoporoidea**), to produce organic build-ups or reef mounds. See **HETEROCORALLIA**; **OCTOCORALLIA**; **RUGOSA**; **SCLERACTINIA**; **TABULATA**; **ZOANTHARIA**.

**anthracite** A type of **\*coal**, relatively hard, jet black, with a metallic **\*lustre**, sub-**\*conchoidal** fracture, unbanded, with less than 10% **\*volatiles** and more than 90% carbon. It burns with intense heat and a non-luminous flame.

**Anthracosauria (Embolomeri)** An order of **\*Carboniferous** and early **\*Permian** labyrinthodont amphibians (**\*Labyrinthodontia**) that includes the ancestors of reptiles. Many of the fossils were found in coal measures and the name means 'coal lizards'. The anthracosaurs were first defined as large, aquatic amphibians, and later as reptile-like amphibians. They are now defined as reptile-like tetrapods (**\*Tetrapoda**), and probably not true amphibians, comprising a **\*paraphyletic** order within the superorder **\*Reptilomorpha**.

**anthropogenic** Applied to substances, processes, etc. of human origin, or that result from human activity.

**anthropogeomorphology** The study of those land-forms and processes that are a direct result of human activity, including accelerated **\*erosion**, channelized river channels (i.e. rivers made to flow along fixed, sometimes concrete-lined, channels), the melting of **\*permafrost**, and ground subsidence due to the extraction of water or minerals. Particular examples include the Norfolk Broads, England, which are essentially flooded peat quarries, and the Zuider Zee, whose damming has had a major impact on the coastal morphology of the Netherlands.

**Anthropoidea** See **SIMIIFORMES**.

**anthrosols** A reference soil group in the **\*World Reference Base for Soil Resources**. Anthrosols have **\*soil horizons** that have been strongly affected

by human activity (see [HORTIC HORIZON](#); [IRRAGIC HORIZON](#); [TERRIC HORIZON](#)).

**anti-** From the Greek *anti*, meaning ‘against’, a prefix meaning ‘against’ (in the sense of opposed to), ‘opposite’, or ‘preventing’.

**Antian** The Lower [\\*Pleistocene](#), temperate, marine deposits that form the upper section of a tripartite division of deposits revealed in a [\\*borehole](#) at Ludham, in eastern England. See also [BAVENTIAN](#); [LUDHAMIAN](#); [PASTONIAN](#); [THURNIAN](#).

**anticlinal trap** A [\\*fold](#) structure with an arch of non-porous rock overlying porous strata ([\\*reservoir rock](#)), providing a trap in which oil, gas, or water may accumulate. In Middle East oilfields, large, upright folds occur in thick, competent [\\*limestones](#) which extend for many kilometres; fracturing along the crests of the [\\*anticlines](#) increases [\\*permeability](#) in the reservoir rocks. See [PETROLEUM](#); [NATURAL GAS](#). Compare [FAULT TRAP](#); [REEF TRAP](#); [STRATIGRAPHIC TRAP](#); [STRUCTURAL TRAP](#); [UNCONFORMITY TRAP](#).

**anticlinal valley** Valley developed along the axis of an [\\*anticline](#). Inward-facing [\\*escarpments](#) are developed where the upper beds of the anticline are relatively resistant, while softer lower materials occupy the valley floor. It is a common land-form in gently folded strata of varying resistance, such as occur in southern Britain.

**anticline** Arch-shaped fold in rocks, closing upwards, with the oldest rocks in the core.

**anticlinorium** A regional antiformal structure (see [ANTIFORM](#)) composed of a series of smaller, higher-order [\\*anticlines](#) and [\\*synclines](#), some of which may be small enough to be viewed in [\\*outcrop](#).

**anticoincidence circuit** A device to minimize errors that may occur when measurements are made to date radiocarbon samples. These measurements must be extremely accurate due to the very low level of activity (see [RADIOCARBON DATING](#)). The error quoted on a radiocarbon age determination is solely an error in counting statistics. Such errors may rise from spurious counts generated by contamination of the sample, cosmic activity detected by the counter, and radioactive contaminants in the equipment being used. Initially the counter was shielded by surrounding it

with large amounts of iron, lead, distilled mercury, or paraffin wax mixed with boric acid. An anticoincidence circuit is an alternative to material absorbers, and consists of a series of tangentially placed **\*Geiger** tubes operated in anticoincidence (i.e. they do not require input signals to arrive within specified intervals in order to be activated). These are positioned within an iron shield, and around the central counting chamber. Radiation from outside, or from within the shield, is detected by this ring of Geiger tubes and can be discounted. Special counters have now been developed in which the anticoincidence counters are built into the same tube as the main counter, so that the same gas is used in the whole system. The wall of such a counter usually consists of a polystyrene foil covered on both sides with aluminium. This is then surrounded by a ring of wires forming the **\*anode** for the anticoincidence circuit.

**anticyclogenesis** Process whereby an **\*anticyclone** or a ridge of high pressure is formed and developed.


**anticyclolysis** Process whereby an **\*anticyclone** or ridge of high pressure is dissipated or weakened.

**anticyclone** Area or system of high atmospheric pressure that has a characteristic pattern of air circulation, with subsiding air and horizontal divergence of the air near the surface in its central region. Winds are generally light because of small pressure gradients; they flow clockwise in the northern hemisphere and anticlockwise in the southern hemisphere. A **\*temperature inversion** is common at the base of the air subsidence, and this restricts the vertical development of cloud. Weather conditions are generally settled. Cold anticyclones form over continents in winter and over polar areas at any time, accompanied by strong inversions: in the clear air, pronounced frosts and very cold surface conditions result. Warm anticyclones (so called because of the warm, subsided air aloft) over land areas typically bring spells of settled and often warm weather. *See also* ANTICYCLONIC GLOOM.

**anticyclonic gloom** Condition of low visibility associated with **\*anticyclones** accompanied, in the colder months of the year, by well-developed **\*temperature inversions** that can trap dust and other pollutants, and often have **\*radiation fog** in the lower layers. The stability of the high-

pressure system can make the resultant reduced visibility very persistent and may establish *\*smog* conditions.

**antidune** The *\*sediment \*bedform* generated by fast, shallow *\*flows* of water with a *\*Froude* number greater than 0.8. Antidunes form beneath *\*standing* waves of water that periodically steepen, migrate, and then break upstream. The antidune bedform is characterized by shallow *\*foresets* (see *CROSS-STRATIFICATION*) which *\*dip* upstream at an angle of about 10°. Their preservation potential is low, but they can be identified by low-angle (less than 10°) foresets, dipping up-current. Normally antidunes show a close association with upper-flow-regime *\*plane beds*.



**Antidune**

**antiferromagnetic** Applied to a *\*ferromagnetic* (in the wide sense) substance in which the magnetic lattices are magnetized in exactly equal and opposite directions. Such a substance does not have an external magnetic field in its pure form, but a distorted lattice may result in a *\*parasitic magnetization*.

**antiform** Arch-shaped rock structure which, by definition, closes (i.e. arches) upward, but in which it may not be possible to determine the oldest rocks. It is frequently observed in complex orogenic regions.

**antigorite** See SERPENTINE.

**antimonite** See STIBNITE.

**antimony, native** Metallic element, Sb; soft, whitish; *\*crystals \*tabular \*hexagonal*; *\*cleavage* basal; occurs native in association with *\*stibnite* (antimonite).

**antimony glance** See STIBNITE.

**antiperthite** See HYPERSOLVUS GRANITE; PLAGIOCLASE FELDSPAR.

**antipode** The diametrically opposite point on the surface of a sphere.

**antithetic fault** A *\*fault*, usually one of a set, which in vertical section shows a sense of *\*slip* opposite to that of the major fault from which it originates. The term derives from the Greek word *antithethemi*, meaning ‘set against’.

**antitrade** Upper wind in low latitudes that flows counter (i.e. west-to-east) to the *\*trade* wind below.

**ANUSat** See ANNA UNIVERSITY MICROSATELLITE.

**anvil** The common name for the supplementary cloud feature known as an *\*incus*.


**apatite** A widely distributed *\*phosphate \*mineral*, with the formula  $\text{Ca}_5(\text{PO}_4)_3(\text{F},\text{Cl},\text{OH})$ ; sp. gr. 3.1–3.3; *\*hardness* 5; *\*hexagonal*; usually a shade of green or grey-green, but may also be white, brown, yellow, bluish, or red; white *\*streak*; *\*vitreous \*lustre*; *\*crystals* commonly hexagonal *\*prisms*, and often *\*tabular*, also occurs *\*massive*, and granular; *\*cleavage*, basal {0001}, imperfect prismatic {1010}; found as an *\*accessory mineral* in *\*igneous* rock, in *\*pegmatites* and high-temperature *\*hydrothermal* veins, and in *\*metamorphic rocks*. It is the principal constituent of fossil bones (see also COLLOPHANE). Apatite is widely used as a phosphate fertilizer, and for the production of phosphoric acid and various other chemicals.

**Apatosaurus** One of several gigantic *\*saurischian \*dinosaurs* recorded from the Upper *\*Jurassic*. *Apatosaurus* was a sauropod, a quadruped with a long neck, the total weight of which reached 30 tonnes. The name *Apatosaurus* is a senior synonym of *Brontosaurus* (i.e. it was the earlier of the two names). Animals of 22 m in length have been recorded from the Morrison Formation of N. America.

**ape** A name originally (in medieval times) applied to the Barbary macaque (*Macaca sylvanus*) of North Africa (as were the Latin word *simia* and Greek *pithecus*) and, by extension, applied to other primates as these were made known in Europe. As long-tailed monkeys (‘tailed apes’, or cercopithecids) became better known, ‘ape’ came to mean primarily ‘tailless ape’, and today commonly denotes a member of the *\*Hominoidea*,


comprising lesser apes (gibbons) and great apes (orang-utan, gorilla, chimpanzee, and, in some usages, human).

**aperture 1. (window)** The portion of a data record which is selected for specifying operators for use on the data set, e.g. operators such as **\*autocorrelation** functions and **\*filters** are applied by apertures on **\*seismic records**. **2.** In a mollusc (**\*Mollusca**) shell, the opening through which the soft parts of the animal emerge. It is often a simple, circular opening, but is modified in some genera. In gastropods (**\*Gastropoda**), where the aperture is circular or elliptical it is said to be 'holostomatous' or 'entire'; where it is notched (to accommodate a **\*siphon**) it is said to be 'siphonostomatous'. In cephalopods (**\*Cephalopoda**), the aperture may be indented or notched at the ventral margin by a hyponomic sinus, which houses the **\*hyponome**. Some compressed ammonites (**\*Ammonoidea**) possess a pair of lateral shell extensions (lappets) on either side of the aperture.



### Aperture

**apex** The first-formed end of a shell, which is usually pointed. The term is most commonly applied to gastropod (**\*Gastropoda**) shells.



**Apex**

**APF** See ABSOLUTE POLLEN FREQUENCY.

**aphanitic** An *igneous* rock *texture* characterized by mineral grains which are too small to be identified without a petrological microscope. These extremely fine-grained, crystalline *fabrics* are formed when a *magma* solidifies in response to a very rapid loss of heat and dissolved gases. Emplacement in high-level *dykes* or eruption on to the surface can result in the development of aphanitic fabrics.

**Aphebian** A *stage* (2500–1600 Ma ago) of the lowermost *Proterozoic* of Canada, preceded by the Neoproterozoic era and overlain by the *Helikian* stage.

**aphelion** Point in the Earth's elliptical orbit at which the planet is farthest from the Sun. See also PERIHELION.

**aphyric** An *igneous* rock *texture* characterized by a fine-grained *aphanitic* *groundmass* and by an absence of any *phenocrysts*. Aphyric texture forms by the rapid crystallization of *melts* lacking large suspended

**\*crystals**, and thus these melts must have been very close to their **\*liquidus** temperature (the temperature at which initial crystallization takes place in a cooling melt).

**API gravity** A measure of the density of **\*petroleum** devised by the American Petroleum Institute (API). If the API gravity is less than 10 the petroleum is denser than water and will sink; if it is greater than 10 the petroleum is less dense than water and will float. Although the value is dimensionless, it is expressed in degrees (° API).

**aplite** A light-coloured, fine-grained, equigranular **\*igneous** rock composed of **\*subhedral** to **\*anhedral** grains of **\*quartz** and **\*alkali** feldspar, and found as late-stage veins in **\*granite** bodies. The quartz–alkali feldspar composition corresponds to the lowest temperature **\*melts** in granite **\*magma** systems, suggesting that they are residual melts formed by the differentiation of granite **\*magma** (see **MAGMATIC DIFFERENTIATION**). The lack of any hydrous minerals and the fine **\*grain** size points to the aplites crystallizing from dry residual melts.

**apodeme** See **EXOSKELETON**.

**apogee** The point in the orbit of the Moon (or any artificial satellite) that is most distant from the Earth. Compare **PERIGEE**.

**Apollo 1**. A **\*solar** system asteroid (No. 1862), diameter 1.6 km; approximate mass  $2 \times 10^{12}$  kg; rotational period 3.063 hours, orbital period 1.81 years. Its orbit crosses that of earth. **2**. The name of the **\*NASA** manned lunar programme that ran from 1963 to 1972.



<http://nssdc.gsfc.nasa.gov/planetary/lunar/apollo.html>

- A NASA programme of missions from 1963 to 1972 to land humans on the Moon and return them to Earth.

**apomorphic** Evolutionarily advanced (‘derived’) character state. The long neck of the giraffe is apomorphic; the short neck of its ancestor is **\*plesiomorphic**.

**apomorphic** Applied to features possessed by a group of biological organisms that distinguish those organisms from others descended from the

same ancestor. The term means 'new-featured' and refers to 'derived' characters that have appeared during the course of evolution.

**apophysis** (*pl.* **apophyses**) **1.** An irregular or sheet-like *\*vein* or *\*dyke* which originates from a larger *\*igneous* rock body. **2.** See VERTEBRA.

**Appalachian orogenic belt** A 3200 km long, *\*Palaeozoic*, *\*orogenic* belt extending from Newfoundland to Alabama and interpreted as the consequence of the closure of the *\*Iapetus Ocean*. Extensions of the belt along *\*strike* include the Caledonian orogenic belt (see CALEDONIAN OROGENY) in north-western Europe, now separated by the Atlantic Ocean. The deformation in the Appalachian belt ranges from late *\*Precambrian* to *\*Permian*, divided into four major *\*orogenies* (*\*Avalonian*, *\*Taconic*, *\*Acadian*, and *\*Alleghanian*), with the transport of *\*thrust-\*nappes* predominantly north-westwards. *\*Seismic* reflection *\*profiling* by *\*COCORP* has supported, at least for the southern Appalachians, recent interpretations of a thin-skinned tectonic style in which a few kilometres' thickness of highly deformed material has been thrust westward for at least 200 km.

**apparent age** See ABSOLUTE AGE.

**apparent cohesion** Cohesion of grains due to surface tension in the surrounding pore water.

**apparent conductivity** ( $\sigma_a$ ) The inverse of *\*apparent resistivity*; the units are *\*siemens*/metre.

**apparent dip** The dip of the trace of a plane, which is measured from the horizontal in any section non-perpendicular to the *\*strike* of the plane. Compare TRUE DIP. See also DIP, ANGLE OF.

**apparent polar wander (APW)** The line traced on each of the continents by joining together the positions of the *\*palaeomagnetic* poles at different times. The geometry of the line differs for each continent and appears to suggest a path followed by poles that changed their positions over time. In fact, the changing positions of the poles reflect the latitudinal movements and rotations of lithospheric *\*plates*, rather than movements of the poles.

**apparent resistivity** ( $\rho_a$ ) A measurement of resistivity which is calculated as the product of the measured resistance ( $R$ ) and a **\*geometric factor** ( $K_g$ ) such that  $\rho_a = K_g R$ , in units of ohms/m. It is important to note that the apparent resistivity is not an explicit measurement of the electrical resistivity of a material and should not be interpreted as being diagnostic of a given material.

**apparent velocity** ( $v_a$ ) The velocity which a wave-front exhibits along a line of **\*geophones**. If the wavefront approaches the geophone **\*array** at an angle  $\theta$ , then the true velocity of the wavefront,  $v$ , is given by  $v = v_a \cos\theta$ . In **\*refraction** seismology,  $v_a$  is the reciprocal of the gradient of a straight-line segment of a **\*time–distance** graph.

**apparent wavelength** ( $\lambda_a$ ) If a wave train approaches a line of **\*geophones** at an angle  $\theta$ , then the distance between corresponding points on successive cycles of the wave form as detected by those geophones is an apparent and not a true wavelength. True wavelength  $\lambda = \lambda_a \sin\theta$ .

**appinites** Medium- to coarse-grained **\*plutonic** rocks that occur as **\*intrusions** of **\*hornblende** in a **\*matrix** of **\*intermediate rock** and/or **\*alkali \*plagioclase feldspar**. They are named for the region around Appin in western Scotland.

**AprizeSat-3 and -4** Two **\*minisatellites**, each a cube of 25 cm sides weighing 13 kg, belonging to Aprize Satellite, Inc., of Fairfax, Virginia, that aim to demonstrate an automatic identification system for detecting ships on the ocean. The minisatellites were launched on 29 July 2009, from Baikonur Cosmodrome, Kazakhstan, into a Sun-synchronous near-circular orbit.

**apsacline** A condition of the **\*interarea** of a strophic brachiopod (**\*Brachiopoda**) shell where the inclination of the **\*cardinal** area relative to the plane of **\*commissure** is 90–180°. It is one of the most commonly occurring conditions. *Compare* ANACLINE.

**aptation** A character that suits its possessor to its **\*environment**; it may be an **\*abaptation**, **\*adaptation**, or **\*exaptation**.

**Aptian** 1. An Early *\*Cretaceous* *\*age*, preceded by the *\*Barremian*, followed by the *\*Albian*, and dated at 125–112 Ma ago (Int. Commission on Stratigraphy, 2004). 2. The name of the corresponding European *\*stage*, which is named after Apt in France. *See also* **NEOCOMIAN**.

**aptychus** A calcitic plate associated with *\*Mesozoic* *\*ammonites*. Normally these plates occur in pairs. Aptychi are shaped like bivalves (*\*Bivalvia*) and have an ornamented outer surface. Since they have been found inside the aperture of ammonoid shells they were originally interpreted as *\*opercula*, protecting the ammonoid body when it was withdrawn into the shell. It has now been shown that aptychi are probably the lower jaws of ammonoids and that the horny upper jaw is rarely preserved.

**APW** *See* **APPARENT POLAR WANDER**.

**APXS** *See* **ALPHA–PROTON–X-RAY SPECTROMETER**.

**Aqua** A *\*NASA* satellite, launched on 4 May 2002, that carries six Earth-observing instruments to measure almost all aspects of the *\*hydrologic cycle*, as well as radiative energy fluxes, *\*aerosols*, terrestrial vegetation cover, marine phytoplankton and dissolved organic matter, and air, land, and water temperatures. It is one of the four satellites that comprise the *\*A-Train* and was still functioning on 31 July 2018.



[http://www.nasa.gov/mission\\_pages/aqua/](http://www.nasa.gov/mission_pages/aqua/)

- A NASA mission launched in 2002 to study the Earth's water.

**aquamarine** *See* **BERYL**.

**Aquarius** An instrument built and owned by NASA that was launched on 10 June 2011 on a Delta II vehicle, carried on the Argentine Satellite de Aplicaciones Cientificas-D (SAC-D) of the Comisión Nacional de Actividades Espaciales (CONAE), the Argentine space agency. Aquarius is in a *\*Sun-synchronous orbit* at a height of 657 km, and it measures sea-surface salinity.



<http://aquarius.nasa.gov/>

- A joint CONAE and NASA mission to measure sea-surface salinity.

**aquiclude (aquifuge)** A rock with very low values of hydraulic conductivity which, although it may be saturated with *\*groundwater*, is almost impermeable with respect to groundwater flow. Such rocks will act as boundaries to *\*aquifers* and may form confining strata. See PERMEABILITY.

**aquic moisture regime** The moisture balance of humid climates and soils, where annual precipitation exceeds the combined actual evaporation and transpiration, and where the soil-moisture status is normally above *\*field capacity*.

**aquifer** A body of permeable rock, for example, unconsolidated gravel or sand *\*stratum*, that is capable of storing significant quantities of water, is underlain by impermeable material, and through which *\*groundwater* moves. An unconfined aquifer is one in which the *\*water table* defines the upper water limit. A confined aquifer is sealed above and below by impermeable material. A perched aquifer is an unconfined groundwater body supported by a small impermeable or slowly permeable unit. See PERMEABILITY.

**aquifer test** See PUMPING TEST.

**aquifuge** See AQUICLUDE.

**Aquitanian 1.** The earliest *\*age* (23.03–20.43 Ma ago) in the *\*Miocene* epoch, preceded by the *\*Chattian* (*\*Oligocene*), followed by the *\*Burdigalian* (Int. Commission on Stratigraphy, 2004). **2.** The name of the corresponding European *\*stage*, which is roughly contemporaneous with the upper *\*Zemorrian* and lower *\*Saucesian* (N. America), parts of the *\*Otaian* (New Zealand), and the upper *\*Janjukian* and lower *\*Longfordian* (Australia). The *\*stratotype* is in the Aquitanian Basin, France. The Aquitanian is itself characterized by the appearance of the planktonic *\*foraminiferid* *Globigerinoides primordia*.

**aquitard** A *\*rock* with low values of *\*hydraulic* conductivity, which allows some movement of water through it, but at rates of flow lower than those of

adjacent **\*aquifers**. Compare **AQUICLUDE**.

**Arabian Plate** One of the present-day minor lithospheric **\*plates**, separated from the **\*African Plate** by the spreading **\*Red Sea**, the continuation of the **\*Carlsberg Ridge** into the Gulf of Aden, and the Dead Sea transform system, while the boundary with the **\*Indo-Australian Plate** is the Owen Fracture Zone; the plate is colliding with the Iran Plate.

**arachnid** See **ARACHNIDA**.

**Arachnida (arachnids: mites, scorpions, spiders, etc.)** Class of terrestrial chelicerates (**\*Chelicerata**) which have book lungs or tracheae derived from gills, indicating their aquatic derivation. Of recent terrestrial animals, the arachnids are probably the oldest known class, scorpions having been recorded from the **\*Silurian** period. A Silurian scorpion, *Palaeophonus nuncius*, was perhaps the first terrestrial animal. The first fossil spiders are known from the **\*Devonian**. The class is extremely diverse, but except in the mites the body is in two portions: the prosoma (anterior portion) which bears the four pairs of legs, the eyes, the pedipalps (second pair of appendages), and the chelicerae (first pair of appendages, usually pincer-like); and the opisthosoma (posterior portion) which contains most of the internal organs and glands. The two portions may be broadly jointed, or connected by a stalk or pedicel. The prosoma has a dorsal shield or carapace, and the opisthosoma is segmented in most orders, but not in spiders and mites and only very weakly in harvestmen. The number of eyes varies, and can be as many as twelve in some scorpions.

**arachnid structure (arachnoids)** A strange pattern of ridges, gathered in braids and belts, which merge with radial and concentric ridges to give a 'spiders and cobwebs' appearance on the surface of Venus, particularly in the area between Sedna Planitia and Bell Regio (about 43° N, 19° E). The ridges are large, 100–200 km long and up to 20 km wide.

**arachnoids** See **ARACHNID STRUCTURE**.

**aragonite** **\*Mineral**, CaCO<sub>3</sub>; sp. gr. 2.9; **\*hardness** 3.5–4.0; **\*orthorhombic**; colourless, white, grey, or yellowish; white **\*streak**; **\*vitreous** **\*lustre**; **\*crystals** normally **\*prismatic**, often **\*acicular**, sometimes **\*tabular** and pseudo-hexagonal; also occurs fibrous and stalactitic; **\*cleavage** imperfect {010}; occurs in hot springs and in association with **\*gypsum**, also in veins

and cavities, and in the oxidized zone of ore deposits with other **\*secondary minerals**. Aragonite is a **\*polymorph** of **\*calcite**, from which it is distinguished by its lack of cleavage and its higher specific gravity. Calcite is the more stable form of  $\text{CaCO}_3$ , and many fossil shells that were made originally of aragonite have either converted to calcite or undergone replacement by some other mineral. Present-day mollusc shells are formed of aragonite crystals. The name is derived from the Aragon province of Spain. *See also* **CARBONATES**.

**aragonite compensation depth (ACD)** The depth in the ocean at which the rate of dissolution of **\*aragonite** is equal to the rate at which aragonite accumulates. In sea water, aragonite is less stable than **\*calcite**. Consequently, the ACD is shallower than the **\*carbonate compensation depth (CCD)**. The ACD is typically at 2–3 km.

**aragonite mud** A fine **\*carbonate** mud, with particles less than 4  $\mu\text{m}$ , composed mainly of **\*aragonite** needles. The aragonite is generally believed to have been deposited from the break-up of calcareous green algae (**\*Chlorophyta**).

**Aratauran** A New Zealand **\*stage** (199.6–188 Ma ago) of the Early **\*Jurassic \*epoch**, preceded by the **\*Otapirian** and followed by the **\*Uroan**.

**arborescent** *See* **DENDRITIC**.

**arc** *See* **ISLAND ARC**.

**archae-** (**arche-**) Prefix, from the Greek *arkhaios* ('ancient'), itself derived from *arkhe* ('beginning'). It adds the meaning 'ancient', with the implication 'first', to words to which it is attached.

**Archaea** The **\*domain** comprising what were formerly known as the archaeobacteria. What used to be the kingdom Archaeobacteria has been split into two kingdoms: **\*Crenarchaeota** and **\*Euryarchaeota**. The domain Archaea contains the **\*phenotypes**: **\*methanogens**, **\*sulphate-reducing organisms**, and **\*extremophiles**.

**Archaean (Archaean, Azoic)** One of the three subdivisions of the **\*Precambrian**, lasting from about 4000 to about 2500 Ma ago. Now formally an **\*eon**.

**archaeobacteria** (domain *\*Archaea*) Organisms belonging to the kingdoms *\*Crenarchaeota* and *\*Euryarchaeota*; formerly these were grouped together as the kingdom Archaeobacteria.

***Archaeocalamites radiatus*** First described by A. *\*Brongniart* in 1828, this species is one of the earliest recorded equisetaleans or 'horse-tails', a group of plants noted for the jointed nature of the stem. At each joint there is a ring of short branches. See *SPHENOPSIDA*.

**Archaeoceti (ancient whales)** (cohort Mutica, order Cetacea) An extinct suborder comprising the oldest and most primitive cetaceans, which flourished in the *\*Eocene* and may have originated in Africa. Most were comparable in size with modern porpoises, had an elongated snout, and nostrils on top of the skull. The brain case was long and low. The front teeth were peg-like, the cheek teeth *\*heterodont* and characteristic of primitive carnivores. There were 44 teeth in all. The hind legs in most were reduced to vestiges, but in some early genera (*\*Ambulocetus*, *\*Basilosaurus*) still protruded from the body wall. They were fish-eating carnivores that had adopted an aquatic life to which they were more highly adapted than, for example, modern seals. The term archaeocete really means any primitive cetacean and probably does not designate a natural, *\*monophyletic* group.

**Archaeocyatha** Extinct phylum of reef-forming organisms known only from the *\*Cambrian*. They were cup-like, usually 10–30 mm in diameter and up to 50 mm in height. In some respects they were similar to both *\*sponges* and *\*corals*, and may represent a true advance in evolutionary *\*grade* over the former. It is possible that they lived in a symbiotic relationship with some *\*trilobites*. The cause of their extinction is not known. There were two classes: *\*Regulares* and *\*Irregulares*.

**Archaeogastropoda** An order of *\*Gastropoda* which appears in the Lower *\*Cambrian* and also includes the extant *Patella vulgata* (limpet). Gastropods may be subdivided according to their respiratory structures; archaeogastropods, the most primitive gastropods, possess just two gills. Some forms possess a marginal slit near the *\*aperture* to facilitate the removal of exhaled water and wastes. This may extend posteriorly for some distance but eventually it is filled with shell material as the animal grows and extends its shell. The filled slit (selenizone) is usually completely

plugged. In some species (e.g. *Haliotis*) the selenizone is represented by a linearly arranged series of apertures (tremata, sing. trema).

**archaeological geology** The study of archaeological sites in order to obtain data relevant to geological investigations.

**archaeomagnetism** The study of the magnetic properties of objects and materials from an archaeological context. Such studies include **\*magnetic dating**, reconstruction of objects and structures, sourcing artefacts, past firing temperatures, etc.

**Archaeopteris** Early **\*progymnosperm** found first in the **\*Frasnian** stage of the **\*Devonian** period. It was identified by its fronds, and is the earliest known representative of the Archaeopteridales.

**Archaeopteryx lithographica** Only seven specimens of this species, the first bird, are known. *Archaeopteryx* is recorded solely from the Lithographic Limestone of the Solnhofen region of Bavaria, Germany. It was first described by H. von Meyer in 1861 and is of Middle **\*Kimmeridgian** or Upper **\*Jurassic** age. Recent work on this species by several palaeontologists tends to support the theory that the birds, through *Archaeopteryx*, evolved from **\*coelurosaur** **\*dinosaurs** similar to *Compsognathus*. The species *A. lithographica* possesses several primitive characters such as teeth, as well as specialized features such as feathers and hollow bones. It is a good example of a connecting species which exhibits a mosaic of evolutionary features.

**archaeopyle** See DINOPHYCEAE.

**Archaeosperma arnoldii** The earliest (**\*Devonian**) seed-like structure.

**Archaeosphaeroides** A coccoid cyanophyte (see CYANOBACTERIA) from the Figtree (see SWAZIAN) cherts of S. Africa, which are **\*Archaean** and date from perhaps 3000 Ma ago.

**Archaeozoic** See ARCHAEOAN; CRYPTOZOIC.

**archaic sapiens** Humans (i.e. *Homo sapiens*) that share features with earlier (non-Homo) species, relating mainly to cranial capacity and the robustness of the teeth and skeleton, that are lost in anatomically modern humans. The

archaic species are classified as *Homo sapiens*, but not as belonging to the subspecies of modern humans, *Homo sapiens sapiens*.

**arche-** Alternative spelling for the prefix **\*archae-**.

**archetype** A hypothetical ancestral form in which all the basic characteristics of a taxonomic group occur, although they are not specialized in any one direction. Thus the modern primitive mollusc *Neopilina* is perhaps close to the molluscan archetype.

**Archie's law** An empirical law which relates, for a clay-free sediment, the electrical **\*resistivity**  $\rho$  of a porous rock containing water and cement to the fraction of the pore space that is filled with water.  $\rho = \rho_0^{-m} s^{-n}$ , where  $\rho_0$  is the resistivity of the water and  $s$  is the fraction of pore space filled with water. The exponent  $n$  is usually about 2.0 if one-third of the pore space is filled with water, and exponent  $m$  usually varies between 1.3 for unconsolidated sediments to approaching 2.0 for a well-cemented sediment. In the oil-bearing rocks, the remaining pore space is considered to be filled with either oil or gas and the law is usually expressed as  $S_w = (R_w/R_t)^{0.5} \phi$ , where  $S_w$  is the fraction of pore space filled by water with a resistivity  $R_w$  in a rock with a true resistivity of  $R_t$  and a **\*porosity**  $\phi$ .

**arching 1.** In an underground **\*excavation**, a small inward movement of the sides, roof, and floor of the cavity caused by *in situ* stress around the excavation. This reduces the permeability of the rock in the immediate vicinity of the excavation. **2.** Masonry or steel support in underground workings.

**archipelago** Group of islands; a sea containing many scattered small islands.

**architecture of sandbodies** The large-scale form and arrangement of **\*sandstone** beds. See **SANDBODY**.

**archosaur** See **ARCHOSAURIA**.

**Archosauria (archosaurs)** Infraclass of **\*diapsid** reptiles, including the **\*crocodiles**, **\*dinosaurs**, **\*pterosaurs**, and **\*thecodontians**. Thecodontians were ancestral to the other groups and appeared at the base of the **\*Triassic**. The term 'archosaur' is from the Greek words *arkhi* ('chief' or 'leading') and *saura* ('lizard').

**Arcsecond Space Telescope Enabling Research in Astrophysics (ASTERIA)** A 6-unit **\*CubeSat** space telescope that aims to achieve arcsec-level line of sight pointing error and highly stable focal plane temperature control, to allow precision photometry. The mission is sponsored by the Massachusetts Institute of Technology (MIT) and **\*NASA** and was launched on 14 August 2017, from the Kennedy Space Center, Florida.

**arctic air** Very cold **\*air mass**, generally formed north of the Arctic Circle. As air from this source moves southwards, it cools the regions in which it arrives; but being itself heated in the process it becomes convectively unstable. Polar **\*lows** sometimes form and the accompanying wintry precipitation is often heavy.

**arctic bottom water (ABW)** Cold, dense water that sinks, to a maximum depth of about 6000 m, from the subpolar **\*gyre** in the Greenland and Norwegian Seas, fills the basins of those seas, then spills as an intermittent southward flow through narrow channels in the ridge between Scotland, Iceland, and Greenland.


**arctic front** Frontal boundary between cold, arctic air and warmer **\*air masses**, usually lying to the south of it. Many depressions originate on it. In north-western Canada in winter, for example, the frontal zone incorporates cold, dry, continental polar air and modified maritime arctic air from the Gulf of Alaska to the north of continental tropical air.

**Arctic Ocean** Smallest and shallowest of the major oceanic areas, the shallowness being caused by the surrounding wide continental shelves (up to 1700 km wide). For much of the year the surface is covered by floating pack-ice.

**arctic sea smoke (frost smoke) \*Fog** appearing in very cold air from the arctic-ice or frozen-land regions, when it comes over the warmer water of open parts of the **\*Arctic Ocean**. The rapid heating induces convection currents which rise in the air: these carry moisture upwards from the water surface, and this becomes visible as the moisture quickly condenses again in the very cold surrounding air. Thus a fog of rising columns of condensing water vapour is formed. The fog is usually fairly shallow, wispy, and smoke like. This, and its common occurrence in coastal seas around cold land masses (e.g. Labrador, Greenland, and Norway), gave rise to the name.

Similar steam fogs may be seen in winter over the open water of rivers when the air is 10 °C or more colder than the water.

**arc-trench gap (fore-arc)** The region between an oceanic **\*trench** and the adjacent volcanic **\*island arc**. The arc-trench gap is at least 100 km wide in nearly all cases, and up to 570 km at the eastern end of the Aleutian arc. The width of the arc-trench gap increases with time through the growth of an **\*accretionary wedge** and the development of a **\*fore-arc basin**.



### Arc-trench gap

**arcus 1.** From the Latin *arcus* ('arch'). A cloud feature having a rolled appearance, with fragmented edges on the leading surface of **\*cumulonimbus** and occasionally **\*cumulus**. When well developed, the feature has a prominent arch-like form. *See also* CLOUD CLASSIFICATION. **2.** A curved or arc-shaped feature on the surface of an extraterrestrial body, e.g. **\*Titan**.

***Ardipithecus ramidus*** The earliest known member of the human lineage, discovered in 1993 by Tim White, Gen Suwa, and Berhane Asfaw at Aramis, Ethiopia, and dated to 4.4 million years BP. The canine teeth are somewhat reduced from the primitive ape-like condition, but not so much as in **\*Australopithecus**; the **\*enamel** on the teeth is thin; the deciduous molars are intermediate between those of a human and a chimpanzee. The postcranial skeleton indicates that it was, at least to some degree, bipedal.

**Arduino, Giovanni** (1714–95) Arduino was a Venetian mining engineer, who devised a classification of the rocks of northern Italy, later adopted by T. O. *Bergman*, A. G. *Werner*, etc. He distinguished between the Primary (mountain mica-slates), Secondary (mountain limestones, with marine fossils), and Tertiary rocks (fossiliferous valley sediments).

**areal erosion** Erosion by an *ice* sheet of an area too large to be visible as an erosion feature and identifiable only through mapping at a continental scale.

**arenaceous** Sandy, or sand-like in appearance or texture. The term is applied to *clastic sedimentary rocks* with a grain size of 0.0625–2.00 mm. Three main groups of arenaceous rocks are recognized: quartz sandstones (*quartzites*), which contain 95% *quartz*; *arkoses*, which have greater than 25% *feldspar*; and *greywackes*, which essentially are poorly sorted sediments with rock (lithic) fragments in a mud *matrix*.

**Arenicolites** An *ichnogenus* of *domichnia*.

**Arenig** A *series* (478.6–471.8 Ma ago) of the Lower *Ordovician*, underlain by the *Tremadocian* and overlain by the *Llanvirn*.

**arenite** See DOTT CLASSIFICATION; SANDSTONE.

**arenosols** A reference soil group in the *World Reference Base for Soil Resources*. Arenosols are weakly developed soils with a coarse texture.

**areology** The scientific study of Mars. Derived from the name of the Greek god Ares, known to the Romans as Mars, the god of war, and *logos* (the Greek for word or discourse). It is analogous to geology as the study of the Earth.

**arête** Knife-edged, steep-sided ridge found in upland areas that have been or are being glaciated, and formed by the meeting of adjacent *cirque* headwalls. It may be diversified by ‘gendarmes’ (abrupt rock pinnacles that have resisted frost shattering).

**arfvedsonite** A sodium- and iron-rich *amphibole*.

**argentite (silver glance)** Ag<sub>2</sub>S, *ore mineral* for silver; sp. gr. 7.2–7.4; *hardness* 2.0–2.5 (can be cut with a knife); *cubic*; black and opaque; slimy black *streak*; *metallic lustre*; *crystals* commonly cubes or

octahedra, but can be *\*massive*; *\*cleavage* poor, cubic; occurs in *\*hydrothermal* veins in association with native silver and as a weathering product of primary silver sulphides; unstable below 179 °C and replaced by the *\*monoclinic form*, acanthite.

**argic horizon** In the *\*World Reference Base for Soil Resources*, a *\*soil horizon* showing signs of clay enrichment.

**argillaceous** Applied to rocks which are *\*silt-* to *\*clay-sized* *\*sediments* (grain size less than 0.0625 mm in diameter). They account for more than 50% of *\*sedimentary rocks* and most have a very high *\*clay mineral* content. Many contain a high percentage of organic material and can be regarded as potential *\*source rocks* for *\*hydrocarbons*.

**argillaceous limestone** See MARLSTONE.

**argillans** See CUTAN.

**argillic horizon** Subsurface B *\*soil horizon* that is identified by the illuvial (see ILLUVIATION) accumulation of silicate *\*clays*. The amount of clay necessary is defined in comparison with the quantity in the overlying eluvial (see ELUVIATION) horizon, but is at least 20% more. *\*Cutans* or clay skins may be used to help identify an argillic horizon. It is at least one-tenth as thick as the overlying horizons.

**argillite (lutite)** A well-compacted, non-fissile rock, containing *\*clay-* and *\*silt-sized particles*; more indurated than a *\*mudstone*.

**Argo** An international programme that began in 2000 to monitor temperature, salinity, currents, and biological properties of the world's oceans through more than 3900 floats which drift with the currents and are now distributed throughout all oceans. They are parked 1 km below the surface. Every 10 days, by altering their buoyancy, they sink to 2 km then rise to the surface to transmit the data their instruments have acquired.



[www.argo.net](http://www.argo.net)

- Argo Programme home page.

**argon–argon dating** A method for **\*radiometric dating** in which a sample is irradiated to convert  $^{39}\text{K}$  to  $^{39}\text{Ar}$ . The sample is then heated in stages, at each stage yielding argon with a particular  $^{40}\text{Ar} : ^{39}\text{Ar}$  ratio. The age of the sample is calculated by comparing that ratio with the ratio in a sample of known age. The method provides more detail than **\*potassium–argon dating** and it can be used with very small samples.



<http://www.diggles.com/bturrin/40Ar.html>

- A tutorial on the  $^{40}\text{Ar}/^{39}\text{Ar}$  step-heating dating technique.

**argon-40** See **POTASSIUM–ARGON METHOD**.

**Argos DCS (Data Collecting System)** A spaceborne system that collects data for studying and protecting the environment. It comprises polar-orbiting weather satellites linked to ground stations equipped with sensors and transmitters. The moving satellites allow for the use of **\*Doppler** shift calculations. The programme began in 1974 and since 1978 has been flown on all TIROS-N (**\*Television and Infrared Observation Satellite**) satellites.

**aridic moisture regime** The moisture balance of arid climates and soils, where the annual precipitation is less than the potential evaporation and transpiration, and where soil moisture status is normally less than **\*field capacity**.

**aridisols** In the US Soil Taxonomy, an order of soils found in arid environments. These soils have very little organic matter in their surface **\*horizons**, but may contain calcium carbonate or **\*gypsum**, and/or soluble-salt accumulations.

**aridity index** Indication of moisture deficit. All climatic classifications include arid categories, defined either by quantitative or, more usually, by mainly subjective criteria. C. W. Thornthwaite first used the term ‘aridity index’ and calculated it as 100 times the water deficit, divided by the potential evaporation. See also **RADIATION INDEX OF DRYNESS**; **THORNTHWAITE CLIMATE CLASSIFICATION**.

**Ariel (Uranus I)** One of the major satellites of **\*Uranus**, with a radius of  $581.1 \times 577.9 \times 577.7$  km; mass  $13.53 \times 10^{20}$  kg; mean density 1670

kg/m<sup>3</sup>; albedo 0.34. Surface features are visible, including fissures, craters, and flows.

**aristogenesis** Theory, like those of entelechy, homogenesis, and orthogenesis, that claims that evolution proceeds along a determined path. Today, however, most people accept that natural selection does not direct evolution towards any particular kind of organism or physiological attribute.

**Aristotle's lantern** The jaw apparatus present in **\*regular echinoids**. Five strong jaws, each with a single **\*calcite** tooth, come together into a lantern-shaped structure which is suspended within the mouth of the echinoid. The teeth are used in scraping algae and other food from the sea floor.

**arkose** **\*Arenaceous** rock containing **\*quartz** and 25% or more of **\*feldspar**. The feldspar is easily destroyed during chemical change or transportation, implying that arkoses were deposited rapidly under fairly arid environmental conditions. Most were deposited near to land, probably in close proximity to a granitic area.

**arkosic arenite** A **\*sandstone** comprising more than 25% **\*feldspar** and with more feldspar than rock fragments, with less than 15% mud **\*matrix**. The matrix is defined as material less than 30 µm in grain size. See **DOTT CLASSIFICATION**.

**arkosic wacke (feldspathic wacke, feldspathic greywacke)** A **\*sandstone** comprising more than 5% **\*feldspar**, and more feldspar than rock fragments, with more than 15% **\*matrix**. The matrix is defined as material less than 30 µm in grain size. See **DOTT CLASSIFICATION**.

**arls** A collective name for **\*marl**, **\*sarl**, and **\*smarl**.

**Arnsbergian** A **\*stage** (326–325 Ma ago) of the **\*Serpukhovian** epoch, underlain by the **\*Pendleian** and overlain by the **\*Chokierian**.

**Arowhanan** A New Zealand **\*stage** (95–93 Ma ago) of the Late **\*Cretaceous** **\*epoch**, preceded by the **\*Ngaterian** and followed by the **\*Mangaotanean**.

**array** A geometrical distribution or pattern. A line of **\*geophones** or **\*shot** points constitutes an array, as does a line of electrodes for a **\*resistivity**

survey. *See also* ELECTRODE CONFIGURATION.

**Array of Low-Energy X-Ray Imaging Sensors (ALEXIS)** A US demonstration mission of \*microsatellite technology by Los Alamos National Laboratory that was launched in April 1993 and was decommissioned in April 2005.

**arrow worms** *See* CHAETOGNATHA.

**arroyo** Gully found along valley floors in an arid or semi-arid region and possessing steep or vertical walls cut in fine-grained cohesive sediments. The floor is flat and usually sandy. Found especially in the south-western United States, parts of India, S. Africa, and around the Mediterranean.

**arsenopyrite** (known historically as **mispickel**) \*Mineral, FeAsS; sp. gr. 5.9–6.2; \*hardness 5.5–6.0; \*monoclinic; silver-grey to white, often with a tarnish; dark greyish or black \*streak; \*metallic \*lustre; \*crystals \*prismatic, often striated, can also be \*massive and granular; \*cleavage distinct {101}; occurs in high- to moderate-temperature mineral veins in association with \*gold, ores of tin, tungsten, \*galena, and \*quartz, and also disseminated in \*limestones, \*dolomites, \*gneisses, and \*pegmatites. It is the principal source of arsenic compounds used for pest control, for the manufacture of dyes and chemicals, and in leather treatment.

**artesian water** \*Groundwater that is confined in an \*aquifer, but which may overflow on to the land surface via artificial \*boreholes or, sometimes, natural \*springs, because of the high \*hydraulic head that may be developed in a confined aquifer. Artesian conditions are common when the aquifer has a \*syncline form. The London Basin, England, provided artesian water during the 19th century from a chalk aquifer sealed by clays. The term is derived from the Artois region of north-western France.

**artesian well (overflowing well)** A well that flows at the surface without pumping, because it is sunk into a confined aquifer whose \*hydraulic head (sometimes called the potentiometric or piezometric head) lies above ground level. *See* AQUIFER; ARTESIAN WATER.

**Arthrodira (arthrodiriformes)** (class \*Placodermi) Group or order of fossil (\*Devonian) fish with a body covered with bony plates, including a heavily armoured \*head shield, the gills opening between head and body armour.

**Arthropoda** A highly diverse phylum of jointed-limbed animals, which includes the crustaceans, arachnids, and insects as the major components, as well as the classes Symphyla, Paupoda, Chilopoda (centipedes), Diplopoda (millipedes), and the extinct *\*Trilobita* and eurypterids (see *CHELICERATA*; *MEROSTOMATA*). Arthropods first appeared in the *\*Cambrian*, already well diversified with such forms as the trilobites, trilobitoids, *\*Ostracoda*, and crabs present, implying an earlier, hidden history, reaching back into the Precambrian. Embryological evidence shows that they are derived either from primitive *\*Polychaeta*, *\*Annelida*, or from ancestors common to both. Arthropods share with annelid worms a metamerically segmented body (see *METAMERIC SEGMENTATION*), at least in the embryo, a dorsal heart, a dorsal anterior brain, and a ventral nerve cord that has segmental, ganglionic swellings. The limbs of all arthropods are paired, jointed, and segmental, and the body has a chitinous *\*exoskeleton*. Primitively, the limbs and cuticular plates correspond to the metamerically segmentation of the body, but in many groups there is considerable loss and/or fusion of segments.

**arthropods** See *ARTHROPODA*.

**Articulata** 1. (phylum *\*Brachiopoda*) A class of brachiopods, existing from the Lower *\*Cambrian* to the present day, in which the shell is calcareous and comprises valves hinged by teeth in one valve and sockets in the other. The *\*pedicle* is of a horny material. Their main radiation was in the early *\*Ordovician*; of the seven orders present in the *\*Palaeozoic*, three are still extant. 2. (*sea lilies*) (subphylum Crinozoa, class *\*Crinoidea*) Subclass of sea lilies (crinoids) in which the basal plates are small or strongly reduced. Articulation between the radial and brachial plates, and in the majority of brachials, is muscular, with a well-developed fulcral ridge. The arms are always *\*uniseriate*. All post-*\*Palaeozoic* crinoids belong to the Articulata.

**artificial freezing** Method of controlling *\*groundwater* and improving strength of ground by pumping a refrigerant, e.g. calcium chloride or liquid nitrogen, through tubes in the ground; as the ground freezes around closely spaced tubes, a continuous frozen zone may be formed. This is an expensive technique, but suitable for mixed *\*strata* as the results are more predictable than *\*dewatering* or *\*grouting*.

**artificial rain** Rain, or increased rain, produced by seeding clouds artificially with 'dry ice' (frozen carbon dioxide), silver iodide, or other appropriate particles, which act as condensation nuclei.

**artificial recharge** A process whereby the amount of water in an **\*aquifer** is supplemented by engineered, as opposed to natural, means. Artificial recharge may be through **\*boreholes**, purpose-built ponds, or simply by diverting more water on to the surface **\*catchment** of the aquifer. Artificial recharge may be implemented as part of a conjunctive use scheme.

**Artinskian** 1. An **\*age** in the **\*Permian** epoch, preceded by the **\*Sakmarian**, followed by the **\*Kungurian**, and dated at 284.4–275.6 Ma ago (Int. Commission on Stratigraphy, 2004). 2. The name of the corresponding eastern European **\*stage**, which is roughly contemporaneous with the upper **\*Rotliegendes** (western Europe), lower/middle **\*Leonardian** (N. America), and the Bitaunian (New Zealand).

**Artiodactyla (artiodactyls)** (cohort **\*Ferungulata**) Even-toed ungulates, an order of mammals that includes the living camels, pigs, and ruminants. Descended from the **\*Condylarthra**, they underwent a spectacular burst of **\*adaptive radiation** in **\*Eocene** and early **\*Oligocene** times, largely replacing the initially more numerous **\*Perissodactyla**. The ankle bone (astragalus) is specialized in artiodactyls to give better spring. The axis of the foot is paraxonic (passing between the third and fourth digits). In primitive, four-toed types, e.g. the pig, the first digit is absent; in advanced forms the second and fifth digits are also reduced or lost. Early forms had an unspecialized dentition, but in the course of evolution the upper incisors were lost in some species, the lower incisors biting against the hardened gum of the upper jaw, an adaptation to a herbivorous diet.

**Arundian** A **\*stage** (341–339 Ma ago) of the **\*Visean** epoch, underlain by the **\*Chadian** and overlain by the **\*Holkerian**.

**Asaphida** An order of **\*Trilobita** that lived from the Upper **\*Cambrian** to **\*Silurian**. There were six suborders.

**asbestos** Fibrous varieties of **\*amphibole** and **\*serpentine**, including chrysotile (fibrous serpentine), **\*actinolite** (asbestos proper), amosite (a variety of **\*anthophyllite**), and crocidolite (blue asbestos; a variety of **\*riebeckite**). Ancient civilizations referred to asbestos cloth as *amianthus*

(from the Greek word meaning ‘undefiled’), because it could be cleaned by throwing it into a fire and all varieties of asbestos have great heat-resistant properties; varieties with fibres long enough to be spun and woven were formerly used commercially for heat-resistant cements, cladding, and insulation material, and for asbestos corrugated sheets. However, many of its uses, particularly as brake pads for cars and as pipe and ceiling insulation materials, have been banned, because the inhalation of asbestos dust (small, airborne, needle-like fibres) can cause serious lung diseases (e.g. asbestosis) and contribute to pneumoconiosis. Health concerns have led to a worldwide reduction in asbestos production to below 3 million tonnes annually.

**Asbian** A *\*stage* (336.4–336 Ma ago) of the *\*Visean* epoch, underlain by the *\*Holkerian* and overlain by the *\*Chesterian*.

**aseismic** Free from *\*earthquakes*.

**aseismic margin** See PASSIVE MARGIN.

**aseismic ridge** Long, linear, inactive, volcanic, topographic feature found in many deep-ocean basins. An example is the Walvis Ridge in the south-eastern Atlantic, which extends for 3000 km and in places reaches a height of 2 km above the deep-ocean floor.

**ash** *\*Tephra* less than 2 mm in size.

**ash-cloud surge** See SURGE.

**ash cone** See SCORIA CONE.

**ash-flow** See PYROCLASTIC FLOW.

**Ashgill** A *\*series* (443.7–449 Ma ago) of the Upper *\*Ordovician*, underlain by the *\*Caradoc* and overlain by the *\*Llandovery* (*\*Silurian*).

**asiderite** See STONY METEORITE.

**ASIM** See ATMOSPHERE-SPACE INTERACTIONS MONITOR.

**ASNARO** See ADVANCED SATELLITE WITH NEW SYSTEM ARCHITECTURE FOR OBSERVATION.

**asperitas** See CLOUD CLASSIFICATION.

**asphalt** Brown or black, solid or semi-solid, bituminous substance made almost entirely of carbon and hydrogen. It melts between 65 °C and 95 °C and is soluble in carbon disulphide. It is formed by the evaporation of volatile **\*hydrocarbons** and occurs in oil-bearing rocks, e.g. in Trinidad, or as a residue from **\*petroleum** refining.

**asphaltite** See GILSONITE.

**assay** The analysis of **\*minerals** and mine products to determine the concentrations of their components.

**Asselian** 1. An **\*age** (299–294.6 Ma ago) in the Early **\*Permian** epoch, preceded by the **\*Stephanian**, **\*Gzhelian**, and Noginskian (**\*Carboniferous**), followed by the **\*Sakmarian** (Int. Commission on Stratigraphy, 2004). 2. The name of the corresponding eastern European **\*stage** which, because of lower boundary uncertainties, has also been considered as part of the Carboniferous system. It is roughly contemporaneous with the lower **\*Wolfcampian** and lower **\*Rotliegendes** (western Europe), and with the lower Somoholoan (New Zealand). See also SAKMARIAN.

**assemblage zone (coenozone, faunizone)** **\*Biostratigraphic** unit or level of **\*strata** characterized by a particular assemblage of animals or plants. An assemblage zone is named after one or more of the distinguishing **\*fossils** present, which are chosen without regard for their total time ranges, so that the assemblage is of purely environmental significance. Compare CONCURRENT RANGE ZONE.

**assimilation** In **\*petrology**, the processes of melting and solution by which wall rock is incorporated into **\*magma**. Partial or complete melting may occur at **\*contacts** and at depth in detached blocks or **\*xenoliths**. Assimilation causing changes in composition of the original magma may lead to hybrid or contaminated rocks. The term does not imply any particular mechanism. See PARTIAL MELTING.

**assimilation-fractional crystallization (AFC)** An important process in **\*igneous \*petrology** whereby **\*melts** with widely differing isotopic and **\*trace** elements can be produced. When a primitive **\*magma**, such as a **\*basalt**, invades crustal rocks, portions of the **\*country** rock may become detached and included in the magma as **\*xenoliths**. Because of the high temperature and thermal capacity of the basalt, it is capable of melting a

proportion of the country rock. In doing so it loses some of its own heat and thus a proportion of the magma crystallizes. The composition of the resulting magma is determined by the relative amounts of magma and country rock initially present; the rates at which assimilation and crystallization proceed; and by the *\*partition* coefficients of the various elements between solid and liquid.

**astatic magnetometer** See MAGNETOMETER.

**ASTERIA** See ARCSECOND SPACE TELESCOPE ENABLING RESEARCH IN ASTROPHYSICS.

**Asteriacites** An *\*ichnogenus* of *\*cubichnia*.

**asteroid** A small rocky or metallic body orbiting the *\*Sun* in the asteroid belt between the orbits of *\*Mars* and *\*Jupiter* (from about 2.2 to 3.2 AU (see ASTRONOMICAL UNIT), with one family (Trojans) at 5 AU occupying two of the jovian *\*Lagrangian Points*). Two families (Apollo and Aten, as well as some Amor objects) have orbits that cross *\*Earth's* orbit. The largest asteroid is 1 Ceres (diameter  $987 \pm 150$  km). About 30 exceed 200 km in diameter and over 3000 have been identified. *\*Meteorites* are probable samples of the asteroid belt. See APOLLO (1862); CASTALIA (4769); CERES (1); CHIRON (2060); EROS (433); GASpra (951); GEOGRAPHOS (1620); ICARUS (1566); IDA (243); JUNO (3); MCAULIFFE (3352); MATHILDE (253); MIMISTROBELL (3840); NEREUS (4660); PALLAS (2); SHIPKA (2530); TOUTATIS (4179); VESTA (4).

**Asterosoma** A mounded, lobed, somewhat star-shaped *\*trace fossil*. The central tube was occupied by an organism; the surrounding, detrital lobes are the result of feeding processes.

**Asteroxylon** An early lycopsid (*\*Lycopsidea*) plant that occurs in the Rhynie Chert, Aberdeenshire, Scotland (Lower *\*Devonian*). It has dichotomous rhizomes and the aerial extensions bear leaf-like enations (outgrowths).

**Asterozoa (starfish, brittle stars)** (phylum *\*Echinodermata*) Subphylum whose members have *\*radial symmetry* of projecting rays, and a star-shaped body. See also AMBULACRUM.

**asthenosphere** The weak zone within the upper *\*mantle*, underlying the *\*lithosphere*, where the mantle rocks deform by plastic flow in response to applied stresses of the order of 100 MPa. It is commonly considered to be coincident with the upper-mantle seismic *\*low-velocity zone*, but this is probably valid only for the oceanic sectors of the mantle. *\*Viscosity* is of the order of  $10^{21-22}$  poise, i.e. the same as the underlying mantle, but it is much more 'fluid' than the overlying lithosphere. Originally it was recognized as a possible explanation for *\*isostatic* behaviour, and it is generally recognized as a mantle zone within which convective motions take place. The depth of *\*earthquake* foci in *\*subduction zones* suggests that descending convection limbs penetrate to 700 km, just above the upper mantle–lower mantle boundary. Rising limbs of asthenospheric mantle convection are located under surface spreading centres (*\*mid-oceanic ridges*).

**astogenetic heterochrony** Among colonial animals, *\*heterochrony* that affects the colony as a whole. Compare MOSAIC HETEROCHRONY; ONTOGENETIC HETEROCHRONY.

**astra** See ASTRUM.

**astraeoid** A condition that occurs in massive *\*corals*. Massive corals are composed of closely packed *\*corallites* and the individuals become polygonal in shape. In the astraeoid condition the walls of the corallites become reduced or lost but the *\*septa* remain entire. See COMPOUND CORALS.

**astragalus** The ankle bone.

**astrobleme** Literally, 'star-wound'; a terrestrial crater formed by *\*meteorite*, *\*asteroidal*, or *\*cometary* impact. The term is generally used for large craters (more than 10 km diameter), but is not in common usage.

**astrogeology** See PLANETARY GEOLOGY.

**astronomical unit (AU)** The average distance between the *\*Earth* and *\*Sun*; a unit of measurement equal to 149 597 870 km.

**astrum** (*pl.* astra) On *\*Venus*, a surface feature with a radial pattern.

**A-subduction** The movement of one continental lithospheric *\*plate* under another (*\*subduction*) in a *\*collision zone*, with the separation of part or all of the upper *\*crust* from the lower crust and *\*mantle*. The denser material subducts normally whilst the buoyant material can underthrust or overthrust the crust of the overriding plate. A-subduction, named after O. Ampferer, has also been called ‘delamination’, and is contrasted with B-subduction, named after Hugo *\*Benioff*, in which oceanic *\*lithosphere* subducts. A-subduction is postulated to involve *\*shortening* of a maximum of only a few hundred kilometres, whereas B-subduction can recycle thousands of kilometres of *\*oceanic* crust and upper mantle.

**asymmetrical fold** A *\*fold* in which the *\*axial plane* is inclined relative to the median plane, and adjacent limbs *\*dip* in opposite directions.

**asymmetric valley** Valley that has one side steeper than the other, the opposing sides standing at significantly contrasting angles. This may be due to geologic structure, or to variation in the nature and intensity of erosional processes. Such valleys are common in past and present *\*periglacial* environments, where aspect has a significant effect on the nature of frost-based processes.

**Atdabanian** A Russian-Kazakhstanian *\*stage* of the *\*Early Cambrian* (Caerfai epoch), underlain by the *\*Tommotian* and overlain by the *\*Botomian*, and dated at 530–524 Ma ago (Int. Commission on Stratigraphy, 2004).

**Athabasca** See TAR SAND.

**Atlantic** Period in post-glacial times (i.e. post-*\*Devensian* or *\*Flandrian*) from about 7500 to 5000 BP which, according to pollen evidence, was warmer than the present, and moist; with oceanic climatic conditions prevailing throughout north-western Europe. It corresponds to *\*Pollen Zone VIIa*, which throughout north-western Europe is characterized by the most thermophilous (warmth-loving) species found in post-glacial pollen records. The climatic optimum of the post-glacial, or current Flandrian Interglacial, is dated to the early Atlantic period. Compare BOREAL. See POLLEN ANALYSIS.

**Atlantic conveyor** An informal name sometimes applied to the *\*meridional overturning circulation*.

**Atlantic Ocean** One of the main oceanic areas of the world. It is relatively shallow, having an average depth of 3310 m; and it is the warmest (average temperature 3.73 °C) and most saline (average *\*salinity* 34.9 parts per thousand) of the major oceans.

**Atlantic Province (Acado-Baltic Province, European Province)** A subdivision of the early *\*Cambrian* olenellid *\*trilobite* *\*fauna*. The trilobite faunas of the early Cambrian can be divided into two main regional groups: the olenellid fauna found in north-western Europe and N. America, and the redlichiids in Asia, Australia, and N. Africa. The olenellid fauna is also subdivided into two provinces: the Atlantic Province on the southern and eastern flank of the *\*Iapetus Ocean*, and the *\*Pacific* (or American) Province on the northern and western margins. The names Atlantic Province and Pacific Province have also been applied to the *\*Ordovician* trilobite and *\*graptolite* faunas in the same areas.

**Atlantic-type coast (transverse-type coast)** A coast characterized by subsidences and fractures that cut across the grain of the folded mountain formations inland. Typically, such a coast borders a relatively young ocean that is widening because of *\*sea-floor spreading*. This type of coast was first recognized by Eduard *\*Suess*. Compare PACIFIC-TYPE COAST.

**Atlantic-type margin** See PASSIVE MARGIN.

**Atlantis Massif** A dome-shaped *\*massif* approximately 16 km across and rising about 4270 m from the ocean floor at approximately 30.13°N 42.13°W, close to the intersection of the *\*Mid-Atlantic Ridge* and the Atlantis *\*transform fault*.

**Atlas (Saturn XV)** One of the lesser satellites of *\*Saturn*, discovered in 1980 by *\*Voyager* 1, with a radius measuring 18.5 × 17.2 × 13.5 km; visual albedo 0.9.

**atlas vertebra** See VERTEBRA.

**atmometer** An instrument for measuring evaporation. It is normally in the form of an open-ended glass tube from which water can evaporate.

**atmophile** Applied to the gaseous elements most typical of, and concentrated in, the Earth's atmosphere; e.g. H, C, N, O, I, and inert gases. They may occur in an uncombined state or combined as, for example, in

water (H<sub>2</sub>O), carbon dioxide (CO<sub>2</sub>), and methane (CH<sub>4</sub>). *Compare* [BIOPHILE](#); [LITHOPHILE](#); [CHALCOPHILE](#); [SIDEROPHILE](#).

**atmosphere** **1.** Air surrounding the Earth. The atmosphere has no precise upper limit, but for all practical purposes the absolute top can be regarded as being at about 200 km. The density of the atmosphere decreases rapidly with height, and about three-quarters of the mass of the atmosphere is contained within the lowest major layer, the [\\*troposphere](#), whose depth varies between about 10 km and 17 km, being generally smaller further from the equator. **2.** Unit of pressure (abbreviation: atm). Its value is approximately the average pressure of the atmosphere at sea level, the figure adopted being the pressure at sea level in the International Standard Atmosphere (760 mm of mercury, or 1013.25 mb). In SI units, 1 atm = 101 325 Pa. *See also* [ATMOSPHERIC STRUCTURE](#).

**Atmosphere-Space Interactions Monitor (ASIM)** An assembly of instruments from the [\\*European Space Agency](#) (ESA) that observe transient luminous events in the upper atmosphere that are linked to thunderstorms in the lower atmosphere. The assembly was launched on 2 April 2018 to the International Space Station.

**Atmospheric Neutral Density Experiment (ANDE)** A low-cost space mission by the US Naval Research Laboratory that monitors the neutral density of the [\\*thermosphere](#) at a height of about 400 km, in order to improve the determination of resident space objects. The experiment comprises two [\\*minisatellites](#) that were launched into near-circular orbit from the Space Shuttle on 10 December 2006 and retired in 2008. A follow-up mission, ANDE-2, comprising two [\\*microsatellites](#) was launched on 30 July 2009 and ended in 2010.

**atmospheric pollution** Solid and gaseous contaminants in the atmosphere which occur as dust, smoke, or sulphur dioxide and other gases, particularly from the combustion of fossil fuels and certain industrial processes. Air pollution is most marked in urban areas. *See also* [PHOTOCHEMICAL SMOG](#).

**atmospheric pressure** Downward force exerted by the weight of the overlying [\\*atmosphere](#), expressed per unit area in a given horizontal cross-section. Pressure varies throughout the atmosphere, owing to the distribution of mass; there are small diurnal variations of partly tidal and

partly thermal origin, as well as bigger changes associated with the passage of depressions and anticyclones. Atmospheric pressure is measured in pascals or millibars (mb), 1 mb being equal to 100 kilopascals (kPa). Measurements are usually made with a mercury barometer. The overall global average pressure at sea level is 101.325 MPa (1013.25 mb), but as air is readily compressible, pressure decreases exponentially with altitude.

**atmospheric shimmer** The effect observed when light passes through moving air masses which have differing *\*refractive indices*. This results in effects such as star twinkling and ultimately limits the resolution of any *\*remote sensing* system.

**atmospheric structure** The broadly horizontal layering of the *\*atmosphere*, the layers being distinguished by differences in the rate of change of temperature with height, which either favour or discourage the development of vertical exchanges (convection). From the surface of the Earth upwards the layers are: (a) the *\*troposphere*, in which convection is often prominent, especially over warm regions, extending to the *\*tropopause* at a somewhat variable height, generally about 8 km over the poles, 11 km in middle latitudes, and 16 km over the equator; (b) the *\*stratosphere*, in which there is much less vertical motion, and which extends from the tropopause to about 50 km at the *\*stratopause*; (c) the *\*mesosphere*, in which there is once again more convection, extending from the stratopause to a height of about 80 km at the *\*mesopause*; and (d) the *\*thermosphere*, extending from the mesopause to the effective limit of the atmosphere, at about 200 km.

**atmospheric 'window'** The range of wavelengths (about 8.5–11  $\mu\text{m}$ ) at which radiation is only slightly absorbed by water vapour. Terrestrial radiation within this range may escape into space unless it is absorbed by cloud (water droplets can absorb in this range). *See also* GREENHOUSE EFFECT; TERRESTRIAL RADIATION.

**Atokan (Derryan)** A *\*series* (311.7–308 Ma ago) in the *\*Pennsylvanian* of N. America, underlain by the *\*Morrowan* and overlain by the *\*Desmoinesian* series. It is roughly contemporaneous with the Vereiskian and Kashirskian stages of the *\*Moscovian* series.

**atoll** Ring-shaped organic *\*reef* that encloses or almost encloses a *\*lagoon*, and which is surrounded by the open sea. The reef may be built of *\*coral*

and/or calcareous **\*algae**. An atoll is built on an existing structure such as an extinct, submerged volcano.

**atollon** The local name for an **\*atoll** in the Maldiv Islands.

**atomic absorption spectrometry** Instrumental technique for the chemical analysis of material. The basic theory was known as long ago as the mid-19th century, and utilizes the observation that certain wavelengths of radiation emitted by excited atoms are strongly absorbed by unexcited atoms of the same element (emission **\*spectrum**). The radiation will be reduced if it passes through an area containing such unexcited atoms. This reduction can be measured by a detector, and gives the **\*concentration** of the element.

**atomic number** Chemical elements are composed of atoms, all of which are held together by electrical charges. Every atom has a relatively heavy **\*nucleus** that is composed of protons (with a positive charge) and neutrons (neutral particles). Orbiting the nucleus there are a number of exceptionally light **\*electrons**, whose negative charges balance the positive charge provided by the protons. The number of protons in the structure provides the atomic number; atoms having the same atomic number belong to the same chemical element.

**A-train (Afternoon Constellation)** A series of Earth-observing satellites operated by NASA and other national space agencies. All the satellites are in **\*polar orbit** and they cross the equator at about 13.30 local time, within seconds or minutes of each other. There are six satellites in the train: **\*Aqua**, **\*Aura**, **\*CALIPSO**, and **\*CloudSat**, GCOM-W1 “SHIZUKU”, and OCO-2. A seventh satellite, Glory, was lost due to a launch failure on 4 March 2011. **\*PARASOL** was part of the A-train from 2004 until 2009.



<http://atrain.nasa.gov/>

- A NASA mission of several satellites placed in polar orbit to study Earth-system science.

**attapulгите (palygorskite)** An **\*inosilicate** mineral (a magnesium aluminium silicate) that is a type of **\*fuller's** earth. Its name is taken from the town of Attapulgitus, in the extreme south-west corner of Georgia, USA, where the mineral is abundant. It has medicinal uses derived from its

capacity for binding to acids and certain toxic substances in the stomach and digestive tract.

**attenuation** The reduction in amplitude or energy of a signal. Attenuation of *\*seismic waves* (seismic attenuation) occurs as a result of spherical divergence, *\*absorption*, energy losses at interfaces through *\*reflection* and *\*refraction*, and by internal scatterers. For electromagnetic waves see *SKIN DEPTH*.

**Atterberg limits** Series of thresholds which are observed when the water content of a soil is steadily changed. The ‘contraction limit’ occurs when sufficient water is added to a dry soil for contraction cracks to close. The addition of further water leads to plastic deformation at the ‘plastic limit’. The ‘liquid limit’ occurs when just enough water is then added for the soil to behave like a liquid. Knowledge of these limits is important for understanding and predicting hillslope failure. The difference in percentage water content between the liquid limit and the plastic limit is called the ‘plasticity index’. The limits were described by the Swedish soil scientist Albert Mauritz Atterberg (1846–1916).

**attitude** **1.** Of a bed or other planar feature, the disposition with respect to the horizontal and compass bearings; these are obtained by measuring the *\*dip* and *\*strike* of the bed respectively. **2.** Of a *\*fold*, the overall disposition; this is defined by measuring the dip and strike of the *\*axial plane*, and the *\*trend* and *\*plunge* of the *\*hinge* line.

**atritus** Minute particles of plant material that result from attrition by living organisms in vegetable deposits, especially in swamps and bogs.

**AU** See *ASTRONOMICAL UNIT*.

**aubrite** An *\*enstatite* *\*pyroxene*-rich, *\*achondritic* *\*meteorite* that is poor in calcium. The enstatite, which often forms large *\*crystals*, has very low FeO/(FeO + MgO) ratios, Aubrites may contain up to 1.2% elemental silicon included within rare Fe–Ni grains, and appear to have close affinities with the similarly highly reduced enstatite *\*chondrites*.

**augen-gneiss** A medium- to coarse-grained, banded, *\*regional metamorphic* rock composed mainly of *\*quartz* and *\*feldspar* with *\*hornblende* and *\*mica* also present in variable quantities, and characterized by large ovoidal *\*megacrysts* of feldspar known as ‘augen’

(derived from the German for 'eyes'). The banding is due to variations in the modal proportions (see MODAL ANALYSIS) of the \*mineral \*phases constituting the rock. The best-developed augen-gneisses are formed by high-grade \*metamorphism of aluminous sediments. See also GNEISS.

**auger** Tool used primarily for soil sampling, but also for sampling \*peat and other \*unconsolidated sediments. The simplest and most universal form has a screw head to bore the soil or sediment. Alternative auger heads are available for more specialized needs. Standard augers sample to one metre depth, but extension rods can be attached enabling sampling at deeper levels.

**augite** An important member of the \*pyroxene group of silicate minerals, occupying a field of composition within the \*calc-alkaline pyroxenes between the \*diopsides and the \*pigeonite field, and with the formula  $\text{Ca}(\text{Mg,Fe})\text{Si}_2\text{O}_6$  (Na may substitute for Ca and Al for Si; and a large increase in the amount of Na and  $\text{Fe}^{3+}$  (for  $\text{Fe}^{2+}$ ) gives aegirine augite  $(\text{Na,Ca})(\text{Fe}^{3+},\text{Mg,Fe}^{2+})\text{Si}_2\text{O}_6$ , an important pyroxene in \*alkaline rocks); sp. gr. 3.3; \*hardness 6.0; \*monoclinic; greenish-black; \*crystals short, \*prismatic; a common constituent of \*basic igneous rocks but may also occur in high-grade \*metamorphic rocks. See also CLINOPYROXENE.

**augite-minette** See MINETTE.

**Aulacocerida** See BELEMNITIDA.

**aulacogen** A long-lived, \*sediment-filled \*graben oriented at a high angle to either a neighbouring modern ocean or a neighbouring \*orogenic belt. The sediments in an aulacogen are largely characterized by a lack of major deformation, although late \*strike-slip movement on the boundary \*faults is known to deform the sediments extensively. Aulacogens have been interpreted as forming within failed rifts (see RIFT VALLEY) of \*triple junctions and thus are taken to indicate \*plate tectonic activity. They have been identified as far back as the early \*Proterozoic and are of worldwide distribution.

**aulodont** Applied to \*echinoids with a lantern (see ARISTOTLE'S LANTERN) characterized by teeth which are longitudinally grooved and broadly U-shaped in cross-section.

**Aura** A \*NASA satellite, launched into a Sun-synchronous orbit from California on 15 July 2004, as part of the Earth Science Projects Division. Aura (Latin for breeze) carries four instruments that monitor the chemistry and dynamics of the atmosphere from the surface to the \*mesosphere. It is one of the four satellites that comprise the \*A-Train.



[http://www.nasa.gov/mission\\_pages/aura/main/index.html](http://www.nasa.gov/mission_pages/aura/main/index.html)

- A NASA mission to monitor complex scientific interactions affecting the globe.

**aureole** The luminous white or bluish disc, surrounded by a brown ring, sometimes observed directly surrounding the Sun or Moon. The term is also used to describe the bright area with no definite boundary commonly seen surrounding the Sun in a clear sky. *See also* CORONA; CONTACT AUREOLE.

**aurora** Illumination of the sky, sometimes in brilliant colours, as a result of high-speed solar particles entering the \*ionosphere (at a height of 100–130 km) and releasing electrons from air molecules by excitation. The re-establishment of molecules leads to the emission of light, especially red- and green-coloured light, e.g. in arcs or bands over large areas. The effect is called ‘aurora borealis’ or ‘northern lights’ in the northern hemisphere and ‘aurora australis’ or ‘southern lights’ in the southern hemisphere. Such atmospheric disturbances occur in relation to disturbances on the Sun in the course of the sunspot cycle.

**Australian faunal realm** Region distinguished by a unique \*marsupial fauna, including herbivores, carnivores, and insectivores. These evolved in isolation from the placental mammals which now dominate the other continental faunas. In addition to marsupials there are also very primitive mammals (monotremes): the spiny anteater and the platypus; and small rodents which are relatively recent (probably \*Miocene) immigrants.

**australopithecines** Literally, ‘southern apes’, early members of the human lineage that lived from about 4 to about 1 million years ago in Africa. They were adapted for bipedal locomotion, had a high brachial index (forearm:upper arm ratio) relative to other hominids, were sexually dimorphic, were 1.2–1.5 m tall, had a cranial capacity of 350–600 cc, and had large premolar and molar teeth and relatively small incisors and

canines. The so-called ‘robust australopithecines’ are nowadays placed in a separate genus, *Paranthropus*. The other (‘gracile’) australopithecines are also a very diverse group of species, some very primitive and perhaps ancestral to all later hominins, others probably specialized sidelines. The species usually recognized are *Australopithecus anamensis* (3.9–4.1 million years BP) and *A. afarensis* (3.75–3.0 million years BP) from East Africa, *A. bahrelghazali* (about 3.4–3.0 million years BP) from West Africa, and *A. africanus* (about 3.0–2.4 million years BP) from South Africa. Probably some of these species should be placed in different genera.

***Australopithecus*** See AUSTRALOPITHECINES.

**autapomorphy** The possession of an **\*apomorphic** **\*character state** that is unique to a particular species or lineage in the group under consideration.

**authigenic** Applied to materials (**\*minerals**, **\*cements**, etc.) that formed in the rock of which they are a part during, or soon after, its deposition. Compare ALLOGENIC; ALLOCHEMICAL.

**autobrecciated lava** A viscous, commonly silica-rich, **\*lava** flow with a congealed crust which has been broken up and fragmented by the continued movement of molten lava within the flow interior. Stressed and deformed by the movement, the crust may fracture in a brittle manner, producing angular, smooth-faced blocks able to weld together if they are hot enough but which otherwise become incorporated into the moving interior of the flow.

**autochthonous** Indigenous; applied to a material (e.g. **\*dripstone**, **\*coal**) that was formed in its present position. No significant transport has been involved. Compare ALLOCHTHONOUS.

**autoclast** A **\*clast** formed by friction from flowing **\*lava** or autobrecciation (see AUTOBRECCIATED LAVA). Compare ALLOCLAST; EPICLAST; and HYDROCLAST.

**autocorrelation** The **\*correlation** of a wave-form with itself; a special case of **\*cross-correlation**. The autocorrelation function (ACF) is especially useful in the identification of **\*multiples** within a **\*seismic record**. While the ACF contains all the amplitude and frequency information of the original wave-form, it possesses no phase information.

**autocyclic mechanisms** Events responsible for the accumulation of sediments that are part of the sedimentary system itself (e.g. size and configuration of a river channel). *Compare* [ALLOCYCLIC MECHANISMS](#).

**automatic point counter** An electronic control panel designed to count the number of times a particular *\*mineral* is recorded on multiple, closely spaced traverses of a *\*thin* section viewed down a *\*polarizing* microscope. The control panel, which contains a bank of recording buttons, each assigned by the user to a particular mineral, is linked to a thin-section holder mounted on to the *\*stage* of a polarizing microscope. The holder is designed to move the thin section by a predetermined distance (variable from 0.001 to 1 mm) in one direction, after the user has pressed the control panel button assigned to the particular mineral observed under the *\*eyepiece* *\*cross-wires*. At the end of a complete traverse, the holder automatically resets by a predetermined distance to start the next traverse. On completion of all the traverses, the control panel shows the number of times each mineral was recorded and can recalculate this as a percentage of the total number of points. This gives the percentage volume of each mineral recorded in the rock.

**automatic weather station** Meteorological station that records measurements of *\*atmospheric* pressure, temperature, *\*humidity*, and wind, and details of weather conditions, and transmits them automatically to a central base.

**autometamorphism (autometasomatism)** The recrystallization of *\*igneous* rocks as they cool by the action of *\*volatiles* present within the rocks themselves.

**autometasomatism** *See* [AUTOMETAMORPHISM](#).

**AutoNaut** A wave-powered, autonomous unmanned surface vessel (USV), 3.5 m long and 0.43 m beam, with a mass of 100 kg, developed by the *\*European Space Agency's* Business Incubation Center, at Harwell, UK. At either end, the boat has two pairs of foils mounted on struts, and it has a small thruster. It carries sensors to monitor weather, marine life (it can detect cetacean vocalizations over a long distance), pollution, and the marine environment, and it communicates with orbiting satellites, using a lithium-ion battery that recharges from solar cells. It moves at up to 3 km/h.

In March 2018 AutoNaut produced a new range of upgraded and slightly larger USVs.



<http://www.autonautusv.com>

- AutoNaut home page.

**autosuspension** In an active *\*turbidity* current, a feedback mechanism whereby: turbulence maintains the suspended load; the suspended load causes the high density of the suspension; the high density of the suspension causes the flow; the flow causes turbulence.

**autotheca** One of the three types of graptolite *\*thecae*, possibly containing the female zooid. See [GRAPTOLITHINA](#); [GRAPTOLOIDEA](#); [DENDROIDEA](#). Compare [BITHECA](#); [STOLOTHECA](#).

**Autunian** A lesser-used stratigraphic name for the lowest of three divisions of the *\*Permian* of Europe, the others being the *\*Saxonian* (mid) and *\*Thuringian* (upper). Together, the Autunian and Saxonian form the Lower Permian.

**autunite** *\*Secondary mineral*,  $\text{Ca}(\text{UO}_2)_2(\text{PO}_4)_2 \cdot 10\text{--}12\text{H}_2\text{O}$ ; sp. gr. 3.1; *\*hardness* 2.0–2.5; *\*tetragonal*; usually bright lemon-yellow to greenish-yellow; yellow *\*streak*; *\*vitreous* *\*lustre*; *\*crystals* *\*tabular*, square, forming foliated, scaly masses; *\*cleavage* perfect, basal; radioactive; fluorescent; occurs with *\*torbernite* in the oxidized zone of mineral veins.

**auxiliary reference section** See [HYPOSTRATOTYPE](#).

**available nutrients** Any elements or compounds in the soil solution that can be absorbed readily into plant roots and that function as nutrients to growing plants. The available amount is usually much less than the total amount of that plant nutrient in the soil.

**available relief** The part of a landscape that is higher than the floors of the main valleys. It is therefore available for destruction by the agents of *\*erosion*, controlled by the local base level. It is measured by the vertical distance between hilltops and valley floors.

**available water** In soil, the water that can be absorbed readily by plant roots. It is usually taken to be water held in the soil under a pressure of 0.3

to about 15 bars.

**avalanche** See MASS-WASTING.

**avalanche wind** A blast of often very destructive air ahead of a descending avalanche.

**Avalonian orogeny** An episode of mountain building, named after the Avalon Peninsula of Newfoundland, that occurred about 650–500 Ma ago (\*Cambrian to \*Ordovician) as a result of rifting (see RIFT) and \*volcanicity associated with the opening of the Atlantic. Its rocks are found intermittently from Georgia to Newfoundland. See APPALACHIAN OROGENIC BELT.

**aventurescence** See AVENTURINE.

**aventurine (adventurine, Indian jade)** A translucent variety of \*quartz that contains platy fragments, usually of \*mica, that give it a glistening quality, called aventurescence. It is usually green or blue-green, but the inclusion of \*hematite or \*goethite makes it orange or brown. Aventurine can be polished and used in jewellery or to make ornaments.

**average** In statistics, a summary of the data using a single value. The data may be summarized by the \*mean, \*median, or \*mode values. See also VARIANCE.

**average velocity (time-averaged velocity,  $\bar{v}$ )** The ratio of a given depth divided by the travel time to that depth, usually assuming straight ray paths and parallel layering. Thus  $\bar{v} = z_n/t_n$ , where  $z_n$  is the depth of the top  $n$  layers and  $t_n$  the single-travel time through those  $n$  layers. Also,  $\bar{v} = \Sigma z_i/\Sigma t_i$  where  $z_i$  and  $t_i$  are the thickness of and single-travel time through the  $i$ th layer respectively, and  $\Sigma$  is the total thickness and total travel time respectively of the overlying layers.


**Aves (birds)** (subphylum \*Vertebrata, superclass \*Gnathostomata) The class that comprises all the birds. The late \*Jurassic \*Archaeopteryx lithographica is still the best-known \*Mesozoic bird, but others have been described since the 1980s: \*Noguerornis, from the lowermost \*Cretaceous of Spain; the slightly later and more advanced Concornis and \*Iberomesornis, known by complete skeletons, also from Spain; and, also

early Cretaceous, *Sinornis* and *Cathayornis* from China. There were also some curious, specialized, Late Cretaceous birds, such as the flightless *Mononykus* from Central Asia, in which the forelimbs were reduced to stubby claws, and *Hesperornis*, a diving form. All these early birds had teeth and long, bony tails. Birds arose from within the theropod dinosaurs (**\*Theropoda**) and so should properly be classified as a subgroup of them; those closest were the Dromaeosauridae (the family which includes the famous *Velociraptor* of *Jurassic Park*).

**Avicenna** (Abu Ali al-Husayn Ibn Abdallah Ibn Sina) (980–1037) Avicenna was an Arab physician and philosopher. His ‘geological’ ideas were published in *Liber De Mineralibus*, which was attributed to Aristotle, and influential up to about 1500. He wrote about earthquakes, erosion of valleys, the deposition of sediments, etc.

**Avogadro constant (Avogadro number)** The number of molecules, atoms, or **\*ions** in one **\*mole** of a substance:  $6.02252 \times 10^{23}$  per mole. It is derived from the number of atoms of the pure isotope  $^{12}\text{C}$  in 12 grams of that substance and is the reciprocal of atomic mass in grams.

**avulsion** Lateral displacement of a stream from its main channel into a new course across its **\*floodplain**. Normally it is a result of the instability caused by **\*channel aggradation**. The avulsion of a stream into an adjacent valley may explain some cases of apparent **\*river capture**.



**Avulsion**

**Awamoan** A New Zealand *\*stage* (20–17.5 Ma ago) of the Early *\*Miocene*, preceded by the *\*Hutchinsonian* and followed by the *\*Altonian*.

**axial modulus ( $\phi$ )** The ratio of longitudinal stress to uniaxial longitudinal strain, i.e. where there is no lateral strain. This is a special form of *\*Young's modulus*.

**axial plane** The plane that bisects the angle between the two limbs of a *\*fold*.

**axial plane cleavage** The *\*cleavage* within a *\*fold* which is systematically oriented with respect to the *\*axial plane*, and which lies parallel or subparallel to the axial plane, particularly in the *\*hinge* region. Away from this region the cleavage commonly deviates from its parallelism and forms a symmetric, convergent, or divergent fan about the *\*axial surface*.

**axial ratio (intercept ratio)** In the study of crystals, the position of a *\*crystal* face in space is given by the intercepts the face or plane makes on three (or four) imaginary lines, called 'crystallographic axes'. The X-ray crystallographer can measure the 'unit-cell' dimensions in ångstrom units (Å), and the axial ratios express the relative, and not the absolute, lengths of the cell edges corresponding to the crystallographic axes. These ratios (or '*\*parameters*') are often expressed reciprocally as 'indices', e.g. *\*Miller indices*.

	Crystallographic Axes		
	a (x)	b (y)	c (z)
Intercepts of crystal face DEF in ångstroms, on a, b, and c axes, measured from origin.	OD 20Å	OE 10Å	OF 40Å
If b intercept is made equal to 1 the axial ratio is obtained for the crystal.	20 10 2	10 10 :1	40 10 :4
Indices are obtained by dividing the intercepts of face DEF into those of the parametral plane, which is a face of the unit form with intercepts (111).	1 2	1 1	1 4
Miller's indices of face DEF are obtained by removing fractions.	2	4	1

Note that if face DEF is selected as the *\*parametral plane*, then its indices would be  $\frac{2}{2}$ ,  $\frac{1}{1}$ ,  $\frac{4}{4} = 111$ .

**axial rift** See *MEDIAN VALLEY*.

**axial surface** The surface which joins the *\*hinge* lines on adjacent folded surfaces. Where it is planar it is called the *\*axial plane*.

**axial tilt** The angle by which the rotational axis of the Earth differs from a right angle to the orbital plane; this angle varies between 22.1° and 24.5° over a cycle of 41 000 years and at present is about 23.45°. See *MILANKOVITCH CYCLES*.

**axial trace** A line that marks the intersection of the *\*axial surface* of a *\*fold* with any other plane or surface. In practical terms it is the intersection of the axial surface with the surface of the Earth, or with a vertical profile through a fold sequence.

**axial trough** See *MEDIAN VALLEY*.

**axinite** An uncommon borosilicate *\*mineral* with variable composition, generally given as  $(\text{Ca}, \text{Mn}, \text{Fe}^{2+})\text{Al}_2(\text{BO}_3)\text{Si}_4\text{O}_{12}(\text{OH})$ ; sp. gr. 3.26–3.36;

**\*hardness** 7; violet-brown; **\*crystals** normally **\*triclinic**; occurs mainly as a product of **\*contact metamorphism** of calcium-rich **\*sedimentary rocks** (e.g. **\*contact zones** of Cornish **\*granites**).

**axiolic structure** Elongate fibres of **\*alkali** feldspar intergrown with **\*cristobalite** which have nucleated and grown outwards from both sides of a linear fracture within rhyolitic **\*glass**. The fibres, which can be seen clearly only by using a petrological microscope, grow in the solid state during devitrification of the **\*rhyolite** glass, the linear fracture acting as the axis and nucleation point for **\*crystals** defining the structure.

**axis of rotation** The line about which rotation occurs. The Earth's axis of rotation is the line between the North and South geographic Poles, which precesses (see **CHANDLER WOBBLE**; **PRECESSION**). Tectonic **\*plates** rotate about **\*Euler poles**.

**axis of symmetry** See **CRYSTALLOGRAPHIC AXES**; **CRYSTAL SYMMETRY**.

**axis vertebra** See **VERTEBRA**.

**Ayusokkanian** A Russian-Kazakhstanian **\*stage** (501–494.5 Ma ago) of the Late **\*Cambrian** **\*epoch**, preceded by the **\*Mayan** and followed by the **\*Sakian**.

**azimuth** The angle made by a line on the surface of the Earth with magnetic meridian. In **\*radar** terminology, the direction at right angles to the direction of radar propagation.

**azimuthal distribution** The spread of directional data measured from features such as cross-beds (see **CROSS-STRATIFICATION**), **\*ripples**, and oriented **\*fossils**, used to determine the direction of current flow. Azimuthal distributions may be displayed on a **\*rose diagram**, or treated statistically to give vector mean and variance data.

**azimuth resolution** In **\*radar** terminology, the width of ground area illuminated by each pulse of **\*electromagnetic** radiation. The azimuth increases with increasing distance from the radar. The azimuth resolution and the **\*slant-range** resolution govern the resolution of a radar.

**Azoic** See **CRYPTOZOIC**.

**Azores high** Semi-permanent anticyclonic region with subsiding air over the **\*Atlantic Ocean** at around 30° N latitude. Movement of the system poleward in summer has a major impact upon the climate of Europe. The aridity of the Sahara Desert and the adjacent Mediterranean region is due to the subsidence of air in this high-pressure system. See **AIR MASS**; **ANTICYCLONE**.

**azurite** **\*Secondary mineral**,  $\text{Cu}_3(\text{CO}_3)_2(\text{OH})_2$ ; sp. gr. 3.7–3.9; **\*hardness** 3.5–4.0; **\*monoclinic**; various shades of deep azure blue; light-blue **\*streak**; **\*vitreous** **\*lustre**; **\*crystals** often **\*tabular** or short **\*prisms**, and radiating aggregates; **\*cleavage** **\*prismatic** or **\*pinacoidal**; occurs in the oxidized zone of copper deposits, associated with **\*malachite** but less widespread; soluble in nitric or hydrochloric acid, with effervescence. It is a minor ore of copper.

# B

**b** See BAR.

**B** See MAGNETIC INDUCTION.

**Bacillariophyceae (diatoms)** Class of unicellular *\*algae*, usually occurring singly, but may be colonial or filamentous. Cell size ranges from 5 to 2000  $\mu\text{m}$ . The cell wall (*\*frustule*) is impregnated with silica and consists of two *\*valves*, one of which overlaps the other like the lid on a box. The frustule is commonly delicately ornamented and pierced by tiny holes (punctae) which may be covered by porous sieve membranes. There are two *\*orders*. The Centrales (centric diatoms) are circular, with *\*radial symmetry*, and are predominantly marine. The Pennales (pinnate diatoms) are elliptical, with *\*bilateral symmetry*, and dominate freshwater environments. The frustules have formed an important constituent of deep-sea deposits since the *\*Cretaceous*. The oldest known diatom is usually taken to be *Pyxidicula bollensis* from the *\*Jurassic*. See DIATOMACEOUS EARTH; DIATOMITE; DIATOM OOZE.

**back-arc basin** The zone of thickened sedimentation and extensional *\*tectonics* which lies behind an *\*island arc*. Sedimentation in back-arc basins can be very varied, ranging from *\*pelagic* through *\*turbidites* to *\*alluvial fans*, although volcanoclastics (see CLAST) are common. Various convective systems have been postulated to explain extension, which is oriented parallel to the compression in a *\*destructive* margin. Back-arc basins have been classified into *\*retro-arc* basins and *\*inter-arc* basins. Examples include the Japan Sea (between Japan and Korea) and the basins west of the *\*Marianas Trench*.

**back-arc spreading** The formation of the \*oceanic crust of the \*marginal basins, which is thought to occur by a process similar to that of normal oceanic crust, but involving a convective system developed over a \*subducting lithospheric \*plate. In most known instances the injection of new oceanic crust appears to be diffuse, rather than concentrated into \*spreading ridges.

**backing** Anticlockwise shift of the direction of the wind. The reverse change is called veering.

**backreef** Area behind, or to the landward of, a \*reef. This zone usually includes a \*lagoon between the reef and the land.

**backscatter** When a surface is illuminated by a radar beam, a portion of the energy is reflected back to the antenna ('specular reflection'). A further portion is scattered back in the same way as light is scattered from non-reflective surfaces. The proportion scattered is controlled by factors such as the roughness and \*dielectric properties of the surface, and the \*wavelength of the incident beam. Acoustic signals used to study atmospheric waves are also subject to backscatter; the amount is calculated by comparing the strength of the transmitted signal with that of the received signal after allowing for attenuation and absorption of the signal by the medium through which it propagates.

**backshore** The part of a \*beach that is above the level of normal high \*spring tides. This zone is usually dry; only when exceptionally high tides or storms occur does wave action influence this part of a beach.

**backswamp** Area of low, ill-drained ground on a \*floodplain away from the main channel. It stands slightly lower than adjacent \*alluvial fans extending from the valley sides, and is below natural \*levées that rise towards the main channel. It is a site of slow accumulation of silts and clays.

**back thrust** A \*thrust in which displacement is in an opposite direction to that of the main thrust propagation. Back thrusts are thought to form as a result of \*layer-parallel shortening in a late stage of thrust sequences.

**backwash** Seaward return of water down a \*beach. The process is affected by wave height and \*frequency, and by beach properties such as gradient and permeability. The general effect is to steepen the beach profile.

**Bacteria** One of the three taxonomic *\*domains* into which all living organisms are divided, comprising eleven groups of *\*prokaryotes*, most of which are single-celled, have a rigid cell wall, and usually reproduce by binary fission. The groups are: purple (photosynthetic); gram positive; *\*cyanobacteria*; green non-sulphur; spirochaetes; flavobacteria; green sulphur; Planctomyces; Chlamydiales; Deinococci; and Thermatogales. The only feature common to all bacteria is their prokaryotic cellular organization. Mitosis never occurs. Bacteria are almost universal in distribution and may live as saprotrophs, parasites, *\*symbionts*, pathogens, etc. They have many important roles in nature, e.g. as agents of decay and mineralization, and in the recycling of elements (such as nitrogen) in the *\*biosphere*. Bacteria are also important to humans, e.g. as causal agents of certain diseases, as agents of spoilage of food and other commodities, and as useful agents in the industrial production of commodities such as vinegar, antibiotics, and many types of dairy products. The oldest fossils known are of bacteria, from rocks in S. Africa that are apparently 3200 million years old. These must have been heterotrophic bacteria, feeding off organic molecules dissolved in the oceans of that time. The first photosynthetic bacteria, of *\*anaerobic* type, appeared a little later, about 3000 Ma ago.

**bacterial chemosynthesis** See CHEMOSYNTHESIS.

**badlands** Term originally applied to the intricately eroded plateau country of S. Dakota, Nebraska, and N. Dakota, but now widened to refer to any barren terrain that has been similarly intensively dissected. It is most common in areas of infrequent but intense rainfall and little vegetation cover.

**Badr-B (Badr-2)** A Pakistani Earth-observation *\*microsatellite*, weighing 68.5 kg, that was launched on 10 December 2001, from Baikonur, Kazakhstan, into a Sun-synchronous near-circular orbit. It marks the commencement of an indigenous development capability of low-cost satellites and their infrastructure.

**bafflestone** An *\*autochthonous* *\*carbonate* rock, with the original components organically bound during deposition. The organisms acted as baffles to permit the deposition of finer *\*matrix* material. See EMBRY AND CLOVAN CLASSIFICATION.

**baguio** The name given to a **\*tropical cyclone** that forms in the vicinity of Indonesia and the Philippines (Baguio is the name of a town in Luzon, Philippines).

**bahada** See BAJADA.

**Bairnsdalian** A **\*stage** (15–10.5 Ma ago) in the Upper **\*Tertiary** of south-eastern Australia, underlain by the **\*Balcombian**, overlain by the **\*Mitchellian**, and roughly contemporaneous with the **\*Serravallian** stage.

**Bai-u season** The principal rainy season of the south-east monsoon in southern and central Japan. The name means ‘plum rains’.

**bajada (bahada)** Extensive, gently sloping plain of unconsolidated rock debris resting against the foot of a mountain front in a semi-arid environment. Typically it is made of a number of coalescing **\*alluvial fans** laid down by **\*ephemeral** streams as their gradients lessen on leaving the mountain zone. Material is also supplied by the **\*weathering** of the mountain front. Alternatively, it may comprise the alluvial accumulation on the lower part of a **\*pediment**.

**Bajocian** A **\*stage** in the European Middle **\*Jurassic** (171.6–167.7 Ma ago, Int. Commission on Stratigraphy, 2004), preceded by the **\*Aalenian** and followed by the **\*Bathonian**. See also DOGGER.

**balanced sections** A method for reconstructing graphically the original, pre-deformational geometry of folded (see FOLD) or **\*thrust** terrains from their present appearance. Balanced sections allow the amount of regional **\*shortening** to be measured and the sequence of thrusting to be deduced. Sections are usually made parallel to the main axis of compression and originate from a reference line within undeformed strata. Three principal methods are used to balance different properties: line-length balancing; areal balancing; and **\*strain** balancing.

**Balanidae (acorn barnacles)** (order Thoracica, suborder Balanomorpha) Family of radially symmetrical, balanomorph (**\*sessile** symmetrical) barnacles, including the familiar barnacles of the genus *Balanus* exposed at low tide and characteristic of the **\*littoral zone**. *Balanus* has a fossil record extending back to the **\*Eocene**.

**Balcombian** A *stage* (15.5–15 Ma ago) in the Upper *Tertiary* of south-eastern Australia, underlain by the *Batesfordian*, overlain by the *Bairnsdalian*, and roughly contemporaneous with the upper *Burdigalian* and *Langhian* stages.

**Balfour** A *series* in the New Zealand Upper *Triassic*, underlain by the *Gore*, overlain by the *Herangi* (*Jurassic*), and comprising the Oretian, Otamitan, Warepan, and Otapirian *stages*.

**ball and pillow structure** A *sedimentary structure* occurring on the base of some *sandstones* which are interbedded with *mudstones*, and characterized by globular protrusions and isolated pillows of sandstone found in the underlying mudstone. These structures form by the differential settling of the unconsolidated sand into less dense mud below.

**ball clay** A *sedimentary*, usually *lacustrine*, *kaolinitic* *clay*, derived from the intense *weathering* of granitic and other rocks to give a unique clay sediment with both plasticity and strength. Ball clays were originally cut into cubes but by the time they had been hauled from Bovey Tracey, Devon, UK, to Stoke-on-Trent where they were to be used, the cubes had been turned into balls as they jostled together in barges and carts, hence the name.

**balloon sounding** Use of lighter-than-air balloons to establish wind conditions in the upper air. Usually the balloons are tracked by radar, and instruments may be attached to the balloons to record temperature and humidity at given pressure levels. *See also* [RADIOSONDE](#); [RAWINSONDE](#).

**Baltica (Baltoscandia)** The continental mass of north-western Europe (including most of what are now the UK, Scandinavia, European Russia, and Central Europe) that formed the south-eastern margin of the *Iapetus* Ocean. During the *Caledonian* orogenic event, this continental mass was brought into juxtaposition against N. America and Greenland by *subduction* of the Iapetus Ocean during the *Silurian* and Early *Devonian*.

**Baltoscandia** *See* [BALTICA](#).

**band** In *remote* sensing, the range of wavelengths which are examined.

**banded iron formation (BIF)** Finely banded, siliceous, \*hematite deposits found in \*Precambrian rocks, forming \*stratiform units often several hundred metres thick and persistent over 150 km or more. They probably formed by chemical–organic processes during sedimentation in stable, shallow basins with little detritus, and so are \*syngenetic deposits. In their enriched form (40–60% iron) BIFs are mined \*open-cast. They include the world’s most important sources of iron ore, e.g. at Hammersley, Western Australia; Lake Superior, USA; Labrador, Canada; Ukraine; and Brazil. *Compare* GRANULAR IRON FORMATION.

**band filter** A \*filter that either passes over (band-pass filter) a discrete range of \*frequencies with minimal \*attenuation, or rejects (band-reject filter) those frequencies by substantial attenuation. Band-reject filters are the inverse of band-pass filters.

**band-pass filter** *See* BAND FILTER.

**band-reject filter** *See* BAND FILTER.

**band silicate** *See* INOSILICATE.

**bank calving** The toppling of slabs of earth from a river bank that occurs when the stream undermines the bank.

**bankfull flow** Maximum amount of \*discharge (usually measured in m<sup>3</sup>/s) that a stream \*channel can carry without overflowing. Its frequency of occurrence varies between streams, from a few times each year to once every few years. The water height at bankfull discharge is referred to as the ‘bankfull stage’.

**bankfull stage** *See* BANKFULL FLOW.

**banner cloud** Motionless, flag-like cloud, commonly of lenticular (lens-like) shape, forming to the lee (eddy zone) side of a hill or mountain peak. The cloud extends downwind in a strong current of humid air, resembling a banner in the wind. Many distinctive mountain peaks (e.g. the Matterhorn and Table Mountain) are associated with a characteristic banner cloud. *See also* LEE WAVE.


**bar 1. (b)** Unit of pressure approximately equal to one atmosphere (14 lb/in<sup>2</sup>), and precisely equal to 10<sup>5</sup> Pa (10<sup>5</sup> N/m<sup>2</sup>) in SI units. The pressure of

the atmosphere at sea level on average is very approximately one bar, or about 1013 millibars (the bar commonly being divided into one thousand millibars, mb). 2. Geomorphologic term: (a) Low ridge of sand or shingle laid down by marine aggradation in shallow water adjacent to a coastline. There are several varieties: a bay bar joins the two flanks of a bay and may enclose a lagoon; an offshore or barrier bar runs parallel to a coastline and up to 40 km distant. (b) Rocky obstruction across a glaciated valley. See [GLACIAL](#) stairway. (c) Lobate river [\\*bedform](#), typically constructed of gravel, often regularly spaced, and forming a riffle or shallow section. (d) Point bar: a low crescentic shoal on the convex side (inside) of a river bend, consisting of material that has been eroded from an outside bend, either opposite or upstream. Point-bar deposits consist of relatively coarse materials, often showing an upstream [\\*dip](#).

[Baragwanathia longifolia](#) One of the earliest known vascular plants (see [TRACHEOPHYTA](#)). During the [\\*Devonian](#) and [\\*Carboniferous](#), the lycopods (club mosses) reached the peak in their diversification. *B. longifolia* was an early representative of the family Drepanophycaceae (class [\\*Lycopsida](#)) and is known from the Lower Devonian.

[barat](#) Local term for a fierce north-westerly wind common from December to February on the northern coast of Sulawesi, Indonesia.

[barchan](#) (*adj.* [barchanoid](#)) Crescent-shaped mobile [\\*dune](#) in a sand desert in which the wind blows predominantly from one direction. The dune moves by the erosion of sand from the windward slope, and its accumulation on the steeper lee or slip slope, which stands at about 32°. Average velocities of dune movement are 10–20 m/yr.



**Barchan**

**barchanoid** See AKLÉ; BARCHAN.

**barite (baryte)** A *mineral*,  $\text{BaSO}_4$ , which may form a *solid solution* series with *celestite* ( $\text{SrSO}_4$ ); sp. gr. 4.3–4.6; *hardness* 3.0–3.5; *orthorhombic*; colourless to white, often tinged yellow, brown, blue, green, and red; white *streak*; *vitreous lustre*; *crystals* commonly *tabular*, *prismatic*, but can be fibrous, lamellar, and often granular; *cleavage* perfect {001}, present {210}, {010}. Occurs as a vein filling, as a *gangue* mineral with ores of lead, copper, zinc, silver, iron, and nickel, associated with *calcite*, *quartz*, *fluorite*, *dolomite*, and *siderite*, and as a low-temperature mineral which also occurs as a replacement for *limestone*, and as a *cement* in *sandstone*. Insoluble in acid. It is used as a weighting agent in drilling muds, in the chemical industry, in the manufacture of rubber, paper, and high-quality paints, and as an X-ray absorbent.

**barkevikite** An alkali (sodium and potassium), iron-rich *amphibole*, whose colour is very distinctive in *thin* section.

**barnacles** See BALANIDAE; CIRRIPIEDIA.

**baroclinic** 1. Applied to an atmospheric condition in which isobaric and constant-density surfaces are not parallel, e.g. in a frontal zone. 2. Applied to a state in the ocean in which the surfaces of constant pressure intersect surfaces of constant density. In this situation, the water density gradient

depends on water properties (temperature and *\*salinity*) as well as pressure (depth). This can be contrasted with the *\*barotropic* situation.

**baroduric** Capable of withstanding high pressures.

**barograph** *\*Barometer* that gives a continuous recording of air pressure. It is based on an *\*aneroid* instrument with levers attached to the vacuum chambers, and records a trace on a chart mounted around a clock drum.

**barometer** Instrument for the measurement of *\*atmospheric* pressure. The usual type is a mercury barometer, in which the atmosphere's pressure on a small reservoir of mercury supports a column of mercury in a vacuum tube the open end of which is below the surface of the mercury in the reservoir. The column is on average about 76 cm (30 inches) high. Readings must be corrected to compensate for pressure variation due to gravitational anomalies and for thermal expansion or contraction of the mercury; therefore correction to a standard temperature is necessary. *See also* ANEROID BAROMETER; FORTIN BAROMETER; KEW BAROMETER.

**barothermograph** Device for the continuous measurement of both pressure and temperature, recorded on a revolving chart.

**barotropic** Applied to a state in a water or *\*air mass* in which the surfaces of constant pressure are parallel to the surfaces of constant density. In this situation the density gradient depends on depth only, as in an isothermal freshwater lake. *Compare* BAROCLINIC.

**barred basin** A partially restricted *\*sedimentary basin*, where free movement of waters is impeded by the presence of a rock sill or sediment barrier. This restriction often results in *\*anoxic* or oxygen-poor waters, or, in arid areas, in *\*evaporite* deposition.

**barrel** A unit of volume used to measure dry goods or liquids. An oil barrel (bbl) is equal to 42 US gallons = 34.9723 imp. gallons = 158.983 litres = 0.159 m<sup>3</sup>.

**Barrell, Joseph** (1869–1919) Barrell was a teacher of mining engineering at Yale University, who developed (in 1917) a chronology for the *\*Phanerozoic*, based on rates of sedimentation and *\*radiometric* dating. He proposed that the anomalously low age of the Earth based on sedimentation rates could be explained by long gaps in the sedimentary record, called

\*diastems, and that sediments formed only when \*subsidence occurred. Barrell was the first person to use the terms \*lithosphere and \*asthenosphere, based on his observations of \*isostasy.

**barrel trend** See BOW TREND.

**Barremian** 1. An early \*Cretaceous \*age, preceded by the Neocomian \*epoch (but see NEOCOMIAN), followed by the \*Aptian age, and dated at 130–125 Ma ago (Int. Commission on Stratigraphy, 2004). 2. The name of the corresponding European \*stage, for which the \*stratotype is at Angles, France.

**barren interzone** An \*interval zone that is devoid of \*fossils.

**barren intrazone** A measurable breadth of \*strata, occurring within a \*biostratigraphic unit but itself containing no \*fossils.

**barrier** General term for a depositional feature standing on the seaward side of a coastline. See BARRIER BAR; BARRIER BEACH; BARRIER ISLAND; BARRIER REEF; BAYHEAD BARRIER; BAYMOUTH BARRIER.

**barrier bar** A major \*longshore bar of gravel or sand whose surface is below mean still water level. Normally it is formed off a depositional coast of low gradient and with ample unconsolidated sediment. See BARRIER.

**barrier beach** A relatively small, shingle feature that protects a steep coast. See BARRIER.

**barrier island** An elongated ridge that may extend from a few hundred metres to 100 km along a coast forming a segmented \*barrier-bar complex, and found between two tidal inlets. Barrier-island systems have a \*lagoonal area on their landward side, and often have wind-blown dunes and vegetation on the exposed (seaward) side of the \*barrier. There are three main hypotheses to explain the origin of barrier islands: (a) the building up of submarine bars; (b) spit progradation parallel to the coast and segmentation by inlets; and (c) submergence of subaerial coastal beach ridges by a rise in sea level. Barrier islands are most common in areas of low tidal range.

**barrier reef** \*Reef trending parallel to, but separated by a \*lagoon from, a shore. The reef-building organisms build up the structure to approximately

the low tide level. One of the finest examples is the Great Barrier Reef which lies off the north-eastern coast of Australia: it extends for about 1900 km and is 30–160 km in width.

**Barrovian-type metamorphism** A sequence of *\*regional metamorphic \*mineral* reactions recorded by the successive mineral assemblages seen in *\*metapelites* (metamorphosed or sandy shales) from the Barrovian terrain around Glen Esk in north-eastern Scotland and characteristic of medium regional metamorphic gradients of temperature and pressure. George Barrow (1853–1932) in 1912 was the first to recognize this sequence of metamorphic mineral assemblages. In *\*pelites*, Barrovian-type metamorphism is marked by the development of a sequence of *\*index* minerals, starting with *\*chlorite* in the lowest-grade rocks, and passing up through *\*biotite*, *\*garnet*, and *\*kyanite*, to *\*sillimanite* in the highest-grade rocks (see *BARROW'S ZONES*).

**Barrow's zones** The original subdivision by George Barrow (1853–1932) of the sequence of mineral changes seen in rocks of pelitic composition in the Glen Esk region of north-eastern Scotland. Each zone is bounded by two *\*isograds*, each of which marks the appearance, in the direction of increasing *\*metamorphic grade*, of a new *\*index mineral*, and is named after the index mineral seen on the lower-grade boundary. (For instance, the *\*kyanite* zone has the kyanite isograd as its low-grade boundary and the *\*sillimanite* isograd as its high-grade boundary.) Within each zone, no additional minerals appear in the *\*pelites*, the constant mineral assemblage representing equilibrium over a range of metamorphic conditions. See *BARROVIAN-TYPE METAMORPHISM*. Compare *BUCHAN METAMORPHIC ZONES*.

**Bartonian** 1. An *\*age* (40.4–37.2 Ma ago) in the Middle *\*Eocene*, preceded by the *\*Lutetian*, followed by the *\*Priabonian* (Int. Commission on Stratigraphy, 2004). 2. The name of the corresponding European *\*stage*, which is roughly contemporaneous with most of the upper *\*Narizian* (N. America), upper *\*Bortonian* and *\*Kaiatan* (New Zealand), and part of the *\*Aldingan* (Australia). It was originally considered to be a lateral equivalent of the Priabonian stage (Upper Eocene). (The name should not be confused with Bortonian.)

**barycentric reference frame** In *\*geodesy*, a reference frame centred on the centre of mass of the solar system (i.e. the Sun), that is used to express positional coordinates derived from astronomical or satellite measurements.

**baryte** See BARITE.

**basal conglomerate** 1. A *\*conglomerate* deposited at the base of a sedimentary sequence, e.g. at the base of a *\*channel-fill* deposit. 2. A conglomerate deposited above an *\*unconformity* surface.

**basal sliding** The process by which a temperate *\*glacier* moves over its bed. It involves three mechanisms: relatively rapid creep in the basal layers; pressure melting, whereby *\*ice* under pressure melts on the up-glacier side of a small obstacle and the released water freezes on the down-glacier side; and slippage over a layer of water at the bed.

**basalt** A dark-coloured, fine-grained, *\*extrusive*, *\*igneous* rock composed of *\*plagioclase* feldspar, *\*pyroxene*, and *\*magnetite*, with or without *\*olivine*, and containing not more than 53 wt. % SiO<sub>2</sub>. Many basalts contain *\*phenocrysts* of olivine, plagioclase feldspar and pyroxene. Basalts are divided into two main types, *\*alkali basalts* and *\*tholeiites*, with the tholeiites being subdivided into olivine tholeiites, tholeiites, and quartz tholeiites. Petrographically (see PETROGRAPHY), alkali basalts have as their *\*groundmass* pyroxene titanite (an *\*augite* rich in titanium), whereas tholeiites have pigeonite (a calcium-poor pyroxene). Also, for similar concentrations of SiO<sub>2</sub>, alkali basalts have a higher content of Na<sub>2</sub>O and K<sub>2</sub>O than tholeiites. Basalt flows cover about 70% of the Earth's surface and huge areas of the *\*terrestrial* planets, and are therefore arguably the most important of all crustal rocks. They are formed by *\*partial* melting of *\*mantle* *\*peridotite*. Alkali basalts are typically found on oceanic islands and on the *\*continental crust* in regions of crustal upwarping and rifting. Tholeiites are typically found on the ocean floor and on the stable continental crust where they form large basalt plateaux such as the *\*Deccan Traps* of India.

**basal thrust** See SOLE THRUST.

**basaltic meteorites** *\*Achondrite* *\*meteorites* that have been derived from parent bodies sufficiently evolved to have acquired a basaltic crust (in

contrast, for example, to [\\*chondrites](#), which have derived from relatively primitive bodies). [\\*Eucrites](#) and howardites (similar to eucrites, but more [\\*pyroxene](#)-rich) probably formed through magmatic processes on the same parent body (possibly the [\\*Moon](#)) around 4.5 billion years ago. The [\\*shergottite](#)–nakhilite–chassignite class of basaltic meteorite has a distinctly different oxygen-isotope composition (see [OXYGEN-ISOTOPE RATIO](#)), [\\*volatile](#) content, and age (around 1.3 billion years), and originated from a separate parent body, probably the planet [\\*Mars](#).

**basanite** An [\\*extrusive](#), [\\*mafic](#), [\\*igneous](#) rock consisting of a [\\*feldspathoid](#) [\\*mineral](#) ([\\*nepheline](#), [\\*analcite](#), or [\\*leucite](#)), [\\*olivine](#), [\\*plagioclase](#) feldspar, [\\*pyroxene](#), and minor [\\*accessory minerals](#). Essentially, the mineralogy is that of a [\\*basalt](#), with the addition of a feldspathoid mineral and olivine. Varieties of basanites are defined by the type of feldspathoid mineral present to give nepheline-basanite, analcite-basanite, and leucite-basanite. These rocks are found in close association with [\\*alkali](#) basalts. *See also* [TRACHYBASALT](#).

**base** According to the Brønsted–Lowry theory, a substance that in solution can bind and remove hydrogen [\\*ions](#) or protons. The Lewis theory states that it is a substance that acts as an electron-pair donor. A base reacts with an [\\*acid](#) to give a salt and water (a process called neutralization), and has a [\\*pH](#) greater than 7.0. The theory was proposed in 1923 by the Danish physical chemist Johannes Nicolaus Brønsted (1879–1947) and the British chemist Thomas Martin Lowry (1874–1936), and independently by the American theoretical chemist Gilbert Newton Lewis (1875–1946).

**basecourse** **1.** The lowest course of masonry. **2.** *See* [PAVEMENT](#).

**baseflow (dry-weather flow)** In a stream or river, the flow of water derived from the seepage of [\\*groundwater](#), and/or throughflow into the surface watercourse. At times of peak river flow, baseflow forms only a small proportion of the total flow, but in periods of [\\*drought](#) it may represent nearly 100%, often allowing a stream or river to flow even when no rain has fallen for some time. *See also* [INTERFLOW](#); [SUBSURFACE FLOW](#).

**baselap** The discordant relationship marking the lower boundary of a [\\*depositional sequence](#), where upper beds lap out over the underlying surface. There are two types of baselap: [\\*onlap](#), where the overlying beds

thin out updip; and **\*downlap**, where younger, initially inclined, overlying beds thin out downdip. See **DIP**.

**base level** A theoretical plane surface underlying a land mass, denoting the depth below which **\*erosion** would be unable to occur. Sea level provides a base level on a regional scale. Local base levels may be provided by the base of a hillslope, lakes, or by the junction between a tributary and the main river.

**basement** 1. Highly folded, **\*metamorphic** or **\*plutonic** rocks, often unconformably (see **UNCONFORMITY**) overlain by relatively undeformed **\*sedimentary** beds (or cover). In this sense, basement is often, though not necessarily, **\*Precambrian**. 2. The **\*crust** below rocks of interest. In this sense, to a petroleum geologist 'basement' means non-prospective rocks which lie below prospective strata.

**base saturation** The extent to which the exchange sites of the soil's **\*adsorption complex** are 'saturated' (or occupied) by exchangeable basic **\*cations**, or by cations other than hydrogen and aluminium, expressed as a percentage of the total **\*cation-exchange capacity**.

**base station** A station to which reference can be made, to normalize measurements made at out-stations. Geophysical parameters measured at base stations are presumed to be known very accurately and preferably absolutely. For example, in a **\*gravity survey**, the base station is used to determine instrumental **\*drift** and to provide an absolute value to which **\*gravity anomalies** relate. In **\*magnetic** surveying, a continuous-reading **\*magnetometer** may be used at a base station to monitor the **\*diurnal** fluctuations in the Earth's **\*geomagnetic** field and thus aid the interpretation of magnetic survey data.

**base surge** A turbulent, dilute flow of **\*ash** and either water or steam, which expands radially as a collar-like cloud from the base of a vertically venting eruption column generated by the explosive interaction of **\*magma** and water. The surges are commonly cold and wet, consisting of ash mixed with water at temperatures below 100 °C. With a high magma:water mass ratio the surges can become dry and hot, consisting of ash mixed with steam. Base surges can be very hazardous, as was the one that occurred

during the eruption of Taal, Philippines, 1965. Radially expanding basal clouds observed in nuclear explosions are a type of base surge. See [SURGE](#).

**Bashkirian (Morrowan) 1.** The earliest [\\*epoch](#) in the [\\*Pennsylvanian](#), comprising the Kinderscoutian, Marsdenian, and Yeadonian ages (these are also [\\*stage](#) names in British stratigraphy), and the Cheremshanskian and Melekesskian ages (stage names in Russian stratigraphy). The Bashkirian age is preceded by the [\\*Serpukhovian](#), followed by the [\\*Moscovian](#), and has its initial boundary (the [\\*Mississippian](#)–Pennsylvanian boundary) dated at 318.1 Ma ago and its end at 311.7 Ma ago (Int. Commission on Stratigraphy, 2004). **2.** The name of the corresponding eastern European [\\*series](#), which is roughly contemporaneous with the [\\*Namurian](#) B and C plus the [\\*Westphalian](#) A (western Europe), and the Morrowan series (N. America).

**basic 1.** Fundamental. **2.** Adjective derived from [\\*base](#). **3.** See [BASIC ROCK](#).

**basic rock** Rock with a relatively high concentration of iron, magnesium, and calcium, and with 45–53% of [\\*silica](#) by weight. Examples include [\\*gabbro](#), which is a coarse-grained basic [\\*intrusive](#) rock, and [\\*basalt](#), which is a fine-grained basic volcanic ([\\*extrusive](#)) rock. Compare [ACID ROCK](#); [INTERMEDIATE ROCK](#). See also [ALKALINE ROCK](#).

**basic soil** A soil with a [\\*pH](#) greater than 7.0. See [BASE](#). Compare [ALKALINE SOIL](#).

**Basilosaurus** (order Cetacea, suborder [\\*Archaeoceti](#)) One of the best-known archaeocetes; it grew to approximately 20 m in length and lived during the Upper [\\*Eocene](#). *Basilosaurus* retained small hind limbs, useless for locomotion, but perhaps usable as claspers during copulation.

**basin 1.** Depression, usually of considerable size, which may be erosional or structural in origin. The converse of a [\\*dome](#). **2.** See [PERICLINE](#).

**basin-and-range crustal type** A type of [\\*crust](#) named after the type found in the [\\*basin-and-range](#) province of the western United States. Extension has made the crust thin (average thickness 30 km) and highly unstable, resulting in a series of basins separated by upland areas, most of which are heavily eroded. There are strong negative [\\*gravity anomalies](#) and marked volcanic and seismic activity.

**basin-and-range province** 1. Specifically, the structural subdivision of the western USA between the Colorado Plateau to the east and the Sierra Nevada to the west. It is dominated by a series of block-faulted ranges and troughs which dissect it, rising above *\*bajadas* and *\*alluvial* plains which mask down-faulted units. Frequently the ranges are tilted blocks approximately 30–40 km across. Movement occurred initially in *\*Miocene* times, and then during the late *\*Pliocene* to *\*Recent*. 2. Generally, applied to landscapes having a series of tilted *\*fault blocks* that form long, asymmetric ranges separated by broad *\*basins*.

**basin-and-swell sedimentation** A form of sedimentation in a region of differential *\*subsidence*, where thin, *\*condensed*, sedimentary sequences are deposited on slowly subsiding highs or ‘swells’, and thicker, usually muddier sediments accumulate in more rapidly subsiding basins between the swells.

**basin modelling** The computer simulation of the geological evolution of a sedimentary basin, quantifying certain processes within it, with the aim of predicting the distribution and movement of petroleum within the basin and determining the temperatures and pressures.

**Baslean** See *KAZANIAN*; *UFIMIAN*.

**bastnäsite** A *\*mineral*,  $\text{Ce}(\text{CO}_3)\text{F}$ ; average density 4.97; *\*hardness* 4–5; *\*hexagonal* to ditrigonal dipyramidal; *\*cleavage* {1011} imperfect, {0001} indistinct; yellow or reddish brown, translucent to transparent; white *\*streak*; *\*vitreous* to *\*greasy lustre*; crystals *\*prismatic* but it can be massive; named for the Bastnäs mine, Västernmanland, Sweden, it also occurs widely, but never in large concentrations, in *\*karst bauxite* deposits and *\*carbonatites* elsewhere in Europe, and in Mongolia, Malawi, California, and Canada.

**Batesfordian** A *\*stage* (16.5–15.5 Ma ago) in the *\*Miocene* of south-eastern Australia, underlain by the *\*Longfordian*, overlain by the *\*Balcombian*, and roughly contemporaneous with the mid *\*Burdigalian* stage.

**batholith** Large (more than 100 km<sup>2</sup>) *\*igneous* intrusion, which may comprise several *\*plutons* amalgamated at depth. Most batholiths are granitic in composition and their genesis is linked with *\*plate tectonics*.

Generally, batholiths cut across *\*country rocks* and therefore are discordant in nature.

**Bathonian** Middle *\*Jurassic \*stage* (167.7–164.7 Ma ago), preceded by the *\*Bajocian*, followed by the *\*Callovian*, and commonly represented by carbonate sediments in many areas of Europe but not in the North Sea or Scandinavia. It is known to contain an abundant fauna of invertebrates. *See also DOGGER.*

**bathy-** From the Greek *bathus* meaning ‘deep’, a prefix meaning ‘deep’ as applied to the oceans.

**bathyal zone** Oceanic zone at depths of 200–2000 m, lying to the seaward of the shallower *\*neritic zone*, and landward of the deeper *\*abyssal zone*. The upper limit of the bathyal zone is marked by the edge of the *\*continental shelf*. In marine ecology, it is the region of the *\*continental slope* and rise. It may be geologically active, and include *\*trenches* and *\*submarine canyons*, with underwater erosion producing avalanches.

**bathymetry** The measurement of the depth of the ocean floor from the water surface; the oceanic equivalent of topography.

**bathythermograph** An instrument, invented in 1937, that is used to measure changes of temperature with depth in the ocean, down to about 450 m. A bathythermograph is shaped like a bomb. It carries a *\*thermistor* in its lead-weighted nose and also a pressure sensor, and it trails a wire connected to equipment on the deck of the ship. The thermistor provides a continuous record of changing water temperature and the pressure sensor records the depth.

**bats** *See* CHIROPTERA.

**Batyrbayan** A Russian-Kazakhstanian *\*stage* (491.5–488.3 Ma ago) of the Late *\*Cambrian \*epoch*, preceded by the *\*Aksayan* and followed by the *\*Tremadocian*.

**Bauplan** (*pl.* **Baupläne**) The generalized body plan of an archetypal member of a major taxon.

**bauxite** A mixture of three hydrates of alumina, mainly *\*gibbsite*, and also *\*diaspore* and *\*boehmite*, and containing impurities of iron, phosphorus, and titanium; colour is variable from dirty white through grey, yellow,

brown, and red; sp. gr. 2.0–2.55; \*hardness 1–3; it can be compact, \*earthy, concretionary (see CONCRETION), \*pisolitic, or \*oolitic. Bauxite results from the tropical weathering of aluminium silicate rocks under good surface drainage to yield \*clay minerals which are subsequently desilicated. \*Minerals associated with the alumina hydrates in bauxites and \*laterites (ferruginous bauxites) include \*goethite and lepidocrocite, \*hematite, and the clay minerals \*kaolinite and \*halloysite. Bauxite is the main ore of aluminium and to be commercially exploited should contain more than 25–30% aluminium oxide. The main constraint is the amount of available alumina which can be extracted by the Bayer or similar process. It is named after Les Baux de Provence, in southern France; the major producers are Australia and Brazil.


**Baventian** The Lower \*Pleistocene, cold-stage, marine silts and marine clays, that form part of the tripartite division of deposits revealed in a \*borehole at Ludham, in eastern England. *See also* ANTIAN; LUDHAMIAN; PASTONIAN; THURNIAN.

**bay bar** *See* BAR.

**Bayesian** In statistics, applied to the re-evaluation of probabilities based on empirical observation.

**bayhead barrier** A \*barrier beach protecting the head of a bay, but separated from it by a \*lagoon. *See* BARRIER.

**bayhead beach** Sand or shingle \*beach in the low-energy environment at the head of a bay. It is typical of irregular coastlines in which bays and promontories alternate.



**Bayhead beach**

**baymouth barrier** A *barrier* that partially encloses a bay at its entrance.

**bayou** A slow-moving river, marshy lake, or a tidal creek containing brackish water.

**Bazin's average velocity equation** In 1897, when considering flow in open *channels*, H. E. Bazin (1829–1917) proposed that the Chezy discharge coefficient ( $C$ ) could be related to the *hydraulic radius* ( $r$ ) and a channel roughness coefficient ( $k_1$ ) by the formula:  $C = 157.6/[1 + (k_1/r^{1/2})]$ . Other more complex formulae have also been proposed. See **CHEZY**'s formula.

**bbi** See **BARREL**.

**beach** An accumulation of sand and gravel found at the landward margin of the sea or a lake. Along sea coasts, beaches comprise the *backshore* zone (above mean high water) and the *foreshore* zone (between mean high and low tide levels). The angle of slope and the *sedimentary structures* of a beach are related to the grain size of the beach materials, and to the nature of wave activity and other sedimentary processes active in the area.

**beach cusp** One of a series of regularly spaced crescent-shaped structures forming local relief along a *beach*. The horns or 'headlands' of the cusp are composed of coarse *sand* or gravel, and point seaward down the beach. The intervening troughs or 'bays' are made up of finer sand. Beach cusps

are usually several centimetres high, although larger examples have been described. The size and spacing of cusps appears to be related to the nature of the waves breaking on the beach.

**beach drift** The zig-zag progression of *\*sand* and other debris along a *\*beach*. Particles are driven obliquely up a beach by the *\*swash* and are then returned down the steepest gradient of the beach by the *\*backwash*. The combination of these two movements gives the zig-zag progression. *See also* LONGSHORE DRIFT.

**beach rock** *\*Cemented* beach *\*sand* deposit that develops within the intertidal zone by the precipitation of needle-like crystals of *\*aragonite* in the *\*pore space* between the grains. The cementation process is relatively rapid, taking as little as 10 years for a lithified rock to develop. The precipitation of the cement is favoured by a warm climate, and may be aided by algal or bacterial action.

**beaded lightning** *See* PEARL-NECKLACE LIGHTNING.

**beak** *See* SHELL BEAK.

**BEAMS** *See* BRIDGMANITE-ENRICHED MANTLE STRUCTURE.

**beardworms** *See* POGONOPHORA.

**bearing capacity** Maximum *\*load* per unit area a surface can support in safety without shear *\*failure*.

**Beaufort scale** Scale of values, from 0 to 12, for describing wind strength, as defined by Commander (later Rear-Admiral) **Francis Beaufort** (1774–1857) in 1806; it was adopted throughout the Royal Navy in 1838 and by the International Meteorological Committee in 1874. Originally, the scale determined the wind force from the amount of sail a warship needed to carry to attain a specified speed, with no listing of wind speeds. The modern version, finalized in 1939 by the International Meteorological Committee, included wind speeds. Wind force is now recognized by its common effects on objects in the landscape (dust, flags, trees, etc.) and on people in the open, or on the state of the sea surface. *See also* SAFFIR–SIMPSON SCALE; FUJITA TORNADO INTENSITY SCALE; and [appendix C](#).

**Beche, Henry Thomas de la** (1796–1855) Founder of the British Geological Survey, the Museum of Practical Geology, the Mining Records Office, and the School of Mines, all of which are in Britain. De la Beche was a careful observer and skilled cartographer and artist, who emphasized the importance of stratigraphy and pioneered the reconstruction of ancient environments.

**Becke, Friedrich Johann Karl** (1855–1931) An Austrian mineralogist and petrologist from the University of Prague, Becke developed a method for determining the relative *\*refraction* of light in microscopy, later named for him. He also did important work on metamorphic *\*recrystallization*, developing a descriptive terminology and a classification of metamorphic *\*facies*. See [BECKE LINE TEST](#).

**Becke line test** In transmitted-light microscopy, a comparative test used to determine the approximate *\*refractive index* of a *\*mineral*. In *\*plane-polarized light*, when the substage iris diaphragm is partly closed to accentuate grain boundaries, a thin line of light, the ‘Becke white line’, appears. If the microscope tube is then racked up, or the microscope *\*stage* racked down, the light line will move into the medium with the higher refractive index. The medium could be either an adjacent mineral of known refractive index or, more commonly, the mounting medium, usually a cold-setting resin or Canada balsam (refractive index 1.54).

**bed** See [BEDDING](#); [STRATUM](#).

**bedding** Layering of sheet-like units, called beds or strata (the terms are not synonymous, see [STRATUM](#)); a bed or stratum being the smallest distinguishable division within the classification of stratified sedimentary rocks. Cross-bedding is a type of stratification in which some sediment layers have an inclined attitude in relation to those immediately above or below. See [CROSS-STRATIFICATION](#). Compare [LAMINATION](#).

**bedding foliation** A *\*foliation* parallel to the *\*bedding* which is commonly found in limbs of tight and *\*isoclinal folds*, where the *\*axial plane* is parallel or subparallel to the limbs.

**bedding plane** **1.** Well-defined, planar surface that separates one bed from another in *\*sedimentary rock*. Each plane marks a break in deposition. **2.** See [VOIDS](#).

**bedform** The shape of the surface of a bed of granular *\*sediment*, produced by the *\*flow* of air or water over the sediment. The nature of the bedform depends upon the flow strength and depth, and upon sediment *\*grain size*. For fine to medium *\*sand*, the typical sequence of bedforms produced under conditions of constant depth and increasing strength of unidirectional flow is: no movement; *\*ripples*; *\*sand waves*; *\*dunes*; and an upper-flow-regime *\*plane bed*. In coarse sand a lower-flow-regime plane bed develops first, then ripples, followed by sand waves, then dunes, and an upper-flow-regime plane bed. At higher-strength flows, the upper flow regime plane bed is replaced by *\*antidunes*.

**bed load (traction carpet, traction load)** The coarser fraction of a river's total *\*sediment* load, which is carried along the bed by sliding, rolling, and *\*saltation*. It constitutes 5–10% of the total load.

**bed roughness** The surface relief developed at the base of a flowing fluid, comprising *\*bedforms* (form roughness), and particles projecting from the sediment carpet (grain roughness). 'Bed roughness' is the quantifiable factor which expresses the frictional effect that the bed exerts on the flow. Surfaces whose bed-roughness elements do not project through the viscous sub-layer at the base of the flow are said to be smooth. Surfaces whose bed-roughness elements project through the sub-layer are said to be rough.

**Beekmantownian** A *\*stage* of the *\*Ordovician* in the *\*Canadian \*series* of N. America.

**Beestonian** 1. A cold period during the Middle *\*Pleistocene*. 2. Arctic, freshwater bed deposits of *\*sand* and silts represented at Beeston, Norfolk, England. Some gravels in Norfolk, Suffolk, and Essex, England, have also been ascribed to this period.


**belemnites** See BELEMNITIDA.

**Belemnitida (belemnites)** (class *\*Cephalopoda*, subclass Coleoidea) One order of extinct cephalopods in which the shell is internal and composed of a *\*phragmocone*, *\*rostrum*, and pro-ostracum (see SKELETAL MATERIAL). Belemnites appear in the *\*Jurassic*, continue through the *\*Cretaceous*, and a few persist into the *\*Eocene*. The other order, Aulacocerida (*\*Carboniferous* to Jurassic) may have retained a body chamber, but in the Belemnitida this is reduced to the pro-ostracum. The most posterior portion

of the shell is known as the ‘guard’ (*\*rostrum*). This is a bullet-shaped cylinder made up of radiating needles of *\*calcite* with a conical cavity (alveolus) in its anterior end into which fits the phragmocone, a conical, aragonitic (see *ARAGONITE*), septate (see *SEPTUM*) structure, cut by a tiny *\*siphuncle*, that is *\*homologous* to the external shell of other cephalopods. The pro-ostracum is a tongue-like, anterior projection from the phragmocone and perhaps protected the anterior part of the body.

**Belinda (Uranus XIV)** One of the lesser satellites of *\*Uranus*, with a diameter of 34 km. It was discovered in 1986.

**bellerophontiform** Applied to the shape of gastropod (*\*Gastropoda*) shells which resemble those of the genus *Bellerophon*. The coiling is isotrochic (i.e. two faces of the shell are symmetrical in respect of a median plane perpendicular to the axis).



### **Bellerophontiform**

**bell pit** In mining, an obsolete method for winning *\*ore* or *\*coal* from shallow deposits. Material was extracted and dragged to a central shaft, leaving a bell-shaped excavation. The term comes from ironstone working in Derbyshire, England.

**bell trend** See *DIRTYING-UP TREND*.

**bench** 1. Horizontal step along which material is worked in an open pit. 2. Any narrow, flat surface in solid rock.

**Bendigonian** A *\*stage* (475–473 Ma ago) of the Lower *\*Ordovician* of Australia, underlain by the *\*Lancefieldian* and overlain by the

\*Chewtonian.

**beneficiate** To increase the grade of ore after grinding by a concentration process, such as froth flotation or gravity, or by \*magnetic, electrostatic, or other methods. The term also implies the elimination of waste material. See FLOTATION SEPARATION.

**Benguela current** Oceanic water current that flows northward along the west coast of southern Africa between latitude 15° S and 35° S. It is distinguished by an area of cold upwelling water, and is a relatively weak current, flowing at less than 0.25 m/s.

**Benioff, Hugo** (1899–1968) An American seismologist, and professor at the California Institute of Technology from 1950 to 1964, Benioff designed seismographs and other instruments used to study \*earthquakes. In 1954 he published a cross-section showing \*seismicity beneath the Kamchatka Peninsula, which demonstrated that the depths of earthquakes increased with distance away from the trench. This earthquake zone has been named after him. See BENIOFF ZONE.

**Benioff zone (Wadati–Benioff zone)** One of the zones of deep \*earthquake \*hypocentres whose existence was first demonstrated in 1927 by the Japanese seismologist Kiyoo Wadati. The zones were mapped in the 1940s and 50s by Hugo \*Benioff. They dip from near-surface to a maximum depth of approximately 700 km and are associated with oceanic \*trenches, \*island arcs, volcanic chains, and young \*fold mountains, and are thought to indicate active \*subduction. See SUBDUCTION ZONE.

**benmoreite** An \*extrusive \*igneous rock consisting of \*anorthoclase, sodic \*plagioclase, ferroaugite (an iron-rich \*augite), and iron-rich \*olivine, and found as a member of the \*alkali basalt \*magma series. With increasing differentiation (see MAGMATIC DIFFERENTIATION) shown by increasing SiO<sub>2</sub> content, the series is alkali basalt–hawaiite–mugearite–benmoreite–\*trachyte. The type locality after which the rock is named is Ben More on the Isle of Mull, Scotland.

**Bennettiales** Extinct \*gymnosperm order ranging from the \*Triassic into the \*Cretaceous. They resembled \*cycads, and possessed reproductive structures which must have looked more like flowers than cones.

**Benson's flood-peak formula** See FLOOD-PEAK FORMULAE.

**benthic storm** See ABYSSAL STORM.

**benthos** (*adj.* **benthic**) In freshwater and marine **\*ecosystems**, the collection of organisms attached to or resting on the bottom sediments (i.e. **\*epifaunal**), and those which bore or burrow into the sediments (i.e. **\*infaunal**).

**bentonite** **\*Montmorillonite**-rich **\*clay** formed by the breakdown and alteration of volcanic **\*ash** and volcanic **\*tuffs**.

**BepiColombo** A collaborative mission between the European Space Agency (ESA) and the Japan Aerospace Exploration Agency (JAXA) Institute of Space and Astronautical Science (ISAS) to explore **\*Mercury**, named for Giuseppe ('Bepi') Colombo, the Italian astrophysicist who helped plan NASA's Messenger expedition to Mercury, which ended in 2015. It was launched successfully on 20 October 2018, from Kourou, French Guiana. BepiColombo comprises two spacecraft that will separate on arrival; the ESA orbiter will study the planet itself, and the ISAS probe will study the planet's magnetosphere.

**Bergen School** The name used to distinguish the group of meteorologists (led by **Vilhelm \*Bjerknes**, his son Jacob Bjerknes, H. Solberg, and Tor Bergeron) who, working at the Bergen Geophysical Institute, in Norway, between 1917 and 1920, established the existence and role of **\*fronts** and **\*air masses** in the atmosphere.

**Bergeron theory (Bergeron–Findeisen theory)** A theory, proposed around 1930 by T. Bergeron, and subsequently developed by W. Findeisen, that provides a mechanism for the growth of raindrops in ice/water (mixed) cloud. It is based on the differential values for saturation vapour pressure over ice and supercooled-water surfaces. At cloud temperatures of  $-12$  to  $-30$  °C air can be saturated over ice but not over water particles. Consequently, water evaporates from liquid droplets and accumulates on ice crystals by **\*deposition**. When they are large enough, the ice particles can fall through the cloud, melting to form rain drops as they pass through lower, warmer air. The process depends on there being a mixture of ice and water, and so may operate in mid- and high-latitude cloud but not in all

clouds, e.g. not in tropical clouds which are at temperatures above freezing throughout. *See also* COLLISION THEORY; ICE NUCLEUS.

**Bergman, Torbern Olof** (1735–84) Professor of chemistry at Uppsala, Sweden, Bergman made contributions to mineralogy. He was a diluvialist, believing that the Earth had an aqueous origin; some of his work on the formation of rocks was later developed and extended by \*Werner. *See* DILUVIALISM.

**Bergmann's rule** The idea, proposed in 1847 by the German anatomist and physiologist C. G. L. C. Bergmann (1814–65), that the size of \*homoiothermic animals in a single, closely related, evolutionary line increases along a gradient from warm to cold temperatures, i.e. that races of species from cold climates tend to be composed of individuals physically larger than those of races from warm climates. This is because the surface-area:body-weight ratio decreases as body weight increases. Thus a large body loses proportionately less heat than a small one. This is advantageous in a cold climate but disadvantageous in a warm one. *See also* ALLEN'S RULE; GLOGER'S RULE.

**bergschrund** A wide and deep \*crevasse found between a \*cirque glacier and its \*headwall. It forms when the glacier has developed to the stage at which it pulls away from the rock slope on its upper side. A series of small bergschrunds ('bergschrund crevasses') may form instead of the single feature.

**berg wind** A hot, dusty wind, generically of the \*föhn type, which blows across southern Africa, most frequently in winter.

**Beringia** Area comprising the Bering Strait and adjacent Siberia and Alaska. At various times in the late \*Mesozoic and in the \*Cenozoic, the strait was dry land and so provided an important migration route for plants and animals between the Palaearctic and Neartic biogeographical regions.

**Bering land bridge** The intermittent land connection between Siberia and Alaska that operated throughout the \*Cenozoic. This provided the only route into N. America for the mammals, the direct route from Europe to N. America having been interrupted by the developing Atlantic Ocean. *See also* LAND BRIDGE.

**berm** Ridge or nearly flat platform at the rear of a **\*beach** and standing just above the mean high-water mark. Its distinguishing feature is a marked break of slope at the seaward edge.

**Berman balance** See DENSITY DETERMINATION.

**Bermuda high** Anticyclonic cell in the Bermuda region that is a westward extension to, or displacement of, the **\*Azores** high.

**Bernoulli, Daniel** (1700–82) A Swiss mathematician (one of eleven eminent mathematicians his family produced over four generations), whose most important work was in the field of hydrodynamics. In his book *Hydrodynamica* (1738), he showed that the pressure within a flowing fluid depends inversely on its velocity (the greater the velocity, the lower the pressure). This is now known as Bernoulli's principle (see **BERNOULLI EQUATION**). Bernoulli was born at Groningen, the Netherlands, and educated at Basel, Switzerland, where his father had been appointed professor of mathematics on the death of his brother (Daniel's uncle) who held the post previously. Daniel obtained his master's degree at the age of 16 and his doctorate, on the action of the lungs, at 21. In 1725 he was appointed professor of mathematics at St Petersburg Academy, Russia, but left Russia in 1732. In 1733 he became professor of anatomy and botany at the University of Basel; in 1750 he became professor of natural philosophy, a post he held until his retirement in 1777. He died in Basel.

**Bernoulli equation** An equation that describes the conservation of energy in the steady flow of an ideal, frictionless, incompressible fluid. It states that:  $p_1/p_2 + gz + (v^2/2)$  is constant along any stream line, where  $p_1$  is the fluid pressure,  $p_2$  is the mass density of the fluid,  $v$  is the fluid velocity,  $g$  is the **\*acceleration** due to gravity, and  $z$  is the vertical height above a datum level.

**Berriasian** A **\*stage** of the European Lower **\*Cretaceous**, dated at 145.5–140.2 Ma ago (Int. Commission on Stratigraphy, 2004), preceded by the **\*Tithonian**, and followed by the **\*Valanginian**, for which the **\*stratotype** is at Berrias, France. See also **NEOCOMIAN**.

**bertrandite (hessenbergite)** A **\*mineral**,  $\text{Be}_4(\text{Si}_2\text{O}_7)(\text{OH})_2$ ; density 2.59–2.6 g/cm<sup>3</sup>; **\*hardness** 6–7; **\*orthorhombic**: perfect **\*cleavage**; colourless or

pale yellow; white **\*streak**; **\*vitreous** **\*lustre**; crystals thin, tabular, prismatic to needle-like; named after the French mineralogist Emile Bertrand (1844–1909), found in beryllium-bearing **\*pegmatites** in France and Russia, and may result from the alteration of **\*beryl**.

**Bertrand lens** An accessory lens which may be inserted into the light path above the **\*analyser** in a transmitted-light microscope. When determining **\*vibration** directions or **\*interference figures** using parallel or convergent polarized light, the Bertrand lens is inserted to bring the image of the interference figures into focus. Alternatively, if the Bertrand lens is absent, the **\*eyepiece** may be removed and the vibration directions or interference figures observed by looking down the microscope tube. The lens was first used in 1878 by E. Bertrand who adapted an original (1844) design by G. B. Amici.

**beryl** **\*Accessory mineral**,  $\text{Be}_3\text{Al}_2\text{Si}_6\text{O}_{18}$ ; sp. gr. 2.6–2.8; **\*hardness** 7.5–8.0; **\*hexagonal**; normally green, sometimes blue, yellow, or pink, and translucent to transparent; **\*vitreous** **\*lustre**; **\*crystals** hexagonal **\*prisms**, often striated, also occurs **\*massive**; **\*cleavage** perfect basal {001}; occurs extensively in cavities in **\*granites**, **\*pegmatites**, **\*mica** **\*schists** and **\*gneisses**, and associated with **\*rutile**. It is an **\*ore mineral** for beryllium. Transparent green varieties are emeralds, bluish-green are aquamarine, and pink are morganite.

**beta decay** Some unstable atoms decay by emitting a negatively charged beta particle (negatron) from the **\*nucleus**, often accompanied by the emission of radiant energy (**\*gamma rays**). Beta decay may be regarded as the alteration of a neutron into a proton and an **\*electron**. As a result of beta decay the **\*atomic number** of the atom is increased by one, while the neutron number is decreased by one.

**beta diagram** A stereographic diagram which represents the **\*trend** and **\*plunge** of the line produced by the intersection of two planes. In its application in structural geology, the **\*dip** and **\*strike** of a **\*fold** surface are recorded as a **\*great** circle, and two great circles representing both limbs of a fold intersect along a line called a  $\beta$ -axis (in this case the fold axis).

**beta-mesohaline water** See HALINITY.


**bevelled cliff** A sea cliff whose upper part has been trimmed to a relatively low angle by *\*Quaternary* *\*periglacial* processes, while the lower part is still steep as a result of recent marine activity. Such cliffs are common in south-west England. See *COASTAL PROCESSES*.

**Bianca (Uranus VIII)** One of the lesser satellites of *\*Uranus*, with a diameter of 22 km. It was discovered in 1986.

**biaxial interference figure** See *INTERFERENCE FIGURE*.

**Bibymalagasia** An order of mammals described in 1994 by R. D. E. MacPhee, based on the enigmatic genus (see *ENIGMATIC TAXON*) *\*Plesioxycteropus*, known from sub-Recent fossil material from Madagascar. The order is distinguished by having large, perforating, transarcual canals in the neural arches of the lumbar, posterior thoracic, and anterior sacral vertebrae (see *VERTEBRA*); a posteriomedial process on the *\*astragalus*, with a ventral groove for flexor tendons; and large ischial expansions (see *ISCHIUM*). *Plesioxycteropus* was formerly assigned to the order Tubulidentata. It was a ground-dwelling insectivore, sometimes called the Malagasy aardvark, although it was not related to the aardvark.

**biconical** Applied to a gastropod (*\*Gastropoda*) shell where the shell is in the shape of two cones with the bases in contact. The axial whorls make one cone, the last-formed whorl or living chamber the other cone.




**Biconical**

**BIF** See BANDED IRON FORMATION.

**bifurcation** In a phylogenetic (see PHYLOGENY) tree, the dichotomous forking of an ancestral \*branch which indicates a speciation event.

**bifurcation ratio** Dimensionless number denoting the ratio between the number of streams of one order (see STREAM ORDER) and those of the next-higher order in a drainage network. It may be a useful measure of proneness to flooding: the higher the bifurcation ratio, the greater the probability of flooding.



bifurcation  
ratio = 2.08



bifurcation  
ratio = 12

### **Bifurcation ratio**

**big bang theory** The current explanation for the origin of the universe, in which it expands and evolves from an initial very high-temperature condition about 13.7 billion years ago. The expansion time is given from the reciprocal of the Hubble constant (the rate at which galaxies are receding). All all-pervasive background radiation of 3 K is considered to be

residual from the big bang and is the strongest supporting evidence for the theory.



<http://www.big-bang-theory.com/>

- An overview of the big bang theory.

**bilateral symmetry** The condition, found in many organisms, where one half of the body or structure is the mirror image of the other. Sometimes bilateral symmetry is superimposed on another kind of symmetry, e.g. in some echinoids (*\*Echinoidea*) where basic pentamerous (five-sided) symmetry has a bilateral symmetry superimposed upon it.

**Bilateria** The subkingdom that contains all animals (kingdom Animalia) possessing *\*bilateral symmetry* and three embryonic germ layers (the cells that will develop into the organs of the body). It is the largest animal subkingdom. *Compare* **RADIATA**.

**billow clouds** Parallel rolls of cloud with distinct clear areas between the cloud bands; it is a type of the cloud variety *\*undulatus*.

**bilophodonty** In some mammals, a condition in which the four cusps on the molar teeth are joined by two transverse ridges. *See also* **LOPHODONT**.

**bimodal distribution** A distribution of data characterized by two distinct populations. For example, a bimodal grain size will be characterized by two *\*particle size* modes. A bimodal palaeocurrent distribution will exhibit two main current directions (not necessarily opposing directions, which would be termed a 'bipolar' distribution).

**bin** In a range of data broken into a series of equal intervals, a single interval.

**binary system** 1. Chemical system of two components, e.g. MgO–SiO<sub>2</sub>. 2. *See* **STAR PAIR**.

**bindstone** An *\*autochthonous* *\*carbonate* rock, with original components organically bound during deposition. The organisms encrust and bind the finer *\*matrix* *\*sediment* together, but the binding organisms need not

necessarily be preserved, as in the case of *\*stromatolites*. See [EMBRY AND CLOVAN CLASSIFICATION](#).

**Bingham fluid** A viscous fluid that possesses a yield strength which must be exceeded before the fluid will flow. Most *\*lava* flows are examples of Bingham fluids. When an initial *\*shear stress* is applied to a fluid lava (e.g. by increasing the slope angle) it will not begin to flow immediately. The slope angle, and therefore shear stress, must be increased until the yield strength of the fluid is exceeded, after which flow will occur. This contrasts with a Newtonian fluid, which has zero yield strength and will flow on any slope.

**binomial distribution** In statistics, a discrete *\*probability distribution* of the number of outcomes of a particular kind occurring for a set number of trials, where one of two outcomes is possible, each trial is independent, and the probability of a particular outcome is constant.

**bio-** **1.** From the Greek *bios* meaning 'human life', a prefix associating the word to which it is attached with living organisms or processes. **2.** Prefix for *\*allochems* composed of *\*bioclastic* material. See [FOLK LIMESTONE CLASSIFICATION](#).

**biochemical oxygen demand** See [BIOLOGICAL OXYGEN DEMAND](#).

**biochron** The length of time represented by a biostratigraphic *\*zone*.

**biochronology** Measurement of units of geologic time by means of biological events. Biochronologists often derive their correlations from widespread and distinctive events in the biological history of the world, based on the first and last appearances of organisms.

**bioclast** A single, often broken, shell or *\*fossil* fragment.

**biocoenosis** See [LIFE ASSEMBLAGE](#).

**biofacies** Rock unit, or an association of rock units, characterized by the presence of a *\*fossil* assemblage that is restricted to that particular *\*facies* and that is typical of a specific environment. Compare [TAPHOFACIES](#).

**biogenesis** Principle that a living organism can arise only from another living organism, a principle contrasting with concepts such as that of the spontaneous generation of living from non-living matter. The term is

currently more often used to refer to the formation from or by living organisms of any substance, e.g. coal, chalk, chemicals, etc.

**biogenetic law** Law formulated by the embryologist E. K. von Baer (1792–1876) stating that the early stages of development in animal species resemble one another, the species diverging more and more as development proceeds.

**biogenic** Applied to material, processes, or activities of living or once-living organisms.

**biogenic deposit** The formation of rocks, traces (see **TRACE FOSSIL**), or structures as a result of the activities of living organisms.

**biogeochemical cycle** Movement of chemical elements from organism to physical environment to organism, in a more or less circular pathway. They are termed ‘nutrient cycles’ if the elements concerned are essential to life. An element may be solid, liquid, or gaseous, or form different chemical compounds, in the various parts of the cycle. Amounts in the inorganic **\*reservoir pools** are usually greater than those in the **\*active pools**. Exchange between the system components is achieved by physical processes (e.g. **\*weathering**) and/or biological processes (e.g. protein synthesis and decomposition). The latter form the vital negative-feedback mechanisms that regulate the cycles. Cycles may be described as varying from perfect to imperfect. A perfect cycle (e.g. the nitrogen cycle) has a readily accessible abiotic, usually gaseous, reservoir and many negative-feedback controls. By contrast, the phosphorus cycle, which has a sedimentary reservoir accessed only by slow-moving physical processes, has few biological feedback mechanisms. Human activities can disrupt these cycles, leading to pollution. Theoretically, perfect cycles are more resilient than imperfect cycles.

**biogeochemical exploration** See **GEOBOTANICAL EXPLORATION**.

**biogeochemistry** Science concerned with the effects of living things on subsurface geology; or with the distribution and fixation of chemical elements in the biosphere. Its principles are applied to the systematic collection and analysis of plants in the exploration for mineral deposits. It is also the study of the chemistry of organic sediments and of the chemical

composition of fossils and fossil fuels. *See also* [GEOBOTANICAL EXPLORATION](#).

**biogeocoenosis** *See* [ECOSYSTEM](#).

**biogeography** The scientific study of the past and present geographical distribution of plants and animals at different taxonomic levels. Modern biogeography also lays great stress on the ecological character of the world vegetation types, and on the evolving relationship between humans and their environment. *See also* [PALAEOBIOGEOGRAPHY](#).

**biogeomorphology** The study of the interrelationships between [\\*geomorphology](#) and the living organisms that inhabit the resulting land-forms. Organisms influence many geomorphological processes and are affected by them.

**bioglyph** Ornamentation on the side of a [\\*trace fossil](#) burrow.

**bioherm** A build-up of largely *in situ* organisms that produces a [\\*reef](#) or mound of organic origin.

**biohorizon** 1. An interface between [\\*strata](#) at which some [\\*biostratigraphy](#) change has occurred. 2. A biostratigraphic [\\*marker bed](#).

**bioimmuration** A type of [\\*fossilization](#) in which soft-bodied encrusting organisms are overgrown by other encrusting organisms. As the soft-bodied organisms disappear, they leave natural moulds on the underside of the layer above them.

**biointermediate elements** Elements that are partially depleted in surface waters as a result of biological activity. Four such elements are known: Ba, Ca, C, and Ra. *Compare* [BIOLIMITING ELEMENTS](#); [BIOUNLIMITING ELEMENTS](#).

**biolimiting elements** A few elements (N, P, and Si) which are almost totally depleted in surface waters, relative to deep water, by biological activity. As these elements are essential to living organisms, the depletion limits further biological production until the scarce elements are replaced, e.g. by [\\*upwelling](#). *Compare* [BIOINTERMEDIATE ELEMENTS](#); [BIOUNLIMITING ELEMENTS](#).

**biolithite** A *\*carbonate* rock formed of organisms that grew and remained in place, comprising a rigid framework of organisms, together with associated debris. A *\*reef* represents a typical biolithite. See FOLK LIMESTONE CLASSIFICATION.

**biological marker** See BIOMARKER.

**biological oxygen demand (BOD, biochemical oxygen demand)** Indicator of the polluting capacity of an effluent, where pollution is caused by the take-up of dissolved oxygen by micro-organisms that decompose the organic material present in the effluent. It is measured as the weight (mg) of oxygen used by 1 litre of sample effluent stored in darkness at 20 °C for five days.

**biomagnetism** The influence of the magnetic field on living organisms, which is strong in some species. Certain *Spirillum* bacteria orient themselves along the geomagnetic lines of force. Higher organisms, e.g. snails, bees, birds, porpoises, and possibly humans, appear able to sense and utilize *\*geomagnetic field* directions for purposes of orientation. Strong magnetic fields may be detrimental to health in humans, but the effects of medium strength fields have not been studied. Magnetic fields are also used for scanning biological organisms.

**biomarker (biological marker, chemical fossil)** An organic compound derived from living organisms that occurs in trace amounts in *\*fossils* and *\*sediments*.

**biome** See COMMUNITY.

**biomicrite** A *\*limestone* consisting of *\*bioclasts* set in a *\*micrite* *\*matrix*. It is the product of a poorly sorted accumulation of shell fragments and mud. See FOLK LIMESTONE CLASSIFICATION.

**biomineralization** The incorporation of inorganic compounds, such as salts, into biological structures, often to lend them hardness or rigidity. Biomineralization first occurred in, and defines, the *\*Cambrian* period beginning 542 million years ago, in *\*Brachiopoda*, *\*Trilobita*, *\*Ostracoda*, and *\*Graptolithina*. In vertebrates, *\*hydroxyapatite* usually occurs, in invertebrates inorganic minerals are more varied: *\*calcite* and *\*aragonite* (a harder, less stable form of calcite) are common, permeating chitin (see

SKELETAL MATERIAL) to form the hard *\*exoskeletons* of *\*Arthropoda* and also forming the calcareous material of shells; in *\*Radiolaria* and some *\*Porifera*, the skeleton is made of opaline silica; Radiolarians occasionally have a strontium sulphate instead of siliceous skeleton.

**biophile** Applied to those elements required by, or found in, living plants and animals, including C, H, O, N, P, S, Cl, I, Br, Ca, Mg, K, Na, V, Fe, Mn, and Cu. *Compare* *ATMOPHILE*; *CHALCOPHILE*; *LITHOPHILE*; *SIDEROPHILE*.

**biosparite** A *\*limestone* consisting of *\*bioclasts* together with a sparry *\*calcite* *\*cement* (*\*sparite*). It is the product of an accumulation of clean-washed, mud-free shell debris, with diagenetic cement growth (*see* *DIAGENESIS*) in *\*pore spaces*. *See* *FOLK LIMESTONE CLASSIFICATION*.

**biosphere** The part of the Earth's environment in which living organisms are found, and with which they interact to produce a steady-state system, effectively a whole-planet *\*ecosystem*. Sometimes it is termed 'ecosphere' to emphasize the interconnection of the *\*biotic* and *\*abiotic* components.

**biostratigraphic interval zone** *See* *INTERVAL ZONE*.

**biostratigraphic unit** A unit of *\*strata* characterized by a particular content of *\*fossils*; these were deposited at the same time as the *\*sediments* and distinguish the unit from adjacent strata. A biostratigraphic unit may be of *\*chronostratigraphy* or environmental significance.

**biostratigraphic zone** *See* *ZONE*.

**biostratigraphy** Branch of stratigraphy that involves the use of *\*fossil* plants and animals in the dating and correlation of the stratigraphic sequences of rock in which they are discovered. A *\*zone* is the fundamental division recognized by biostratigraphers.

**biostratinomy** The study of the sedimentological processes by which a *\*fossil* assemblage is formed. This includes the transport, deposition, and compaction of the organic material.

**biostrome** A layered, sheet-like accumulation of *in situ* organisms. It differs from a *\*bioherm* in its geometry, lacking a mound-like or *\*reef*-like form.

**biota** The living organisms occupying a place together, e.g. marine biota, terrestrial biota.

**biotic** Applied to the living components of the **\*biosphere** or of an **\*ecosystem**, as distinct from the non-living, **\*abiotic**, physical and chemical components.

**biotic index** Scheme for grading river quality according to the diversity and abundance of the river fauna present, with particular reference to 'key species' with known pollution tolerances. The scale ranges from 10 (clean water with diverse fauna) to 0 (grossly polluted water with no fauna or with only a few **\*anaerobic** organisms).

**biotite** An important rock-forming silicate **\*mineral**; a member of the **\*mica** group  $K_2(Mg,Fe)_6[Si_3AlO_{10}]_2(OH,F)_4$ , in which the Mg/Fe ratio is less than 2:1; biotite forms a series with **\*phlogopite** (of similar composition, but with Mg/Fe greater than 2:1) and may have  $Fe^{3+}$ ,  $Ti^{4+}$ , and  $Al^{3+}$  also present in the formula; sp. gr. 2.7–3.3; **\*hardness** 2–3; **\*monoclinic**; black, dark brown, or greenish-black; **\*vitreous** to **\*sub-metallic** **\*lustre**; **\*crystals** **\*tabular** and pseudo-hexagonal, with lamellar aggregates and flakes; **\*cleavage** perfect basal {001}; occurs in **\*granites**, **\*syenites**, and **\*diorites**, in **\*schists** and **\*gneisses**, in contact **\*metamorphic rocks**, and often in **\*sedimentary rocks**.

**bioturbation** The disruption of **\*sediment** by organisms, seen either as a complete churning of the sediment that has destroyed depositional **\*sedimentary structures**, or in the form of discrete and clearly recognizable **\*burrows**, **\*trails**, and traces (see **TRACE FOSSIL**).

**biounlimiting elements** Elements (e.g. B, Mg, Sr, and S) that show no measurable depletion in surface waters compared to deep waters as a result of biological activity. Compare **BIOLIMITING ELEMENTS**; **BIOINTERMEDIATE ELEMENTS**.

**biozone 1.** The total, global, time-stratigraphic range of a given **\*taxon**, i.e. all the rocks laid down in the time interval during which that taxon existed.  
**2.** A shortened form of 'biostratigraphic zone'. See **ZONE**.

**bipolar distribution** See **BIMODAL DISTRIBUTION**.

**bipolar seesaw** An asymmetry in which climatic cooling at one pole is accompanied by warming at the other.

**bipyramid** In crystallography, a pyramid defines a *\*crystal face* which cuts the vertical 'c' (or 'z') axis and one or two of the horizontal 'a' (or 'x') and 'b' (or 'y') axes. In general, the prefix 'bi-' refers to the repetition of faces about a plane of symmetry; 'di-' usually suggests the presence of an axis of symmetry about which the pyramid may be rotated.

**biramous** Applied to limbs that are two-branched, e.g. in many *\*Arthropoda*.

**bird** 1. See *AVES*. 2. See *AEROMAGNETIC SURVEY*.

**birdseye fabric (fenestral fabric)** A *\*fabric* characterized by irregular, *\*equant* cavities in *\*sediment*, several centimetres across, often filled with sparry *\*calcite* (*\*sparite*). It is thought to form as the result of the entrapment of gas, and desiccation within *\*intertidal* to *\*supratidal* muddy *\*carbonate* sedimentary environments. See *FENESTRAE*.

**bireflectance** In reflected-light or ore microscopy, the ability of a *\*mineral* to reflect different quantities of plane-polarized light in different orientations on rotation of the *\*stage*. This property is often related to reflection *\*pleochroism* and is diagnostic of some minerals, e.g. *\*covellite*.

**birefringence (double refraction)** In optical mineralogy, the ability of *\*anisotropic \*minerals* to split *\*plane-polarized* light into two rays as it passes through them. The rays vibrate at right angles to each other and travel along different path lengths within the mineral, depending on the degree of birefringence encountered. As a result, when the two sets of rays emerge from the mineral one set has travelled farther than the other and is retarded. The retardation shows itself as a range of interference colours when the *\*analyser* is inserted, and may be determined by reference to an interference colour chart. See *MICHEL-LÉVY CHART*.

**birefringence chart** See *MICHEL-LÉVY CHART*.

**BIRPS** Acronym for the British Institutions Reflection Profiling Syndicate. It is mainly concerned with the deep lithosphere.

**Birimian orogeny** A phase of mountain building, affecting **\*Proterozoic \*greenstone belts** in an area of what is now W. Africa.

**bischofite** A hydrated **\*evaporite** mineral,  $MgCl_2 \cdot 6H_2O$ , deposited in environments where the **\*relative humidity** is about 30%.

**bise** The Swiss term for a cold, dry, northerly or north-easterly winter wind blowing from the European Alps and affecting adjacent regions of southern Switzerland and eastern France (at Languedoc the 'bise noire' is associated with heavy cloud).

**biserial** **1.** Applied to those graptoloids (**\*Graptoloidea**) in which **\*thecae** occur side by side on each side of the **\*stipe**. **2.** In crinoids (**\*Crinoidea**), applied to the side-by-side condition of the brachial plates (see **BRACHIA**) in the arms. **3.** See **TRISERIAL**.

**bismuth, native** Metallic element, Bi; sp. gr. 9.7; **\*hardness** 2.5; greyish-white; soft or as **\*rhombohedral \*crystals**; occurs native in veins and **\*granite \*pegmatites** associated with tin, silver, cobalt, and nickel mineralization.

**bismuthinite** **\*Sulphide** mineral,  $Bi_2S_3$ ; sp. gr. 6.5; **\*hardness** 2; greyish-white; **\*metallic** lustre; **\*massive**; occurs in mineral veins associated with **\*granites** and tin mineralization.

**bistatic radar** A technique for studying the electrical properties of a planetary surface by the reflection of radio waves. If the transmitter and receiver are located at different places, the term 'bistatic radar' is used; if in the same place, this is called 'monostatic radar'. In the study of the **\*Moon** with bistatic radar during the Apollo missions, the transmitter was located on the command module orbiting the Moon, while the receiver was located on the Earth.

**bisulcate** Applied to shells that possess two grooves or depressions; the ventral valves of brachiopods (**\*Brachiopoda**) are commonly bisulcate. The term is also applied to the **\*venter**, which has twin grooves, in some ammonoids (**\*Ammonoidea**).

**bit** **1.** The cutting part of a **\*drill** stem, which may either break or crush the rocks as the drill stem rotates. **2.** In computing technology, a binary digit or element of information.

**Bitauanian** See ARTINSKIAN.

**bitheca** One of the three types of graptolite *\*thecae*, possibly containing the male zooid. Compare AUTOTHECA; STOLOTHECA. See GRAPTOLITHINA; DENDROIDEA.

**bitter lake** A saline lake which is rich in sulphates and carbonates, dominated by high concentrations of sodium sulphate.

**bittern salts** A bitter-tasting (hence the name) solution that remains after *\*halite* (NaCl) is removed from sea water or brine, either naturally or as a result of desalination or sea-salt production. It is rich in KCl, MgCl<sub>2</sub>, MgSO<sub>4</sub>, and *\*double salts*.

**bitumen** Naturally occurring, inflammable, solid or semi-solid *\*hydrocarbons*, black or dark brown in colour, with characteristic pitch odour, and burning with a smoky flame. It is also the group name for *\*asphalts*, mineral waxes, and related substances.

**bituminous coal** A type of coal between *\*lignite* and *\*anthracite* in *\*rank*, with 18–35% *\*volatiles*, which softens and swells on combustion, either caking or non-caking, and burns with a smoky flame. It is used in coke manufacture.

**bivalves** See BIVALVIA.

**Bivalvia (bivalves)** (Pelecypoda, Lamellibranchia; phylum *\*Mollusca*) A class of molluscs in which the body is laterally compressed and is enclosed between two oval or elongated valves. The valves are united dorsally by a toothed *\*hinge* and in most species the valves are *\*bilaterally symmetrical* along the plane of junction (*\*commissure*) between them. The valves are opened by a horny, elastic ligament and closed by the action of one or two adductor muscles. Large, modified, ciliated (see CILIUM) gills are involved in food collection and bivalves are entirely aquatic. They are adapted to various modes of life, e.g. boring, burrowing, free-living, and sessile, and these modes of life are generally reflected in the shape of the shell which is modified in various ways. Bivalves first appear in the Lower *\*Cambrian* and are generally of limited abundance in the *\*Palaeozoic*. They become more abundant from the *\*Mesozoic* and they now form the second largest molluscan class, with more than 20 000 species.

**Bjerknes, Vilhelm Frimann Koren** (1862–1951) A Norwegian geophysicist and meteorologist who was born and died in Oslo. As a student, he helped his father, who was professor of mathematics at the Christiania University (now the University of Oslo). He himself held professorships at Stockholm and Leipzig before returning to Norway in 1917 to found the Bergen Geophysical Institute (*see* [BERGEN SCHOOL](#)). During the First World War, Bjerknes established a series of weather stations throughout Norway. Information from these allowed Bjerknes and his colleagues, who included his son Jakob (1897–1975) and Tor Harold Percival Bergeron (1891–1977), to develop their theory of air masses bounded by fronts.

**black body** A body which absorbs electromagnetic energy perfectly. If the body remains at constant temperature then it also radiates *\*electromagnetic radiation* perfectly in equilibrium with that which it absorbs. *See also* [STEFAN–BOLTZMANN LAW](#).

**black earth** *See* [CHERNOZEM](#).

**Blackett, Patrick Maynard Stuart** (1897–1974) A British physicist at Imperial College, London, Blackett worked on *\*magnetometers* during the Second World War, and subsequently developed an instrument capable of detecting very small magnetic fields. This led him to the serious study of *\*palaeomagnetism*. He tried, without success, to demonstrate that magnetism is a property of massive rotating bodies.

**black ice** Type of glazed frost (e.g. on roads or the superstructures of ships), caused when water close to freezing temperature falls on a surface that is below freezing temperature. The thin sheet of ice is transparent and, unlike white hoar frost or rime, may be difficult to see. *See also* [GLAZE](#).

**black jack** *See* [SPHALERITE](#).

**Blackriverian** *See* [COSTONIAN](#).

**black shale** A *\*mudstone* with high concentrations of organic material, which is deposited in *\*euxinic* environments. Such sediments form important *\*hydrocarbon \*source rocks*.

**BlackSky Constellation** A constellation of sixty Earth-observation satellites, each designed for a 3-year mission, planned by BlackSky Global,

of Seattle, Washington, USA, the first of which, a pathfinder *\*microsatellite*, was launched on 26 September 2016, from the Satish Dhawan Space Centre, India, into a Sun-synchronous orbit. The company aimed to have the full constellation in orbit by 2019. The satellites will provide 1-m-resolution colour images of the entire Earth.

**'black smoker'** See *HYDROTHERMAL VENT*.

**blade** A term describing grain shape; bladed grains have ratios of less than 2:3 for both the intermediate:long diameter and the short:intermediate diameter. See *PARTICLE SHAPE*; *ZINGG DIAGRAM*.

**bladed** In *\*mineralogy*, describes the flat and elongated form or shape of a *\*mineral*, rather like a knife blade or lath, e.g. *\*kyanite*.

**Blake** See *BRUNHES*.

**blanket bog** Ombrogenous bog community, typical of flat or moderately sloping areas in very wet, oceanic climates with high humidity. In the British Isles, blanket bogs are widespread on the Pennine summits, in north-west Scotland, and in parts of Ireland.

**-blast** A suffix applied to metamorphic terms and used to indicate the *in situ* growth of crystals during metamorphic *\*recrystallization*. For example, the term 'porphyroblast' refers to a large, well-formed crystal which grew *in situ* as a rock was metamorphosed.

**blasto-** A prefix applied to textural terms and used to indicate the partial or complete replacement of a *\*textural* element in the original rock by crystals which have grown during *\*metamorphic \*recrystallization*. For example, the term 'blastoporphyratic' refers to a porphyritic *\*igneous* rock (see *PORPHYRY*) which has undergone metamorphism, and in which the *\*phenocrysts* remain as recognizable textural relicts even though each has been replaced by an aggregate of new metamorphic minerals.

**blastoporphyratic** See *BLASTO-*.

**Blastozoa** (phylum *\*Echinodermata*) Subphylum of extinct, stalked echinoderms, ranging from the *\*Silurian* to the *\*Permian*. The stem (rarely preserved) was surmounted by a bud-like *\*theca* with five prominent petaloid *\*ambulacra* showing a well-developed *\*pentameral symmetry*. The

theca had thirteen major plates arranged in three circlets. The ambulacra each had a central food groove, with delicate food-gathering appendages called **\*brachioles** to either side. Underlying the ambulacra were the characteristic hydrospires, thought to have functioned in respiration. These were thin-walled, calcareous infoldings (more rarely simple tubes), forming water conduits that communicated with the outer environment via marginal pores in the ambulacra and five apertures around the mouth (the spiracles). Blastozoa are not common as fossils, except locally in shallow-water calcareous rocks, often associated with **\*reef \*limestones**.

**blast ratio** The ratio of the mass of explosive used and the amount of rock broken by an explosion, usually given as tonnes of rock to one kilogram of explosive.

**B-layer** The uppermost part of the **\*Earth's \*mantle**, mostly above the first major phase change at about 370 km depth and below the **\*Mohorovičić discontinuity**.


**bleb** A small, rounded inclusion of one mineral within a large mass of another.

**bleicherde** Ashy-grey soil that forms the **\*leached** layer in a **\*podzol** soil.

**blind hole** A **\*borehole** from which all the **\*drilling** fluid has escaped into the surrounding rocks.

**blind pores** See EFFECTIVE POROSITY (2).

**blind thrust** A **\*thrust fault** that terminates before reaching the surface. There is no evidence of the fault at the surface, so it may remain undetected until it ruptures.



**Blind thrust:** A blind thrust is a thrust fault that terminates below the surface.

**blind valley (steephead valley)** A steep-sided, flat-bottomed valley that terminates in a steep gradient. Such valleys occur in limestone or *\*karst* environments, where a subterranean stream erodes a channel that later collapses, forming the valley bottom, and the stream then flows over the surface from a *\*spring*, disappearing below ground again at the lower end.

**blind zone** A layer of rock that cannot be detected by *\*seismic \*refraction* methods, usually because it has a seismic velocity lower than that of the overlying layer(s).

**blizzard** A storm of blowing snow with high winds and low temperatures. Blizzards are a notable climatic feature of the northern and central parts of the USA in winter, and are related to depression tracks. In the USA, a blizzard is defined by the National Oceanic and Atmospheric Administration (NOAA) as a storm with winds of at least 56 km/h, temperatures below  $-6.7\text{ }^{\circ}\text{C}$ , and enough falling or blowing snow to reduce visibility to less than 0.4 km. In a severe blizzard, wind speed is at least 72.5 km/h, temperatures below  $-12.2\text{ }^{\circ}\text{C}$ , and visibility is close to zero.

**block-and-ash deposit** The deposit formed by a *\*nuée ardente*.

**blockfield (felsenmeer)** Spread of coarse, angular, frost-shattered rock debris resting on a level or gently sloping upland surface, and found in present or former periglacial environments. *See also* FROST WEDGING.

**block glide** The sliding movement of a large block of rock over a surface that has been lubricated. Alternatively, the block may be carried downslope by the \*plastic deformation of underlying material.

**blocking** In synoptic meteorology, the establishment in the mid-latitudes of a high-pressure system that interrupts or diverts for a considerable period the typically eastward movement of \*depressions and other synoptic features in the \*zonal flow. Over western Europe, for example, blocking often forces depressions to move northward toward Scandinavia or southward over southern France and Spain.

**blocking anticyclone** *See* BLOCKING HIGH.

**blocking high (blocking anticyclone)** In synoptic meteorology, an \*anticyclone with deep circulation so placed as to interrupt or divert the eastward movement of the typical succession of low-pressure features (\*depressions) in the \*zonal flow of mid-latitudes. At such times the usual mid-latitude zonal flow is diverted into more \*meridional flow around the high-pressure area, where settled weather can then result. *See also* AIR MASS.

**blocking temperature ( $T_B$ )** The temperature at which a \*thermal remanent magnetization becomes locked in a rock for geologic times.

**blocking volume ( $V_B$ )** The grain volume at which a \*chemical remanent magnetization becomes locked in a rock for geologic times.

**block volume** The natural size of a block of rock bounded by \*joint (or other discontinuity) planes. Blocks are defined by their length of side as: very large (more than 2 m); large (600 mm–2 m); medium (200–600 mm); small (60–200 mm); and very small (less than 60 mm).

**blocky lava** A surface flow of hot, molten \*lava covered in a carapace of crystalline, angular blocks which tend to be smoothly faceted and may have dimensions up to several metres. The blocks, which have the same composition as the flow interior, are formed by fragmentation of the chilled flow surface as lava continues to move within the flow interior. Blocky lava

morphology is usually confined to lavas of high *\*viscosity* and intermediate to high *\*silica* contents.

**bloedite** A non-marine *\*evaporite* mineral,  $\text{Na}_2\text{SO}_4 \cdot \text{MgSO}_4 \cdot 4\text{H}_2\text{O}$ .

**blood rain** Phenomenon of reddish coloured rainfall, caused by dust particles that have been lifted up from arid areas and carried long distances by the winds before they are washed out in precipitation. Saharan red dust sometimes occurs in rainfall over parts of Europe, even as far north as Finland.

**blow-hole** See COASTAL PROCESSES.

**blow-out** Wind-eroded section of a sand *\*dune* that has been largely stabilized by vegetation. *\*Erosion* results from a break in the vegetation cover, due typically to overgrazing or to recreational pressure. A parabolic dune may result.

**blue** A quarrymen's term, applied to stone and meaning 'hard'.

**blueberries** Small mineral spheres observed in 2004 by the Mars Exploration Rover Opportunity. Despite their nickname, the spherules are grey rather than blue in colour, and are only a few millimetres across. The rover's Mössbauer spectrometer (see MÖSSBAUER SPECTROMETRY) revealed that the blueberries are rich in *\*hematite*.

**blue-green algae** See CYANOBACTERIA.

**blue jet** A *\*transient luminous* event, first documented in 1994, in which light flashes from the electrically active core of a thunderstorm, typically propagating upward from the top of the cloud in a cone about 15° wide and disappearing after about 0.2 s at an altitude of 40–50 km.

**Blue John** See FLUORITE.

**blue Moon (blue Sun) 1.** Occasional appearance of the Moon or Sun when partly obscured by large particles in the atmosphere, as in dust storms, or following forest fires or great volcanic explosions. When the Moon or Sun is viewed through dust or smoke trails it usually appears to be very white, but when the suspended particles are predominantly of one size it sometimes appears blue, at other times green or orange. The phenomenon is attributed to diffraction, although no full explanation seems to be known.

The smaller the particles, the more the colour of the Sun or Moon tends to the blue end of the spectrum. The phenomenon is believed to be more common in China than elsewhere. (In the neighbourhood of the 1883 Krakatoa volcanic explosion the Sun was seen as an azure blue sphere.) 2. In 1946 'blue Moon' was defined in the magazine *Sky and Telescope* as the third full Moon in a calendar season in which there are four.

**blueschist** \*Metamorphic rock that has undergone \*regional metamorphism at low temperatures and high pressures. Blueschists contain abundant blue \*glaucophane, which is a blue \*amphibole. Blueschists are usually associated with \*destructive plate boundary environments.

**blueschist facies** See GLAUCOPHANE-SCHIST FACIES.

**blue starter** A \*transient luminous event that is brighter than a \*blue jet but reaches upward to only about 20 km. It is reputed to occur over regions of heavy hailstorms.

**blue Sun** See BLUE MOON.

**bocca** An Italian word meaning 'opening' or 'mouth', used originally by Italian volcanologists when referring to an opening through which \*lava is effusing. The term is used extensively to describe secondary effusion points on the surface of active compound lava flows, as well as primary effusion points at the base of cinder cones lying on major eruptive fissure lines.

**BOD** See BIOLOGICAL OXYGEN DEMAND.

**Bode, Johann Elert** (1747–1826) A German astronomer who popularized the theory, known later as Bode's law or, more correctly, as the \*Titius–Bode law, that there is a simple arithmetical relationship between the distances from the \*Sun to the planets of the solar system. After the discovery of \*Neptune, which did not conform with the 'law', the idea fell into disrepute.

**body chamber** The last-formed chamber of a cephalopod (\*Cephalopoda) shell, which houses the animal's body.

**body plan** See BAUPLAN.

**body wave** A seismic wave which propagates through the body of a medium; \*P-waves and \*S-waves are body waves. These contrast with

\*surface waves.

**boehmite** An aluminium hydrate  $\gamma$ -AlO(OH), boehmite forms a continuous series with its \*polymorph \*diaspore  $\alpha$ -AlO(OH) and is a constituent of \*bauxite and \*laterite; sp. gr. 3.0; \*hardness 4; \*orthorhombic; usually white; normally occurs as microscopic grains, or \*pisolitic aggregates; \*cleavage good {010}; occurs extensively in the weathering profiles of rocks together with \*gibbsite, diaspore, and \*kaolinite. It is used for aluminium production and is named after the 19th-century German chemist J. B. Böhm.

**Bogen structure** See COAL MACERAL.

**boghead coal** \*Sapropelic coal with a greasy, dull \*lustre and no lamination; in appearance and combustible properties it resembles \*cannel coal but with a brown or yellow \*streak. It has a high percentage of algal matter and \*volatiles, and when distilled gives a high yield of tar and oil.

**bog iron ore** An impure, limonitic deposit (see LIMONITE), usually porous, and probably formed as a result of bacterial action in swampy conditions. It is a low-grade iron \*ore which was used extensively in early iron smelting.

**bolide** Large \*meteor that explodes in passing through the Earth's atmosphere. The term is sometimes used synonymously with \*fire-ball, but some people reserve 'bolide' for an exploding meteor, and 'fire-ball' for a less-bright object.

**Bolindian** A \*stage (450–443.7 Ma ago) of the Upper \*Ordovician of Australia, underlain by the \*Eastonian and followed by the \*Keiloran.

**Bølling-Allerød interstadial** A warm period that occurred during the final stages of the \*Devensian glaciation that began abruptly about 14 700 years ago following the Older \*Dryas, ended about 12 700 years ago, and was followed by the Younger Dryas. Formerly the \*interstadial was divided into separate Bølling and Allerød episodes, but it is now apparent that they constituted a single interstadial interrupted by several cooler episodes. Pollen records for the NW European area indicate a cool temperate flora with birch (*Betula* species) widespread.

**bolson** \*Intermontane basin extending from the divide of one block-faulted mountain to the divide of the adjacent mountain. The surface form is made

up of mountain front, *\*bajada*, *\*pediment*, and *\*playa*. It is classically described for the *\*basin-and-range* province of the western USA.

**Boltwood, Bertram Borden** (1870–1927) Professor of radiochemistry at Yale University, in 1907 Boltwood made early determinations of the age of the *\*Earth* using uranium:lead ratios. He was able to show a long geologic timescale of up to 2000 Ma. See *RADIOMETRIC DATING*.

**Boltzmann constant (*k*)** The ratio of the ideal gas constant to the *\*Avogadro* constant:  $k = R/N_A = 1.3805 \times 10^{-23}$  J/K, where *R* is the ideal gas constant (equal to 8.3 J/mol/K), and  $N_A$  is the Avogadro constant.

**bomb** See *VOLCANIC BOMB*; *VOLCANO*.

**bomb sag** The deformation of primary, unconsolidated, volcanoclastic bedding structures by the impact of a large ballistic block. The block, which can be a *\*volcanic bomb* or a fragment of crystalline *\*country rock*, is ejected from its source vent during a period of violent explosive activity. The asymmetry of the deformation structure it produces can be used to locate the position of the source vent.

**bone** The skeletal tissue of *\*vertebrates*. Bone is composed of about 70% inorganic calcium salts, mostly *\*hydroxyapatite* but carbonate, citrate, and fluoride amines are also present. The organic component is mostly made up of the structural protein *\*collagen*.

**bony fish** See *OSTEICHTHYES*.

**Boomer** A marine seismic source in which capacitor plates are highly charged and then allowed to discharge via a transducer in the water. Eddy currents induced in the transducer plates force the plates apart, thus producing a low-pressure region between the plates. Water implodes into this region, generating a pressure wave. The Boomer (the name is a commercial trademark) is used in high-frequency marine surveys as a low-energy (less than 1 kJ), high-resolution seismic source. The term ‘boomer’ is also used colloquially to describe a high-amplitude, low-frequency reflection event associated with a large and distinctive reflector.

**Boomerangian** An Australian *\*stage* (504–501 Ma ago) of the Middle *\*Cambrian*, preceded by the *\*Undillian* and followed by the *\*Mindyallan*.

**bootstrapping** In statistics, an approach to estimating the robustness or degree of error of a statistic when the distribution from which the statistic is drawn is unknown. An estimate of the distribution is constructed based on the data of the sample being analysed, often by repeatedly random sampling from the data itself and recalculating the statistic in order to produce a statistic distribution.

**bora** Regional term for a cold and typically very dry wind from the north-east, blowing down from the mountains on the eastern side of the Adriatic Sea. They are most common in winter on northern Adriatic coasts. The wind is probably a consequence of continental high pressure in central Europe with low pressure to the south in the Mediterranean. It is often accompanied by much precipitation when associated with *\*depressions* in the Adriatic. In other areas it is used as a generic term for cold squalls moving downhill from uplands.

**borax** *\*Evaporite* mineral,  $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$ ; sp. gr. 1.7; *\*hardness* 2.0–2.5; *\*monoclinic*; white, sometimes greyish or tinged blue; white *\*streak*; *\*vitreous* to *\*resinous* *\*lustre*; *\*cleavage* good {100}, {110}; occurs as a sediment of saline lakes, and in association with *\*halite*, *\*sulphates*, *\*carbonates*, and other borates such as *\*colemanite* ( $\text{Ca}_2\text{B}_6\text{O}_{11} \cdot 5\text{H}_2\text{O}$ ) and ulexite ( $\text{NaCaB}_5\text{O}_9 \cdot 8\text{H}_2\text{O}$ ). Soluble in water. It is a source mineral for boric compounds.

**bord and pillar** See *PILLAR AND STALL*.

**border fault** The *\*listric* *\*normal* fault that forms the *\*footwall* on one side of a half-graben (see *GRABEN*).

**bore** Very rapid rise of the tide, in which the advancing flood waters form a wave with an abrupt front. Bores occur in certain shallow *\*estuaries* and river mouths where there is a large tidal range, and suitably funnel-shaped regions, e.g. the Amazon, the Bay of Fundy, the Tsing Kiang River in China, and the Rivers Severn, Trent, and Ouse in England.

**boreal** Pertaining to the north (from Boreas, the Greek god of the north wind).

**Boreal** Period in post-glacial (i.e. post-*\*Devensian* or *\*Flandrian*) times from about 9500 to 7500 BP, which preceded the climatic optimum of early

**\*Atlantic** times. Pollen records typically show an increasing abundance of thermophilous (warmth-loving) tree species and also indicate the drier, more continental conditions that characterized the ensuing Atlantic period. The early Boreal corresponds to late **\*Pollen Zone V**; otherwise the Boreal is linked with **\*Pollen Zone VI**, which is sometimes subdivided to give Zones VIa, VIb, and VIc, according to the most abundant tree pollen represented. For Britain, the Boreal period is significant as the last period in post-glacial times in which Britain was joined to mainland Europe by a **\*land bridge** across the Dover Strait. *See also* **POLLEN ANALYSIS**.

**boreal climate** The climate associated with the boreal (taiga) forest zone of Eurasia, where it extends to 65°–70° N in the west and 50° N in the east, and with N. America, where it extends from the fringe of the tundra southwards to 55° N in the east. Winters are long and cold, with temperatures below 6 °C for 6–9 months, and summers are short, with temperatures averaging more than 10 °C. Precipitation, as snow in winter, typically amounts to 380–635 mm per annum.

**Boreal realm** The name used to denote that a fauna or flora has northern affinities. At certain times in the **\*Cenozoic** this implies a degree of coldness, while in the **\*Mesozoic** it is less specific, being used in comparative terms with the **\*Tethyan Realm**.

**borehole (well, dug well)** A hole drilled into rock, usually by **\*rotary** methods, to enable an assessment to be made of the characteristics of the rock itself and of contained fluids, e.g. **\*groundwater**, **\*natural gas**, or **\*petroleum**. The size may range from a few tens of millimetres in diameter to over 300 mm, and boreholes can be drilled at any angle to depths of several kilometres. Rock samples may be recovered from below the surface, the rock may be examined by down-hole geophysical or other methods, or **\*pumping tests** may be run. Boreholes may also be used as water, gas, or oil production wells. A completed borehole is frequently termed a **\*well**. *See* **BOREHOLE LOGGING; WELL LOGGING, GEOPHYSICAL**.

**borehole effect** The distortion on a **\*well log** resulting from changes in the diameter of the hole or in the thickness of the **\*invaded zone**.

**borehole logging** The recording of geophysical and geologic data recovered from a **\*borehole**. Geophysical borehole logs can be used in conjunction with drill cuttings, sidewall samples, and **\*cores** to establish the

geologic succession in the well. This information can then be interpreted geologically. The logs can be used to estimate formation *\*porosity*, *\*jointing*, fluid properties, hydrocarbon saturation, and formation pressures. Water wells are often logged by devices to detect water flows, levels of inflow, and such water properties as temperature and electrical conductivity.

**borehole sonde** See *SONDE*; *WELL LOGGING*.

**boring** (*noun*) A *\*trace fossil* that penetrates the hard surface of an organism or rock. The word may be used to describe the drill holes made by naticid gastropods (*\*Gastropoda*) into the shells of prey, or the short, tube-like extensions to larger borings made by the *\*siphons* of certain bivalves (*\*Bivalvia*), e.g. *Pholas*.

**bornhardt** Rounded, often isolated hill developed in massive rock and found in the humid tropics. Its shape is controlled by large-scale *\*exfoliation* or sheeting *\*joints*. Such hills are sometimes called ‘sugar-loaf’ hills, after the granitic dome of the Sugar Loaf, Rio de Janeiro, Brazil.

**bornite (peacock ore)** Common and important copper *\*ore mineral*,  $\text{Cu}_5\text{FeS}_4$ ; sp. gr. 5.0–5.1; *\*hardness* 3; *\*cubic*; reddish-brown to various shades of purplish-blue, the colours iridescent on tarnished surfaces; pale grey *\*streak*; *\*metallic lustre*; *\*crystals* rough cubes, rhombododecahedra, can also be *\*massive*; *\*cleavage* poor {111}; occurs in *\*hydrothermal veins*, in zones of secondary enrichment, and in various *\*syngenetic* copper ores, *\*porphyry copper* deposits, *\*skarns*, etc.; soluble in nitric acid.

**bort (bortz)** A compact variety of *\*diamond*, occurring naturally in granular *\*aggregates* and used as an abrasive in drill bits, saws, and gem-cutting wheels. It occurs in *\*ultrabasic*, *\*igneous*, *\*breccia* pipes, particularly in southern Africa.

**Bortonian** A *\*stage* (44–40 Ma ago) in the Lower *\*Tertiary* of New Zealand, underlain by the *\*Porangan*, overlain by the *\*Kaiatan*, and roughly contemporaneous with the upper *\*Lutetian* and lower *\*Bartonian* stages. (The name should not be confused with *\*Bartonian*.)

**bortz** See *BORT*.

**boss** Discordant *\*igneous* intrusion that has roughly vertical sides and a subcircular outcrop, and an area of less than 25 km<sup>2</sup>.

**Bothriocidaroida** (phylum *\*Echinodermata*, class *\*Echinoidea*) Small order of echinoderms which have five double columns of perforate plates and five columns of single imperforate plates. Traditionally they were regarded as the ancestral stock of all echinoids, but this is now considered unlikely. There are only three genera, *Bothriocidaris* (the earliest echinoid), *Neobothriocidaris*, and *Unibothriocidaris*, ranging in age from Middle *\*Ordovician* to Upper *\*Silurian*.

**Botomian** A Russian-Kazakhstanian *\*stage* (524–518.5 Ma ago) of the Early *\*Cambrian \*epoch*, preceded by the *\*Atdabanian* and followed by the *\*Toyonian*.

**botryoidal** In mineralogy, describes the form or shape of a *\*mineral* that occurs as spheroidal aggregations, often as a result of concentric growth patterns during its formation, e.g. *\*azurite*.

**bottleneck** A severe reduction in population size, often leading to a *\*founder effect*. Bottleneck events are commonly followed by rapid population expansion in which the rate of loss of new lineages is greatly reduced, giving rise to *\*star phylogenies*.

**bottomset beds** 1. The part of a cross-bedded set of *\*sediments* that forms at the base of the downcurrent or lee-side of a *\*dune-* or *\*ripple-*form structure. 2. Offshore clays formed at the base of a prograding deltaic sequence. *See also* DELTA; PROGRADATION.

**bottom water** The water mass that lies at the deepest part of the water column in the ocean. It is relatively dense and cold, e.g. the North Atlantic bottom water has a temperature of 1–2 °C. Bottom-water circulation is slow-moving, is greatly influenced by sea-bed topography, and is driven by differences in water density (*\*thermohaline circulation*).

**boudin** *See* BOUDINAGE.

**boudinage** Minor structural feature in which *\*competent \*strata* resemble a series of sausages ('boudins') that form by the stretching of the competent units which, unlike *\*incompetent* ones, cannot deform plastically. Initially, local thickening and thinning occurs ('pinch-and-swell' structures); as deformation proceeds, complete separation between boudins may take place.

**Bouguer, Pierre** (1698–1758) A French mathematician, Bouguer was a member of an expedition to Ecuador, sent to study the shape of the Earth. He made gravity measurements in the Andes, showing how the pull of gravity diminishes with altitude, and has given his name to corrections used in calculating gravitational anomalies. *See also* [BOUGUER ANOMALY](#); [BOUGUER CORRECTION](#); [GRAVITY ANOMALY](#).

**Bouguer anomaly** The gravitational attraction remaining after correcting the measured vertical component of [\\*gravitational acceleration](#) at a point for: (a) the theoretical gravity at that point, usually using the [\\*International Gravity Reference Field](#); (b) the [\\*free-air correction](#); (c) the [\\*Bouguer correction](#); and (d) the [\\*topographic](#) elevation correction, usually correcting to sea level. This anomaly is the fundamental gravity anomaly, reflecting all variations in density away from that expected for a homogeneous Earth.

**Bouguer correction** The correction applied to a measurement of [\\*gravitational acceleration](#), which allows for the attraction of rock between the observing station and some reference height, usually sea level. The correction is  $0.4185 \rho h$ , where  $\rho$  is the [\\*density](#) of the rock in  $\text{kg/m}^3$  and  $h$  is the difference in height between the two points in metres.

**Bouguer gravity map** A map of the gravitational field over an area after correction for theoretical gravity, [\\*free-air](#), solid rock ([\\*Bouguer correction](#)), and topography; i.e. a map showing [\\*Bouguer gravity anomalies](#).

**boulder** *See* [PARTICLE SIZE](#).

**boulder clay** Glacial deposit consisting of boulders of varying size in a clay-dominated matrix, and laid down beneath a valley glacier or ice sheet. Typically it is unstratified and unsorted, and characterized by rock types derived from the country crossed by the depositing glacier. *See also* [TILL](#).

**Bouma sequence** Idealized sequence of sedimentary structures observed in [\\*turbidity-current](#) deposits. It is named after the Dutch-American geologist Arnold H. Bouma (1932–2011), who first emphasized its generality (*Sedimentology of Some Flysch Deposits*, Elsevier, Amsterdam, 1962). The lowest unit, A ([\\*massive](#) or [\\*graded \\*sand](#)) is overlain progressively by the B (lower division of parallel [\\*lamination](#)), C (ripple or convolute laminations), D (upper division of plane parallel laminations), and E

(\*pelagic shale) units. Examples showing the entire sequence are not common. The sequence can be interpreted in terms of deposition under waning current conditions.

**bounce mark** A structure in a \*sedimentary rock that was caused by an object striking the surface of the sediment at a shallow angle and bouncing, leaving a depression that was later filled. *See also* PROD MARK; SKIP MARK; TOOL MARK.

**boundary current** Northward- or southward-directed ocean-water current which flows parallel and close to a continental margin. Such currents are caused by the deflection of eastward- and westward-flowing currents by the continental land masses. Boundary currents on the western margins of ocean basins, such as the \*Gulf Stream and the \*Kuroshio current, are deep, narrow, fast-moving currents carrying warm water; while currents along the eastern boundaries, such as the \*Canaries current and the \*California current, tend to be relatively shallow, broad, diffuse, slow-moving, and carry cool water.

**boundary layer** 1. At the interface between a solid surface and a fluid, a thin fluid layer that is static because of friction between molecules of the fluid and the solid. 2. *See* PLANETARY BOUNDARY LAYER.

**boundary stratotype** A specified rock section (*see* STRATOTYPE) within which the time-line ('golden spike', boundary zonal) occurs that marks the standard demarcation between \*chronostratigraphic units. In practice such time-lines are usually based on either the appearance or the disappearance of a key \*species (*see* INDEX FOSSIL) or other \*taxon. Associated \*faunas and \*sediments may transgress zonal boundaries. The term 'boundary stratotype' has also been used in the sense of the time-line itself.

**boundary wave** A seismic wave which travels along a boundary between two media of contrasting properties, rather than propagating through the bodies of the media (*compare* BODY WAVE). Where the boundary is a free surface (e.g. ground–air, sea bed–sea water) these waves are called \*surface waves and give rise to \*ground roll and \*mud roll respectively. Boundary waves are of two types: \*Love waves and \*Rayleigh waves.

**boundary zonal** *See* BOUNDARY STRATOTYPE.

**boundstone** A general term for *\*autochthonous* *\*carbonate* deposits in which the sediments are bound during deposition by organisms such as corals (*\*Anthozoa*) or *\*algae*. Boundstones are further subdivided into *\*bafflestone*, *\*bindstone*, and *\*framestone*. See also EMBRY AND CLOVAN CLASSIFICATION.

**ourn (bourne)** 1. A small stream, especially an *\*intermittent stream* in a chalk landscape. 2. The southern English version of *\*burn*.

**bourne** See BOURN.

**ournonite (cogwheel ore)** *\*Sulphide* mineral,  $\text{CuPbSbS}_3$ ; sp. gr. 6; *\*hardness* 3; grey; *\*sub-metallic* *\*lustre*; *\*massive* or rarely as cogwheel *\*aggregates*, in which the twinning habit (see TWIN LAW) causes *\*crystals* to form a rough cross with pitted, deeply striated edges; occurs in *\*hydrothermal* veins associated with copper and lead mineralization. The mineral was first identified in 1804 by the French mineralogist and crystallographer Count Jacques Louis de Bournon (1751–1825).

**ow-and-arrow rule** A rule stating that the best estimate of the direction of movement of a *\*thrust* is given by a line drawn at right angles to the *\*chord* joining the ends of the curving *\*fault* trace. The fault trace is approximately the shape of a bow, the chord joining its ends resembles the bowstring, and the line at right angles to the chord resembles the arrow.

**Bowen, Norman Levi** (1887–1956) A Canadian geochemist and experimental petrologist at the Carnegie Institution, Washington, DC, Bowen's classic work was on the chemistry of *\*igneous* rocks, published in *The Evolution of the Igneous Rocks* (1928). Later (1940) he turned to the study of *\*metamorphism* in *\*limestones* and *\*dolomites*. See BOWEN'S REACTION PRINCIPLE.

**Bowen's ratio** The ratio of sensible heat to latent heat transport from the ground to the atmosphere, which is generally calculated from the ratio of the vertical gradients of vapour pressure and temperature. It is used in the assessment of evaporation. It is named for the American physicist and astronomer Ira Sprague Bowen (1898–1973).

**Bowen's reaction principle** A concept, first propounded in 1928 by Norman *\*Bowen*, which explains how *\*minerals* can respond to the

changing equilibrium conditions when a *\*magma* is cooled, by either a continuous, diffusion-controlled exchange of elements with the magma or discontinuous melting of the mineral. In a *\*continuous* exchange or reaction, *\*solid-solution* minerals such as *\*feldspar* adjust their composition during cooling by a continuous diffusion of elements between magma and mineral, whilst in a *\*discontinuous* reaction, minerals such as *\*olivine* undergo melting at a specific temperature during cooling (the peritectic point) at the same time as a new mineral in equilibrium with the magma begins to crystallize (in this case *\*pyroxene*). Bowen suggested a series of these reactions that might take place during the cooling of a *\*tholeiitic* basalt magma, the so-called Bowen's reaction series, but pointed out that the series was a simplification of very complex reactions and could be misleading if taken at face value. The specific reaction series for tholeiitic magmas was never intended to become a general reaction series for all magmas.

**Bowen's reaction series** See BOWEN'S REACTION PRINCIPLE.

**bow shock** The shock wave in front of an object moving through a gas or liquid medium. The most familiar *\*solar-system* example is the shock-wave front between the magnetic field of a planet and the *\*solar* wind. It is the outer boundary of a planetary *\*magnetosphere*, where the solar wind is slowed from supersonic to subsonic velocities. It forms upwind of the *\*magnetopause* (the boundary where the planet is shielded from the solar wind).

**bow-tie reflection** A concave-upwards depression in a reflector with a curvature greater than that of an incident wave front is represented on a brute stack by a 'bow-tie' event. This results from there being three reflection points on the reflector for each surface location. A high degree of reflector curvature is required for each 'bow-tie' event to feature on normal incidence traces, with a reduced degree of curvature needed for a similar effect on offset traces. Consequently, 'bow-tie' events are more likely on long-offset traces and at greater depth within the *\*seismic section*. Curvature of the reflector out of the plane of the section can cause off-section 'bow-tie' *\*ghosts*.

**bow trend (barrel trend, symmetrical trend)** In the reading from a *\*wireline* log (see WELL LOGGING) an expression from a *\*gamma-ray* sonde

showing a gradual decrease, then increase in gamma radiation, indicating an increase followed by a decrease in clastic sedimentation within a basin.

**boxcar trend (cylindrical trend)** In the reading from a *\*wireline* log (see WELL LOGGING) an expression from a *\*gamma-ray* sonde showing no change and indicating a river channel bed, *\*turbidite*, or wind-blown sand.

**box classification** A remote sensing *\*classification* system in which a rectangular box is placed around *\*digital* number parameter space in a graph defining a certain *\*training* area (such as road). If the digital number values of the different *\*bands* plot to within that box then the pixel is assigned to that classification. See also MINIMUM-DISTANCE-TO-MEANS CLASSIFICATION; MAXIMUM-LIKELIHOOD CLASSIFICATION.

**box fold** A rectangular, *\*conjugate fold* in which the inter-limb angles approximate to right angles. The fold has two angular *\*hinges* and three limbs.

**BP** Initials which stand for 'before present' and relate to dates before the present day (taken conventionally to be 1950). The term should not be confused with 'bc', which relates to dates prior to the birth of Christ.

**brachia** (*sing.* **brachium**) **1.** In brachiopods (*\*Brachiopoda*), a pair of feathery structures forming part of the lophophore. The brachia are twisted up within the shell in various ways and serve to filter food particles from the water. **2.** In crinoids (*\*Crinoidea*), long, plated, flexible arms made up of a series of articulated *\*ossicles* containing a water vessel from which the *\*tube feet* arise. When feeding, the tube-feet are extended; at other times they are covered by 'brachial plates'. Small arms coming from the main brachia are called 'pinnules' and are made up of 'pinnular plates'.


**brachial plates** See BRACHIA.

**brachidium** The calcified support for the lophophore in brachiopods (*\*Brachiopoda*). It may be of different shapes in different groups (loops, spires, etc.). Some groups do not possess this calcified structure and the support for their lophophores is entirely hydrostatic.

**brachiole** The food-gathering arm of a blastoid or a cystoid (phylum *\*Echinodermata*, subphylum *\*Blastozoa*). The arms are *\*biserial* and are

simple versions of the arms that occur in crinoids (*\*Crinoidea*). They are rarely preserved.

**Brachiopoda (lampshells)** A phylum of solitary, benthic, marine, bivalved, *\*coelomate*, invertebrate animals that have existed from the Lower *\*Cambrian* to the present day. Brachiopods are commonly attached posteriorly to the sea bed by a stalk (*\*pedicle*), but may be secondarily cemented, or free-living (e.g. the fossil form *\*Productus* which, like many productids, was spinose, thick-shelled, and lived partly buried in the mud of the sea bed). Usually they consist of two unequal *\*valves*: a larger pedicle (ventral) valve and a brachial (dorsal) valve, lined by reduplications (*\*mantle* lobes) of the body wall which enclose the large mantle cavity. They are *\*bilaterally symmetrical* about the posterior–anterior mid-line of the valves (i.e. through the valves, *compare BIVALVIA*). The characteristic feeding and respiratory organ, the lophophore, surrounds the mouth and is covered by ciliated tentacles. It may be a simple horseshoe but more often forms two ciliated arms or brachia that project through the gape (thus giving the phylum its name). The alimentary canal is divided into oesophagus, stomach, and intestine, with or without an anus. The nervous system consists of a circum-oesophageal ring with a small aggregation of nerve cells on the ventral side. The excretory organs are one or two pairs of nephridia (excretory tubules) also acting as gonoducts (for the release of eggs and sperm). The circulatory system is open, with the contractile vesicle (heart) near the stomach. Considerably more than 3000 fossil species are known and about 330, in ten genera, are alive today; these are widely distributed and occur at all depths. Brachiopods are divided into three classes: *\*Lingulata*, *\*Inarticulata*, and *\*Articulata*.



## Brachiopoda

***Brachiosaurus*** See SAUROPODA.

**brachium** See BRACHIA.

**brachydont** Applied to teeth in which the crowns are low or short and the roots well developed, with narrow canals.

**bradyseism** A slow *\*earthquake*; i.e. gradual, differential motions of parts of the Earth's *\*crust* that do not suddenly release seismic energy.

**bradytelic evolution** One of three types of evolutionary rates within groups. In this instance the rate is exceedingly slow, and is manifest by slowly evolving lineages which survive much longer than would normally be expected. Living fossils, e.g. the *\*coelacanth*, represent the slow end of the bradytelic range; while the better-known groups that have remained relatively stable with time include opossums and crocodiles.

**Bragg, William Lawrence** (1890–1970) An English physicist who, with his father William Henry Bragg, evolved the technique of X-ray crystallography, used to determine the atomic structure of crystals. He was

an early supporter of the concept of *\*continental drift*, and is thought to have been responsible, in 1922, for arranging the translation into English and publication of *\*Wegener's work*. See also *BRAGG'S LAW*.

**Bragg equation** See *BRAGG'S LAW*.

**Bragg's law** In crystallography, the law that describes how an X-ray beam is reflected or diffracted in a crystal lattice, given by the Bragg equation  $n\lambda = 2d\sin\Theta$  where  $n$  is any integer,  $\lambda$  is the wavelength of the incident-beam X-ray,  $d$  is the spacing between crystal planes ( $d$  spacing), and  $\Theta$  is the angle between the crystal plane and the diffracted beam (the Bragg angle). Used in X-ray diffraction for the identification of minerals.

**braided stream (braided channel, braided river)** Stream whose plan form consists of a number of small channels separated by *\*bars*. The latter may be vegetated and stable (e.g. the eyots (small islets) in the River Thames, England), or barren and unstable (as at glacial margins, where rapid changes occur).

**branch** The graphical representation of an evolutionary relationship in a phylogenetic (see *PHYLOGENY*) tree.

**branchial arch** A single gill arch which supports a single pair of gill slits. Together the gill arches form the branchial basket which is part of the visceral skeleton.

**branchial basket** See *BRANCHIAL ARCH*.

**branching decay (dual decay)** The decay of an isotope by different methods to two or more different end-members. For example,  $^{40}\text{K}$  decays either to  $^{40}\text{Ar}$  (12%) by positron emission and *\*electron capture*, or to  $^{40}\text{Ca}$  (88%) by emission of a negative beta particle.

**Brandenburg** One of a series of *\*moraines* which mark the southern limit of *\*Weichselian* ice, extending some 500 km across the N. German Plain. A Russian equivalent extends a further 2000 km into European Russia.

**Brandon** An episode in the *\*Upton Warren Interstadial*, which occurred within the *\*Würm (\*Weichsel) Glacial*.

**braunerde** See *BROWN EARTH*.

**braunite** Oxide *\*mineral*  $Mn_2O_3$ , but with about 10% by weight of silica, so the mineral may be considered to have the composition  $3Mn_2O_3 \cdot MnSiO_3$ ; sp. gr. 4.7–4.8; *\*hardness* 6.0–6.5; *\*tetragonal*; brownish-black to steel-grey; brown-black *\*streak*; *\*sub-metallic* *\*lustre*; *\*crystals* pyramidal, can also occur *\*massive* or granular; occurs in *\*hydrothermal* veins together with other manganese oxides, as a *\*secondary* mineral, and as a result of the *\*metamorphism* of manganese-bearing sediments. It is very similar to other manganese minerals, but is soluble in hydrochloric acid, when it leaves a residue of silica. It is an ore of manganese and is used in steel-making.

**Bravais lattice** A lattice is a framework, resembling a three-dimensional, periodic array of points, on which a *\*crystal* is built. The smallest array which can be repeated is the ‘unit cell’. In 1850, M. A. Bravais showed that identical points can be arranged spatially to produce 14 types of regular pattern. These 14 space lattices are known as ‘Bravais lattices’.

**Bravais law** See BRAVAIS RULE.

**Bravais rule (Bravais law)** *\*Crystals* are formed by the repetition in three dimensions of a unit-cell structure defined by *\*lattice* points in space. The Bravais rule (proposed by M. A. Bravais) states that the density of these lattice points, or spacing between lattice planes, is proportional to the relative importance of crystal forms. It follows that the faces most likely to be found on a crystal are those parallel to lattice planes of greatest point density.


**Brazil current** Warm water, forming the oceanic *\*boundary* current that flows southward along the Brazilian coastal margin. It is marked by its high *\*salinity* (36.0–37.0‰), its low velocity, and its shallowness (100–200 m) in comparison to its northern hemisphere equivalent, the *\*Gulf Stream*.

**Brazilian CubeSat Project-1 (NanoSatC-Br1)** A Brazilian *\*CubeSat* mission that monitors the *\*magnetosphere* by measuring the *\*geomagnetic field* over Brazil and studies magnetic phenomena of the South Atlantic Anomaly and the Equatorial Electrojet. The 1-unit *\*CubeSat* was launched on 19 June 2004, from the Yasny Cosmodrome, Russia, into a Sun-synchronous orbit at an altitude of 630 km.

**bread-crust bomb** A ballistic mass of *\*lava*, usually greater than 64 mm in size, ejected from a source vent. The exterior of the lava rapidly chills to form a fine-grained, glassy crust, which is then cracked open by continued vesiculation (see *VESICLE*) and the expansion of the hot lava interior. The cracking of the crust resembles that sometimes seen on loaves of bread baked in the traditional manner.

**break** The onset of an event, in particular the *\*first* break. A sudden onset of new energy giving rise to a noticeably different amplitude event. A time break is the shot-instant time mark on a *\*seismic record*. See also *UPHOLE TIME*.

**break-back thrust** A *\*thrust \*fault* sequence in which the first fault increases in length until the amount of *\*stress* needed to move the volume of rock exceeds the breaking strength of the rock. The *\*hanging wall* then fractures and the thrust produces a newly active fault behind the first one (rather than ahead of the first one as in a *\*piggyback thrust sequence*). This sequence may be repeated several times.



**Break-back thrust**

**breaker** *\*Wave* that is collapsing or breaking as a result of the wave approaching the shore and reaching shallower water. The decreasing water depth causes the wavelength and speed to decrease and the wave height to increase. Consequently wave steepness increases and the wave becomes unstable: it breaks when the wave height is about 0.8 times the water depth. Several types of breaker have been described, e.g. spilling breakers (in which the wave breaks forward, with broken water spilling down the front of the wave), and plunging breakers (in which the wave crest curls over a large air pocket and falls vertically into the trough).

**breccia** Coarse, \*clastic, \*sedimentary rock, the constituent clasts of which are angular. 'Breccia' literally means 'rubble' and implies a rock deposited very close to the source area. The term may also be applied to angular volcanic rocks from a volcanic vent (vent breccia).

**breccio-conglomerate** A \*conglomerate with \*clasts which are both rounded and angular; it is intermediate in character between a \*breccia and a \*conglomerate.

**brecciola** From the Italian, literally meaning 'little \*breccia', normally-graded beds of small, angular, \*limestone \*clasts.

**breeze** Relatively light wind, often of convective origin, in the \*Beaufort scale blowing at 6.5 km/h (light breeze) to 50 km/h (strong breeze). The term also includes particular local air movements, e.g. mountain, land, and sea breezes.

**breviconic** Applied to a \*cephalopod conch when it is short and expands quickly. The \*aperture is quite wide.

**brick** Shaped block of baked \*clay used for construction. Usually obloid in shape with plane faces and parallel edges; fine uniform texture with no fissures, air bubbles, or pebbles; some are deep-salmon colour from the iron content of the clay. Small proportions of CaCO<sub>3</sub> and \*silt may be added for strength. The type of clay preferred has a low water content, to limit shrinkage on drying, and is high in organic matter to reduce fuel costs, e.g. the \*Jurassic Oxford Clay of England. Furnace, or refractory, bricks are made from \*kaolinite, etc.

**brickearth** Fine-grained silty deposit occurring in south-eastern England. Of complex origin, it probably resulted from the reworking of \*loess, either by hillslope washing or by redeposition in standing water.

**Bridgerian** A N. American \*stage of the \*Eocene (50.5–45.4 Ma ago), preceded by the \*Wasatchian and followed by the \*Heretaungan.

**Bridgman, Percy Williams** (1882–1961) A physicist who studied the effects of extreme high pressures on materials and their thermodynamic behaviour. He was awarded the Nobel Prize for Physics in 1946. Bridgman was born in Cambridge, Massachusetts, and spent his entire career at Harvard University, where he was appointed Hollis Professor of

Mathematics and Natural Philosophy in 1926 and Higgins University Professor in 1950.

**bridgmanite** A magnesium silicate mineral ( $\text{MgSiO}_3$ ) in an *\*orthorhombic* *\*perovskite* structure that, with *\*ferropericlase*, is the predominant material in the Earth's lower *\*mantle* and, therefore, the most abundant solid phase in the Earth's interior, accounting for 38% of the Earth's volume. Bridgmanite was discovered in the Tenham L6 *\*chondrite* and the name approved in 2014. It is named for Percy W. *\*Bridgman*.

**bridgmanite-enriched mantle structure (BEAMS)** *\*Mantle* material that forms because *\*bridgmanite* is two to three orders of magnitude more viscous than *\*ferropericlase*. BEAMS survive for long periods, forming reservoirs of material that does not mix with adjacent *\*mantle plumes* or subducting crust.

**Brigantian** The final *\*stage* (336–326.4 Ma ago) of the *\*Visean* epoch, followed by the *\*Serpukhovian*.

**bright spot** A characteristic phase reversal on a reflection event, usually indicating the presence of gas. The *\*phase* reversal is caused by a marked decrease in the *\*acoustic* impedance of gas-filled material which causes a strong negative reflection from the top of the gas-filled unit. Often this is accompanied by a *\*flat* spot below.

**Brillouin spectroscopy** A technique for measuring the elastic properties of single crystals in materials, providing data that allow the calculation of *\*seismic velocities* and changes in density that occur in materials under extreme pressure and/or temperature. It is based on the inelastic scattering of photons (light) by acoustic vibrations (phonons) induced by changes of temperature. This type of scattering was first described in 1922 by the French physicist Léon Nicolas Brillouin (1889–1969).



[http://serc.carleton.edu/NAGTWorkshops/mineralogy/mineral\\_physics/brillouin.html](http://serc.carleton.edu/NAGTWorkshops/mineralogy/mineral_physics/brillouin.html)

- Explains and describes Brillouin spectroscopy.

**brine** A concentrated solution of inorganic salts, formed by the partial evaporation of saline waters. *See also* [GEOTHERMAL BRINE](#).

**Brioverian** A \*stage of the Upper \*Proterozoic of Brittany, from about 626 to 575 Ma ago, underlain by the \*Pentevrian.

**bristlecone pine** See PINUS LONGAEEVA.

**British classification (of particle size)** See PARTICLE SIZE.

**British Geological Survey (BGS)** The organization responsible for advising the British government on all aspects of Earth science and for providing impartial geological advice to industry, academia, and the public. The BGS was founded in 1835, making it the oldest of all national geological surveys. It is a component part of the Natural Environment Research Council (NERC), and incorporates the former Overseas Geological Surveys and Mineral Resources Department of the Imperial Institute.



<http://www.bgs.ac.uk/>

- Home page of the BGS.

**brittle** See TENACITY.

**brittle behaviour** The manner in which \*competent rocks lose their internal cohesion along certain surfaces when the \*elastic limit is exceeded under an applied \*stress. Such behaviour gives rise to \*fractures (either \*faults or \*joints) and is common in upper crustal regions where temperatures and pressures are relatively low.

**Brøgger, Waldemar Christofer** (1851–1940) A Norwegian geologist and petrologist, Brøgger was a professor at the University of Oslo from 1890 to 1917. He had interests in many fields, including the geologic mapping of Norway and studies of \*pegmatites, but his main work was concerned with the theory of \*magmatic differentiation.

**Brongniart, Alexandre** (1770–1847) Brongniart was professor of mineralogy in the Paris Museum of Natural History, and collaborated with \*Cuvier in his work on the mapping of the Paris basin. He demonstrated that lower taxonomic orders of animals were found lower in the \*stratigraphic column, claiming this gave evidence of progression in the succession of life.

**Brontosaurus** See APATOSAURUS.

**bronzite** Old-fashioned name for an \*orthopyroxene (an \*orthorhombic \*pyroxene mineral) with a composition in the range  $(\text{Mg}_{1.4}\text{Fe}_{0.6})\text{Si}_2\text{O}_6$  to  $(\text{Mg}_{1.8}\text{Fe}_{0.2})\text{Si}_2\text{O}_6$ ; occurs in \*basic and \*ultrabasic \*igneous rocks and some rare \*metamorphic rocks.

**brookite** \*Mineral,  $\text{TiO}_2$ ; sp. gr. 4.1; \*hardness 5.5–6.0; \*orthorhombic; reddish-brown to brownish-black; white \*streak; \*metallic \*lustre; \*crystals normally \*platy and \*tabular; \*cleavage imperfect prismatic; occurs as an \*accessory mineral in \*igneous and \*metamorphic rocks, and in \*hydrothermal veins; Brookite, \*anatase, and \*rutile are all \*polymorphs of titanium dioxide. It is named after the British mineralogist H. J. Brooke.

**Brørup (Loopstedt)** An \*interstadial that occurred during the last (\*Weichselian) glacial. Named from a place in Jutland, it dates from about 60 000 years BP and is perhaps the equivalent of the \*Chelford Interstadial of the British Isles. Estimated mean July temperatures for this period are between 15 and 20 °C.

**brown clay** See RED CLAY.

**brown coal** Brown to brownish-black \*fossil fuel of low \*rank, intermediate between \*peat and \*lignite. It is soft, with a dull lustre and earthy texture, and contains the remains of plants; has a high moisture content (45–66%), and decomposes and darkens on dehydration. Its ash content is 1–5%, and its calorific value varies from low in soft, high-moisture types to high in those resembling \*lignites. It comes mainly from east and central Europe, where it is mined from \*open-cast pits.

**brown earth** Freely draining, and only slightly horizonated, \*soil-profile type. It has a \*mull \*humus in the surface \*horizon and very little differentiation of horizons below. Brown earths are well weathered and slightly \*leached soils, with a \*cambic horizon in the middle part of the profile (also known as braunerde and now included in the \*inceptisols).

**brown forest soil** Little-used \*soil-profile term that has been applied to both acid \*brown earths and \*brown podzolics.

**Brownian motion** Random movement of small particles dispersed in a colloidal solution or suspension.

**brown podzolic soil** Freely draining, \*leached \*soil profile that has developed acid surface \*horizons, a \*mor surface \*humus, and a clearly visible enrichment of translocated iron oxide in the middle, or B, horizon. The profile has been leached to the early stage of \*podzolization, identified by the movement down-profile of iron and aluminium compounds.

**brown soil** A major soil group in the classification of the Soil Survey of England and Wales, comprising well-drained soils with no gleyed features (see GLEY) within 40cm of the surface. These are good agricultural soils, divided into subgroups on the basis of their characteristics below 40 cm depth.

**brucite** \*Mineral,  $Mg(OH)_2$ ; sp. gr. 2.4; \*hardness 2.5; \*trigonal; white, shades of pale grey, blue, or green; white \*streak; pearly to waxy \*lustre; \*crystals normally broad and \*tabular, but also occurs fibrous, \*massive, and foliated; \*cleavage perfect basal {0001}; occurs in \*metamorphosed \*limestone, also in \*hydrothermal veins together with \*calcite, \*talc, and \*serpentine; soluble in hydrochloric acid. It is named after the 19th-century American mineralogist A. Bruce.

**Brückner cycle** Tendency for the cyclical recurrence of runs of wet years at about 35-year intervals, with warmer and drier years in between, reported in 1890 by A. Brückner (1862–1921). Much attention was paid to it for a time, but its operation is obscured by the greater magnitude of the irregular year-to-year variations, and by cyclical recurrences of other periods.

**Brunhes** The final normal \*polarity chron in the \*Quaternary, preceded by the Matuyama, and \*radiometrically dated from 0.78 Ma ago to the present. The Brunhes chron contains several brief \*reversed \*polarity subchrons of which the Blake (c.110 000 years BP), Lake Mungo (c.30 000 years BP), Laschamp (c.18 000 years BP), and Gothenburg (c.13 000 years BP) are the best, but still poorly, established.

**Bruun rule** The idea, proposed in 1962 by P. Bruun, that a marine shoreline maintains an equilibrium profile with a depth and slope determined by the current and wave regime.

**Bryophyta (bryophytes)** A division of plants, which includes only the mosses, with approximately 10 000 extant species. In earlier classifications the division also included the hornworts and liverworts, but that grouping is

now thought to be *\*paraphyletic*. Although bryophytes lack differentiated water-conducting vessels and rely largely or entirely on water absorbed from rain falling on the plants or from a moist atmosphere, they may have simple water-conducting cells in some larger species. They lack true roots, but possess root-like rhizoids which anchor them to a substrate and which can absorb water and minerals. The plants show an alternation of generations, with a green vegetative gametophyte (the familiar plant) and a sporophyte which typically takes the form of a (usually stalked) capsule and which is partially or wholly parasitic on the gametophyte. Bryophytes are found world-wide in a range of habitats. The earliest fossil moss is *Musites polytrichaceus* from the Upper *\*Carboniferous* of France. There is no evidence to link bryophytes with either the green algae (*\*Chlorophyta*) or the more advanced *\*pteridophytes*.

**Bryozoa (moss-animals)** Phylum of small, aquatic, colonial animals, related to the *\*Brachiopoda*; many colonies possess a well-developed, *\*calcite* skeleton which comprises microscopic, box-like divisions, each housing an individual animal possessing ciliated tentacles and a *\*coelom*. Food is collected by the tentacles which surround the mouth and are borne on a ridge called the 'lophophore'. Reproduction takes place by (a) asexual budding, and (b) the release of larvae which give rise to new colonies. There is a possible bryozoan fossil from the Late *\*Cambrian*, and bryozoans have occurred abundantly from the *\*Ordovician* to the present day. Branched colonies are common fossils in some rocks. They were important *\*reef* builders and binders in the *\*Phanerozoic*, and underwent several great radiations. See *ECTOPROCTA*.

**B-subduction** See *A-SUBDUCTION*.

**bubble pulse** When an explosion or *\*airgun* is triggered in water, the resulting gas bubble oscillates with decreasing energy with each oscillation generating a pressure pulse, known as the 'bubble pulse'. Unwanted bubble pulses can be prevented by setting off the source sufficiently close to the surface to allow the bubble to vent before it collapses, a method which, although spectacular, is inefficient. Where an *\*array* of tuned airguns is fired at depth, the individual airgun bubble pulses destructively interfere to provide the desired source signature. A knowledge of the bubble pulse period,  $T$ , is thus required for each source and this can be calculated using

the Rayleigh-Willis formula such that:  $T = (0.0452Q^{1/3})/(D/0.3048) + 33)^{5/6}$ , where  $Q$  is the source energy in joules and  $D$  is the water depth in metres to the bubble centre.

**bubnoff unit** A standard measure for describing the rates of geologic and geomorphologic erosional processes. One bubnoff unit (B) is equal to the removal of one micrometre per year, or one millimetre of surface material per thousand years. The unit is named after S. von Bubnoff (1888–1957).


**Buchan metamorphic zones** Areas in which a series of mineral assemblages occur, with a \*biotite zone surrounded by zones of \*cordierite, \*andalusite, and \*sillimanite, in increasing order of metamorphic grade. The zones are named after the region of north-east Scotland (north of Aberdeen) in which they occur. Compare BARROW'S ZONES.

**Buchan spells** Several periods of the year, 'more or less well defined', when the normal seasonal rise or fall of temperature is halted or reversed for a time, e.g. 9–14 May is a cold period and 3–14 December a warm period. They are named after the Scottish meteorologist Alexander Buchan (1829–1907), whose analysis of temperature records for Scottish stations in the middle of the 19th century revealed them. Six cold and three warm periods of this type were identified by Buchan in 1869 from his examination of records covering several years. See also SINGULARITY.

**Buckland, William** (1784–1856) The first reader in Geology and Mineralogy at the University of Oxford, Buckland was appointed Dean of Westminster in 1846. He developed the English school of historical geology with its emphasis on the progressive (but not evolutionary) nature of Earth history. Initially, he was a diluvialist and his *Reliquiae Diluvianae* (1823) is based on studies of fossil deposits in Kirkdale Cavern and other caves. He subsequently abandoned \*diluvialism in favour of Agassiz's \*glacial theory which he was the first to introduce into British geology. *Buckland's* Bridgewater Treatise (1836) was not only an attempt to reconcile geology and \*natural theology but also an up-to-date manual of historical geology. See also CATASTROPHISM.

**buckle folding** The sideways deflection from a median line of \*competent layers in a less competent matrix, due to mechanical 'buckling' instabilities set up by \*layer-parallel shortening. A regular dominant wavelength is

created as a function of layer thickness, layer *\*rheology*, and contrasts in competence.



**Buckle folding**

**Buffon, Georges Louis Leclerc, Comte de** (1707–88) Buffon was an extremely influential French natural philosopher whose extensive works included one of the last speculative cosmogonies: *Les Époques de la Nature* (1778). Beginning with the formation of the Earth, it covered seven epochs in analogy to the seven days of Creation. He proposed that the Earth was hot at its creation and, from the rate of cooling, calculated its age to be more than 100 000 years.

**Buganda-Toro-Kibalian orogeny** A phase of mountain building that occurred about 2075–1700 Ma ago, affecting what is now an area of Central and East Africa stretching from northern Zambia to Katanga Province in the Democratic Republic of Congo, and the borders of Uganda. Thrusting in a north-westward direction produced a belt now trending SW–NE and forming the north-eastern section of the Kibaride belt.

**bulb of pressure** Bulb-shaped lines of equal vertical stress below a footing or foundation.

**Bulitian** A *\*stage* (55.8–53 Ma ago) in the Lower *\*Tertiary* of the west coast of N. America, underlain by the *\*Ynezian*, overlain by the *\*Penutian*, and roughly contemporaneous with the upper *\*Thanetian* stage.

**bulk composition of Earth (whole Earth composition)** Information on the composition of the Earth as a whole has been deduced from: (a) cosmochemical models, which assume that the compositions of all members of the *\*solar* system are related, and that the bulk composition of the Earth can be inferred from the abundances of non-volatile elements in the Sun and some primitive *\*meteorites* (see [CHONDRITIC EARTH MODEL](#)); and (b) geophysical evidence, e.g. *\*seismic* data, *\*density determinations*, and *\*magnetic surveys*. The three layers, *\*core*, *\*mantle*, and *\*crust*, of which the *\*Earth* is formed differ markedly in their composition. The core and mantle make up more than 99% of the Earth's mass but their compositions can be only inferred, unlike that of the crust (see [CRUSTAL ABUNDANCE OF ELEMENTS](#)). The density and *\*magnetic field* of the Earth, information from seismic surveys, and the existence of iron–nickel meteorites lead to the conclusion that the core is predominantly iron, with a small proportion of lower-density element(s). Nickel is largely excluded because it would make the core too dense. The nature of the light component is controversial, but may be sulphur, carbon, oxygen, silicon, and potassium. From seismic evidence the mantle appears to be composed of *\*dunite*, *\*peridotite*, and *\*eclogite*, rocks similar in composition to *\*chondritic* meteorites. The upper mantle is probably formed from dense *\*silicates* of iron and magnesium, with silicon and magnesium oxides becoming commoner with depth. Information regarding the bulk composition of the Earth is fundamental to resolving such questions as the relationship between the Earth and the *\*Moon*. Perhaps the most critical single feature of the Earth's bulk composition is its content of K, U, and Th, since the radioactive isotopes of these elements control radioactive heat production, and therefore the thermal and geologic history of the Earth. See also [COSMIC ABUNDANCE OF ELEMENTS](#); [METEORITIC ABUNDANCE OF ELEMENTS](#); [SOLAR ABUNDANCE OF ELEMENTS](#).

**bulk density** Mass per unit volume of soil (sampled as a clod or core), dried to constant weight at 105 °C.

**bulking** **1.** Increase in volume of material when it absorbs water; dry *\*sand* or *\*clay* may swell as much as 50%. **2.** Increase in volume of solid rock or soil when broken.

**bulk minerals** See [AGGREGATE](#).

**bulk modulus (incompressibility modulus;  $K$ )** The stress:strain ratio. When a simple hydrostatic pressure  $p$  is applied to a cube the resulting volume strain is equal to the ratio of the change in volume  $\delta V$  to the original volume  $V$ . Thus:  $K = (\text{volume stress } p)/(\text{volume strain } \delta V/V)$ .

**Bullard, Edward Crisp** (1907–80) A geophysicist of the University of Cambridge, England, Bullard worked on many of the geophysical techniques used to gather data in support of *\*plate tectonic* theory, including studies of gravity, heat flow, and *\*palaeomagnetism*. He was an early supporter of plate tectonic theory, publishing in 1964 a computer-generated map showing the matching of *\*continental shelves* on either side of the Atlantic.

**bumpiness** Unevenness experienced in the flight of an aircraft caused by convection or turbulence in conditions of atmospheric instability or over orographic barriers. *See also* CLEAR-AIR TURBULENCE.

**Buntsandstein** *See* CHANGHSINGIAN.

**buoyancy** Condition arising from the difference between the density of a given *\*parcel* of fluid and that of the surrounding fluid. It is this that makes a hydrogen- or helium-filled balloon or airship float or rise through the atmosphere. If an air parcel is warmer (therefore less dense) than its surroundings, the density difference implies an upward-directed force acting upon that parcel of air and it will rise, with positive buoyancy. When a parcel of air is colder (denser) than its surroundings the reverse condition causes it to sink, with downward (negative) buoyancy. If the fluid density is equal to that of the surrounding medium the fluid will experience neutral buoyancy and remain at the same level. Buoyancy per unit mass of a fluid is given by  $(\rho' - \rho)/\rho$ , where  $\rho'$  is the density of the surrounding medium and  $\rho$  is the density of the parcel of fluid. *See* THERMAL.

**buran** Regional wind in Russia and central Asia which blows strongly from the north-east in both winter and summer. In winter it is very cold and is associated with much snow. *See also* PURGA.

**Burdigalian** 1. An *\*age* in the Early *\*Miocene*, preceded by the *\*Aquitanian* and followed by the *\*Langhian*. 2. The name of the corresponding European *\*stage*, dated radiometrically as about 20.43–15.97 Ma ago, which is roughly contemporaneous with the upper *\*Saucesian* (N.

America), upper *\*Otaian* and *\*Altonian* (New Zealand), and the upper *\*Longfordian* and *\*Batesfordian* (Australia). The type section is at Le Coquillat, France. Mammals, including elephants, are important in the stratigraphic definition of this stage.

**Burgers model** A model of the *\*stress–strain–time* behaviour of compressible *\*viscoelastic* materials, e.g. soil, *\*asphalt*, *\*mantle* rock, that combines the four *\*Maxwell* and *\*Kelvin–Voight* coefficients by connecting them in series. It is named for the Dutch physicist Jan Martinus Burgers (1895–1981). *Compare* *KELVIN–VOIGHT MODEL*; *MAXWELL MODEL*.

**Burgess Shale** A *\*horizon* from the *\*Cambrian* of British Columbia which has yielded an exceptionally preserved fauna. Originally discovered by C. D. Walcott in 1909, it has been redescribed by H. B. Whittington and other authors (1967–8). Apparently the fauna was deposited in deep water on or near a submarine fan. *\*Arthropods* of various types account for more than 30% of the fauna, but other groups are also represented, many of them bizarre. It is probable that in terms of its diversity (140 species in 119 genera) this was a typical Cambrian fauna, but it is unusual in that the soft-bodied forms are preserved.

**burial metamorphism** A term first used by D. S. Coombs in 1961 to describe metamorphic *\*recrystallization* during *\*epeirogenic* as opposed to *\*orogenic* earth movements. Sediments and volcanic rocks in a developing *\*basin* gradually become buried during sagging of the crust in response to the weight of the accumulating rock column above, so that temperatures, even at great depth, are much lower than those experienced during plate *\*collision*, when forcible depression of the rock mass to regions of much higher temperature and pressure cause metamorphic changes characteristic of *\*regional metamorphism*. *See* *METAMORPHIC ROCK*.

**buried soil** Soil covered by an *\*alluvial*, *\*colluvial*, *\*aeolian*, glacial, or organic deposit, and that is a product of a former period of *\*pedogenesis*. In US usage a buried soil is defined as lying beneath 300–500 mm if the covering layer is more than half the thickness of the buried soil, otherwise beneath more than 500 mm.

**buried topography** An *\*overstepped* erosion surface (*\*unconformity*) which shows topographic relief of hills and valleys. Such relief has a strong effect on the thickness and *\*attitudes* of the overstepping strata.

**burn (bourn, bourne)** A small stream, from the Old English *burna*.

**burner reactor** Type of nuclear reactor using  $^{235}\text{U}$  as fuel. *\*Enriched* uranium is used to increase efficiency. To prolong the *\*fission* reaction, fast neutrons are slowed down with a moderator (graphite or heavy water) and the rate of reaction is adjusted by control rods (boron) which can absorb neutrons. Some neutrons react to form plutonium, but in smaller amounts than the original uranium, hence the term burner reactor. *Compare* FAST BREEDER REACTOR.


**Burnet, Thomas** (1635?–1715) A natural philosopher whose *Sacred Theory of the Earth* (1680, 1689) was an early diluvialist work. He tried to correlate the seven days of creation with Earth history, describing the Earth as a giant shell from which flood waters gushed when it was broken by God in the Deluge. The broken fragments of the crust formed mountains. *See* DILUVIALISM.

**burrow** A *\*trace fossil* formed by an animal during feeding, migration, or in the creation of a resting place. Burrows are formed in soft *\*sediments* and may occur on the surface or be the result of subsurface activities.

**‘burst of monsoon’** Applied to the onset of a marked change in weather conditions in the Indian subcontinent and south-east Asia, associated with the arrival of humid south-westerly winds which displace the hot, dry, pre-monsoon regime. The changed surface-level wind pattern is related to the establishment of a high-level easterly *\*jet stream*.

**Burzyan** A *\*system* of the Middle *\*Proterozoic*, extending from about 1400 Ma ago to about 1375 Ma ago (Int. Commission on Stratigraphy, 2004), of western Russian origin.

**butte** Small, isolated, flat-topped hill resulting from the erosion of near-horizontal strata. The diameter of the *\*cap* rock is less than the height of the land-form above the surrounding country. Buttes are commonly found in semi-arid regions dominated by *\*duricrust* horizons. *See also* MESA.



**Butte**

**Byes Ballot's law** Law, enunciated in 1857 by the Dutch meteorologist Christoph Hendrick Diderik Buys Ballot (1817–90), that in the northern hemisphere the winds blow anticlockwise around centres of low pressure and clockwise around centres of high pressure. In the southern hemisphere both these directions are reversed.

**Byerlee's rule (Byerlee's law)** A rule, derived experimentally in 1978 by the American geophysicist James Byerlee, that determines the relation between *\*shear stress* ( $\tau$ ) and *\*normal stress* ( $\sigma$ ) at which rocks will *\*fracture* at a *\*fault*. Byerlee found that  $\tau = 0.85\sigma$  for  $\sigma$  up to 200 MPa and  $\tau = 50 + 0.6\sigma$  for  $\sigma$  above 200 MPa.

**by-product** In mining, subsidiary material worked from *\*ore* deposits in which other materials are dominant. In some cases the revenue from the by-products can exceed that from the major type, e.g. gold from *\*porphyry* copper deposits. *Compare* CO-PRODUCT.


**byssate** Applied to the condition found in some bivalves of being attached to the substrate or to objects by strands of byssus (usually tough, horny threads).

**byssus** *See* BYSSATE.

**byte** A group of 8 *\*bits* of digital data in binary form. A byte can represent values of 0–225.

**bytownite** *See* PLAGIOCLASE FELDSPAR.

**b-zone** A biostratigraphic *\*zone* distinguished by *\*benthonic \*fossils*, e.g. *\*brachiopods* and *\*trilobites*. The term was proposed by T. G. Miller in 1965. *Compare P-ZONE.*



**<sup>14</sup>C** See RADIOMETRIC DATING.

**cable drilling** Drill in which a weight is repeatedly lifted and dropped to create a hole. Only shallow holes can be made in this manner. *See also* ROTARY DRILLING.

**cadicone** See INVOLUTE.

**CADRE** A 3-unit \*CubeSat space-weather mission of the University of Michigan, Ann Arbor, \*NASA, and the Naval Research Laboratory that studies the properties of the \*thermosphere and \*ionosphere. It was launched on 6 December 2016 to the International Space Station and deployed from there on 16 May.

**Caerfai epoch** See EARLY CAMBRIAN EPOCH.

**Cainozoic** See CENOZOIC.

**cairngorm** See QUARTZ.

**cake** See MUD-CAKE.

**Calabrian** Early \*Pleistocene \*stage (1.806–0.781 Ma ago), preceded by the Gelasian and followed by the Middle Pleistocene, that is noted for major evolutionary changes in mammalian faunas. *See also* CASTLECLIFFIAN; QUATERNARY.

**calamine** See SMITHSONITE.

**Calamites cistiiformes** Jointed-stemmed calamitids were an important component of \*Carboniferous swampland floras. The generic name *Calamites* was first given to the ridged and furrowed casts of the pith

cavity, which are commonly encountered fossils (see [FORM-GENUS](#)). Unlike their smaller modern counterparts (*\*Equisetum* species) some Carboniferous species grew to 18 m tall. *Calamites cistiiformes* was the first representative of the family Calamitaceae. It was first described by Stur in 1877. See also [SPHENOPSISIDA](#).

**calc-alkaline** The name given to a suite of rocks comprising the volcanic association *\*basalt–\*andesite–\*dacite–\*rhyolite*, or the *\*plutonic* association *\*gabbro–\*diorite–\*granodiorite–\*granite*. The suite is defined chemically using a graph depicting the variation of CaO and (Na<sub>2</sub>O + K<sub>2</sub>O) with SiO<sub>2</sub> (wt. %), and is classified as calc-alkaline if the SiO<sub>2</sub> value at the intersection of the CaO and (Na<sub>2</sub>O + K<sub>2</sub>O) trends is between 56 and 61 SiO<sub>2</sub> wt. %. (Less than 51 SiO<sub>2</sub> wt. % and the rock suite is classed as belonging to the alkalic series, 51–56 SiO<sub>2</sub> wt. % to the alkali-calcic series, and more than 61 SiO<sub>2</sub> wt. % to the calcic series.) Calc-alkaline rocks are typically developed on the continental side of *\*plate \*subduction zones*, well-known examples occurring in the Andes of S. America and in Japan.

**Calcarea (Calcispongea)** (phylum *\*Porifera*) A class of sponge, ranging from *\*Cambrian* to Recent, in which the skeleton is made entirely of calcareous *\*spicules* which are commonly of a tuning-fork shape. The sponges are sometimes associated with *\*reefs* in the *\*Jurassic*, or they may form widespread sponge beds.

**calcarenite** Calcareous sediment in which a high percentage of the *\*clasts* can be of *\*quartz* within a calcareous *\*matrix* or, more generally, a *\*clastic \*limestone* in which both clasts and matrix are calcareous. Calcarenites are characterized by a particle range of 0.006–2.00 mm in diameter.

**calcareous ooze** Deep-sea, fine-grained, *\*pelagic* deposit containing more than 30% calcium carbonate. The calcium carbonate is derived from the skeletal material of various *\*planktonic* animals and plants, e.g. *\*foraminiferan \*tests* and *\*coccoliths* (which are calcitic), and pteropod tests (which are aragonitic). Calcareous ooze is the most extensive deposit on the ocean floor but is restricted to water depths less than about 3500 m. See also [CARBONATE COMPENSATION DEPTH](#).

**calcareous soil** Soil that contains enough free calcium carbonate to effervesce visibly, releasing carbon-dioxide gas, when treated with cold 0.1 N hydrochloric acid, and which could also be regarded as a **\*basic** or **\*alkaline soil**.

**calceolid** See SOLITARY CORALS.

**Calcichordata** See HOMALOOZOA.

**calcichordates** A group of early **\*Palaeozoic** animals with calcareous **\*exoskeletons**, interpreted by some as **\*carpoid** echinoderms (**\*Echinodermata**), and by others as ancestors of the **\*Chordata**.

**calcic horizon** Mineral **\*soil horizon** with evidence of secondary calcium-carbonate deposition which is more than 150 mm thick, with a calcium-carbonate content of more than 15% by weight, and with 5% more calcium carbonate than is in the parent material or horizons below it.

**calcic series** See CALC-ALKALINE.

**calcification** Process of redeposition of secondary calcium carbonate from other parts of the **\*soil profile** which, if sufficiently concentrated, may develop into a calcrete, **\*caliche**, kunkar, or **\*calcic** horizon (all similar and usually comprising more than 15% by weight of calcium carbonate in more than 150mm soil thickness). Calcification involves limited upward and lateral movement of calcium salts in solution, and downward movement in less-wet periods. When occasionally **\*leaching** is deeper, some but not all of it may redissolve.

**calcilutite** A fine-grained **\*limestone** consisting of **\*silt** and clay-sized **\*carbonate** particles (less than 63 $\mu$ m in **\*grain** size).

**calcirudite** A coarse-grained **\*limestone** consisting of limestone **\*clasts**, more than 2 mm in diameter, set in a **\*carbonate** **\*matrix**. The clasts may be angular or rounded.

**calcisiltite** A fine-grained **\*limestone** consisting of **\*silt-sized** (but not **\*clay-sized**) **\*carbonate** particles (more than 2 but less than 63  $\mu$ m sizes).

**calcisols** A reference soil group in the **\*World Reference Base for Soil Resources** classification scheme. Calcisols have a **\*calcic** horizon within 125 cm of the surface.

**calcispheres** Small *\*calcite* spheres, up to 500 µm in diameter, commonly found in *\*Palaeozoic \*limestones* and believed to be of algal origin. They consist of a *\*micrite* wall enclosing an interior which is hollow or filled with sparry-calcite (*\*sparite*).

**Calcispongea** See CALCAREA.

**calcite** Very common, widespread, rock-forming carbonate *\*mineral*, one of two *\*polymorphs* of CaCO<sub>3</sub>, the other being *\*aragonite*; sp. gr. 2.7; *\*hardness* 3; *\*trigonal*; usually colourless or white, but may be shades of yellow, grey, green, red, or even brown or black; white *\*streak*; *\*vitreous \*lustre*; *\*crystals* common, often *\*tabular*, *\*prismatic*, or rhombohedral, but fibrous aggregates and granular masses may also occur; *\*cleavage* perfect rhombohedral {10 $\bar{1}$ 1}; cleavage rhombs exhibit *\*double refraction*; a major constituent of calcareous *\*sedimentary rocks*, e.g. *\*marbles*. Calcite can be precipitated from sea water and is a common constituent of invertebrate shells, and late-stage hydrothermal solutions (see HYDROTHERMAL ACTIVITY) may precipitate calcite in cavities in some *\*igneous* rocks such as *\*basalts*. It is soluble in dilute hydrochloric acid. It is used as a flux, in cement-making and fertilizers, and as a building stone. See also CARBONATES.

**calcium feldspar** See ALKALI FELDSPAR.

**calcrete** See CALICHE.

**calcrete uranium** In arid climates prolonged evaporation may cause hard surface *\*crusts* to form; if CaCO<sub>3</sub> predominates the crust is called *\*caliche* (calcrete). Where the bedrock contains high levels of uranium the calcrete is locally uraniferous and may constitute a workable *\*ore*, e.g. in western Australia and Namibia.

**calcsilicate** A group of *\*minerals* whose bulk composition consists of calcium silicates. Common calcsilicates include the minerals *\*wollastonite* (CaSiO<sub>3</sub>), the calcium *\*garnet* grossularite, and the *\*pyroxene \*diopside* (CaMgSi<sub>2</sub>O<sub>6</sub>). Calcsilicates are commonly formed by the *\*metamorphism* of *\*limestones* and *\*dolomites*.

**calc-sinter** See TRAVERTINE.

**caldera** A roughly circular topographic and structural depression, on Earth varying in diameter from about 1 to 100 km and averaging 68 km on **\*Venus**, 40 km on **\*Io**, and 48 km on **\*Mars**, formed by the foundering and collapse of a **\*magma chamber** roof into its underlying magma body (e.g. Crater Lake, Oregon, formed by the eruption of Mt Mazama about 6000 years ago). Caldera is Spanish for cauldron or cooking pot. Caldera collapse is commonly preceded or accompanied by rapid explosive evacuation of magma from the chamber in the form of surface **\*pyroclastic flows**. This leaves the chamber roof unsupported by magma pressure and collapse follows. Slumping and erosion of the caldera walls may enlarge the topographic rim of the depression well beyond the structural rim. Later injection of magma into the chamber can cause doming of the caldera floor to create a resurgent caldera. Tobu caldera, Sumatra, almost 100 km in longest dimension, is the largest terrestrial caldera.

**Caledonian orogeny** Major mountain-building episode which took place during the Lower **\*Palaeozoic** era. The orogeny affected Greenland, Ireland, Scotland, and Scandinavia, and was associated with the closure of the **\*Iapetus** Ocean between the old continents of **\*Laurentia**, **\*Gondwana**, and **\*Baltica**.

**Caliban (Uranus XVI)** A lesser satellite of **\*Uranus** with a radius of 36 km and a visual **\*albedo** of 0.07.

**calibration graph** A plot of the line intensities (relative to the **\*internal standard**) of a set of samples with known intensities, used in the calculation of the concentration of elements (e.g. in emission spectrometry) from the line intensity of the sample being examined.

**caliche (calcrete)** **\*Carbonate** **\*horizon** (the K horizon) formed in a soil in a semi-arid region, under conditions of sparse rainfall (20–60 mm/yr) and a mean annual temperature of about 18 °C, normally by the precipitation of calcium carbonate carried in solution. The **\*soil profile** develops over several thousand years, initially in the form of nodules (glaebules), more mature caliches taking a massive, laminar form. It may become **\*cemented** and **\*indurated** on exposure, when it gives rise to a tabular landscape. See also **CALCIC**; **DURICRUST**; **DURIPAN**; **PETROCALCIC**.

**calichnia** A type of **\*trace fossil** of a structure that was made for breeding purposes.

**California bearing ratio (CBR)** 1. Strength tests of *\*subgrades* and construction material; also used in the design of flexible *\*pavements* to meet specifications. 2. Measure of soil resistance to penetration under controlled conditions. See *PENETRATION TESTS*.

**California current** Southward-flowing, eastern *\*boundary* current carrying cool water from the North Pacific current to join the North *\*Equatorial* current. This slow-moving, diffuse water mass flows down the west coast of N. America.

**caliper log** The record produced by a spring-loaded caliper that continually adjusts itself to the size of a *\*borehole* as it is pulled to the surface. The log is a direct measure of the diameter of the borehole as a function of depth.

**CALIPSO** See *CLOUD-AEROSOL LIDAR AND INFRARED PATHFINDER SATELLITE OBSERVATIONS*.

**Callisto (Jupiter IV)** The third largest of the *\*Galilean* satellites, with the lowest density, the body is believed not to have differentiated into a core and mantle, consisting throughout of a mixture of rock and ice. Its surface is the darkest of the Galileans (albedo 0.20; though twice as bright as that of the *\*Moon*) and Callisto is the most heavily cratered body in the solar system. There is believed to be an almost complete absence of geologic activity at the surface, which has an age of about 4 billion years and shows no sign of any extensive resurfacing. The surface is of dirty ice and there are no large mountains. Surface craters and rings are shallow; the largest structures are Valhalla, a bright patch about 600 × 3000 km, and Asgard, a ring about 1600 km across. The surface temperature is about 134 K. Callisto was discovered on 7 January 1610 by Galileo. Its diameter is 4820 km; mass  $1.076 \times 10^{23}$  kg; density 1.8344 kg/m<sup>3</sup>; surface gravity 0.126 g (Earth = 1); mean distance from Jupiter  $1.883 \times 10^6$  km; mean distance from Sun 5.203 AU; orbital period 16.68902 days; rotational period 16.68902 days.

**Callovian** A *\*stage* in the European Middle *\*Jurassic* dated at 164.7–161.2 Ma ago (Int. Commission on Stratigraphy, 2004), preceded by the *\*Bathonian*, followed by the *\*Oxfordian*. See also *DOGGER*.

**calm** Condition of general lack of wind, indicated by a wind speed of less than 1 knot (0.5 m/s). *See also* [BEAUFORT SCALE](#).

**Calorian** The third and most recent period of the *\*Mercurial* geologic timescale, lasting from 3850 Ma ago to 3000 Ma ago and comprising the Early (3850–3800 Ma ago) and Late (3800–3000 Ma ago) Calorian epochs. The period is named for the Caloris Basin, and during it the *\*late heavy bombardment* ended.

**calving** Process whereby portions of a glacier's leading edge break off as icebergs into an adjacent body of water.

**calvus** From the Latin *calvus* meaning 'bald' or 'stripped'; a species of *\*cumulonimbus* cloud in which upper protrusions form a more amorphous mass than appears from the cumuliform outlines. *See also* [CLOUD CLASSIFICATION](#).

**calyce** The skeletal surface of a coral that in life was in contact with the basal ectoderm of the polyp. It can be quite variable in shape and dimensions, but is commonly saucer- or cup-shaped.

**Calymmian** A period of the *\*Proterozoic* eon that began 1600 Ma ago and ended 1400 Ma ago. Its limits are set by fixed dates. The Calymmian followed the *\*Statherian* period and was succeeded by the *\*Ectasian* period.

**Calypso (Saturn XIV)** One of the lesser satellites of *\*Saturn*, discovered in 1980 by *\*Voyager 1*, with a radius measuring  $15 \times 8 \times 8$  km; visual albedo 0.6.

**Calyptomatida** (phylum *\*Mollusca*) Extinct *\*Palaeozoic* class of marine molluscs, with *\*bilateral* symmetry, in which the shell shape is variable: most are subtrigonal in cross-section. The walls are generally thick, and composed of laminated calcium carbonate. The apertural end is the widest; the narrowest is the closed apex, which may be pointed or blunt. The juvenile portion is chambered with *\*imperforate* *\*septa*. Externally the shells are smooth, or ornamented with growth lines and/or longitudinal and transverse ridges. The *\*aperture* is usually protected by an *\*operculum* possessing paired muscle scars. Individuals are 1–150 mm long. The class ranges in age from Lower *\*Cambrian* to Middle *\*Permian*, and contains two orders.

**calyx** 1. Cup-shaped, plated body of *\*pelmatozoan* *\*echinoderms*. The calyx is made up of several rows of plates. If there is no stem, a centrodorsal plate is succeeded by a ring of five basal plates and then by a further ring of radial plates above them. Above these two rows of plates come the plates making up the arms. Where these two rows of plates are present the calyx is said to be 'monocyclic'. In some species a third row of plates, the infrabasals, is present beneath the basals and the calyx is then said to be 'dicyclic'. In some cases radial plates may be compounded (i.e. split transversely into two plates) and then, depending on their position in the calyx, they may be known as 'infraradial' or 'supraradial plates'. 2. Bowl-shaped depression at the top of a calcareous coral skeleton, usually formed by the upper edges of the *\*septa*.

**calyx drilling** Method of *\*core* drilling performed by the rotation of a steel cylinder and cutting with chilled shot, about 2.4 mm diameter, which cuts a formation core. Circulating water carries the cuttings up to a basket-like space at the top of the *\*core barrel*. The core is wedged into the barrel and pulled up one barrel-length at a time. Shafts up to 2 m diameter can be drilled, and holes to a depth of more than 300 m.

**camara** See CAMERA.

**camaral** See CAMERA.

**cambering** Consistent *\*dip* of strata towards local valley floors, in conflict with the general regional dip. It is well displayed in the English Midlands, where *\*ironstone* beds above clays are cambered down as much as 30 m below their original level. It is probably due to large-scale structural disturbance when *\*permafrost* thawed and when the plastic clays allowed overlying massive beds to flow towards valleys.

**cambic horizon** Weakly developed mineral *\*soil horizon* of the middle part (B horizon) of *\*soil profiles*, and one that has few distinguishing morphological characteristics except for evidence of *\*weathering* and sometimes of gleying. It is found in *\*brown earths* and *\*gleys*.

**cambisols** A reference soil group in the *\*World Reference Base for Soil Resources* classification scheme. Cambisols have a *\*cambic* horizon or *\*mollic* horizon above a subsoil with a *\*base saturation* lower than 50% in

the uppermost 100 cm, or an *\*andic horizon*, *\*vertic horizon*, or *\*vitric horizon* commencing 25–100 cm below the surface.

**Cambrian** The first of six periods of the *\*Palaeozoic* era, which began about 542 Ma ago and ended about 488.3 Ma ago. Sediments deposited during the period include the first organisms with mineralized skeletons. Common fossils include *\*brachiopods*, *\*trilobites*, *\*ostracods*, and, late in the period, *\*graptolites*. Trilobites are important in the stratigraphic subdivision of the period.

**camera (camara; adj. cameral or camaral)** One of the chambers within a chambered mollusc, e.g. a nautiloid (*\*Nautiloidea*) or ammonoid (*\*Ammonoidea*). In living cephalopods (*\*Cephalopoda*), e.g. *Nautilus*, the chambers contain gas at pressures ranging from about 0.3 to 1 atm (30–100 kPa). Some of the chambers may contain ‘cameral fluid’ which can be extracted by the *\*siphuncle*, allowing the animal to adjust its density according to the depth at which it is living.

**cameral** See CAMERA.

**cameral fluid** See CAMERA.

**Camerata** (class Crinoidea) Subclass of echinoderms (*\*Echinodermata*), with thecal plates typically united by rigid structures composed of numerous small, polygonal plates, and some lower brachial plates, all the plates firmly sutured together. The *\*tegmen* is plated, some of the plates roofing over the mouth and *\*ambulacral* grooves. The Camerata are known from the Lower *\*Ordovician* to Upper *\*Permian*.

**camouflage** In *\*geochemistry*, the way in which a *\*trace* element can substitute for a common element of the same *\*valency* and nearly the same *\*ionic radius* in a crystal lattice, e.g.  $\text{Hf}^{4+}$  for  $\text{Zr}^{4+}$ . The trace element is said to be camouflaged by the more common element. Compare CAPTURE.

**Campanian (Piripauan)** A *\*stage* in the European *\*Cretaceous* (83.5–70.6 Ma ago, Int. Commission on Stratigraphy, 2004), preceded by the *\*Santonian*, followed by the Maastrichtian. See also SENONIAN.

**Campbell–Stokes sunshine recorder** Device that utilizes a glass sphere to concentrate the Sun’s rays on to a calibrated paper, where the resulting

burns register the time and duration of sunshine. The instrument must be pre-set for a given latitude.

**camptonite** A dark-coloured, medium-grained **\*igneous** rock composed of **\*plagioclase** feldspar, barkevikite (a sodium-bearing **\*amphibole**) and/or **\*augite**. Camptonites are one variety of a group of **\*intrusive** rocks collectively known as **\*lamprophyres** which characteristically form **\*dykes**. The original camptonite was named from Campton Falls in New Hampshire, USA.

**Canada balsam resin** A naturally occurring resin, distilled from the bark of *Abies balsamea* (the balsam fir) and other N. American *Abies* species, which, when heated to 160 °C, becomes liquid and is used to cement **\*mineral** or **\*rock** chips to glass slides as part of the process of preparing **\*thin** sections. Its use has been largely superseded by warm- or cold-setting epoxy resins which combine low **\*viscosity** (about 100 centipoise) with high shear strength (about  $11.7 \times 10^6$  Pa), good adhesion, and a **\*refractive index** of 1.54.

**Canadian** A **\*series** (488.3–471.8 Ma ago) of the Lower **\*Ordovician** of N. America, equivalent to the **\*Tremadoc** and **\*Arenig**.

**‘canali’** See MARTIAN CANALS.

**canalizing selection** The elimination of **\*genotypes** that render developing individuals sensitive to environmental fluctuations. Genetic differences may be revealed in organisms by placing them in a stressful environment, or if a severe **\*mutation** stresses the developmental system.

**Canaries current** Cool, oceanic water current, which flows south along the continental margin of Spain, Portugal, and W. Africa. This slow-moving eastern **\*boundary** current is the cause of frequent sea fogs off north-west Spain and Portugal. The water is further cooled by **\*upwelling** of cold water off W. Africa.

**cancrinite** A member of the **\*feldspathoid** group of silicate **\*minerals** with a complex formula, approximately  $(\text{Na,Ca,K})_{6-8}[\text{AlSiO}_4]_6(\text{CO}_3,\text{SO}_4,\text{Cl})_{1-2} \cdot 1-5\text{H}_2\text{O}$ , it may be an alteration product of **\*nepheline**, and is related to vishnevite, the sulphate-rich variety; normally **\*massive**; occurs in nepheline-syenites.

**Canidae** See CARNIVORA.

**Caniformia** See CARNIVORA.

**cannel coal** Fine-grained, tough, compact and uniform *\*bituminous* coal, which is dark grey to black with a dull, greasy *\*lustre* and *\*conchoidal* fracture. It has a high percentage of *\*volatiles*, high ash content, and burns with a smoky, luminous flame. It is a *\*sapropelic coal* formed mainly of *\*lycopod* spores and *\*algae* with fossils of water-dwelling animals. It is found chiefly in Lancashire, England; the local name was 'cannel' meaning 'candle coal'.

**cannel shale** See TORBANITE.

**cannonball bomb** See VOLCANIC BOMB.

**canyon** A deep, steep-sided gorge cut by a river, generally into bedrock. See also SUBMARINE CANYON.

**cap** See PILEUS.

**capacity (of stream)** Maximum load of solid particles that a stream is capable of carrying. It is largely a function of particle size, in that a decrease in the size of particles involves an increase in the total load that can be transported. Ultimately a heavily loaded stream merges imperceptibly with a mud flow.

**capillarity** See CAPILLARY ACTION.

**capillary action (capillarity)** The process by which *\*soil moisture* may move in any direction through the fine (i.e. capillary) pores of the soil, under the influence of *\*surface tension* forces between the water and the soil particles. It is analogous to the rise of water in a cylindrical glass capillary tube whose end is placed in a bath of water in the laboratory, except that the pores of the soil are neither cylindrical, nor smooth-walled, nor clean. See also CAPILLARY MOISTURE.

**capillary fringe** See CAPILLARY ZONE.

**capillary moisture (capillary water)** Moisture that is left in the soil, along with hygroscopic moisture and water vapour, after the gravitational water has drained off. Capillary moisture is held by *\*surface tension* (known in

the USA as ‘water potential’) as a film of moisture on the surface of soil particles and *\*peds*, and as minute bodies of water filling part of the pore space between particles. Curved water surfaces or *menisci* (singular: *meniscus*) form bridges across the pores at the boundaries between their water-filled and air-filled parts. Capillary moisture may move through the soil under the influence of surface tension forces (see [CAPILLARY ACTION](#)) and is available for removal by plant roots.

**capillary water** See [CAPILLARY MOISTURE](#).

**capillary wave** Water wave whose wavelength is less than 1.7 cm and in which the primary restoring force is the *\*surface tension* of the water. The slightest of breezes may cause slight ‘puckering’ of the water surface, and the capillary waves so produced will be smoothed and flattened by the effects of surface tension.

**capillary zone (capillary fringe)** The zone immediately above the *\*water table*, into which water may be drawn upward as a consequence of *\*capillary* action. A typical height for the capillary fringe in clay with a pore radius of 0.0005 mm might be 3 m, compared with less than 10 cm in a fine sand with a pore radius of 0.02 mm.

**capillatus** From the Latin *capillatus* meaning ‘with hair’, a species of *\*cumulonimbus* cloud with fibrous cirriform appearance in its upper parts. It is often associated with an anvil or other protrusion in which wisps of cloud trail from the tip to give a hair-like appearance. Typically this cloud type brings squalls with showers or thunderstorms. See also [CLOUD CLASSIFICATION](#).

**Capitanian** A *\*stage* (265.8–260.4 Ma ago) of the Middle *\*Permian* epoch, preceded by the *\*Wordian* and followed by the *\*Wuchiapingian*.

**cap rock** Layer of hard, impervious rock which lies immediately above a *\*source rock* and which, because of its impervious nature, acts as a barrier, preventing the migration of hydrocarbons or water up-sequence. Such impervious rocks include clay-rich *\*sandstones*, *\*limestones*, and *\*evaporite* deposits associated with salt *\*diapirs*. Where it lies above an oil trap it prevents the upward migration of *\*petroleum*. Above a *\*salt dome*, the layer is usually composed of *\*anhydrite* and *\*gypsum*. In coal mining the term refers to a sandstone layer above shale overlying a coal seam.

**Captorhinomorpha** (order \*Cotylosauria) Suborder of \*reptiles, which appeared in the \*Carboniferous and became extinct after the early \*Permian. The suborder includes the earliest known reptiles, retaining some \*amphibian features, but distinctly reptilian in the form of their skulls, and differing from amphibians in the structure of the pelvic girdle. Because they were ancestral to all later reptile groups, the cotylosaurs are known as 'stem reptiles'.

**capture 1.** Substitution in a crystal lattice of a trace element for a common element with lower \*valency or larger \*ionic radius, e.g. Ba<sup>++</sup> for K<sup>+</sup>. There is often a higher concentration of captured trace elements relative to common elements in the mineral than in the liquid from which it crystallized. *Compare* CAMOUFLAGE. **2.** See RIVER CAPTURE.

**capuliform** Applied to a cap-shaped gastropod (\*Gastropoda) shell, similar to those of the genus *Capulus*.

**Caradoc** A \*series (460.9–449 Ma ago) of the Upper \*Ordovician, underlain by the \*Llandeilo and overlain by the \*Ashgill.

**carbon** Non-metallic element, chemical symbol C, which is unique in the number of compounds it is able to form that contain chains or rings of carbon atoms. This ability to form large, complex molecules in which other elements are bonded to carbon atoms is exploited by all living organisms. The discipline of organic chemistry is essentially the study of cyclic carbon compounds. Carbon is extracted from gaseous carbon dioxide by plants during \*photosynthesis, is incorporated in living matter, and when organic matter decomposes its carbon is oxidized and so returned to the atmosphere as carbon dioxide. Pure carbon occurs naturally as \*diamond, \*graphite, fullerene, and graphene and as the amorphous carbon black. \*Charcoal, produced by the destructive distillation of organic matter, is also a pure form of carbon. In the Earth sciences, carbon is also important in the form of carbonates, as in \*limestones.

**carbon-14** See RADIOCARBON DATING.

**carbonaceous chondrite** Dull, black \*stony meteorite, with little or no metal and abundant carbon; iron occurs as sulphide, silicate, or oxide. Carbonaceous \*chondrites show very little \*metamorphism but display evidence of chemical alteration by water, which continued after their

formation, suggesting the parent body was rocky material mixed with ice. They contain a varied suite of organic compounds, including amino acids and a high content of inert gases. Carbonaceous chondrites have very primitive compositions, comparable to that of the *\*Sun's* atmosphere and the nebula from which the *\*solar* system formed. *See also* METEORITE. *Compare* ACHONDRITE.

**carbonate compensation depth (CCD)** The depth in the sea at which the rate of dissolution of solid calcium carbonate equals the rate of supply. Surface ocean waters are usually saturated with calcium carbonate, so calcareous materials are not dissolved. At mid-depths the lower temperature and higher CO<sub>2</sub> content of sea water cause slow dissolution of calcareous material. Below about 4500 m, waters are rich in dissolved CO<sub>2</sub> and able to dissolve calcium carbonate readily. Carbonate-rich sediments are common in waters less than 3500 m depth, but are completely absent below about 6000 m. *See also* CALCAREOUS OOZE.

**carbonate lump** *See* INTRACLAST.


**carbonate platform** A structure that forms in shallow water where there is extensive deposition of carbonate that is not mixed with *\*siliciclastic* sediments. Carbonate platforms were extensive at times when large areas of the continents lay beneath shallow seas. Today they are much narrower. Changing sea levels have produced long-lived carbonate platforms with structures showing complex cyclical changes. Carbonate platforms are classified as either carbonate ramps or rimmed shelves. A carbonate ramp has a gently sloping surface extending seaward from the coastline, with no sharp break between shallow and deep water. Rimmed shelves have a fairly sharp break where the water depth increases. In some places *\*oolitic* sand has accumulated to form *\*barrier islands*. In others reef-building organisms have built reefs along the rim (e.g. the Great Barrier Reef).

**carbonate ramp** *See* CARBONATE PLATFORM.

**carbonates** 1. Group of *\*minerals* found mostly in *\*limestones* and *\*dolomites*. *\*Calcite* (CaCO<sub>3</sub>) is the most abundant and most important. *\*Aragonite* has the same formula as calcite but is less stable and shells composed of aragonite change to calcite through geological time. Dolomite (or pearl-spar) is the magnesium-bearing carbonate commonly found as a

rock-forming mineral,  $\text{CaMg}(\text{CO}_3)_2$ . 2. The term 'carbonate' is frequently used with reference to those **\*sedimentary rocks** with 95% or more of either **\*calcite** or **\*dolomite**, and is synonymous with **\*limestone**.

**carbonation** Chemical **\*weathering** process involving a reaction between dilute carbonic acid, derived from the solution in water of free atmospheric and soil-air carbon dioxide, and a mineral. The best-known example is the reaction between **\*limestone** (calcium carbonate) and carbonated water (carbonic acid) which yields calcium and bicarbonate ions in solution:



**carbonatite** Unusual **\*igneous** rock, rich in **\*calcite** and other **\*carbonate** minerals (including **\*dolomite** and **\*ankerite**), which is considered to be **\*mantle**-derived. Carbonatites occur as intruded masses, **\*dykes**, and as **\*cone sheets**, and rarely as **\*lavas** and **\*tephra**; and are found in association with **\*alkali**-rich igneous rocks, notably those erupted by the volcanoes of the East African Rift. Rare elements, including the REEs (**\*rare-earth** elements), barium, niobium, thorium, and phosphorus, are often enriched in comparison with many crustal and mantle-derived igneous rocks.

**carbon 'burning'** 'Burning' (see **NUCLEOSYNTHESIS**) that occurs in large stars, due to the intense heat of the stellar core, following **\*hydrogen** 'burning' and **\*helium** 'burning'. During this stage of stellar evolution  $^{12}\text{C}$  nuclei undergo nuclear reactions at temperatures around  $8 \times 10^8$  K to produce elements such as  $^{20}\text{Ne}$ ,  $^{23}\text{Na}$ ,  $^{23}\text{Mg}$ , and  $^{24}\text{Mg}$ . See **OXYGEN 'BURNING'**; **SILICON 'BURNING'**.

**carbon cycle** The movement of carbon through the surface, interior, and **\*atmosphere** of the Earth. Carbon exists in atmospheric gases, in dissolved **\*ions** in the **\*hydrosphere**, and in solids as a major component of organic matter and **\*sedimentary rocks**, and is widely distributed. Inorganic exchange is mainly between the atmosphere and hydrosphere. The major movement of carbon results from **\*photosynthesis** and respiration, with exchange between the **\*biosphere**, atmosphere, and hydrosphere. Rates of exchange are very small, but over geologic time they have concentrated large amounts of carbon in the **\*lithosphere**, mainly as **\*limestones** and **\*fossil fuels**. This carbon was probably present as  $\text{CO}_2$  in the primordial

atmosphere. The burning of fossil fuels and the release of CO<sub>2</sub> from soil air through the clearance of tropical forests may eventually change the balance of the carbon cycle, although the climatic effects may be partly mitigated by the buffering action of the oceans; it is estimated that about 200 billion tonnes of CO<sub>2</sub> have been added to the atmosphere in this way since 1850.

See [GREENHOUSE EFFECT](#).

**carbon dating** See [RADIOCARBON DATING](#).

**Carbon Dioxide Monitoring Mission (MicroCarb)** A French [\\*microsatellite](#) mission that will monitor fluxes of CO<sub>2</sub> at the surface between the atmosphere, oceans, and vegetation. It is scheduled for launch in 2020.

**Carboniferous** Penultimate period of the [\\*Palaeozoic](#) era, preceded by the [\\*Devonian](#) and followed by the [\\*Permian](#). It began about 359.2 Ma ago and ended about 299 Ma ago. In Europe the lower part of the [\\*system](#) is termed the [\\*Dinantian](#). It is divided into two [\\*series](#) and is characterized by marine [\\*limestones](#) with a rich [\\*coral](#)–[\\*brachiopod](#) fauna. In contrast the upper part, the [\\*Silesian](#), which is subdivided into three series, is noted for the deposition of terrestrial and freshwater sediments. The vast forests of the Upper Carboniferous gave rise to the rich coal measures of south Wales, England, Scotland, and many other areas worldwide. N. American geologists subdivide the Carboniferous system into two sub-systems. Of these the lower (359.2–318.1 Ma ago) is named the [\\*Mississippian](#) and is the equivalent of the Dinantian sub-system plus the lower part of the Silesian sub-system. The upper sub-system, the [\\*Pennsylvanian](#) (318.1–299 Ma ago), is the equivalent of most of the Silesian.

**carbon isotopes** Natural carbon is composed of three [\\*isotopes](#): <sup>12</sup>C making up about 98.9%; <sup>13</sup>C about 1.1%; and <sup>14</sup>C whose amount is negligible, but which is detectable because it is radioactive. The relative abundance of these isotopes varies and the study of this variation is an important tool in geologic research, especially [\\*radiometric](#) dating. Carbon-isotope dating is a method of radiometric age-dating using the amount of the heavy, radioactive isotope carbon-14 remaining in organic matter. Carbon-14 has a [\\*half-life](#) of 5730 ± 30 years and the amount of the isotope present can be used to date materials up to about 70 000 years old (see

RADIOCARBON DATING). In diagenetic studies (see DIAGENESIS), measurement of the ratio of carbon-13 to carbon-12 allows the recognition of \*carbonate precipitated from a variety of different sources.

**carbonization** See FOSSILIZATION.

**cardinalia** A collective term that describes the structures found at the posterior end of the interior of the brachial valve of a brachiopod (\*Brachiopoda). In its simplest form, as a cardinal process, it is merely the site of a series of muscle bases. In other species it may be much more complex.

**cardinal septum** One of the initial \*septa (vertical plates within the coral skeleton) which occur in rugose corals (\*Rugosa). When the coral is very young a single proseptum develops in the \*calyce. This soon separates into two, one called the 'cardinal septum', the other the 'counter-cardinal septum'. Other septa then follow to make up the arrangement of septa.

**cardinal tooth** Large \*hinge tooth present in some bivalves which is immediately below the \*umbo. More than one may be present on each \*valve.

**Caribbean current** Warm, ocean water current which flows westward through the Caribbean Sea, passes into the \*Florida current, and thus contributes to the \*Gulf Stream. Its flow velocity averages 0.38–0.43 m/s.

**Caribbean Plate** One of the present-day minor lithospheric \*plates, which is subducting beneath the Antilles in the east, with the \*Cocos Plate subducting beneath it on the western side of Central America. The northern and southern boundaries are both \*faults: the Bartlett Fault separates it from the \*N. American Plate and the Bocono Fault from the \*S. American Plate.

**carina** See KEEL.

**Carlsbad twin** The \*mineral \*orthoclase feldspar forms \*crystals which may twin according to several laws. The most common of these is the Carlsbad law, with a \*composition plane on 010 and with the vertical 'c' (or 'z') axis as twin axis. It is commonly an \*interpenetrant (penetration) twin.

**Carlsberg Ridge** The slow-spreading oceanic \*ridge which separates the \*African Plate and the \*Indo-Australian Plate.

**Carme (Jupiter XI)** One of the lesser satellites of **\*Jupiter**, with a diameter of 30 km; its orbit is **\*retrograde**.

**carnallite** **\*Mineral**,  $\text{KMgCl}_3 \cdot 6\text{H}_2\text{O}$ ; sp. gr. 1.6; **\*hardness** 1–2; **\*orthorhombic**; normally white, occasionally yellowish and reddish; greasy **\*lustre**; **\*massive** and granular, the pseudo-hexagonal **\*crystals** being rare; no **\*cleavage**; occurs in **\*evaporite** deposits and is soluble in water; bitter taste. It is used as a fertilizer, and is named after the 19th-century mining engineer R. V. Carnall.

**carnassial** In many **\*Carnivora**, a modification of premolar or molar teeth, commonly the lower first molar and the upper last premolar, giving them a scissor-like shearing action used for cutting flesh.

**Carnian (Karnian) 1.** A **\*Triassic** **\*age**, preceded by the **\*Ladinian**, followed by the **\*Norian**, and dated at 228–216.5 Ma ago (Int. Commission on Stratigraphy, 2004). **2.** The name of the corresponding European **\*stage**, roughly contemporaneous with the Banan (China), and Oretian and Otamitan (New Zealand).

**Carnivora** (cohort Ferungulata, superorder Ferae) An order that comprises the modern carnivorous placental mammals and their immediate ancestors. It used to be divided into two suborders, the Fissipedia (mainly land-dwelling) and Pinnipedia (seals, sea lions, walrus), but a more modern classification is into Caniformia (dog-like) and Feliformia (cat-like), with the ‘pinnipeds’ belonging to the former. The carnivores are descended from a single stock of the probably insectivorous, placental mammals of the early **\*Cretaceous**, the change being reflected in their dentition. Strong incisors for biting, and canines for piercing, were retained from the insectivorous forms, but in general carnivores acquired modified cheek teeth (carnassials) specialized for shearing. These subsequently became reduced in those carnivores which adopted a herbivorous diet. Hoofs have rarely developed, as claws are used for seizing prey, and digits are never greatly elongated (and, apart from the pollex and hallux, they are not reduced). The first true carnivores were the weasel-like Miacidae of the **\*Palaeocene**, which had diverged by the end of the **\*Eocene** to give the Canidae (dogs) and Mustelidae (weasels and their allies) as one branch and the Viverridae (Old World civets) and Felidae (cats) as another. According to some authors, the Mustelidae later branched again to give the Phocidae (seals); and the

Canidae diversified widely to produce such forms as the Amphicyonidae ('dog-bears'), Otariidae (sea lions), Procyonidae (raccoons and pandas), and, ultimately, Ursidae (bears); but molecular studies seem to indicate that the Phocidae and Otariidae are descended from a single ancestor which was related to the mustelid–ursid–procyonid stem. Finally, the Hyaenidae (hyenas) emerged in the late **\*Miocene** from viverrid stock; this is the youngest of the carnivore families.

**carnosaur** **\*Saurischian** (lizard-hipped), carnivorous **\*dinosaur**. Carnosaurs were bipedal, powerfully built, and possessed large, dagger-like teeth. **\*Tyrannosaurus rex**, of the Upper **\*Cretaceous**, represented the culmination of the carnosaur line.

**carnotite** **\*Mineral**,  $K_2(UO_2)_2(VO_4)_2 \cdot 1-3H_2O$ ; sp. gr. 4.5; **\*hardness** 2.0–2.5; **\*monoclinic**; normally bright yellow to greenish-yellow; pearly **\*lustre**; **\*cleavage** perfect, basal {001}; strongly radioactive; occurs in the weathered zone of sedimentary **\*uranium** ores, especially in **\*sandstones** enriched with organic matter, normally as coatings and masses in rocks. It is an important ore for uranium.

**carpal** One of the bones of the wrist that articulate with the digits.

**Carpentarian** A **\*stage** (1800–1300 Ma ago) of the Lower–Middle **\*Proterozoic** of south-eastern Australia, underlain by the **\*Nullaginian** and overlain by the **\*Adelaidean**.

**Carpenter, William Benjamin** (1813–85) A surgeon and comparative anatomist, who used microscopy to investigate the structure of fossil shells, especially **\*foraminifera** and **\*crinoids**. He took part in several scientific voyages, including the **\*Challenger** expedition, as a result of which he became interested in marine physics and developed a theory about oceanic circulation.

**carpoids** (phylum Echinodermata) Informal collective term describing the homalozoan classes Homoiostelea, Homostelea, and Stylophora. See **HOMALOZOA**.

**carrier element** **1.** Major element in a **\*mineral**, for which a trace amount of another element may substitute, e.g. Mg (the carrier element) replaced by Ni (the **\*trace** element) in **\*olivine**. **2.** Inactive material, isotopic with a radioactive transmutation product, which is added to act as a carrier for

active material in subsequent chemical reactions. Carrier elements are used in analyses of small samples, as the mass of radioactive material produced in a nuclear reaction is usually too small to be suitable for ordinary analytical procedures such as precipitation and filtration.

**Cartesian projection** A technique for the mapping of space in which every plane in the area being mapped is projected on to a plane in the map and every line on to a line. Each point in the area under study is identified by three values, representing its location in relation to three mutually perpendicular axes (the Cartesian coordinates of the point). These coordinates are transformed mathematically into a homogeneous set of four coordinates which can then be plotted to produce a graphic representation (a map). The word 'Cartesian' is derived from the name of René Descartes (1596–1650).

**cartilage** In vertebrates, flexible skeletal tissue formed from groups of rounded cells lying in a matrix containing **\*collagen** fibres. It forms most of the skeleton of embryos and in adults is retained at the ends of bones, in intervertebral discs, and in the pinna of the ear; in **\*Elasmobranchii**, calcified cartilage rather than true **\*bone** provides the entire skeleton.

**cartilaginous fish** Fish in which the skeleton, including the skull and jaws, consists entirely of **\*cartilage** and never, even in the adult stage, comprises bony tissue. Sharks and rays (**\*Chondrichthyes**) have a cartilaginous skeleton, as have other lower vertebrates, e.g. the jawless lampreys and hagfish (**\*Agnatha**).

**CartoSat** A series of Indian satellites that provide high-resolution optical imagery. CartoSat-2 was launched on 10 January 2007, -2A on 28 April 2008, -2B on 12 July 2010, -2D on 15 February 2007, and -2E on 23 June 2017, all from the Satish Dhawan Space Centre, India.

**Cascade SmallSat and Ionospheric Polar Explorer (CASSIOPE)** A programme of **\*minisatellites** by the Canadian Space Agency and Industrial Technologies Office that carries an experimental payload (Cascade) to demonstrate the world's first commercial digital courier service able to store and forward file transfer techniques, eight instruments (Enhanced Polar Outflow Probe) to measure the interaction between the upper atmosphere and **\*solar** wind, and a test bed for bus and subsystem functions. It was launched on 29 September 2013, from California.

**cascading system** In geomorphology, a type of dynamic system characterized by the transfer of mass and energy along a chain of component sub-systems, such that the output from one sub-system becomes the input for the adjacent sub-system. An example is the **\*valley glacier**, where the inputs of snowfall and rock debris from the slopes above, and potential energy (derived from elevation), are cascaded through a sequence of climatic environments with a progressive reduction in mass and dissipation of energy, the output from the glacier being sediment and water which form the input to the **\*proglacial** sub-system.

**casing** Steel tubes, usually screwed together, which line a **\*borehole** to prevent caving-in of the side walls. The gap between the casing and the natural rock is often filled with concrete. Casings are used mainly where the borehole passes through clays.

**Cassadagian** See CHAUTAUQUAN.

**Cassini** A mission to **\*Saturn** launched in 1997 and operated jointly by **\*NASA**, **\*ESA**, and the Italian Space Agency to study Saturn's atmosphere, rings, and satellites, including a fly-by of **\*Enceladus**. The mission was launched on 15 October 1997, entered Saturn's orbit on 1 July 2004, and ended on 15 September 2017, when it entered Saturn's atmosphere and burned up. See also HUYGENS.



<http://saturn.jpl.nasa.gov/>

- A joint ESA and NASA mission that studied Titan, moon of Saturn, and carried the Huygens probe.

**Cassini Division** See RESONANCE.

**CASSIOPE** See CASCADE SMALLSAT AND IONOSPHERIC POLAR EXPLORER.

**cassiterite (tinstone)** **\*Mineral**, SnO<sub>2</sub>; sp. gr. 6.8–7.1; **\*hardness** 6–7; **\*tetragonal**; usually reddish-brown to nearly black, but it can be yellowish and ruby; white to grey **\*streak**; **\*adamantine** lustre; **\*crystals** often pyramidal and **\*prismatic**, may also be **\*massive** and granular; **\*cleavage** prismatic {100}, {110}; occurs typically in high-temperature **\*hydrothermal** veins, with **\*granites** and **\*pegmatites**; associated minerals:

\*topaz, \*quartz, \*tourmaline, \*mica, \*chlorite, and high-temperature metallic \*ores; also worldwide in \*alluvial deposits because of its resistance to chemical and physical attack. Cassiterite is the only important ore for tin.

**cast** The preserved sediment infill of an impression or mould made in the top of a bed of soft \*sediment (e.g. \*flute cast, \*load cast, \*tool cast, casts of footprints or \*trails).

**Castalia** A \*solar system near-Earth asteroid (No. 4769), measuring  $1.8 \times 0.8$  km; approximate mass  $10^{11}$  kg; orbital period 0.41 years. It is a double-lobed object, each lobe about 0.75 km in diameter.

**castellanus** From the Latin *castellum*, meaning ‘castle’, a cloud species commonly associated with the upper parts of \*altocumulus, \*stratocumulus, \*cirrus, and \*cirrocumulus. Cumuliform, turreted protrusions extend in linear fashion from the cloud top, producing a crenulate form. *See also* CLOUD CLASSIFICATION.

**Castlecliffian** A \*series (1.1–0.01143 Ma ago) in the \*Pleistocene epoch of New Zealand, underlain by the \*Nukumaruan, overlain by the \*Holocene, and roughly contemporaneous with the uppermost Calabrian and Emilian \*stages, and the subsequent Upper \*Pleistocene series.

**castle koppie** *See* KOPPIE.

**Castlemainian** A \*stage (471–470 Ma ago) of the Middle \*Ordovician of Australia, underlain by the \*Chewtonian and overlain by the \*Ypeenian.

**CAT** *See* CLEAR-AIR TURBULENCE.

**cataclasis** *See* CATACLASITE.

**cataclasite** Rock that has been deformed by the process of shearing and \*granulation (cataclasis). Cataclasites are the products of dislocation \*metamorphism and \*tectonism. *See also* MYLONITE.

**catagenesis** Following \*diagenesis, in which sedimentary material is compressed and undergoes chemical changes, a phase in the formation of \*petroleum and \*natural gas during which continuing sedimentation and subsidence produce temperatures of 50–150 °C and \*kerogen is produced. *Compare* METAGENESIS.

**cataracta** See CLOUD CLASSIFICATION.

**cataractagenitus** See CLOUD CLASSIFICATION.

**catarrhine** In primates, applied to nostrils that are close together and open downwards. Old World monkeys, apes, and humans have catarrhine nostrils.

**catastrophic evolution (catastrophic speciation)** Theory proposing that environmental stress can lead to the sudden rearrangement of chromosomes, which in self-fertilizing organisms may then give rise sympatrically to a new species (see SYMPATRIC EVOLUTION). Recent research suggests that at best this explanation applies only to some special cases.

**catastrophism** Theory that associates past geologic change with sudden, catastrophic happenings. Early geologists, including \*Cuvier, \*Buckland, and \*Sedgwick, claimed that catastrophism was a sound scientific theory. Although it met with considerable scorn in more recent times, many modern geologists would describe themselves as ‘neocatastrophists’.

**catazone** See KATAZONE.

**catchment** The area from which a surface watercourse or a \*groundwater system derives its water. Catchments are separated by \*divides. A surface catchment area may overlie an \*aquifer system, but may be unconnected with the aquifer rock itself if there are intervening impermeable \*aquicludes. In US usage, a catchment is often termed a ‘watershed’.

**catena** 1. Topographic sequence of soils, of the same age and usually on the same parent material, that is repeated across larger landscape transects. Individual \*soil-profile types are related to site conditions and to position on a slope. The term was introduced in E. Africa in the 1930s, and is mainly applicable in certain non-glaciated landscapes, particularly those with small, hilly relief, e.g. \*loess areas. 2. A chain of craters on the surface of an extraterrestrial body.

**cateniform** Applied to the form of the \*corallum which occurs in the tabulate corals (\*Tabulata). The \*corallites are elongated and joined side by side to make fence-like structures. In end section they resemble chain links,

hence the name (from the Latin *catena* meaning ‘chain’). See [COMPOUND CORALS](#).

**cathode** See [CATION](#).

**cathodoluminescence** The luminescence induced by the bombardment of [\\*minerals](#) in polished [\\*thin](#) section with electrons. The technique is particularly useful in the identification of mineral [\\*cements](#) and overgrowths produced in successive episodes, or for distinguishing between particles and [\\*authigenic](#) overgrowths. The particles and each separate generation of cement will be of slightly different chemical composition, and so will luminesce with slightly different colours. See [QUARTZ OVERGROWTH](#).

**cation** A positive [\\*ion](#), i.e. an atom, or complex of atoms, that has lost one or more electrons and is left with an overall positive electric charge, e.g.  $\text{Na}^+$ ,  $\text{Mg}^{2+}$ ,  $\text{NH}_4^+$ . The name is derived from the fact that when an electric current is passed through a conducting solution the positive ions present in the solution move towards the cathode (the negative electrode). Compare [ANION](#).

**cation exchange** Process in which [\\*cations](#) in solution are exchanged with cations held on the exchange sites of mineral and organic matter, particularly on the surfaces of [\\*colloids](#) of [\\*clay](#) and [\\*humus](#).

**cation-exchange capacity (CEC)** The total amount of exchangeable [\\*cations](#) that a particular material or soil can adsorb at a given [\\*pH](#). Exchangeable cations are held mainly on the surface of [\\*colloids](#) of [\\*clay](#) and [\\*humus](#), and are measured in milligram-equivalents per 100 g of material or soil.

**cation ordering** The phenomenon, the extent of which is temperature dependent, in which a [\\*cation](#) shows preference for the site it occupies because it provides greater chemical stability.

**CATS** See [CLOUD-AEROSOL TRANSPORT SYSTEM](#).

**cat's eye** See [RIEBECKITE](#).

**cauda** See [CLOUD CLASSIFICATION](#).

**caudal** Pertaining to the tail.

**caudal vertebra** See VERTEBRA.

**cauldron-subsidence** Collapse of a volcanic **\*crater** due to the evacuation of a large **\*magma chamber** (cauldron) and marked by a **\*ring fracture**, or **\*ring-dyke**. There are many examples of cauldron-subsidence (e.g. Glencoe, Scotland), although the various mechanisms of their formation are not necessarily identical.

**Cautleyan** A **\*stage** (447.5–446.5 Ma ago) of the **\*Ordovician** in the **\*Ashgill**, underlain by the **\*Pusgillian** and overlain by the **\*Rawtheyan**.

**cavate** 1. Applied to **\*spores** where the **\*exine** layers are separated by a cavity. 2. Applied to dinoflagellate (**\*Dinophyceae**) cysts where there is a space between the periphragm and the endophragm.

**cavern porosity** See CHOQUETTE AND PRAY CLASSIFICATION.

**cavitation** The collapse of a region because it is at a lower pressure than the ambient environment. The collapse of a gas bubble, in water, causes an **\*implosion** that may be detectable as a seismic energy source. If the hydrostatic pressure in a rock exceeds the hydrostatic pressure of the **\*drilling** fluid, this can cause a collapse of the walls around the hole, or a ‘blow-out’ of the drill stem. Erosion by cavitation occurs in waterfalls, rapids, and subglacial river channels, and can produce pot-holes.

**cavum** See CLOUD CLASSIFICATION.

**cavus** (*pl.* **cavi**) An irregular, steep-sided depression or hollow on the surface of an extraterrestrial body.

**cay** Small, flat, marine island formed from coral-reef material or **\*sand**. The term is applied, for example, to the low-lying, sparsely vegetated islands off the coast of southern Florida.

**Cayugan** A **\*series** (421.3–416 Ma ago) of the **\*Silurian** of N. America equivalent to the Lower **\*Ludlow** to terminal **\*Pridoli**, preceded by the **\*Gorstian**, followed by the **\*Lochkovian**.

**Cazenovian** See ERIAN.

**CBERS** See CHINA-BRAZIL EARTH RESOURCES SATELLITE.

**CBR** See CALIFORNIA BEARING RATIO.

**CCD** 1. See CARBONATE COMPENSATION DEPTH. 2. See CHARGE-COUPLED DEVICE.

**CCL** See CONVECTIVE-CONDENSATION LEVEL.

**CCN** See CLOUD CONDENSATION NUCLEI.

**<sup>14</sup>C dating** See RADIOCARBON DATING.

**CDP** See COMMON DEPTH POINT.

**CDP stack** See COMMON-DEPTH-POINT STACK.

**CEC** See CATION-EXCHANGE CAPACITY.

***Celastrphyllum circinerve*** One of the earliest flowering plants known to palaeobotanists. Flowering plants of the family Celastraceae began with *C. circinerve* in the Early \*Cretaceous, and survive today in the form of *Euonymus europaeus*, the spindle tree. See ANGIOSPERM.

**celerity** Velocity with which a wave advances. The celerity ( $c$ ) of an ideal wave is related to its wavelength ( $\lambda$ ) and frequency ( $f$ ) by the wave equation  $c = f\lambda$ . The wave frequency ( $f$ ) is the number of waves ( $n$ ) passing a point in unit time ( $t$ ), i.e.  $f = n/t$ . In deep water the wave celerity may be calculated by the equation:  $c = (g\lambda/2\pi)^{1/2} = 1.25\sqrt{\lambda}$ , where  $\lambda$  is the wavelength in metres and  $g$  is the \*acceleration due to gravity (9.81 m/s<sup>2</sup>). The speed of shallow water waves may be calculated by the equation:  $c = (gd)^{1/2} = 3.13\sqrt{d}$ , where  $d$  is the depth of water in metres.

**celestial reference frame** In \*geodesy, a reference frame in which the equations of motion of satellites are related to the relatively fixed positions of distant stars. It is used to express positional coordinates derived from astronomical or satellite measurements. A celestial reference frame cannot be perfectly accurate, because all stars exhibit some \*proper motion with respect to each other.

**celestite** \*Mineral SrSO<sub>4</sub> which may form a \*solid solution series with \*barite; sp. gr. 3.9–4.0; \*hardness 3.0–3.5; \*orthorhombic; faint blue to colourless, sometimes stained red; white \*streak; \*vitreous \*lustre;

**\*crystals** normally very **\*tabular**, resembling barite, but can also occur fibrous and granular; **\*cleavage** perfect basal {001}, present {210}, {010}; often fluorescent; occurs in **\*sedimentary rocks**, particularly **\*dolomites**, as a cavity-lining in association with barite, **\*gypsum**, **\*anhydrite**, and **\*halite**, with gypsum and anhydrite in **\*evaporite** deposits, also as a **\*gangue** mineral in hydrothermal veins with **\*galena** and **\*sphalerite**, and it will form **\*concretionary** masses in clays and marls. It is the main **\*ore mineral** for strontium and strontium compounds.

**cement** 1. Manufactured powder made from limestone and clay which sets to a solid mass when mixed with water. Commercial cements have to fulfil certain defined standards. Combined with **\*aggregate** it forms concrete. 2. Material, e.g. **\*calcite**, that fills open **\*pore space** in fragmental and organic **\*sediments**.

**cementation** Process by which **\*sedimentary rock** particles or fragments are cemented together after deposition. Cementing materials are deposited from the mineral-rich waters that percolate through the open **\*pore space** of the rock. The percentage of cement depends on the amount of pore space and on the mud content within a given rock.

**cemented** Applied to massive, infilled, and **\*indurated** mineral soil: such soil has a hard and often brittle consistency because soil particles are joined together by cementing substances, e.g. calcium carbonate, silica, iron and aluminium oxide, or **\*humus**. Cemented soil usually appears as a highly distinctive and resistant horizon. *Compare* CEMENT; CEMENTATION.

**Cenomanian** 1. A Late **\*Cretaceous** **\*age**, preceded by the **\*Albian**, followed by the **\*Turonian**, and dated at 99.6–93.5 Ma ago (Int. Commission on Stratigraphy, 2004). 2. The name of the corresponding European **\*stage** for which the **\*type locality** is near Le Mans, France.

**Cenozoic (Cainozoic, Kainozoic)** **\*Era** of geologic time extending from about 65.5 million years ago to the present. It includes the **\*Palaeogene** and **\*Neogene** periods: the so-called ages of mammals and man. **\*Molluscs** and **\*microfossils** are used in the stratigraphic subdivision of the era. The **\*Alpine–Himalayan** orogeny reached its climax during this period of geologic time.



<https://ucmp.berkeley.edu/cenozoic/cenozoic.php>

- The Cenozoic Era.

**centipedes** See MYRIAPODA.

**Centrales** See BACILLARIOPHYCEAE.

**Central European Sea** See PARATETHYS.

**central limit theorem** In statistics, the theorem stating of a series of data sets drawn from any *\*probability distribution*, that the distribution of the *\*means* of those data sets will follow a normal distribution.

**central vent volcano** A point source of the Earth's surface through which *\*lavas*, *\*pyroclastics*, and gas are erupted. The eruptive products have their thickest accumulation around the point source and build up a low-angle shield or higher-angle cone topography constituting the volcanic pile. Later eruptions may occur from fissures within the volcanic pile, which radiate and are fed from the point source. Continuous release of gas (explosive or quiet) from the point source ensures the existence of an open vent at the top of the volcanic pile. This vent can often be enlarged by the peripheral collapse of its walls. See also VOLCANO.

**centre of curvature** The point at which arcs of the circle which forms the surfaces of a typical parallel-concentric *\*fold* have a common origin. The radii of such arcs represent the constant *\*orthogonal* thickness of layers typical of this class of fold.

**centre of symmetry** See CRYSTAL SYMMETRY.

**centric diatoms** See BACILLARIOPHYCEAE.

**centrifugal pump** See PUMP.

**centripetal drainage pattern** See DRAINAGE PATTERN.

**Centroceratida** (class *\*Cephalopoda*, subclass *\*Nautiloidea*) Order of generally *\*evolute*, *\*gyroconic* cephalopods in which the *\*siphuncle* is sub-central in position. Some forms are *\*nautiliconic*. *\*Sutures* are tri-lobed. The order ranges in age from Lower *\*Devonian* to Upper *\*Jurassic*.

**centrum** See VERTEBRA.

**Cephalaspida** See OSTEOSTRACI.

**Cephalaspis** (order \**Osteostraci*) One of the best-known osteostracans. The bony head shield tapered posteriorly and laterally to form two 'horns'. The body segments were freed from the head shield, allowing greater mobility than in earlier genera.

**cephalic** Pertaining to the head.

**cephalic spine** A spine occurring on the \**cephalon* (head) of a trilobite (\**Trilobita*).

**cephalic suture** In trilobites (\**Trilobita*), the structure that includes the facial and ventral cephalic sutures. The facial sutures situated on the dorsal part of the \**cephalon* are of three main types: protoparian, where the suture runs from the anterior border of the cephalon to a position anterior to the \**genal* angle; opisthoparian, where the suture runs from the anterior margin to cut the posterior margin posterior to the genal angle; and marginal, where the suture runs along the edge and is not visible on the upper surface. In one family the facial suture runs directly through the genal angle and is 'gonatoparian'. On the ventral side of the cephalon the facial sutures are joined and continue as the ventral cephalic suture. An elongated rostral plate (see **ROSTRUM**) in the anterior mid-line of the ventral surface has a rostral suture anterior to it and a hypostomal suture posterior to it. In some genera the lateral ventral sutures are combined as a single median suture.

**Cephalochordata (Acrania)** Subphylum of \**Chordata*, containing only *Amphioxus* (lancelet), probably the most primitive of living chordates, although some soft-bodied \**Cambrian* fossils are dubiously referred to the group. Cephalochordates have a \**notochord* extending into the head, gill slits, and segmented muscle blocks, and are thought to resemble the ancestors of vertebrates.

**cephalon** The anterior or head region of a trilobite (\**Trilobita*), which consists of at least five fused segments and is generally semicircular in shape. The glabella (the raised median part of the cephalon) is very variable in size and structure and it is probable that in life the stomach lay beneath it. Between the lateral margins of the glabella and the facial suture (see **CEPHALIC SUTURE**) is the fixed cheek (fixigena) and between the facial suture and the lateral margin is the free cheek (librigena). The glabella and

the fixed cheeks together are known as the ‘cranidium’. Above the visual surface of the eye is the palpebral lobe. In **\*Cambrian** trilobites the eyes are commonly connected to the glabella by narrow ocular ridges. The glabella may be indented by pairs of furrows and a transverse furrow separates the posterior occipital segment from the remainder of the glabella. In some families (e.g. the Trinucleidae) the border of the cephalon is developed into an extensive, pitted, cephalic fringe. The dorsal **\*exoskeleton** of the cephalon also continues on to the ventral surface as the ‘doublure’.

**Cephalopoda (cephalopods)** Literally ‘head- foot’, a class of **\*Mollusca**, exclusively marine, related to the **\*Bivalvia** and **\*Gastropoda**. The class includes the **\*Nautiloidea** (nautiloids), Sepioidea (cuttlefish), Teuthoidea (squids), Octopoida (octopuses), and the extinct **\*Ammonoidea** (goniatites, ceratites, ammonites) and **\*Belemnitida** (belemnites). The earliest forms belonged to the Nautiloidea and date from the Upper **\*Cambrian**.

**ceratites** See **AMMONOIDEA**.

**ceratoid** See **SOLITARY CORALS**.

**Ceratopsia** Horned, **\*ornithischian** (bird-hipped) **\*dinosaurs**, which had beak-like jaws, from the Upper **\*Cretaceous**. The head accounted for about one-third of the total length of the body because of the development of a large, bony frill which protected the neck and shoulders. *Triceratops* is perhaps the best-known member of the group. It was 5–6 m long and had three forward-projecting horns, one over each eye and the third over the nose.

**Ceres** A **\*solar** system **\*dwarf** planet, diameter 974 km, approximate mass  $10^{21}$  kg; rotation period 9.078 hours; orbital period 4.6 years. Ceres was discovered in 1801 by G. Piazzi. In 2007, the spacecraft *Dawn* took off on a mission to study Vesta and Ceres. It arrived at Ceres in March 2015 to map the surface in detail and take close-up images of a pair of bright spots on the planet believed to be deposits of water ice or mineral salts. In 2018 it was still in orbit about Ceres, propelled by an **\*ion engine**.



<https://dawn.jpl.nasa.gov/mission/>

- Dawn Mission Overview.

**cerioid** Applied to those corals in which the individuals comprising the colony are packed together and **\*corallites** are polygonal in section, each individual corallite retaining its wall. See **COMPOUND CORALS**.

**cerussite** **\*Mineral**,  $\text{PbCO}_3$ ; sp. gr. 6.4–6.6; **\*hardness** 3.0–3.5; **\*orthorhombic**; usually white or grey; white **\*streak**; **\*adamantine** lustre; crystals often **\*prismatic** or **\*tabular**, also **\*acicular**, but can also occur granular, **\*massive**, and compact; **\*cleavage** good {110}, {021}; of **\*secondary** origin in the oxidized zone of lead veins, associated with **\*anglesite**, **\*galena**, **\*smithsonite**, **\*pyromorphite**, and **\*sphalerite**; soluble (with effervescence) in warm, dilute nitric acid. It is an **\*ore mineral** of lead.

**cervical vertebra** See **VERTEBRA**.

**CFESat** See **CIBOLA FLIGHT EXPERIMENT SATELLITE**.

**CFOSAT** See **CHINESE-FRENCH OCEANOGRAPHY SATELLITE**.

**c.g.s. system** A set of units of measurement derived from the metric system and based on the centimetre, gram, and second. It has now been largely replaced by the SI (Système International d'Unités) system.

**Chadian** A **\*stage** of the **\*Visean** epoch, dated at 345.3–341 Ma ago (Int. Commission on Stratigraphy, 2004), preceded by the **\*Tournaisian** and followed by the **\*Arundian**.

**Chaetognatha** Phylum comprising the arrow worms, first encountered in the fossil record in **\*Carboniferous** rocks.

**chain-former** Class of element oxides which form chain structures in **\*silicate \*melts**, e.g.  $\text{SiO}_2$ ,  $\text{KAlO}_2$ ,  $\text{NaAlO}_2$ ,  $\text{Ca}_{1/2}\text{AlO}_2$ , and  $\text{Mg}_{1/2}\text{AlO}_2$ .

**chain lightning** See **PEARL-NECKLACE LIGHTNING**.

**chain-modifier** Class of element oxides which modify or disrupt chain structures in **\*silicate \*melts**, e.g.  $\text{CaO}$ ,  $\text{K}_2\text{O}$ ,  $\text{Na}_2\text{O}$ ,  $\text{MgO}$ ,  $\text{FeO}$ ,  $\text{TiO}$ , and  $\text{Al}_2\text{O}_3$ .

**chain silicate** See **INOSILICATE**.

**chalcedony** The group name for *\*cryptocrystalline* varieties of silica composed of minute crystals of *\*quartz* with submicroscopic pores and with composition ranging from  $\text{SiO}_2$  to  $\text{SiO}_2.n\text{H}_2\text{O}$  and including the *\*minerals* *\*agate*, *\*chert*, *\*opal*, *\*onyx*, *\*jasper*, and *\*flint*; sp. gr. 2.50–2.67; habits variable from stalactitic to massive; commonly white, greyish-white, or grey, and occasionally yellow.

**chalcocite (copper glance)** *\*Mineral*,  $\text{Cu}_2\text{S}$ ; sp. gr. 5.5–5.8; *\*hardness* 2.5–3.0; *\*orthorhombic*; dark lead-grey tarnishing to black; black *\*streak*; *\*metallic* *\*lustre*; *\*prismatic* and *\*tabular* *\*crystals* are rare, it is usually found *\*massive*, or as powdery coatings; *\*cleavage* prismatic, poor; can occur in *\*hydrothermal* veins in a *\*primary* state, but more usually found in the zones of *\*supergene* enrichment of copper orebodies. It is an important *\*ore mineral* for copper.

**chalcophile** Applied to elements having a strong affinity for sulphur, which concentrate in sulphides and are typical of the Earth's *\*mantle* rather than its *\*core*. Commonly found in *\*sulphide* minerals and *\*ores*. Typical chalcophile elements are Cu, Zn, Pb, As, and Sb. *Compare* *ATMOPHILE*; *BIOPHILE*; *LITHOPHILE*; *SIDEROPHILE*.

**chalcopyrite (copper pyrites)** Most common copper mineral,  $\text{CuFeS}_2$ ; sp. gr. 4.1–4.3; *\*hardness* 3.5–4.0; *\*tetragonal*; brass-yellow, often with an iridescent tarnish; greenish-black *\*streak*; *\*metallic* *\*lustre*; sometimes *\*massive*, crystals usually *\*tetrahedra*; *\*cleavage* imperfect {011}; *\*primary mineral* found in *\*igneous* rocks and *\*hydrothermal* veins, in association with *\*pyrite*, *\*pyrrhotite*, *\*cassiterite*, *\*sphalerite*, *\*galena*, *\*calcite*, and *\*quartz*, an important mineral in *\*porphyry* copper deposits, and also occurs in quartz *\*diorite*, and in *\*pegmatites*, crystalline *\*schists*, *\*porphyry* copper deposits, *\*syngenetic* copper ores and *\*skarns*, and contact *\*metamorphic zones*. It is deeper in colour than pyrite, more brittle and harder than *\*gold*, is soluble in nitric acid, and its alteration products are *\*secondary* copper minerals. It is a major *\*ore mineral* for copper.

**chalk** Porous, fine-grained rock, predominantly composed of the calcareous skeletons of micro-organisms, e.g. *\*coccolithophores* and *\*foraminifera*. The Chalk Formations of the Upper *\*Cretaceous* of Europe form the White Cliffs of Dover and the cliffs south of Calais.

**Challenger expedition** (1872–5) The first expedition to explore the deep oceans was led by John **\*Murray** (1841–1914), in the British naval ship HMS *Challenger*. With a staff of biologists, chemists, and geologists, the expedition surveyed the Atlantic, Indian, Antarctic, and Pacific Oceans, taking soundings and collecting specimens in dredges. The extent of the **\*Mid-Atlantic Ridge** was first demonstrated by the crew of the *Challenger*.

**Challenging Minisatellite Payload (CHAMP)** A German **\*minisatellite** mission to study global long- to medium-wavelength recovery of the static and time-variable Earth gravity field from orbit perturbation, for use in solid-Earth and **\*geodesy** research and in studying ocean currents and climate. The satellite was launched on 15 July 2000, but after 3718 days of supplying data it burnt up on 19 September 2010 over the Sea of Okhotsk.

**chalybdite** See **SIDERITE**.

**chalybeate** Applied to natural waters containing iron.

**Chamberlin, Thomas Chrowder** (1843–1928) Professor of geology at Chicago University and Head of the Glacial Division of the US Geological Survey, in which capacity he mapped the ice deposits of Wisconsin, Chamberlin developed a theory of successive episodes of mountain building, around an ancient continental **\*craton**. With Moulton, he proposed the **\*planetismal hypothesis** of the formation of the Earth.

**Chambers, Robert** (1802–71) The author of *Chambers's Encyclopaedia*, who in 1844 published anonymously a book called *Vestiges of a Natural History of Creation*, in which he revived the idea of evolution first proposed by **\*Lamarck** 30 years earlier. The book's popularity and notoriety refocused attention on this issue and so paved the way, among the general public, for **\*Darwin's** *Origin of Species*.

**chamosite** Silicate **\*mineral** (see **SILICATES**) and member of the **\*chlorite** group and particularly of the septichlorite group which also includes the minerals amesite, greenalite, and cronstedite; composition  $\text{Fe}^{2+}_{10}\text{Al}_2[\text{Si}_3\text{AlO}_{10}]_2(\text{OH})_{16}$ ; soft but massive; occurs in sedimentary ironstones where it may form by alteration from the original **\*siderite** ore, and is found in association with **\*chert** and **\*clay**.

**Chamovnicheskian** A *\*stage* (306–305 Ma ago) in the *\*Kasimovian* epoch, preceded by the *\*Krevyakinskian* and followed by the *\*Dorogomilovskian*.

**CHAMP** See CHALLENGING MINISATELLITE PAYLOAD.

**Champlainian** A *\*series* of the Middle *\*Ordovician* of N. America, generally equivalent to the *\*Llanvirn* and Middle *\*Caradoc*.

**Chandler wobble** The free oscillation of the *\*Earth's* pole of rotation, discovered in 1891 by the American astronomer Seth Carlo Chandler (1846–1913). This wobble in the Earth's rotation has a 435-day periodicity and appears to have a decay time of the order of 40 years. The cause (excitation) is unknown: atmospheric effects appear to be on too short a timescale; *\*earthquake* activity has been proposed, but has not been established. The major cause appears to be related to the Earth's *\*core* and its magnetic coupling with the lower *\*mantle*. In 2005 the wobble changed phase by 180°.

**Changhsingian** A *\*stage* in the Late *\*Permian \*epoch*, preceded by the *\*Wuchiapingian*, followed by the *\*Induan (\*Triassic)*, and dated at 253.8–251 Ma ago (Int. Commission on Stratigraphy, 2004). It is roughly contemporaneous with the Bundsandstein (western Europe), upper Amarasian (New Zealand), and upper *\*Ochoan* (N. America).

**Changlangpuan** A Chinese *\*stage* (523–518 Ma ago) of the Early *\*Cambrian*, preceded by the *\*Qungzusian* and followed by the *\*Longwangmiaoan*.

**channel** 1. The preferred linear route along which surface water and *\*groundwater* flow is usually concentrated (although water can flow across wide, flat surfaces as sheet flow). It is commonly a linear, concave-based depression (e.g. river channel, submarine fan channel). The geometry may be sinuous, anastomosing, or straight, and with a widely variable width-to-depth ratio. See BRAIDED STREAM; MEANDER. 2. A narrow sea-way connecting two wider bodies of water (e.g. the English Channel). 3. In remote sensing, the range of wavelengths recorded by a single detector to form an image.

**channel and vug porosity** See CHOQUETTE AND PRAY CLASSIFICATION.

**Channel Deposits** See MARTIAN TERRAIN UNITS.

**channel fill** The \*sediment infill of a \*channel, produced either by the accretion of sediment transported by water flowing through the channel, or by the infilling of an abandoned channel. See also MULTISTOREY SANDBODY.

**channel wave** An \*elastic wave which travels through a layer with a lower velocity than that of the surrounding layers. The wave tends to be confined to that layer because of repeated internal reflection and the refraction of escaping waves back towards the channel owing to the increasing velocities away from the channel. See also HIDDEN LAYER.

**chaos** 1. A state of disorder which is governed by simple and precise laws, but where the outcome is unpredictable and may change greatly with slight variations in starting conditions. Most real systems, such as weather patterns and satellite orbits, display chaotic behaviour. See also FRACTAL. 2. An area of jumbled or broken terrain on the surface of an extraterrestrial body.

**char** Solid carbonaceous residue of high calorific value, derived from incomplete burning of organic material. It may be formed into briquettes and burned for fuel; if pure it can be used as a filter medium. Charcoal is made from wood or bone; coke, another char, is derived from \*coal. See PYROLYSIS.

**character** Any detectable attribute or property of the \*phenotype of an organism. Defined heritable differences in the character may exist between individuals within a species.

**character states** Particular versions of a \*character. Thus, the character 'horns' may have the character states 'straight', 'curly', etc. The proper elucidation of character states and their \*polarity is one of the major concerns of \*cladistic analysis.

**chara marl** A \*marl that consists of the calcified oogonia (female sex organs) of charophytes (\*Charophyceae), together with the stems of aquatic plants coated with calcium carbonate, mixed with \*siliciclastic detritus.

**charcoal** See CHAR.

**charge 1. (explosive)** The combination of detonator and main explosive. The effective energy released is a function of the nature and weight of the main explosive material, the type of detonator used, and how the charge is fired. **2. (electrical)** A source of electric field forces; the transfer of such charge through a conducting medium is measured as electric current. Electric charge comprises whole-number multiples of electronic charge, of which the electron constitutes a negative charge.

**chargeability ( $M$ )** One of the units of measurement of *\*induced* polarization in the *\*time domain*. True chargeability is the ratio of the over- or secondary voltage,  $V_s$ , to the observed voltage,  $V_0$ , applied by way of an *\*electrode* array so that  $M = V_s/V_0$ , expressed as a percentage or as millivolts per volt; this quantity is independent of topographic effects and of electrode geometry and is thus a good measure of induced polarization. In reality, what is measured is the apparent chargeability ( $M_a$ ) which is the area ( $A$ ) beneath the voltage-time decay curve over a defined time interval ( $t_1$  to  $t_2$ ) and normalized by the supposed steady-state primary voltage,  $V_p$ , such that  $M_a = A/V_p \times \int_{t_1}^{t_2} V(t)dt$ , in units of mVs/V.

**charge-coupled device (CCD)** A light-sensitive semiconductor used to enhance images obtained from faint objects. It accumulates and temporarily stores charge at particular *\*pixel* locations when struck by photons. The pixels can then be moved, allowing them to be pieced together to form an image.

**charged-body potential method** See MISE-À-LA-MASSE METHOD.

**Charnian** A *\*stage* of the Upper *\*Proterozoic* of Charnwood Forest, England. The Charnwood sequence contains impressions of very early *\*Plantae* or *\*Animalia*.

**Charniodiscus** See EDIACARAN FOSSILS.

**charnockite** A light-coloured, medium- to coarse-grained *\*igneous* rock containing *\*quartz* and microcline feldspar (see ALKALI FELDSPAR) as major components with, in order of decreasing abundance, oligoclase feldspar (see PLAGIOCLASE FELDSPAR), *\*hypersthene*, *\*biotite*, and *\*magnetite*. Despite its low abundance, hypersthene (a calcium-poor, iron-rich *\*pyroxene*) is a distinctive feature of this rock. Rocks with a charnockite mineralogy can

also be formed by the **\*metamorphism** of quartzofeldspathic rock under dry **\*granulite** facies conditions. The type location for the rock is in Tamil Nadu, India.

**charnockitic gneiss** See GRANULITE.

**Charon** The satellite of **\*Pluto**, discovered in 1979, which orbits at a mean distance of 19 405 km, with an orbital period of 6.387 days. Its equatorial radius is 586 km; mass  $1.7 \times 10^{21}$  kg; mean density 1800 kg/m<sup>3</sup>; surface gravity 0.21 (Earth = 1); albedo 0.375. Some consider Charon and Pluto to comprise a double **\*dwarf** planet system.

**Charophyceae (charophytes)** A class (in some classifications a division: Charophyta) of **\*algae** that in some ways resemble bryophytes (**\*Bryophyta**). They occur in fresh and brackish water and their calcified fructifications are ornamented with spiral striae. Neglected by geologists until 1959, they are now used as stratigraphic markers in **\*Cenozoic** strata. See GYROGONITE.

**Charophyta** See CHAROPHYCEAE.

**Charpentier, Jean de** (1786–1855) A Swiss superintendent of mines, Charpentier made extensive field studies in the Alps. Attending a lecture by the naturalist Ignace Venetz (1788–1859) in 1821, Charpentier was persuaded that Swiss glaciers had once been much more extensive. Using evidence of erratic boulders and **\*moraines** to support this idea, he finally managed to overcome the resistance of **\*Agassiz** by taking him to visit the Aar Glacier. Agassiz then developed further the theory of a former ice age.

**chart datum** Datum, or plane, to which depth measurements on a chart are referred.

**chasma** (*pl.* **chasmata**) Originally, a very large canyon on Mars, generally of structural origin. Such structures are distinguished from smaller, sinuous channels (*valles*, see VALLIS) probably of fluvial origin, and from fossae (*see* FOSSA) which are linear depressions analogous to terrestrial **\*grabens**. The term is now used for any similar large valley on a planetary or **\*satellite** surface.

**chatoyance** See CHATOYANCY

**chatoyancy (chatoyance)** The shimmering reflectance seen in a gemstone, from the French *oeil de chat*, cat's eye.

**chattermark** Small (less than 5 mm) crescentic scar typically found on the surface of rocks, of rock particles, and of rounded beach pebbles. It is a percussion fracture, produced when particles are thrown together in wind or water environments.

**Chattian** 1. The final *\*age* in the *\*Oligocene* epoch, preceded by the *\*Rupelian* (Stampian), followed by the *\*Aquitainian*, and dated at 28.4–23.03 Ma ago (Int. Commission on Stratigraphy, 2004). 2. The name of the corresponding European *\*stage*, which is roughly contemporaneous with the upper *\*Zemorian* (N. America), upper *\*Whaingaroan*, *\*Duntroonian*, and *\*Waitakian* (New Zealand), and most of the *\*Janjukian* (Australia). It is dated by means of *\*sea-floor* spreading rates, small mammals, and *\*plankton*.

**Chautauquan** A *\*series* (370–359.2 Ma ago) in the Upper *\*Devonian* of N. America, underlain by the *\*Senecan*, overlain by the *\*Kinderhookian* (*\*Mississippian*), and comprising the Cassadagian and Bradfordian stages. It is roughly contemporaneous with the *\*Famennian* (Upper Devonian) and, possibly, the Hastarian (*\*Tournaisian*) stages of Europe.

**Chazyan** A *\*stage* (464–460.9 Ma ago) of the *\*Ordovician* in the Middle *\*Champlainian* *\*series* of N. America, preceded by the *\*Llanvirn* and followed by the *\*Ordovician V*.

**Chebotarev sequence** An idealized sequence of chemical changes in *\*groundwater*. As groundwater moves through rock its chemical composition normally changes. In general, the longer groundwater remains in contact with the *\*aquifer* rocks the greater the amount of material it will take into solution. Changes in composition also occur with increasing depth of travel, as bicarbonate *\*anions*, which dominate in many shallow groundwaters, give way to sulphate and then chloride anions, and calcium is exchanged for sodium.

***Cheirolepis trilli*** Early representative of the primitive *\*bony* fish, the *\*palaeoniscids*. It is known from the Middle *\*Devonian*.

**chelation** Equilibrium reaction between a metallic ion and an organic molecule in which more than one bond links the two components. The

metallic ion is termed the complexing agent, the chelating organic molecule the **\*ligand**. Chelation is a naturally occurring mechanism in soils, useful since it removes heavy-metal ions that are in solution in simple inorganic form, in which state they may be directly toxic to plants or may interfere with the uptake of essential nutrients. Heavy-metal toxicity will tend to be reduced by the application of organic material.

**Chelford** 1. An **\*interstadial** that occurred 65 000–60 000 years BP, during the **\*Devensian** glaciation. 2. Sections of **\*alluvial \*sands** and organic muds containing tree remains from the Chelford Interstadial, overlaid and underlaid by **\*till**, that are exposed in sand pits between Chelford and Congleton, in Cheshire, England.

**cheliceræ** See ARACHNIDA; ARTHROPODA; CHELICERATA.

**Chelicerata** (phylum **\*Arthropoda**) Subphylum comprising a diverse group of animals which possess an anterior prosoma (fused head and thorax), a posterior opisthosoma (abdomen), and a pair of jointed pincers. The subphylum includes the spiders, mites, and scorpions, as well as the king-crab (*Limulus*) and the eurypterids (**\*Palaeozoic** water scorpions).

**Cheltenhamian** A **\*stage** (5–4.3 Ma ago) in the **\*Pliocene epoch** of south-eastern Australia, underlain by the **\*Mitchellian**, overlain by the **\*Kalimnan**, and roughly contemporaneous with the mid **\*Zanclean** (Tabianian) stage.

**chemical demagnetization** The washing of permeable sediments in weak acid, usually 10% hydrochloric, in order to remove **\*cement** with its associated **\*remanent** magnetism. See DEMAGNETIZATION.

**chemical fossil** See BIOMARKER.

**chemical garden** A structure, comprising an array of buds, tubes, and branches, that forms when soluble metal salts are added to an aqueous solution, commonly of calcium silicate, containing **\*anions**, commonly of silicate, borate, phosphate, or carbonate. The colours vary according to the ingredients used. The classic demonstration is being performed on the International Space Station to determine the effect of gravity on the growth of nanotubes.

**chemical layering** See EARTH COMPOSITIONAL LAYERING.

**chemical oxygen demand (COD)** Indicator of water or effluent quality, which measures oxygen demand by chemical (as distinct from biological) means, using potassium dichromate as the oxidizing agent. \*Oxidation takes two hours and the method is thus much quicker than an assessment of biological oxygen demand (BOD), which takes five days. Since the BOD:COD ratio is fairly constant for a given effluent, COD is used more frequently than BOD for the routine monitoring of an effluent once this ratio has been determined.

**chemical potential** See WATER POTENTIAL.

**chemical remanent magnetization (crystalline remanent magnetization; CRM)** The magnetization acquired as \*ferromagnetic minerals grow through their \*blocking volume.

**chemical weathering** The action of a set of chemical processes operating at the atomic and molecular levels to break down and re-form \*rocks and \*minerals. The results of chemical weathering are frequently new substances of reduced particle size, greater plasticity, lower density, and increased volume, compared with the original materials. Some of the important processes are \*solution, \*hydration, \*hydrolysis, \*oxidation, \*reduction, and \*carbonation.

**chemocline** In a \*meromictic lake, the transition between the upper \*mixolimnion and lower \*monimolimnion layers, marked by a change from aerobic to anaerobic conditions.

**chemofossil** A \*fossil that consists only of chemicals remaining from the decomposition of a living organism.

**chemostratigraphy** The branch of \*stratigraphy in which inorganic chemical data are used to characterize and correlate strata.

**chemosymbiosis** A symbiotic association between a multicelled animal which provides a protected environment and \*bacteria that oxidize hydrogen sulphide, using the energy this releases to fix carbon and synthesize carbohydrates and enzymes which sustain the metabolism of the host.

**chemosynthesis** The pathway by which bacteria in \*hydrothermal vent communities synthesize complex organic molecules from hydrogen

sulphide gas and dissolved carbon dioxide:  $4\text{H}_2\text{S} + \text{CO}_2 + \text{O}_2 \rightarrow \text{HCHO} + 4\text{S} + 3\text{H}_2\text{O}$ . HCHO is methanal.

**chenier** Beach ridge, or sandy, linear mound that is built on a marsh area. It is at least 150 m broad, up to 3 m high, and up to 50 km long, and is typical of the Gulf Coast of America. A chenier is formed by the reworking of river-derived materials by waves. There are usually muddy, marshy zones to the front and rear of the chenier.

**chenier plain** Area of marine *\*aggradation* consisting of sandy ridges (*\*cheniers*) separated by clay-rich depressions. Occasionally the ridges may be vegetated, as on the Gulf Coast of the USA. Chenier plains may be large: that on the north coast of S. America is about 2250 km long and up to 30 km wide.

**Cheollian-1** See COMS.

**Cheremshanskian** A Russian *\*stage* (314.5–313.4 Ma ago) of the *\*Pennsylvanian* epoch, and substage of the *\*Bashkirian* stage, preceded by the *\*Yeadonian* and followed by the *\*Melekesskian*.

**chernozem (black earth)** Freely draining, dark coloured *\*soil profile* whose name is the Russian word for 'black earth'. Chernozems are associated with grassland vegetation in temperate climates, and identified by the deep and even distribution of *\*humus* and of exchangeable *\*cations* (calcium and magnesium) through the profile (now included in *\*Mollisols* in the US *\*soil taxonomy*). In the *\*World Reference Base for Soil Resources* chernozems are soils in which the A *\*soil horizon* is a *\*mollic* horizon extending to a depth of more than 20 cm.

**chert** 1. Chalcedonic (see CHALCEDONY) variety of *\*cryptocrystalline* silica,  $\text{SiO}_2$ , that occurs as nodules or irregular masses in a sedimentary environment, often in association with black *\*shales* and *\*spilites*. 2. A fine-grained rock consisting of beds of cryptocrystalline silica, usually of *\*biogenic*, volcanogenic, or diagenetic (see DIAGENESIS) origin.

**Chesterian** A *\*series* (333–318.1 Ma ago) in the *\*Mississippian* of N. America, underlain by the *\*Meramecian* and overlain by the *\*Bashkirian*. It is roughly contemporaneous with the Brigantian stage of the *\*Visean* series

and the **\*Serpukhovian** series. The Springerian is an alternative name for the upper part of the Chesterian.

**chevron fold (accordion fold, zig-zag fold)** A type of **\*fold** which shows characteristically long, planar limbs with a short, angular **\*hinge** zone. Ideal chevron folds have inter-limb angles of 60°. Chevron folds occur in sequences of regularly bedded layers of alternating **\*competent** and **\*incompetent** material which deform by **\*flexural slip** and ductile flow respectively.

**chevron marks** A type of **\*sole structure** characterized by a linear pattern of small, V-shaped ridges formed by the dragging of an object over the surface of a viscous mud. The V-shapes close in the down-current direction, enabling the chevron marks to be used as a palaeocurrent indicator. See **PALAEOCURRENT ANALYSIS**.

**Chewtonian** A **\*stage** (473–471 Ma ago) of the Lower **\*Ordovician** of Australia, underlain by the **\*Bendigonian** and overlain by the **\*Castlemainian**.

**Chezy's formula** An empirical formula relating river **\*discharge** ( $Q$ ) to **\*channel** dimensions and water surface slope.  $Q = AC\sqrt{rS}$ , where  $A$  is the cross-sectional area of the river,  $C$  is the Chezy discharge coefficient,  $r$  is the **\*hydraulic** radius, and  $S$  is the slope of the water surface. This formula is useful for extending river-flow rating curves. The formula was devised by the French hydrologist Antoine Chézy (1718–98).

**chiastolite** See **ANDALUSITE**.

**Chibis-M** A **\*microsatellite** mission of the Space Research Institute of the Russian Academy of Sciences to study processes in the **\*ionosphere** related to space weather as part of the search for universal laws governing the transformation and dissipation of plasma-wave energy in the **\*magnetosphere-ionosphere** system. The minisatellite reached the International Space Station (ISS) on 2 November 2011, and on 25 January 2012 it was separated from the ISS and transferred to its final orbit at a height of 513 km.

**chickenwire structure** A closely packed, nodular mass of **\*gypsum** or **\*anhydrite**, with thin, interconnected fingers of mud between. It is commonly found in **\*sabkha** salt deposits.

**Chicxulub** A small town on the Yucatán Peninsula, Mexico, that has been identified as lying within the crater formed by a major impact event dated at about 65 Ma ago. The crater is about 195 km wide, has a multi-ring structure, and could have been made by an object about 12 km in diameter, the impact believed to be the event associated with the mass extinction marking the end of the *\*Cretaceous* period.

**Chile Rise** The oceanic *\*ridge* which separates the *\*Nazca* Plate and the *\*Antarctic* Plate.

**Chile saltpetre** See SODA NITRE.

**chilidial plates** See NOTOTHYRIUM.

**chilidium** See NOTOTHYRIUM.

**chilled edge** See CHILLED MARGIN.

**chilled margin (chilled edge)** A fine-grained or glassy carapace found around crystalline *\*magma* bodies. The fine grain size is produced by the rapid loss of heat from the external surface of the magma body, causing severe undercooling of the magma which generates many crystal-nucleation sites producing small crystals, or suppression of nucleation sites altogether, producing glass.

**Chilopoda** See MYRIAPODA.

**China–Brazil Earth Resources Satellite (CBERS)** A cooperative programme between China and Brazil to provide high-resolution surface images, which it supplies free. The first three satellites in the series either failed or were retired. CBERS-4 (also called Ziyuan 1-04) was launched on 7 December 2014, from the Taiyuan Satellite Launch Center, China.

**china clay (kaolin)** See CHINASTONE.

**China Seismo-Electromagnetic Satellite (CSES; Zhangheng 1)** A Chinese-Italian satellite mission to monitor the Earth's electromagnetic field and waves, *\*plasma* and particles, and perturbations of the atmosphere, *\*ionosphere*, and *\*magnetosphere*. It was launched on 2 February 2018 into a Sun-synchronous orbit from the Jiuquan Satellite Launch Center, China. Zhang Heng (78–139 CE) was a polymath, astronomer, and seismologist.

**chinastone** A light-coloured, hydrothermally altered, *\*igneous* rock composed of *\*quartz*, strongly kaolinized orthoclase (see *ALKALI FELDSPAR*), fresh *\*euhedral* albite (see *PLAGIOCLASE FELDSPAR*), and minor *\*muscovite*, *\*topaz*, and *\*fluorite*. The rock is essentially a *\*granite* which is in an arrested state of alteration by the process of *\*kaolinization*. The orthoclase, which shows strong kaolinization, can be veined by secondary quartz and fluorite, both products of the kaolinization process. Superb examples are found in the St Austell granite of Cornwall, England, where the product of granite kaolinization, china clay (kaolin), is extracted for use in the ceramic, paper, pharmaceutical, and other industries. *See also* *KAOLINITE*.

**chine** Precipitous ravine found along eroding coastlines. It is typically developed in the soft *\*Mesozoic* and *\*Cenozoic* sediments of southern England. It results from the rapid incision that occurs when a stream responds by down-cutting to the rejuvenating effect of coastline retreat.

**Chinese–French Oceanography Satellite (CFOSAT)** A joint mission of the Chinese and French space agencies to monitor ocean surface winds and waves and to acquire data on related oceanic and atmospheric processes. It is planned for launch in 2018 in China into a Sun-synchronous near-circular orbit.

**chinook** Warm, dry, westerly wind of the *\*föhn* type which blows on the eastern side of the Rocky Mountains. The quick onset of the wind and the sudden large rise in temperature is associated in spring with rapid melting of the snow.

**chip sampling** *See* *SAMPLING METHODS*.

**Chiron** A *\*solar* system asteroid (No. 2060), *\*comet* (95P), or minor planet, diameter 180 km (148–208 km); approximate mass  $4 \times 10^{18}$  kg ( $2 \times 10^{18}$ – $10^{19}$  kg); rotational period 5.9 hours; orbital period 50.7 years; *\*perihelion* date 14 February 1996; perihelion distance 8.46 AU. It is in a chaotic eccentric orbit near *\*Saturn* and *\*Uranus*. It was discovered in 1977 by Charles Kowal (1940–2011).

**Chiroptera (bats)** (class *\*Mammalia*) Order comprising the only true flying mammals, possessing features parallel to those of birds, e.g. active metabolism and economy of weight. Insectivores possibly ancestral to the

bats are known from the **\*Palaeocene**. The first undoubted bats, however, are preserved in Middle **\*Eocene** deposits in both Europe and N. America. Differentiation of the modern lineages was far advanced by the **\*Eocene–\*Oligocene** transition.

**chi-squared test ( $\chi^2$ )** In statistics, a **\*hypothesis** test used to determine the goodness of fit of a particular data set with that expected from a theoretical distribution. The test statistic is a function of the difference between observed and expected values, which is compared to the chi-squared distribution. The chi-squared distribution is a distribution of sample variance based on a single parameter, the **\*degrees** of freedom.

**chitin** See SKELETAL MATERIAL.

***Chitinodendron franconianum*** A primitive allogrominid **\*foraminiferid** with an external chitinoid membrane. It is from the Upper **\*Cambrian** of Wisconsin, USA. Single-celled organisms are known from **\*Precambrian** strata, but *C. franconianum* is one of the first **\*protozoans** with an external skeleton.

**chloralgal** Applied to an association of green algae (**\*Chlorophyta**), living in sea water more saline than corals could tolerate, that forms a characteristic calcareous sediment. Compare CHLOROZOAN; FORAMOL.

**chlorinity** Measure of the chloride content, by mass, of **\*sea water**. It is defined as the amount of chlorine, in grams, in 1 kg of sea water (bromine and iodine are assumed to have been replaced by chlorine). Chlorinity and **\*salinity** are both measures of the saltiness of sea water. The relationship can be expressed mathematically, as salinity is equivalent to 1.80655 times the chlorinity. There is a constant ratio of dissolved chloride to total dissolved salts in all sea water.

**chlorite** Important group of **\*phyllosilicate** (sheet silicate) **\*minerals** with the general composition  $(\text{Mg,Fe,Al})_6[(\text{SiAl})_4\text{O}_{10}](\text{OH})_8$  and related to the **\*micas**; sp. gr. 2.6–3.3; soft and green; platy or tabular **\*habit**; occur in low-grade **\*metamorphic rocks** of **\*greenschist facies** and as an alteration product of **\*ferromagnesian** minerals in **\*igneous** rocks. The group includes specific minerals, e.g. clinocllore, delessite, penninite, and thuringite. **\*Chamosite** and greenalite are septichlorites which are chemically similar to the chlorites.

**chloritoid (ottrelite)** A member of the *\*nesosilicates* with the formula  $(\text{Fe}^{2+}, \text{Mg})(\text{Al}, \text{Fe}^{3+})\text{Al}_3\text{O}_2[\text{SiO}_4]_2(\text{OH})_4$  and an important metamorphic *\*index mineral*; sp. gr. 3.51–3.80; *\*hardness* 6.5; *\*monoclinic* or *\*triclinic*; *\*crystals* *\*tabular*, pseudo-hexagonal; dark green to black; occurs in *\*regionally metamorphosed* *\*pelites* with a high  $\text{Fe}^{3+}:\text{Fe}^{2+}$  ratio at low metamorphic grades; it develops at the same time as *\*biotite* and changes to *\*staurolite* with increasing temperature and pressure.

**Chlorophyta (chlorophytes, green algae)** Superdivision of *\*algae* which are typically green in colour. In common with land plants, green algae include chlorophylls *a* and *b* among their principal pigments, have cellulose as the main constituent of cell walls, and form food reserves of starch. Consequently it is believed that the ancestors of land plants must have belonged to this group. The organisms take many forms, ranging from unicellular to relatively complex multicellular plants. They are found today mainly in freshwater habitats and their distribution is cosmopolitan. They are known from the *\*Precambrian* onwards, and the earliest *\*eukaryotes* were probably of this taxon. Marine, lime-secreting green algae have contributed to algal *\*limestone reefs* since the *\*Cambrian*. Fossil genera include *Palaeoporella* (similar to modern *Halimeda*), and *Coelosphaeridium* (similar to modern *Acetabularia*). *See also* CHAROPHYCEAE.

**chlorozoan** Applied to an association of calcareous green algae (*\*Chlorophyta*), *\*hermatypic* corals, and molluscs (*\*Mollusca*) that lives in low latitudes, in seas where the temperature is always more than 20 °C and salinity 32–40 ‰, and produces a characteristic carbonate sediment. *Compare* CHLORALGAL; FORAMOL.

**Choanichthyes (\*Sarcopterygii)** Term used by some zoologists to group together *\*Crossopterygii* (lobe-finned fish) and *\*Dipnoi* (lungfish). Members of this group or subclass were thought to share functional lungs, external and internal nares (nostrils), and narrow-based, paired fins with fleshy lobes.

**Chokerian** *See* CHOKIERIAN.


**Chokierian (Chokerian)** A *\*stage* (325–324.5 Ma ago) of the *\*Mississippian* epoch, underlain by the *\*Arnsbergian* and overlain by the *\*Alportian*.

**Chondrichthyes** Class of vertebrate animals characterized by a cartilaginous endoskeleton, a skin covered by *\*placoid* scales, the structure of their fin rays, and the absence of a bony *\*operculum*, lungs, and swim bladder. It includes the subclasses *\*Elasmobranchii* (sharks, rays) and Holocephali (ghostfish). The group extends back to the Upper *\*Devonian*.  
*See also* CARTILAGINOUS FISH.

**chondrite** *\*Stony* meteorite; the majority of chondrites are characterized by the presence of *\*chondrules*, and they constitute about 86% of meteorite *\*falls*. The principal minerals they contain are *\*olivine*, *\*pyroxene*, *\*plagioclase* feldspar, *\*troilite*, and the iron-nickel minerals kamacite and taenite. Chondrites are grouped according to their petrological type (texture, crystal structure, etc.) into six classes. On the basis of their chemical composition they fall into five main groups: enstatite chondrites (highly reduced, iron in metallic form); high-Fe (H) chondrites; low-Fe (L) chondrites; low-Fe/low-metal (LL) chondrites (some iron in *\*silicate* form); and carbonaceous chondrites (relatively high *\*oxidation* levels, containing *\*volatiles*). C1 chondrites are often used as a chemical model for the *\*bulk* composition of the Earth, but there are notable discrepancies, e.g. in the proportions of volatile elements.

**chondrite model (chondritic Earth model)** Hypothesis that the *\*bulk* composition of the Earth is close to that of *\*carbonaceous* chondrites.

**Chondrites** With *Zoophycus*, an *\*ichnoguild* of many-branched, radial, *\*trace fossils* probably made by a worm that moved back and forth through the sediment, each branch of its burrow exploring a new area.



## Chondrites

**chondritic Earth model** See CHONDRITE MODEL.

**chondritic unfractionated reservoir (CHUR)** Undifferentiated material from which **\*chondrites** are believed to have formed and in which elemental abundances are essentially the same as those in the solar atmosphere (see METEORITIC ABUNDANCE OF ELEMENTS; SOLAR ABUNDANCE OF ELEMENTS). CHUR therefore provides a starting point for discussion of the chemical (and in particular the isotopic) evolution of the **\*solar** system.

**chondrocranium** See CRANIUM.

**chondrodite** Member of the humite group of minerals. The group includes humite, clinohumite, and norbergite; general formula is  $n\text{Mg}_2[\text{SiO}_4]\text{Mg}(\text{OH},\text{F})_2$ , with  $n = 1$  in norbergite, 2 in chondrodite, 3 in humite, and 4 in clinohumite; related to the **\*olivine** group; occurs in contact metamorphic zones (see CONTACT METAMORPHISM) of **\*limestones** and **\*skarns**.

**chondrophore** In certain genera of **\*bivalves** which possess **\*desmodont** dentition, an internal process with a depressed surface that supports the ligament.

**chondrosteans** See CHONDROSTEI.

**Chondrostei (chondrostean fish)** Group of *\*bony* fish belonging to the class *\*Actinopterygii*. Often ranked as a superorder, the chondrosteans have a partly cartilaginous skeleton (see *CARTILAGE*), a *\*heterocercal* tail, a spiracle, and an intestinal spiral valve. The group includes the extant orders *Acipenseriformes* and *Polypteriformes*, and the extinct order *Palaeonisciformes* (*\*palaeoniscids*, or ‘ancient fish’).

**chondrule** Small (0.1–2.0 mm), glassy, spherical to subspherical droplets, diagnostic of chondritic *\*meteorites*. They are thought to have been produced by the melting and sudden quenching of pre-existing silicate material. See *CHONDRITE*.

**Choquette and Pray classification** A widely used, descriptive classification of *\*porosity* types developed in *\*carbonates*. Primary porosity types are classified as: interparticle (pores between the grains); framework (pore spaces between the rigid framework of carbonate skeletons, as in a reef); shelter (pore spaces preserved when curved shell fragments or irregularly shaped grains prevent intergranular spaces from becoming completely occupied by the mud *\*matrix*); intraparticle (pores within the skeletal material which do not become filled with diagenetic (see *DIAGENESIS*) *\*cement*); and *\*fenestral* (pores within carbonate muds provided by *\*fenestrae*). Secondary porosity types, produced by diagenetic and tectonic (see *TECTONISM*) effects, are classified as: intercrystalline (pores produced by *\*dolomitization* or by preferential dissolution of mineral cements); moldic (pore spaces produced by the dissolution of grains, e.g. *\*ooids*, shell fragments); channel and *\*vug* (both produced by the dissolution of the rock by formation waters to generate linear (channel) or patchy (vug) cavities); cavern (the large-scale dissolution of the rock to yield man-sized or larger pores); and fracture (formed by tectonic stresses within the rock).

**chord** A straight line joining the ends of an arc.

**Chordata (chordates)** Large phylum comprising the animals that possess a rod of flexible tissue (*\*notochord*), which is protected in higher forms by a vertebral column (see *VERTEBRA*). The phylum includes the *\*Craniata*, the *\*Urochordata*, and the *\*Cephalochordata*. The first chordates and the

earliest **\*vertebrates** (Craniata) are both found in **\*Cambrian** rocks. See **PIKAIA**.

**chroma** One of the three variables of colour (hue and value being the other two). Chroma measures the strength, wavelength purity, or saturation of colour. See **MUNSELL COLOUR**.

**chromatid** One of the two daughter strands of a **\*chromosome** that has undergone division. Chromatids are joined together by a single centromere, usually positioned in the centre of the pair as they lie beside one another. When the centromere divides during the third stage (anaphase) of cell division (mitosis or meiosis), the sister chromatids become separate chromosomes.

**chromatography** Analytical technique for the separation of the components of complex mixtures, based on their repetitive distribution between a mobile phase (of a gas or liquid) and a stationary phase (of solids or liquid-coated solids). The distribution of the different component molecules between the two phases is dependent on the method of chromatography used (e.g. **\*gel-filtration**, or **\*ion-exchange**), and on the movement of the mobile phase (which results in the differential migration and therefore separation of the components along the stationary phase).

**chrome diopside** See **DIOPSIDE**.

**chromite** Member of the chromite group of **\*minerals** in the **\*spinel** group,  $\text{Fe}^{2+}\text{Cr}_2\text{O}_4$ , along with magnesiochromite ( $\text{MgCr}_2\text{O}_4$ ); sp. gr. 5.1; black; **\*sub-metallic** **\*lustre**; **\*crystals** **\*cubic**, but normally it is **\*massive** or a granular **\*aggregate**; occurs in **\*basic** and **\*ultrabasic** rocks. It is the primary source of chromium.

**chromosome** A protein thread composed of DNA and histone, usually associated with RNA, occurring in the nucleus of a cell. Although chromosomes are found in all animals and plants, bacteria and viruses contain structures that lack protein and contain only DNA or RNA: these are not chromosomes, though they serve a similar function. Chromosomes occur in pairs. Each species tends to have a characteristic number of pairs of chromosomes (23 in humans), found in most nucleated cells within most organisms. The presence of pairs of homologous chromosomes is referred to as the diploid state and is normal for the sexual phase of an organism.

Gametes (reproductive cells), and cells of the gametophyte (gamete-producing phase) of plants, however, have only one member of each pair in their nuclei (the haploid state).

**chron** 1. A small unit of geologic time, equivalent to the *\*chronostratigraphic unit* *\*chronozone*, usually based on *\*fossil* zonation (see *BIOCHRON*). When used formally (e.g. Gilbert Chron) the initial letter is often capitalized. 2. A single time interval of constant polarity of the *\*geomagnetic* field (*\*polarity* chron).

**chronohorizon** See *MARKER BED*.

**chronomere** A term for any *\*geologic-time* unit. See *CHRONOSTRATIGRAPHY*.

**chronometry** Measurement of time, which in a geologic sense is associated with the measurement of *\*absolute* ages.

**chronosequence** Sequence of related soils that differ in their degree of profile development because of differences in their age. Chronosequences can be found in evolving landscapes such as those produced by deglaciation, volcanic activity, wind deposits, or sedimentation.

**chronosome** A sedimentary rock unit bounded by planes defined by their time of formation.

**chronospecies (evolutionary species)** According to one view of *\*evolution* (that of *\*phyletic gradualism*), a new organism may be derived from its ancestor by a process of slow, steady evolutionary change. Conceivably, the descended organism might not be regarded as a member of the same species as its ancestor, in which case it would constitute a new species, in particular a chronospecies.

**chronostratic scale** An abbreviation for *\*chronostratigraphic scale*.

**chronostratigraphic correlation chart** A graphic display summarizing the *\*stratigraphy* of a particular area. The vertical scale represents geologic time, the horizontal scale represents distance, and the time-range and geographic extent of known *\*stratigraphic units* are plotted against these two variables. A range of information, obtained from *\*seismic sections*, *\*well log*, outcrops, etc., is shown on the chart. This may include *\*lithology*, bounding *\*unconformities* at the top or bottom of *\*depositional*

sequences (see [BASELAP](#); [DOWNLAP](#); [ONLAP](#); [TOPLAP](#)), the position of known [\\*lithostratigraphic](#) units, [\\*facies](#) changes, and the location of wells. See also [SEISMIC STRATIGRAPHY](#).

**chronostratigraphic horizon** See [MARKER BED](#).

**chronostratigraphic scale (chronostratic scale)** Ideally, a timescale in which the sequence of [\\*geologic-time](#) units and their corresponding time-rock divisions ([\\*chronostratigraphic units](#)) are defined by standard and internationally agreed reference points within [\\*boundary stratotypes](#). In practice, boundary stratotypes have been agreed only for parts of the scale and the chronostratigraphic units within it continue to be defined by [\\*biostratigraphic](#) means. See [STANDARD STRATIGRAPHIC SCALE](#); [STRATIGRAPHIC SCALE](#).

**chronostratigraphic unit (time-stratigraphic unit, time-rock unit)** The sequence of rocks formed during a discrete and specified interval of geologic time. Chronostratigraphic units are ranked, according to the length of time they record, into [\\*erathems](#) (the longest), [\\*systems](#), [\\*series](#), [\\*stages](#) (the basic working unit), and [\\*chronozones](#) (the shortest). Each unit comprises a number of units of lower rank, e.g. a system would consist of a number of series, and, similarly, a number of stages would constitute a series. All the rocks formed anywhere in the world, regardless of [\\*lithology](#) or local thickness, would be referred to the chronostratigraphic unit appropriate to their time of formation, e.g. all rocks laid down in the [\\*Cambrian](#) period belong to the Cambrian system. In the traditional [\\*stratigraphic scales](#), however, note that chronostratigraphic units, and the [\\*geologic-time](#) units to which they correspond, have been defined on the basis of a type section (see [STRATOTYPE](#)), so historically it is the chronostratigraphic unit that has determined the geologic-time unit and not vice versa. See [CHRONOSTRATIGRAPHY](#); [STRATIGRAPHIC NOMENCLATURE](#).

**chronostratigraphy** Branch of stratigraphy linked to the concept of time. In chronostratigraphy, intervals of geologic time are referred to as chronomeres. These may be of unequal duration. Intervals of geologic time are given formal names and grouped within a Chronomeric Standard hierarchy. The formal terms are: [\\*eon](#), [\\*era](#), [\\*period](#), [\\*epoch](#), [\\*age](#), and [\\*chron](#). The last four of these are the equivalent of system, series, stage, and

chronozone in the Stratomeric Standard hierarchy. The formal terms are often written with initial capital letters when accompanied by the proper names of the intervals to which they refer. Some geologists hold that the term 'chronostratigraphy' is synonymous with '\*biostratigraphy', but most agree that the two branches are separate.

**chronozone** The lowest-ranking \***chronostratigraphic unit**. The duration of a chronozone is defined by a \***type** section which may be based on a \***biostratigraphic zone** (in which case it would include all rocks laid down during that time interval regardless of \***fossil** content or rock type), or it may be defined on the basis of the time span of an existing \***lithostratigraphic** unit. With increasing knowledge, chronozone boundaries may be found not to coincide with the boundaries of the \***stages** to which they belong, and stages may more precisely be divided into substages. The \***geologic-time** unit equivalent to a chronozone is a \***chron**.

**chrysoberyl** \***Mineral**,  $\text{BeAl}_2\text{O}_4$ ; sp. gr. 3.5–3.8; \***hardness** 8.5; \***orthorhombic**; various shades of green or greenish-yellow; white \***streak**; \***vitreous lustre**; \***crystal** form \***tabular**; \***cleavage** \***prismatic** {110}; occurs in \***granites**, \***pegmatites**, and \***mica schists**, also in \***alluvial sands** and gravels. The emerald-green variety is used as a \***gemstone** (alexandrite).

**chrysocolla** A hydrated silicate of copper, with the formula  $\text{CuSiO}_2 \cdot 2\text{H}_2\text{O}$ ; sp. gr. 2.0; \***hardness** variable; greenish-blue; variable \***lustre**; normally forms from the weathering of copper-rich mineral deposits, and when sufficiently hard it can be cut and polished to make jewellery.

**Chrysophyceae** (**golden algae, golden-brown algae**) Class of predominantly unicellular \***algae** in which the chloroplasts contain large amounts of the pigment fucoxanthin, giving the algae their brown colour. Many are flagellated, having one flagellum of the tinsel type, with or without a second flagellum of the whiplash type. The silicoflagellates (20–100  $\mu\text{m}$ ) form a siliceous skeleton of rings, rods, and spines. They are marine forms with a fossil record going back to the Early \***Cretaceous**, and are valuable palaeoclimatic indicators. In other chrysophyceans, which are found mainly in freshwater habitats, only the cell walls of the cysts or resting spores contain silica.

**chrysotile** See ASBESTOS; SERPENTINE.

**ChubuSat-1** A Japanese \***microsatellite**, developed by Nagoya and Daido Universities, that carries an optical camera and infrared camera to observe the Earth's surface with a 10-m resolution. It was launched on 6 November 2014, from Yasny Cosmodrome, Russia, together with the microsatellites \***QSat-EOS**, **Hodoyshi-1**, and **Tsubame**.

**CHUR** See CHONDRITIC UNFRACTIONATED RESERVOIR.

**Churchillian orogeny** A \***Proterozoic** phase of mountain building that affected an area in what are now northern Saskatchewan and northern Manitoba, Canada. It was probably caused by the collision of a microcontinent (see **MICROPLATE**) within a subducting oceanic \***plate**, moving northwards, with a \***fore-arc**, and may have started about 1900 Ma ago and ended about 1850 Ma ago.

**Cibola Flight Experiment Satellite (CFESat)** A satellite mission of the Los Alamos National Laboratory, New Mexico, to survey portions of the VHF and UHF radio spectra, detecting and measuring impulse events that occur in a complex background. The satellite was launched on 9 March 2007, from Cape Canaveral, Florida, into a non-Sun-synchronous near-circular orbit.

**CICERO** See COMMUNITY INITIATIVE FOR CONTINUOUS EARTH REMOTE OBSERVATION.

**cilia** See CILIUM.

**cilium** (*pl.* **cilia**) Short, hair-like appendage, normally 2–10  $\mu\text{m}$  long and about 0.5  $\mu\text{m}$  in diameter, usually found in large numbers on those cells that have any at all. In certain \***protozoa**, cilia function in locomotion and/or feeding. They generate currents in the fluid surrounding the cell by beating in a co-ordinated manner.

**Cincinnatian** A \***stage** (451–443.7 Ma ago) of the Late \***Ordovician** of N. America, generally equivalent to the Middle \***Caradoc** to terminal \***Ashgill**, preceded by the \***Mohawkian** and followed by the \***Rhuddanian**.

**cinder cone** See SCORIA CONE.

**CINEMA** See CUBESAT FOR IONS, NEUTRALS, ELECTRONS & MAGNETIC FIELDS.

**cingulum** See DINOPHYCEAE.

**cinnabar** The commonest mercury *\*mineral*, HgS; sp. gr. 8.0–8.2; *\*hardness* 2.0–2.5; *\*trigonal*; scarlet-red to brownish-red; vermilion *\*streak*; *\*adamantine* *\*lustre*; *\*crystals* rhombohedral or thick, *\*tabular* plates, but can occur *\*massive* or granular; *\*cleavage* perfect *\*prismatic* {1011<sup>-</sup>} sedimentary rocks, in fractures in areas of volcanic activity, and around hot springs, associated with *\*pyrite*, stibnite, and *\*realgar*. It is the only major ore of mercury.

**CIPW norm calculation** Combination of the oxide components of an *\*igneous* rock into a set of water-free, standard, *\*mineral* compounds (termed ‘normative constituents’) according to a rigidly prescribed order that originally was thought to be the order of mineral crystallization in most *\*magmas*. The calculation, developed in the late 19th century by the American petrologists W. Cross, J. P. Iddings, L. V. Pirsson, and H. W. Washington, provides a basis for comparing and classifying rock types independently of their modal mineral assemblages. Rocks of similar chemistry may develop contrasting modal mineral assemblages if they crystallize under contrasting pressure,  $P_{\text{total}}$ , and  $P(\text{H}_2\text{O})$  conditions. Use of the CIPW norm calculation eliminates these effects, allowing a comparison based on an ideal mineral assemblage controlled only by the original magma chemistry.

**CIRAS** See CUBESAT INFRARED ATMOSPHERIC SOUNDER.

**circalittoral zone** The area of the *\*continental-shelf* sea bed that lies below the zone of periodic tidal exposure. It is approximately equivalent to the *\*sublittoral* zone. See also LITTORAL ZONE.

**circularity index** 1. The ratio between the area of an inscribed circle to the area of a circumscribed circle fitted to the outline of the crest of a crater (see CRATER 2). For fresh lunar craters, the average circularity index reaches a maximum value of about 0.85–0.90 for craters about 10 km in diameter. 2. See DRAINAGE BASIN SHAPE INDEX.

**circularity ratio** See DRAINAGE BASIN SHAPE INDEX.

**circular polarization** In optical *\*mineralogy*, the polarization that may occur if light passing through a *\*mineral* emerges with a velocity that is uniform in all directions, thus producing a spherical wave front. If the mineral splits the light into two rays, one of them may have circular polarization; if the light is unsplit, all of it may have circular polarization.

**circulation index** See ZONAL FLOW.

**circum-oral canal (ring canal)** Part of the water-vascular system in echinoderms (*\*Echinodermata*). The *\*madreporite* allows water into a canal that feeds the circum-oral canal, located close to the upper surface of *\*Aristotle's* lantern. Five radial water vessels extend from the circum-oral canal up the centre of each *\*ambulacrum* and from these vessels the *\*tube feet*, used in locomotion and respiration, occur at intervals.

**cirque (corrie, cwm)** Half-open, steep-sided hollow in a mountain region that has been or is being glaciated. Its form is due to a combination of glacial scouring (which deepens the floor and often produces a reversed long-profile gradient), and glacial erosion by basal sapping and *\*gelifraction* (which acts on the cirque's head and side walls).

**cirque glacier** A relatively small body of *\*ice*, *\*firn*, and snow, occupying an armchair-shaped hollow in bedrock. It is generally wide in relation to its length. It is actively supplied by drifting snow and therefore shows vigorous behaviour, involving rotational sliding.

**cirri** Plural of *\*cirrus*. See also CIRRIPEDIA.

**Cirripedia (cirripedes, barnacles)** (phylum *\*Arthropoda*, subphylum *\*Crustacea*) Class of crustaceans comprising the familiar barnacles that settle on rocks, submerged timbers, corals, shells, and the undersides of ships. They are entirely marine. The name literally means 'comb-foot'. After a free-swimming larval stage the bivalved cypris larva attaches itself to the substratum by means of cement secreted from glands in the first antennae. The body rotates and the thoracic appendages (usually six) are modified to form cirri (flexible feeding appendages) that point upwards or sideways through the gape in the carapace. The carapace persists in the adult barnacle as an inner mantle which is covered externally by calcareous plates. In the sessile (unstaked) barnacles (e.g. *Balanus*), the plates are

large and heavy, making the organism very well adapted to the high-energy conditions of a rocky, intertidal zone. Two orders of cirripedes have fossil records: the Thoracica and Acrothoracica. The Thoracica includes stalked forms (e.g. the modern Lepas or goose barnacle) and unstalked forms (e.g. Balanus). The stalked form is thought to be the more primitive. The ancestral form of the cirripedes probably resembled the cypris larva. The undoubted fossil record of the Thoracica began in the Upper **\*Silurian** with Cyprolepis (a stalked form), but fragments and disarticulated plates of **\*Cambrian–\*Ordovician** age have also been assigned to this group. The Acrothoracica are the smallest cirripedes. They bore into calcareous material such as shells and corals. Their distinctive burrows have been recorded as **\*trace fossils** from **\*Devonian** rocks.

**cirrocumulus** Cloud genus comprising shallow, high-level cloud, made entirely from ice crystals. The form is sheeted or layered with small-scale billows, or ripples, or spherical masses called elements, each with an apparent width of about 1°. *See also* **CLOUD CLASSIFICATION**.

**cirrostratus** Cloud genus comprising clouds made entirely from ice crystals forming a semi-transparent veil of fibrous or smooth appearance, often covering the sky, and identified by **\*halo** phenomena. When associated with a cold front, the cloud sheets have distinct edges. *See also* **CLOUD CLASSIFICATION**.

**cirrus** (*pl. cirri*) **1.** From the Latin *cirrus*, meaning a tuft or lock of hair. A cloud genus comprising high-level, banded clouds, made entirely from ice crystals, in wispy fibrous filaments aligned approximately along their line of movement. *See also* **CLOUD CLASSIFICATION**. **2.** In certain ciliate **\*protozoa**, an organelle formed by the fusion of a group of **\*cilia**, which usually functions in locomotion or feeding.

**citrine** *See* **QUARTZ**.

**clade** Term derived from the Greek *klados*, a ‘twig’ or ‘branch’. In **\*cladistics**, or phylogenetic systematics, it refers to a lineage branch that results from splitting in an earlier lineage. A split produces two distinct new **\*taxa**, each of which is represented as a clade, or branch, in a phylogenetic diagram.

**cladism** *See* **CLADISTICS**.

**cladistic analysis** The method of analysis which aims to discover **\*clades** and their interrelationships. For each taxon of the group being analysed, **\*character** states are ordered by their **\*polarity**, to find which taxa are united by most **\*derived** states. Only the sharing by two taxa of derived character states is evidence that they belong to the same clade.

**cladistics (cladism, phylogenetic systematics)** Special **\*taxonomic** system, founded by W. **\*Hennig** (1966), and applied to the study of evolutionary relationships. It proposes that common origin can be demonstrated by the shared possession of derived characters, characters in any group being either **\*primitive** or **\*derived**. In the branching diagrams (**\*cladograms**) used to portray these relationships, it is assumed that **\*cladogenesis**, or splitting of an evolutionary lineage, always creates two equal daughter **\*taxa**: the branching is **\*dichotomous**. Thus each pair of daughter taxa constitutes a **\*monophyletic** group with a common stem taxon, unique to the group, and a parent taxon always gives rise to two daughter taxa which must be given different names from each other and from the parent, so the parent species ceases to exist. A cladogram is therefore synonymous with a classification. A shortcoming of the method would seem to be that usually it takes no account of the time dimension.

**cladogenesis** In **\*cladistics**, the derivation of new taxa that occurs through the branching of ancestral lineages, each such split forming two (possibly more) equal sister taxa that are often considered taxonomically separate from the ancestral taxon, though this is no longer considered obligatory.

**cladogram** Diagram that delineates the branching sequences in an evolutionary tree.

**Cladoselache** Late **\*Devonian**, shark-like fish recorded from Europe and N. America. *Cladoselache* ranged between 0.5 and 1.2 m in length and is noted for the presence of a large ventral fin. Numerous specimens of this genus have been collected from the Cleveland Shales, Upper Devonian, in N. America.

**Cladoselachiformes** (class **\*Chondrichthyes**, subclass **\*Elasmobranchii**) Order of fossil sharks with an elongate body, the two dorsal fins each with a spine, e.g. **\*Cladoselache**. They lived from the **\*Devonian** to the **\*Carboniferous**.

**Clapeyron–Clausius equation** Equation that relates the pressure and temperature at which a **\*phase** change occurs in a closed system.  $dP/dT = \Delta H_v/T\Delta V$ , where  $P$  is the pressure,  $T$  the absolute temperature,  $\Delta H_v$  the heat absorbed per mole during the phase change, and  $\Delta V$  the change in molar volume.

**Clapeyron equation** Equation that relates the temperature ( $T$ ) at which a mineralogical **\*phase** change occurs, with associated change in volume ( $\Delta V$ ), to the pressure.  $\Delta H/dT/dp = T\Delta V$ , where the thermal gradient as a function of pressure is  $dT/dp$  (usually assumed to be **\*adiabatic** within the Earth) and  $\Delta H$  is the heat of fusion.

**clarain** See COAL LITHOTYPE.

**Clarence** A **\*series** in the Lower **\*Cretaceous** of New Zealand, 108–95 Ma ago, underlain by the **\*Taitai**, overlain by the **\*Raukumara**, and comprising the Urutawan, Motuan, and Ngaterian **\*stages**. It is roughly contemporaneous with the upper **\*Albian** and lower **\*Cenomanian**.

**Clarke, Frank Wigglesworth** (1847– 1931) As Chief Chemist to the US Geological Survey from 1884 to 1925, Clarke systematized the collection and chemical analysis of **\*minerals**, **\*rocks**, and **\*ores**, and also did important work on the composition of the Earth's **\*crust**. The first of five editions of his book *The Data of Geochemistry* was published in 1908.

**Clarke, William Branwhite** (1798–1878) A clergyman and amateur scientist, who emigrated from Britain to Australia in 1839. He is credited with the first discovery of gold in Australia (1841). He took part in the geologic investigation of the New South Wales coalfield, and worked on the sedimentary structure of that state.

**Clarke orbit** See GEOSTATIONARY ORBIT.

**Clarkfordian** A N. American **\*stage** of the **\*Palaeocene** (56–55.5 Ma ago), preceded by the **\*Tiffanian** and followed by the **\*Wasatchian**.

**classification 1.** Any scheme for structuring data that is used to group individuals. In ecological and taxonomic studies numerical classification schemes have been devised, but various hierarchical or non-hierarchical classificatory strategies have also been used. In taxonomy, the fundamental unit is the species. Among living forms species are groups of individuals

that look alike and can interbreed, but cannot interbreed with other species. In *\*palaeontology*, where breeding capability cannot be determined, species are defined according to morphological similarities. In formal nomenclature, taxonomists follow the binomial system developed by the Swedish naturalist Carolus Linnaeus (1707–78). In this system each species is defined by two names: the generic (referring to the genus) and the specific (referring to the species). Thus various related species may share a common generic name. Genera (sing. genus) may be combined with others to form families, and related families combined into an order. Orders may be combined into classes, and classes into phyla (sing. phylum) or divisions in the case of *\*Plantae* (although this level is omitted in modern angiosperm classification). For example, the brachiopods comprise some eleven orders split between two classes and these two classes are the major subdivisions of the phylum *\*Brachiopoda*. The basic groupings, the phyla, are combined together into kingdoms, e.g. *Plantae* (the plants), *\*Fungi*, and *\*Animalia* (animals), and the kingdoms into *\*domains*. Some workers have tackled the uncertainties arising from subjectivity in classification by using numerical methods. In their view, if enough characters were measured and represented by cluster statistics, the distances between clusters could be used as a measure of difference. Even so, the worker has to decide (subjectively) how best to analyse the measurements, and so objectivity is lost. Other workers emphasize those features shared by organisms that show a hierarchical pattern (see *CLADISTICS*). 2. In *\*remote* sensing, the computer-assisted recognition of surface materials. The process assigns individual *\*pixels* of an image to categories (e.g. vegetation, road) based on spectral characteristics compared to spectral characteristics of known parts of an image (training areas). Assignment of pixels is not always possible when the parameter space of different training areas overlaps. In such cases a *\*principal* component analysis prior to classification may be used to allow better separation of training areas by increasing the overall parameter space. See also *BOX CLASSIFICATION*; *MINIMUM-DISTANCE-TO-MEANS CLASSIFICATION*; *MAXIMUM-LIKELIHOOD CLASSIFICATION*.

**clast** Particle of broken-down rock. These fragments may vary in size from boulders to *\*silt-sized* *\*grains*, and are invariably the products of *\*erosion* followed by deposition in a new setting. See *CLASTIC ROCK*. See also *BIOCLAST*.

**clastic (fragmental)** Applied to the texture of fragmental *\*sedimentary rocks*.

**clastic dyke** An intrusion of *\*sediment* that cuts across the sedimentary layers.

**clastic rock** *\*Sediment* composed of fragments of pre-existing rocks (*\*clasts*). Consolidated clastic rocks include *\*conglomerates*, *\*sandstones*, and *\*shales*.

**clastogenic flow** See FIRE-FOUNTAIN.

**clathrate** A compound in which molecules of one substance, commonly a noble gas, are completely enclosed within the crystal structure of another substance. Typical examples are Kr and Xe encapsulated in *\*zeolites* structures, or Ar, Kr, and Xe trapped in water ice.

**Clausius–Clapeyron equation** See CLAPEYRON–CLAUSIUS EQUATION.

**Clavatipollenites** A miospore from the *\*Barremian* (Lower *\*Cretaceous*); one of the oldest known *\*angiosperm* *\*pollen* grains. It is oval and monosulcate (see SULCUS).

**clay 1.** In the Udden–Wentworth scale, particles less than 4  $\mu\text{m}$  in size. See PARTICLE SIZE. **2.** In pedology, a soil separate comprising mineral particles less than 2  $\mu\text{m}$  in diameter according to the Atterberg and *\*USDA* classifications. **3.** Class of soil texture, irrespective of particle diameter but usually containing at least 20% by weight of clay particles. Compare CLAY MINERALS.

**clay dune** See LUNETTE.

**C-layer** The lower part of the upper *\*mantle* of the Earth, between about 370 and 720 km depth, within which there are several strong seismic-velocity gradients thought to correspond with mineralogical phase changes.

**clay films** See CUTAN.

**clay minerals** Members of the *\*phyllosilicates* (sheet silicates) with related chemistry, all are hydrous aluminium silicates with layered structure; layers of  $[\text{SiO}_4]$  tetrahedra of composition  $[\text{Si}_4\text{O}_{10}]^{4-}$  are joined to Al-O layers (gibbsite-type layers) or (Mg,Fe)-O layers (brucite-type layers). 1:1 sheet

silicates have one Si-O layer coupled to one *\*brucite* or *\*gibbsite* layer and include the *\*serpentine* group and the *\*kaolinite* or kandite group of clays; 2:1 sheet silicates have two Si-O layers joined to one brucite or gibbsite layer and include the *\*smectite* and *\*illite* groups of clays, *\*bentonite* and *\*montmorillonite*, as well as *\*talc* and the *\*mica* group; 2:2 sheet silicates have two Si-O layers joined to two brucite or gibbsite layers and include the *\*chlorite* group. It is difficult to distinguish clay minerals by hand or under the microscope, so sophisticated techniques of *\*X-ray* diffraction and *\*scanning electron microscopy (SEM)* are used to determine the precise clay mineral under investigation.

**clay pan** See PAN.

**clayskins** See CUTAN.

**claystone** A compacted, non-fissile, fine-grained, *\*sedimentary rock* composed predominantly of clay-sized (less than 4 µm grain size) particles. Compare MUDSTONE.

**cleaning, magnetic** See DEMAGNETIZATION.

**cleaning-up trend (tunnel trend)** In the reading from a *\*wireline* log (see WELL LOGGING) an expression from a *\*gamma-ray* sonde showing a progressive upward decrease in the gamma reading, indicating a change in the clay-mineral content.

**clear-air turbulence (CAT)** The variable pattern of up- and down-draughts, or turbulence, sometimes occurring in the absence of any cloud. It is caused by strong wind shear, especially associated with *\*jet streams* in the upper *\*troposphere* and lower *\*stratosphere*. The phenomenon is significant for aircraft.

**clear ice (glaze)** Layer of transparent ice formed on objects near the ground, or on aircraft in flight, by freezing rain.

**cleat** System of joints in most coal seams, along which the coal parts. There are usually two systems at right angles, one better developed and with a more shiny surface than the other. The orientation and intensity of cleats may influence the direction of mine workings.

**cleavage** 1. In *\*minerals*, cleavage is evident when crystals split along planes of weakness inherent in the structure of their atomic lattices. Cleavage is described by an adjective, e.g. good, poor, etc., and by referring to its crystallographic direction, plane, and degree of perfection, the resulting digits being contained in braces ({} ) to distinguish them from descriptions of crystals. See **MILLER INDICES**. 2. The formation of a set of fractures along closely spaced, parallel surfaces in a rock (the term is usually applied to low-grade *\*metamorphic rocks*) by the alignment of various mineralogical and structural elements during *\*metamorphism* and deformation, e.g. in *\*slates*, where cleavage is due to a parallel arrangement of minerals. The *\*fabric* generally gives rise to a preferred direction of fracturing, broadly analogous to mineral cleavage. Rock cleavages may be divided into two groups: (a) continuous cleavages, e.g. ‘*\*slaty cleavage*’ (synonymous with *\*schistosity* and *\*foliation* in high-grade *\*metamorphic rocks*, see **METAMORPHIC GRADE**) which, with further deformation, may be superimposed and cross-cut by a secondary crenulation cleavage; (b) spaced cleavages, either crenulation or disjunctive, e.g. *\*fracture cleavage*. Crenulation cleavages form by the microfolding of a pre-existing *\*anisotropic* fabric. Disjunctive cleavages require no such primary fabric. Compare **FOLIATION**.

**cleavage refraction** The change in orientation (analogous to wave *\*refraction*) of *\*cleavage* planes as they are traced into and through layers of varying *\*lithology* and therefore differing *\*competence*. Angles between the cleavage plane and the bedding are largest where the lithology is most competent; in *\*graded* beds there is a gradational change in orientation.

**Clementine** A lunar mapping mission on behalf of *\*NASA* and the US Department of Defense, launched in January 1994. It achieved lunar orbit, completed its photographic tasks, and left orbit on 3 May, heading for asteroid *\*Geographos*. On 7 May a housekeeping computer malfunction led to the depletion of the attitude-control propellant and the abandonment of the asteroid mission. Clementine entered an Earth orbit that passed repeatedly through the *\*Van Allen belts*, from which the spacecraft continued to transmit data from its sensors.



<http://www.lpi.usra.edu/lunar/missions/clementine/>

- A joint NASA and USDD mission launched in 1994 to map the Moon.

**Cliffdenian** A *\*stage* (16.5–15 Ma ago) in the *\*Miocene* epoch of New Zealand, underlain by the *\*Altonian*, overlain by the *\*Lillburnian*, and roughly contemporaneous with the *\*Langhian* stage.

**CLIMAP** See CLIMATE–LEAF ANALYSIS MULTIVARIATE PROGRAM.

**CLIMAPP** See CLIMATE/LONG-RANGED INVESTIGATION MAPPING AND PREDICTIONS PROJECT.

**climate classification** The grouping of climates into broad types according to their shared characteristics. There are three principal approaches to the task. (a) Generic classification is based on levels of temperature and aridity as these relate to vegetation boundaries. Aridity is usually expressed as ‘effective precipitation’, which is calculated as the ratio of rainfall to temperature. Climatic types are defined by the response of flora to them. The *\*Köppen* system, with its modifications, uses this approach. (b) Classifications based on the moisture budget and ‘potential evapotranspiration’ (i.e. the maximum moisture that will be transferred from the ground to the atmosphere, provided that sufficient moisture is available) which do not rely on vegetation boundaries. The *\*Thornthwaite* system uses this approach. (c) Genetic (i.e. pertaining to its origin) classification, based on factors related to the atmospheric circulation of major winds and *\*air* masses, and on other factors that cause climate, is used in the systems of H. Flohn (1950) and A. N. *\*Strahler*.

**Climate–Leaf Analysis Multivariate Program (CLIMAP)** An approach to the estimation of mean annual temperatures in the past based on a suite of 29 characters found in the leaves of dicotyledonous plants known to have been present at the site. Compare LEAF MARGIN ANALYSIS.

**Climate/Long-ranged Investigation Mapping and Predictions Project (CLIMAPP)** An integrated project to study the climatic history of the *\*Quaternary*, conducted by a team of scientists engaged in Earth and ocean research. Since 1971 the administrative base for the project has been at Columbia University, New York.

**climatic geomorphology** That branch of *\*geomorphology* which deals with the effects of climate on geomorphological processes and consequently on the character of land-forms. It has included the identification of climatically controlled zones and has attempted to define provinces with distinctive denudational processes.

**climatic optimum** Period of highest prevailing temperatures since the last ice age, in most parts of the world about 4000–8000 years ago.

**climatic station** Place where the basic climatic elements are regularly observed and recorded.

**climatic zone** Region or zone characterized by a generally consistent climate. Climatic zones approximate to distinct latitude belts around the Earth. The principal ones are the: humid tropical; subtropical arid and semi-arid; humid temperate; boreal (northern hemisphere) or sub-arctic/sub-antarctic; and polar zones.

**Climatiiformes** One of the larger orders of fossil fish belonging to the class *\*Acanthodii*. They possessed bony jaws and body skeleton, *\*ganoid* scales, a *\*heterocercal* tail, and stout spines located before the dorsal, anal, pectoral, and pelvic fins. One of the better-known forms, *Climatius*, carried five additional pairs of fins along the belly and reached a length of only 8 cm. These ‘spiny sharks’ may actually be half-way between the sharks and the bony fish (*\*Osteichthyes*).

**climatostratigraphy** The study of geologic–climatic units, which are climatic episodes defined in *\*Quaternary* rocks. They are similar to *\*chronostratigraphic units* (i.e. they are inferential), but their boundaries are *\*diachronous*. They record the effects of climate (i.e. the type of biota, soils, etc.), but not the climate itself, and therefore complications arise when planning boundaries. It would be difficult to envisage a stratigraphic scheme which did not recognize the influence of climate in the Quaternary.

**climax trace fossil** A *\*trace fossil* produced by a member of a climax *\*community*.

**climbing dune** An irregular *\*sand \*dune* that forms on the windward side of a large obstacle, where sand accumulates.

**climbing-ripple cross-lamination** See RIPPLE-DRIFT CROSS-LAMINATION.

**climosequence** Sequence of *\*soil profiles*, usually on the same parent material, that differ from each other in their profile development because of local or site differences in climatic conditions. Climosequences can be found along mountain slopes in certain highland areas.

**cline** Gradual change in *\*gene* frequencies or *\*character* states within a *\*species*, across its geographic distribution.

**clino-** Prefix derived from the Greek *klino*, meaning 'sloping' or 'inclined'.

**clinocllore** See CHLORITE.

**clinof orm** A sloping depositional surface of a major morphological feature giving seismic expression, e.g. a *\*delta front*.

**clinohumite** See CHONDRODITE.

**clinometer** An instrument for measuring the angle between an inclined surface and the horizontal, using a pendulum or spirit level and a calibrated circular scale. A clinometer may be incorporated into a magnetic compass, as a 'compass clinometer'. Clinometers are used to measure the inclination of bedding, *\*cleavage*, *\*lineations*, *\*fault* planes, etc.

**clinopyroxene (cpx)** A group of *\*pyroxene* minerals which crystallize in the *\*monoclinic* crystal system, including the calcium-bearing clinopyroxenes and the sodium-bearing pyroxenes. Important members of the group include *\*augite*, *\*diopside*, *\*pigeonite*, and *\*aegirine*.

**clinosequence** Sequence of soils in which *\*soil-profile* development is related to the angle of slope of the soil surface. Clinosequences can be found on land-forms such as *\*escarpments* and *\*drumlins* with varying surface angles of slope.

**clinothem** Rock units generated by *\*strata* which gently *\*prograde* seawards into deeper water.

**clinozoisite** A member of the *\*epidote* group of *\*minerals*,  $\text{Ca}_2\text{Al}_3(\text{SiO}_4)_3\text{OH}$ ; sp. gr. 3.3; *\*hardness* 6.5; greyish-white to yellowish-white; crystallizes in the *\*monoclinic* system (unlike *\*zoisite* which is *\*orthorhombic*) hence the prefix 'clino-'; *\*massive*; occurs as an alteration product of *\*plagioclase* feldspars in *\*igneous* and *\*metamorphic* rocks, particularly *\*anorthosites*, where the rock is said to be 'saussuritized' when

feldspars are hydrothermally altered to zoisite or clinozoisite. See SAUSSURITIZATION.

**clint** Joint-bounded surface unit of a hard, almost horizontally bedded \*limestone. The diameter varies typically from 1 to 2 m. It may be shallowly dissected by smooth gutters, or runnels, called \*grikes. It is a repeating component of a limestone pavement (an extensive surface exposed by erosion).

**Clinton ironstone** Red, fossiliferous iron ore from the Middle \*Silurian Clinton Formation of east central USA. It often occurs in lenses and may be \*oolitic in texture. The \*ore mineral is usually \*hematite.

**clipped trace** A wave-form display in which amplitudes greater than a certain level are not shown. For example, clipped \*seismic traces have a truncated wiggle with a flat top instead of the usual rounded wiggle.

**clitter** Local term for the gently sloping spread of coarse, often angular rock debris surrounding many of the tors of Dartmoor, south-west England. It was produced when \*blockfields developed on the Dartmoor granite during the periglacial phases of the \*Pleistocene and the exposure of \*tors, but the rock debris is now stable and largely vegetated.

**clod** Compact, coherent block of soil, found *in situ* when soil is broken up by digging or ploughing. Clods are of varied sizes.

**closed form** The \*crystal faces of a \*mineral which occupy repeat positions in relation to its symmetry elements can be grouped together as forms. A closed form is one which totally encloses a space, e.g. a cube or tetrahedron. Compare OPEN FORM.

**closed system** A system that is able to exchange energy, but not matter, with its surroundings.

**close fold** A \*fold in which the inter-limb angle is between 30° and 70° (as defined by M. J. Fleuty (1964), in a classification of folds based on their degree of tightness).

**closure age** The time at which diffusion within a mineral or rock of a \*daughter product of a \*radioactive decay system becomes negligible compared to accumulation, so that the system becomes 'closed', usually due to a fall in temperature.

**closure temperature** The temperature at which a rock that is cooling ceases to exchange the products of **\*radioactive** decay with the surrounding material. It is the temperature at which the rock becomes a **\*closed system**, and since the rock henceforth retains radioactive decay products it can be dated radiometrically from the time it cooled to the closure temperature.

**Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO)** An environmental satellite operated jointly by **\*NASA** and the French Centre National d'Études Spatiales (CNES) carrying three instruments that monitor clouds and **\*aerosols** and their effects on weather, climate, and air quality 24 hours a day. It was launched together with **\*CloudSat** on 28 April 2006 into a **\*Sun-synchronous** orbit at approximately 750 km altitude, and flies in formation with four other satellites, known as the **\*A-Train**. In May 2017 the mission was extended for a further three years.



<http://www-calipso.larc.nasa.gov/>

- A joint CNES and NASA mission to study the role that clouds and atmospheric aerosols play in regulating the Earth's weather and climate.

**Cloud-Aerosol Transport System (CATS)** An investigation using a LIDAR (Light Detection And Ranging) instrument to measure atmospheric clouds and aerosols. It was launched on 10 January 2015 to the International Space Station.

**cloud amount** The proportion of the sky seen to be covered by cloud. Nowadays it is commonly measured in 'oktas' or eighths of the sky covered, but sometimes it is quoted as a percentage, or as tenths.

**cloud base** The undersurface of cloud, representing the **\*condensation** level of water droplets or ice.

**cloudburst** Term popularly applied to brief but exceptionally heavy precipitation, of shower or thunderstorm type. It occurs when cold down-draughts suppress the warm up-draughts in a **\*cumulonimbus** cloud, causing a breakdown in the convection cells that makes the cloud dissipate, releasing all of its moisture as it does so.

**cloud classification** Clouds have been classified by various systems according to form, altitude, and the physical processes generating them. The World Meteorological Organization (*International Cloud Atlas*, 2017 online) classifies 10 genera in three major groups (cumulus or heap clouds, stratus or sheet clouds, and cirrus or fibrous clouds) by criteria essentially based on cloud form. Some of the genera are subdivided according to variations in internal shape and structure, to give fifteen species. Additional or supplementary features, e.g. transparency, arrangement, and characteristics of growth, are defined by Latin names as variants and accessory types of cloud. The cloud genera, with their abbreviations, are: cirrus (Ci), cirrocumulus (Cc), cirrostratus (Cs), altocumulus (Ac), altostratus (As), nimbostratus (Ns), stratocumulus (Sc), stratus (St), cumulus (Cu), and cumulonimbus (Cb). Clouds can also be referred to, according to their composition, as water or ice clouds; combinations of these are called mixed cloud. Clouds are also classified by the average height of their bases as: high (3–8 km, including Ci, Cc, and Cs); middle (2–4 km, including Ac, As, and Ns); and low (0–2 km, including St, Sc, Cu, and Cb). **\*Cloud bases** of high and medium clouds are highest at the equator and decrease with increasing latitude.

The 2017 additions to the *International Cloud Atlas* are:

**asperitas** A supplementary cloud feature comprising large, wave-like features resembling exaggerated undulations in the cloud base, as seen from below. It occurs in association with **\*altocumulus** and **\*stratocumulus**.

**cataracta** See CATARACTAGENITUS.

**cataractagenitus (cataracta)** **\*Stratus** cloud that forms above a waterfall, where water is broken into spray.

**cauda (tail cloud)** A supplementary cloud feature comprising a tail-shaped extension from the main precipitation region of a **murus**, seen in association with a **\*cumulonimbus**.

**cavum** An **\*accessory cloud** comprising a thin veil with small cloud elements, seen in association with **\*cirrocumulus**, **\*altocumulus**, and **\*stratocumulus**.

**flamma** See FLAMMAGENITUS.

**flammagenitus (flamma)** **\*Cumulus** cloud that develops in air rising by **\*convection** above a forest fire, wildfire, or volcanic eruption.

**fluctus** A short-lived supplementary cloud feature comprising a formation resembling curls or breaking waves on the upper surface of clouds. It occurs in association with *\*cirrus*, *\*altocumulus*, *\*stratocumulus*, *\*stratus*, and *\*cumulus*.

**flumen (inflow band)** An *\*accessory cloud* feature associated with a *\*supercell* *\*cumulonimbus* marking the region where air is flowing into the cloud.

**homogenitus** A variety of *\*cirrus* cloud formed from a condensation trail that has persisted for at least 10 minutes.

**homomutatus** A variety of *\*cirrus* cloud that develops when strong winds modify the form of long-lasting cirrus **homogenitus**.

**inflow band** See **FLUMEN**.

**muris (wall cloud)** A localized, persistent, supplementary cloud feature that often develops abruptly from the base of a *\*supercell* or multicell *\*cumulonimbus*. It comprises a lowering of the base in the rain-free portion of the cloud. If the muris exhibits significant rotation it may produce *\*funnel clouds* and *\*tornadoes*.

**silva** See **SILVAGENITUS**.

**silvagenitus (silva)** *\*Stratus* cloud that forms above a forest canopy due to high *\*humidity* caused by evaporation and *\*transpiration*.

**tail cloud** See **CAUDA**.

**wall cloud** See **MURIS**.

**wedge tornado** A *\*tornado* in which the funnel is at least as wide as the distance from the ground to the cloud base.

**cloud condensation nuclei (CCN)** Small airborne particles on to which water vapour condenses to form cloud droplets; in the absence of CCN it is possible for the *\*relative humidity* to exceed 100%. The most effective CCN are 0.2–2.0 µm in diameter and are hygroscopic. *\*Hygroscopic* nuclei include a variety of substances, including dust, smoke, sulphate, and salt (NaCl). There are typically 5–6 million CCN per litre of air over land and about 1 million per litre of air over the oceans far from land.

**cloud discharge** Electrical discharge in a thunder cloud. *See also* **LIGHTNING**.

**cloud droplet** The liquid component of clouds, occurring as water droplets with an average size of 10  $\mu\text{m}$ . In a non-rain cloud the droplets are suspended at a near-constant level, because air friction approximately balances the gravitational force. *See also* [RAINDROP](#).

**CloudSat** A [\\*NASA](#) satellite that monitors clouds and their effect on weather and climate. It was launched together with [\\*CALIPSO](#) on 28 April 2006 into a [\\*Sun-synchronous](#) orbit at approximately 750 km altitude, and flies in formation with four other satellites, known as the [\\*A-Train](#). On 23 February 2018, managers lowered the satellite's orbit following the loss of one of its reaction wheels, which control its orientation. The mission continues but is no longer part of the [\\*A-train](#).



<http://cloudsat.atmos.colostate.edu/home>

- A NASA mission to further knowledge of atmospheric science.

**cloud seeding** Process of introducing nuclei, e.g. silver-iodide crystals or solid carbon dioxide (dry ice), into clouds composed of supercooled water droplets, in an attempt to induce precipitation. Dry ice introduced (at  $-80\text{ }^{\circ}\text{C}$ ) from the air into cloud lowers the air temperature so that (particularly at temperatures below  $-40\text{ }^{\circ}\text{C}$ ) some of the supercooled water droplets are converted into ice crystals which then grow by collisions with further droplets. Silver iodide (which has a crystal structure similar to that of ice), introduced from the air or ground, is the substance most commonly used in seeding: its crystals act as ice nuclei. Other substances, e.g. common salt or fine water droplets, may also be used to encourage coalescence. Natural seeding may be significant in cases where ice crystals from a high 'releaser' cloud (e.g. [\\*altostratus](#) or [\\*cirrostratus](#)) fall into a supercooled water 'spender' cloud (e.g. [\\*nimbostratus](#)) and encourage ice-crystal growth.

**cloud street** Band (or bands) of (usually) [\\*cumulus](#) cloud parallel to the wind direction in a sky otherwise more or less clear of cloud. Streets may form in an [\\*air](#) mass at a sharply demarcated convection layer, with separate thermals producing parallel streets.

**Cluster** A collaborative, multisatellite mission between the [\\*European Space Agency](#) and [\\*NASA](#) to study the interaction between the two [\\*plasma](#) environments of the [\\*solar](#) wind and Earth's [\\*magnetosphere](#). The

first mission was launched on 16 July and the second on 9 August 2000, both from Baikonur, Kazakhstan.

**cluster analysis** In statistics, the classification of observations into subsets based on a criterion of similarity.

**CMP (common mid-point)** See [COMMON DEPTH POINT](#).

**Cnidaria (Coelenterata)** Phylum comprising the sea anemones, jellyfish, and corals, which is known from the late [\\*Precambrian](#). The [\\*Ediacaran](#) fauna from southern Australia, from between 630 and 542 Ma ago, includes clearly identifiable jellyfish and their allies. Corals occur for the first time in [\\*Ordovician](#) rocks. Cnidarians are all aquatic and mostly marine. The body plan is characterized by a single internal cavity, the enteron (or gut), and the body wall is composed of two layers of cells separated by a gelatinous mass (the mesogloea). There are two basic body shapes, the sessile polyp (e.g. sea anemones, corals) or the free-swimming medusa (e.g. jellyfish). The mouth is surrounded by tentacles armed with stinging cells (nematocysts) that are unique to cnidarians. Typically, the body approaches [\\*radial symmetry](#), though some forms are [\\*bilaterally symmetrical](#). See [ANTHOZOA](#); [HYDROZOA](#); [SCYPHOZOA](#).

**CNMMN** Commission on New Minerals and Mineral Names. See [COMMISSION ON NEW MINERALS, NOMENCLATURE, AND CLASSIFICATION](#).

**CNMNC** Commission on New Minerals, Nomenclature, and Classification. See [COMMISSION ON NEW MINERALS, NOMENCLATURE, AND CLASSIFICATION](#).

**C/NOFS** See [COMMUNICATION/NAVIGATION OUTAGE FORECAST SYSTEM](#).

**co-** See [COM-](#).

**co-adaptation** Development and maintenance of advantageous genetic traits, so that mutual relationships can persist. Predator–prey and flower–pollinator relationships often exhibit examples of co-adaptation, which is an aspect of [\\*co-evolution](#).

**Coahulian** A N. American [\\*stage](#) of the Early [\\*Cretaceous](#), 145.5–127 Ma ago, preceded by the [\\*Tithonian](#) and followed by the [\\*Comanchean](#).

**coal** Carbon-rich mineral deposit formed from the remains of fossil plants. These are deposited initially as *\*peat*, but burial and increase in temperatures at depth bring about physical and chemical changes. The process of 'coalification' results in the production of coals of different ranks ('coal series'), from peat, through the bituminous coals and lignite, to anthracite. Each rank marks a reduction in the percentage of *\*volatiles* and moisture, and an increase in the percentage of carbon. They are termed 'woody' or 'humic' coals if formed from fragments of trees or bushes. If the major constituents of coal are pollen grains and/or finely divided plant debris, the term 'sapropelic coal' is used.

**coalescence** Process of cloud water droplet enlargement, as larger drops (more than 19 µm) fall and collide with smaller drops. Ice crystals in frozen cloud tops may also fall and grow by coalescence with drops below, ultimately melting into raindrops. In cumuliform cloud, turbulence may encourage this process of drop growth by the upthrust of small droplets to overtake and coalesce with other droplets. Repetition of the process can eventually produce rain-sized droplets. Factors favouring coalescent growth are high moisture content, water cloud of large vertical extent, and strong upward turbulence. *See also* BERGERON THEORY; COLLISION THEORY.

**coalification** *See* COAL.

**coal lithotype** Type of *\*coal*, the nature of which depends on original plant structure. Clarain has a semi-bright, shiny *\*lustre*, is finely laminated, with smooth or irregular fracture, and is banded parallel to bedding. Durain is grey to brownish-black, banded, dull, with a granular and rough surface; harder than vitrain and more common. Fusain is sooty black, with silky lustre; it is fibrous and friable like charcoal, and occurrences are usually thin and impersistent. Vitrain is black, with brilliant, glassy lustre, conchoidal fracture, and cubic *\*cleavage*. It is clean and structureless, and occurs in thin bands or lenses.

**coal maceral** Elementary and microscopic constituent of *\*coal*. There are a number of different types. Alginite is formed from algal remains; typical of *\*boghead* coal. Collinite derives from cell infillings, is structureless and falls within the vitrinite *\*coal-maceral group*. Cutinite consists of plant *\*cuticles*, usually the hard outer coat of the epidermal cells of leaves. Fusinite comes from woody material; it has high reflectivity and

microhardness (its individual grains are hard), well-defined cellular structure (Bogen structure), and is high in carbon. Micrinite has high reflectivity, medium hardness, is opaque, granular, with grain size less than 10 µm, and has no cell structure. Resinite is formed of small ellipsoids or spindles of resin, a cell-infilling material. Sclerotinite describes variously sized round or oval bodies of irregular structure which may have been fungi or *\*spores*. It is similar in hardness and reflectivity to fusinite. Sporinite is formed of spore *\*exines*, usually flattened parallel to stratification. Telinite is that part of the vitrinite group deriving from cell-wall material.

**coal-maceral group** One of a particular assemblage of *\*coal macerals*. Exinite (liptinite) is a group consisting of *\*spores*, *\*cuticles*, resins, and waxes, rich in hydrogen and typical of attrital coals. It includes sparain, *\*cutan*, *\*alginite*, *\*resinite*, and liptodetrinite. Inertinite is charcoal-like, the result of bacterial action or forest fires; high in carbon, relatively inert during carbonization. It includes *\*micrinite*, semifusinite, *\*fusinite*, *\*sclerotinite*, and detritinite. Vitrinite is characteristic of vitrain coal derived from *\*humus*; with medium reflectivity. It is subdivided into telinite with visible residual cell-wall material, and collinite which is structureless.

**coal measures** A *\*lithostratigraphic* term for coal-bearing rock strata of the Upper *\*Carboniferous*. The term is also used informally for coal-bearing strata of other ages (e.g. *\*Permian* in Australia).

**coal series** See COAL.

**coarsening-upward succession** A vertical change in a *\*facies* in which the grain size increases with height above the base. Compare FINING-UPWARD SUCCESSION.

**coastal onlap** The deposition and *\*onlap* of coastal non-marine and *\*littoral* deposits further and further inland, due to a relative rise in sea level. If, subsequently, relative sea level falls, the *\*base level* is lowered and *\*erosion* probably occurs at the top of the sequence. At the next relative sea level rise, coastal onlap will recommence but from a lower level. This downward shift in coastal onlap is thus an indication that there has been a relative fall in sea level. See also SEISMIC STRATIGRAPHY.

**coastal processes** The set of mechanisms that operate along a coastline, bringing about various combinations of *\*erosion* and deposition. A cliffed

coastline is affected by *\*slope processes* and by wave activity. Both agents give rise to distinctive land-forms, including the *\*geo*, the *\*bevelled* cliff, and the 'blow-hole' (a chamber with a relatively narrow exit at the top of the cliff, from which water and spray are forced when waves are driven against the coast). A low coastline is largely affected by processes in the *\*surf* zone, where most work is done by shoaling and breaking waves, and in the offshore zone, where tidal currents are the chief agents of sediment transfer.

**coastal toplap** *See* TOPLAP; SEISMIC STRATIGRAPHY.

**coated stone** *See* AGGREGATE.

**co-axial correlation** A graphic method of correlation used in the prediction of storm runoff volumes.


**cobalt glance** *See* COBALTITE.

**cobaltite (cobalt glance)** *\*Mineral*, CoAsS; sp. gr. 6.0–6.5; *\*hardness* 5–6; *\*cubic*; from white to dark grey or greyish-black, all with a reddish tinge; grey-black *\*streak*; *\*metallic* *\*lustre*; crystals octahedra, cubes, and pentagonal-dodecahedra, also occurs *\*massive*, and in irregular grains; *\*cleavage* perfect cubic; occurs mainly in *\*hydrothermal* veins and contact *\*metamorphic zones*, and is associated with *\*chalcopyrite*, *\*sphalerite*, *\*chlorite*, *\*tourmaline*, and *\*apatite*. It is one of the principal *\*ore minerals* for cobalt.

**cobble** *See* PARTICLE SIZE.

**Coble creep** A form of *\*diffusion* creep in which atoms migrate along grain boundaries. *See also* NABARRO–HERRING CREEP.

**coccolithophorids** (phylum Prymnesiophyta) A group of unicellular, marine, planktonic *\*algae* which are, at least at some stage in their life cycle, covered in calcareous plates (*\*coccoliths*) embedded in a gelatinous sheath. They are spherical or oval, and less than 20 µm in diameter. They range in age from the Upper *\*Triassic* to *\*Holocene*, although dubious examples have been described from the Upper *\*Precambrian* and *\*Palaeozoic*. Because of their abundance, with large seasonal blooms, and the preservation of coccoliths in sediments, coccolithophorids are believed to play a major role in the global carbon cycle.



**Coccolithophorid**

**coccoliths** The microscopic calcareous plates or discs, often oval and commonly intricately patterned and ornamented, that occur as part of the protective covering of a group of the unicellular *\*algae* called *\*coccolithophorids*. Coccoliths are a major component of the modern deep-sea calcareous oozes, and were especially abundant in the *\*Mesozoic*, particularly the *\*Cretaceous* period, in which they became a major component of the Chalk lithology.

**Cochiti** A normal *\*polarity subchron* which occurs within the *\*Gilbert* reversed *\*chron* (*\*Pliocene*).

**COCORP** (**C**onsortium for **C**ontinental **R**eflection **P**rofilng) A project operated from Cornell University, USA, whose objects are to map the *\*basement* rocks of the USA and to focus on specific geologic problems. In areas of interest the practice is to run long profiles, typically 50–200 km, and to record data to 18–20 seconds *\*two-way* travel time (one second of two-way travel time is equivalent to about 3 km of crystalline basement).

**Cocos Plate** One of the present-day minor lithospheric *\*plates* that lies beneath the *\*Pacific Ocean*. A remnant of the *\*Farallon Plate*, it is subducting under the *\*North American Plate* and *\*Caribbean Plate* and has constructive boundaries with the *\*Pacific Plate* (along the *\*E. Pacific Rise*) and the *\*Nazca Plate* (along the *\*Galápagos Rise*).

**COD** See CHEMICAL OXYGEN DEMAND.

**coefficient of compressibility** Amount by which a *\*stratum* of given thickness will compress under increasing load pressure. It is measured as the change in *\*void ratio* per unit increase in pressure.

**coefficient of consolidation** Factor governing the rate by which compression can occur in a particular soil. The rate and amount of compression in soils varies with the rate at which pore water is lost, and therefore depends on *\*permeability*. The coefficient is defined as: coefficient of permeability  $\times$  (1 + initial void ratio)/ coefficient of compressibility  $\times$  density of water.

**coefficient of nivosity** A measure that indicates the amount of snow melt contributing to warm-season river flows.

**coefficient of sliding friction** See ANGLE OF INTERNAL FRICTION.

**coefficient of variation** A measure of variability within a sample, representing a population or a species; it is calculated as the *\*standard deviation*  $\times$  100 divided by the mean. Experience shows that within a homogenous population of mature individuals the coefficient of variation rarely exceeds 10.

**coelacanth** See COELACANTHIFORMES.

**Coelacanthiformes** (class *\*Osteichthyes*, subclass *\*Crossopterygii*) Order of *\*bony* fish thought to have been extinct for 50 million years (since the end of the *\*Mesozoic* era) until the discovery of the coelacanth (*Latimeria chalumnae*) off S. Africa in 1938. Both fossil and living coelacanths are bulky marine fish with a diphyccercal or tri-lobed tail fin; the second dorsal, anal, pectoral, and pelvic fins are very unusual as they are supported by movable stalks or lobes. The much smaller fossil species lived from the *\*Devonian* to the *\*Cretaceous*. They were initially freshwater fish, but in the *\*Triassic* marine representatives also evolved.

**Coelenterata** Name formerly applied to a phylum comprising both **\*Cnidaria** and Ctenophora. Today, when these two groups are universally separated in different phyla, the term 'Coelenterata' is sometimes used as a synonym for Cnidaria alone.

**coelom** Principal body cavity in most animals, forming the cavity around the gut in many **\*Annelida**, and in **\*Echinodermata** and **\*Vertebrata**. In **\*Arthropoda** and **\*Mollusca**, the main body cavity is an expanded part of the blood system (a haemocoel) and the coelom is small.

**coelomate** Applied to animals possessing a coelom.

**Coelophysis** One of the first flesh-eating **\*dinosaurs**, recorded from the Late **\*Jurassic** of N. America. It was a slender, bipedal **\*coelurosaur** less than 2 m in length, weighing approximately 23 kg. It had a small skull with sharp serrated teeth, a long neck and tail, and hands with long, grasping fingers.

**Coelurosauria (coelurosaurs)** (order **\*Saurischia**, suborder **\*Theropoda**) Infra-order of carnivorous, bipedal **\*dinosaurs**, which were the most persistent of the infra-ordinal dinosaur groups, extending from **\*Triassic** to **\*Cretaceous** times. The largest measured about 3 m long, although many were much smaller.

**coelurosaurs** See COELUROSAURIA.

**Coenopteridales** See PRE-FERNS.

**coenosteum** In stromatoporoids (**\*Stromatoporoidea**), the name given to the skeleton. It is a laminated structure and the broad laminae or latilaminae seen in vertical section probably record growth periods in the organism.

**coenozone** See ASSEMBLAGE ZONE.

**coercivity, magnetic** The magnetic field (direct or alternating) required to reduce the external magnetization of a **\*ferromagnetic** substance to zero. It depends on the composition, grain size, and temperature of the substance.

**coesite** A variety of **\*quartz** (SiO<sub>2</sub>) produced at high pressures (greater than 4 GPa) and found in rocks subjected to impact by large **\*meteorites**.

**co-evolution** Complementary evolution of closely associated species. The interlocking adaptations of many flowering plants and their pollinating

insects provide some striking examples of co-evolution. In a broader sense, predator–prey relationships also involve co-evolution, with an evolutionary advance in the predator, for instance, triggering an evolutionary response in the prey. *See also* [CO-ADAPTATION](#).

**cognate lithic** *See* [LITHIC FRAGMENT](#).

**cogwheel ore** *See* [BOURNONITE](#).

**coherence** Two wave trains which are *\*in* phase. Coherence is also an indication of the similarity of two functions and is the *\*frequency-domain* analogy with *\*correlation* in the *\*time domain*.

**cohesion** Ability of particles to stick together without dependence on interparticle friction. In soils, cohesion is due to the *\*shearing strength* of the *\*cement* or film of water that separates individual grains. In *\*powder technology*, cohesion refers to the forces of attraction by which the particles are held together either by *\*compaction* or a binding substance.

**cohort** Group of individuals or *\*taxa* of the same age.

**coign** The top and bottom corners of a *\*crystal* when the crystal is held between thumb and forefinger in order to examine its symmetry elements. However, any three-dimensional ‘corner’ in a crystal, such as the corners of a cube, may be called a coign.

**co-ignimbrite breccia** *See* [LAG BRECCIA](#).

**coiling** In many univalve and bivalve molluscs (*\*Mollusca*) the shells are coiled. The condition is most noticeable among gastropods (*\*Gastropoda*) and cephalopods (*\*Cephalopoda*), where it is obvious that the shell is a hollow cone, coiled up to a greater or lesser extent. These rolled-up cones grow at the apertural end only and form a logarithmic spiral. Since the shell is a hollow cone, coiling about a vertical axis and growing at the apertural end, it is possible to generate a number of shapes. The shape of the tube in section (known as the ‘generating curve’) when expanding and coiling around and away from the vertical axis in a single plane defines a ‘planispirally coiled’ shell. If the coiling does not remain in a single plane but moves down the axis (translation), a helically coiled shell results. Where the translation down the axis is in a clockwise direction the coiling is termed ‘dextral’; where it is anticlockwise it is termed ‘sinistral’. In most

cases the coils (whorls) remain in contact during expansion and coiling, but in some cases they do not, resulting in a loosely coiled or 'disjunct' shell.

**coke** See CHAR.

**col 1.** High pass or saddle in a ridge. It may mark the line of a former stream valley or of a former glacier, and so provides evidence of an early stage in the development of the landscape. **2.** The saddle region of the atmospheric-pressure field between two high-and two low-pressure centres.

**colatitude** The latitudinal distance from the pole, i.e.  $90^\circ$  minus the latitude.

**cold front** Boundary between dense, cold air and the warmer, less dense air ahead of it, which the cold air tends to undercut as it advances. The gradient of the cold front may be steep, e.g.  $2^\circ$ . Air is forced to rise along this steep front, commonly triggering the development of \*cumuliform cloud, often with precipitation. Weather changes occur with the passage of a cold front, sometimes including a pronounced temperature fall, a rise in pressure, and a change in wind direction. The passage of the front commonly brings clearer, brighter weather, but the unstable air in the \*cold sector can produce showers which may be heavy, with thunder and lightning.

**cold-front clearance** Clearing of the sky after the passage of a \*cold front, the cold air displacing the warm, moist air of the warm sector as a depression moves away. The process is usually marked by a veering of the wind (in the northern hemisphere, backing in the southern hemisphere), rising pressure, and a fall in temperature. The clearing of the cloud may be due to the subsidence of air in the \*cold sector behind the front.

**cold glacier** See GLACIER.

**cold low (cold pool)** Basically non-frontal depression, comprising cold air surrounded by warmer air at a higher pressure. It is typically associated with circulation in cold \*air masses in the mid-\*troposphere over north-eastern USA and north-eastern Siberia, though sometimes occurring also over the oceans in air that has emerged from the arctic, and it is often persistent. Such depressions are marked by more or less concentric isotherms around the core of the low. Such lows may originate along the arctic coast as a result of strong vertical uplift and adiabatic cooling in \*occlusions; they do not necessarily influence surface weather conditions,

but when they occur over warmer surfaces in middle latitudes, convection develops strongly. *See also* [CUT-OFF LOW](#).

**cold pole** The area or point in each hemisphere that has the lowest mean temperatures. Verkhoyansk in north-eastern Siberia is in the region of one northern-hemisphere cold pole, with an annual mean of  $-17.2$  °C and an absolute minimum of  $-67$  °C. There is a second northern-hemisphere cold pole at Snag, Yukon, where the annual average temperature is  $-5.8$  °C and  $-63$  °C (the lowest temperature ever recorded in N. America) was registered in February 1947. The southern-hemisphere cold pole is at Vostok Station, Antarctica, where the average annual temperature is  $-55.1$  °C and where  $-89.2$  °C (the lowest temperature ever recorded on the Earth's surface) was measured on 21 July 1983.

**cold pool** *See* [COLD LOW](#).

**cold sector** Zone of colder air surrounding the narrowing wedge of warm air, the warm sector, in a developing depression. At the [\\*occlusion](#) stage the whole of the surface layer forms a cold sector, with the warm air lifted off the surface.

**cold seep** Hypersaline brine or a mixture of hydrocarbons that seeps from the ocean floor in a manner similar to that of a [\\*hydrothermal vent](#), but at about the ambient temperature. Cold seeps are rich in methane and support large and diverse communities sustained by the primary productivity of chemosynthetic bacteria; methane is taken up by the gills of the dominant mussels (*Bathymodiolus*), and is oxidized by intracellular bacteria to release energy.

**cold wave** The conditions associated with air of continental polar origin, often dominated by an [\\*anticyclone](#) behind a [\\*cold](#) front, that moves south into central and eastern parts of the USA. Cold waves are defined as a fall of  $11$  °C or more to a minimum base ( $-18$  °C in northern, central, and northeastern regions) within a 24-hour period. In southern states (Florida, California, and the Gulf Coast) the minimum fall is  $9$  °C and the base minimum  $0$  °C.

**cold working** A mechanism of low-temperature deformation which involves the development of intracrystalline distortions by bending and

twin-gliding. Cold working typically produces considerable strain-hardening.

**colemanite** A hydrated calcium borate, with the formula  $\text{Ca}_2\text{B}_6\text{O}_{11}\cdot 5\text{H}_2\text{O}$ ; sp. gr. 2.4; \*hardness 4.5; \*monoclinic; crystals short, \*prismatic, also \*massive; occurs in saline lake deposits in arid regions.

**coliform bacteria** Rod-shaped, Gram-negative \*bacteria that occur widely in soil, on vegetation, and in water. They are abundant in the faeces of birds and mammals.

**coliform count** A count made of the numbers of \*coliform bacteria present as part of most standard analyses of water intended for potable use. The number of organisms present is normally expressed per 100 ml of water.

**collagen** Fibrous scleroprotein of high tensile strength which is a major constituent of connective tissue and the organic material in \*bone. It may represent up to 6% of the total body weight. When boiled, collagen yields gelatin.

**colles** An area of small, clearly defined hills on the surface of an extraterrestrial body.

**collinite** See COAL MACERAL.

**collision theory** Theory to account for the growth of water droplets in cloud to produce raindrops, based on the mechanisms of collision, coalescence, and 'sweeping'. It holds that larger drops, with terminal velocities increasing in proportion to their diameter, fall faster than smaller drops, and collide with them. The probability of collision depends on the spacing of the drops in the cloud (i.e. on the mean free path) and on the relative sizes of droplets. For example, if some drops are up to 50  $\mu\text{m}$  diameter in cloud consisting mainly of droplets smaller than this, collisions can be frequent. Such collisions can lead to coalescence, and an overall increase in size to produce particles of raindrop size. 'Sweeping' is an ancillary process whereby small drops that are swept into the rear of larger drops may be absorbed. These mechanisms are believed to be entirely responsible for rainfall from tropical convection cloud, as well as playing a part in other clouds, including those of mid-latitudes. See also BERGERON THEORY.

**collision zone** A type of *\*convergent* margin in which two continents or *\*island* arcs have collided. The zone may be marked by young fold mountains, a suture, *\*ophiolites*, scattered, mostly shallow *\*earthquakes*, and *\*faults* with relatively short lengths of *\*outcrop*.

**colloform banding** A texture, often found in certain types of *\*mineral* deposits, where *\*crystals* have grown in a radiating and concentric manner which may reflect underlying geochemical controls. Lead–zinc deposits of sedimentary origin often show colloform banding of *\*pyrite* and *\*sphalerite* (e.g. at Silvermines, Ireland).

**colloid** 1. Substance composed of two homogeneous phases, one of which is dispersed in the other. 2. Soil colloids are substances of very small particle size, either mineral (such as *\*clay*) or organic (such as *\*humus*), which therefore have a large surface area per unit volume. Colloids usually provide surfaces with high *\*cation* exchange capacity, and also exhibit an instability controlled by soil chemistry.

**collophane** The name often given to a *\*cryptocrystalline* variety of *\*apatite* having the same formula ( $\text{Ca}_5(\text{PO}_4)_3(\text{OH},\text{F},\text{Cl})$ ), which commonly occurs in phosphatic sediments and as the principal mineral in fossilized bone and fish scales.

**colluvial** Applied to weathered rock debris that has moved down a hill slope either by *\*creep* or by surface wash.

**colonization window** The period of time during which organisms are able to colonize a substrate in which *\*trace fossils* will be left.

**colonnade** Sets of regular columns that form the lower part of a thick lava flow showing *\*columnar* *\*jointing* and lie beneath an *\*entablature*.

**Colorado Student Space Weather Experiment (CSSWE)** A 3-unit *\*CubeSat* mission by students at the University of Colorado to investigate the relationship between solar flares and energetic particles. It was launched on 13 September 2012.

**colorimetric analysis** A method of chemical analysis in which reagents are added to a rock solution to form coloured compounds with specific elements. The intensity of the colour, measured on a *\*spectrophotometer*, is proportional to the concentration of the element.


**colour** See PIXEL COLOUR.

**colour index** The total volume per cent of \*mafic (coloured) \*silicate \*minerals in an \*igneous rock. The mafic minerals would include the \*olivine group, \*pyroxene group, and \*amphibole group of minerals, as well as \*biotite. Colour index is one of many methods used for classifying igneous rocks. If a rock has 60% \*modal mafic minerals it has a colour index of 60. Rocks with colour indices between 50 and 90 are termed ‘\*mafic’ rocks (e.g. \*basalts).

**Columbia** See NUNA.

**columbite (niobite, tantalite)** A \*mineral oxide  $(\text{Fe,Mn})(\text{Ta,Nb})_2\text{O}_6$ ; when  $\text{Ta} > \text{Nb}$  the mineral is called tantalite, but when  $\text{Ta} < \text{Nb}$  it is called columbite; sp. gr. 5.2–6.5, increasing with increasing Ta content; \*hardness 6.0–6.5; \*orthorhombic; weakly magnetic; black or brownish-black; very dark red \*streak; \*sub-metallic \*lustre; \*crystals normally \*tabular and short \*prismatic forms, and can also be blade-like; \*cleavage pinacoidal; occurs in \*granites and \*pegmatites in association with high-temperature minerals, e.g. \*cassiterite, \*wolframite, \*tourmaline, \*quartz, and \*feldspar. It is a major \*ore mineral of niobium.

**columella** 1. In corals, a rod-like structure formed from the swollen axial end of the counter-\*septum that may occur as a central structure within the skeleton. 2. In gastropods (\*Gastropoda), a spiral, rod-like structure formed by the fusion of the inner shell surfaces of the whorls where \*coiling around the vertical axis is very tight.



### **Columella**

**columnal** One of the plates that form the stem (stalk) of crinoids (*\*Crinoidea*). Commonly cylindrical, columnals have a variety of forms and may produce an articulated stem of variable length. *See also* OSSICLE.

**columnar** Applied to the *\*habit* or shape of a *\*crystal* that is rod-like, with parallel faces, giving it the appearance of a column. Rocks such as *\*basalts* and *\*dolerites* may cool and develop a polygonal system of *\*joints*, usually at right angles to the cooling surfaces of the *\*igneous* body; the joints are well exposed by *\*weathering*. This feature is known as 'columnar jointing'.

**columnar joint** *See* COLUMNAR; JOINT.

**columnar section** A simplified graphic representation of the succession of rocks in a particular area, shown in the form of a column with the oldest rocks at the bottom. Each rock unit is distinguished by an appropriate *\*lithologic* symbol, and the thicknesses of the units are drawn to scale. Columnar sections from across a region are often shown side by side to demonstrate lateral variations in *\*lithology* and *\*stratigraphy*.

**com-** (**co-**, **con-**) From the Latin *cum* or *com-*, meaning ‘with’, a prefix meaning ‘with’ or ‘jointly’. ‘Com-’ is generally used before words beginning with b, m, and p, and sometimes before words beginning with vowels; ‘con-’ before words beginning with other consonants; and ‘co-’ before most words beginning with a vowel and before words beginning with h, gn, l, and r.

**COM** See COMMISSION ON ORE MINERALOGY.

**coma** The diffuse shell of gas, typically about 150 000 km in diameter, which surrounds the nucleus of a **\*comet**. The coma and the nucleus form the ‘head’ of the comet.

**Comanchean** A **\*series** (127–99.6 Ma ago) in the N. American Lower **\*Cretaceous**, underlain by the **\*Coahulian**, followed by the **\*Cenomanian**, and containing the **\*Aptian** and **\*Albian**.

**comber** Deep-water wave that has a breaking crest blown forward by a strong wind. The term is also applied to a long-period **\*spilling** breaker.

**combe rock** See HEAD.

**combination trap** Oil, gas, or water trap combining structural and stratigraphic features. See also STRUCTURAL TRAP; STRATIGRAPHIC TRAP.

**combined plate margin** A **\*plate margin** which has some combination of convergence or divergence with tangential movement, e.g. a **\*leaky** transform fault.

**Comely epoch** See CAERFAI EPOCH.

**comet** A small body composed of meteoric dust and frozen ices (H<sub>2</sub>O, CO<sub>2</sub>, CO, HCHO) in a highly elliptical or parabolic orbit around the Sun. The average **\*perihelion** distance is less than 1 AU, and the average **\*aphelion** distance is about 10<sup>4</sup> AU. Comets are derived from the **\*Oort** cloud and have average lifetimes of about 100 passages. Comet nuclei are irregular in shape, a few kilometres in diameter and have a low density (100–400 kg/m<sup>3</sup>). Due to solar radiation, they emit gas and dust, forming the characteristic tail, when within a few astronomical units of the Sun; the dust composition appears to resemble that of primitive **\*carbonaceous** chondrites.

**Comet Nucleus Tour** A \*NASA mission to map the cores of comets and analyse dust streaming from them. It was launched in July 2002, but broke apart shortly after launch and the mission was lost.

**comfort zone** Range of atmospheric temperature and humidity characteristics within which the human body feels and works comfortably and efficiently. The typical zone is delimited by a temperature range of 19–24 °C. Beyond the limits corrective adaptations are necessary to produce bodily comfort.

**comminution** The liberation of valuable \*minerals from their \*ores by crushing and grinding the ore to a particular grain size so that the residue is a mix of relatively clean particles of ore minerals and waste. Comminution is carried out initially to make excavated material easier to handle and of a specific size. Comminution theory is concerned with the relationships between energy input and particle size from a given feed size.

**Commission on Classification of Minerals** See COMMISSION ON NEW MINERALS, NOMENCLATURE, AND CLASSIFICATION.

**Commission on New Minerals and Mineral Names (CNMMN)** See COMMISSION ON NEW MINERALS, NOMENCLATURE, AND CLASSIFICATION.

**Commission on New Minerals, Nomenclature, and Classification (CNMNC)** A body formed in July 2006 and charged with developing coherent guidelines for mineral nomenclature. The CNMNC resulted from the merger of the International Mineralogical Association (IMA), the Commission on New Minerals and Mineral Names (CNMMN), and the Commission on Classification of Minerals. The CNMMN was established in 1959 with the purpose of regulating the introduction of new minerals and mineral names, and of rationalizing mineralogical nomenclature. The CNMNC consists of representatives appointed by national mineralogical bodies, and an executive committee comprising a chair, two vice-chairs, and a secretary.

**Commission on Ore Mineralogy (COM)** An organization formed in 1962 by the International Mineralogical Association that provides a short course on ore mineralogy and collects, evaluates, and publishes quantitative data on ore minerals.



<http://projects.gtk.fi/com/association/history.html>

- Forty years of COM.

**commissure** A line or plane of junction, e.g. between the two valves of the shell in a brachiopod (*\*Brachiopoda*) or bivalve (*\*Bivalvia*).

**common canal** In graptoloids (*\*Graptoloidea*), a passage into which all the *\*thecae* open (the thecae being short tubes arranged in overlapping series along the *\*stipe*).

**common depth point (CDP)** In multichannel *\*reflection* profiling, the unique point on an individual reflector from which seismic reflection information is recorded in traces at different *\*offsets*. A set of traces containing information for one CDP is called a ‘CDP gather’, and the midway position between the shots and their respective detectors for a CDP gather is called the ‘common mid-point’ (*CMP*). *See also* [FOLD](#).

**common-depth-point stack (CDP stack, horizontal stack)** The sum of the traces of seismic *\*reflection* data, acquired from a *\*roll-along* survey, which correspond to the same *\*common depth point* but which originate from different seismic *\*profiles* and different *\*offsets*. The technique reduces substantially the amplitude of incoherent noise, *\*multiples* with their different normal *\*moveouts*, diffractions, etc., compared with the seismic reflections.

**common lead** Lead which differs from primeval lead by the addition of *\*radiogenic* lead resulting from the decay of uranium and thorium (*see* [DECAY SERIES](#)), but which occurs in minerals whose U:Pb and Th:Pb ratios are so low that its isotopic composition does not change appreciably with time. *See also* [ANOMALOUS LEAD](#).

**common mid-point (CMP)** *See* [COMMON DEPTH POINT](#).

**common strontium** *See* [INITIAL STRONTIUM RATIO](#); [RUBIDIUM–STRONTIUM DATING](#).

**Communication/Navigation Outage Forecast System (C/NOFS)** A US space mission, launched on 16 April 2008, that ended on 15 December 2015, when the satellite burned up in the atmosphere during a planned re-

entry. The mission studied the **\*ionosphere**, yielding data that enable improvements to models used to predict satellite trajectories, orbital drag, and uncontrolled re-entry.

**Communication, Ocean and Meteorological Satellite (COMS, Cheollian-1, GEO-KOMPSAT-1)** A S. Korean **\*geostationary** meteorological satellite programme to verify already-developed communications technology, monitor ocean colour, and provide meteorological observations. ‘Cheollian’ means ‘thousand li view’, i.e. long distance. The satellite was launched into **\*geostationary orbit** on 26 June 2010, from Kourou, French Guiana. Further GEO-KOMPSAT missions are planned to continue the programme.

**community** In ecology, a general term applied to any grouping of populations of different organisms found living together in a particular environment; essentially, the **\*biotic** component of an **\*ecosystem**. The organisms interact (by competition, predation, mutualism, etc.) and give the community a structure. Globally, the climax communities characteristic of particular regional climates are called ‘biomes’.

**Community Initiative for Continuous Earth Remote Observation (CICERO)** A constellation of 24 6-unit **\*CubeSats** in low Earth orbit that provide radio occultation data to measure atmospheric temperature, humidity, density, and pressure, as well as ocean and ice properties, in order to contribute to climate research, improve weather forecasting, and contribute to the study of disruptive storms in the **\*ionosphere** (space weather). Launches began on 23 June 2017 from the Satish Dhawan Space Centre, India.



<https://digitalcommons.usu.edu/cgi/viewcontent.cgi?article=2936&context=smallsat>

- Cicero – A Distributed Small Satellite Radio Occultation Pathfinder Mission.

**compaction** Physical process that reflects the increase in pressures brought upon sediments as a result of deeper and deeper burial. The individual grains or particles of the sediment are packed closer and closer together, thus reducing pore space. Pore waters are forced out and ultimately

compaction will result in grain-to-grain suturing. *See also* [PRESSURE SOLUTION](#).

**compaction test** A field trial to assess the type of equipment required to compact soil, the number of passes needed, and the thickness and moisture content of the soil layer, in order to achieve a high soil density for engineering works.

**COMPASS-1** An imaging project developed by eight students at the University of Applied Science, Aachen, Germany, employing a [\\*picosatellite](#) comprising a standard [\\*CubeSat](#). It was launched into a Sun-synchronous near-circular orbit on 28 April 2008 and retired on 3 March 2012.

**COMPASS-2** *See* [COMPLEX ORBITAL MAGNETO-PLASMA AUTONOMOUS SMALL SATELLITE-2](#).

**compass clinometer** *See* [CLINOMETER](#).

**compensation level** *See* [PRATT MODEL](#).

**competence** **1.** The maximum size of rock particle transported by a particular flow of ice, water, or air. Ice has a high competence, because of its viscosity; flowing water has a lower competence, though this increases sharply as velocity increases. Wind has the least competence. **2.** The relative rheological (*see* [RHEOLOGY](#)) properties of different rock types that are adjacent to one another. Typically, a competent rock is more viscous than an incompetent rock, more prone to [\\*fracture](#), and maintains its thickness on deformation. An incompetent rock is usually more ductile than a competent rock and therefore flows more easily.

**competent rock** *See* [COMPETENCE](#).

**complex** *See* [LITHOSTRATIGRAPHIC UNIT](#).

**complexation** The formation in solution of complexes of minerals and organic matter carrying opposite charges.

**Complex Orbital Magneto-Plasma Autonomous Small Satellite-2 (COMPASS-2)** A Russian [\\*microsatellite](#) mission to develop coordinated monitoring methods to observe pre-earthquake phenomena, monitor man-

made catastrophes, and study the dynamic coupling of the atmosphere, *\*ionosphere*, and *\*magnetosphere*. It was launched on 26 May 2006, from the submarine *Ekaterinburg* in the Barents Sea, and terminated in July 2007.

**complex twins (compound twins)** A twin *\*crystal* in which twinning has taken place on two or more *\*twin laws*.

**component-stratotype** One of the individual sections of *\*strata* constituting a *\*composite-stratotype*.

**composite fault-line scarp** See SCARP.

**composite intrusion** An intrusion consisting of two or more *\*igneous* rocks of different composition.

**composite-stratotype** A *\*stratotype* (type section) comprising more than one designated standard section of *\*strata*. Such a designation may be made: (a) if an entire *\*lithostratigraphic* unit is not completely exposed as a discrete section at any one locality, in which case only one section can be the *\*holostratotype*, the other or others being subsidiary *\*parastratotypes*; or (b) in the case of *\*chronostratigraphic units*, when those of higher rank (e.g. *\*systems*) may be composite-stratotypes of constituent lower-rank stratotypes, (e.g. *\*series*).

**composite volcano** See STRATO-VOLCANO.

**compositional immaturity** See COMPOSITIONAL MATURITY.

**compositional maturity** The condition of a *\*siliciclastic \*sediment* that consists almost wholly of *\*quartz* and *\*clay minerals*, which are the end products of *\*weathering*. A sediment containing large amounts of rock fragments and *\*silicates* that have not decomposed is said to be compositionally immature.

**composition plane (composition surface)** The plane along which two halves of a twinned *\*crystal* are joined (see CRYSTAL TWINNING). It may or may not be coincident with the reflection or twin plane.

**composition surface** See COMPOSITION PLANE.

**compound corals** Colonial corals that form a compound *\*corallum* (skeleton). Where the *\*corallites* are packed so closely as to be polygonal in section the corallum is said to be 'massive'. Other shapes are described as *\*dendroid*, *\*cateniform*, *\*fasciculate*, phaceloid, *\*cerioid*, and *\*astraeoid*.

**compound twins** See COMPLEX TWINS.

**compressed air** Air under pressure. This may be used in tunnels to balance the pressure of water in the ground and so control *\*groundwater*. Compressed air is also widely used to drive machinery in mining and tunnelling operations, e.g. in rock *\*drills* and ventilation fans.

**compressibility ( $\beta$ )** Ability of a substance to decrease in specific volume and increase in density under pressure; the reciprocal of *\*bulk modulus*. In *\*powder technology*, the ratio of the volumes of loose to compact powder.

**compressional wave** See P-WAVE.

**compressive stress** The stress produced by forces acting on a unit area of rock, which can be resolved into a *\*normal stress* and a *\*shear stress*, acting perpendicularly and parallel to a plane respectively. Compressive stress is measured in pascals (Pa) or kilobars (kb).

**COMS** See COMMUNICATION, OCEAN AND METEOROLOGICAL SATELLITE.

**con-** See COM-.

**concealed coalfield** Coal deposits buried beneath younger strata, e.g. the Kent coalfield, south-eastern England.

**concentration** 1. In chemistry, the number of molecules or *\*ions* in a given volume of a substance, expressed as *\*moles* of solute per litre of solution (molarity). 2. In mineral processing, the production of a concentrate from its *\*ore*, or the process of increasing concentration by evaporation, etc.

**concentration factor** The amount by which an element must be increased above its normal *\*crustal* abundance in an *\*ore* to make it commercially extractable.

**concentration-Lagerstätte** See LAGERSTÄTTE.

**concentric fold** A *\*parallel fold* in which individual layers maintain a uniform *\*orthogonal* thickness and ideal fold surfaces are arcs of circles

sharing a common centre. Due to geometric constraints, the fold profile must change continuously upwards and downwards. *See also* [FOLD](#).

**conchoidal fracture** Curved fracture pattern characteristic of some siliceous minerals, e.g. [\\*quartz](#), fine-grained [\\*igneous \\*rocks](#), and volcanic glasses, especially [\\*obsidian](#).

**concordant** **1.** Applied to an [\\*igneous \\*intrusion](#) that has been emplaced parallel with the structure ([\\*bedding](#), [\\*foliation](#), etc.) of the invaded [\\*country](#) rock. [\\*Sills](#) are examples of concordant intrusions. **2.** Applied to a relationship in which adjacent [\\*strata](#) are structurally concurrent with, or parallel to, each other. *Compare* [DISCORDANT](#). *Contrast* [CONFORMABLE](#).

**concordant age** If the [\\*minerals](#) in a rock have remained geochemically undisturbed since their formation, it is possible for the ages obtained by different radiometric methods (for example, the decay of  $^{40}\text{K}$  to  $^{40}\text{Ar}$ ,  $^{87}\text{Rb}$  to  $^{87}\text{Sr}$ ,  $^{238}\text{U}$  to  $^{206}\text{Pb}$ ,  $^{235}\text{U}$  to  $^{207}\text{Pb}$ , and  $^{232}\text{Th}$  to  $^{208}\text{Pb}$ ) to agree within the levels imposed by the accuracy of their known half-lives (*see* [DECAY CONSTANT](#)). If all the methods employed give a reasonable level of agreement, then the age is concordant. *See* [RADIOMETRIC DATING](#).

**Concordia diagram** A plot of  $^{206}\text{Pb}:^{238}\text{U}$  against  $^{207}\text{Pb}:^{235}\text{U}$  for concordant samples of various ages should define a single curve, named 'Concordia' by G. W. Wetherill (1956). If the ratios plotted for samples fall below this [\\*concordant age](#) pattern expected for the rock body, this produces a [\\*discordant age](#) pattern. A straight line drawn through two or more such points should intersect the Concordia curve at two points. One of these will give the true age of the rock, while the younger intersect should give the date of any [\\*lead loss](#) (which is what causes the discrepancy).

**concrete** Building material composed of [\\*cement](#), [\\*aggregate](#), and water in varying proportions according to use; when mixed together the material hardens to a rock-like consistency.

**concrete dam** Dam made of Portland cement or cyclopean concrete (i.e. with large stones). A dense material is important in its construction to reduce percolation and ensure permanence; a relatively dry mix with careful tamping is required.

**concrete minimum temperature** An alternative to the *\*grass* minimum temperature, producing more uniform results, and relevant to the likely freezing on road surfaces, etc. The temperature is recorded by a standard thermometer exposed in contact with a concrete surface.

**concretion** **1.** Roughly spherical or ellipsoidal body, produced as a result of local early *\*cementation* within a sediment. It is often found with a *\*fossil* as a 'nucleus'. The size ranges from approximately 1 mm to more than 1 m, and concretions are generally monomineralic. **2.** In pedology, the localized concentration of material, e.g. calcium carbonate or iron oxide, in the form of a nodule; such nodules are of various sizes, shapes, and colours.

**concretionary** See NODULAR.

**concurrent range zone (overlap zone)** A body of *\*strata* which is characterized by the overlapping stratigraphic range of two or more *\*taxa*, selected as diagnostic, and after which the zone is named. Concurrent range zones are very widely used in the time *\*correlation* of strata. **2.** See OPPEL ZONE. Compare ASSEMBLAGE ZONE; PARTIAL RANGE ZONE.

**condensation level** The atmospheric level at which condensation occurs as a result of convection, the lifting of air (e.g. *\*orographic* lifting), or vertical mixing. See also LIFTING CONDENSATION LEVEL; CONVECTIVE CONDENSATION LEVEL.

**condensation nucleus** Small particle of an atmospheric impurity (e.g. salt, dust, or smoke), which provides surfaces for the condensation of water. Condensation nuclei vary in size from about 0.1  $\mu\text{m}$  to more than 3  $\mu\text{m}$ . Some nuclei, e.g. salt and acid particles, can encourage condensation at a relative humidity well below 100%. See also AITKEN NUCLEUS; HYGROSCOPIC NUCLEI.

**condensation trail (contrail)** Broadening trail of water droplets or ice particles behind an aircraft, caused by the release of water vapour in the engine exhaust and its condensation or deposition as it enters cold air. See also DISSIPATION TRAIL; HOMOGENITUS; HOMOMUTATUS.

**condensed bed** A series of *\*strata* which show a much thinner succession than their lateral equivalents due to greatly reduced sedimentation rates. A condensed bed was formerly known as 'remanié beds'.

**condenser** In microscopy, a hemispherical lens or series of lenses which helps to direct the optimum amount of light through varying sizes of aperture in objectives of various magnifications. Condensers are usually placed on the microscope below the *\*stage*, between the *\*polarizer* and the *\*mineral* specimen. In the control of illumination, the area of light from the illuminator is normally cut by the condenser to equal the field of view, thus minimizing glare due to interference from marginal light.

**conditional instability** Atmospheric condition in which the *\*environmental lapse rate* (ELR) is greater than the *\*saturated adiabatic lapse rate* (SALR) but lower than the *\*dry* adiabatic lapse rate (DALR), and stable, saturated air (i.e. air at the same temperature and density as the surrounding air, and so possessing neutral *\*buoyancy*) is forced to rise, e.g. to cross a mountain barrier. As the air rises it cools at the SALR, but since this is less than the ELR the rising air is warmer than the surrounding air at every height and, therefore, more buoyant, and it continues to rise. It is then unstable, its *\*instability* being conditional on the air being forced to rise. The air ceases to rise when condensation has removed sufficient moisture for it to cease to be saturated, when it will cool at the DALR, which is less than the ELR, so the air once more becomes neutrally buoyant and stable.

**Condobolinian** A *\*stage* in the *\*Devonian* of Australia, underlain by the *\*Cunninghamian*, overlain by the *\*Hervyan*, and roughly contemporaneous with the *\*Givetian* and, possibly, *\*Frasnian* stages of Europe.

**conductance** ( $\kappa$ ) In a direct current circuit, the reciprocal of the circuit's *\*resistance*; in an alternating current circuit, the circuit's resistance divided by the square of its impedance, the result being the real component of admittance. In both cases, the SI unit is the siemens (S), formerly known as the mhos (reciprocal ohms).

**Condylarthra (condylarths)** (class *\*Mammalia*) Extinct order comprising seven families of primitive *\*ungulates*, including some resembling the original eutherian (see EUTHERIA) stock so closely that their classification has presented many difficulties. Condylarths appear to be transitional between insectivores and true ungulates. They ranged from the late *\*Cretaceous* to the latter part of the *\*Miocene* and were a major part of the *\*Palaeocene* fauna. Some forms possessed claws and may have been arboreal (Hyopsodontidae), others became highly specialized, but the

development of hoofs and ungulate dentition can be traced to families in the order.

**condylarths** See [CONDYLARTHRA](#).

**cone-in-cone structure** A *\*secondary* sedimentary structure consisting of small cones nested one inside another and most commonly made from calcium carbonate. They are believed to form by the growth of fibrous crystals in the sediment while this is still plastic.

**cone of depression** The region, shaped like an inverted cone, in which the *\*water* table is drawn down, or depressed, in the vicinity of a *\*borehole* from which *\*groundwater* is being abstracted by pumping.

**cone penetrometer** Instrument for testing the relative *\*shear* strength of soil. A standard cone is pressed into the ground under known *\*load* and depth of penetration measured. See [PENETRATION TESTS](#).

**cone sheet** A *\*dyke* shaped in cross-section like a cone dipping inwards to a central *\*pluton*. This characteristic form is usually explained by reference to a stress field, shaped like an inverted umbrella, generated by the parent pluton at depth.

**Conewangan** See [FAMENNIAN](#).

**confidence interval** In statistics, a range of values based on the observed data which are likely to contain the true unknown value for a specified proportion of the time (confidence level) usually expressed as a percentage.

**confidence level** See [CONFIDENCE INTERVAL](#).

**confined aquifer** See [AQUIFER](#).

**confining pressure** The combined *\*hydrostatic* stress and *\*lithostatic* stress; i.e. the total weight of the interstitial pore water and rock above a specified depth.

**confluence** Of air flow, convergence of adjacent streamlines, which increases air velocity and reduces pressure. See also [CONVERGENCE](#).

**conformable** Applied to a sequence of *\*strata* deposited in an apparently continuous succession. Contrast [CONCORDANT](#).

**conglifraction** See FROST WEDGING.

**congeliturbation** See GELITURBATION.

**congestus** From the Latin *congestus* meaning ‘piled up’, a species of \*cumulus cloud with deep bulging form and an upper part having a cauliflower-like appearance. See also CLOUD CLASSIFICATION.

**conglomerate (puddingstone)** Coarse grained (\*rudaceous) rock with rounded \*clasts that are greater than 2 mm in size. Conglomerates may be termed ‘intraformational’ if formed of local, recently deposited clasts, or ‘extraformational’ if the clasts are derived from outside the area of deposition. The term ‘polygenetic’ is used to describe a conglomerate rock that has been produced under a variety of conditions or processes.

**congruent dissolution** Transition from a solid substance to a liquid of the same composition.

**congruent solution** Solution of a double salt (i.e. a salt formed by the crystallization of two or more components, e.g. \*dolomite) to yield quantities of component \*ions in the same proportions as existed in the solid.

**Coniacian** A \*stage in the European Upper \*Cretaceous, for which the \*stratotype is at Cognac, France, preceded by the \*Turonian and followed by the \*Santonian. It is dated (Int. Commission on Stratigraphy, 2004) at 89.3–85.8 Ma ago. See also SENONIAN.

**Coniferales** In some classifications, the order comprising the conifers. See CONIFEROPHYTA.

**Coniferophyta (Pinophyta)** The biggest division of \*gymnosperms, with a long fossil history, comprising trees and shrubs; most are resinous. The leaves are often needle- or scale-like. Fertile parts occur in unisexual cones, variously containing sterile scales. The ovule and seed are naked and borne on a scale. They first appear as fossils in \*Carboniferous rocks.

**Coniferopsida** In some classifications, a class of \*gymnosperms comprising 4 orders: Cordaitales; Coniferales (conifers); Ginkgoales (ginkgos); and Taxales (yews). The Cordaitales appeared in the early \*Carboniferous and became extinct at the end of the \*Permian. The

Coniferales appeared in the Upper Carboniferous, the Ginkgoales in the early Permian, and the Taxales in the *\*Jurassic*. These three are extant.

**conifers** See CONIFERALES.

**conjugate fault set** A cross-cutting set of *\*fault* planes which ideally intersect at angles of 60° and 120°, and have both left-handed and right-handed *\*shear senses*. The line of intersection is parallel to the direction of intermediate *\*principal stress* ( $\sigma_2$ ). The maximum principal stress bisects the acute angle and the minimum principal stress bisects the obtuse angle.

**conjugate fold** A set of paired, asymmetric *\*folds* whose *\*axial planes* *\*dip* towards one another. Limbs are commonly straight, and *\*hinge* zones short and angular. Conjugate folds are thought to be formed during the final stages of deformation.

**conjunct** Applied to the distribution of populations that have overlapping ranges, allowing DNA to be exchanged between them. Compare DISJUNCT.

**connate water** From the Latin *connatus* meaning ‘born together’, water that has remained trapped in a *\*sedimentary rock* since the original sediments were laid down in that water, prior to *\*lithification*. Connate water may be very old and saline.

**Conodontophora** The category into which *\*conodonts* were formerly placed.

**conodonts** Small, phosphatic, *\*fossil* teeth, common in rocks from the *\*Cambrian* to *\*Triassic* (and formerly placed in the category *\*Conodontophora*) that belonged to elongated, fish-like animals that were probably chordates (*\*Chordata*), possibly vertebrates, and lived as active predators. Two eyes were located in lobe-shaped structures at the anterior end, a *\*notochord* ran down the length of the worm-shaped body, there were muscular fins at the posterior end, and the feeding apparatus comprised the only hard parts.

**conoscopic** Applied to the converging light in a transmitted-light microscope, when the addition of a condenser below the microscope stage through which *\*plane-polarized* light is passed converges the light on to the stage to produce *\*interference* figures and optical effects characteristic of individual *\*minerals*. Compare ORTHOSCOPIC.

**Conrad discontinuity** A boundary within the Earth's **\*continental crust** that can be detected seismically at about 10–12 km depth, although exploratory deep drilling has failed to locate it. The boundary separates the crust into a lower, basic layer and an upper, granitic layer.

**consequent stream** Stream whose course is consequent upon the shape of a newly emerged land surface. Its course shows no necessary relationship with the underlying geologic structure, although older usage tended to restrict the term to a stream flowing in a downdip direction across gently inclined strata.

**conservation-Lagerstätte** See LAGERSTÄTTE.

**conservative margin** The zone between two lithospheric **\*plates** in which **\*crust** is being neither created nor destroyed, but the plates are sliding tangentially relative to each other along a **\*transform fault**.

**consistence (consistency)** The resistance of soil to physical impact such as ploughing, digging, or handling. It is controlled by the degree of adhesion between soil particles. It is described when dry as loose, soft, or hard, when moist as loose, friable, or firm, and when wet as sticky, or plastic.

**consistency** See CONSISTENCE.

**consistency index** In **\*cladistic** analysis, a measure of **\*homoplasy** in a phylogenetic tree (or **\*cladogram**), calculated as the number of steps (i.e. **\*character** state changes) in the cladogram divided by the smallest possible number of steps. The index therefore runs from 0 to 1. A low consistency index (less than 0.5) tends to indicate that much homoplasy has occurred.

**consolidation** In geology, any process by which loose earth materials become compacted, including **\*cementation**, **\*diagenesis**, **\*recrystallization**, **\*dehydration**, and **\*metamorphism**. In soil mechanics, the term implies a slow reduction in volume and an increased density of saturated soil under **\*load**, e.g. beneath buildings. Rate of consolidation depends on the rate of pore water escape, therefore on the **\*permeability** of the soil.

**constant head permeameter** See PERMEAMETER.

**constant offset** A constant separation between a geophysical source and a receiver (see also OFFSET). Constant-offset profiling (COP) is a specialized

method of marine seismic profiling using two ships, one shot-firing and the other recording, which travel along a profile at a constant offset. COP is used for mapping variations in crustal structures over large areas.

**constant-separation traversing (CST)** An electrical technique in which an *\*electrode* array (commonly a *\*Wenner* or *\*dipole–dipole* array) is moved along a survey line, keeping the electrode separations constant. Similar survey techniques are used in *\*electromagnetic* surveying, where a dual-coil method is employed; in this case constant coil separations used.

**constant-velocity gather (CVG)** In a *\*seismic velocity* analysis, a method that assumes a constant velocity for the entire path of a ray, allowing the *\*normal moveout* (NMO) to be calculated for each *\*seismic* trace as a function of two-way travel time (TWTT). Where the correct velocity has been applied for a given reflector, the NMO will appear horizontal on the CVG; if the velocity applied is too high or too low, it will appear curved. CVG is a necessary precursor for *\*stacking*.

**constant-velocity stack (CVS)** A method similar in principle to a *\*constant-velocity gather* (CVG), in which the *\*normal* move-out is applied as in a CVG and up to ten gathers are stacked (see *STACKING*) and displayed as CVS velocity panels. The correct velocity is that which produces the strongest event for a given reflector. CVS tends not to produce such precise values for velocity as the CVG method.

**Constellation Observing System for Meteorology, Ionosphere and Climate** See FORMOSAT-3.

**constructive boundary** See CONSTRUCTIVE MARGIN.

**constructive margin (constructive boundary)** The zone between two lithospheric *\*plates* which are diverging and consequently where new *\*crust* is being formed. Constructive *\*plate margins* are associated with shallow-focus earthquakes, high heat flow (up to ten times the average), and tholeiitic basalt (see *THOLEIITE*). Constructive plate margins develop oceanic *\*ridges*. See SEA-FLOOR SPREADING.

**constructive wave** Wave that leads to the build-up of a beach, due to the *\*swash* of the wave being more effective in moving material than the

**\*backwash.** Usually, constructive waves are associated with low-energy conditions and a gentle offshore gradient.

**contact** The depositional, **\*intrusive**, or faulted surface along which two different rock types are juxtaposed. The term is especially applicable to situations where **\*plutonic \*igneous** rocks intrude into **\*country** rocks; in this context ‘contact’ also refers to the effect on country rocks of conductive or convective heat transfer (i.e. **\*contact metamorphism**).

**contact aureole (metamorphic aureole)** A region in which **\*country** rocks surrounding an **\*igneous** intrusion have been **\*recrystallized** in response to the heat supplied by the intrusion. The widths of contact aureoles are quite variable and partly depend on the size of the igneous intrusion—the larger the intrusion, the wider the aureole. For intrusions of similar size, an aureole formed in response to simple conduction of heat will, in general, be thinner than an aureole formed in response to the more efficient convective transfer of heat in escaping mineralizing fluids. *See also* CONTACT METAMORPHISM.

**contact goniometer** *See* GONIOMETRY.

**contact metamorphism** The **\*recrystallization** of rocks surrounding an **\*igneous** intrusion in response to the heat supplied by the intrusion. Since there is no significant increase in the pressure gradient around an intrusion, recrystallization processes in the surrounding **\*country** rocks are a response only to an increase in the thermal gradient around the intrusion. Hence contact metamorphism is also known as ‘thermal metamorphism’. **\*Metasomatism** often takes place during contact metamorphism, unlike **\*regional metamorphism**. *See also* CONTACT AUREOLE.

**contact resistance** The measured resistance between an earthed electrode and the ground, or between a polarizable electrode and a rock specimen, or between contacts in an electrical circuit.

**contact twin** Twinned **\*crystals** (*see* CRYSTAL TWINNING), where the two individuals comprising the twins are in contact along a **\*composition** plane.

**contained fragments** *See* PRINCIPLE OF INCLUDED FRAGMENTS.

**contessa del vento** Type of lenticular cloud (see [LENTICULARIS](#)) typically with a rounded base and bulging upper surface. Sometimes a number of separate discs of this cloud-form extend one above another. Such cloud occurs on the lee side of distinct mountains within an eddy zone. In the case of Mt Etna the cloud is related to a westerly air stream.

**continental air** A very dry [\\*air mass](#) that forms over a continent.

**continental crust** That portion of the Earth's surface overlying the [\\*Mohorovičić](#) discontinuity, and with an average density of 2700–3000 mg/m<sup>3</sup>. The thickness is variable, mostly 30–40 km, except for areas of recent mountain building where the thickness can be 70 km. The crust comprises two layers, but the boundary between them is poorly defined. The upper layer is of granitic composition with surface enrichment in [\\*radiogenic](#) and other [\\*lithophile](#) elements, and has an average density of 2700 mg/m<sup>3</sup>. The lower layer was previously considered to be of gabbroic composition but is now thought to be quartz [\\*diorite](#) in [\\*granulite](#) metamorphic grade.

**continental drift** Hypothesis proposed around 1910 to describe the relative movements of continental masses over the surface of the Earth. A major theorist of continental drift, and certainly the one who gave the hypothesis scientific plausibility, was [Alfred \\*Wegener](#) (1880–1930). His work was based on qualitative data, but has been vindicated in recent years by the development of the [\\*plate tectonics](#) theory, which has provided geologists with a viable mechanism to account for continental movements. *See also* [POLAR WANDER](#).

**continental freeboard** The average level of the sea surface relative to the continents.

**continentality** A measure of how the climate of a place is affected by its remoteness from the oceans and oceanic air. The difference between the average temperatures prevailing in January and July is most often quoted as an indicator of this.

**continental margin** Zone that consists of the [\\*continental shelf](#), [\\*continental slope](#), and [\\*continental rise](#). It extends from the shoreline to the deep-ocean floor at a depth of 2000 m. The zone is underlain by [\\*continental crust](#). Continental margins have been divided into [\\*active](#)

margins or **\*passive** margins depending on their coincidence, or otherwise, with **\*plate margins**.

**continental rise** Smooth-surface accumulations of sediment which form at the base of the **\*continental slope**. The surface of the rise is gently sloping with gradients between 1:100 and 1:700. The width of the rise varies but is often several hundred kilometres. Two types of deposit lead to the formation of rises: **\*turbidites** laid down by **\*turbidity** currents flowing down the continental slope, and **\*contourites** laid down by **\*contour currents** flowing along the rise at the base of the **\*continental margin**.

**continental shelf** Gently seaward-sloping surface that extends between the shoreline and the top of the **\*continental slope** at about 150 m depth. The average gradient of the shelf is between 1:500 and 1:1000 and, although it varies greatly, the average width is approximately 70 km. Five major types of shelves may be recognized: (a) those dominated by tidal action; (b) those dominated by wave and storm action; (c) those dominated by carbonate deposition; (d) those subject to modern glaciation in polar areas; and (e) those floored by **\*relict sediments** which constitute up to 50% of the total shelf area.

**continental-shelf waves** **\*Vorticity** waves produced in a **\*continental-shelf** area where there is a sea-bed slope. In the northern hemisphere, if a water column is displaced into shallower water it develops negative relative vorticity, or anticyclonic motion; if displaced into deeper water it will develop positive relative vorticity, or cyclonic motion. The net result is for shelf waves to progress in a poleward direction along the west coasts of continents, or equatorwards along east coasts.

**continental shield crustal type** A region composed of highly deformed **\*metamorphic rocks** and **\*plutons** that are **\*Hadean**, **\*Archaean**, or **\*Proterozoic** in age and have not been affected by **\*tectonism** or covered by **\*sediments** since the Proterozoic. See **CRATON**.

**continental slope** The relatively steeply sloping surface that extends from the outer edge of a **\*continental shelf** down to the **\*continental rise**. The total relief is substantial, ranging from 1 km to 10 km, but the slope is not precipitous and ranges from 1° to 15° of slope (average 4°). Along many coasts of the world the slope is furrowed by deep submarine canyons, terminating as fan-shaped deposits at the base.

**continuation** The use of one set of measurements of a potential field (usually gravity or magnetic) over one surface to determine the set of values the field would have over another surface, usually at a different elevation. *See also* UPWARD CONTINUATION; DOWNWARD CONTINUATION.

**continuous distribution** Data that yield a continuous spectrum of values. Examples are measurements of the wing lengths of birds, or the weights of mammals, or the heights of plants.

**continuous profiling** A seismic technique in which the \*geophone and shooting patterns provide 100% subsurface coverage and which can be applied in reflection surveys and, with more practical difficulty, in refraction surveys. In a refraction survey the refractor must be monitored, especially if it is not planar, and this may necessitate laying out the geophones in an irregular pattern.

**continuous reaction series** The continuous change in composition of a \*solid-solution \*mineral in order to maintain a state of equilibrium with a cooling \*magma. The mineral changes its composition by continuously exchanging \*cations with the cooling magma in which it floats. The \*plagioclase feldspar solid-solution mineral series is a good example of a continuous reaction series. During the cooling of a magma, the plagioclase continuously exchanges Ca and Al for Na and Si in order to maintain an equilibrium composition. The principle of continuous reaction was first presented by Norman \*Bowen in 1928. *See also* BOWEN'S REACTION PRINCIPLE.

**continuous velocity logging (sonic log)** A technique which provides a record of the single travel time a sound wave takes to pass through the sidewall of a \*borehole from a \*sonde source to a receiver, when these are a constant distance apart and on the same down-hole tool. As the tool is drawn up the borehole to the surface a record of the variations in travel time is plotted as a function of down-hole depth. Measured in units of  $10^{-6}$  feet per second, it provides the inverse of the formation velocity.

**CONTOUR** The abbreviated name of the failed \*Comet Nucleus Tour, launched in 2002 to visit three comets.

**contour current** Undercurrent typical of the \*continental rise which flows along the western boundaries of ocean basins. Such currents occur

particularly in regions in which density stratification is strong because of the supply of cold waters originating near the poles. A well-known example is the Western Boundary Undercurrent which hugs the continental rise of eastern N. America. Contour currents are persistent, slow-moving (velocity 5–30 cm/s) flows capable of transporting mud, *\*silt*, and *\*sand*.

**contour diagram** A stereographic *\*equal-area* net on which orientation data (i.e. the *\*azimuths* of structural features) are plotted as lines or points and then joined to form contours linking areas of equal density of data, thus providing a visual description of the range and concentration of the data. There are several ways to construct such diagrams manually or by computer, and there are statistical tests for evaluating the significance of the densities revealed.

**contourites** Sediments that have been deposited by *\*contour currents* on the *\*continental rise*. The sediments are thin-bedded *\*silts*, *\*sands*, and muds. The sands are well sorted, laminated, or cross-laminated, with many internal erosion surfaces, and there are concentrations of heavy minerals. Both bottom and top contacts of the beds are sharp, and the beds lack great lateral continuity.

**contracting Earth hypothesis** The hypothesis stating that mountain building is caused by the contraction of the Earth. Conventionally, the wrinkles that form on a drying apple are used as an analogy for the formation of mountains on Earth. The hypothesis was prevalent in the 19th and early 20th centuries, prior to the discovery of *\*radiogenic heating* within the Earth. The Earth has probably reduced by some 5% in volume since its original formation, but mountains mainly result from *\*plate tectonic* motions.

**contractional kink band (reversed kink band)** *\*Kink bands* that form when a compressive *\*stress* acts at an angle to the layering of the *\*foliation*. The angles on either side of each kink are approximately equal. *Compare CONJUGATE FOLDS.*

**contraction limit** *See ATTERBERG LIMITS.*

**contrail** *See CONDENSATION TRAIL.*

**contrast** In *\*remote* sensing, the ratio of energy emitted or reflected by an object to that emitted or reflected by its immediate surroundings.

**contrast stretching** In *\*remote* sensing, the artificial increase in range of *\*digital* number values of *\*pixels* in an image in order to increase *\*contrast*. *See also* DECORRELATION STRETCHING; EDGE ENHANCEMENT.

**control system** In *\*geomorphology*, a *\*process–response system* where human interference changes the natural functioning of the system. For example, the trapping of sediments that occurs where beach *\*groynes* are built may lead to *\*erosion* within the adjacent sub-system and so to a change in the geometry of the overall process–response system.

**convection** **1.** Vertical circulation within a fluid that results from density differences caused by temperature variations. Convection currents occur in the oceans when a water mass that is denser than the water below it sinks and is replaced by lighter, warmer water. **2.** In meteorology, the process in which air, having been warmed close to the ground, rises. The convective uplift of *\*air* parcels is one of the main processes leading to condensation and cloud formation. *See also* DISH-PAN EXPERIMENT; FORCED CONVECTION; HADLEY CELL; INSTABILITY; STABILITY. **3.** Within the Earth, the *\*radiogenic* heat release results in convective motions causing tectonic *\*plate motions*. The location and configuration of the *\*convective* cells is uncertain, but they appear to be *\*mantle*-wide and marked by most heat loss along the *\*mid-ocean* ridges. The difference in temperature between upgoing and downgoing convective limbs within the mantle may be only 1–2 °C. In the upper *\*oceanic* crust, heat loss is mainly by convective circulation systems combined with thermal conduction. *See also* CREEP MECHANISMS; GEOTHERMAL GRADIENT; HEAT FLOW; NUSSELT NUMBER; RAYLEIGH NUMBER.

**convection current** *See* CONVECTION (1).

**convective cell** The pattern formed in a fluid when local warming causes part of the fluid to rise, and local cooling causes it to sink again elsewhere. The atmosphere in low latitudes forms such cells, known as *\*Hadley* cells, as warm equatorial air rises, moves away from the equator and cools, and then descends over the subtropics.

**convective condensation level** The level at which surface air will become saturated when rising by convection. *See also* LIFTING CONDENSATION


LEVEL.

**convective instability** See POTENTIAL INSTABILITY.

**convergence** 1. Situation in which, over a given lapse of time, more air flows into a given region than flows out of it. It is commonly accompanied by confluence of the streamlines, but may be caused by differences of velocity, e.g. where the wind comes against a coast or a mountain wall. Surface friction can produce convergence. Compare DIVERGENCE. 2. The point, line, or region where two oceanic water masses or surface currents meet. This leads to the denser water from one side sinking beneath the lighter water of the other side.

**convergent evolution** The development of similar external morphology in organisms which are unrelated—except through distant ancestors—as each adapts to a similar way of life. Sharks (fish), dolphins (mammals), and **\*ichthyosaurs** (extinct reptiles) provide good examples of convergence in the aquatic habitat.

**convergent margin** A boundary between two lithospheric **\*plates** which are moving towards each other. Some such boundaries involve **\*subduction** of **\*oceanic** crust and are called '**\*destructive**' boundaries; others are **\*collision** zones between continents where it is thought that all the oceanic crust has been subducted and that the ocean basin has reached the final stage of the **\*Wilson** cycle.




**Convergent margin**

**converted wave** See S-WAVE.

**convex slope** See SLOPE PROFILE.

**convolute** In a coiled gastropod (\*Gastropoda), applied to the condition in which the outer whorls embrace the inner ones so that the latter are nearly or partly invisible. In nautiloid cephalopods (\*Nautiloidea) this type of \*coiling is sometimes called 'nautilicone'.



## Convolute

**convolute lamination** A *sedimentary structure* consisting of a series of upright or overturned *folds* whose intensity dies out both upwards and downwards within a single bed. The convolutions are formed by the expulsion of water from rapidly deposited sediments by an external shock, e.g. an earthquake, the effect of large waves, or by the rising and falling of the *water* table through the sediment.

**convolution** A mathematical operation (symbol  $*$ ) to define the change in the shape of a wave-form caused by its passage through a *filter*. If an *impulse* response function  $f$  is convolved with an input  $g$ , then the output,  $h$ , is given by:  $h = g*f = \sum^j \times g(t - j) \times f(j)$ .

**Conybeare, William Daniel** (1787–1857) An English clergyman, Conybeare is best known as co-author, with William Phillips, of *Outline of the Geology of England and Wales* (1822), one of the most influential textbooks on stratigraphy of the period. He also described and reconstructed *saurian* fossils from the Lyme Regis area of England. As a friend and collaborator of **William Buckland**, Conybeare was an influential member of the Oxford School of Geology.

**Cook, James** (1728–1779) A navigator, surveyor, and explorer, who captained three expeditions to the Pacific between 1768 and 1779. He surveyed the coasts of New Zealand and eastern Australia, and explored part of the seaboard of Antarctica. He was able to show that the supposed ‘Great Southern Continent’ did not exist.

**Cooksonia hemispherica** (order Rhyniopsida) Extremely primitive plant, a few centimetres high, from the Upper \*Silurian and Lower \*Devonian. Members of the family did, however, possess an epidermis (protective outer layer) and stomata (specialized pores) to control the passage of gases. They also had an underground rooting portion, the rhizome, and they branched \*dichotomously. *C. hemispherica* is the first representative of the Rhyniaceae, and is known from the Silurian of Europe. See also PSILOPHYTALES; RHYNIA.

**cooling age** See FISSION-TRACK DATING.

**cooling joint (shrinkage joint)** See JOINT.

**coombe rock** See HEAD.

**coordinated stasis** The idea, proposed in 1992 by Gordon Baird, that certain groups of species remain unaltered for tens of millions of years, then experience an episode of rapid extinction and the formation of new species. This resembles \*punctuated equilibrium acting at the level of communities and may occur because the species interact so closely they cannot evolve, instead responding to environmental change by moving as a group to a more hospitable location. The fossil record of animals dwelling in ocean-bottom muds in the \*Silurian to middle \*Devonian appears to support the idea, but it is not accepted by all palaeontologists.

**coordination number** Number of atoms, ions, molecules, or groups of these that surround a given atom in a complex molecule. In geology the most common groups are oxygen atoms surrounding a \*cation, e.g. in silicate minerals (see SILICATES) silicon is always surrounded by four oxygen atoms.

**COP (constant-offset profiling)** See CONSTANT OFFSET.

**Copernican System** See LUNAR TIMESCALE.

**Copernicus** A space mission, formerly known as GMES (Global Monitoring for Environment and Security), headed by the European Commission and the \*European Space Agency, to coordinate the delivery of data from thirty satellites, known as Sentinels. Sentinel-1A and -1B, launched in April 2014 and April 2016, respectively, provide all-weather day and night radar images. Sentinel-2A, launched on 23 June 2015, provides high-resolution optical images for land services. Two Sentinel-3 satellites, launched on 16 February 2016 and 25 April 2018, provide data on ocean and land. Sentinel-4 and -5, planned for launch in 2021, -4 into \*geostationary orbit and -5 into \*Sun-synchronous orbit, will monitor atmospheric trace gases and \*aerosol over Europe. Sentinel-5P, launched on 13 October 2017 into Sun-synchronous orbit, monitors atmospheric composition and ozone. Sentinel-6, also called Jason-CS (Jason Continuity of Service), provides high-resolution radar altimetry to complement Sentinel-3 and continues the missions of \*Jason-2 and -3.



[https://www.esa.int/Our\\_Activities/Observing\\_the\\_Earth/Copernicus/Overview3](https://www.esa.int/Our_Activities/Observing_the_Earth/Copernicus/Overview3)

- Copernicus, Observing the Earth.

**Cope's rule** In 1871, the American palaeontologist Edward Drinker Cope (1840–97) noted a phylogenetic trend towards increased body size in many animal groups, including mammals, reptiles, arthropods, and molluscs. This came to be known as Cope's rule. It remained unchallenged until a study of more than 1000 insect species in 1996 and was finally disproved in 1997, by a study in which David Jablonski made more than 6000 measurements on 1086 species of Late \*Cretaceous fossil molluscs spanning 16 million years and found as many lines led to decreased size as increased. Evolutionary lineages show no overall tendency to greater size, but if the extant survivor happens to be larger than its immediate ancestor (e.g. the horse) this coincidence appears to validate Cope's rule.

**copper, native** A malleable metallic element, Cu; sp. gr. 8.9; \*hardness 2.5; \*cubic, but normally found \*massive, or as \*platy flakes infilling cracks or \*vesicles in volcanic rocks, or as a result of \*weathering of copper-rich \*mineral deposits.

**copper glance** See CHALCOCITE.

**copper pyrites** See CHALCOPYRITE.

**coppice dune** A sand \*dune that forms around and in the lee of a clump of vegetation.

**co-product** Many \*ore deposits are worked for more than one element. Where those elements are of almost equal economic significance they are called 'co-products'. Compare BY-PRODUCT.

**coprolite** A fossilized dropping or excreta (i.e. a fossilized faecal pellet). Coprolites may have distinctive shapes or markings which can provide information regarding the structure of the animal's alimentary canal, and analysis of the contents may also reveal its diet.

**coquina** \*Clastic or detrital \*limestone that contains a high proportion of coarse shell debris cemented by calcium carbonate.

**coral** See ANTHOZOA; HETEROCORALLIA; HYDROZOA; MILLEPORINA; OCTOCORALLIA; RUGOSA; SCLERACTINIA; STYLASTERINA; TABULATA; ZOANTHARIA.

**coral growth lines** All corals that have a calcified outer wall display minute growth lines on their outer surfaces. The carbonate is secreted by symbiotic \*algae (zooxanthellae) which, responding to day and night, create a series of diurnal growth increments. Studies on \*Devonian isolate corals indicate a 400-day year and, therefore, a 22-hour day. Post-Devonian data confirm a near-linear deceleration of the Earth's rotational velocity towards the present 24-hour day. The number of daily growth lines can therefore be used in a crude way to calibrate the geologic record.

**coralline limestone** See LEIGHTON—PENDEXTER CLASSIFICATION.

**corallite** The skeleton formed by an individual coral polyp, which may be either solitary or part of a colony.

**coralloid** Literally, coral-like, but used to describe the circular, curved, or irregular shapes that sometimes result from the chemical precipitation of minerals, e.g. \*marcasite.

**corallum** The skeleton of a colonial coral, made up from individual skeletons (**\*corallites**) each secreted by one polyp. Distinctive corallum shapes are given a variety of names.

**Cordaitales** Extinct **\*gymnosperm** order, included in the **\*Coniferopsida**, which appeared in early **\*Carboniferous** times and disappeared towards the end of the **\*Permian**. The cordaitaleans were trees up to 30 m high, with strap-like leaves and primitive cones. Some cordaitaleans developed stilt roots, probably lived in swamp habitats, and were analogous to modern mangroves. Fossils of stems, leaves, roots, and cones are locally abundant in coal. The term *Cordaites* is a **\*form-genus** name, and strictly applies to the leaves of cordaitaleans, but it has often been used informally for the whole plant. Other form-genera include *Cordaianthus* for cordaitean cones, *Amelyon* for the roots, and *Dadoxylon*, *Araucarioxylon*, *Mesoxylon*, and *Pennsylvanioxylon*, according to type or age, for stem fragments and wood.

**Cordelia (Uranus VI)** One of the lesser satellites of **\*Uranus**, with a radius of 13 km. It was discovered in 1986.

**cordierite** **\*Silicate** mineral  $\text{Al}_3(\text{Mg,Fe})_2 [\text{Si}_5\text{AlO}_{18}]$  which can be iron-rich and is a member of the **\*cyclosilicate** group; sp. gr. 2.5–2.8; **\*hardness** 7; **\*orthorhombic**; dark blue or greyish-blue, translucent to transparent; **\*vitreous** **\*lustre**; **\*crystals** rare, **\*prismatic** or pseudo-hexagonal, but usually occurs **\*massive**; **\*cleavage** imperfect {010}, parting {001}; occurs in aluminous rocks that have been subjected to thermal or regional **\*metamorphism**, in **\*hornfels**, **\*schists**, and **\*gneisses**, in association with **\*andalusite**, **\*spinel**, **\*quartz**, and **\*biotite**. Fine dark blue examples are used as **\*gemstones**. It is named after the 19th-century French geologist P. L. A. Cordier (1777–1861).

**cordillera** **1.** Broad assemblage of mountain ranges belonging to **\*orogenic** belts of different ages which formed originally at **\*destructive** plate margins. **2.** System of mountain ranges, together with their related plateaux and **\*intermontane** basins. For example, the Cordillera of N. America includes all the mountain ranges and plateaux west of the Great Plains and of the Mexican lowlands. **3.** Subsidiary complex of ranges within a mountain system, e.g. the eastern and western cordillera of the Andes. **4.** Individual range, e.g. the Cordillera Patagonica of the southern Andes.

**core 1.** The central zone or unit of the Earth. It is composed of iron, with a lighter element, probably sulphur, and accounts for 16% of the Earth's volume and 32% of its mass. The core is separated into inner and outer units. The inner core is a solid with a radius of about 1220 km and the outer core, which does not permit the passage of shear waves (*\*S-waves*), is liquid. Other planets have mass distributions that suggest they possess cores, e.g. *\*Mars*, *\*Venus*, and *\*Mercury*. The *\*Moon* may have a small core. *\*Saturn* has magnetic fields interpreted to indicate a metallic core, probably of liquid hydrogen. **2.** A rock or ice specimen obtained by *\*drilling*.

**core barrel** The part of *\*borehole* drilling equipment in which the core is collected. Usually the barrel is 50–100 mm in diameter with the drill *\*bit* at one end. Sometimes an inner tube is present, linked by a swivel to prevent its rotation (double core barrel), and even a triple core barrel may be used to aid the recovery of weak rock.

**core-logging** Geologic description of a *\*core* obtained from a *\*borehole*. This includes lithological, palaeontological, and structural information, often with reference to engineering properties. *See also* ACOUSTIC LOGGING; CONTINUOUS VELOCITY LOG; INDUCED POTENTIAL LOG; NEUTRON LOG; RESISTIVITY LOG; SELF-POTENTIAL LOG.

**core recovery** Length of *\*core* recovered from a *\*borehole*, compared with depth of hole cored, expressed as a percentage. *See* TOTAL CORE RECOVERY (TCR).

**core slicer 1.** A diamond-edged cutter that can be used inside a *\*borehole* to cut triangular slices 2–3 cm wide and up to 1 m long from the rock walls. **2.** A diamond-tipped saw blade used to slice specimens from a rock *\*core* drilled in the field.

**corestone (woolsack)** Rounded boulder, occurring individually or in piles at the ground surface, or in exposed sections. It results from an initial phase of subsurface *\*chemical weathering*, of a *\*joint-bounded* block, followed or accompanied by surface *\*erosion* that exposes the corestone.

**core wall** *See* DIAPHRAGM WALL.

**Cor F** *See* CORIOLIS EFFECT.

**Coriolis effect (Cor F)** An apparent force (hence the abbreviation to Cor F) acting on moving objects that results from the Earth's rotation. It causes objects in motion, including oceanic and atmospheric currents, to be deflected to the right in the northern hemisphere and to the left in the southern hemisphere by an amount that varies with latitude and the speed of the moving body. The changing magnitude of the Cor F with latitude is given by the Coriolis parameter:  $2\Omega \sin\phi$ , where  $\Omega$  is the angular velocity of the Earth ( $7.29 \times 10^{-5}$  rad/s) and  $\phi$  is the latitude. Since  $\sin 0 = 0$  and  $\sin 90 = 1$ , the Cor F varies from zero at the equator to a maximum at the poles. When the speed of the body is taken into account,  $\text{Cor F} = 2\Omega \sin\phi v$ , where  $v$  is the speed. It was first described in 1835 by the French physicist and engineer Gustave de Coriolis (1792–1843).

**Coriolis parameter** See CORIOLIS EFFECT.

**Coriolis/WindSat** A space mission by the US Department of Defense to develop and validate measurements of wind vectors and to monitor solar activity for better prediction of solar storms. The satellite was launched on 6 January 2003, from California, into a Sun-synchronous orbit.

**corona** 1. White or occasionally coloured rings of lights, typically from blue inside to red outside, that sometimes appear to surround the Sun or Moon. The effect is created by diffraction of light by spherical water drops in such clouds as *\*altocumulus*. Compare HALO. 2. Concentric zones of one or more *\*minerals* surrounding a core mineral. Coronas can be formed in a number of ways. (a) The *\*discontinuous* reaction of minerals with a *\*magma* can be preserved as coronas around the original high-temperature mineral if the cooling rate is fast enough to prevent the reactions going to completion. (b) Late-stage fluids may react with an earlier *\*primary mineral* to develop a corona of *\*secondary* minerals. (c) Two minerals may undergo sub-*\*solidus* reactions (reactions occurring after the rock has solidified) to maintain equilibrium as a rock mass cools, developing a corona of lower-temperature minerals. These types of *\*texture* are also known as 'reaction rims'. See also CRYSTAL ZONING. 3. One of the large, circular features (150–600 km diameter) of uncertain origin, comprising up to 10–12 subconcentric ridges and grooves, which surround an inner region of irregular relief, found on the surface of Venus, mainly in a latitudinal belt

between 55° N and 80° N along the borders of Ishtar and Tethus Regio. Most are associated with what appear to be lava flows.

**corrasion** See ABRASION.

**correlated progression** The hypothesis that evolutionary change of \*characters occurs by correlated response, i.e. a change in one character may influence change in another, such that the rate of change of the two characters is not independent. Compare MOSAIC EVOLUTION.

**correlation** 1. In stratigraphy, correlation is the establishment of a correspondence between stratigraphic units. It depends on the similarities that exist in terms of \*lithology or \*fossil content. Isolated stratigraphic units, or successions, may be either 'correlated', i.e. they were once physically continuous, or time-correlated, i.e. equated in terms of time. 2. In geostatistics, correlation is a technique used to determine the degree of association between two data sets. 3. In geophysics, the comparison of one wave-form with another in the \*time domain. It is analogous to \*coherence in the \*frequency domain. See AUTO-CORRELATION; CROSS-CORRELATION.

**correlation diagram** Diagram illustrating probable stratigraphic equivalence from place to place.

**correlogram** A graph showing the strength of correlations in data at different time intervals and thereby exposing the existence and phases of cycles.

**corridor dispersal route** As originally defined by the American palaeontologist G. G. \*Simpson (1940), a corridor is a migration route that allows more or less uninhibited faunal interchange. Thus many or most of the animals of one \*faunal realm can migrate to another one. A dispersal corridor has long existed between western Europe and China via Central Asia.

**corrie** See CIRQUE.

**corundum** \*Mineral, Al<sub>2</sub>O<sub>3</sub>; sp. gr. 3.9–4.1; \*hardness 9; \*trigonal; the two main varieties blue and green, but can be yellow, or brown to almost black, and transparent; \*adamantine to \*vitreous \*lustre; \*crystals usually rough and barrel-shaped, tapering, and also flat and \*tabular; no cleavage, partings {0001}, {0112<sup>-</sup>}; occurs in silica-poor rocks such as \*nepheline syenites

and undersaturated (see [SILICA SATURATION](#)) alkali [igneous](#) rocks, in [contact](#) aureoles in thermally altered alumina-rich [shales](#) or [limestones](#), in aluminous [xenoliths](#) found within [basic](#) igneous rocks in association with [spinel](#), [cordierite](#), and [orthopyroxene](#), in metamorphosed [bauxite](#) deposits and in [emery](#) deposits, and in [alluvial](#) deposits because of its hardness and resistance to [abrasion](#) along with [muscovite](#), [hematite](#), and [rutile](#). Flawless crystals are the [gemstones](#) blue sapphire, red ruby, and green oriental emerald. The main use of corundum is based on its hardness. It is made into grinding wheels and discs, emery paper, and powders for grinding and polishing.

**Corynexochida** An order of [Trilobita](#) that lived from the Lower [Cambrian](#) to Middle [Devonian](#). The glabella (see [CEPHALON](#)) is variable, but usually with parallel sides or wider anteriorly. There are three suborders.

**coseismic period** See [SUBDUCTION EARTHQUAKE CYCLE](#).

**coset** See [CROSS-STRATIFICATION](#).

**COSMIC** See [FORMOSAT-3](#).

**cosmic abundance of elements** Based on data from the [Sun](#) and other stars, hydrogen and helium are by far the most abundant elements of the cosmos (e.g. the Sun's atmosphere may contain 70% hydrogen and 28% helium by mass). In general, elements show an exponential decrease in abundance with increasing atomic number ( $Z$ ), up to about 45; the abundances of heavier elements thereafter appears fairly constant. Other regularities are superimposed on this general pattern: elements of even atomic number are more abundant than adjacent elements of odd atomic number ([Oddo–Harkins rule](#)); there is a pronounced peak at atomic number 26 (the 'iron' peak); and isotopes whose mass numbers are multiples of four, i.e. multiples of the alpha-particle (helium nucleus) mass, e.g. carbon, oxygen, neon, magnesium, silicon, sulphur, and iron, have enhanced abundances. Cosmic abundances of elements constrain the way in which the [solar](#) system evolves and dictate the composition of its members (which include the Earth and ourselves on it). See also [NUCLEOSYNTHESIS](#).

**cosmic dust** Particles with a wide range of masses ( $10^{-2}$ – $10^{-18}$  g) and velocities occurring in interplanetary, circumstellar, and interstellar space. They are typically porous with low densities and a ‘cluster of grapes’ morphology, but many are compact with densities of about  $2000 \text{ kg/m}^3$ . Most particles collected from the *\*stratosphere* are comprised of layer lattice *\*silicates*, *\*olivine*, or *\*pyroxene*. In the terrestrial neighbourhood, they are mostly derived from *\*comets* and *\*asteroids*.

**cosmic radiation** Ionizing radiation from space, comprised principally of protons, alpha particles, and 1–2% heavier atomic nuclei, as well as some high-energy photons and electrons. On encountering the Earth’s atmosphere, secondary radiation is produced, mainly gamma rays, electrons, pions, and muons. Three sources are identified: (a) galactic cosmic rays, from outside the *\*solar* system, with energies in the range 1–10 GeV per nucleon; (b) solar cosmic rays, mainly associated with *\*solar* flares, with energies in the range 1–100 MeV per nucleon; and (c) *\*solar wind*, with energies of about 1000 eV per nucleon.

**cosmic-ray track** The interaction of ionizing cosmic rays (see *COSMIC RADIATION*) with mineral surfaces produces solid-state damage which, when etched with acid, is revealed as tracks of varying length. Most tracks are produced by iron group nuclei (nuclear charge (*Z*) 18–28, VH ions). Track lengths due to galactic cosmic rays may extend to depths of 20cm. In contrast, *\*solar flare* tracks are mostly less than one millimetre in length. Track densities, typically about  $10^{12}/\text{m}^2$  on unshielded lunar mineral surfaces, provide a measure of *\*exposure* age.

**cosmine** See *COSMOID SCALE*.

**cosmochemistry** The study of the chemical, isotopic, and mineralogical composition of materials originating in outer space. These include cometary and interplanetary dust particles, lunar materials, and *\*meteorites*.

**cosmoid scale** Type of scale found only in fossil lung-fish (*\*Dipnoi*), and in *\*Crossopterygii*, including the living *\*coelacanth*. The thick scales are composed of layers of vitrodentine (harder than dentine and containing little organic matter), followed by cosmine (a type of *\*dentine* perforated by branching canals), and finally by layers of vascular and laminated bone underneath.

**cosmology** The study of the origin and evolution of the universe. The current **\*big bang cosmology** derives the observable universe from a singular event 15–20 billion years ago. Previous hypotheses include the steady-state theory, in which the expansion of the universe was due to the continuous creation of matter. The most celebrated of earlier world views was the Ptolemaic system, in which the Earth was the centre of the universe; this was superseded by the Copernican revolution, which displaced the Earth from its central position.

**cosmopolitan distribution (pandemic distribution)** Distribution of an organism that is worldwide or ‘panendemic’. Apart from weeds, commensal animals, and some of the lower groups of cryptogams (plants reproducing by **\*spores** or gametes, rather than seeds), there are relatively few organisms that occur on all six of the widely inhabited continents. *See also* FAUNAL REALM.

**Cosmorhaphe** A complex grazing **\*trail** found in deep-water **\*flysch** deposits. It is a single, sinuous trail formed by a **\*deposit** feeder which exhibited efficient feeding behaviour in an area of low productivity.

**COSMO-SkyMed** A constellation of four **\*microsatellites** for global Earth observation, developed by the Agenzia Spaziale Italiana and funded by the Italian Ministries of Research and Defence, that carry **\*synthetic-aperture** radar instruments capable of operating at high resolution in all visibilities and in real time. The satellites were launched on 8 June 2007, 9 December 2007, 25 October 2008, and 6 November 2010. A second generation of two satellites, to improve the quality of the imaging service, is to be launched in 2018 and 2019.

**costa** **1.** In corals, an external projection of a **\*septum** beyond a **\*corallite** wall, forming a rib. **2.** A rib-like thickening of a shell, extending from the **\*umbo** to the margins in brachiopod (**\*Brachiopoda**) and lamellibranch (**\*Bivalvia**) shells. In gastropods (**\*Gastropoda**) the thickening of the shell usually runs either axially or spirally, and thus there may be axial or spiral costae.

**Costonian (Blackriverian)** A **\*stage** (460.9–459 Ma ago) of the **\*Ordovician** in the Lower **\*Caradoc**, preceded by the **\*Darriwilian** and overlain by the **\*Harnagian**.

**cotectic curve** In a system of three mineral components represented by a block diagram, the **\*liquidus** surface forms a trough which defines the composition of the **\*melt** having the lowest freezing temperature. The cotectic curve is the axis of this trough.

**cotectic surface** Curved surface in a **\*quaternary system** which defines the temperature range over which two or more solid **\*phases** crystallize simultaneously from a liquid. Analogous to a cotectic line in a **\*ternary system**.

**coterminous** With shared boundaries. In stratigraphy, applied, for example, to a time-line that is common to **\*stratigraphic units** of different **\*rank**.

**cotidal line** Line joining points at which given tidal levels (such as mean high water or mean low water) occur simultaneously. The lines are shown on certain hydrographic charts. The same information is contained in tide tables, where the data are given as differences from the times of high or low water at a 'standard port'.

**Cotylosauria** **\*Anapsid** 'stem reptiles' which appeared in the **\*Carboniferous**, having diverged from **\*labyrinthodont** **\*amphibians**. They flourished in the remainder of the **\*Palaeozoic**, but rapidly dwindled to extinction in the **\*Triassic**.

**coulée flow** A very thick, and relatively short, blocky **\*lava** flow formed from viscous lava, usually **\*rhyolite** or **\*dacite** in composition, which has been extruded from a **\*volcanic cone** or **\*fissure**. The short flows can be up to 100 m thick and hence have a morphology lying between that of a **\*dome** and a flow. (When used in the French literature, the term 'coulée' is applied to lava flows in general.)

**Coulomb failure criterion** The simplest relationship between **\*shear** stress ( $\tau$ ) acting parallel to a plane and **\*normal stress** ( $\sigma$ ) acting perpendicular to the plane at the point of failure. The relationship  $\sigma = \sigma_0 + \tau \tan \phi$ , where  $\sigma_0$  is the normal stress after failure and  $\phi$  is the angle of failure, implies that  $\sigma_2$  (see **PRINCIPAL STRESS AXES**) does not affect brittle fracture strength.

**Coulomb–Terzaghi shear strength equation** The equation that is used to calculate the total **\*shear** strength ( $s$ ) of the material composing a slope, taking account of the **\*pore-water** pressure. The shear strength is then equal

to the sum of the **\*normal stress** ( $\sigma$ ) and **\*cohesion** ( $c$ ). The Coulomb–Terzaghi equation is:

$$s = c + \sigma \cdot \tan \phi,$$

where  $\tan \phi$  is the coefficient of plane sliding friction, which describes the packing, surface roughness, and hardness of the materials constituting the slope.

**coulter counter** An electronic measuring device used mainly for determining the grain-size distribution in silt- and clay-sized **\*sediments** (see **PARTICLE SIZE**). The coulter counter measures the volume of the individual particles in a suspension and is particularly useful for determining the size of mud flocculates (see **FLOCCULATION**). It can also be used for sand-sized sediments.

**counterpoised beam balance (Walker's steelyard)** A long, graduated, horizontal beam which is pivoted near one end on a vertical shaft, permitting the beam (arm) to swing freely. It is used to weigh objects, by attaching them to the short end of the arm and balancing their weights with counterweights which can be moved along the graduated arm, and it can also be used to determine the densities of minerals.

**country rock** The rock which has been intruded into and/or surrounded by a **\*plutonic \*igneous** intrusion.

**couple** Two equal, parallel forces which act in opposite directions in the same plane but not along the same axis.

**coupled substitution** The substitution of two or more elements in a crystalline compound, which maintains electrical neutrality. In forming a **\*solid solution** series, ionic size is more important than **\*ionic charge**, as this can be compensated for elsewhere. For instance, when a **\*plagioclase** feldspar solid solution series forms, in passing from albite ( $\text{NaAlSi}_3\text{O}_8$ ) to anorthite ( $\text{CaAl}_2\text{Si}_2\text{O}_8$ ),  $\text{Al}^{3+}$  replaces  $\text{Si}^{4+}$ . This leaves a negative charge that is balanced by the (coupled) substitution of  $\text{Ca}^{2+}$  for  $\text{Na}^+$ .

**covalent bond** Bond in which a pair (or pairs) of electrons is shared between two atoms. The bond is often represented by drawing a single line between the symbols of the two atoms that have bonded together. Sometimes the bonding is between atoms of different elements (e.g.

hydrogen chloride, H–Cl), and sometimes between atoms of the same element (e.g. fluorine, F–F). The name ‘molecule’ is used to describe any uncharged particle containing covalently bonded atoms. *See also* [HYDROGEN BOND](#); [IONIC BOND](#); [METALLIC BOND](#).

**covalent compound** A compound with [\\*covalent](#) bonds.

**covalent radius** The atomic radius of an atom (e.g. carbon), determined in a covalent compound by a technique which can be used only for structures in which that atom (carbon) is [\\*covalently bonded](#). Measurement of the distance between planes of atoms (the ‘d-spacing’) gives the sum of the radii (the bond length) of the two covalently bonded atoms in two planes. If all the atoms in the compound are of the same element this gives the value of the radius of an atom in each plane (e.g. the bond length in [\\*diamond](#) (pure carbon) is twice the radius of an individual carbon atom). If atoms of different elements are covalently bonded, the bond length of one of the atoms can be derived from the sum of the two different bond lengths only if the bond length of the other atom is known.

**covariance** In statistics, a measure of the association between two variables. Covariance is calculated as the difference between the average product of corresponding values in the two data sets and the product of the [\\*means](#) of the two data sets. A value of zero indicates no relationship between the data sets. The covariance value can be used to calculate [\\*linear](#) regression and [\\*principal](#) components.

**covellite** [\\*Mineral](#), CuS; sp. gr. 4.6; [\\*hardness](#) 2; dark blue; [\\*sub-metallic](#) [\\*lustre](#), may show an iridescent tarnish; found [\\*massive](#) or in thin plates; occurs in the zone of [\\*secondary](#) enrichment overlying copper-rich mineral deposits.

**cow-dung bomb** A [\\*volcanic](#) bomb whose name is derived from its characteristic shape.

**coxa** The uppermost joint of the leg of an insect.

**cpx** *See* [CLINOPYROXENE](#).

**crabs** *See* [MALACOSTRACA](#).

**crachin** Condensation as low cloud or fog, often with drizzle, which is frequent in spring in coastal areas of the Gulf of Tonkin and southern China.

The weather results from the **\*advection** of warm air over a cold surface, or the mixing of **\*air** masses at the surface.

**cracking** Breaking down or decomposition of large, more complex molecules, particularly **\*hydrocarbons**, into small simple molecules, usually by heating. The process can occur naturally, resulting in older deposits containing lighter crude oil than those of younger fields. Commercially, cracking is carried out in oil refineries.

**crag** Shelly sand.

**crag and tail** Land-form consisting of a small rocky hill (crag) from which extends a tapering ridge of unconsolidated debris (tail). The crag is a residual feature left by selective glacial erosion, while the tail is drift-deposited by ice on the lee side of the obstacle.

**Craniata (Vertebrata, vertebrates)** (phylum **\*Chordata**) The subphylum of animals that have a bony or cartilaginous skull and a dorsal vertebral column. It includes the fish, **\*amphibians**, **\*reptiles**, **\*birds**, and **\*mammals**, which appear successively in the fossil record, starting in the **\*Ordovician**. See also **BONE**; **CARTILAGE**.

**cranidium** See **CEPHALON**.

**cranium (skull)** In the axial skeleton of vertebrates, the bony structure that encases the brain. The cranium comprises three parts: dermatocranium; chondrocranium; and splanchnocranium. The dermatocranium includes the roof of the cranium, the area around the orbits, and the jaw. The chondrocranium includes the floor of the cranium. The splanchnocranium gives rise to the visceral skeleton (gill arches and derivatives such as larynx and trachea).

**crater** **1.** General term for a circular, funnel-shaped depression, up to 1 km in diameter, produced by volcanic processes by which gases, **\*tephra**, and **\*lava** are or have been ejected. Several types are recognized: a crater at the summit of a volcanic cone marks the site of **\*magma** degassing and ejection of material; a **\*maar**, often occupied by a lake, results from explosive activity; and a **\*caldera** is a large volcanic depression greater than 1 km in diameter. **2.** Near-circular depression produced by the impact of an extraterrestrial body, e.g. Meteor Crater, Arizona. **\*Meteorite** craters are formed by the explosion outward and upward of material compressed and

heated strongly by the energy of impact, and so usually are circular at the time they form. They are characterized by topographically raised rims and by *\*ejecta* blankets which show inverted stratigraphy with respect to the target rocks. *See also* SHATTER CONES.

**crater counting** *See* CRATER DENSITY STUDIES.

**crater density studies (crater counting)** The establishment of the relative age of a portion of a planetary or *\*satellite* surface from the observed density of *\*meteorite* impact craters. If a calibration is available for the meteoric flux (e.g. from dated lunar surfaces), *\*absolute* ages may be obtained by this technique. Problems with the method include the identification of primary from secondary craters, and whether the surface is saturated with craters.

**Cratered Plains** *See* MARTIAN TERRAIN UNITS.

**Cratered units** *See* MARTIAN TERRAIN UNITS.

**crateriform** Having a crater-like form: a circular depression with a raised rim, surrounded by an *\*ejecta* blanket.

**craton (shield, adj. cratonic)** Area of the Earth's *\*crust*, invariably part of a continent, which is no longer affected by orogenic activity. This stability has existed for approximately 1000 Ma. A classic example is the Canadian Shield.

**creep** 1. Slow downslope movement of the *\*regolith* over hillslopes, due to gravity. The necessary disturbance of the regolith may be due to freezing and thawing; to expansion and contraction (resulting from temperature change or from wetting and drying); to additional weight and lubrication by water; or to the activities of burrowing animals. 2. *See* CREEP MECHANISMS. 3. The behaviour of *\*minerals* under low differential *\*stress* over long periods of time. Typically there is an initial stage of transient creep (*\*primary* creep) with viscoelastic strains, which changes progressively to a state of purely viscous strain, until the mineral ruptures in a final (tertiary) stage of creep.

**creep mechanisms** Mechanisms by which materials deform at the Earth's surface or, more commonly, at depth. These can be: (a) cataclastic, in which individual grains or fragments physically rotate or glide past one another

(see **CATACLASIS**); (b) dislocation creep, by a gliding motion along crystalline dislocations; (c) grain-boundary sliding; (d) **\*recrystallization**; and (e) **\*diffusion** of individual atoms. Each process is dependent on the stress, temperature, and duration of the stress.

**creep strength** The strength of a rock which is undergoing long-term **\*creep** processes. It is the threshold value, beyond which creep gives way to permanent rupture, and is virtually synonymous with **\*fundamental strength**.

**Crenarchaeota** (domain Archaea) The less derived (see **APOMORPH**) of the two kingdoms of the Archaea, composed principally of extreme **\*thermophiles** and **\*psychrophiles**. Members of the Crenarchaeota show a greater genetic similarity to those belonging to the **\*domains \*Eukaryota** and **\*Bacteria** than do those of the **\*Euryarchaeota**.

**crenulation cleavage** See **CLEAVAGE**.

**Creodonta** (class **\*Mammalia**) The more ancient of the two placental mammalian carnivorous orders, an extinct order comprising two families (Oxyaenidae and Hyaenodontidae), which appeared in the late **\*Cretaceous** and dwindled to extinction in the **\*Pliocene**. Oxyaenids (e.g. *Oxyaena*) were rather weasel-like, although some of them were large (*Patriofelis* was the size of a bear). Hyaenodonts had narrower skulls, longer legs, and well-developed **\*carnassials**, and diversified into forms reminiscent of the dogs, cats, and hyenas. Only the hyaenodonts survived the Eocene, and filled the role of scavengers until they were displaced by the modern hyena. Creodonts were small-brained and slow-moving, and are not closely related to modern carnivores.

**creodont-like teeth** Teeth resembling those of the **\*Creodonta**, carnivorous mammals of the early **\*Cenozoic** (although the first representatives appeared in the late **\*Cretaceous**), in which **\*carnassials** were formed by the molars, rather than premolars and molars as in modern **\*Carnivora**.

**crepuscular rays** Beams of sunlight made visible by haze in the atmosphere, and seen where rays penetrate gaps in clouds such as **\*stratocumulus**; this effect is called 'Jacob's ladder'. In other cases rays from a low Sun diverge upwards above cumuliform cloud.

**crescent-and-mushroom** A two-dimensional \*outcrop pattern, produced by the superposition of two \*fold systems, whose forms resemble alternating crescents and mushrooms. The pattern is sometimes called 'dome–crescent–mushroom'. See INTERFERENCE PATTERN.

**Cressida (Uranus IX)** One of the lesser satellites of \*Uranus, with a diameter of 33 km. It was discovered in 1986.

**crest** See CRESTAL PLANE.

**crestal plane** A plane or surface which contains the highest points (crests) of all beds within a folded sequence.

**crest line** A line on a folded layer which is common to and links the highest points on that layer. The term is employed most usefully in respect of plunging \*folds where the \*axial trace and crest line occupy different positions.

**Cretaceous** Third of the three \*periods included in the \*Mesozoic era. It began 146 Ma ago and ended 65.5 Ma ago. It is noted for the deposition of the chalk of the White Cliffs of Dover, England, and for the mass extinction of many invertebrate and vertebrate stocks. Among these were the \*dinosaurs, mosasaurs, \*ichthyosaurs, and \*plesiosaurs. See K/T BOUNDARY EVENT.



<https://ucmp.berkeley.edu/mesozoic/cretaceous/cretaceous.php>

- The Cretaceous Period of the Mesozoic Era.


**crevasse** 1. Deep fissure in the surface of a glacier, caused when tensile stress overcomes the shear strength of ice in the brittle upper few metres. Appropriate tensile stresses are typically developed when a glacier moves over a convex-up slope. 2. A breach in a \*levée along the bank of a river through which flood water may flow and produce \*crevasse deposits (2) (crevasse splay).

**crevasse deposit** 1. The gravelly or sandy sediment infill of a \*crevasse in glacial ice. 2. (crevasse splay) The deposit generated by a river crevasse event. It is sheet-like in geometry, thinning away from the side of the breach

in the river bank, and characterized by rapidly deposited sands, fining upwards to a muddy top, produced by the waning flow of a flood event.

**crevasse splay** See CREVASSE DEPOSIT.

**Crinoidea (crinoids)** (subphylum Crinozoa; phylum \*Echinodermata) The most primitive living class of echinoderms, whose members are either stalked (sea lilies) or unstalked (feather stars). The body is contained within a cup-like \*calyx, composed of regularly arranged plates, consisting of a lower dorsal cup which is covered by a dome (the \*tegmen). There are usually five plated and branching arms (brachial processes, or \*brachia) that articulate freely with the calyx. The upper surface contains the mouth and anus. There are tube feet along each arm with a median food groove between them leading to the mouth. The stem, when present, consists of a column of calcite discs (ossicles or \*columnals) each with a central hole (lumen) for extensions of the soft parts. All \*Palaeozoic forms were stemmed (sometimes of considerable length), but most modern forms are free swimming. They first arose in the Lower \*Ordovician, and fossil crinoids are an important constituent of Palaeozoic \*limestones.



**Crinoidea**

**crinoids** See CRINOIDEA.

**crystalite** A high temperature form of **\*quartz**  $\text{SiO}_2$  which has a stability field at atmospheric pressure above 1470 °C and below 1713 °C, at which latter temperature crystalite melts and the quartz **\*liquidus** is reached; sp. gr. 2.32; occurs as fine aggregates in cavities in **\*basalts** and in some thermally metamorphosed (see **CONTACT METAMORPHISM**) **\*sandstones**.

**critical angle ( $\Theta_c$ )** The **\*angle of incidence** at which a refracted ray grazes the interface between two media whose velocities are  $V_1$  and  $V_2$ , such that, according to **\*Snell's law**,  $\sin\Theta_c = V_1/V_2$ , provided that  $V_2$  is greater than  $V_1$ .

**critical damping ( $\mu_c$ )** The minimum amount of **\*damping** which will stop oscillation or movement.

**critical erosion velocity** See **CRITICAL VELOCITY**.

**critical flow** Critical flow occurs when the flow velocity in a **\*channel** equals the wave velocity generated by a disturbance or obstruction. In this condition the **\*Froude** number ( $Fr$ ) = 1. When the wave velocity exceeds the flow velocity ( $Fr$  is less than 1) waves can flow upstream, water can pond behind an obstruction, and the flow is said to be subcritical or tranquil. When  $Fr$  is greater than 1 waves cannot be generated upstream and the flow is said to be supercritical, rapid, or shooting. In this condition a standing wave is formed over obstructions in the river bed. In nature, supercritical flow is found only in rapids and waterfalls, but it is often created artificially by **\*weirs** and **\*flumes** with the aim of measuring **\*discharge**.

**critical reflection** The reflection, normally at a very large angle, observed when the **\*angle of incidence** and **\*angle of reflection** of a ray incident on a surface are both equal to the **\*critical angle**.

**critical temperature** The temperature above which a gas cannot be liquefied, regardless of the pressure, and becomes a supercritical fluid with the molecular freedom of a gas, but the density of a liquid. The critical temperature for water is 374°C.

**critical velocity (critical erosion velocity)** The minimum velocity of a flowing fluid that is required in order to entrain a particle. See **HJULSTRÖM RELATIONSHIP**.

**CRM** See CHEMICAL REMANENT MAGNETIZATION.

**crocidolite** See ASBESTOS; RIEBECKITE.

**Crocodylia** (class *\*Reptilia*) Order of the ‘ruling reptiles’ (*\*Archosauria*) which includes the crocodiles, alligators, caimans, and gavials (gharials). They are derived from *\*thecodontian* ancestors and are closely related to the *\*dinosaurs* and *\*pterosaurs*. They are first recorded from *\*Triassic* rocks and were the only *\*archosaurs* to survive the *\*Mesozoic* era, although strictly speaking the birds (*\*Aves*) are also archosaurs.

**Croixian** An *\*epoch* (501–488.3 Ma ago) of the Upper *\*Cambrian* of N. America, equivalent to the *\*Merioneth*, preceded by the Middle Cambrian, and followed by the *\*Ordovician*.

**Cromerian** 1. A northern European *\*interglacial* *\*stage* dating from about 0.6–0.55 Ma ago. It coincides approximately with the *\*Günz/Mindel interglacial* of the Alps. 2. Temperate deposits of Middle *\*Pleistocene* age found at West Runton, Norfolk, England. Estuarine *\*sands* and silts, and freshwater peat (Upper Freshwater Bed), with a temperate-forest flora, are succeeded by glacial deposits. It is not possible to correlate them with continental European deposits.

**Crommelin** A *\*comet* with an orbital period of 27.89 years; *\*perihelion* date 1 September 1984; perihelion distance 0.743 AU.

**Cromwell current** See EQUATORIAL UNDERCURRENT.

**cronstedite** See CHAMOSITE.

**cross-bedding** See BEDDING; CROSS-STRATIFICATION.

**cross-correlation** In comparing one wave-form with another, the *\*correlation* of two digital traces (i.e. wave-forms that have been *\*digitized*) which are similar but not the same, with one being delayed in time with respect to the other. The operator slides one trace past the other in small time steps (called delays or lags) and at each step the elements of the traces are multiplied together, term by term, and the products added. The maximum value (almost equal to unity) of this cross-correlation is obtained when the two traces are in closest alignment with each other. A value of  $-1$  means the wave-forms are identically matched but opposite in phase; a

value approaching zero indicates low degrees of similarity. The method is extremely useful for detecting wave-forms swamped by noise and, in particular, in the analysis of *\*Vibroscis* records. *See also* [AUTO-CORRELATION](#).

**crosscut** In mining, 1. A tunnel that links two seams, passing through or across the intervening material. 2. A tunnel at right angles to the *\*orebody*. 3. A ventilation tunnel between the mine entrance and one or more air courses.

**cross-cutting relationships** *See* [LAW OF CROSS-CUTTING RELATIONSHIPS](#).

**cross-dating** The matching of tree-ring width patterns and other properties among the trees and fragments of wood from a particular area. This enables the year in which each ring was formed in living trees and recent stumps to be determined accurately, the presence of false rings or the absence of rings in individual specimens being made apparent. By matching ring series from living specimens with those from older (e.g. constructional) timbers, the chronology may be extended backward in time.

**crossed nicols** *See* [CROSSED POLARS](#).

**crossed polars (crossed nicols; XPL, xpols)** The situation in a reflected- or transmitted-light microscope when the pieces of *\*Polaroid* of both the *\*analyser* and substage *\*polarizer* are inserted into the light path through the microscope. The pieces of Polaroid are aligned at right angles to each other so that no light is transmitted to the observer, but a *\*mineral* (or rock) *\*thin* section present on the microscope stage interferes with the light rays into which the light is split by the mineral and produces interference leading to *\*interference colours*. *See also* [BIREFRINGENCE](#).

**cross-hairs** *See* [CROSS-WIRES](#).

**cross-lamination** The smallest structure formed of inclined laminations (*see* [LAMINA](#)), with thicknesses measured in millimetres. *See* [CROSS-STRATIFICATION](#).

**Crossopterygii** (class *\*Osteichthyes*) Subclass of *\*bony* fish comprising both fossil and living lobe-finned or tassel-finned fish, including the *\*Coelacanthiformes* and *\*Rhipidistia*. The former are well known from

\*Palaeozoic and \*Mesozoic rocks, but were thought to have become extinct by the end of the \*Cretaceous until living specimens were netted last century in the Indian Ocean. The Rhipidistia did become extinct, although not before they gave rise, in the \*Devonian, to the \*amphibians. The Crossopterygii are characterized by the fact that all fins (except the tail-fin) are based on movable stalks or lobes. The tail fin is either \*heterocercal or \*diphycercal.

**cross-over distance** ( $x_c$ ) The distance on a seismic \*refraction survey \*time–distance chart at which the \*travel times of the \*direct and refracted waves are the same. This distance also marks the point when the refracted wave overtakes, and thus arrives before, the direct wave. The cross-over distance is related to the refractor depth,  $h$ , and the velocities of the overlying medium and the refractor,  $V_1$  and  $V_2$  respectively, such that  $x_c = 2h[(V_2 + V_1)/(V_2 - V_1)]^{1/2}$ . The value of  $x_c$  will always be greater than twice the refractor depth.

**cross set** See CROSS-STRATIFICATION.

**cross-stratification** A family of primary \*sedimentary structures formed by the migration of the slip-faces of rippled bedforms or of \*bars. It is characterized by inclined laminations (\*foresets) bounded by planar surfaces (planar or tabular cross-stratification), or by scoop-shaped surfaces (trough cross-stratification). The foresets \*dip at the \*angle of repose of the sediment on the \*ripple slip-face and are oriented in the direction of migration of the ripple (see PALAEOCURRENT ANALYSIS). Tabular cross-stratification is produced by the migration of straight-crested, asymmetrical ripples or \*sand waves. Trough cross-stratification is generated by the migration of \*linguoid ripples or \*dunes. The term ‘cross-lamination’ is applied to cross-stratification formed by the migration of ripples; ‘cross-bedding’ is used for cross \*strata formed by the migration of large-scale forms such as dunes, sand waves, or bars. The term ‘cross set’ is used to define the cross-stratification preserved between any upper and lower bounding surface. Where the original bedform which produced the cross set is preserved and forms the upper bounding surface to the set, the term ‘form set’ is used. A number of cross sets preserved within a single bed are called a ‘coset’.

**cross-well seismic** A technique, used in prospecting for crude oil and natural gas, in which a powerful sound is produced at different levels in one well and its vibrations recorded in one or more other wells. The character of the received vibrations provides information about the rock structures between the wells.

**cross-wires (cross-hairs)** Two very thin, black lines or pieces of wire set at right angles to each other in the *\*eyepiece* of a microscope. They are normally positioned N–S and E–W, parallel to the alignment of the *\*polarizer* and *\*analyser*. Many optical properties, e.g. *\*extinction* angles, are measured with respect to the position of the cross-wires.

**crotoquina** See KROTOVINA.

**crown group** In *\*cladistic* analysis, the extant taxa descended from a common ancestor. *Compare* STEM GROUP.

**crude oil** See PETROLEUM.

**Crudinian** The basal *\*stage* in the *\*Devonian* of Australia, underlain by the *\*Silurian*, overlain by the *\*Merionsian*, and roughly contemporaneous with the *\*Gedinnian* stage in Europe.

**crumb structure** Type of soil structure in which the structural units or *\*peds* have a spheroidal or crumb shape. Crumb structure is more often found in porous than granular organo-mineral surface *\*soil horizons*, and provides optimal pore space for soil fertility.

**crura** See CRUS.

**crus** (*pl.* **crura**) Part of the calcified brachial support (*\*brachidium*) in *\*brachiopods* which attach the brachium to the interior *\*shell* beak region of the brachial valve. See BRACHIA.

**crushing** In mineral processing, the breaking up of large particles by compression between two surfaces. Primary crushing produces pieces 100–50 mm in diameter; secondary crushing reduces those to 10 mm.

**crust 1.** The thin outermost solid layer of the Earth. It represents less than 1% of the Earth's volume, and varies in thickness from approximately 5 km beneath the oceans to approximately 60 km beneath mountain chains. Most of the *\*terrestrial* planets have a solid surface, generally considered to be of


different composition to the underlying, higher-density rocks, and regarded as crust. *See also* CRUSTAL ABUNDANCE OF ELEMENTS; CONTINENTAL CRUST; OCEANIC CRUST. 2. A surface soil layer, sometimes slightly cemented with calcium carbonate, silica, or iron oxide, which may be from a few millimetres to many tens of millimetres in thickness but which is always harder and more compact than the soil below. Crusts are now being produced by mechanical action or \*pedogenesis, mainly in arid environments; less commonly they are relict or fossil features exhumed on the soil surface.

**Crustacea** (crustaceans, crabs, lobsters, shrimps, slaters, woodlice, barnacles) (phylum \*Arthropoda) Diverse subphylum of mandibulate arthropods, the body usually divided into three parts: head, thorax, and abdomen. In some crustaceans (e.g. crayfish) the head and thorax may be joined to form the cephalothorax. The head bears two pairs of antennae, one pair of \*mandibles, and two pairs of maxillae. The limbs are \*biramous, and are adapted for a wide range of functions. Closely placed \*setae on the limbs function as filters in filter-feeding species. Respiratory gills are situated on the appendages, but vary greatly in location and number; they are absent only in very small species. In addition to the antennae, sense organs include a pair of compound eyes, and a small, dorsal, median, nauplius eye, comprising three or four closely applied ocelli (clusters of photoreceptors). The nauplius eye, characteristic of crustacean larvae, is absent in many adults; and some groups lack the compound eyes. Mainly marine, but there are many freshwater species, and a relatively small number have invaded the land. Four classes of crustaceans have an important fossil record. The Malacostraca (crabs, lobsters, woodlice, etc., \*Cambrian to Recent) includes the earliest crustaceans of the subclass Phyllocarida. The Branchiopoda (similar to modern water fleas, Lower \*Devonian to Recent) are valuable \*index fossils in non-marine strata. The \*Cirripedia (barnacles) occur from Upper \*Silurian to Recent, and the \*Ostracoda from Lower \*Cambrian to Recent. The living class Cephalocarida (e.g. *Hutchinsonella*) is thought to be closest to the ancestral crustacean stock, but the group is without any unequivocal fossil representative. Hexapods (\*Hexapoda) are believed to be descended from a crustacean, and hexapods and insects are grouped in the \*clade Pancrustacea.

**crustaceans** See CRUSTACEA.

**crustal abundance of elements** The Earth's \*crust has an average density of 2800 kg/m<sup>3</sup> and a thickness varying from about 30 km below the continents (up to 60 km beneath some mountains) to 5 km beneath the oceans. By mapping the major rock types and averaging their composition the abundance of elements can be estimated. The crust is enriched in \*incompatible elements (e.g. K and Rb) as well as \*lithophile elements, but a few elements predominate, especially in \*silicate minerals, while some \*ore metals are rare (e.g. Cu and Sn). Because the crust was formed from material extruded from the Earth's mantle, it is to be expected that the mantle is depleted in 'crustal' components. Oxygen (O) constitutes almost 50% of the Earth's crust by weight and is the most abundant element. Other major elements include: silicon (Si), which is the second most abundant, constituting 27.72% of the crust by weight; aluminium (Al) third; sodium (Na); magnesium (Mg); calcium (Ca); and iron (Fe). Other elements, including such desired metals as gold (Au), silver (Ag), and platinum (Pt), are rare in the crust.

**Cruziana** The name given to the \*tracks made by an animal (some by Trilobita) as it crawled over the sea floor. The tracks are bilobed in appearance, divided along the mid-line, and each half is scored by fine grooves created by the action of the walking limb. See REPICHNIA.



**Cruziana**

**cryergic** Applied to the work of ground \*ice, and so includes \*frost heaving, \*frost wedging, and the thaw processes. The term has been used as

a synonym for **\*periglacial**.

**crylic** See **PERGELIC**.

**cryoconite** A layer of windblown debris and **\*aerosol** that accumulates as dust on the surface of an **\*ice** sheet or **\*glacier**. The dust may contain nutrients and microbial cells, allowing biological processes to occur, and it may also contain small amounts of soot and other dark-coloured matter that lowers the **\*albedo**, causing the snow or ice beneath the dust to melt, forming holes called cryoconite holes. Cryoconite was first described in 1870 by the Finnish geologist and explorer Nils Adolf Erik Nordenskiöld (1832–1901) while travelling over the Greenland ice cap.

**cryoconite hole** See **CRYOCONITE**.

**Cryogenian** A period of the **\*Proterozoic** eon that began 850 Ma ago and ended 630 (+5, –30) Ma ago. Its start is a fixed date and its end is defined by the termination of the Varinger (Marinoan) glaciation and a change in the <sup>13</sup>C content of rocks. The Cryogenian followed the **\*Tonian** period and preceded the **\*Ediacaran** period.

**cryogenic** 1. Applied to: features or materials generated by the action of ice; 2. instruments operating a few degrees above absolute zero (0 K = –273.15 °C).

**cryogeyser** A jet-like eruption of **\*volatiles** with entrained dust and ice particles, but without liquid, that occurs on several bodies in the **\*solar system**, including **\*Enceladus**, **\*Europa**, **\*Mars**, and **\*Triton**.

**cryolite** **\*Mineral**, Na<sub>3</sub>AlF<sub>6</sub>; sp. gr. 3.0; **\*hardness** 2.5; **\*monoclinic**; colourless to white, sometimes brown to reddish; white **\*streak**; **\*vitreous** to greasy **\*lustre**; **\*crystals** rare, but cubic-like and hexahedral, also occurs **\*massive**; no **\*cleavage**, basal parting and poor prismatic parting; occurs in **\*pegmatites** that have been enriched with fluorine, and in association with **\*siderite**, **\*quartz**, **\*galena**, **\*chalcopyrite**, **\*fluorite**, and **\*cassiterite**. The powder becomes almost invisible in water due to its low **\*refractive index**. Synthetic cryolite is used as a flux in the production of aluminium and enamels.

**cryonival** Applied to the set of geomorphological processes comprising **\*cryergic** and **\*nival** mechanisms.

**cryopediment** A bench-like land-form, cut indiscriminately across bedrock, and confined to past or present **\*periglacial** environments. Its position on the lower part of a hillslope is the main criterion for distinguishing it from a **\*cryoplanation** terrace. It is the periglacial analogue of the warm desert **\*pediment**.

**cryoplanation** The reduction of relief to a gently undulating land surface under periglacial conditions. Equivalent to extensive **\*altiplanation**.

**Cryosat-2** See **EARTH EXPLORER OPPORTUNITY MISSION-2**.

**cryosols** A reference soil group in the **\*World Reference Base for Soil Resources** classification scheme. Cryosols have **\*permafrost** horizons within 100 cm of the surface.

**cryosphere** That part of the Earth where the surface is frozen, comprising the area covered by **\*ice** sheets and **\*glaciers**, **\*permafrost** regions, and sea areas covered by ice, at least in winter.

**cryoturbation** See **GELITURBATION**.

**cryovolcanism** Volcanism occurring at low temperatures, where liquid erupts through an overlying crust of ice. The liquid probably consists of brine containing magnesium and sodium sulphates; at extremely low temperatures ammonia may also be present. Cryovolcanism is known to be active on **\*Enceladus**, **\*Europa**, and **\*Ganymede**.

**Cryptic** A **\*period** of the **\*Priscoan**, dated from about 4567.17–4150 Ma ago (Int. Commission on Stratigraphy, 2004).

**cryptocrystalline** Applied to a very fine aggregate of crystals in an **\*igneous** rock, and to minerals in which the individual crystals are too fine to be distinguished even under a petrological microscope. They will produce a diffraction pattern with X-rays. Rapid cooling of a **\*magma** as it is extruded as a lava often produces cryptocrystalline aggregates of minerals.

**cryptodome** An uplifted area, approximately dome-shaped, caused by the intrusion of viscous **\*magma**.

**Cryptodonta** (phylum **\*Mollusca**, class **\*Bivalvia**) Subclass of bivalves most of which have thin, equivalve shells composed of **\*aragonite**. They

have an *\*amphidetic* to *\*opisthodontic* external ligament. The *\*hinge* plate is narrow or absent, with most forms having a toothless hinge margin; some members are *\*taxodont*. They have an *\*infaunal* mode of life. They first appeared in the *\*Ordovician*.

**cryptoperthite** See PERTHITE.

**cryptotephra** Microscopic fragments of volcanic ejecta.

**Cryptozoic (Archaean, Azoic)** A period of time equivalent to the combined *\*Hadean*, *\*Archaean*, and *\*Proterozoic* i.e. the former *\*Precambrian*, 4567.17–542 Ma ago (Int. Commission on Stratigraphy, 2004).

**crystal** A homogeneous, ordered solid, having naturally formed plane faces and a limited chemical composition. Crystals have definite geometric forms that reflect the arrangement in *\*lattices* of the atoms of which they are composed. See CRYSTAL CLASS; CRYSTAL SYMMETRY.

**crystal class (point group)** Crystals are formed by the repetition in three dimensions of a *\*unit* cell structure, defined by lattice points in space. There are only 32 ways to arrange the space lattices in terms of symmetry elements, and these are called the ‘crystal classes’.

**crystal face** One of the relatively flat surfaces by which a crystal is bounded. Faces are produced naturally during the process of crystal growth. Cut and polished *\*gemstones* are bounded by plane faces which are often produced artificially and which, therefore, are not crystal faces.

**crystal-field theory** A theory describing the behaviour of elements with partly filled d- or f-orbitals. In *\*geochemistry* it is particularly concerned with the crystal-field (*\*ligand*-field) effects on the first transition metal *\*ions*, i.e. Sc, Ti, V, Cr, Mn, Fe, Co, Ni, and Cu. If the *\*anions* surrounding a *\*cation* are considered as point charges then the electrostatic interactions between them will vary according to: (a) ionic distance; (b) the strength of the charge; and (c) the *\*coordination* number (how anions are distributed around the cation). Some of the five d-orbitals will have maximum electron density nearer to the anion than others. Interactions between cation and anion will affect the energies of the d-orbitals and cause a separation of energy levels between orbitals (crystal-field splitting).

**crystal group** See CRYSTAL SYSTEM.

**crystalline** A general term, applied to \*metamorphic or \*igneous rocks formed by the process of crystallization from solid or liquid precursors (although certain \*sedimentary rocks, e.g. some \*limestones, may also be made up of crystalline grains, cemented by crystalline \*cement).

**crystalline carbonate** A \*carbonate rock whose original sedimentary texture has been recrystallized (see RECRYSTALLIZATION). See DUNHAM CLASSIFICATION.

**crystalline limestone** See DUNHAM CLASSIFICATION.

**crystalline remanent magnetization** See CHEMICAL REMANENT MAGNETIZATION.

**crystallite** A microscopic, often skeletal \*crystal which represents the initial form of crystalline material just after \*nucleation has taken place in a \*magma. Crystallites are usually preserved in volcanic \*glass, which represents a quenched magma. \*Quenching is the only way in which crystallites can be preserved.


**crystalloblastic** A metamorphic \*texture characterized by the mutual interference of polygonal grains which meet at approximately 120° triple junctions. The texture is produced during solid-state \*crystal growth as a result of increasing the pressure on or temperature in the rock system. As the grains grow, they have to compromise in their competition for space and hence form the typical polyhedral grains with triple junctions. The texture can be '\*isotropic' (no grain alignment) where there is no directed \*stress during \*metamorphism, or '\*anisotropic' (with grain alignment) where there is a directed stress during metamorphism.

**crystallographic axes** Axes used to define the position of a \*crystal face in space by the intercepts of the face on three (or four) imaginary lines.

**crystallography** The study of crystals, including their form, structure, habit, and symmetry.

**crystal symmetry** In well-formed crystals, the symmetrically arranged faces reflect the internal arrangement of atoms. The symmetry of individual crystals is determined by reference to three elements. The plane of

symmetry (also called the ‘mirror plane’ or ‘symmetry plane’) is a plane by which the crystal may be divided into two halves which are mirror images of each other. The axis of symmetry is a line about which a crystal may be rotated through  $360^\circ/n$  until it assumes a congruent position;  $n$  may equal 2, 3, 4, or 6 (but not 1), depending on the number of times the congruent position is repeated. These correspond respectively to 2-fold (diad), 3-fold (triad), 4-fold (tetrad), and 6-fold (hexad) axes. The centre of symmetry is a central point which is present when all faces or edges occur in parallel pairs on opposite sides of the crystal. Using these elements of symmetry, crystallographers have recognized 32 \*crystal classes and seven \*crystal systems. Symmetry is highest (high symmetry) in the \*cubic system, where many elements are repeated, and lowest (low symmetry) in the \*triclinic system, where only a centre of symmetry may be present (i.e. there may be no plane or axis of symmetry).



Crystal symmetry

**crystal system (crystal group)** One of the seven systems or groups into which crystals can be placed by reference to their *\*crystallographic* axes. It is possible to arrange the 14 *\*Bravais* lattices into seven systems (groups) each characterized by a primitive (*p*) lattice.

**crystal twinning** Feature, common to many single crystals, where the crystal *\*lattice* is differently orientated in two or more parts of the crystal, e.g. twin plane, twin axis, etc., and simple or multiple twins. See ALBITE TWIN; CARLSBAD TWIN; COMPLEX TWINS; CONTACT TWIN; DEFORMATION TWINNING; GENICULATE TWIN; GROWTH TWINNING; INTERPENETRANT TWIN; MIMETIC TWINS; NORMAL TWINS; PARALLEL TWINS; PERICLINE TWINNING; SWALLOWTAIL TWINNING; TRANSFORMATION TWINNING; TWIN AXIS; TWIN LAW; TWIN PLANE.

**crystal zoning** A *\*texture* developed in *\*solid-solution* minerals and characterized optically by changes in the colour or *\*extinction* angle of the mineral from the core to the rim. This optical zoning is a reflection of chemical zoning in the mineral. For example, a *\*plagioclase* can be zoned from a Ca-rich core to an Na-rich rim. Zoning results from the mineral's inability to maintain chemical equilibrium with a *\*magma* during rapid cooling; the zonation represents a frozen picture of the *\*continuous reaction series* for that mineral. Zoning can be of three types, the first two applying mostly to plagioclase feldspars. (a) Normal zoning is where the mineral is zoned from a high-temperature core composition to a low-temperature rim composition. (b) Reverse zoning is where a mineral is zoned from a low-temperature core composition to a high-temperature rim composition. (c) Oscillatory zoning is where the mineral chemistry continuously oscillates between high- and low-temperature compositions going from the core to the rim. Compare CORONA.

**CSES** See CHINA SEISMO-ELECTROMAGNETIC SATELLITE.

**CSSWE** See COLORADO STUDENT SPACE WEATHER EXPERIMENT.

**CST** See CONSTANT-SEPARATION TRAVERSING.

**CTD** An instrument for measuring sea-water conductivity (from which *\*salinity* can be calculated), temperature, and depth (actually, pressure). A sensor unit is lowered through the water on the end of an electrical

conductor cable which transmits the information to indicating and recording units on board ship.

**cube** A hexahedral crystal shape or form which can be referred to three axes of equal length which intersect at right angles.

**CubeRRT** See CUBESAT RADIOMETER RADIO FREQUENCY INTERFERENCE TECHNOLOGY.

**CubeSat** A miniaturized space satellite made from multiples of cubic units 10×10×10 cm in size, with a mass of no more than 1.33 kg per unit.

**CubeSat for Ions, Neutrals, Electrons & Magnetic fields (CINEMA)** An international 3-unit \*CubeSat mission developed by the CINEMA Consortium comprising teams from universities in the US, UK, and South Korea to supply space weather measurements from low Earth orbit. The \*nanosatellite was launched on 13 September 2012, from California.

**CubeSat Infrared Atmospheric Sounder (CIRAS)** A \*CubeSat mission developed by the Jet Propulsion Laboratory, California Institute of Technology, to measure upwelling \*infrared radiation, which can be used to retrieve lower-tropospheric temperature and water vapour for weather and climate studies. The mission is scheduled to be launched in late 2018.



<https://www.jpl.nasa.gov/cubesat/missions/ciras.php>


- Mission information.

**CubeSat – Launch 1** The first multiple launch of \*CubeSats, involving about six CubeSats and a primary commercial payload. It was launched on 30 June 2003, from Plesetsk, Russia. It was followed by Launch 2, comprising three CubeSats, on 27 October 2005, from Plesetsk, and Launch 4, of seven CubeSats, on 17 April 2006, from Baikonur, Kazakhstan. Launch 3 failed shortly after launch on 26 July 2006.

**CubeSat Radiometer Radio Frequency Interference Technology (CubeRRT)** A collaborative \*CubeSat mission between Goddard Space Flight Center (NASA), Ohio State University, and the Jet Propulsion Laboratory (NASA) to develop techniques for detecting and discarding

radio-frequency interference. It was launched on 21 May 2018, from Virginia, USA.


**cubic (isometric)** One of the seven *\*crystal* systems and the one with the highest number of symmetry elements present. It is characterized by four triad axes of symmetry, and requires three *\*crystallographic* axes of equal lengths intersecting at right angles.



**Cubic**

**cubichnia** A category of *\*trace fossils* that mark the temporary resting places of various organisms.

**cuesta** Asymmetric land-form consisting of a steep scarp slope, and a more gentle dip (or back) slope. It is typical of areas underlain by strata of varying resistance that are dipping gently in one direction, and is intermediate between the flat-topped *\*mesa* and *\*butte*, and the more symmetric ridge form of the *\*hog's* back.



**Cuesta**

**culmination** The highest antiformal point (see [ANTIFORM](#)) of a *\*crest* line along all non-*\*cylindroidal* folds.

**cumec** See [DISCHARGE](#).

**cummingtonite** A *\*monoclinic*, Mg-rich member  $(\text{Mg,Fe})_2(\text{Mg,Fe})_5[\text{Si}_4\text{O}_{11}]_2(\text{OH,F})_2$  of the calcium-poor *\*amphiboles*, along with grunerite ( $\text{Mg} < \text{Fe}$ ) with which it forms a series, similar to *\*anthophyllite*–gedrite; sp. gr. 3.1–3.6; *\*hardness* 5.5; dark green; occurs in amphiboles and some *\*intermediate \*igneous* rocks.

**cumulate** Applied to *\*igneous \*intrusive* rocks formed by the accumulation of crystals as a result of *\*gravity settling*. It is typical of layered intrusions and common in some differentiated *\*meteorites*. The early-formed minerals are called ‘cumulus’ minerals and show a regular variation in composition with their height in the intrusion. See [MINERAL LAYERING](#).

**cumulative percentage curve** A graphical plot in which size classes are plotted against the percentage frequency of the class plus the sum of the percentages in preceding size classes. When plotted on normal graph paper, the cumulative frequency curve resembles an S-shape. When plotted on a normal-probability scale, the cumulative percentage data appear in a series of straight-line segments, each with a different gradient.

**cumuliform** Applied to clouds resembling *\*cumulus* or *\*cumulonimbus*.

**cumulonimbus** From the Latin *cumulus* meaning ‘heap’ and *nimbus* meaning ‘cloud’, a genus of clouds of bulging, dense form, often towering to great height in unstable air. Young clouds have distinctive fibrous or lined features; older, glaciated types, with abundant ice crystals, are lustrous. Typically, the upper parts are spread into *\*incus* or plume features. The cloud base is dark and usually gives rise to precipitation, often with *\*virga*. See also CLOUD CLASSIFICATION.

**cumulus** 1. From the Latin *cumulus* meaning ‘heap’, a genus of dense, isolated, and clearly defined clouds with vertical growth in bulges or domes, and a flattened, darker base. Sharply outlined, bulging cloud tops indicate vigorous growth. Occasionally a more ragged form occurs. See also CLOUD CLASSIFICATION. 2. See CUMULATE.

**Cunninghamian** A *\*stage* in the *\*Devonian* of Australia, underlain by the *\*Merionsian*, overlain by the *\*Condobolinian*, and roughly contemporaneous with the *\*Emsian* and *\*Eifelian* stages of Europe.

**Cupid (Uranus II)** A lesser satellite of *\*Uranus* with a radius of 5 km.

**cupola** A small, dome-shaped, satellite intrusion (see INTRUSIVE) projecting upwards from the main body of a larger intrusion or *\*batholith*.

**cuprite (red copper ore)** *\*Mineral*,  $\text{Cu}_2\text{O}$ ; sp. gr. 5.8–6.1; *\*hardness* 3.5–4.0; *\*cubic*; red to nearly black; brownish-red *\*streak*; *\*adamantine* to *\*sub-metallic* *\*lustre*; crystals usually octahedral and *\*acicular*, but can be granular and *\*rhombohedral*; *\*cleavage* poor {111}; usually occurs in the oxidized zone of copper deposits, as a *\*secondary* mineral, and associated with *\*malachite* and *\*azurite*. It is a minor *\*ore mineral* for copper.

**Curie point** See CURIE TEMPERATURE.

**Curie symmetry principle** A principle formulated in 1894 by Pierre Curie (1859–1906) stating that: (a) If certain causes yield known effects, the symmetry elements of the causes should be contained in the effects generated; (b) If there is an absence of symmetry elements in the known effects, that absence should be contained in the causes which generated those effects; (c) The effects may have higher symmetry than the causes

which generated them, therefore the converse of (a) and (b) is not true. See also SANDER'S SYMMETRY PRINCIPLE.

**Curie temperature (Curie point)** The temperature at which thermal vibrations prevent quantum-mechanical coupling between atoms, thereby destroying any \*ferromagnetism (in the wide sense). Typical Curie temperatures are 675 °C for \*hematite and 575 °C for \*magnetite.


**Curie–Weiss law** A law that describes the \*magnetic susceptibility ( $X$ ) of a ferromagnet (see FERROMAGNETISM) at temperatures above the \*Curie temperature.  $X = C/(T - \theta)$  where  $C$  is a constant for each material,  $T$  is the temperature in kelvins, and  $\theta$  is the Curie temperature in kelvins. The law was formulated by the French physicists Pierre Curie (1859–1906) and Pierre-Ernest Weiss (1865–1940).

**current electrode** An electrode by which an electrical current enters or leaves a conducting medium. Two current electrodes are used in a variety of \*electrode configurations in \*resistivity surveys (\*vertical electrical sounding and \*constant-separation traversing) and \*induced-polarization surveys.

**current meter** An instrument for measuring the speed of flow in a watercourse. The most common type of current meters relate current speed to the rate at which an impeller is rotated by the flowing water.

**Curvolithus** A genus of \*trace fossils comprising short, horizontal burrows close to the surface.

**cusate foreland** Large, triangular area of coastal deposition, which is dominated by many shingle ridges, and is often terminated landward by poorly drained terrain. It is the result of a long episode of local marine aggradation under wave advance from two dominant directions. Dungeness, on the south coast of England, is a typical example.



### Cusped foreland

**cuspidate** Having a sharp tip or point.

**cutan** 1. (clay films, clayskins, argillans, tonhäutchens) Deposited skin or coating of material on the surfaces of **\*peds** and stones, which is usually composed of fine, **\*clay**-like soil particles which have been moved down through the soil. 2. A complex, insoluble biopolymer with waterproofing qualities found in the **\*cuticle** of plants.


**cut and fill** Levelling of material by excavation in one place and its deposition in an adjacent place, to produce a uniform height for roads, railways, canals, etc. In mining, the back-filling of excavations by waste materials.

**cuticle** 1. The impervious covering to the outer walls of the epidermal cells of aerial plant organs, composed of **\*cutin**, **\*cutan**, or a mixture of both. 2. The outer layer of an insect, secreted by epidermal cells; it has a complex structure that varies according to species. See **SKELETAL MATERIAL**.

**cutin** A complex biopolymer comprising a mixture of fatty-acid derivatives with waterproofing qualities found in the **\*cuticle** of plants.

**cutinite** See **COAL MACERAL**.

**cut-off (ox bow)** Section of a river channel that no longer carries the main discharge. Its abandonment results from meander development associated with lateral channel migration across a **\*floodplain**. Channel length is shortened by contact at the neck of a loop, which becomes a cut-off.



**Cut-off**

**cut-off grade** Lowest grade or assay value of ore in a deposit that will recover mining costs; the cut-off grade determines the workable tonnage of an ore.

**cut-off high** **\*Anticyclone** isolated from the main subtropical belt of high pressure and around which the main flow of the upper westerlies is diverted, causing a blocking situation in middle latitudes.

**cut-off low** **\*Cold low** in mid-latitudes (occasionally almost in subtropical latitudes) where it is cut off from the main subpolar belt of low pressure. Sometimes a cut-off low occurs with a **\*cut-off high** over the higher

latitudes, typically in **\*blocking** situations. In summer, weather associated with such slow-moving lows is unsettled and thundery.

**cut-off trench** See DIAPHRAGM WALL.

**cutting bar** See CUTTING BOOM.

**cutting boom (cutting bar)** Part of a cutting machine used to undercut or overcut coal or other stratified material. The cutting machine consists of a rubber-tired base and a cutter bar 3 m long and 380 mm wide, with a chain running along its edge and holding the **\*bits**. It is driven by an electric motor.

**Cuvier, Chrétien Frédéric Dagobert ('Georges'), Baron** (1769–1832) *Professor* at the National Museum of Natural History in Paris, where he developed the discipline of comparative anatomy and applied it to the study of fossil quadrupeds in his *Recherches sur les ossements fossiles de quadrupèdes* (1812). Its *Discours préliminaire* (1811), in which he outlined a theory of multiple catastrophes, became one of the most influential scientific treatises of the early 19th century. With **\*Brongniart** he produced a map of the Paris basin, having used his fossil studies to work out the stratigraphy. Cuvier opposed Lamarckian evolution, arguing that species were stable and became extinct because of 'revolutions' (changes in sea level, etc.). See CATASTROPHISM; DILUVIALISM; LAMARCKISM.

**CVG** See CONSTANT-VELOCITY GATHER.

**CVS** See CONSTANT-VELOCITY STACK.

**cwm** See CIRQUE.

**cyanobacteria** A large and varied group of bacteria which possess chlorophyll *a* (carried on specialized membranes (thylakoids) within the cells) and carry out **\*photosynthesis** in the presence of light and air with concomitant production of oxygen. They do not have chloroplasts. Cyanobacteria were formerly regarded as **\*algae** (division Cyanophyta) and were called 'blue-green algae'. Fossil cyanobacteria have been found in rocks almost 3000 Ma old and they are common as **\*stromatolite** colonies in rocks 2300 Ma old. They are believed to have been the first oxygen-producing organisms and to have been responsible for generating the oxygen in the atmosphere, thus profoundly influencing the subsequent

course of evolution. The organisms may be single-celled or filamentous, and may or may not be colonial. Some are capable of a gliding motility when in contact with a solid surface. Many species can carry out the fixation of atmospheric nitrogen.

**Cyanophyta** See CYANOBACTERIA.

**cyanophyte** Cyanobacterium (blue-green alga). See CYANOBACTERIA.

**cycad** See CYCADOPSIDA.

**Cycadaceae** A monotypic family comprising the genus *Cycas*, with a fossil record from the \*Permian. They are the most widely distributed of all extant cycads.

**Cycadales** In some classifications, an order of \*gymnosperms comprising the cycads. These are included in the order \*Cycadopsida.

**Cycadopsida (cycads)** A class of \*gymnosperms comprising plants with leaves and habit similar to those of palm trees, although some species are quite small. Most cycads bear large, coloured, female or male cones. Pollen grains have motile spermatozoa within them, which is a very primitive feature. Formerly they were much more important and, following their appearance in the \*Permian, remained important members of the world's \*Mesozoic floras. Their reduction was particularly marked in the Late \*Cretaceous as they were progressively displaced by \*angiosperm trees. The survivors are regarded as 'living fossils'.

**cycle of erosion** See DAVISIAN CYCLE.

**cyclic sedimentation** A style of sedimentation where the sequence of sedimentary \*facies develops in a regular and repeated manner. The cycle of sediments may consist of two or more facies, and the cycles may be symmetrical or asymmetrical (e.g. A → B → C → D, A → B → C → D; or A → B → C → D → C → B → A, A → B → C → D → C → B → A). The cyclicity may be the product of repeated episodes of marine \*transgression and \*regression linked to climatic cycles (see MILANKOVITCH CYCLES), or of the lateral shifting of depositional environments, or due to repeated cycles of tectonic activity in source areas.

**cycling pool** See ACTIVE POOL.

**cyclogenesis** Formation and strengthening of cyclonic air circulation, tending to form or deepen depressions. The process is associated with upper-air divergence over or near the frontal zone.

**cyclolysis** The processes of dissipation of cyclonic air circulation around a depression, or the weakening of lesser cyclonic features.

**cyclone** **1.** The name given to a **\*tropical cyclone** that develops in the Indian Ocean and Bay of Bengal. Cyclones usually travel north, on tracks that carry them over Bangladesh. **2.** See **DEPRESSION**.

**Cyclone Global Navigation Satellite System (CYGNSS)** A constellation of eight **\*microsatellites** forming part of the **\*NASA** Earth System Science Pathfinder programme that aim to improve predictions of extreme weather through studying inner core processes in **\*tropical cyclones**. The satellites were launched on 15 December 2016, from Cape Canaveral, Florida.

**cyclopean concrete** See **CONCRETE DAM**.

**cyclopel** A laminated sediment, formed in a glaciomarine environment, that consists of layers of silt and mud. *Compare* **CYCLOPSAM**.

**cyclopsam** A laminated sediment, formed in a glaciomarine environment, that consists of layers of sand and mud. *Compare* **CYCLOPEL**.

**cyclosilicate (ring silicate, metasilicate)** Applied to the 'ring' structure of linked  $\text{SiO}_4$  **\*tetrahedra** in **\*silicate** minerals. The ratio of Si to O is 1:3 and there may be three, four, or six  $\text{SiO}_4$  tetrahedra linked together. Examples of cyclosilicates include **\*tourmaline**, **\*beryl**, **\*axinite**, and **\*cordierite**.

**cyclostratigraphy** The study of stratified rocks (**\*stratigraphy**) in relation to cycles of formation and destruction. First-order cycles, over periods of 200–400 million years, are linked to the formation and breaking apart of supercontinents. Second-order cycles (super cycles), lasting 10–100 million years, are linked to **\*plate tectonics**. Third-order cycles (mesothems), over 1–10 million years, are linked to plate movements and glacial and interglacial episodes. Fourth-order cycles (**\*cyclothem**s) are linked to the **\*Milankovitch** solar radiation curve. Fifth-order cycles (minor cycles), over 0.01–0.2 million years, are linked to the Milankovitch curve and astronomic forcing.

**cyclothem** Unit, or given set of deposits, laid down as a result of either cyclic or rhythmic sedimentation. In a cyclic sequence the cyclothem would represent the whole succession, e.g. 1234321, whereas in a rhythmic sequence the cyclothem units would be repeated, e.g. 12341234. See CYCLOSTRATIGRAPHY.

**CYGNSS** See CYCLONE GLOBAL NAVIGATION SATELLITE SYSTEM.

**cylindrical** See SOLITARY CORALS.

**cylindrical trend** See BOXCAR TREND.

**cylindroidal fold** A *fold* which maintains its two-dimensional profile in a third dimension, like a cylinder (as compared to a circle).

**cymatogeny** Large-scale regional uplifting of the *crust* that produces *domes* and *basins*.

**cyrenoid** Applied to a type of *heterodont dentition* that occurs in bivalve molluscs (*Bivalvia*), in which there are three *cardinal* teeth present in each valve. Compare LUCINOID.

**cyrtconic** Applied to the conch of a *cephalopod* when it is a curved, tapering cone.

**Cystoidea (cystoids)** (subphylum Blastozoa, phylum *Echinodermata*) Extinct group of echinoderms, ranking in some classification schemes as a class or superclass, and ranging from the Lower *Ordovician* to the Upper *Devonian*. The body is spherical or ovoid, covered with a plated *theca*, and was usually attached to the substratum either directly or by a short stalk. The thecal plates are calcitic and perforated by characteristic pore structures that functioned in respiration. The plates are arranged in circlets but vary in number from around 13 to more than 100, giving an irregular appearance. The mouth, on the upper surface, is surrounded by *ambulacral* food grooves that in some cystoids were equipped with food gathering appendages called *brachioles*. These are rarely preserved. In many cases *radial symmetry* is poorly developed.

**cystoids** See CYSTOIDEA.

**Cytherean** An alternative name for 'venusian'. In Greek mythology, Cythera was an alternative name for Aphrodite, the goddess of love,

identified with the Roman Venus. It is derived from Cythera, in Cyprus, near the place where, according to the legend, Aphrodite was born in the sea.

# D

**dacite** A light-coloured, fine-grained *\*igneous* rock containing 63–70 wt. % SiO<sub>2</sub>, as well as *\*plagioclase* feldspar, *\*alkali* feldspar, *\*quartz*, *\*biotite*, and *\*hornblende* as *\*essential* minerals, and *\*sphene*, *\*apatite*, and *\*magnetite* as *\*accessory minerals*. Plagioclase feldspar dominates over alkali feldspar in a 2:1 ratio. Dacites are the volcanic equivalents of *\*granodiorites* and are found as members of the *\*calc-alkaline* series, typically developed as eruptive products on the continental side of oceanic-continental *\*plate* *\*subduction zones*. They are often surprisingly coarse-grained for lavas.

**Dactyl** See *IDA*.

***Dactylodites ottoi*** A feeding *\*trace fossil* (*\*fodinichnia*) in the form of a J-shaped burrow.

***Daedalus*** See *SPREITEN*.

***dagalas*** See *KIPUKA*.

**Dalmatian-type coast** A *\*Pacific-type* coast that has been partially drowned by a rise in sea level, producing many long, narrow islands lying parallel to the shore.

**Dalradian** The last, or youngest, stratigraphic unit of the *\*Precambrian* of Scotland and Ireland.

**Dalslandian** A *\*stage* of the Middle–Upper *\*Proterozoic*, from about 1600–650 Ma ago, and equivalent to the *\*Jotnian* (Van Eysinga, 1975). The name is not used in the 2004 revised timescale of the Int. Commission on Stratigraphy.

**Dalslandian orogeny** (**Gothian orogeny**, **Gothic orogeny**, **Sveconorwegian orogeny**) An eastward continuation of the **\*Proterozoic \*Grenvillian** phase of mountain building that affected what are now southern Sweden and southern Norway. It occurred about 1050–1100 Ma ago and may have been caused by **\*subduction** along the north-western margin of the **\*Iapetus** Ocean, by the closing of an earlier ocean, or by rifting (see **RIFT**) that ended before producing significant **\*sea-floor** spreading.

**Daly, Reginald Aldworth** (1871–1957) A mathematician who turned to geology and the study of **\*igneous** rocks and **\*volcanoes**, Daly was a professor at Harvard University. In the 1920s he was one of the very few supporters in America of the theory of **\*continental drift**. He proposed that drift was caused by continental ‘downsliding’ due to gravity, above a molten **\*mantle** which had the properties of glass. It was Daly who first suggested, in 1936, that **\*submarine** canyons may be excavated by suspension currents. *See also* **TURBIDITY CURRENTS**.

**DAMPE** *See* **DARK MATTER PARTICLE EXPLORER**.

**damping** A slowing down or prevention of oscillation due to the dissipation of the kinetic energy of oscillation. Friction will dampen a mechanical system, and electromagnetic damping uses eddy currents to oppose motion. *See also* **CRITICAL DAMPING**.

**Dana, James Dwight** (1813–95) American mineralogist and geologist, responsible for the term ‘geosyncline’, who proposed that the Earth is contracting as it cools, causing deformation which is concentrated at continental margins on the limbs of geosynclines, thus forming mountains, metamorphic belts, etc.

**Danian** **1.** The earlier of two **\*ages** in the **\*Palaeocene** epoch, preceded by the **\*Maastrichtian** (Late **\*Cretaceous**), followed by the **\*Selandian**, and dated at 65.5–61.7 Ma ago (Int. Commission on Stratigraphy, 2004). **2.** The name of the corresponding European **\*stage** which, in Denmark, is characterized by chalky **\*limestone** rich in **\*reef**-dwelling organisms. It is roughly contemporaneous with the Montian (Belgium), Danian (see (3) below) and lower **\*Ynezian** (N. America), lower **\*Teurian** (New Zealand), and **\*Wangerriplan** (Australia). In the past, because of its chalk **\*facies**,

some authors considered the Danian to be Upper Cretaceous, and succeeded in the Lower *\*Tertiary* by the Montian. **3.** The basal stage in the Lower Tertiary of the west coast of N. America, overlain by the Ynezian and roughly contemporaneous with just the lower part of the Danian stage in Europe.

**Dansgaard–Oeschger event (DO event)** A brief warm period that occurred during the *\*Devensian* glaciation. There were many DO events, lasting from hundreds to several thousand years, and they may also have occurred during the *\*Ipswichian* interglacial. The existence of DO events was discovered in the early 1980s by the Danish climatologist Willi Dansgaard (1922–2011) and the Austrian climatologist Hans Oeschger (1927–98).

**darcy** The unit of intrinsic *\*permeability*, used particularly in the oil industry. One darcy is equal to  $0.987 \times 10^{-12} \text{ m}^2$ .

**Darcy's law** A description of the relationships among factors that determine *\*groundwater* flow, expressed as an equation. At its simplest, Darcy's law states that  $Q = kIA$ , where  $Q$  is the ground water flow,  $k$  the *\*hydraulic* conductivity of the rock,  $I$  the hydraulic gradient (i.e. gradient of *\*hydraulic* head), and  $A$  the cross-sectional area through which flow occurs. The law was formulated in 1856 by the French engineer Henry Philibert Gaspard Darcy (1803–58).

**Dark Matter Particle Explorer (DAMPE)** A Chinese space mission to detect electrons and photons in the range of 5 GeV–10 TeV at a resolution of 1.5% at 100 GeV in order to identify possible dark matter signatures. The satellite was launched on 17 December 2015, from Jiuquan Satellite Launch Centre, China, into a *\*Sun-synchronous orbit* at 500 km altitude.

**d'Arrest** A *\*comet* with an orbital period of 6.51 years; *\*perihelion* date 1 August 2001; perihelion distance 1.346 AU.

**Darriwilian** A *\*stage* (468.1–460.9 Ma ago) of the Middle *\*Ordovician* of Australia, underlain by the *\*Ypeenian* and overlain by the *\*Gisbornian*.

**Darwin, Charles Robert** (1809–82) English naturalist, who is remembered mainly for his theory of evolution, which he based largely on observations made in 1832–6 during a voyage around the world on HMS *Beagle*, which was engaged on a mapping survey. After receiving a letter from **Alfred**

**Russel Wallace** (1823–1913) outlining the similar conclusions he had reached, on 1 July 1858 **Sir Charles \*Lyell** and Sir Joseph Dalton Hooker (1817–1911) presented separate papers by Darwin and Wallace at a meeting of the Linnean Society (neither Darwin nor Wallace was present). In 1859 Darwin published a longer account in his book, *On the Origin of Species by Means of Natural Selection*. In this he presented powerful evidence suggesting that change (evolution) has occurred among species, and proposed natural selection as the mechanism by which it occurs. His theory may be summarized as follows: (a) the individuals of a species show variation; (b) on average, more offspring are produced than are needed to replace their parents; (c) populations cannot expand indefinitely and, on average, population sizes remain stable; (d) therefore there must be competition for survival; (e) it is the best-adapted variants (the fittest) that survive and reproduce. Since environmental conditions change over long periods of time, a process of natural selection occurs which favours the emergence of different variants and ultimately of new species (the ‘origin of species’). This theory is known as Darwinism. The subsequent discovery of chromosomes and genes, and the development of the science of genetics, have led to a better understanding of the ways in which variation may be caused. Modified by this modern knowledge, Darwin’s theory is called ‘neo-Darwinism’.

**darwin** A measure of evolutionary rate (introduced by J. B. S. Haldane in 1949), given in units of change per unit time. *See also* HALDANE.

**Dasycladales** (class Ulvophyceae) An order of green **\*algae** in which the thallus contains a single nucleus, becoming multinucleate prior to reproduction. The thallus consists of an erect axis with branches, the whole sharing radial symmetry. The group appeared in the **\*Ordovician**, where elaborate and ornate forms occurred. They were more diverse in the **\*Permian** and **\*Jurassic**, but became reduced in variety during the **\*Cretaceous** and survive now only in the tropics.

**data** *See* DATUM.

**dating errors** *See* ERRORS; ISOCHRON.

**dating methods** During the last century geologists constructed a relative timescale based on **\*correlation** of palaeontological and stratigraphic data.

Depositional rates of *\*sediments* have also been employed as a dating method, but only recently has absolute dating been made possible through the use of radioactive *\*isotopes*. Of the various methods the last is obviously the most precise, but *\*fossils*, *\*lithologies*, and cross-cutting relationships do enable the geologist to give an approximate *\*relative* age in field studies. *See also* ABSOLUTE AGE; RADIOACTIVE DECAY; RADIOMETRIC DATING; ISOTOPIC DATING; RADIOCARBON DATING; DENDROCHRONOLOGY; GEOCHRONOLOGY; GEOCHRONOMETRY; VARVE ANALYSIS.

**Datsonian** A *\*stage* (488.3–485 Ma ago) of the Lower *\*Ordovician* of Australia, underlain by the *\*Payntonian* (Upper *\*Cambrian*) and overlain by the *\*Warendian*.

**datum** (*pl. data*) Something that is known or assumed to be true. *See also* DATUM LEVEL; MARKER BED.

**datum level** A surface or level which is regarded as a base from which other levels can be counted (i.e. a *\*datum*). For example, sea level is often used as a datum level against which the height of land and depth of the sea bed are measured.

**daughter** *See* CLADISTICS; RADIOACTIVE DATING; RADIOACTIVE DECAY.

**daughter minerals** *See also* FLUID INCLUSION.

**Davis, William Morris** (1850–1934) An American geologist from Harvard University, Davis initiated the study of *\*geomorphology*. He evolved the concept of the cycle of erosion (the *\*Davisian* cycle), and described the role of rivers in the evolution of landscape.

**Davisian cycle (cycle of erosion)** Orderly series of stages through which land-forms were believed to pass from their initiation following uplift to their final planation by erosion. The main stages were those of youth, when hillslopes were steep and river profiles irregular; of maturity, when river profiles were smoothly concave-up and incision had markedly slowed; and of old age, when the landscape was reduced to a gently undulating surface or *\*peneplain*. This framework for land-form studies has now largely dropped out of use.

**day degrees (degree days)** The departure of the average daily temperature from a defined base, e.g. 4 °C, the minimum recognized temperature for the

growth of many plant species, or 18 °C, which determines the need for heating in buildings. It is calculated by multiplying the number of days on which the mean temperature is above or below the datum level over a specified period (usually one month) by the sum of the number of degrees by which it deviates during that period. The number of day degrees may be totalled to assess the accumulated warmth of a particular year's growing season for crops. *See also* ACCUMULATED TEMPERATURE; AGROMETEOROLOGY; MONTH DEGREES.

**day length** *See* EARTH ROTATION.

**dB** *See* DECIBEL.

**D days** (geomagnetic) *See also* DISTURBED DAYS.

**dead end** In the tide-dominated region of a *\*delta*, a sinuous tidal channel that has a single connection to the network of channels.

**death assemblage (thanatocoenosis)** An assemblage of *\*fossils* of organisms that were not associated with one another during their lives. The remains were brought together after death, often by the action of currents.

**débâcle** Break-up of river ice in spring over northern Eurasia and N. America. The onset of thaw begins in March in low latitudes and later further north.

**debris fall** *See* TOPPLE.

**debris flow** Slow-moving, sediment gravity flow composed of large *\*clasts* supported and carried by a mud-water mixture. Debris flows occur as overland and submarine mudflows, and as submarine deep-sea deposits. The deposits are poorly sorted and internally structureless; typically they have a pebbly, *\*mudstone* texture. Debris-flow deposits cover many thousands of square kilometres of the *\*abyssal plain* after originating on the *\*continental slope* as slumped material.

**debris slide** A shallow landslide within rock debris, characterized by a displacement along one or several surfaces within a relatively narrow zone. It may take place as a largely unbroken mass, or may be disrupted into several units, each consisting of rock debris.

**debrite** A deposit formed by a *\*debris* flow.

**decalcification** A stage in the development of a soil during which soluble calcium ions ( $\text{Ca}^{2+}$ ) leach from the *soil profile*.

**decay constant** *Radioactive* decay involves only the *nucleus* of the parent atom, and thus the rate of decay is independent of all physical and chemical conditions (e.g. pressure, temperature, etc.). The decay was shown by *Rutherford* to follow an exponential law. The fundamental equation describing the rate of disintegration may be written as:  $-(dN/dt) = \lambda N$ , where  $\lambda$  is the decay constant, representing the probability that an atom will decay in unit time  $t$ , and  $N$  is the number of radioactive atoms present. It is a fundamental assumption in geochronology that  $\lambda$  is a constant and that the only alteration in the amount of daughter or parent in the system is due to radioactive decay. The constant  $\lambda$  is usually expressed in units of  $10^{-10}$  per year (e.g.  $^{235}\text{U}$  is 9.72,  $^{40}\text{K}$  is 5.31,  $^{87}\text{Rb}$  is 0.139, and  $^{238}\text{U}$  is 1.54). The total lifetime of a radioactive parent in a given system cannot be specified; in theory it is infinite. It is a simple matter, however, to specify the time for half of the radioactive parent atoms in a system to decay. This is called the 'half-life' ( $T$ ), which is related to the decay constant by the expression  $T = 0.693/\lambda$ . See also [DECAY CURVE](#).

**decay curve** A graphical representation of the exponential rate at which radioactive disintegration occurs (see [RADIOACTIVE DECAY](#)). If half the parent *nuclide* remains after one time increment, one-quarter will remain after the next (identical) time increment, and so on. A plot of the surviving parent atoms against time in half-lives (see [DECAY CONSTANT](#)) gives a decay curve that approaches the zero line asymptotically. In theory it should never attain zero. The number of surviving parent atoms  $N(t)$  at the end of a number of half-lives ( $n$ ) is simply  $N_0/2^n$ . Plotted as a function of time  $t$ , survivors form a characteristic decay curve, the equation of which is  $N_t = N_0 e^{-\lambda t}$ , where  $e = 2.718$  and  $\lambda$  is the *decay* constant.

**decay index** In *cladistic* analysis, the number of additional steps required to dissolve a given *clade*.

**decay series** *Radioactive* decay of a parent nuclide through a sequence of radioactive daughter nuclides to a final, stable daughter nuclide. Uranium has three naturally occurring *isotopes*:  $^{238}\text{U}$ ,  $^{235}\text{U}$ , and  $^{234}\text{U}$ ; all are radioactive. Thorium exists mainly as one isotope ( $^{232}\text{Th}$ ) which is also

radioactive. In addition, five radioactive isotopes of thorium occur in nature as short-lived, intermediate daughters of  $^{238}\text{U}$ ,  $^{235}\text{U}$ , and  $^{232}\text{Th}$ . Each of these isotopes is the parent of a chain (decay series) of radioactive daughters, each ending with stable isotopes of lead. Decay of  $^{238}\text{U}$  gives rise to the 'uranium series' which includes  $^{234}\text{U}$  as an intermediate daughter and ends in stable  $^{206}\text{Pb}$ . The decay of  $^{235}\text{U}$  gives rise to the 'actinium series' which ends in stable  $^{207}\text{Pb}$ . The decay of  $^{232}\text{Th}$  results in the emission of six alpha and four beta particles (see [ALPHA DECAY](#); [BETA DECAY](#)) leading to the formation of stable  $^{208}\text{Pb}$ .

**Deccan Traps** A large expanse of *\*flood basalt*, comprising many layers that together are more than 2 km thick, that covers 500 000 km<sup>2</sup> of west-central India. The layers are exposed as step-like formations known as traps, from a Sanskrit word for step, and the layers of *\*sedimentary rock* between the basalt layers are rich in fossils.

**decibel (dB)** One-tenth of a bel (named after Alexander Graham Bell) and the unit in which two power levels are compared. It is used most commonly in acoustics and in describing electrical signals. The decibel difference ( $N$ ) between the largest ( $A_{\text{max}}$ ) and smallest ( $A_{\text{min}}$ ) measurable amplitudes is given by  $N = 20\log_{10}(A_{\text{max}}/A_{\text{min}})$ . The ratio of values for two power levels,  $P_1$  and  $P_2$ , is given by  $N = 10\log_{10}(P_1/P_2)$ .

**declination** **1.** The angle between magnetic North and true (geographic) North. **2.** The angle between a celestial object and the celestial equator.

**declined** See [STIPE](#).

**décollement** Literally, 'unsticking', a concept formulated by geologists studying the structure of the Swiss Jura, and dealing with the sliding and buckling of the *\*Mesozoic* cover over the crystalline *\*basement*. *\*Triassic* *\*evaporites* act as the 'lubricant' of materials in the Jura environment.

**décollement plane** See [DETACHMENT HORIZON](#).

**decompression melting** The melting of *\*mantle* material that occurs when the material rises into a region of lower pressure, allowing it to cross from its *\*solidus* to its *\*liquidus*. Decompression melting takes place in *\*mantle*

plumes and regions where plate movements stretch the crustal rocks, making them thinner.

**deconvolution** Inverse *\*filter*, or the action of undoing the effect of a previous *\*convolution* process. The convolution of the Earth's *\*impulse* response  $R$  (where  $R$  is the set of *\*reflection coefficients* characteristic of particular layered strata) with a wavelet  $W$  results in the observed *\*seismic* trace  $S$ , such that  $S = R*W$ . A spike signal (*\*Dirac function*) ( $\delta$ ) at time zero is the convolution of the wavelet  $W$  with some designed operator  $D$  such that  $D*W = \delta$ . Thus if the operator  $D$  is applied to the seismic trace  $S$ ,  $R$  can be obtained:  $D*S = D*R*W = D*W*R = \delta*R = R$ . (It should be noted that convolution of a spike function  $\delta$  leaves the function unaltered, hence  $\delta*R = R$ ).

**decorrelation stretching** In *\*remote* sensing, a type of *\*contrast stretching* that results in the artificial enhancement in colour of an image. The spread of multispectral data is increased along the natural maximum as identified by *\*principal* component analysis.

**decussate** Applied to a *\*texture* in which crystals have no preferred orientation, but have grown in a random arrangement.

**dedolomite** *\*Limestone* formed by the replacement of *\*dolomite* by *\*calcite* in a *\*carbonate* rock. This replacement occurs mainly when a dolomite is flushed through by *\*meteoric* or sulphate-rich waters. A characteristic feature of dedolomite is the presence of dolomite-shaped, rhombohedral crystal forms now occupied by calcite *\*pseudomorphs*.

**deepening** In meteorology, a lowering of the central pressure in a depression. *Compare* FILLING.

**deep scattering layer (DSL)** Sound-reflecting layer in ocean waters, consisting of a stratified, dense concentration of zooplankton and fish. Such organism-rich layers, which cause scattering of sound as recorded on an echo sounder, may be 50–200 m thick.

**Deep Sea Drilling Programme (DSDP)** An international programme initiated in 1963, which resulted in more than 500 *\*boreholes* being drilled in the sea bed of the *\*Atlantic*, *\*Pacific*, and *\*Indian* Oceans and the Mediterranean Sea. Until 1975, the programme was financed mainly by the National Science Foundation of America, but subsequently it has received

support from the United Kingdom, France, Germany, Japan, and Russia. The managing institution was the Scripps Institute of Oceanography, and the drilling ship used was the *Glomar Challenger*. The programme evolved into the more ambitious [\\*International Programme of Ocean Drilling \(IPOD\)](#) and now the Integrated Ocean Drilling Programme (ODP).

**deep-sea fan** Fan-shaped body of sediment that accumulates at the lower end of a submarine canyon, either at the foot of the [\\*continental slope](#) or on the [\\*continental rise](#).

**deep-sea geomorphology** The study, including mapping, of the shape and surface structures of the ocean floor that result from [\\*tectonic](#) processes, [\\*volcanicity](#), and sedimentation. ‘Deep sea’ is often taken to mean those areas of the ocean floor lying beyond the [\\*continental shelves](#).

**deep-sea trench** Narrow, elongate, steep-sided and often rock-walled depression that is 3000 m or more deeper than the adjacent deep-sea floor. Trenches are the deepest parts of the ocean, the greatest known depth being that of the Challenger Deep in the [\\*Marianas Trench](#) where a depth of 11 022 m has been recorded. Trenches usually mark the position of a destructive [\\*plate margin](#), where the sea floor is slowly being destroyed by [\\*subduction](#).

**Deep Space Climate Observatory (DSCOVR)** A [\\*NASA/NOAA](#) mission, formerly known as Triana (after Rodrigo de Triana, the lookout who first spotted land on Christopher Columbus’s first mission), to observe the sunlit Earth as a continuous full disk from a satellite placed at the first [\\*Lagrangian point \(L1\)](#). The satellite was launched on 11 February 2015, from Cape Canaveral, Florida.

**Deerparkian (Oriskanyan)** See [ULSTERIAN](#).

**Defense Meteorological Satellite Program (DMSP)** A long-term, low-altitude satellite mission, managed by the US Air Force on behalf of the Department of Defense, to provide daily data on worldwide cloud cover, together with oceanographic and solar-geophysical environmental data for use in weather forecasting. The four spacecraft were launched from California on 12 December 1999, 4 November 2006, 18 October 2009, and 3 April 2014.

**deflation** The removal of material from a land surface by *\*aeolian* processes. It is most effective where extensive unconsolidated materials are exposed, e.g. on beaches, and on dry lake and river beds. The very large, enclosed hollows of many *\*deserts* (e.g. the Qattara Depression of the Egyptian Sahara) may be due to deflation.

**deflation hollow** Enclosed depression produced by wind erosion. It may be found both in hot deserts, where wind may scour a hollow in relatively unconsolidated material, and in more temperate regions, where a protective vegetational cover has been removed from a sand *\*dune*.

**deflexed** See *STIPE*.

**deformation lamellae** A feature that may be present in twinned *\*crystals* where rows of atoms are offset in response to directed pressure, giving *\*lamellar* gliding twins. Examples include *\*sphalerite*, *\*calcite*, and *\*pyroxenes*.

**deformation twinning (lattice gliding)** When *\*crystals* are stressed they may be deformed plastically by gliding or sliding along planes between rows of atoms within the crystal structure. This may take the form of ‘translation gliding’, whereby one or more rows of atoms may be displaced laterally along the glide plane, or ‘twin gliding’, where a smaller displacement is taken up by each row within the *\*lattice*. Twins produced in this way are called ‘gliding twins’ or ‘deformation twins’.

**de Geer moraine** A type of *\*moraine* landscape that consists of a series of separate, narrow ridges trending parallel to a former ice front, and which can form annually. The ridges may be up to 300 m apart and up to 15 m high. They consist typically of a *\*till* core, capped by a layer of partly rounded boulders. This landscape may have formed beneath the grounded part of an *\*ice* sheet that extended into a lake or sea. It is named after the Swedish geologist Baron Gerard de Geer (1858–1943).

**degree days** See *DAY DEGREES*.

**degrees of freedom** 1. When a substance is heated its kinetic energy increases. Kinetic energy is made up from the translation and rotation of particles, and the vibration of atoms which constitute the molecules of a substance. A substance may, therefore, absorb heat energy supplied to it in several ways, and is said to possess a number of degrees of freedom. In

general, a molecule consisting of  $N$  atoms will have  $3N$  degrees of freedom; thus for a diatomic molecule there will be six degrees of freedom: three will be translational, two rotational, and one vibrational. In a **\*phase diagram**, describing, for example, a three-phase system (such as ice–water–vapour), pressure and/or temperature can be altered independently, in an area where only one phase exists, without altering the one-phase condition. Along the line separating two areas, if temperature is altered then pressure must alter accordingly, or vice versa, to maintain the two-phase equilibrium. At a point where three phases are in equilibrium, alteration of either temperature or pressure will cause one phase to disappear. The system thus possesses (a) two degrees of freedom in the area; (b) one degree of freedom along the line; and (c) no degrees of freedom at the point. **2.** In statistics, the number of independent variables involved in calculating a statistic. This value is equal to the difference between the total number of data points under consideration, and the number of restrictions. The number of restrictions is equal to the number of parameters which are the same in both observed data set and theoretical data set, e.g. total cumulative values, means.

**dehydration** Removal of water, especially from a chemical compound by heat, sometimes in the presence of a catalyst or dehydrating agent, e.g. sulphuric acid. Also refers to removal of water from crystals, oil, etc., commercially by distillation, chemicals, or heat. In **\*metamorphism**, **\*prograde** metamorphic reactions commonly involve dehydration of hydrous minerals.

**dehydration curve (vapour pressure curve)** The equilibrium curve for dehydration reactions on graphs showing increasing total pressure (vertically) against increasing temperature (horizontally). The slope of the dehydration curve varies with **\*vapour pressure** and depends on the relationship with total pressure. This is important in **\*prograde** metamorphism when many reactions involve dehydration reactions as increasing temperature produces water vapour. **\*Retrograde metamorphism** is more limited, as most of the water has been driven off previously and the breakdown of existing grains, nucleation, and growth are much slower in the absence of water.

**Deimos** **1.** See **MARS**. **2.** An international collaborative **\*microsatellite** mission to provide 600-km-swath, multispectral, high-resolution surface imagery. The Deimos-1 and UK-DMC-2 satellites were launched on 29 July

2009, from Baikonur, Kazakhstan. Deimos-2, a follow-on mission, was launched on 19 June 2014, from the Yasny Cosmodrome, Russia.

**delamination** 1. An increasingly accepted hypothesis according to which the lower part of the continental or oceanic \***lithosphere** becomes sufficiently dense, usually by cooling and mineralogical phase changes, that it separates from the upper lithosphere and descends into the \***mantle**, being replaced by warmer, less dense mantle rock. These physical and chemical changes introduce a new tectonic regime. 2. *See also* A-SUBDUCTION.

**Delaware effect** The distortion of lines of force by a change of resistivity between adjacent layers as \***laterolog** or \***induction sondes** are pulled through a \***borehole**. The change of \***lithology** correlates with the onset of the change of signal and not half-way through the change-over.

**delayed flow** Water that reaches river channels after flowing through subsurface routes and also from \***groundwater**. It contrasts with \***surface runoff**; together, surface runoff (quick flow) and delayed flow make up the total of river flow.


**delay time** A time gap between the shot-instant and the start of recording by a \***seismograph** to avoid long, blank sections on a record. It is used particularly in \***reflection** work over deep water where there is no geologic reason to record while the energy is travelling through the water column, and the recording starts just before the expected \***first** arrival. It is also used in \***time-domain** \***induced** polarization surveying to allow for the dissipation of transient voltages which have no direct relation to the \***over-voltage**.

**Dellingr** A 6-unit \***CubeSat** mission by \***NASA's** Goddard Space Flight Center to study the \***ionosphere**. Dellingr is the Norse god of the dawn. The CubeSat was launched on 14 August 2017, from the Kennedy Space Center, Florida.

**Delmontian** A \***stage** (7.5–2.9 Ma ago) in the Upper \***Neogene period** of the west coast of N. America, underlain by the \***Mohnian**, overlain by the \***Repettian**, and roughly contemporaneous with the upper \***Messinian**, \***Zanclean** (Tabianian), and lower \***Piacenzian** stages.

**delta** A discrete protuberance of \***sediment** formed where a sediment-laden current enters an open body of water, at which point there is a reduction in


the velocity of the current. This results in rapid deposition of the sediment, which forms a body, for example at the mouth of a river where the river discharges into the sea or a lake. There is a characteristic coarsening upwards of sediments. A river provides the sediments to form a delta; but the shape and nature of a delta is controlled by a variety of factors including climate, water discharge, sediment load, rate of subsidence of the sea or lake floor, and the nature of the river-mouth processes (particularly tidal and wave energy). One classification of delta types, based on variations in transport patterns on the delta, subdivides deltas into three classes: (a) river-dominated, e.g. the Mississippi and Po; (b) wave-dominated, e.g. the Rhône and Nile; (c) tide-dominated, e.g. the Ganges and Mekong. *See also* [GILBERT-TYPE DELTA](#).



**Delta**

**delta front** The sloping portion of a *\*delta*, developed off shore from the bar at the mouth and passing at its toe into the *\*pro-delta*. Delta fronts are the site of active, and often rapid, sedimentation, frequently characterized

by *\*growth faulting* and *\*slump* structures generated by over-steepening of the sediment pile.



**Delta front**

***Deltatheridium*** Small, probably carnivorous mammal from the Upper *\*Cretaceous*. It was once thought to be an insectivore, but the possession of certain dental characters indicate that it was a likely ancestor or sister group to the creodonts (*\*Creodonta*) and *\*Carnivora*.

**$\Delta T$  method** In seismic investigations, a method of velocity analysis associated with the normal *\*moveout correction* ( $\Delta T$ ), which enables the *\*root-mean-square* velocity ( $v_{\text{rms}}$ ) to be calculated down to a particular *\*reflector* for which the normal incidence two-way travel time ( $t_0$ ) and  $\Delta T$  are known for a given *\*offset* ( $x$ ) such that  $v_{\text{rms}} = x/(2t_0\Delta T)^{1/2}$ . Once  $v_{\text{rms}}$  is known, the depth ( $z$ ) to the reflector can be calculated using  $z = v_{\text{rms}}t_0/2$ .

***delthyrium*** Subtriangular opening or slit-like notch beneath the apex of the pedicle valve of some *\*brachiopods*, for the passage of the *\*pedicle*.

***deltidial plate*** Calcareous deposit at the side of the *\*delthyrium* which serves to constrict or close the opening.

**DEM (digital elevation map)** See INTERFEROGRAM.

**demagnetization (cleaning, magnetic)** Geologic and archaeological samples are usually partially demagnetized in a series of incremental steps to determine their *\*coercivity* and/or *\*blocking-temperature* spectra. *\*Viscous* remanences are more readily removed than either *\*thermal* or *\*chemical remanence*, so it is often possible to isolate the original (primary)

magnetization acquired when the samples were first formed. This is known as **\*alternating** magnetic field demagnetization or thermal ‘cleaning’. **\*Chemical demagnetization** can also be used on permeable sedimentary samples in which the **\*cement** is usually most readily removed by acid washing, thereby preferentially removing the chemical remanence associated with the cement and isolating the **\*detrital remanence** acquired during deposition. Direct magnetic fields can be applied to reduce the observed remanence to zero, this field corresponding to the effective mean coercivity of the total remanence. New developments include the use of tuned **\*microwaves** to demagnetize magnetic minerals without much heating and chemical change.

**demagnetizer** One of the various instruments used to remove gradually the **\*remanent** magnetization of specimens. There are three main types of instrument, based on **\*alternating** magnetic fields, **\*thermal** treatment in zero magnetic field, **\*chemical** treatment, and the use of **\*microwaves**.

**de Maillet, Benoît** (1656–1738) De Maillet was a French diplomat and traveller, whose posthumous work *Telliamed* (his name spelled backward) was published in 1748. A cosmogony, covering the creation and pre-history of the Earth, it argued that all rocks and all forms of life had their origin in a diminishing ocean. He considered the Earth to be more than 2 billion years old. De Maillet’s belief in mermaids and mermen made his book an object of easy ridicule by later naturalists.

**Demospongea (Demospongiae)** (phylum **\*Porifera**) A class of sponges that first appears in the **\*Cambrian**. The soft tissue of the skeleton is supported by siliceous **\*spicules**, each spicule consisting of either a single ray (monaxon) or four rays (tetraxon) diverging at 60° or 120°. Most **\*fossil** species are represented by their spicules only, but they can usually be referred to modern families.

**Demospongiae** See DEMOSPONGEA.

**demultiplexing** The reordering of data streams to allow a computer to commence the processing of multiplexed (see MULTIPLEXING) data. The data samples are rearranged into trace-sequential format, i.e. all the samples for channel 1 are followed by those for channel 2, ..., followed by those for channel *n*.

**dendritic (arborescent)** **1.** Shaped or marked in a pattern reminiscent of the branching of a tree (from the Greek *dendron*, 'tree'). **2.** Applied to the shape or *\*habit* of a *\*crystal* that is deposited or precipitated in tree-like, slender branches, often along narrow joint planes, e.g. manganese oxide.

**dendritic drainage** A *\*drainage pattern* whose shape resembles the pattern made by the branches of a tree or veins of a leaf, which may develop on homogeneous rock.

**dendrochronology (tree-ring analysis)** **1.** The science of dating by means of tree rings. **2.** All aspects of the study of annual growth layers in wood.

**dendroclimatology** Branch of *\*dendrochronology* dealing with the relationships between the annual growth increment in tree trunks and climate, and especially with the reconstruction of past climates from dated tree-ring series. It is assumed that by studying present tree-ring patterns in relation to climate, older tree-ring chronologies may be used to indicate the climatic conditions experienced before detailed climate and weather records were kept.

**dendrogeomorphology** The use of dated tree-ring series to study landforms and geomorphological processes.

**dendrogram** A diagram that represents relationships among groups of taxa, with the highest taxon at the base of a vertical line from which lower taxa branch at appropriate levels. There are two principal types: (a) the *\*phenogram*, which is based solely on similarities in *\*phenotypes*; and (b) the *\*cladogram*.

**Dendrograptidae** See DENDROIDEA.

**dendrohydrology** The use of dated tree-ring series to study hydrological questions, especially relating to the periodicity of river flow and flooding.

**dendroid** **1.** In corals (see COMPOUND CORALS), applied to a colony formed by the irregular branching of *\*corallites*. The individual corallites are separated from one another but may be connected by tubules. **2.** In graptolites (*\*Graptolithina*), applied to a bushy colony formed by irregular branching of the *\*stipes*.

**Dendroidea** (subphylum Stomochordata, class *\*Graptolithina*) An order of graptolites that existed from the Middle *\*Cambrian* to the Lower

**\*Carboniferous.** Most lived attached to the sea bed and were upright and bushy in appearance. They are many-branched, with numerous small thecae (see **THECA**), in some connected by **\*dissepiments**. The **\*stipes** bear two kinds of thecae, **\*autothecae** and **\*bithecae**, opening from a continuous, closed **\*stolotheca**. There are concentric sheets of tissue in the lower part of the **\*rhabdosome**. The order contains three families: Acanthograptidae, Dendrograptidae, and Ptilograptidae.

**dendroid graptolites** See **DENDROIDEA**.

**Denekamp (Zelzate)** A Middle **\*Devensian** **\*interstadial** from the Netherlands, dating from about 30 000 years BP, when the estimated mean July temperature for the Netherlands was 10 °C. The Denekamp Interstadial, together with the Moorshoofd and **\*Hengelo** Interstadials, is perhaps the equivalent of the **\*Upton** Warren Interstadial of the British Isles.

**Dense-medium separation (heavy-medium separation, HMS)** The simplest of the **\*gravity separation** processes used to concentrate **\*minerals** before their final grinding and liberation. It is also used to separate **\*coal** from the heavier **\*shale**. Heavy media of suitable density are used, in which lighter minerals float and denser minerals sink. In industry, the medium is usually a thick pulp made from a heavy mineral in water. Dense-medium separation is suitable for any **\*ore** which has enough difference in specific gravity between waste and mineral particles to repay the cost of treatment.

**density** Mass per unit volume, expressed in SI units as kilograms per cubic metre ( $\text{kg/m}^3$ ). It is determined directly, or indirectly using **\*gravity** or **\*seismic** velocity measurements. Typical densities in unconsolidated, wet sediments range between 1200 and 2600  $\text{kg/m}^3$ , and 1000 to 2000  $\text{kg/m}^3$  if dry. Consolidated sediments range between 1600 and 3200  $\text{kg/m}^3$ . (See also **NAFE—DRAKE RELATIONSHIPS** for seismic determinations of the density of sediments.) **\*Basic** **\*igneous** rocks range between 2300 and 3170  $\text{kg/m}^3$ , and **\*metamorphic rocks** are mostly in the range 2400–3100  $\text{kg/m}^3$ , with **\*eclogite** being one of the denser of the commoner metamorphic rocks, between 3200 and 3540  $\text{kg/m}^3$ . The density of the upper **\*mantle** increases with depth from about 3330 to 4000  $\text{kg/m}^3$ , below which the lower mantle systematically increases to about 5400  $\text{kg/m}^3$  above the **\*core**. The outer

core density is about 10 000 to 12 000 kg/m<sup>3</sup>; the inner core is more uniform, with a density of about 13 000 kg/m<sup>3</sup>.

**density current** Current that is produced by differences in density. Where a flow of sea water has a greater density than that surrounding it, the more dense water will dive beneath the less dense water. Density of sea water is affected by temperature, *\*salinity*, and the content of suspended sediment. A turbidity current is a gravity-controlled density current, in which the density contrast is due to the high suspended-sediment content. Most deep, and bottom, ocean currents are density currents.

**density–depth profile** Within the Earth, the relationship between density and depth is determined primarily using *\*seismic waves*, as their velocity is dependent on density, complemented by gravitational-field and *\*free oscillation* measurements. The upper *\*mantle* is complicated by the presence of *\*phase* transitions, particularly at depths of around 300 km and 650 km. Within the lower mantle, the density increase with depth is essentially *\*adiabatic*, except in the region immediately overlying the *\*core*. The outer core density (about 10 000 kg/m<sup>3</sup>) is almost twice that of the lower mantle (about 5000 kg/m<sup>3</sup>), reflecting a compositional change. The core density increases with depth, with a small jump from the outer to inner core of 12 000 kg/m<sup>3</sup> to 13 000 kg/m<sup>3</sup>.

**density determination (specific gravity determination)** The determination of the density of a substance. In the case of *\*minerals* several techniques are available. (a) Using a Jolly's spring balance, the mineral is first weighed in air and then immersed in water and reweighed. The density is calculated by comparing the two weights. (b) The Berman balance is a very sensitive torsion balance used for determining the density of small fragments of minerals, based on the same principle as the Jolly balance. (c) A pycnometer is used to determine the density of soils or powdered minerals. It consists of a small bottle fitted with a ground-glass stopper with a capillary opening. The specimen is placed in the pycnometer, weighed, and then the bottle is filled with water and reweighed. (d) Heavy liquids can also be used. Liquids which are relatively dense, such as bromoform (sp. gr. 2.89) and methylene iodide (sp. gr. 3.33), may be mixed with acetone (sp. gr. 0.79) to produce a series of liquids of known density. If a mineral is introduced into one of these liquids and it neither rises nor sinks its density

is the same as that of the liquid. These heavy liquids are often toxic, and great care must be taken when using them. Gloves and face masks should always be worn. *See also* [COUNTERPOISED BEAM BALANCE](#).

**density log** A [\\*well log](#) (record) of the density of the rocks penetrated by a drill hole. Normally it is based on the output from a [\\*gamma–gamma sonde](#), with compensation for the effects of the [\\*mud cake](#). It is particularly useful for the identification of [\\*evaporites](#), and when examined in conjunction with the [\\*neutron-log](#) it can provide an indication of the presence of gas zones.

**denticle** In many fish, a scale composed of [\\*dentine](#), with a pulp cavity, which resembles a tooth. In [\\*Elasmobranchii](#) denticles cover the entire body.

**dentine** [\\*Bone](#)-like substance, lacking cell bodies and consisting mainly of calcium phosphate in a fibrous matrix.

**dentition** In bivalve [\\*Mollusca](#), the articulating tooth-and-socket system in the shell [\\*hinge](#) system. Various types may be recognized, e.g. [\\*desmodont](#), [\\*dysodont](#), [\\*heterodont](#), [\\*isodont](#), [\\*pachyodont](#), [\\*schizodont](#), and [\\*taxodont](#).

**denudation** From the Latin *denudare*, to strip bare, the removal of surface material through the processes of [\\*weathering](#), transportation, and [\\*erosion](#).

**denudation chronology** Branch of land-form studies that deals with the historical development of landscapes by [\\*denudation](#), especially during pre-[\\*Quaternary](#) time. Evidence for developmental stages is provided by studies of erosion surfaces and their mantling deposits, [\\*drainage patterns](#), stream long-profiles, and geologic structures.

**depleted mantle** The residue that remains after a given element has been removed from [\\*peridotite](#) to form a [\\*basalt](#) melt. The [\\*incompatible](#) elements (e.g. Rb, U, and [\\*rare-earth](#) elements) are preferentially partitioned into a [\\*melt](#), and during crustal formation these elements in particular have been removed from the [\\*mantle](#), leaving the mantle depleted in incompatibles. *Compare* [UNDEPLETED MANTLE](#).

**depocentre** The site of maximum deposition within a sedimentary basin, where the thickest development of the sedimentary sequence will be found.

**deposit feeder** An animal that lives on or in the *\*sediment* of the sea floor and ingests mud rich in organic material in order to obtain nutriment.

**deposit gauge** Device for collecting solid and liquid atmospheric pollutants.

**deposition** The formation of ice on a surface directly from water vapour, without passing through a liquid phase. *See* [SUBLIMATION](#).

**depositional remanent magnetization (DRM)** The magnetization acquired by a sediment during deposition. This is predominantly *\*detrital remanent magnetization*, but used more widely, the term covers all magnetizations acquired at such a time. *See also* [POST-DEPOSITIONAL MAGNETIZATION](#).

**depositional sequence** A discrete succession of *\*strata*, deposited more or less continuously, and bounded at top and bottom by either an *\*unconformity* surface or the equivalent, correlative *\*disconformity* or conformity surfaces (*see* [CONFORMABLE](#)). The ancient stratal surfaces can be traced laterally by seismic *\*reflection \*profiling*. A depositional sequence is the basic, operational, *\*stratigraphic unit* in *\*seismic stratigraphy*. Where top and bottom boundaries can be traced to conformity surfaces a depositional sequence becomes of *\*chronostratigraphic* significance, as it represents a specific interval of geologic time during which the particular unit was deposited. *See also* [BASELAP](#); [DOWNLAP](#); [ONLAP](#); [TOPLAP](#).

**depositional sequence model** One of the two schools of *\*sequence stratigraphy*, devised at Exxon Production Research ([EPR](#)), that uses *\*unconformities* and their correlated conformities as the boundaries of genetically related strata. *Compare* [GENETIC STRATIGRAPHIC SEQUENCE MODEL](#).

**depositional system** An assemblage of *\*lithofacies* (*see also* [FACIES](#); [FACIES ASSOCIATION](#)) formed within a particular depositional environment (e.g. *\*fluvial*).

**depositional systems tract** An association of related *\*depositional systems*, e.g. a systems tract comprising fluvial, deltaic (*see* [DELTA](#)), and *\*continental-shelf* systems.

**depression** **1.** Enclosed area of low pressure revealed by the pattern of pressure distribution. Depressions are also described as cyclones, or cyclonic systems, and have a characteristic pattern of wind circulation (anticlockwise around low pressure areas in the northern hemisphere). Mid-latitude depressions are associated with the convergence of polar and tropical *\*air* masses along a frontal zone: this commonly becomes deformed, and each air mass in turn advances over parts near (especially south of) the depression path, bringing first a warm and then a cold front. **2.** The downward convexing of a *\*crest* line in a non-*\*cylindroidal* fold.

**depression angle** In *\*radar* terminology, the angle between the horizontal plane passing through the radar antenna and the line between the antenna and object. *Compare* LOOK ANGLE.

**depth point** A subsurface point on a reflector which contributes to a reflection event on a *\*seismic* trace. *See also* COMMON DEPTH POINT.

**deranged drainage** An original *\*drainage pattern* that has been deranged or disturbed by the intervention of external factors such as tectonic activity and glaciation.

**derived** **1.** As applied to *\*fossils*, a term implying that material has been fossilized elsewhere, eroded from its original site, and then incorporated in another, younger, geological horizon. The terms 'reworked' and 'remanié' are sometimes used as alternatives. **2.** *See* APOMORPH. *See also* CLADISTICS.

**dermal bone** *See* BONE.

**dermal denticle** *See* PLACOID SCALE.

**dermatocranium** *See* CRANIUM.

**Derryan** *See* ATOKAN.

**desalination** A series of processes whereby salt water is rendered potable.

**Desdemona (Uranus X)** One of the lesser satellites of *\*Uranus*, with a diameter of 29 km. It was discovered in 1986.

**desert** A term which has no precise definition. Deserts may be thought of in terms of biomes in which evaporation exceeds the average precipitation. The rate of evaporation varies with temperature, but desert conditions are

likely to develop wherever precipitation is less than 250 mm/yr. Typically, precipitation in deserts is very erratic. Plants and animals are either absent or sparsely distributed, and are adapted to long droughts or to a lack of access to free water.

**desertification** The spread of *\*desert*-like conditions, particularly in arid or semi-arid areas, due to the influence of human activity and climatic change.

**desert pavement** A thin covering of gravel and stones found in many desert areas, left after erosion by wind and water has removed the finer soil materials.

**desert rose** A radiating series of petal-shaped *\*calcite* or *\*gypsum* *\*minerals*, sometimes resembling the form of a rose, developed in the early stages of sand *\*diagenesis* in arid regions, and particularly in *\*sabkhas*.

**desert varnish** Thin, dark surface veneer of iron and manganese oxides which coats exposed rocks, especially in hot deserts. It results from the surface precipitation of minerals released from the rock by *\*chemical weathering*.

**desiccation** Long-term loss of water associated with regional climatic change.

**desiccation cracks (mud-cracks, shrinkage cracks, sun-cracks, syneresis cracks)** The polygonal-shaped cracks developed in mud which has dried out in a terrestrial environment. They are most often preserved when loose sand infills the cracks and then buries the desiccated mud surface.

**Desmarest, Nicolas** (1725–1815) French encyclopaedist and amateur geologist who did field-work and mapping in the Auvergne, where he recognized that *\*basalt* was associated with lava flows. He realized that the hexagonal cracks in basalt were the result of cooling.

**desmodont** Applied to a type of *\*hinge* condition found in certain *\*bivalves* in which the teeth are very small or lacking, and ridges may have replaced them. The ligament may be supported by a *\*chondrophore*.

**Desmoinesian** A *\*stage* (308–306.5 Ma ago) in the *\*Pennsylvanian* of N. America, underlain by the *\*Atokan*, followed by the *\*Missourian*, and

roughly contemporaneous with the Podolskian and Myachkovskian stages of the *\*Moscovian* series.

**Despina (Neptune V)** A satellite of *\*Neptune*, with a diameter of 148 km; visual albedo 0.06.

**desquamation** See WEATHERING.

**destructive margin** The contact between two lithospheric *\*plates* which are moving towards each other and where *\*oceanic* crust is being destroyed by *\*subduction*. Destructive margins, a type of *\*convergent* margin, are marked by shallow- to deep-focus *\*earthquakes* and typically andesitic *\*volcanicity*, and most are also marked by an oceanic *\*trench*. See also PLATE MARGIN.

**destructive wave** Relatively high-energy shallow-water wave that causes degradation of a beach by moving more material seawards than landwards, thus having a net erosional effect on the adjacent beach. It is characterized by high frequency, which implies that the *\*swash* is impeded by the backwash from the preceding wave. The backwash is more effective than the swash in moving material, and waves having this character are usually steep and associated with onshore winds. The erosional effect is also enhanced by the near-vertical plunge on breaking. It is favoured by a relatively steeply sloping offshore zone.

**detachment** Where crustal extension causes *\*simple* shear, a zone of deformation that penetrates the crust to the top of the *\*asthenosphere*, where the material is ductile. A detachment forms where the crust is weakest.

**detachment control** The properties of the materials forming a hillslope that determine the ease with which they will become detached from the underlying rock and move downslope.

**detachment fault** A low-angle *\*normal fault*, formed due to the gravitational instability of an uplifted block, along which there is considerable horizontal displacement.

**detachment horizon (décollement plane)** A surface along which overlying rocks have moved in the course of deformation.

**detrital** Applied to material derived from the mechanical breakdown of rock by the processes of *\*weathering* and *\*erosion*.

**detrital remanent magnetization (DRM)** The magnetization of a sediment acquired as *\*ferromagnetic* detrital grains become aligned by the *\*geomagnetic* field as they fall through the water and become part of the sediment on the bottom of a lake, pond, river system, or the sea. *See also* POST-DEPOSITIONAL MAGNETIZATION.

**detritivore (detritus feeder)** An organism that feeds on *\*detritus*, typically of plant origin but including some small animals and other organisms mixed with it. *Compare* SCAVENGER.

**detritus** 1. Loose material resulting from the *\*weathering*, *\*abrasion*, or *\*erosion* of rock that has been transported away from the place where it was produced. 2. Small fragments of organic material, including faecal matter and its accompanying bacteria, on which *\*deposit* feeders subsist.

**detritus feeder** *See* DETRITIVORE.

**deuteric alteration** Textural and mineralogical changes occurring within an *\*igneous* rock during the final crystallization stages of the molten rock. It usually occurs at well above room temperature, but some changes can be delayed for a few years after solidification. *See also* DEUTERIC REACTION.

**deuteric reaction** The ‘simmering’ of an *\*igneous* rock in its own ‘juices’. After the last drop of *\*silicate \*melt* has crystallized and the rock is, in effect, solid, residual water-rich vapour, concentrated by the crystallization of a large proportion of non-water-bearing minerals, can permeate along crystal boundaries and through crystal fractures and react with the *\*primary minerals*. Reaction of the low-temperature vapour with high-temperature magmatic minerals (*see* MAGMA) is termed ‘deuteric alteration’.

**Devensian** The last glacial stage in Britain, lasting from around 70 000 years BP (possibly earlier) to about 10 000 years BP. It is approximately synchronous with the *\*Wisconsinian* stage in N. America, the Weichselian Glaciation in northern Europe and the Würm Glaciation in the Alps. The Devensian was preceded by the *\*Ipswichian* Interglacial, and followed by the *\*Flandrian* (i.e. present) Interglacial. *See also* BÖLLING-ALLERØD INTERSTADIAL; CHELFORD INTERSTADIAL; DRYAS; LATE-GLACIAL; LOCH

LOMOND STADIAL; UPTON WARREN INTERSTADIAL; WINDERMERE INTERSTADIAL.

**deviatoric stress** A *\*stress* component in a system which consists of unequal *\*principal stresses*. There are three deviatoric stresses, obtained by subtracting the mean (or *\*hydrostatic*) stress ( $\bar{\sigma}$ ) from *each principal stress* (i.e.  $\sigma_1 - \bar{\sigma}$ ,  $\sigma_2 - \bar{\sigma}$ , and  $\sigma_3 - \bar{\sigma}$ ). Deviatoric stresses control the degree of body distortion.

**devitrification** The crystallization of minerals following the melting of a glass; this most commonly occurs in glassy *\*ignimbrites*, where the glass is replaced by *\*microcrystalline \*cristobalite* and *\*alkali* feldspar.

**Devonian** The fourth of the six *\*periods* of the *\*Palaeozoic* era and the first of the Upper Palaeozoic sub-era. It began about 416 Ma ago and ended about 359.2 Ma ago. In Europe there are both marine and continental *\*facies* present, the latter being commonly known as the *\*Old Red Sandstone*. Although originally described from the type area in Devon, the marine Devonian is subdivided stratigraphically into *\*stages* established in the exceptionally fossiliferous deposits of the Ardennes in Belgium. These stages are the Lochkovian (416–411.2 Ma ago), Praghian (411.2–407 Ma ago), and Emsian (407–397.5 Ma ago) of the Lower Devonian, the Eifelian (397.5–391.8 Ma ago) and *\*Givetian* (391.8–385.3 Ma ago) of the Middle Devonian, and the *\*Frasnian* (385.3–374.5 Ma ago) and Famennian (374.5–359.2 Ma ago) of the Upper Devonian. The subdivision of the marine deposits is based on lithologies and the presence of an abundant invertebrate fauna including *\*goniatites* and *\*spiriferid* brachiopods. The continental Old Red Sandstone deposits contain a fauna of jawless fish and plants belonging to the primitive *\*psilophyte* group. As a result of the *\*Caledonian* orogeny of late *\*Silurian* times, much of the British Isles was covered with continental *\*red-bed* facies.



<https://ucmp.berkeley.edu/devonian/devonian.php>

- The Devonian Period.

**dewatering** The removal of *\*groundwater* to reduce flow-rate or diminish pressure. Methods used depend on the *\*permeability* of the ground,

proximity of hydrogeologic boundaries (see [HYDRAULIC BOUNDARY](#)), \*[storage](#) coefficient of the soil, pressure, and \*[hydraulic](#) gradient. Methods used include \*[abstraction](#) by \*[wells](#), \*[electro-osmosis](#), sumps and drains, vertical drains, or exclusion by \*[grouting](#), \*[compressed](#) air, or freezing techniques (see [ARTIFICIAL FREEZING](#)). Dewatering is usually undertaken to improve conditions in surface \*[excavations](#) and to help construction work at or near the surface.

**dew-point** The temperature, with constant pressure and water-vapour content, to which air must be cooled for saturation to occur. Dew-point may be determined from the dew-point hygrometer which indicates initial condensation on a cooled surface.

**dew-point hygrometer** See [DEW-POINT](#); [HYGROMETER](#).

**dextral coiling** See [COILING](#).

**dextral fault (right lateral-fault)** The sense of displacement in \*[strike-slip](#) fault zones where one block is displaced to the right of the block from which the observation is made. See also [FAULT](#).

**D:H ratio** Ratio between deuterium (heavy hydrogen,  $^2\text{H}$ ) and hydrogen ( $^1\text{H}$ ) in natural waters and other fluids, and in water combined in hydrous \*[minerals](#). This ratio yields information about the origin and geologic history of the fluid, and about fluid/rock interactions. See also [ISOTOPE FRACTIONATION](#).

**diabase** See [DOLERITE](#).

**diachronous** Applied to a lithologic unit that differs in age from place to place.

**diad** See [CRYSTAL SYMMETRY](#).

**diadochy** See [IONIC SUBSTITUTION](#).

**diagenesis** All the changes that take place in a sediment at low temperature and pressure after deposition. With increasing temperature and pressure, diagenesis grades into \*[metamorphism](#). Diagenetic processes such as \*[compaction](#), dissolution (see [PRESSURE SOLUTION](#)), \*[cementation](#), replacement, and \*[recrystallization](#) are the means by which an

unconsolidated, loose *\*sediment* is turned into a *\*sedimentary rock*, e.g. *\*sand* into a *\*sandstone*, or *\*peat* into *\*coal*. See also CARBON ISOTOPES; OXYGEN ISOTOPES.

**diagnostic horizon** Soil layer containing a combination of characteristics typical of that kind of soil.

**diallage** A variety of the *\*pyroxenes* *\*diopside* or *\*augite* which has a characteristic lamellar structure due to a parting along {100}.

**diamagnetism** When a magnetic field is applied to *\*electrons* orbiting a *\*nucleus*, the individual electron spins precess and result in a field in the opposite direction to that applied. All atoms and molecules show this form of magnetization, usually with a negative susceptibility of the order of  $10^{-5}$  or less, but it can have *\*paramagnetism* and/or *\*ferromagnetism* superimposed upon it.

**diamict** A general term describing both *\*diamicton* and *\*diamictite*.

**diamictite** A lithified, conglomeratic, *\*siliciclastic* rock which is unsorted, with *\*sand* and/or coarser particles dispersed through a mud *\*matrix*. The term is commonly used today in preference to 'tillite', which has clear genetic connotations.

**diamicton** The unlithified equivalent of a *\*diamictite*.


**diamond** Crystalline form of carbon that is the hardest naturally occurring material (*\*hardness* 10 on *\*Mohs's* scale); sp. gr. 3.5; *\*cubic*; white or colourless, sometimes yellow, green, red, and rarely blue or black; crystals octahedral; cleavage perfect {111}; of *\*igneous* origin and frequently associated with *\*kimberlites*.

**diamond drilling** The drilling of a *\*borehole* using diamond-studded *\*bits*, usually for *\*core recovery* in exploration.

**diaphragm wall (core wall, cut-off trench)** A nearly impervious wall or core at the base of a dam or similar structure which reduces percolation. *\*Bentonite* suspensions are used in support of deep trench excavations formed for the construction of cast-in-place concrete diaphragm walls.

**diaphoresis** See RETROGRADE METAMORPHISM.


**diapir** Upward-directed, dome-like intrusion of a lighter rock mass, e.g. \*salt or \*granite, into a denser cover. The process is termed diapirism.



### Diapir

**diapirism** See DIAPIR.

**diapsid** In \*Reptilia, describes a skull with two temporal openings behind the eye. This type of skull is characteristic of two subclasses of reptiles: the Lepidosauria, represented today by lizards, snakes, and the \*rhynchocephalian *Sphenodon* (tuatara); and the \*Archosauria, which includes the extinct thecodonts, \*pterosaurs, \*dinosaurs, and the living crocodiles. The earliest known diapsids were lepidosaurs (e.g. *Youngina*) from the Upper \*Permian and Lower \*Triassic of S. Africa. Formerly, many authorities grouped all such reptiles in the subclass Diapsida, but the arrangement has been largely abandoned because the more primitive and more advanced forms are not clearly related to one another. \*Birds are considered to be closely related to the archosaurs. Although the skulls of birds have only one temporal opening, this is apparently derived from the fusion of the two diapsid apertures.



## Diapsid

**Diapsida** See DIAPSID.

***Diarthrognathus broomi*** Of the many mammal-like reptiles (see SYNAPSIDA) recorded from the \*Triassic, this species is one of the closest to mammalian ancestry. It is assigned to the Ictidosauria and is characterized by a number of advanced cranial features. The skull of *Diarthrognathus* is only 4–5 cm long and its dentition is not as specialized as that of certain other mammal-like reptiles from the Upper Triassic.

**diaspore** An aluminium hydrate  $\alpha\text{-AlO(OH)}$ , forming a continuous series with its \*polymorph \*boehmite  $\gamma\text{-AlO(OH)}$ ; sp. gr. 3.3–3.5; \*hardness 6.5–7.0; white; \*massive or foliaceous or as \*orthorhombic prismatic crystals; it is a constituent of \*bauxite and \*laterite deposits along with \*emery and \*corundum.

**diastem** A very small break in a \*conformable succession of \*strata, indicated only by a \*bedding plane, and representing a brief interruption in the deposition of \*sediments, with little or no \*erosion having occurred. Diastems may be very localized, with deposition having occurred elsewhere. Compare NON-SEQUENCE.

**diastrophism** The deformation of the Earth's \*crust on a large scale. \*Faulting, \*folding, and \*plate motion can be involved, to produce mountain ranges, \*rift valleys, continents, and ocean floor.

**diatom** See BACILLARIOPHYCEAE.

**diatomaceous earth (kieselguhr)** Deposit composed of *\*fossil* diatoms (see *BACILLARIOPHYCEAE*), which is mined for many industrial uses: as a mild abrasive in metal polishes, as a filtering medium (e.g. in sugar refineries), for insulation of boilers and blast furnaces, etc., and under the name 'kieselguhr' as a vehicle for explosives. Vast deposits of diatomaceous earth (called '*\*diatomite*') are mined at Lompoc, California.

**diatomite** Diatom-rich *\*sediment*, which has been laid down in a lacustrine or deep-sea environment. The diatom cell wall is made from *\*silica*, therefore the sediment is siliceous. See also *BACILLARIOPHYCEAE*; *DIATOMACEOUS EARTH*; *RADIOLARIAN EARTH*.

**diatom ooze** Soft, siliceous, deep-sea deposit composed of more than 30% (by volume) *\*diatom* cell walls (see *BACILLARIOPHYCEAE*). Diatomaceous *\*sediments* predominate in the high latitudes both around the coast of Antarctica and in the N. Pacific, but in the N. Atlantic the diatom content is overwhelmed by the *\*terrigenous* sediment derived from the adjacent continents. See also *RADIOLARIAN OOZE*.

**diatreme** Carrot-shaped volcanic vent that has formed by explosive action: it is often filled with coarse, angular fragments injected by gas *\*fluidization*, e.g. *\*kimberlites*. Diatremes typically cut through non-volcanic *\*basement* rocks, and contain fragments derived from great depths.

**DICE** See *DYNAMIC IONOSPHERE CUBE SAT EXPERIMENT*.

**dichograptid** A family of graptolites (order *\*Graptoloidea*), 'dichograptid' being the name given to a fauna of graptoloids occurring in the early *\*Ordovician*.

**Dichograptina** See *GRAPTOLOIDEA*.

**dichotomize** To split into two equal parts.

**dichroic** See *DICHROISM*.

**dichroism** (*adj.* **dichroic**) The property some *\*minerals* have of absorbing more light in one *\*vibration* direction than in another, and consequently of giving different colours in two different *\*vibration* directions. It is a form of *\*pleochroism* and is expressed by giving the colours for each of the two vibration directions.

**Dickinsonia** See EDIACARAN FOSSILS.

**dickite** See KAOLINITE.

**Dictyonema flabelliforme** One of the best known of all \*graptolite species, *D. flabelliforme* is characteristically conical or bell-shaped in outline, with numerous branches connected by horizontal crossbars (\*dissepiments). *D. flabelliforme* is a dendroid graptolite which was probably \*planktonic in its mode of life. It ranged from the \*Cambrian into the \*Carboniferous.

**diductor muscle** See DIVARICATOR MUSCLE; MUSCLE SCAR.

**dielectric** A material which does not conduct electricity, but in which an applied electric field displaces \*charge rather than causing it to flow.

**dielectric constant (relative permittivity, symbol  $\epsilon_r$ )** Measure of the polarity of a medium. The force ( $F$ ) between two electric charges ( $e$ ) at a distance ( $d$ ) apart in a vacuum is expressed as:  $F = e^2/d^2$ . In any other medium:  $F = e^2/\epsilon_r d^2$  where  $\epsilon_r$  is the dielectric constant. Typical values are: 1.0 for air; 1.013 for steam; 15.5 for liquid ammonia; 80.36 for water at 20°C; usually 3.18 for ice, although at low frequencies it can be up to two orders of magnitude higher; 5–19 for granite; and 3–105 for dry to moist sand. The dielectric constant is temperature dependent, increasing as temperature increases, and strongly frequency dependent with low-frequency (less than 100 Hz) values being up to 30% higher than those of high frequency (more than 100 Hz). It is analogous to the \*magnetic \*permeability. The complex relative permittivity,  $\epsilon^*$  is given by  $\epsilon^* = \epsilon' - j\epsilon''$  where  $\epsilon'$  and  $\epsilon''$  are the real component (relative permittivity) and the imaginary component (\*dielectric loss factor) respectively, and  $j = \sqrt{-1}$ .

**dielectric loss factor ( $\epsilon''$ )** Related to the complex \*relative permittivity,  $\epsilon''$  is a measure of the loss of energy in a \*dielectric material through conduction, slow polarization currents, and other dissipative phenomena. The peak value for a dielectric with no direct-current conductivity occurs at the \*relaxation frequency, which is temperature related. The maximum value can be used as an important measure of the dielectric properties of rocks and ice.

**dielectric permittivity ( $\epsilon_0$ )** The 'absolute' permittivity of a \*dielectric, being the ratio of the electric displacement to the electric field strength at

the same point. Its value for free space is  $8.854185 \times 10^{-12}$  F/m. The real component of the complex *\*relative* permittivity is an important diagnostic parameter when considering the high- and low-frequency values for zero *\*dielectric* loss. It relates to grain size, packing variations, and density of the material. *See also* DIELECTRIC CONSTANT.

**Dienerian** A *\*stage* of the Early *\*Triassic* epoch, 250.4–249.7 Ma ago, preceded by the *\*Griesbachian* and followed by the *\*Olenekian*.

**Dietz, Robert Sinclair** (1914–95) An American naval oceanographer, Dietz worked on *\*magnetic* anomaly patterns in the north-eastern Pacific, and in 1961 he coined the term *\*sea-floor* spreading for his theory that new ocean floor was created at ocean *\*ridges*. He also worked on lunar cratering, and recognized craters on Earth as being of meteoric origin.

**differential settlement** Unequal settling of material; gradual downward movement of *\*foundations* due to compression of soil which can lead to damage if *\*settlement* is uneven.

**differential stress** *See* MOHR STRESS DIAGRAM; STRESS DIFFERENCE.

**differentiation** *See* MAGMATIC DIFFERENTIATION.

**differentiation index** The sum of the normative constituents (*see* CIPW NORM CALCULATION)  $Q + Ab + Or + Ne + Kp + Lc$  in an *\*igneous* rock, where  $Q =$  *\*quartz*,  $Ab =$  *\*albite*,  $Or =$  *\*orthoclase*,  $Ne =$  *\*nepheline*,  $Kp =$  *\*kaliophilite*, and  $Lc =$  *\*leucite*. The index, defined in 1960 by two American petrologists, C. P. Thornton and O. F. Tuttle (1916–83), seeks to quantify the degree of differentiation a rock has undergone. The greater the degree of differentiation, the more enriched the rock is in *\*felsic* minerals and hence the higher the differentiation index.

**diffraction** The radial scattering of any wave (light, radio, seismic, water, etc.) incident upon an abrupt discontinuity in accordance with *\*Huygens' principle*. A *\*fault* plane, *\*angular unconformity*, small isolated objects (e.g. boulders, fragments of wrecked ships, etc.) will all give rise to the diffraction of incident seismic energy. The quasi-hyperbolic curvature of a seismic diffraction event is related to the velocity within the media through which the diffracted wave travels. In media with slow velocities, the hyperbola is strongly curved, the curvature decreasing as velocity increases.

**diffuse reflection** The reflection of \*electromagnetic radiation from a surface in all directions evenly. *Compare* SPECULAR REFLECTION.

**diffusion** 1. Movement of molecules or \*ions from a region of higher to one of lower solute concentration as a result of their random thermal movement. For example, ions diffuse through a solution or \*melt towards growing \*crystals as their incorporation into the solid phase reduces their concentration in the immediately adjacent liquid. 2. (in crystals) (a) Self diffusion involves the movement of a unit of a given composition through a crystal \*lattice of the same composition. (b) Volume diffusion, the movement of atoms or ions through the crystal lattice. It includes simple self diffusion and more complex situations where ions of a certain species migrate through a lattice containing a variety of ions of different sizes or charges and in various configurations. Self diffusion leads only to change in shape or texture, volume diffusion leads to changes in composition.

**diffusion coefficient (*D*)** In diffusion calculations, the square of the dimension distance divided by time; this varies with the type of particles, the diffusion medium, and temperature.

**diffusion-controlled growth** \*Crystal growth in a \*melt or \*hydrothermal fluid where growth rate is controlled by volume \*diffusion in the liquid. This generally occurs in multi-component systems at large degrees of \*supercooling.

**diffusion creep** Deformation as a result of the migration of atoms along a stress gradient. It usually becomes important as the temperature rises close to the melting temperature. *See also* COBLE CREEP; CREEP MECHANISMS; NABARRO—HERRING CREEP.

**digenite** A copper sulphide  $\text{Cu}_9\text{S}_5$ , closely related to \*chalcocite  $\text{Cu}_2\text{S}$ ; sp. gr. 5.7; \*hardness 3; greyish-blue; \*sub-metallic \*lustre; occurs as irregular aggregates in association with other copper sulphides, e.g. \*chalcopyrite and \*bornite, in mineral deposits.

**digital elevation map (DEM)** *See* INTERFEROGRAM.

**digital image** In \*remote sensing, an image in which the continuous variation in the scene being sensed is converted to discrete variation in the

form of a finite range of integer values (**\*pixels**) each assigned a **\*digital number**. *See also* [ANALOGUE IMAGE](#).

**digital number** In **\*remote** sensing systems, a variable assigned to a **\*pixel**, usually in the form of a binary integer in the range of 0–255 (i.e. a **\*byte**). The range of energies examined in a remote sensing system is broken into 256 **\*bins**. A single pixel may have several digital number variables corresponding to different **\*bands** recorded.

**digitize** **1.** To translate graphical information into a series of numbers suitable for processing by (digital) computer. For example, topographic detail can be taken from a map and digitized to produce a computer-generated topographic cross-section. The superimposition of an orthogonal coordinate system on to an image and the recording of the data in a machine-readable form assumes that a two-dimensional image is being analysed. It can be done three- dimensionally if **\*stereophotographs** are analysed. Fully automated systems also exist for this type of work. **2.** To convert analog data into digital form by sampling the continuous record at discrete sample intervals. *See also* [ALIASING](#); [SAMPLING FREQUENCY](#).

**dihedron** *See* [SPHENOID](#).

**dike** *See* [DYKE](#).

**dilatancy** In rocks, the increase in volume during deformation. This is achieved by an increase in pore volume, rotation of grains, grain boundary slippage, and microfracturing.

**dilatation (dilation)** The process in which an applied **\*stress** leads to a state of **\*strain** involving a change in volume. The change in volume is given by:  $\Delta_v = (V_f - V_o)/V_o$ , where  $V_o$  is the initial volume and  $V_f$  the final volume.

**dilatational wave** *See* [P-WAVE](#).

**dilation** *See* [DILATATION](#).

**diluvialism** Theory of the Earth that used the biblical account of the Noachian Deluge to explain major geologic phenomena. Before about 1800 the biblical flood was frequently cited as the cause of fossilization, sedimentation, deposits, stratifications, etc. In the 19th century the theory

was extended by *\*Cuvier*, *\*Buckland*, and others, who believed in a series of deluges, which had caused the extinction of species and the breaks between major formations. The term is also sometimes (incorrectly) used to describe *\*Werner's* theories.

**dimension stone** Building stone quarried and cut as regular blocks according to specified dimensions, e.g. rectangular, columnar, tabular. Common dimension stones are *\*limestone* and *\*granite*; nowadays often cut quite thinly and used as cladding panels on top of existing *\*concrete* structures, etc. *Compare* FREESTONE.

***Dimetrodon angelensis*** Large, specialized, meat-eating *\*reptile* from the Lower *\*Permian* of North America. *Dimetrodon* had a *\*synapsid*-type skull and is grouped amongst the *\*pelycosaurs*. It grew to over 3m in length and is noted for the presence of a large sail on its back. The skull of this carnivore was high and narrow, with the teeth differentiated and well adapted to predatory habits.

**dimorphism 1.** The presence of one or more morphological differences that divide a species into two groups. Many examples come from sexual differences of particular traits, such as body size (males are often larger than females), or plumage (male birds are usually more colourful than females). *See also* SEXUAL DIMORPHISM. **2.** *See also* POLYMORPHISM.

**dim spot** In a seismic reflection event, a reduction in amplitude caused by the presence of *\*hydrocarbons* which lessen the contrast between the *\*acoustic* impedance of the overlying material and that within the reservoir. *Compare* BRIGHT SPOT; FLAT SPOT.

**dimyarian** *See* MUSCLE SCAR.

**Dinantian** The Lower *\*Carboniferous* sub-system in western Europe, overlain by the *\*Silesian* and comprising the *\*Tournaisian* and *\*Viséan* series. It is dated at 359.2–326.4 Ma ago (Int. Commission on Stratigraphy, 2004) and is roughly contemporaneous with the Carboniferous Limestone (Britain), *\*Kinderhookian*, *\*Osagean*, *\*Meramecian*, and lower *\*Chesterian* (N. America).

**dinocyst** *See* DINOPHYCEAE.

**dinoflagellates** *See* DINOPHYCEAE.

**Dinophyceae** A class of Pyrrophyta, comprising \*algae that are unicellular and have two flagella (thread-like structures) of unequal length. Most dinoflagellates belong to this class and the cysts (dinocysts) are useful in \*biostratigraphy. The organisms have two biological stages. (a) In the motile (thecate) stage (see **THECA**) the organism may have either a flexible cell wall or a rigid, armoured one, and it maintains itself in the water by active movement of the flagella. The cell surface bears two furrows, each holding one flagellum. The transverse furrow is called the ‘cingulum’, the longitudinal one the ‘sulcus’. The cingulum divides the cell into an anterior epitheca and a posterior hypotheca. The apex of the epitheca is sometimes extended to form an apical horn. (b) In the cyst stage the organism is dormant. When encystment occurs a two-layered cyst wall (phragma) is formed. Proximate cysts develop in the cyst wall, in contact with the wall from the motile stage. Chorate cysts develop deeper in the motile cell and are linked to the cell by processes. Cavate cysts are those where the two layers of the wall are separated by cavities. The organism leaves the cyst by an opening (archaeopyle) when conditions change. There are three important orders: Gymnodiniales, Peridinales, and Dinophysiales.

**Dinophysiales** See **DINOPHYCEAE**.

**dinosaurs** Literally the name means ‘terrible lizards’, but in fact the dinosaurs were not lizards. They were \*diapsid reptiles whose closest living relatives are the \*crocodilians and \*birds. Dinosaurs first appeared in the Middle \*Jurassic and produced an astonishing array of different types and sizes before becoming extinct at the end of the \*Cretaceous. The two groups of dinosaurs, \*Saurischia and \*Ornithischia, are not usually thought to be more closely related to each other than to other \*archosaurs, so the concept of ‘dinosaur’ is a heterogeneous one.

**diogenites** A type of \*achondrite composed mainly of \*orthopyroxene and \*hypersthene, which probably formed by crystal separation from a parent \*magma.



<https://www.cefns.nau.edu/geology/naml/Meteorite/Diogenite.html>

- Diogenites.


**Dione (Saturn IV)** One of the major satellites of **\*Saturn**, with a radius of 560 km; mass  $10.52 \times 10^{20}$  kg; mean density  $1440 \text{ kg/m}^3$ ; visual albedo 0.7. It was discovered in 1684 by G. D. Cassini.

**diopside** An important **\*pyroxene**  $\text{CaMgSi}_2\text{O}_6$ , and member of the **\*clinopyroxenes**, forming a continuous series with **\*hedenbergite**  $\text{CaFeSi}_2\text{O}_6$ ; this series may be called diopside solid solution series and denoted  $\text{di}_{\text{ss}}$  (or  $\text{Di}_{\text{ss}}$ ); sp. gr. 3.22 (di) to 3.44 (hed); **\*hardness** 5.5–6.0; **\*monoclinic**; normally a pale, dirty green or grey (or, rarely, colourless); **\*vitreous \*lustre**; **\*crystals** usually short, **\*prismatic**, columnar grains, but the bulk are irregular grains; **\*cleavage** prismatic good {110}; a common constituent of magmatic, **\*basic**, and **\*ultrabasic** rocks, e.g. **\*pyroxenites**, **\*peridotites**, **\*gabbros**, and diabases, and also occurs in **\*basalts** and **\*dolerites**; it is common in many **\*metamorphic rocks**, particularly metamorphosed **\*dolomites** and calcareous sediments. Chrome diopside is a bright green, chromium-rich (1–2%  $\text{Cr}_2\text{O}_3$ ) variety commonly found in **\*kimberlite** pipes.

**diopase** A rare mineral, formula  $\text{CuSiO}_2(\text{OH})_2$ ; sp. gr. 3.3; **\*hardness** 5; **\*trigonal**; emerald-green, transparent to translucent; **\*vitreous \*lustre**; crystals usually short, **\*prismatic**, and terminated by rhombohedra, but also occurs **\*massive**; **\*cleavage** perfect rhombohedral; occurs in the oxidized zone of copper-sulphide deposits.

**diorite** **\*Intermediate**, coarse-grained **\*igneous** rock with up to 10% **\*quartz**. **\*Plagioclase** feldspars are of the **\*oligoclase-\*****andesine** varieties; and **\*ferromagnesian** minerals, e.g. **\*pyroxenes** or **\*hornblende**, are present.

**dip** **1.** The angle of inclination of a planar feature measured from a horizontal **\*datum**. The true dip is always measured in a vertical plane perpendicular to the **\*strike** of the plane of either **\*bedding** or **\*cleavage**. The angle of dip measured in any plane not perpendicular to the strike is an apparent dip and will always be less than the true dip. **2. (magnetic)** The **\*inclination** of the Earth's magnetic field from the horizontal. Positive dips are downwards, negative are upwards from the horizontal plane.



## Dip

**dip circle magnetometer** See MAGNETOMETER.

**dip fault** A *fault* which *dips* parallel to the regional dip of the bed it faults.

**dip-isogon method** A method developed by J. G. Ramsay (1969) for distinguishing three fundamental classes of  *folds* based on their relative thicknesses and the curvatures of their surfaces, as indicated by the inclination (dip) of *isogons*.

**diphycercal tail** Possibly the original type of tail fin in fish, in which the body axis divides the fin into equal *dorsal* and *ventral* sections.

**Diplichnites** A *trace fossil* (*repichnia*) made by an animal as it crawled across a surface.

**Diplocraterion** A permanent, U-shaped dwelling *burrow* created by an animal that lived in the *sediment* but was a suspension feeder. The two parallel tubes are circular in section. Some organisms were capable of responding to rapid deposition or erosion. This response is reflected in the presence of concentric laminae on either side, either inside the two arms of

the burrow or below the outer curve (or, in the form *Diplo craterion yoyo*, in both positions).

**diplograptids** A family of graptoloids (order *\*Graptoloidea*) comprising a fauna that spans the period from the *\*Llanvirn* to the lowermost *\*Silurian*.

**Diplopleurozoa** (phylum *\*Cnidaria*) Extinct, primitive class, known only from the Lower *\*Cambrian*, whose members have a bell-shaped body of elliptical outline, showing distinct *\*bilateral* symmetry. There is a median furrow along the main axis of the body, from which arise numerous, flat, simple segments which reach the periphery and are separated by narrow grooves. The margins are scalloped into short lappets, each bearing a simple tentacle.

**Diplopoda** See MYRIAPODA.

**Diploporita** (subphylum Blastozoa) Extinct class of *\*cystoids* which have *\*uniserial* *\*brachioles*, *\*theca* composed of a large number of irregularly arranged plates, and paired pores (diplopores) present on all plates. They are known from the Lower *\*Ordovician* to Middle *\*Devonian*.

**dipmeter log** A down-hole geophysical log designed to measure the *\*dip* and dip direction of dipping surfaces in a *\*borehole*. The logging tool consists of four resistivity logging devices set at 90° to one another, and held against the side of the borehole (see RESISTIVITY LOG; RESISTIVITY METHODS). When wound back to the surface they respond instantly to layers of differing electrical resistivity (e.g. a *\*clay* *\*horizon* or porous sands) associated with the bedding, while dipping beds produce a response with a time delay related to the dip of the horizon. Computer processing of the data yields a *\*tadpole* plot of dips and dip directions in the well. It is claimed that dipmeter data can be used to identify tectonic as well as sedimentary structures. In practice, the data are often equivocal and must be interpreted with great caution.

**dip moveout** See MOVEOUT.

**Dipneusti** See DIPNOI.

**Dipnoi (Dipneusti)** (class *\*Osteichthyes*) Literally, 'double breathing'. Often ranked as a subclass, the group includes the extant lung-fish and their fossil relatives (e.g. the Middle *\*Devonian* *Dipterus* and the *\*Triassic*

*Ceratodus*). Early forms have an elongated body, a well-ossified internal skeleton, \*heterocercal tail, fleshy-lobed fins, and \*cosmoid scales. Teeth are absent, but one of the commonest fossils is the broad fan-shaped tooth plate that served for shearing and crushing small invertebrates. Dipnoans first appear in Lower \*Devonian rocks and were common in freshwater habitats in the late \*Palaeozoic and the \*Triassic; thereafter their fossil remains are very sparse. There are just three surviving genera (*Neoceratodus*, *Protopterus*, and *Lepidosiren*), all tropical.

**dipole field** The field resulting from the presence of two oppositely polarized magnetic poles. The term is usually applied to the \*geomagnetic dipole field or that associated with a \*magnetic anomaly.

**dip pole** A point, usually on the Earth's surface, where the magnetic vector is vertical, i.e. the \*inclination is 90°. A positive (downward) inclination is a positive (north) dip pole, and a negative (upward) inclination is a negative (south) dip pole. Many dip poles exist and these are not the same as the two \*geomagnetic poles.

**dip shooting** A field procedure in seismic-reflection studies where, commonly, split-\*spread and/or cross-spread shooting is used to determine the structural \*dip of a reflector. The asymmetry of the \*normal moveout (NMO) associated with a split-spread shot is diagnostic of a dipping reflector, with the up-dip side having the smaller NMO corrections. The true structural dip can be determined from the dip information provided by a cross-spread shot.

**dip-slip fault** A fault with a relative displacement parallel to the \*dip of the \*fault plane. In practice many dip-slip faults have some component of \*strike-slip in their displacement. *See also* FAULT.

**dip slope** A geomorphological topographic surface which \*dips in the same direction, and often by the same amount, as the \*true dip or \*apparent dip of the underlying strata. Dip slopes are commonly found in \*cuesta and vale topography.

**dipyramid** *See* BIPYRAMID.

**Dirac function ( $\delta(t)$ )** A spike \*impulse which has an amplitude at only one instant in time and one unit (however measured) of energy.

**direct circulation** Circulation system in which lighter air rises, while denser air descends, leading to the conversion of potential energy to kinetic energy. Land and sea breezes are examples of such circulation, as are the \*Hadley and polar cells in the \*general circulation of the atmosphere. See also THREE-CELL MODEL.

**direct current** A unidirectional current provided by a source of electrical energy. It is used in \*induced polarization surveys and in some electrical \*resistivity surveys, although very-low-frequency alternating current sources are being used increasingly to overcome problems of \*electrode and ground polarization.

**directional fabric** The alignment of linear elements, such as crystals, elongate \*xenoliths, or \*bedding–\*cleavage intersections in a rock. In \*igneous rocks, directional \*fabrics can be imposed by the flow of the \*silicate \*melt aligning \*phenocrysts or elongate xenoliths. In \*metamorphic rocks, directional fabrics can result from directional stresses acting on a rock during metamorphic \*recrystallization, resulting in the alignment of elongate metamorphic minerals such as \*hornblende. During \*folding, directional fabrics can result from the intersection of two planar surfaces such as bedding and cleavage to give an ‘intersection lineation’.

**directional filter** In \*remote sensing, a type of \*spatial-frequency filter which enhances edges within an image which run in selected compass directions.

**direct problem** See FORWARD PROBLEM.

**direct wave** A seismic wave which travels through the ground directly from the source to the detectors without being reflected off or refracted by a subsurface layer.

**dirtying-up trend (bell trend)** In the reading from a \*wireline log (see WELL LOGGING) an expression from a \*gamma-ray sonde showing a progressive upward increase in the gamma reading, indicating a change in the clay-mineral content.

**Disaster Monitoring Constellation for International Imaging (DMCII)** An international programme using eight low-cost \*microsatellites that form a network that provide a daily global imaging capability at 30–40 m resolution in 3–4 spectral bands to assist in rapid-response disaster

monitoring and mitigation. The spacecraft were constructed by Surrey Satellite Technology Ltd., UK, and are operated on behalf of the Algerian, Nigerian, Turkish, British, and Chinese governments. The satellites were launched between 2002 and 2011; six completed their missions, leaving two operational in 2018.

**discharge** A measure of the water flow at a particular point, e.g. a river **\*gauging** station, sewage works, or **\*groundwater** abstraction well. Various units of measurement are in common use depending on the nature of the discharge being measured. River flow may be expressed in cubic metres per second (misleadingly called 'cumecs' in some literature), while **\*borehole** flows may be more conveniently expressed as litres per second.

**discharge hydrograph** See **HYDROGRAPH**.

**discoïd** See **SOLITARY CORALS**.

**disconformity** Unconformity such that the beds above and below the surface are parallel. Some disconformity surfaces are highly irregular, whereas others have no obvious relief.

**discontinuity** **1.** A boundary or layer at depth, marked by a significant change in the speed of transmission of **\*seismic waves**, e.g. **\*Mohorovičić discontinuity**. **2.** In meteorology, sharp changes in temperature, wind, or humidity across a frontal boundary. **3.** A break in sedimentation.

**discontinuous reaction series** A sequence of mineral reactions each of which takes place at a specific temperature during the cooling of a **\*magma**. A high-temperature mineral will remain in equilibrium with a cooling magma until it reaches its reaction temperature. At this point the mineral completely reacts with the **\*melt** and dissolves away to produce a new mineral in equilibrium with the melt. This second mineral remains in equilibrium with the magma as the magma continues to cool until its reaction temperature is also reached. The process then repeats itself. The principle of discontinuous reaction was first advanced by Norman **\*Bowen** in 1928. In sub-alkaline magmas the discontinuous reaction **\*olivine-\*pyroxene-\*amphibole-\*biotite** is commonly observed.

**discordant** **1.** Applied to the cross-cutting relationship of an **\*igneous** intrusion, e.g. a **\*dyke**, when it intersects the **\*bedding planes** or **\*foliation** in the host rock. See **LAW OF CROSS-CUTTING** relationships. **2.** Applied to a

lack of parallelism between the bedding planes (or tectonic *\*fabric*) of one rock unit and others adjacent to it. *Compare* [CONCORDANT](#).

**discordant age** The situation in which a series of age determinations of a single rock body do not yield the same value (*compare* [CONCORDANT AGE](#)). Discordant ages indicate that some transfer of materials has taken place into or out of the rock or mineral, and this transfer could have affected either the parent or daughter *\*nuclides*, or both.

**discordant drainage** A *\*drainage pattern* that runs across (i.e. is discordant to) the geologic structure.

**discordant intrusion** *See* [BATHOLITH](#); [BOSS](#); [DYKE](#).

**Discovery Program** A series of missions planned by *\*NASA*, based on small-scale spacecraft that are designed to proceed from development to flight within less than three years for a cost of less than \$150 million.



<https://science.nasa.gov/solar-system/programs/discovery>

- A NASA programme of missions to study asteroids, comets, interstellar dust, Mars, the Moon, and solar wind.

**disharmonic fold** A *\*fold* which exhibits sharp changes in its geometric properties (i.e. wavelength, symmetry, and shape) when compared with other folds in adjacent layers. It is due to the interlayering of competent and incompetent beds (*see* [COMPETENCE](#)) which buckle on different wavelengths.

**dish-pan experiment** Method for simulating the convective cell and other global atmospheric motions. A shallow layer of water in a round vessel is heated at the edge, to represent the equator, and cooled at the centre, to represent the polar region. As the dish-pan is rotated, various motions are produced in response to the radial temperature gradient and to the speed of rotation. At lower speeds, the flow pattern is *\*zonal*, but with faster rotation large waves develop and incorporate closed circulations. *See also* [HADLEY CELL](#); [ROSSBY WAVE](#).

**dish structure** A *\*sedimentary structure* seen in *\*sandstones*, and characterized by repeated horizons comprising a series of concave-upwards,

dish-like *\*laminations*. The dishes are commonly lined with thin (0.2–2.0mm) *\*clay laminae* and are separated by vertical *\*pillar structures*. Both the dish structures and associated pillars are formed by *\*dewatering* of the pore water from unconsolidated sands.

**disjunct** Applied to the distribution of populations that have distinct ranges separated from one another by a distance large enough to ensure DNA cannot be exchanged between them. *Compare CONJUNCT*.

**disjunct shell** See COILING.

**dislocation** 1. The relative displacement to either side of a fracture. 2. Imperfections developed along a line of atoms within a crystal structure, which may displace or add a row of atoms into the regular structure. Edge dislocations add rows of atoms; screw dislocations displace rows of atoms along a plane.

**dislocation creep** See CREEP MECHANISMS.

**dismicrite** A *\*micrite*-dominated *\*limestone* disrupted by an abundance of *\*fenestrae* or *\*birdseye* structures. Dismicrites are characteristic of *\*intertidal* to *\*supratidal* environments. *See also FOLK LIMESTONE CLASSIFICATION*.

**dispersion** 1. The spreading of a body of water as it flows. Lateral dispersion is the widening of the path taken by *\*groundwater* as it flows from a known point of origin through a rock *\*matrix*, due to its movement around individual mineral grains within the main rock body. Unless it flows in well-defined fissures or fractures, water does not travel through a rock in a straight line but is forced to flow across a widening front because of the granular nature of the rock matrix. Longitudinal dispersion is the spreading out of a body of water along its own flow path, due to the differences in water velocities in larger and smaller pores of the rock. Both modes of dispersion are normally observed by means of *\*tracers*. Lateral and longitudinal dispersion also occur in river *\*channels*, where they are due to differences in *\*flow* velocity across the channel and between the water surface and the bed, and also to random fluctuations in velocity caused by turbulent eddies. 2. The process of separating soil particles (as in *\*aggregates*) from each other so that they may react as individual particles. Aggregates or *\*peds* of soil particles are destroyed by dispersion (and their

formation is initiated by **\*flocculation**). **3.** The distortion of a wave train that occurs when velocity varies with frequency and consequently the **\*phase velocity** is not the same as the **\*group velocity**. There is minimal dispersion for most **\*body waves** (although electromagnetic body waves are subject to considerable dispersion) but it is important for **\*surface waves**, especially **\*Love waves**, particularly in the presence of velocity-layering near the surface. **4.** In **\*mineral optics**, a measure of the difference in **\*refractive index**, determined by direct observation of the biaxial **\*interference figure** using the two extreme wavelengths of the visible spectrum (red and violet). Both **\*isogyres** are observed for red and violet tints. **5.** *See also SWELL.*

**displaced terrane** *See TERRANE.*

**displacement** The relative movement on either side of a **\*fault** plane; it may be in any direction parallel to the plane. The finite displacement on a fault plane is defined by a straight line connecting the positions of the initial and final points.

**displacive transformation** *See POLYMORPHIC TRANSFORMATION.*

**di<sub>ss</sub>** *See DIOPSIDE.*

**disseminated deposit** Deposit in which usually fine-grained **\*ore minerals** are scattered throughout the rock. Large, disseminated deposits form important sources of ore, e.g. **\*porphyry** copper deposits.

**dissepiment** **1.** In **\*corals**, one of the small, horizontal, domed plates which form cyst-like enclosures around the edge of a **\*corallite** and which do not extend right across the corallite. **2.** In **\*graptolites**, a strand of chitinous material which connects adjacent branches in a **\*dendroid** colony.

**dissepimentarium** The part of the interior of a **\*corallite** that is occupied by small plates (dissepiments) which are convex towards the axis of the corallite.

**dissipation trail (distrail)** Phenomenon of cloud evaporation caused by exhaust heat in the rear of an aircraft flight path. *See also CONDENSATION TRAIL.*

**dissolved load** The part of a river's total load that is carried in solution. Five *\*ions* normally constitute approximately 90% of the dissolved load: chloride (Cl<sup>-</sup>), sulphate (SO<sub>4</sub><sup>2-</sup>), dissolved bicarbonate (HCO<sub>3</sub><sup>-</sup>), sodium (Na<sup>+</sup>), and calcium (Ca<sup>2+</sup>). Generally the load is at its maximum concentration during low-discharge conditions when *\*groundwater* is the main source of flow.

**dissolved-oxygen level** The concentration of oxygen held in solution in water. Usually it is measured in mg/l (sometimes in ml/l) or expressed as a percentage of the saturation value for a given water temperature. The dissolved-oxygen level is an important first indicator of water quality. In general, oxygen levels decline as pollution increases.

**distal** Applied to a depositional environment sited at the furthest position from the source area, and generally characterized by fine-grained sediments. *Compare* PROXIMAL.

**distributary channel** A natural stream channel that branches from a trunk stream which it may or may not rejoin. It occurs typically on the surface of an *\*alluvial fan* or *\*delta*, where it may be part of a complex, fan-shaped network that distributes the discharge and sediment load of the main channel among many small distributary channels between which a variable assemblage of bays, lakes, *\*tidal flats*, or marshes may exist. The larger distributary channels show *\*crevasse* splays and *\*levées*. *See also* ANABRANCHING CHANNEL.

**distribution coefficient** *See* PARTITION COEFFICIENT.

**disturbance** General term used to denote low-pressure features, e.g. a depression or trough. Disturbances commonly appear as waves in the major air flows in the mid-*\*troposphere*, e.g. the equatorial easterlies, the prevailing westerlies over middle latitudes, and the trade winds.

**disturbed days (D days)** (geomagnetic) The five days in each month when the disturbance of the geomagnetic field is greatest, determined from observatory records of the *\*diurnal variation*.

**Ditomopyge** During the *\*Carboniferous* and *\*Permian*, the number of *\*trilobite* genera fell consistently until the extinction of the group at the end of the *\*Palaeozoic*. *Ditomopyge* was one of the last representatives of the

family Phillipsiidae. It was comparatively small, with an opisthoparian **\*suture** (see **CEPHALIC SUTURE**), well-developed eyes, and a fused **\*pygidium** or tail area. *D. scitula* is known from the Upper **\*Carboniferous** of N. America.

**Dittmar's law** A principle formulated in 1884 by the Scottish chemist William Dittmar (1833–92) during the **\*Challenger** expedition, stating that within a very narrow range the relative proportions of the major chemical constituents of sea water are constant throughout all the oceans.

**diurnal temperature variation** Daily variations in temperature at a particular place, related to the local radiation budget. In mid-latitudes, for example, maximum temperatures usually occur after noon and minimum temperatures in the early morning. The range varies according to location, with high variation in continental areas, and low variation in maritime areas. The diurnal range in equatorial areas exceeds the annual variation in average temperature.

**diurnal variation** (geomagnetic) Daily variations in the **\*geomagnetic** field, determined from hourly measurements made at observatories, of the horizontal component (*H*), **\*declination**, and vertical component (*Z*) of the field. Most of these variations (Sq variations) are attributed to ionospheric currents induced by solar radiation, but some are associated with lunar effects (Lq variations).

**divariant assemblage** Two or more metamorphic minerals in equilibrium over a range of pressure and temperature (see **METAMORPHISM**). Varying the pressure or temperature within defined limits will not initiate reactions within the equilibrium assemblage of minerals. Since two independent variables, pressure and temperature, can be changed or varied without upsetting the equilibrium, the assemblage of minerals is said to be in divariant equilibrium.

**divariant equilibrium** See **DIVARIANT ASSEMBLAGE**.

**divaricator muscle (diductor muscle)** In **\*Brachiopoda**, the muscle that opens the shell. It is attached at one end to the cardinal process (see **CARDINALIA**) and at the other to the floor of the other valve. See also **MUSCLE SCAR**.

**divergence** **1.** In meteorology, a situation in which, over a given time, more air flows out of a given area than flows in. Some subsiding motion can be expected to take place in the air over this region. **2.** A horizontal flow of water in different directions away from a common centre or line. A particular example of divergence in the oceans is seen in areas of **\*upwelling**. Compare **CONVERGENCE**. **3.** Evolutionarily, genetic segregation and differentiation within a **\*taxon** to the extent that distinct derivative taxa result. The divergence may be at the species, genus, family, order, or higher level (see **CLASSIFICATION**). Thus it is possible to refer for instance to the divergence of **\*reptiles** and **\*mammals** from a stem group, to the divergence of mammal orders, and to the divergence of a breeding population into two related species.

**divergent margin** A boundary between two lithospheric **\*plates** that are moving apart.

**diversification** Increase in the **\*diversity** of distinct types in one or more taxonomic categories (i.e. species, genus, etc., see **CLASSIFICATION**). **\*Phanerozoic**, marine invertebrates with well-developed hard parts provide an illustration: their diversity at phylum level remains much the same throughout, whereas at family level there is a peak at the mid-**\*Palaeozoic**, and a trough at the Permo-**\*Triassic** boundary; after this there is a steady increase to a second, higher peak in the **\*Cenozoic**.

**diversity** Most simply, the **\*species** richness of a **\*community** or area, though it provides a more useful measure of community characteristics when it is combined with an assessment of the relative abundance of species present.

**divide** The boundary between separate **\*catchment** areas or drainage basins. It is normally marked topographically by high ground. In British usage, a divide is sometimes called a watershed, but watershed has a different sense in US usage.

**divining (dowsing, water-witching)** The use of hand-held hazel sticks, pendulums, copper wires, etc., to detect the presence of **\*groundwater**. When groundwater is present the hand-held instrument is reputed to move or twitch. There is no conclusive evidence to show that divining can locate

groundwater any better than a random sinking of wells, but it is widely used nevertheless.

**Diwata-1 (PHL-Microsat-1, Philippine Scientific Earth Observation Microsatellite-1)** The first of two Philippines *\*microsatellites* in Sun-synchronous orbits that provide observations to assist weather forecasting, agriculture, fisheries, forest protection, mining, and the protection of cultural and historic sites. The microsatellite was launched on 23 March 2016, from Cape Canaveral, Florida.

**Dix formula** The equation by which the *\*interval velocity* ( $v_{\text{int}}$ ) can be calculated for a zone between two depths on a *\*seismic section*. For two reflectors with reflected-ray travel times  $t_1$  and  $t_2$ , and *\*root-mean-square* velocities  $v_{\text{rms1}}$  and  $v_{\text{rms2}}$  respectively, then:  $v_{\text{int}} = [(t_2 v_{\text{rms2}}^2 - t_1 v_{\text{rms1}}^2)/(t_2 - t_1)]^{1/2}$ . The stacking velocities are normally assumed to be true root-mean-square velocities but there may be occasions when it is necessary to correct them for the effects of *\*dip*.

**D-layer** The lower *\*mantle*, from a depth of about 720 km to the *\*core* boundary at 2886 km depth, in which the seismic velocities increase as a function of adiabatic compression.

**DMCII** See DISASTER MONITORING CONSTELLATION FOR INTERNATIONAL IMAGING.

**DMO** See MOVEOUT.

**DMSP** See DEFENSE METEOROLOGICAL SATELLITE PROGRAM.

**Dnepr-Samarovo** Glacial deposits occurring in European Russia which may be the equivalent of the *\*Riss* Stadial of Alpine terminology. They are Lower *\*Saalian* in age.

**doctor, the** Local term in equatorial regions for the extremely dry W. African *\*harmattan* wind, which brings welcome relief from the high humidity.

**documentation map** A map showing the total documentary evidence relating to a proposed engineering site, e.g. mining rights, owners of property, etc.

**dodecahedron** A *\*crystal* form composed of twelve rhomb-shaped faces, each face intersecting two of the *\*crystallographic* axes and being parallel to the third. It belongs to the *\*cubic* (isometric) system.

**DO event** See DANSGAARD—OESCHGER EVENT.

**Dogger** **1.** An alternative name for the Middle *\*Jurassic* *\*epoch*, which is dated at 175.6–161.2 Ma ago (Int. Commission on Stratigraphy, 2004). It is preceded by the Early Jurassic, followed by the Late Jurassic, and comprises the *\*Aalenian*, *\*Bajocian*, *\*Bathonian*, and *\*Callovian* *\*stages*. Some authors do not recognize the Aalenian as a separate age, including it instead within an enlarged Bajocian age. **2.** The name of the corresponding European *\*series*, which is roughly contemporaneous with the uppermost *\*Herangi* and lower *\*Kawhia* (New Zealand). **3.** A *\*lithostratigraphic* unit in the Aalenian of Yorkshire, England.

**dog-tooth spar** See SCALENOHEDRON.

**Dokuchayev, Vasily Vasilyevich** (1840–1903) A Russian soil scientist, Dokuchayev was Director of the Kharkov Institute of Agriculture and Forestry. He studied the formation of soils, especially the *\*chernozem*, developing a soil classification and making soil maps.

**doldrums** The oceanic equatorial zone, which has low pressure and light, variable winds. The zone moves seasonally north and south of the equator.

**dolerite (diabase, microgabbro)** A dark-coloured, medium-grained *\*igneous* rock which contains *\*plagioclase* feldspar of labradorite composition and *\*pyroxene* of *\*augite* or titanaugite composition as *\*essential* minerals, and *\*magnetite*, titanomagnetite, or *\*ilmenite* as *\*accessory minerals*. Where *\*olivine* also occurs as an additional mineral, the rock is termed an ‘olivine dolerite’. Where *\*quartz* occurs as an additional mineral in the *\*groundmass*, the rock is termed a ‘quartz dolerite’. Dolerites are the medium-grained equivalents of *\*basalts* and, like the basalts, can be divided into alkali and tholeiitic types (see also ALKALI BASALT; THOLEIITE). Dolerites are commonly found in shallow level *\*intrusions* such as *\*dykes*, *\*sills*, or *\*plugs*.

**Dolgellian (Trempealeauan)** A *\*stage* of the Upper *\*Cambrian*, underlain by the *\*Maentwrogian* and dated at 492.5–488.3 Ma ago (Int. Commission

on Stratigraphy, 2004), preceded by the *\*Festiniogian* and followed by the *\*Tremadocian*.

**doline (swallow-hole, sink-hole)** Steep-sided, enclosed depression in a *\*limestone* region. It is normally located at a site of increased *\*joint* density, which focuses drainage passing vertically through the rock. It enlarges by solution (*\*carbonation*) and by collapse, thus allowing so-called solution dolines to be distinguished from collapse dolines. A shaft may lead from its floor to a cave system.

**Dollo, Louis Antoine Marie Joseph** (1857–1931) Born in France, Dollo became a professor at the University of Brussels. He was a palaeontologist whose major interest was in the development and evolution of fossil reptiles. He also worked on other fossil vertebrates and modern *\*marsupials*, especially the structure of their limbs.

**Dollo's law** Evolutionary irreversibility: once regarded as inevitable, but now considered to apply mainly in special cases. The potential for further useful mutation may well be very limited in highly specialized organisms, since only those mutations that will allow the organism to continue in its narrow niche will normally be functionally possible. In such cases there is therefore a self-perpetuating, almost irreversible, evolutionary trend, so much so that it is regarded virtually as a law, 'Dollo's law' (after the palaeontologist *Louis \*Dollo*). The trend results from steady directional selective pressure, or *\*orthoselection* reinforced by specialization, or developmental canalization (see CANALIZING SELECTION).

**dololithite** A *\*dolomite* composed of *\*detrital* dolomite fragments derived from the *\*weathering* and *\*erosion* of pre-existing dolomites.

**Dolomieu, Déodat de Gratet de** (1750–1801) French explorer and mineralogist, who became Professor of Mineralogy at the Jardin du Roi in Paris. He gave his name to the mineral *\*dolomite* and to the Dolomite Alps.

**dolomite 1. (pearlspar)** Widely distributed rock-forming *\*mineral*,  $\text{CaMg}(\text{CO}_3)_2$ ; sp. gr. 2.8–2.9; *\*hardness* 3.5–4.0; *\*trigonal*; usually white or colourless, but can be yellowish and brown; white *\*streak*; *\*vitreous \*lustre*; crystals are usually rhombohedral with curved, composite faces, also occurs *\*massive* and granular; *\*cleavage* perfect rhombohedral {1011}; usually *\*secondary*, having formed by the action of magnesium-

bearing solutions on **\*limestones** (**\*dolomitization**), also occurs as a **\*gangue** mineral in **\*hydrothermal** veins particularly associated with **\*galena** and **\*sphalerite**. It dissolves very slowly in cold, dilute acid, but effervesces very readily when warmed. It is used as a building stone and in the manufacture of bricks for furnaces. **2. (dolostone)** A **\*sedimentary rock** type, usually formed by the **\*dolomitization** of **\*limestones**, and commonly occurring interbedded with them. Most limestones contain some magnesium carbonate and strictly the term 'dolomite' refers to rocks containing 90% or more of the mineral dolomite (see 1). Dolomite that has formed soon after deposition tends to be fine-grained and to have preserved the original **\*sedimentary structures**, whereas **\*recrystallization** in late-diagenetic dolomites produces a coarser-grained rock, a loss of sedimentary structures, and an increase in **\*porosity**. See also ANKERITE.

**dolomitization** The transformation of **\*limestone** to **\*dolomite** (1), by the conversion of  $\text{CaCO}_3$  to  $\text{CaMg}(\text{CO}_3)_2$ . This occurs by the addition of magnesium to the sediment or rock and may take place soon after deposition or at various stages during **\*diagenesis**. Dolomitization is thought to occur in **\*supratidal** areas where sea water is drawn into the sediment and evaporation results in a high ratio of magnesium to calcium in the pore waters. Deeply buried limestone may become dolomitized if a mixture of fresh water and sea water passes through the rock. Dolomitization results in a 13% increase in **\*porosity**, making dolomites important **\*reservoir rocks**.

**domain** The primary taxonomic division of living systems. There are three domains: **\*Archaea**, **\*Bacteria**, and **\*Eukarya**.

**domal uplift** The upwarping of an area in the form of a dome. Domal uplifts are usually about 1 km high, and  $10^2$ – $10^3$  km across, and are characterized by negative **\*gravity anomalies** and by alkaline **\*volcanicity**. Domal uplifts may also have a radiating system of **\*rift valleys**, and many are thought to be precursors of continental rifting, leading to the development of an ocean basin, and thus represent the first stage of the **\*Wilson** cycle.

**dome** **1. \*Anticlinal** structure which plunges in all directions. **2. (volcanic dome, tholoid)** A mound of viscous **\*lava**, usually **\*rhyolite** in composition, which has grown and built up over a vent. The mound of solid

lava is covered by coarse, angular blocks which form by chilling and brecciation of the growing dome's surface. The blocks accumulate around the growing dome to produce a \*scree slope of crumble \*breccia. Domes can grow by repeated injection of \*magma into the dome body (endogenous dome) or by repeated eruption of small volumes of \*magma from the surface of the dome (exogenous dome). **3. (salt dome)** A circular or elongate plug, 1–2 km in diameter but extending downwards for many kilometres, formed by the upward movement of buoyant and less dense evaporitic material (commonly \*halite) into denser overlying rocks. The diapiric movement (see DIAPIR) may be initiated by tectonic thickening. **4.** A special form of \*crystal development characterized by two roof-like faces symmetrical about a plane of symmetry. The faces are repeated once only about an \*axis of symmetry. **5.** See PERICLINE. **6.** See also ICE DOME.

**dome and basin** See INTERFERENCE PATTERN.

**dome–crescent–mushroom** See CRESCENT-AND-MUSHROOM; INTERFERENCE PATTERN.

**domichnia** A category of traces (see TRACE FOSSIL) made by animals in the creation of a permanent dwelling structure. The \*borings of bivalves (\*Bivalvia) such as *Pholas* are included in this category.

**Dominian Reef** See RANDIAN.

**Donau** A period of glaciation which occurred at about the beginning of the \*Pleistocene. It may correspond to the \*Nebraskan stage in N. America. Evidence of it has largely been removed by later glacial episodes.

**Donau/Günz Interglacial** An \*interglacial \*stage of the Alpine areas that may be equivalent to the \*Waalien of northern Europe and the \*Aftonian of N. America.

**Doppler Orbitography and Radiopositioning Integrated by Satellite (DORIS)** A microwave tracking system based on the Doppler principle that uses a global network of ground-based tracking stations and an unlimited number of satellites. It measures the \*Doppler shift in radio signals sent by the ground stations to satellites carrying DORIS receivers to provide precise measurements of the orbit and data applicable to \*geodesy. The service is operated by the French Centre National d'Études Spatiales.



<https://ids-doris.org>

- International DORIS Service.

**Doppler radar** A device that measures the **\*Doppler shift** in a radar beam reflected from water droplets on either side of a rotating **\*mesocyclone** or **\*tornado**. Angular velocity can be calculated from the extent of the red shift on one side and blue shift on the other.

**Doppler shift** The apparent change in frequency of waves whose source is moving towards or away from an observer. It was first described by the Austrian physicist Christian Doppler (1803–53).

**DORIS** See **DOPPLER ORBITOGRAPHY AND RADIOPOSITIONING INTEGRATED BY SATELLITE**.

**Dorogomilovskian** A Russian **\*stage** (305–303.9 Ma ago) in the **\*Pennsylvanian** epoch, preceded by the **\*Chamovnicheskian** and followed by the **\*Klazminskian** (**\*Gzhelian** epoch).

**dorsa** See **DORSUM**.


**dorsal** Towards the upper surface of an organism (in vertebrates the side of the animal closest to the spine); the opposite of **\*ventral**.

**dorsum** **1.** The inner margin of an ammonoid (see **AMMONOIDEA**) shell; the opposite margin to the **\*venter**. **2. (pl. dorsa, wrinkle ridge)** An elongated area of high elevation (i.e. ridge) on the surface of an extraterrestrial body.

**dot chart** A semi-circular, transparent overlay with points proportional to the distance from a central point. It is used to determine potential effects, particularly for assisting in calculating the **\*topographic** correction in gravity surveys. The elevation and density are determined for each point and assumed to be typical for the area immediately surrounding the point.

**Dott classification** A widely used classification of **\*sandstone** types which divides sandstones into arenites (less than 15% of rock is mud **\*matrix**) and wackes (more than 15% but less than 75% of the rock is mud matrix). The arenites are subdivided into quartz arenite (more than 95% **\*quartz** grains), arkosic arenite (more than 25% **\*feldspar** and more feldspar than rock fragments), subarkose arenite (5–25% feldspar grains and more feldspar

than rock fragments), lithic arenite (more than 25% rock fragments and more rock fragments than feldspar), and sublitharenite (5–25% rock fragments and more rock fragments than feldspar). The wackes are divided into quartz wacke (more than 95% quartz grains), lithic wacke (more than 5% rock fragments and more rock fragments than feldspar), and feldspathic (arkosic) wackes (more than 5% feldspar and more feldspar than rock fragments).



**Dott classification**

**double core barrel** See CORE BARREL.

**double couple (Type II earthquake source)** A seismic wave pattern, consisting of four lobes (for \*P-waves) and four lobes (for \*S-waves) of alternate compression and dilation, which is generated by movement along two \*fault planes at right angles to each other. Compare SINGLE COUPLE.

**double planation** In tropical areas, \*weathering below the surface at the basal \*weathering front is one level of planation, whereas on the land surface weathered material can be modified by wind and water processes and this provides a second level of planation. Climate and local conditions determine the relative rates of development of these two levels of planation, and they can contribute to the production of an \*etchplain.

**double refraction** See BIREFRINGENCE.

**double salt** A salt that contains more than one \*cation or \*anion, formed when two different salts crystallize in the same ionic lattice.

**double sulphides** See SULPHIDES.

**double zig-zag** See INTERFERENCE PATTERN.

**doublure** See CEPHALON.

**down-hole hammer drilling** A method of drilling a \*borehole using a pneumatic, percussive hammer drill. The rock is fragmented by repeated impaction presented directly to it. This produces a medium- diameter hole, but drilling speed is slow so that operating costs tend to be high. The method is often used in drilling holes for a pre-split surface, since such holes have to be precisely located. *See also* PERCUSSION BORING.

**downlap** A discordant relationship at the lower boundary of a \*depositional sequence where younger, initially inclined \*strata thin out down-dip across an underlying inclined or horizontal surface. *Compare* ONLAP. *See also* BASELAP; SEISMIC STRATIGRAPHY.

**downthrow** The relative downward \*displacement on one side of a \*fault.

**Downtonian** In British stratigraphy, a \*stage at the base of the \*Devonian roughly contemporaneous with the lower \*Gedinnian stage. Some authors extend it into the uppermost \*Silurian.

**downward continuation** A technique, which must be used with extreme caution, in which measured values of a potential field (usually gravity or magnetic) at one surface are used to determine the values that field would have at a lower surface. However, the computed field is often erratic and unreliable as a result of the noise contained in the original measurements becoming exaggerated with downward continuation. The method may be useful in resolving anomalies which overlap at the surface where measurements are made, provided the depth to which the field is continued is not below that of the causative bodies themselves. If that depth is exceeded, the computed field may deteriorate completely and become meaningless.

**downwelling (sinking)** The downward movement of surface waters caused by the convergence of different water masses in the open ocean, or where surface waters flow towards the coast. An example of the latter is found along the Washington–Oregon coast in winter. The *\*Antarctic convergence zone* is a major downwelling region in the Southern Ocean and is the source of *\*Antarctic intermediate waters*.

**dowsing** See *DIVINING*.

**draa** High (more than 300 m) sand ridge or chain of *\*dunes* in the Sahara, lying about 0.5–5 km from its nearest neighbour and moving at 2–5 cm per year. It is the largest land-form of the *\*erg*, or sand desert. A star-shaped dune, or ‘rhourd’, is developed at the site where two draa chains cross.

**drag** 1. The flexuring of bedding and *\*cleavage* traces along the margins of a *\*fault* plane produced by the *\*displacement* of either side. Cleavage and bedding traces are seen to *\*dip* asymptotically into and out of the plane of faulting. 2. Ductile deformation of features towards and into *\*shear* zones.

**drainage** 1. The passage of water over and through the land surface, ultimately towards the sea. See *DENDRITIC DRAINAGE*; *DERANGED DRAINAGE*; *DISCORDANT DRAINAGE*; *DRAINAGE DENSITY*; *DRAINAGE PATTERN*; *INCONSEQUENT DRAINAGE*; *SUPERIMPOSED DRAINAGE*. 2. Process of removing the gravitational water from soil, using artificial or natural conditions, such that freely moving water can drain, under gravity, through or off soil. See also *MOLE DRAIN*; *TILE DRAIN*.

**drainage basin morphometry** The measurement of the surface form of a drainage basin, and of the arrangement and organization of the associated river network. Properties such as area, shape, gradient, and relief are important elements of form (see *DRAINAGE BASIN SHAPE INDEX*; *DRAINAGE BASIN RELIEF RATIO*), while the stream network is investigated through a study of its components and of the ways in which they are related. See also *DRAINAGE NETWORK ANALYSIS*.

**drainage basin relief ratio** An index ( $Rh$ ) of the relief characteristics of a drainage basin. It is expressed as  $Rh = H/L$ , where  $H$  is the difference in height between the highest and lowest points in the basin and  $L$  is the horizontal distance along the longest dimension of the basin parallel to the

main stream line. The ratio can provide a measure of the rate of sediment loss from a basin, with which it tends to be positively correlated.

**drainage basin shape index** A measure of the shape of a drainage basin, normally expressed as the ratio between two dimensions of the basin being considered. One such measure is the circularity index (or ratio),  $C$ , expressed as  $C = A_b/A_c$ , where  $A_b$  is the area of the basin and  $A_c$  is the area of a circle with the same length of perimeter as the basin. Another index is the form factor,  $F$ , expressed as  $F = A/L^2$ , where  $A$  is the area of the basin and  $L$  is its length. Such indices may help in forecasting the flood potential of a basin.

**drainage density** A measure of the average spacing between the streams draining an area. It is obtained by dividing the total length of streams by drainage area. Its magnitude is affected by factors such as the amount of rainfall, permeability of the ground surface, and age. *See also* [DRAINAGE](#).

**drainage morphometry (morphometric analysis, Horton analysis)** Calculation of a range of dimensionless drainage-network relationships (*see* [BIFURCATION RATIO](#) for example), based on a system of stream ordering, i.e. the numerical ranking of channel segments within a channel network. Proposed by R. E. Horton in 1945.

**drainage network** *See* [DRAINAGE PATTERN](#).

**drainage-network analysis** The study of the way in which the pattern of streams in a drainage basin is organized. Classical work focused on the relations between the components of a network, e.g. on the link between the importance (or 'order') of a stream segment and its frequency. Certain 'laws' of drainage network composition were derived. The modern approach emphasizes the importance of random processes in an explanation of these 'laws' and is more concerned with the density of the drainage network (drainage density = total length of channels/drainage area).

**drainage pattern** The spatial relationship between individual stream courses in an area. The resulting pattern often reflects the underlying rock type and structure, and several varieties are recognized. A *\*dendritic* pattern is the most common, characterized by a randomly branched arrangement. It is not structurally controlled, and is developed on a homogeneous rock, e.g. *\*clay*. A trellis pattern consists of subparallel streams, usually aligned along

the geologic *\*strike*, and joined at right angles by tributaries. A rectangular pattern is dominated by right-angled bends, and reflects control by *\*joints* or *\*faults*. A centripetal pattern consists of stream courses converging into a central depression. The drainage network is the drainage pattern viewed geometrically.

**drainage-sediment survey** See STREAM-SEDIMENT ANALYSIS.

**drainage wind** See KATABATIC WIND.

**drained test** See TRIAXIAL COMPRESSION TEST.

**dravite** See TOURMALINE.

**drawdown** The lowering of the *\*water-table* or *\*potentiometric* surface, normally as a result of the deliberate extraction of *\*groundwater*.

**dreikanter** See VENTIFACT.

**Dresbachian** See MAENTWROGIAN.

**dribble cone** See SPATTER CONE.

**drift** **1.** Any sediment laid down by, or in association with, the activity of glacial ice. The term is often widened to include related submarine and lacustrine deposits. The British Geological Survey has used it to refer to all superficial (i.e. drift) deposits. It was introduced by C. *\*Lyell* (1797–1875), who suggested that glacial deposits were laid down by melting icebergs which drifted across an ice-age sea covering Britain. This old term is now largely superseded by more recent classifications. **2. (instrumental)** The change in the output of a recording device due to internal factors. Systematic drift can be compensated for by repeat readings at a *\*base station*. **3.** In mining, an underground horizontal passage.

**drifter** See PERCUSSION BORING.

**drift map** A geologic or geomorphological map which shows the distribution of more recent glacial, *\*fluvial*, fluvio-glacial, *\*alluvial*, and marine *\*sediments*. Depending on the distribution and extent of *\*drift* the map may show a combination of solid and drift exposures.

**drill hole** See BOREHOLE.

**drilling** The process of making a hole, usually into the Earth. Two common methods are *\*cable drilling* and *\*rotary drilling*. *See also* BIT; CALYX DRILLING; DIAMOND DRILLING; DOWN-HOLE HAMMER DRILLING; PERCUSSION BORING.

**drilling bit** *See* BIT.

**drilling mud** *See* MUD, DRILLING.

**drill string** *See* STRING.

**dripstone (flowstone, speleothem)** Calcium carbonate rock deposited in caves by the precipitation of *\*calcite* from water as excess dissolved carbon dioxide is diffused into the atmosphere. Dripstone takes various forms, including *\*stalactites*, helictites (having spiral form), curtains, ribbons, and *\*stalagmites*.

**drizzle** Precipitation of very small (<0.5 mm) water droplets generated by *\*coalescence* at the base of cloud such as *\*stratus*. *See also* CRACHIN.

**DRM** 1. *See* DEPOSITIONAL REMANENT MAGNETIZATION. 2. *See also* DETRITAL REMANENT MAGNETIZATION.

**dromaeosaurid** Member of a *\*coelurosaurid* branch of the *\*theropod* *\*dinosaurs*, which developed from the genus *Dromaeosaurus* (the ‘emu reptile’), known from the early Upper *\*Cretaceous* of Canada. Dromaeosaurids had relatively large brain cases and may have been among the most intelligent reptiles ever to have lived.

**drop ball** A simple and economical method for breaking up large stones after quarry blasting by dropping a heavy weight from a crane on to the boulder.

**dropstone** 1. A *\*clast*, released by melting from the base of a floating *\*ice* sheet or *\*glacier*, which falls through the water body to settle in muddy sediment. Dropstones can often be recognized as isolated, coarse clasts set in fine, laminated sediments. 2. A volcaniclastic bomb (*see* VOLCANIC BOMB), ejected from a *\*volcano*, which falls into water and settles in muddy sediment on the lake or sea floor.

**drought** A relative term denoting a period during which rainfall is either totally absent or substantially lower than usual for the area in question, so that there is a resulting shortage of water for human use, agriculture, or natural vegetation and fauna.

**drought cycle** Temporary and repetitive phase of drier conditions in an otherwise favourable environment, e.g. the 22-year drought cycles of N. American grasslands.

**drumlin** Smooth, streamlined, oval-shaped land-form, one end of which is blunt and the other tapered. Drumlins may occur singly, but more commonly they are found within a large group, called a 'drumlin field' or 'drumlin swarm'. Usually they are composed of **\*boulder clay**, but occasionally they are made of solid rock (hence 'rock drumlin'). They are believed to be formed beneath the outer zone of an expanding **\*ice** sheet, during a major advance: they result from the selective deposition of material that is then streamlined by the moving ice. The long axis of a drumlin lies parallel to the direction of the advance.

**drumlin field** See DRUMLIN.

**drumlin swarm** See DRUMLIN.

**druse** (*adj.* **drusy**) A cavity (**\*vugh**) in an **\*igneous** rock or a **\*mineral** vein into which **\*euhedral** (well-formed) **\*crystals** of the rock or mineral vein project, or the crystals themselves. The cavity represents a volume of late-stage, vapour-rich **\*magma** trapped by the rock crystallizing around it. Crystals can grow freely into this medium and hence crystallize in perfect forms (e.g. smokey quartz in granite) to give well-shaped **\*crystal** faces. The word is German, *Druse* meaning decayed or weathered ore.

**drusy** See DRUSE.

**dry adiabatic lapse rate** The rate at which dry (i.e. unsaturated) air cools **\*adiabatically** when rising through the atmosphere as a result of the utilization of energy in expansion. It is 9.8 °C/km. See also **CONDITONAL INSTABILITY**; **ENVIRONMENTAL LAPSE RATE**; **INSTABILITY**; **SATURATED ADIABATIC LAPSE RATE**; **STABILITY**.

**dry air** Air with low **\*relative humidity**; i.e. It is unsaturated air.

**Dryas** Part of the characteristic *\*late-glacial* sequence of climatic change and associated deposits following the last (*\*Devensian*) ice advance and prior to the onset of the markedly warmer conditions of the current (*\*Flandrian*) *\*Interglacial*. The type sequence was first described for Allerød in Denmark, and shows upper- and lower-clay deposits rich in remains of *Dryas octopetala* (mountain avens), and between them deposits of lake mud with remains of cool-temperate flora, e.g. tree birches. The colder Dryas phases mark times of cold, tundra-like conditions throughout what is now temperate Europe and climatic effects that were felt more widely (e.g. extreme aridity to the east of the Mediterranean). The three-fold Older Dryas–Allerød–Younger Dryas sequence forms *\*Pollen Zones* I, II, and III of the widely accepted late and post-glacial chronology of Europe. The basal, Older, Dryas deposit forms Zone I (approximately 15 500–14 700 years ago); the Allerød Zone II; and the Younger Dryas Zone III (12 900–11 600 years ago). The sequence of cool and warm episodes constitutes the *\*Bølling-Allerød* *\*interstadial*.

**dry-bulb thermometer** Thermometer that registers normal air temperature. It may be used in conjunction with a *\*wet-bulb thermometer* in a *\*psychrometer*. See also *HYGROMETER*.

**dry continental morphoclimatic zone** A *\*morphoclimatic* region that produces moderate to low rates of *\*weathering*. *\*Mechanical* weathering, especially due to *\*frost*, occurs in winter and *\*fluvial processes* are active during the wet season. Wind action is moderate locally. Moderate *\*mass-wasting* occurs intermittently.

**dry ice** Solid carbon dioxide, which sublimates at 194.65 K. It is used in *\*cloud seeding* to cool air in supercooled clouds by *\*sublimation* at low temperatures. This can generate many ice crystals for further ice nucleation.

**dry melt** See *MELT*.

**dry season** Period each year during which there is little precipitation. In tropical climates, e.g. over much of India, the dry period is often in the winter season. In very low latitudes, two dry seasons may occur each year, between the northward and southward passage of the equatorial rains. In subtropical, Mediterranean, and west-coast climates, the dry season is in the summer.

**dry valley** Linear depression that lacks a permanent stream but that shows signs of past water *\*erosion*. It is a common land-form in areas underlain by permeable rock, e.g. the Chalk of southern England. The dry valley was eroded during an episode of surface drainage, perhaps due to *\*permafrost* conditions, to greater precipitation, or to a higher *\*water* table.

**dry-weather flow** See BASEFLOW.

**DSCOVR** See DEEP SPACE CLIMATE OBSERVATORY.

**DSDP** See DEEP SEA DRILLING PROGRAMME.

**DSL** See DEEP SCATTERING LAYER.

**d-spacing** See COVALENT RADIUS.

**dual decay** See BRANCHING DECAY.

**DubaiSat** A *\*minisatellite project* of the Mohammed Bin Rashid Space Center, United Arab Emirates, that provides high-resolution panchromatic and multispectral optical images from a spacecraft in a Sun-synchronous low Earth orbit. DubaiSat-1 was launched on 29 July 2009, from Baikonur, Kazakhstan, and completed its mission in March 2017. DubaiSat-2 was launched on 21 November 2013, from Yasný Cosmodrome, Russia.

**ductile behaviour** The response to *\*stress* of certain materials which undergo permanent deformation without fracturing. This produces permanent strain marked by smooth variations within the deformed rock. Ductile behaviour is enhanced where high *\*confining* pressures are combined with high temperatures and low rates of strain, conditions characteristic of deeper crustal levels.

**dug well** See BOREHOLE.

**dump structure** A conical mound of sediment formed from a large amount of debris released when an iceberg overturns or breaks up.

**dune** A land-form produced by the action of wind on unconsolidated sediment, normally sand. Aeolian dune forms range from small *\*ripples* less than 1 cm in height, to the *\*draa* forms of the Sahara which rise to more than 300m. Such dunes may be divided into three basic categories: *\*barchans*; longitudinal or 'seif' dunes, which parallel the wind direction;

and transverse dunes which are aligned normally to the dominant wind. Transverse dunes are initial forms on sandy coastlines in temperate regions. They migrate inland and may be eroded locally by the wind to form a damp hollow or ‘\*dune slack’. The enclosing crescentic dune is a ‘parabolic’ dune whose form reverses that of the barchan. *See also* [AKLÉ DUNE](#); [COPPICE DUNE](#); [DUNE BEDFORM](#); [STAR DUNE](#).

**dune bedform** (‘megaripples’) Mounds or ridges of \*sand which are asymmetrical, and are produced subaqueously by flowing water. The external morphology is similar to the smaller ‘\*ripple’ and larger ‘\*sand wave’, with a gently sloping, upstream side (stoss), and a steeper downstream side (lee). The crestline elongation extends transverse to the flow direction and is sinuous or lunate in plan. The height varies between 0.1 m and 2 m, while the wavelength (spacing) between dunes is 1–10 m. Size and growth are limited by water depth and, in general, dune height is less than one-sixth of the flow depth. The down-current migration of dune bedforms leads to the formation of cross-bedding in sediments.

**dune slack** Flat-bottomed, hollow zone within a sand-dune system that has developed over impervious strata. The slack may result from erosion or \*blow-out of the \*dune system, and the flat base level is therefore close to or at the permanent \*water-table level. Characteristically, dune slacks have rich, marshy flora, with *Salix* species (willows) as typical woody colonizers.

**Dunham classification** A widely used \*limestone classification, proposed by Robert Dunham in 1962, which divides limestones on the basis of their texture and mud content. For limestones which retain their original, depositional texture, the main subdivisions are: lime mudstone (limestone with less than 10% grains in a mud-supported sediment); lime wackestone (limestone with more than 10% grains in a mud-supported sediment); lime packstone (grain-supported limestone with mud \*matrix between the grains); lime grainstone (grain-supported limestone with no mud matrix); and lime boundstone (limestone whose original components were bound together (e.g. by corals or algae) during deposition). For limestones in which the depositional texture has been destroyed by \*recrystallization, Dunham defines two types: crystalline limestone (recrystallized limestone with a fine texture); and sucrosic limestone (recrystallized limestone with a coarse texture). The original Dunham classification does not subdivide

limestones with particles coarser than 2 mm, or differentiate between different types of organically bound limestone. These categories of limestone are defined by Embry and Clovan in their modifications to the Dunham classification. *See also* [EMBRY AND CLOVAN CLASSIFICATION](#).

**dunite** Coarse-grained, [\\*igneous](#) rock, consisting mainly of [\\*olivine](#). It was first described from the Dun Mountain Range, New Zealand.

**Dunroonian** A [\\*stage](#) (28–27 Ma ago) in the [\\*Oligocene](#) of New Zealand, underlain by the [\\*Whaingaroan](#), overlain by the [\\*Waitakian](#), and roughly contemporaneous with the mid [\\*Chattian](#) stage.

**duplex** A series of [\\*horses](#) bounded by a [\\*roof thrust](#) and a [\\*floor thrust](#). There are three main types of compressional duplexes: hinterland dipping; foreland dipping; and antiformal stacks. There are also extensional duplexes in normal [\\*dip-slip](#) regimes. Duplex terminology has also been applied to [\\*strike-slip](#) terrains where smaller [\\*en échelon](#) features bound by two continuous, major [\\*fault](#) zones form a strike-slip complex. In strike-slip regimes extensional and compressional duplexes may coexist.

**duplicatus** From the Latin *duplicatus* meaning ‘doubled’, a variety of cloud with overlapping layers at different heights, typified by such genera as [\\*stratocumulus](#), [\\*altocumulus](#), [\\*altostratus](#), [\\*cirrostratus](#), and [\\*cirrus](#). *See also* [CLOUD CLASSIFICATION](#).

**durain** *See* [COAL LITHOTYPE](#).

**duric horizon** In the [\\*World Reference Base for Soil Resources](#) classification, a [\\*soil horizon](#) consisting of cemented [\\*silica](#).

**duricrust** Deposit of the [\\*weathering zone](#), especially in subtropical environments, which may ultimately develop into a hardened mass. A range of types occurs, each distinguished by a dominant mineral: ferricrete and alcrete are dominated by [\\*sesquioxides](#) of iron and aluminium respectively; silcrete by silica; and [\\*calcrete](#) (caliche) by calcium carbonate.

**duripan** Mineral diagnostic [\\*soil horizon](#) which is [\\*cemented](#) by silica and so will not slake or fall apart in water or hydrochloric acid. It may contain secondary cement, e.g. carbonates and iron oxide. Where duripans are exposed on the soil surface, they are called [\\*duricrust](#). *Compare* [CALICHE](#).

**durisols** A reference soil group in the *\*World Reference Base for Soil Resources* classification scheme. Durisols have a *\*duric* horizon within 100 cm of the surface.

**durophagic** Adapted to the eating of hard materials, such as the diet of many benthic dwellers, comprising shelled invertebrates. Skates and rays, for instance, have a tough dentition and protrusible mouth capable of powerful suction which may be used to dislodge shellfish from rock faces.

**dust** Solid particles, the size of clay and silt particles (see *PARTICLE SIZE*), that can be raised and carried by the wind.

**dust-bowl** An area of the Great Plains region, USA, where a combination of drought and inappropriate farming practices, especially an expansion of wheat production, led to severe *\*deflation* and soil erosion during the middle 1930s. More generally, any region where deflation of cultivable land occurs.

**dust detection instrument** An instrument carried by the *\*Galileo* Orbiter that was designed to measure the movement of the dust stream in the vicinity of *\*Jupiter*. The dust detector was able to measure the distribution, spatial extent, and orbital trajectories of the submicrometre-sized dust particles and to determine if there is indeed a dust wedge, hypothesized to extend for about 700 000km about 10° above and below the jovian equator and to result from volcanic activity on *\*Io*. The instrument also measured the electrical charge on larger particles entering it.

**dust devil (dust whirl, sand pillar)** Very localized *\*whirlwind* in a desert area, where strong convection uplifts dust and sand often to a height of a few tens of metres.

**dust storm** A wind carrying sufficient dust for visibility to be reduced to less than 1 km.

**Dutch cone** Type of *\*cone penetrometer* which may be used to give a continuous log of layered sequence in a *\*soil profile*.

**Du Toit, James Alexander Logie** (1878–1948) Du Toit was a South African geologist who made field studies of the provinces of *\*Gondwana*, and found extensive evidence for *\*continental drift*. He published his ideas in *Our Wandering Continents: An Hypothesis of Continental Drift* (1937).

**Dutton, Clarence Edward** (1841–1912) An officer in the US Army, Dutton was seconded to the US Geological Survey. He introduced the term ‘\*isostasy’, a concept for which he saw evidence in the Colorado plateau. Interested also in \*volcanology and \*seismology, he investigated the Charleston (USA) earthquake of 1886, and in 1904 published a textbook on \*seismology.

**dwarf planet** A body in orbit about a star that is neither an \*asteroid nor a true planet. It is sufficiently massive for its own gravity to compress it into a \*spheroid shape but not to accumulate and so clear other material along its orbital path.

**dyke (dike)** \*Discordant, or cross-cutting, tabular intrusion. Most dykes are vertical or near vertical, having pushed their way through the overlying \*country rock. *See also* DYKE SET; DYKE SWARM; RADIAL DYKE.

**dyke set** A suite of \*dykes whose alignment is parallel or subparallel, reflecting their emplacement from a common source and under a common \*stress regime.

**dyke swarm** A collection of many subvertical \*radial dykes around a central \*intrusion; or many parallel to subparallel \*dykes occurring over a large regional area (\*dyke set).

**dynamic correction** *See* STATIC CORRECTION.

**dynamic correlation** A \*cross-correlation process which involves traces of different offsets, and the adding together of the cross-correlations for similar pairs of traces over a number of adjacent \*depth points. The cross-correlations for successive differences of offset are squared, displayed, and alignments selected to calculate the residual \*normal moveout and \*stacking velocity for each alignment. It is a GSI (Geophysical Service Inc.) process.

**dynamic equilibrium** *See* EQUILIBRIUM.

**Dynamic Ionosphere CubeSat Experiment (DICE)** A US mission to investigate the \*ionosphere involving twin identical \*microsatellites (‘Yahtzee’ and ‘Farkle’), each measuring 10 × 10 × 15 cm and weighing 2 kg, that were launched from California in October 2011.



<https://www.astraspace.net/news/astra-dice-cubesat-update-successfully-captures-first-ever-sed-observations-by-a-cubesat/>

- ASTRA mission update.

**dynamic metamorphism** Fragmentation and *\*recrystallization* of rocks in narrow zones such as *\*faults* or *\*thrusts* where strong deformation has occurred. Rocks are ground to a fine powder in the zone of deformation and, because of their fine *\*grain* size, recrystallize efficiently under the extreme directional stress and release of frictional heat during deformation. The fine-ground powder recrystallizes to a flinty rock which often surrounds fragments of uncrushed *\*country* rock in the deformation zone to form a *\*mylonite*. During extreme deformation all the fragments in the deformation zone are ground down to powder and recrystallize to form a fine-grained, banded rock known as an ‘ultramylonite’, and rocks can melt to form *\*pseudotachylite*.

**dynamic viscosity** In a flowing fluid, a measure of the resistance of the fluid to changing its shape, defined as the ratio of the sheer stress to the rate of deformation sustained across the fluid.

**dysaerobic (poikiloaerobic)** Applied to a depositional environment with 0.1–1.0 ml of dissolved oxygen per litre of water. *Compare* **AEROBIC** and **ANAEROBIC**.

**dysodont** Applied to a type of hinge *\*dentition*, found in certain *\*bivalves*, where teeth are simple, small, and situated very close to the *\*dorsal* margins of the *\*valves*.

**E1P-2** See HRBE CUBESAT MISSION.

**Early Cambrian epoch (Caerfai epoch, Georgian epoch, (US usage) Waucoban epoch)** The earliest *\*epoch* of the *\*Cambrian \*period* in the *\*chronostratigraphic scale*, lasting approximately from 542 to 513 Ma ago and comprising the *\*Tommotian*, *\*Atdabanian*, and *\*Lenian \*ages*. This epoch has also been termed the Comley epoch by some authors and the rocks deposited during this time the Comley *\*series*, underlain by the *\*Ediacaran (\*Precambrian)* and overlain by the *\*St David's*.

**EARS** See EUMETSAT ADVANCED RETRANSMISSION SERVICE.


**Earth** The third planet in the *\*solar system*, outwards from the *\*Sun*. The mean distance of the Earth from the Sun is  $149.6 \times 10^6$  km. This distance provides the standard 'astronomical unit' (*AU*) of measurement. The Earth has a mean radius of 6371 km, density of  $5517 \text{ kg/m}^3$ , and a mass of  $5.99 \times 10^{27}$  g. The *\*oceanic* (5–7 km thick) and *\*continental* (40 km thick) crusts are separated by the *\*Mohorovičić discontinuity* from the silicate *\*mantle*, which extends to the *\*Gutenberg* discontinuity at 2900 km depth, and overlies a molten, iron-rich *\*core*. The oldest rocks are about 3980 million years old, and the Earth formed about 4600 million years ago.

**EarthCARE** See EARTH CLOUDS, AEROSOLS AND RADIATION EXPLORER.

**Earth Clouds, Aerosols and Radiation Explorer (EarthCARE)** A joint European and Japanese mission to observe the vertical profiles of aerosols, and liquid water and ice, and to retrieve profiles of atmospheric radiative heating and cooling. The *\*European Space Agency (ESA)* provides the spacecraft, three instruments, and the launch; the *\*Japan Aerospace Exploration Agency (JAXA)* and the National Institute of Information and

Communications Technology (NICT) provide one instrument. ESA and JAXA provide the ground segment. The mission is planned for launch in August 2019.

**Earth compositional layering (chemical layering)** The solid \*Earth consists of distinct layers, each with a characteristic chemical composition. The most abundant \*crustal elements, in order of abundance, are oxygen, silicon, aluminium, sodium, magnesium, calcium, and iron. The \*mantle is rich in magnesium, iron, aluminium, silicon, and oxygen. The outer \*core contains iron and sulphur. The inner core is iron, with some nickel.



Earth compositional layering

**Earth Explorer Opportunity Mission-2 (CryoSat-2)** A satellite mission, forming part of the \*European Space Agency's Living Planet Programme, to monitor the thickness of land and sea ice. It replaces CryoSat, which was lost in 2005 due to launch failure. The CryoSat-2 spacecraft was launched on 8 April 2010, from Baikonur, Kazakhstan.

**earth fall** See TOPPLE.

**earthflow** Flow of unconsolidated material down a hillslope, normally resulting from an increase in pore-water pressure, which reduces the friction between particles. Flow velocities vary from slow, when behaviour is plastic, to rapid, when behaviour is more liquid, and reflect variations in water content. Dry flows may occur when an earthquake shock breaks intergranular bonds.

**Earth Observing System (EOS)** A series of coordinated satellites in \*polar orbit, owned and operated by NASA, carrying instruments that monitor the solid Earth, land surface, atmosphere, oceans, and biosphere.



<http://eosps0.gsfc.nasa.gov/>

- A NASA series of polar-orbiting satellites to study the Earth's land, atmosphere, and oceans.

**earth pillar** See HOODOO.

**earthquake** Motion of the Earth. Tectonic earthquakes result from the release of accumulated \*strain when brittle failure occurs. This failure coincides with the release of \*stress on the rocks that actually break. Earthquakes are usually classified in terms of their depth: shallow are less than 70 km depth; intermediate 70–300 km; and deep more than 300 km. No earthquakes are known below 720 km depth. Earthquakes may also be caused by volcanic activity or induced explosions (e.g. A-bombs) to which the elastic model of tectonic earthquakes does not apply. The energy released is not stored kinetic energy, but chemical/physical energy which imposes a sudden stress that locally exceeds the strength of the rocks and no significant accumulated strain is involved as the rocks yield to the imposed stresses.

**earthquake accelerometer** See ACCELEROGRAPH.

**earthquake energy** The amplitude of a seismic signal is proportional to the energy ( $E$ ) released in an earthquake. There is no consensus on the actual equation. This is based on the \*moment magnitude scale or the \*Richter magnitude scale but also depends on the frequency of the wave, distance from the source, etc., and is of the form  $\log E = aM + b$ . The annual energy dissipation by this means is estimated to be more than  $10^{11}$  watts, of which 75% is released in shallow earthquakes and 3% in deep.

**earthquake intensity** See EARTHQUAKE MAGNITUDE.

**earthquake magnitude (earthquake intensity)** The magnitude of an \*earthquake can be estimated from its destructiveness using the \*Mercalli scale. As this measure is also dependent on the local geologic and building

context, it has been largely replaced by the **\*Richter scale**, based on the amplitude of **\*seismic waves**, and more recently by the **\*moment magnitude scale**, based on the **\*seismic moment**.

**earthquake mechanisms** Natural, artificial, or induced events that cause earthquakes. Natural mechanisms include rock falls and slides, spontaneous rock-bursts, volcanic explosions, and tectonic plate motions. Artificial and induced earthquakes can result from explosions (quarry blasts, pressure release below dam sites, nuclear bombs, etc.) or rock-bursts associated with pressure release due to mining, etc. Generally such stress releases are sudden, resulting in the release of seismic energy, but **\*bradyseisms** gradually release stress and thus do not result in an earthquake.

**earthquake prediction** Most predictions are based on attempts to determine a stress increase prior to rock rupture. This may involve **\*geodetic** measurements to monitor relative motions, changing elevation, etc., or phenomena resulting from stress accumulation (changes in the magnetization, temperature, gas release, etc.), some of which may affect animals. So far, most methods indicate only an increasing probability of seismic activity and cannot be used to predict an actual occurrence, other than the use of **\*foreshocks**, often only minutes prior to a major main **\*earthquake**, but such small earthquakes do not necessarily lead to major activity. Quiescence within an active seismic area can indicate either a gradual increase in stress or that stress release is taking place gradually. See **FOCAL MECHANISM**; **EARTHQUAKE MAGNITUDE**.


**Earth Remote Observation System (EROS)** A constellation of three high-resolution, commercial, imaging satellites, primarily for intelligence and national security applications, by ImageSat International, NV, incorporated in the Netherlands Antilles and built by Israeli Aircraft Industries Ltd. The first satellite (EROS-A) was launched on 6 December 2000, EROS-B on 24 April 2006, and EROS-C is scheduled for launch in 2019, all from the Svobodny Cosmodrome, Russia.



<https://www.imagesatintl.com/eros-sat/>

- EROS Satellite Overview.

**Earth rheological layering (mechanical layering)** The solid **Earth** forms distinct layers. The rocky **lithosphere** comprises **continental crust** 30–40 km thick (but 80 km thick beneath Tibet) and is composed mainly of up to 1 km sediments, 1–30 km **granite** and **granodiorite**, beneath that **gabbro**, and **oceanic crust** a total 7 km thick, comprising up to 0.5 km sediments, 0.5–1 km **basalt**, and beneath that gabbro. The somewhat plastic **asthenosphere** lies beneath the lithosphere and extends to a depth of about 200 km. The upper **mantle** extends to a depth of about 670 km and the lower mantle to about 2885 km. The liquid outer **core** extends to a depth of about 5150 km, and the solid inner core to the centre of the Earth, at a depth of 6360 km.



### Earth rheological layering

**Earth rotation (day length)** Astronomical observations, mainly based on **eclipses** of the **Sun**, suggest a deceleration of the Earth's rotation by about 41 seconds of arc per century. Fossil coral growth rings indicate rotation rates corresponding to about 400 days/year some 400 million years ago. Rotation is affected by periodic and irregular events. Most rapid irregular changes are meteorological or oceanic in origin and depend on the mechanical coupling between the **atmosphere**, oceans, and solid Earth, but they also affect the Earth's moment of inertia. The major long-term change is due to the slowing of the Earth's rotation by the tidal drag of the **Moon** and, to a much lesser extent, by the Sun and planets, most of it attributed to the  $M_2$  **ocean** tide. Irregular fluctuations of largely unknown origin also

occur, possibly associated with electromagnetic coupling between the Earth's *\*core* and *\*mantle*. See also [CHANDLER WOBBLE](#).

**Earth Science Enterprise (ESE)** A *\*NASA* research programme that aims to develop a scientific understanding of the Earth system and its response to natural and human-induced changes to enable improved prediction of climate, weather, and natural hazards.



<https://www.hq.nasa.gov/office/codez/plans/99nsp/12.html>

- Earth Science Enterprise.

**Earth tides** All terrestrial tides are caused by an imbalance between the centrifugal forces operating on the *\*Earth* and the changing gravitational fields of the *\*Moon*, *\*Sun*, and planets. The *\*oceanic* tide is thus a manifestation of the same phenomenon as that causing internal ('solid' Earth) tidal effects, particularly in the outer, liquid *\*core* of the Earth. The tides are seen at the surface mostly as resonance effects between the tidal components of similar harmonic frequencies as the Earth's *\*free oscillation* periodicities.

**earthy** Applied to the non-*\*metallic* *\*lustre* of porous aggregates of *\*minerals*, e.g. *\*clays*, *\*laterites*, and *\*bauxite*.

**East Australian current** Oceanic water current that flows along the east coast of Australia. This narrow (100–200 km wide) current forms the westerly part of the anticyclonic circulation in the S. Pacific. The flow velocity varies in the range 0.3–0.5 m/s. It is an example of a western *\*boundary* current.

**easterly wave** Type of weak *\*trough* in a tropical easterly airflow, which has a wavelength generally of 2000–4000 km. Such waves occur, for example, over W. Africa and in the Caribbean area where they develop in summer and autumn with a weak or absent *\*trade-wind inversion*, and in the central Pacific, when the equatorial trough is displaced northward. Disturbances in such waves often lead to tropical storms, which vary in intensity.

**Eastonian** A *\*stage* (456–450 Ma ago) of the Upper *\*Ordovician* of Australia, underlain by the *\*Gisbornian* and overlain by the *\*Bolindian*.

**East Pacific Rise** The oceanic *\*ridge* which separates the *\*Pacific Plate* from the *\*Antarctic*, *\*Nazca*, and *\*Cocos Plates*. The East Pacific Rise is a fast-spreading ridge with a maximum *\*half-spreading* rate calculated at 4.4 cm/yr, and its topographic profile is relatively smooth compared with slow-spreading ridges, e.g. the *\*Mid-Atlantic Ridge*.

**East Wind Drift** See ANTARCTIC COASTAL CURRENT.

**ebb tide** Falling *\*tide*: the phase of the tide between high water and the succeeding low water. Compare FLOOD TIDE.

**ebullient** Applied to a gas undergoing *\*ebullition*.

**ebullition** Bubbling, as when a gas escapes into water from a *\*seep* and rises to the surface, or when a liquid is agitated as it boils.

**ebullitive** Applied to a gas prone to *\*ebullition*.

**Eburonian** A northern European *\*stage* dating from about 1.8 to 0.9 Ma ago, associated with a period of glaciation at about the beginning of the *\*Pleistocene*. It may correspond to the *\*Donau* stage in the Alpine area and the *\*Nebraskan* stage in N. America.

**Ecardines** Alternative name for the class *\*Inarticulata* of the phylum *\*Brachiopoda*.

**ecdysis** Periodic shedding of the *\*exoskeleton* by some invertebrates, or of the outer skin by some *\*Amphibia* and *\*Reptilia*.

**Echinodermata (echinoderms)** Phylum of 'spiny-skinned' invertebrate animals which are entirely marine. They are characterized by an internal skeleton of porous calcite plates; a *\*pentameral symmetry* (although a *\*bilateral symmetry* is often superimposed upon this radial plan, especially in many modern *\*Echinoidea*); and the presence of a water-based vascular system, a complex internal apparatus of fluid-containing tubes and bladders which pass through pores in the skeleton and are seen from the outside as *\*tube feet*. The phylum is varied, and includes *\*Ophiuroidea* (brittle stars), Asteroidea (starfish), *\*Echinoidea* (sea urchins), *\*Holothuroidea* (sea cucumbers), and *\*Crinoidea* (sea lilies); and the extinct Edrioasteroidea, *\*Blastoidea*, *\*Cystoidea*, and members of the subphylum *\*Homalozoa* (*\*carpoids*). *Tribrachidium* from the late *\*Precambrian* of Australia is probably an echinoderm, close to the stock from which the other groups

evolved in the *\*Cambrian* and *\*Ordovician*. Echinoderms first appeared in the Lower Cambrian, but of the 20 classes known in the *\*Palaeozoic*, only six survive into the *\*Mesozoic* and on to the present day.

**Echinoidea** (echinoids; sea urchins, sand dollars, heart urchins) (phylum *\*Echinodermata*) Class of free-living echinoderms in which the body is enclosed in a globular, cushion-shaped, discoidal, or heart-shaped *\*test* built of meridionally arranged columns of interlocking, calcareous plates, which bear movable appendages (spines, pedicellariae, and spheridia). The test is composed of 20 vertical rows of plates arranged in five double rows of perforate (*\*ambulacral*) plates and five double rows of imperforate (interambulacral) plates. *\*Tube* feet, connected to the internal water-based vascular system, emerge through the pores of the ambulacra. The apical system on the upper surface consists of five ocular plates and up to five genital plates. In all *\*regular* echinoids the anus is enclosed within the apical system, but in many *\*irregular* echinoids it is in the posterior interambulacrum. The mouth is always on the lower surface and may be central or anterior in position. The class first appeared in the *\*Ordovician*; underwent a great *\*adaptive* radiation in post-*\*Palaeozoic* times, when rigid tests were evolved; experienced a marked reduction in the *\*Permian* and *\*Triassic*; and thereafter resumed its diversification, which has continued until the present day. Fossils of the inner skeletons of these echinoderms are common in both *\*Mesozoic* and *\*Cenozoic* sediments. There are about 125 Palaeozoic, 3 670 Mesozoic, 3 250 Cenozoic, and more than 900 extant species.

**Echmatocrinus** A very ancient crinoid (*\*Crinoidea*), or perhaps an ancestor of the crinoids, that lived in the *\*Cambrian* seas. It resembled crinoids, having a long stalk, cup-like *\*calyx*, and feathery arms with which it combed food from the water.

**echo dune** An elongated *\*sand \*dune* that lies approximately parallel to, but separated from, the windward side of a large obstruction. It results from the rotating airflow on the upwind side of the obstruction (the same mechanism that produces a *\*rotor cloud*).

**echo-sounding** A method for determining water depth by using the elapsed time (*t*) between the transmission of a pulse of high-frequency sound and the return of the echo from a reflector, e.g. the sea floor. Knowing the

velocity of sound in water ( $v_w$ ), the depth ( $h$ ) to the reflector is given by  $h = v_w t/2$ .

**eclipse** The partial or complete obscuration of one heavenly body by another, as perceived by an observer on one of the bodies. The proper description of an eclipse also refers to the period of time involved.

**ecliptic** The plane of the orbit of the **\*Earth** around the **\*Sun**. It forms an angle of  $23^{\circ}27'$  with the Earth's equator. The orbits of the planets all lie within  $3.4^{\circ}$  of this plane, except for those of dwarf planet **\*Pluto** ( $17.2^{\circ}$ ) and **\*Mercury** ( $7^{\circ}$ ).

**eclogite** Very rare, coarse-grained, **\*igneous** rock with a chemical composition similar to that of **\*basalt**, but noted for the presence of the rare, bright-green **\*pyroxene** omphacite, and red, almandine–pyrope **\*garnets**. Eclogites may be basalt metamorphosed by high temperatures and pressures.

**eclogite facies** A set of metamorphic **\*mineral** assemblages produced by **\*metamorphism** of **\*basic \*igneous** rocks under high-pressure, moderate-temperature conditions and typically characterized by the development of the mineral assemblage pyrope **\*garnet** and omphacite (an Na-rich, high-pressure **\*pyroxene**). Other rocks of contrasting composition, e.g. **\*shales** or **\*limestones**, are not known to show mineral assemblages characteristic of the high pressures and moderate temperatures involved. Eclogite facies can form when a slab of **\*oceanic** crust is **\*subducted** into the **\*mantle**. The basic igneous rocks of the slab can then be converted to a typical **\*eclogite** mineral assemblage.

**ecologic reef** A name proposed in 1970 by R. J. Dunham to describe a **\*reef** that is a rigid, wave-resistant, sediment-binding structure actively built by organisms. *Compare* **STRATIGRAPHIC REEF**.

**ecology** The scientific study of the interrelationships among organisms and between organisms, and between them and all aspects, living and non-living, of their environment. Ernst Heinrich Haeckel (1834–1919) is usually credited with having coined the word 'ecology' in 1869, deriving it from the Greek *oikos*, meaning 'house' or 'dwelling place'.

**economic basement** Rocks below which the chance of finding economic mineral resources is minimal. For instance, oil seldom occurs in economic

quantities below 6–7 km.

**economic geology** The study of *\*mineral* deposits that contain some valuable commodity in order to determine how they were formed. The aim is to discern those particular processes that result in the formation of valuable deposits rather than ordinary minerals of no economic importance.

**ecophenotype** See *ECOPHENOTYPY*.

**ecophenotypic effects** Non-heritable modifications of a *\*phenotype*, produced in response to factors in the environment or habitat, that become preserved in the *\*fossil*.

**ecophenotypy** The divergence of *\*phenotypes* due to developmental changes induced by local environmental conditions, producing distinct ecophenotypes. Such divergence is not heritable, but when found in the fossil record can be mistaken for speciation.

**ecosphere** See *BIOSPHERE*.

**ecostratigraphy** The study of the occurrence and development of *\*fossil \*communities* throughout geologic time, as evidenced by *\*biofacies*, with particular reference to its relevance in stratigraphic *\*correlation* and other fields such as biogeography and basin analysis.

**ECOSTRESS** See *ECOSYSTEM SPACEBORNE THERMAL RADIOMETER EXPERIMENT ON SPACE STATION*.

**ecosystem (ecological system)** Term first used by A. G. Tansley (1935) to describe the interdependence of *\*species* in the living world (the biome or *\*community*) with one another and with their non-living (abiotic) environment. Fundamental concepts include the flow of energy via food-chains and food-webs, and the cycling of nutrients biogeochemically (see *BIOGEOCHEMICAL CYCLE*). Ecosystem principles can be applied at all scales—thus principles that apply to an ephemeral pond, for example, apply equally to a lake, an ocean, or the whole planet. In Russian and central European literature ‘biogeocoenosis’ describes the same concept.

**ECOSystem Spaceborne Thermal Radiometer Experiment on Space Station (ECOSTRESS)** A high-resolution, multiple wavelength, imaging spectrometer from NASA’s Jet Propulsion Laboratory that studies the

effectiveness of water use by vegetation. It was launched on 29 June 2018, to the International Space Station.

**ecotone** Narrow and fairly sharply defined transition zone between two or more different communities.

**Ectasian** A period of the **\*Proterozoic** eon that began 1400 Ma ago and ended 1200 Ma ago. Both its commencement and its end are defined by fixed dates. The Ectasian followed the **\*Calymmian** period and was followed by the **\*Stenian** period.

**ectocochlear** Applied to those cephalopods (see **CEPHALOPODA**) in which the shell is totally external to the body. Cephalopods whose shell is encased in soft tissue are said to be 'endocochlear'.

**Ectoprocta** 1. (phylum **\*Bryozoa**) The major subphylum of bryozoans, in which the lophophore (feeding and respiratory organ) surrounds the mouth but not the anus (in contrast to the **\*Entoprocta**). Many possess a calcite skeleton. Ectoprocts are mostly, but not exclusively, marine, and have an extensive fossil record from the **\*Ordovician** to the present. 2. Alternative name for Bryozoa.

**ectotherm** See **HOMOIOTHERM**.

**edaphic** Of the soil, or influenced by the soil. Edaphic factors that influence soil organisms are derived from the development of soils and are both physical and biological (e.g. mineral and humus content, and **\*pH**).

**eddy** Motion of a fluid in directions differing from, and at some points contrary to, the direction of the larger-scale current. In air, eddies vary in size from small-scale turbulence (which can transport dust and diffuse pollutants) to large-scale movements (e.g. cyclone and **\*anticyclone** cells) within the general global circulation of the atmosphere. See **GENERAL CIRCULATION**.

**eddy currents** The **\*alternating** current induced in a conductor when it is subjected to a time-varying magnetic field in accordance with **\*Lenz's law**. The eddy currents then generate their own secondary **\*electromagnetic** field which is of considerable importance in **\*electromagnetic** exploration techniques.

**eddy viscosity** A coefficient relating the average shear stress within a **\*turbulent** flow of water or air to the vertical gradient of velocity. The eddy viscosity depends on the fluid density and distance from the river bed or ground surface. The concept of eddy viscosity is fundamental to the von **\*Karman–Prandtl** description of the velocity profile in **\*turbulent** flow, and is important in determining rates of evaporation or cooling by wind, and the shear stress exerted by rivers on moving particles on their beds.

**Edenian** A **\*stage** (454–449 Ma ago) of the **\*Ordovician** in the Lower **\*Cincinnatian** **\*series** of N. America, preceded by the **\*Shermanian** and followed by the **\*Richmondian**.

**edenite** See HORNBLENDE.

**edentulous** 1. Applied to a condition found in some **\*Bivalvia** in which hinge teeth are absent. 2. In mammals, toothless, either naturally or as a result of tooth loss.

**edge** 1. A **\*crystal** consists of a three-dimensional stacking of a **\*unit** cell defined by a space lattice. In **\*crystallography**, the three edges of the lattice are labelled *a*, *b*, and *c* (or *x*, *y*, and *z*), and they define both the edges of crystals in the seven **\*crystal** systems and their **\*crystallographic** axes. 2. In **\*remote** sensing, a boundary between area of different tones. See also EDGE ENHANCEMENT.

**edge dislocation** See DISLOCATION.

**edge enhancement** The process of increasing the **\*contrast** between adjacent areas of different tones in an image in order to define **\*edges** more clearly.

**Ediacaran** The final period of the **\*Proterozoic** eon. It began 630 (+5, –30) Ma ago and ended 542 (± 0.3) Ma ago. It commenced with the end of the Varinger (Marinoan) glaciation and a distinct change in the <sup>13</sup>C content of sedimentary deposits, and ended with the last appearance of **\*Treptichnus pedum**, a **\*trace fossil**. The Ediacaran followed the **\*Cryogenian** period and was followed by the **\*Cambrian** period, at the base of the **\*Phanerozoic** eon, and was followed by the **\*Cambrian**.



<https://ucmp.berkeley.edu/vendian/ediacaran.php>

- The Ediacaran Period.

**Ediacaran fossils** Late *\*Proterozoic* *\*fossils*, first described from Ediacara, Australia, dated to the *\*Ediacaran* period. They come from a shallow, littoral, marine environment and the animals appear to have been stranded on mudflats or in tidal pools. About 30 genera are known and include: medusoids (jellyfish), e.g. *Medusina mawsoni* and *Medusinites*; pennatulaceans (soft corals), e.g. *Charniodiscus*; annelid worms, e.g. *Spriggina*; and *Dickinsonia*, up to 1 m long, which may have resembled an anthozoan polyp, but has been variously assigned to the annelids, medusoids, and to a phylum of its own. There are also ovoid or discoid forms of unknown affinity. Since the fossils were first described by M. F. Glaessner in 1961 similar faunal types of equivalent age have been found elsewhere in the world.

**Eemian** An *\*interglacial* *\*stage* in northern Europe, dating from about 100 000 years BP to about 70 000 years BP, which may be the equivalent of the *\*Riss/Würm interglacial* of the Alpine area and the *\*Ipswichian* of the East Anglian succession.

**effective porosity** **1.** That proportion of the total *\*pore space* in a rock which is capable of releasing its contained water. *\*Clay*, for example, may have a total porosity of 50% or more, but little if any of the water contained in these pores may be released, because of the retentive forces (e.g. surface tension) that hold it within the rock. **2.** The proportion of the pore space through which *\*groundwater* flow occurs. For example, in fractured rocks the majority of flow occurs in the fractures, and intergranular pore water may be almost static. In porous rocks, some pores may have only one connection with the general pore space ('blind' pores) and so contain only static water.

**effective precipitation** Net precipitation after losses by evaporation. As higher temperatures increase evaporation, an index of effective precipitation derived from a temperature:precipitation ratio has been used as a criterion for some systems of climate classification (e.g. those of *\*Köppen* in 1936, *\*Thornthwaite* in 1948).

**effective stress** In soil, pressure between grains at points of contact; it is at equilibrium in saturated soil. Effective stress equals total pressure minus the neutral pressure of water in *\*pores*. During *\*consolidation*, effective stress increases and reaches maximum at complete consolidation before shear *\*failure* occurs.

**effective temperature ( $T_e$ )** **1.** The temperature of a planetary surface in the absence of an atmosphere. The effective temperature of Earth is some 35–40 °C lower than the actual Earth surface temperature ( $T_s$ ): the latter is approximately 15 °C owing to the *\*greenhouse* effect of the Earth's atmosphere. **2.** The temperature of saturated air that, with a wind speed no greater than 0.2 m/s, would produce the same sensation of comfort in a sedentary person wearing indoor clothes as the air to which that person is actually exposed.

**EGS** See EXPERIMENTAL GEODETIC SATELLITE.

**EGT** See EUROPEAN GEOTRAVERSE.

**Egyptian jasper** See JASPER.

**$E_H$**  See REDOX POTENTIAL.

**Eifelian (Couvinian)** **1.** A *\*stage* in the Middle *\*Devonian* *\*epoch* preceded by the *\*Emsian*, followed by the *\*Givetian*, and dated at 397.5–391.8 Ma ago (Int. Commission on Stratigraphy, 2004). **2.** The name of the corresponding European *\*stage*, which is roughly contemporaneous with the upper *\*Cunninghamian* (Australia) and upper Onesquethawian (N. America).

**Eildonian** A *\*stage* (433–428.2 Ma ago) of the Early *\*Silurian* of southeastern Australia, underlain by the *\*Keiloran* and overlain by the *\*Melbournian*.

**einkanter** See VENTIFACT.

**ejecta blanket** The blanket of debris surrounding an impact *\*crater*. It is composed of material ejected from the crater during its formation, and is laid down with stratigraphy inverted from that of the bedrock. There is typically a star-shaped distribution of ejecta around the crater rim. In addition to rock fragments excavated from the crater, and melted material,

surface material from outside the crater may be incorporated by **\*base surge** erosion or excavated by secondary craters caused by large ejected blocks. There is a chaotic size distribution of ejected material. Many martian craters are surrounded by fluidized ejecta blankets which have flowed across the surface. In contrast, lunar ejecta blankets are due mostly to ballistic sedimentation.

**Ekman depth** See **EKMAN SPIRAL**.

**Ekman spiral** Theoretical model to explain the currents that would result from a steady wind blowing over an ocean of unlimited depth and extent. In the northern hemisphere the surface layer of the water would flow at an angle of 45° to the right of the wind direction. Water at increasing depths would flow in directions more to the right, until, at a depth known as the Ekman depth, the water would move in a direction opposite to that of the wind. The Ekman depth varies with latitude and wind speed, ranging from about 45 m in latitude 45° with a wind speed of 5 m/s, to 300 m at latitude 15° with a wind speed of 20 m/s. The velocity of the water flow decreases with depth throughout the spiral. In the northern hemisphere, the net water transport is at 90° to the right of the wind direction, and is known as the Ekman transport.

**Ekman transport** See **EKMAN SPIRAL**.

**Elara (Jupiter VII)** One of the lesser satellites of **\*Jupiter**, with a diameter of 80 km ( $\pm 20$  km).

**Elasmobranchii** Subclass of shark-like fish. They have 5–7 gill slits, fairly rigid fins, **\*placoid scales**, a spiracular opening behind the jaws, numerous teeth, and claspers on the ventral fins of the male. Elasmobranchs have a cartilaginous skeleton and their fossil record consists mainly of teeth and fin rays. Well-known fossil genera include *Cladoseleche* (**\*Devonian**), and *Ctenacanthus* (Upper Devonian to Lower **\*Permian**). See **CARTILAGE**; **CARTILAGINOUS FISH**.

**elastic constants** For an **\*isotropic** material which obeys **\*Hooke's law**, where **\*strain** is linearly proportional to **\*stress**, there are only two independent elastic constants, those of stress and strain. The elastic properties of such materials are defined by elastic moduli, notably the **\*bulk modulus**, **\*shear modulus**, **\*Young's modulus**, **\*Lamé's constant**, and

**\*Poisson's ratio.** **\*Seismic wave** velocities are governed by the elastic moduli and the densities of the media through which they travel.

**elastic deformation** Temporary deformation, from which material recovers, caused by an applied **\*stress**, such that on release of the stress the body reverts to its former, unstrained condition. In purely elastic materials such deformation is described by a linear stress–strain relationship (see **HOOKE'S LAW**). In rocks, ideal elastic strain is combined with viscous components.

**elastic limit (yield point)** The point at which a material achieves its maximum elastic **\*strain** and beyond which strain is no longer linearly related to **\*stress**. Beyond the elastic limit flow or rupture induces permanent deformation.

**elastic rebound theory** Theory which holds that accumulated potential energy, stored as elastic **\*strains**, is released by faulting (i.e. when the material ruptures). Zones adjacent to the **\*fault** plane 'rebound' elastically, leaving them relatively unstrained.

**elastic wave** An acoustic or **\*seismic wave**.

**elastoviscous behaviour** See **VISCOELASTIC BEHAVIOUR**.

**E-layer** **1.** The outer **\*core** of the Earth between about 2886 km and 5156 km depth, mainly defined on **\*seismic velocities**. **2.** The **\*ionospheric** layer at about 100 km height from which 3.5 MHz radio waves are strongly reflected.

**elbaite** See **TOURMALINE**.

**elbow of capture** See **RIVER CAPTURE**.

**electrical charge** See **CHARGE**.

**electrical conductivity ( $\sigma$ )** The ease with which an electrical current will pass through a material, in units of siemens or reciprocal ohms (mhos) per metre. In materials which are assumed to be **\*isotropic**, conductivity is equal to the inverse of **\*resistivity** ( $\rho$ ), therefore  $\sigma = 1/\rho$ . See also **APPARENT CONDUCTIVITY**.

**electrical conductivity (in Earth)** This is determined at shallow depths by various **\*electrical** and **\*electromagnetic** survey methods. Upper **\*mantle**

conductivity is studied mainly by using geomagnetic variations, particularly the *\*diurnal variation*, and *\*magnetotelluric soundings*. Lower mantle conductivities are estimated by the influence on geomagnetic *\*secular* variations.

**electrical drilling** See ELECTRICAL SOUNDING.

**electrical sondes** Various *\*well-logging* devices for measuring the electrical resistivity (1/conductivity) of the rocks through which a *\*borehole* passes. See INDUCTION SONDE; MICROLOG SONDE; SELF-POTENTIAL SONDE.

**electrical sounding (vertical electrical sounding (VES), electrical drilling)** In electrical profiling, a technique in which the spacing of the *\*electrodes* or *\*coils* is expanded in order to increase the depth below the surface station from which information is obtained. A variety of *\*electrode* configurations is used for electrical *\*resistivity* sounding and *\*induced* polarization surveys. In resistivity sounding, apparent resistivity data are recorded on a log–log plot as a function of their respective current-electrode separations. *\*Master* curves are often used to provide preliminary interpretation of the curves, and followed by more detailed and sensitive computer analysis. For electromagnetic (EM) sounding the *\*coil* spacing is increased or, in *\*time-domain* EM surveys, the frequency is varied. Compare CONSTANT-SEPARATION TRAVERSING.

**electrical tomography** A geophysical technique, sometimes used in archaeology, in which metal electrodes are inserted into the ground and a current passed between them, through the subsoil. The resistivity is measured and many such measurements, taken at different depths and in different directions, allow a three-dimensional image of the site to be constructed.

**electrode array** See ELECTRODE CONFIGURATION.

**electrode configuration (electrode array)** A geometrical pattern of electrodes used in *\*electrical sounding*, *\*constant-separation traversing*, and *\*induced polarization* surveys. Usual configurations comprise two *\*current electrodes* and two *\*potential electrodes* whose separations are known and defined by a *\*geometric factor*. Common configurations include the *\*dipole–dipole*, *\*Schlumberger*, *\*square*, and *\*Wenner* arrays.

**electrode potential** See OXIDATION POTENTIAL; POTENTIAL ELECTRODE.

**electrolyte** A chemical compound which, while molten or in solution, is decomposed by the conduction of an electrical current through it. The current is moved by the passage of **\*ions** rather than, as in a metal, free **\*electrons**.

**electrolytic conduction** The conduction of an electrical current by the movement of **\*ions**, as in an **\*electrolyte**. Compare OHMIC CONDUCTION.

**electrolytic polarization** The dissociation of an **\*electrolyte** by **\*electrolytic conduction**, which drives **\*ions** of constant polarity towards a particularly-charged electrode. For example, if a current is applied to a dilute solution of sulphuric acid, positively charged hydrogen ions move towards the **\*cathode** and negatively charged sulphate ions towards the **\*anode**; after a time the solutions adjacent to the electrodes have an abundance of ions of appropriate charge and the solution is said to be polarized.

**electromagnetic methods** A range of methods by which magnetic or electrical fields generated in subsurface conductors by artificially induced **\*eddy** currents are measured. The term includes **\*radar**, electromagnetic ground **\*conductivity**, and **\*VLF**, but excludes those in which induction is insignificant, for example, **\*resistivity** and **\*induced** polarization, and those with insignificant depth penetration, for example, microwave sensing. The **\*AFMAG** method, which uses naturally generated electrical signals, is included, but **\*magnetotelluric** methods are not.

**electromagnetic radiation (EMR)** The range in radiation extending from wavelengths of less than  $10^{-12}$  m to more than  $10^3$  m. In order of increasing wavelength are included cosmic ray photons, gamma rays, X-rays, ultraviolet radiation, visible light (violet to red), infrared radiation, microwaves, radio waves, and electric currents.

**electromagnetic spectrum** The range of frequencies or wavelengths of **\*electromagnetic radiation**.

**electromagnetic wave** A wave comprising an electrical and a magnetic component at right angles to one another but which are in phase and have the same frequency. The electrical component represents the electrical field strength,  $E$ , and the magnetic component the magnetic flux density,  $B$ .

Electromagnetic waves travel at the speed of light (about  $2.998 \times 10^5$  km/s in free space), such that the wave velocity  $v = 1/(\mu_0 \epsilon_0)^{1/2}$ , where  $\mu_0$  and  $\epsilon_0$  are the magnetic permeability ( $4\pi \times 10^{-7}$  H/m) and **\*dielectric permittivity** of free space, respectively.

**electrometer** See MASS SPECTROMETRY.

**electron** Elementary particle of mass  $9.11 \times 10^{-31}$  kg and negative electrical charge of  $1.602 \times 10^{-19}$  C (coulombs). Electrons can exist independently, or in groups around the **\*nucleus** of an atom. Experiments show that electrons in an atom may occur at a range of distances from the nucleus but are most likely to exist in certain low-energy orbits or shells, and within these shells there are further subshells, the configuration being such that no two electrons in any one atom have identical properties. When an electron moves from one subshell to another of lower energy, electromagnetic radiation is given off; if an electron moves to a subshell of higher energy, electromagnetic radiation is absorbed. An electron moves about the nucleus in a circular or elliptical orbit and also spins on its axis.

**electron capture** A mechanism by which a **\*nucleus** can decrease its proton number and increase its neutron number by capturing one of its extranuclear **\*electrons**. The process can be envisaged as the reaction between an extranuclear electron and a proton in the nucleus to form a neutron and a neutrino, the neutrino then being emitted from the nucleus. The event may leave the product nucleus in an excited state, and this is then followed by the emission of a **\*gamma ray**. Removal of an extranuclear electron from the k-shell, or from higher-energy shells, leaves a vacancy that is subsequently filled by other electrons that fall into the vacant position. In the process these electrons emit a series of X-rays that can be detected.

**electronegativity** **1.** Tendency to form negative **\*ions**, measured by combining **\*ionization-potential** and electron-affinity values for an element to find the degree to which its atoms attract **\*electrons**. **2.** The ability of an atom to attract electrons, usually in non-metallic, acid-forming elements. Elements with sharply contrasting electronegativities tend to form ionic compounds (see IONIC BOND), e.g. NaCl, where Na and Cl have electronegativities of 0.9 and 3.0 respectively. Elements with similar

electronegativities are likely to form **\*covalent** bonds, e.g. CH<sub>4</sub> (methane), where C and H have electronegativities of 2.5 and 2.1 respectively. See **VALENCY**.

**Electron Losses and Fields Investigation (ELFIN)** A 3-unit **\*CubeSat** mission to study space weather developed by the University of California Los Angeles. It explores the mechanisms responsible for the loss of relativistic electrons in the radiation belts. The CubeSats were launched on 15 September 2018, into a circular, near-polar orbit at about 496 km, and by early 2019 their instruments were operational.

**electron microscope** A microscope that uses beams of high-energy electrons rather than photons to form images. The momentum ( $p$ ) of an electron depends on the voltage accelerating it, and the wavelength ( $\lambda$ ) of the electron beam depends on the momentum of the electrons:  $\lambda = h/p$ , where  $h$  is Planck's constant. A beam of high-energy electrons at 100 keV has a wavelength of approximately 3.7 pm, compared with the wavelength of about 500 nm for green visible light. Consequently, an electron beam is able to resolve details many times smaller than is possible with a light microscope. See **SCANNING ELECTRON MICROSCOPE**; **TRANSMISSION ELECTRON MICROSCOPE**.

**electron-probe microanalyser** An instrument used to determine the chemical composition of a 1  $\mu\text{m}$  diameter specimen of **\*mineral** or **\*glass** at the surface of a polished rock or mineral slice. A narrow beam of electrons is focused on to the polished surface of the specimen to cover a 1  $\mu\text{m}$  diameter spot. The electrons excite the atoms in the specimen to emit X-rays, whose wavelength is characteristic of the elements present and whose intensity at a given wavelength is proportional to the relative concentration of the element corresponding to that wavelength. By comparing the X-ray intensity at any one wavelength with that in a standard sample of known composition, the absolute concentration of the element in the sample can be deduced.

**electro-osmosis** Phenomenon whereby some fine-grained sediments with low **\*permeability** expel **\*pore** water when an electric current is passed through them. This is sometimes exploited to reduce **\*groundwater** by passing currents between **\*anodes** and **\*cathodes**.

**electrophoresis** The migration, under the influence of an electric field, of charged particles within a stationary liquid. The liquid may be a normal solution or held upon a porous medium (e.g. starch, acrylamide gel, or cellulose acetate). The rate at which migration occurs varies according to the charge on the particle and also its size and shape. The phenomenon is exploited in a variety of analytical and preparative techniques employed in studies of macromolecules.

**electropositive element** An element whose **\*electrode** potential is more positive than that of the standard hydrogen electrode which is assigned an arbitrary value of zero. Electropositive elements tend to lose **\*electrons** and form positive **\*ions**, e.g. the univalent alkali metals  $\text{Li}^+$ ,  $\text{Na}^+$ ,  $\text{K}^+$ , etc., and the divalent alkaline-earth metals  $\text{Be}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ . *Compare* ELECTRONEGATIVITY.

**Elektro** The name of a Russian meteorological satellite that was launched into **\*geostationary orbit** in 1994 to monitor conditions over the Indian Ocean. It failed to transmit any data and was shut down in 1998.

**Elektro-L** A Russian meteorological satellite that was launched into **\*geostationary orbit** from Baikonur Cosmodrome, Kazakhstan, on 20 January 2011, carried on a Zenit rocket. Designed to remain operational for ten years, Elektro-L provides data to help with weather forecasting, analysis of ocean conditions, and monitoring of space 'weather'.



<http://www.russianspaceweb.com/elektro.html>

- A Roskosmos mission launched in 2011 to enable local and global weather forecasting and measure the Earth's magnetic field.

**eleutherozoan** Applied to unattached echinoderms. Used informally, and in some classifications Eleutherozoa is a subphylum of the **\*Echinodermata**. *Compare* PELMATOZOAN.

**elevation correction** 1. The correction applied to a measurement of gravitational attraction to allow for the distance to the station from the theoretical reference surface and for the attraction of the rocks in this zone. It is a combination of the **\*free-air** and **\*Bouguer** corrections. 2. **(static**

**correction)** The correction which is made to seismic travel-time data to compensate for irregular topography and to reduce the data to a common **\*datum**. Land ‘statics’ are extremely important in seismic surveying on land. As part of a ‘statics’ survey, refraction shooting is undertaken to determine the depth and velocity of the weathered zone (the low velocity layer, or LVL). This is done in order to design more effective seismic source parameters (e.g. ideally, the shot depth should be below the weathered layer) thereby optimizing the seismic reflection surveys which are undertaken after an LVL survey. In marine surveys, marine ‘statics govern the source– **\*hydrophone** geometry and provide a correction for the finite **\*offset** between the source **\*arrays** and the hydrophone **\*streamer**.

**elevation head** See ELEVATION POTENTIAL ENERGY.

**elevation potential energy (elevation head)** The energy possessed by a mass, e.g. a body of water, by virtue of its being raised above a particular datum point, usually taken as either sea level or local ground level. The energy may be released when the mass is allowed to fall to a lower level, and may be harnessed, e.g. in the case of water by powering a turbine in a hydroelectric scheme. Elevation head is the energy possessed by a unit weight of water at a point, due to this cause. See also BERNOULLI EQUATION; DARCY’S LAW; PRESSURE HEAD.

**elf** A **\*transient luminous event** that consists of a rapidly expanding luminous disc, lasting less than 0.001 second, high above a strong cloud-to-ground lightning stroke.

**ELFIN** See ELECTRON LOSSES AND FIELDS INVESTIGATION.

**Élie de Beaumont, Léonce** (1798–1874) A French geologist who made a detailed study of European folded rocks, and concluded that these showed evidence of distinct mountain building episodes. His overly geometric treatment of the directions of mountain systems was criticized by his contemporaries, but his theory that the Earth is cooling, and therefore shrinking, found wider support.


**elite** In **\*ichnology**, applied (a) to a structure constructed within a sediment by living organisms that indicates a high level of biological activity because of its high content of oxygen or organic matter, and (b) to a **\*trace fossil** that is more conspicuous than others nearby because some feature of its

structure or composition has been emphasized by the processes of **\*diagenesis**.

**elliptical polarization** In optical mineralogy, light passing through a mineral may be split into two rays. The velocity of one ray is uniform in all directions, to give a spherical wavefront; the other ray varies in velocity with direction, to give an ellipsoidal wavefront. Elliptical sections so produced are used to determine the optical properties of minerals.

**ellipticity** See **PRINCIPAL STRAIN RATIO**.

**El Niño** A weakening, or in extreme cases reversal in direction, of the prevailing south-easterly **\*trade winds** over the equatorial South Pacific Ocean that is associated with a change in the distribution of air pressure over the South Pacific known as a **\*southern oscillation**. A similar phenomenon may also occur in the Atlantic. Approximately once every seven years, during the Christmas season (the name refers to the Christ child), prevailing trade winds weaken and the **\*Equatorial** countercurrent strengthens. Warm surface waters, normally driven westward by the wind to form a deep warm pool off Indonesia, flow eastwards to overlie the cold waters of the **\*Peru current**. In exceptional years, e.g. 1891, 1925, 1953, 1972–3, 1982–3, 1986–7, 1994–5, and 1997–8, the extent to which the upwelling of the nutrient-rich cold waters is suppressed causes the death of a large proportion of the plankton population and a decline in the numbers of surface fish. El Niño conditions bring heavy rain to the ordinarily arid western coast of S. America, drought to Indonesia, and produce climatic effects throughout the Pacific region and beyond. See **TELECONNECTIONS**.



**El Niño**

**elongation index** ( $I_E$ ) The percentage by weight of particles whose long dimension is greater than 1.8 times the mean dimension measured with a standard gauge. The elongation,  $n$ , is length divided by breadth and the elongation ratio is  $1/n$ .

**elongation ratio** See [ELONGATION INDEX](#).

**ELR** See [ENVIRONMENTAL LAPSE RATE](#).

**Elsasser, Walter Maurice** (1904–91) A German-born physicist, Elsasser became a US citizen in 1940. He worked on the structure of the nucleus and on terrestrial magnetism. In 1946, he proposed that a ‘\*self-exciting dynamo’ in the Earth’s outer \*core might be the cause of the magnetic field.

**Elsonian orogeny** A \*Proterozoic phase of mountain building that occurred about 1500–1400 Ma ago, affecting what are now the Central and W. Nain Provinces of the Canadian Shield. It was preceded by the \*Hudsonian and succeeded by the \*Grenvillian orogenies.

**Elsterian** A glacial period in northern Europe, dating from about 0.5 Ma ago to 0.3 Ma ago, that is probably equivalent to the \*Mindel glaciation of the Alpine area.

**elutriation** A method of \*grain-size analysis in which the finer \*grades are separated from coarser and heavier particles by the use of a rising current of air or water, which carries the light particles upwards and allows the heavier grains to sink. By controlling the velocity of the flow, grains of different sizes can be separated.

**eluvial deposit** A residual accumulation of \*ore minerals that occurs above the \*source rock and has experienced no transport. In eluvial deposits concentrations of economic minerals are usually increased above the value of those in the underlying rocks by the removal of soluble elements from the host material, e.g. the tin deposits of Rondonia, Brazil. *Compare ALLUVIAL DEPOSITS.*

**eluviation** Removal of soil materials in suspension or in solution from surface \*horizons, and with partial deposition in the lower horizons of \*soil profiles. Removal in solution is called \*leaching, and hence the term 'eluviation' is often limited in use to removal in suspension.

**elvan** See QUARTZ PORPHYRY.

**embankment dam** Earth or gravel wall above ground level to confine water in \*reservoirs or channels, to prevent flooding, to restrain \*tailings in a pond, or to carry roads and railways. On weak or unconsolidated deposits the \*load is spread by wide embankments with a core of impervious \*concrete or rolled clay and with an outer layer of crushed rock or soil.

**Embolomeri** See ANTHRACOSAURIA.

**Embry and Clovan classification** A \*limestone classification, based on textural principles, which expands the \*Dunham classification to include conglomeratic limestones and different types of organically bound limestone. E. F. Embry and J. E. Clovan proposed their system in 1971.

They retain the Dunham terminology of mudstone, wackestone, packstone, and grainstone for limestones with particles less than 2 mm in size, but for limestones containing particles more than 2 mm in size they define two new terms: floatstone (*\*matrix-supported* limestone in which more than 10% of the *\*clasts* are larger than 2 mm in size); and rudstone (clast-supported limestone in which more than 10% of the clasts are coarser than 2 mm in size). For *\*autochthonous* limestones whose original components were organically bound during deposition Embry and Clovan replace the Dunham term 'boundstone' with three new terms: bafflestone (autochthonous limestone whose original components were bound during deposition by organisms which act as baffles, permitting sediment to be trapped in the lee of the baffles); bindstone (autochthonous limestone whose original components were bound during deposition by organisms which encrust and bind, e.g. algae); and framestone (autochthonous limestone whose original components were bound during deposition by organisms which built a rigid framework, e.g. corals (*\*Anthozoa*) in a *\*reef* structure).

**embryophytes** Land plants, including those from which all later terrestrial groups are descended. Gene sequences and comparative morphology suggest land plants form a *\*monophyletic* group, descended from *\*Charophyceae*. Embryophyte fossils occur in rocks of middle *\*Ordovician* (Llanvirn, about 470 Ma ago) age.

**emerald** See BERYL.

**emery** A greyish-black variety of *\*corundum* (Al<sub>2</sub>O<sub>3</sub>) which can be crushed and powdered for use as an abrasive for polishing hard surfaces.

**Emilian** See CASTLECLIFFIAN; QUATERNARY.

**emission spectrum** See SPECTRUM.

**emissivity** The ratio of *\*exitance* of a body to the exitance of a *\*black* body at the same temperature.

**EMR** See ELECTROMAGNETIC RADIATION.

**Emsian** 1. An *\*age* in the Early *\*Devonian* epoch preceded by the *\*Praghan*, followed by the *\*Eifelian*, and dated at 407–397.5 Ma ago (Int. Commission on Stratigraphy, 2004). 2. The name of the corresponding

European **\*stage**, which is roughly contemporaneous with the lower **\*Cunninghamian** (Australia) and part of the Onesquethawian (N. America).

**emu** Electromagnetic units in the **\*c.g.s.** units (gauss, oersteds, etc.). These have now been replaced by SI units of **\*ampere** per metre, weber per metre, and **\*tesla**.

**En** See ENSTATITE.

**en-** From the French *en* meaning 'in', a prefix meaning 'in', 'into', or 'inside'.

**enamel** Crystals of a calcium phosphate-carbonate salt, containing 2–4% of organic matter, which are formed from the epithelium of the mouth and provide a hard outer coating to **\*denticles** and the exposed part of teeth.

**enantiomorphy** The property of different structural forms of the same substance, e.g. certain crystals, of appearing to mirror each other (as do the right and left hands). No rotation or inversion can bring one into coincidence with the other. A good example of this property is found in **\*quartz**.

**enargite** A **\*sulphide** **\*mineral**  $\text{Cu}_3\text{AsS}_4$  and **\*end-member** of an **\*isomorphous** series, the other end-member being famatinite  $\text{Cu}_3\text{SbS}_4$ ; sp. gr. 4.5; **\*hardness** 3; greyish-black; **\*metallic** **\*lustre**; occurs as irregular grains in association with other sulphides in copper-rich mineral deposits.

**enation theory** The theory that accounts for the origin of the fern leaf by suggesting that it arose from the development of simple outgrowths (enations). Any such theory has to account for the large, branching fronds of a fern with branching veins (a 'megaphyll') and also for the small leaves ('microphylls') with a single median vein.

**Enceladus (Saturn II)** A major satellite of **\*Saturn**, discovered in 1789 by Sir William Herschel. It has the brightest surface of any body in the **\*solar** system (albedo more than 0.9), composed of clean, fresh ice (not necessarily water ice). The surface is cratered, but there are also smooth plains and long linear cracks and ridges. The surface appears to be young, probably less than 100 Ma old, indicating it has been geologically active until very recently, and possibly is still active with some kind of water volcanism. This activity may make Enceladus the source of the material

comprising the tenuous E ring of Saturn. Enceladus is much too small to be heated by radioactive decay; the heat would have dissipated long ago. The orbit of Enceladus is locked in a 1:2 **\*resonance** with **\*Dione**, which may provide some **\*tidal heating**, but probably not enough to melt water ice. Enceladus is 238 020 km from Saturn; its radius is 249.1 km; mass  $0.73 \times 10^{20}$  kg; mean density 1120 kg/m<sup>3</sup>; visual albedo 1.0. The **\*Cassini** mission to Saturn includes a fly-by of Enceladus.

**Encke** A **\*comet** with an orbital period of 3.3 years; **\*perihelion** date 11 June 1997; perihelion distance 0.339 AU.

**Endeavour hydrothermal vents** A group of **\*hydrothermal vents** located in the northeastern **\*Pacific Ocean** about 260 km west of Vancouver Island, Canada, at a depth of 2250 m, on the **\*Juan de Fuca Ridge**. Due to its rich biodiversity, the Endeavour hydrothermal vents are designated a marine protected area under the Canada Oceans Act.

**endemism** Situation in which a species or other taxonomic group (*see* **CLASSIFICATION**; **TAXONOMY**) is restricted to a particular geographic region, due to factors such as isolation or response to soil or climatic conditions. Such a taxon is said to be endemic to that region. The size of the region in this context will usually depend on the status of the taxon: thus a family will be endemic to a much larger area than a species, all other things being equal. Reference is frequently made to ‘narrow endemics’, i.e. taxa with markedly restricted ranges. Some of these are evolutionary relics, such as the maidenhair tree (*Ginkgo biloba*), the only surviving species of the **\*Ginkgoales**, confined to Chekiang Province, China, where it was discovered in 1758.

**endichnia** *See* **TRACE FOSSIL**.

**end-member 1.** One of two or more simple substances forming the extreme ends of a **\*solid solution** series, e.g. **\*albite** (NaAlSi<sub>3</sub>O<sub>8</sub>) is one end-member, and **\*anorthite** (CaAl<sub>2</sub>Si<sub>2</sub>O<sub>8</sub>) the other end-member, in the **\*plagioclase series**. **2.** In palaeontology, one of the two distinct forms between which gradual variation occurs.

**end-member textural classification** A method of classifying **\*sediments** consisting of a mixture of **\*grain** sizes which defines the sediment type by plotting the relative proportions of three end-members (e.g. **\*sand**, **\*silt**,

\*clay; or \*gravel, sand, mud). There are many end-member classifications that have been proposed, but all are based on the relative proportions of three end-member components.

**end moraine** See MORAINE.

**endobiont** An organism that lives beneath the surface of a substrate, e.g. the bed of a sea or lake.

**endobyssate** Applied to the habit of specific bivalves (\*Bivalvia) that live in \*sediment. In contrast to \*epibyssate forms, the byssus (see BYSSATE) of these animals is used to anchor the animal within a burrow or boring.

**endochondral bone** See BONE.

**endocochlear** See ECTOCOCHLEAR.

**endocone** Apically pointed, conical layers of calcareous material that may fill the \*siphuncle in some groups of nautiloids (\*Nautiloidea). The layers build forward from the rear of the conch.

**endogenetic processes** Processes which originate below the Earth's surface; particularly applied to Earth movements (by faulting and \*earthquakes) and volcanic activity.

**endogenous dome** See DOME (2).

**endolith** An \*ooid that has been infested by boring micro-organisms (e.g. fungi or \*cyanobacteria).

**endopunctate** See PUNCTATE.

**endorheic lake** A lake that loses water only by evaporation (i.e. no stream flows from it). Compare EXORHEIC LAKE.

**endoskeleton** A skeleton that is contained within the body. In vertebrates, the endoskeleton comprises the axial skeleton and the appendicular skeleton. In \*Echinodermata, the skeleton lies beneath the body surface and technically it is therefore an endoskeleton. Compare EXOSKELETON.

\*\*\*\*\*

**endotherm** See HOMOIOOTHERM.

**endrumpf** In the model of landscape evolution proposed by [Walther](#) [\\*Penck](#), the final stage of [\\*erosion](#) that follows the disappearance of [\\*inselbergs](#) and the [\\*pediments](#) surrounding them. The endrumpf is a landscape of low-angle, concave slopes that are slowly retreating.

**en échelon** The parallel or subparallel alignment of separate structural features (e.g. tension fractures) which are arranged obliquely to a specific directional axis.

**energetic particles detector (EPD)** An instrument carried on the [\\*Galileo](#) spacecraft that provided data on the jovian [\\*magnetosphere](#). The EPD measured the angular distribution and energy of [\\*ions](#) and [\\*electrons](#), the composition of ions, and the elements from helium to iron.

**energy budget** See [RADIATION BUDGET](#).

**energy of activation** See [ACTIVATION ENERGY](#).

**engineering geophysics** The application of geophysical survey techniques to site investigations for civil engineering purposes. Common investigations include: finding the depth to bedrock for foundation studies; determining the [\\*rippability](#) of material for excavations; measuring the degree of fracturing of rock; the detection of underground fissures, cavities, and mineshafts; the location of pipes; and, in a marine environment, the determination of the strength of sediment infill and the detection of dangerous near-surface gas pockets before drilling with a jack-up rig.

**englacial** Contained within the interior of a glacier, as opposed to being at its base (subglacial) or on its surface (supraglacial). Normally the term is applied to meltwater or [\\*drift](#).

**Enhanced Fujita Scale** A 6-point scale used to report the intensity of a [\\*tornado](#) based on the wind speed and damage caused. It is derived from the [\\*Fujita Tornado Intensity Scale](#), which it replaced in the USA in 2007 and in Canada in 2013. See [APPENDIX C](#).

**enhancement seismograph** See [SEISMOGRAPH](#).

**enigmatic taxon** A genus or higher [\\*taxon](#) of restricted diversity, comprising only one or a few species with affinities that are poorly

understood. Many species belonging to enigmatic taxa are known from very few specimens; some have not been seen since they were first described.

**EnMAP** See ENVIRONMENTAL MONITORING AND ANALYSIS PROGRAM.

**enriched uranium** Uranium containing up to 3%  $^{235}\text{U}$ . It is produced industrially for use in some nuclear reactors. Natural uranium contains as little as 0.7%  $^{235}\text{U}$ .

**ensialic belt** An *\*orogenic* belt developed on sialic *\*continental crust*. Ensialic belts are thought to have developed by mechanisms involving little horizontal movement and so are unrelated to *\*plate tectonics*. Many *\*Proterozoic* belts have been classed as ensialic belts, although increasingly these have also been interpreted as *\*collision zones* of *\*plates*.

**ENSO event** The full cycle of an *\*El Niño*, followed by *\*La Niña*, linked to a *\*southern oscillation*.

**enstatite (En)** An important *\*orthorhombic pyroxene* (*\*orthopyroxene*, or opx)  $\text{Mg}_2\text{Si}_2\text{O}_6$  and *\*end-member* of an *\*isomorphous* series, the other end-member being orthoferrosilite (Fs)  $\text{Fe}_2\text{Si}_2\text{O}_6$ ; this series forms the base of a compositional equilateral triangle whose third apex is *\*wollastonite* ( $\text{CaSiO}_3$ ) and into which the common pyroxene minerals (*\*augite*, *\*diopside*, *\*pigeonite*) can be plotted, as well as *\*olivine*; sp. gr. 3.2 (En) to 3.96 (Fs); *\*hardness* 5.5; colourless or greyish-white, with a green tinge, or brownish-green; *\*vitreous lustre*; *\*crystals* are *\*prismatic* or *\*tabular*, but normally forming very irregular grains; *\*cleavage* good prismatic {110}, poor {010}, {100}; occurs in magnesium-rich magmatic rocks (e.g. *\*peridotites*, *\*gabbros*, *\*norites*, and *\*basalts*) and in contact *\*metasomatic* zones associated with these rocks; and in *\*partial* melt envelopes around very high-temperature *\*basic igneous* intrusions.

**entablature** An irregular arrangement of *\*columnar jointing* that lies above a *\*colonnade* in a thick lava flow.

**entelechy** See ARISTOGENESIS.

**enterolithic structure** Irregular, tight to open folding, developed particularly in *\*evaporite* sequences due to volume changes in the rock brought about by chemical transformations of the salts. Such structures are

commonly formed by the swelling of *\*anhydrite* during its *\*hydration* to *\*gypsum*.

**enteron** See ANTHOZOA.

**enthalpy (*H*)** Heat content per unit mass of a substance measured as the internal energy plus the product of its volume and pressure.

**entisols** In the US *\*soil taxonomy*, an order of embryonic mineral soils that have no distinct pedogenic horizons. Representing only the initiation of *\*soil-profile* development, Entisols are common on recent *\*floodplains*, steep eroding slopes, stabilized sand *\*dunes*, and recent deep ash or wind deposits.

**Entoprocta** (phylum *\*Bryozoa*) Subphylum of freshwater bryozoans which entirely lack a mineralized skeleton. The lophophore surrounds both the anus and the mouth. Many fossil forms are known, but the subphylum is known only from the *\*Cenozoic*. Formerly the Entoprocta was classified as a separate animal phylum, although its many resemblances to the Bryozoa were recognized.

**entrainment** Process by which air from the environment outside a growing cloud is caught into the rising convective current within the cloud and mixed with the cloudy air. This is significant in that it reduces the buoyancy of the rising current and causes cloud growth by reason of the cooler, drier air which is introduced. This also causes some evaporation of cloud droplets. When very dry air is introduced, entrainment can produce rapid dissipation of the cloud.

**entrenched meander** See MEANDER.

**entropy** 1. Measure of disorder or unavailable energy in a thermodynamic system; the measure of increasing disorganization of the universe. 2. See LEAST-WORK PRINCIPLE; LEAST-WORK PROFILE.

**environmental geology** Study of the problems resulting from natural hazards and human exploitation of the natural environment. The geologic techniques used include those of engineering geology, economic geology, hydrogeology, etc., as applied to waste disposal, water resources, transport, building, mining, and general land use.

**environmental lapse rate (ELR)** The actual rate at which the air temperature changes with height in the atmosphere in a particular place at a particular time. The overall average rate is a decrease of about 6.5 °C/km, but the rate varies greatly from time to time and place to place. Where the lapse rate of temperature is negative (temperature increases with height), an inversion is said to exist.

**Environmental Monitoring and Analysis Program (EnMAP)** A German satellite programme to monitor rocks and soils, vegetation, and inland and coastal waters on a global scale. It is scheduled for launch in 2020.

**EnviSat** See ENVIRONMENTAL SATELLITE.

**Environmental Satellite (EnviSat)** A space mission of the \*European Space Agency (ESA) to study and monitor the Earth's environment and the management of its resources. The satellite was launched on 1 March 2002, into a Sun-synchronous near-circular orbit.

**Eoarchaeon** The earliest \*era of the \*Archaean \*eon, 3800–3600 Ma ago, preceded by the \*Hadean eon and followed by the \*Palaeoarchaeon era. During this era, for the first time the Earth had a solid crust.

**Eobacrites sandbergeri** First recorded from the Lower \*Ordovician of Bohemia, Czech Republic, *E. sandbergeri* is the species thought by some workers to be the earliest representative of the cephalopod molluscs, the \*Ammonoidea. *Eobacrites* was a straight-shelled form with no internal deposits related to buoyancy control.

**Eocambrian** Little-used term employed in the description of sequences of unfossiliferous rocks that were deposited at the end of the \*Precambrian.

**Eocene** \*Palaeogene \*epoch which began at the end of the \*Palaeocene (55.8 Ma ago) and ended at the beginning of the \*Oligocene (33.9 Ma ago). It is noted for the expansion of mammalian stocks (horses, bats, and whales appeared during this epoch), and the local abundance of \*nummulites (marine protozoans of the \*Foraminiferida). The Eocene epoch is divided into the \*Ypresian, \*Lutetian, \*Bartonian, and \*Priabonian \*ages.



<https://ucmp.berkeley.edu/tertiary/eocene.php>

- The Eocene Epoch.

**Eocrinoidea** (subphylum Blastozoa) Extinct class of \*cystoid-like \*echinoderms, with \*radial symmetry, which range in age from Lower \*Cambrian to Middle \*Silurian (Lower \*Palaeozoic). Their globular or flattened \*thecae are composed of numerous, irregularly arranged plates that lack the thecal pores typical of cystoids, and their sutural pores (i.e. those at plate margins) are unlike those of cystoids. Food grooves are extended on an exothecal skeleton of erect, unbranched, \*biserial, food-gathering appendages (called \*brachioles), which are not homologous with the arms of \*crinoids. The eocrinoids are the earliest blastozoans and may have included ancestors to other cystoid groups, possibly to crinoids also, but the question is still unresolved.

**eocrinoids** See EOCCRINOIDEA.

**Eodelphis** Known from the Upper \*Cretaceous of N. America, *Eodelphis* is a genus of early opossums, \*marsupials with long prehensile tails. Opossums are arboreal in habit and are thought to represent the ideal stem stock from which evolved the whole marsupial group. Several species of *Eodelphis* are known from the Milk River Formation (Upper Cretaceous) of Alberta. The teeth of *Eodelphis* are sharply cusped and somewhat primitive in form.

**Eoembryophytic** An epoch in the evolution of plants, proposed in 1993 by J. Gray, that lasted from the early Llanvirn (about 471.8 Ma ago) in the middle \*Ordovician to the late Llandovery (about 428.2 Ma ago) in the Early Silurian. Fossil tetrads comprising four spores bound by a membrane make their first appearance during this epoch and occur over a wide geographic range.

**'Eohippus'** See EQUIDAE; HYRACOTHERIUM.

**eolian** See AEOLIAN.

**eon** 1. The largest \*geologic-time unit, incorporating a number of \*eras. The equivalent \*chronostratigraphic unit is the \*eonothem. Originally, two eons were proposed in 1930 by G. H. Chadwick. The younger was the \*Phanerozoic eon (time of evident life), comprising the \*Cenozoic, \*Mesozoic, and \*Palaeozoic eras, and this term is still used. The term suggested for the preceding eon was the \*Cryptozoic (time of hidden life).

This time has also been called the **\*Archaeozoic** (time of most ancient life), but it is now known as the **\*Proterozoic** (2500–542 Ma ago). Earlier time has most commonly been known simply as the **\*Precambrian**, but now comprises the **\*Hadean** (4567.17–3800 Ma ago) and **\*Archaean** (3800–2500 Ma ago) eons. **\*Hadean** is an informal name and the Archaean starting date is not defined stratigraphically. The term ‘Precambrian’ is still in frequent use but is **\*informal**. **2.** A time unit of  $10^9$  present Earth years.

**eonothem** The **\*chronostratigraphic unit** equivalent to the **\*geologic-time unit ‘\*eon’**. At present it is a little-used term.

**EOS** See **EARTH OBSERVING SYSTEM**.

**Eosimias** A fossil primate, known mainly by jaws and teeth, from the Middle **\*Eocene** of southern China, described in 1994 by C. K. Beard, Tao Qi, M. R. Dawson, Banyue Wang, and Chuankuei Li. According to its describers, it has the features of a very primitive member of the Simiiformes (monkeys, apes, and humans) and may stand at the base of this group; the name means ‘dawn monkey’. The species Beard et al. discovered is now known as *E. sinensis*. Three more species have since been recognized: *E. centennicus* and *E. dawsonase*, both discovered in 1995, and *E. paukkaungensis*, first described in 2010.

**Eosphaera** A tiny, sphaeroidal **\*fossil** that is associated with other **\*microfossils** in the Gunflint Chert, a rock, some 2 billion years old, from western Ontario, Canada. It is suggested that these fossils are the remains of the first photosynthetic organisms.

**Eotracheophytic** An epoch in the evolution of plants, proposed in 1993 by J. Gray, that lasted from the latest Llandovery (about 428.2 Ma ago) in the Early **\*Silurian** until the mid-**\*Lochkovian** (about 413 Ma ago) in the Early **\*Devonian**. During this epoch simple plant spores, dispersed individually, become increasingly common in several plant groups, including early vascular plants, and the earliest undoubted plant megafossils have been dated to this time.

**Eötvös effect** The gravitational effect due to the vertical component of the **\*Coriolis** effect when the **\*gravimeter** is in motion, e.g. while at sea. It depends on the velocity and direction of the motion.

**EPD** See **ENERGETIC PARTICLES DETECTOR**.

**epeiric sea (epicontinental sea)** Shallow sea which extends far into the interior of a continent, e.g. Hudson Bay and the Baltic Sea. The term also denotes shallow sea areas that cover the **\*continental shelf** and are partially enclosed, e.g. the North Sea.

**epeirogenesis** The large-scale upward or downward movements of continental or oceanic areas. Epeirogenic movements should not be confused with the more dynamic mountain-building episodes of an **\*orogeny**.

**ephemeral stream** A stream which flows only after rain or snow-melt and has no **\*baseflow** component. A desert **\*wadi** may form an ephemeral stream.

**ephemerides** See EPHEMERIS.

**ephemeris (pl. ephemerides)** The position in the sky of an astronomical object or artificial satellite at a specified date and time.

**epi-** From the Greek *epi* meaning ‘upon’, a prefix meaning ‘upon’, ‘in addition to’, or ‘above’.

**epibenthos** The organisms living on the surface of the sea bed or bed of a lake.

**epibole** See ACME ZONE.

**epibyssate** Applied to animals that use the byssus (see BYSSATE) to anchor themselves to rock or seaweed. Compare ENDOBYSSATE.

**epicentral angle ( $\Delta$ )** The angular distance between an **\*earthquake \*focus** and a seismic station.

**epicentre** The point on the Earth’s surface immediately overlying an **\*earthquake \*focus** (the **\*hypocentre**).

**epichnia** See TRACE FOSSIL.

**epiclast** A **\*clast** produced by the chemical or mechanical **\*weathering**, or other surface processes, of volcanic rock. Compare ALLOCLAST; AUTOCLAST; HYDROCLAST.

**epicontinental sea** See EPEIRIC SEA.

**epicratonic** Applied to processes which are active on the surface of a **\*craton**, and the products of such processes.

**epidote (pistacite)** A rock-forming **\*mineral**  $\text{Ca}_2(\text{Al}_2\text{Fe}^{3+})\text{Si}_3\text{O}_{12}(\text{OH})$  and a member of the epidote group of minerals, which includes **\*zoisite**, **\*clinozoisite**, and **\*allanite**; sp. gr. 3.4–3.5; **\*hardness** 6.5; **\*monoclinic**; normally pistachio-green, but also various shades of green, yellow, grey, or black; **\*vitreous lustre**; **\*crystals** are **\*prismatic** parallelograms, often rod-like and isometric, often showing radial, fibrous, and columnar masses; **\*cleavage** perfect {001}; it occurs in **\*hydrothermal** formations, especially associated with altered, **\*basic**, **\*igneous** rocks, and also in **\*contact** metamorphic zones with **\*quartz**, **\*chlorite**, **\*calcite**, and **\*sulphides**, and can also replace various minerals, such as **\*amphiboles**, which break down under late-stage hydrothermal alteration.

**epifaunal** Applied to benthic organisms that live on the surface of the sea bed, either attached to objects on the bottom or free-moving. They are characteristic of the **\*intertidal zone**. *Compare* INFAUNAL.

**epigene** Produced or occurring at the Earth's surface. The term is used especially in relation to the processes of **\*weathering**, **\*erosion**, and deposition. '**\*Epigenetic** drainage' is sometimes used as a synonym for 'superimposed drainage'.

**epigenesis** The hypothesis that an organism develops by the new appearance of structures and functions. An alternative hypothesis (termed 'preformation') is that the development of an organism occurs by the unfolding and growth of characters already present in the egg at the beginning of development.

**epigenetic drainage** Superimposed drainage. *See also* DRAINAGE; EPIGENE.

**epigenetic ore** Deposit later in origin than the host rock.

**epilimnion** The upper, warm, circulating water in a thermally stratified lake in summer. Usually it forms a layer that is thin compared to the **\*hypolimnion**.

**Epimetheus (Saturn XI)** One of the lesser satellites of **\*Saturn**, discovered in 1979 by **\*Pioneer II**, with a radius measuring  $69 \times 55 \times 55$  km; mass  $0.0055 \times 10^{20}$  kg; mean density  $630 \text{ kg/m}^3$ ; visual albedo 0.8.

**Epiphyton** See RHODOPHYCEAE.

**episodic evolution** The fossil record is characterized by extinction events and succeeding phases of rapid evolutionary innovation. The overall picture is thus one of episodic **\*evolution**. However, the term has recently acquired other connotations, and tends to be linked with **\*punctuated equilibrium**.

**epitaxy** The overgrowth of one **\*crystal** on another such that the structural orientations of the two **\*minerals** are specifically related. The minerals themselves do not have to be closely similar in their structures.

**epitheca** 1. The outer wall of a **\*corallite** in the subclass **\*Zoantharia**. 2. See DINOPHYCEAE.

**epithermal** Vein deposit formed within about a kilometre of the Earth's surface by hot (50–200 °C), ascending solutions which often produce shatter zones. Typical minerals are **\*stibnite**, **\*cinnabar**, **\*gold**, and **\*silver**.

**epoch** One of the intervals of geologic time recommended by the International Subcommittee on Stratigraphic Terminology. An epoch is ranked as a third-order time unit, and is the equivalent of the chronostratigraphic unit **\*series**. Several epochs form a **\*period**; several periods an **\*era**. Epochs are themselves subdivided into **\*ages**. When used formally, the initial letter is often capitalized, e.g. Early Devonian Epoch.

**e-process** See EQUILIBRIUM PROCESS.

**epsilon cross-bedding** One of a series of cross-bedding types proposed by J. R. L. Allen in 1963. Epsilon cross-bedding is the type of cross-bedding formed by the lateral accretion of a river **\*point** bar. Other terms (nu, gamma, beta cross-bedding, etc.) proposed in the same scheme are not in current usage.

**epsomite (Epsom salts)** **\*Mineral**,  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ ; sp. gr. 1.7; **\*hardness** 2.5; **\*orthorhombic**; colourless to white; **\*vitreous** to earthy **\*lustre**; crystals rare except locally, and show a fibrous structure; **\*cleavage** one direction, perfect; occurs as encrusting masses in caves and old mine workings, and also in the oxidized zones of **\*pyrite** deposits in arid regions. It is readily soluble in water and has a bitter taste. It is named after mineral springs at Epsom, England.

**Epsom salts** See EPSOMITE.

**equal-area net (Schmidt–Lambert net)** A stereographic net (i.e. a system of coordinates providing a two-dimensional representation of a sphere) used to display structural data in graphic form. The net is divided into 2° blocks of equal area bounded by \*great and \*small circles, and the data plotted on it refer to the orientation of features. The range of orientations and any tendency towards preferred orientation may be observed.

**equant** Applied to a \*clast whose dimensions are broadly similar on each of its long, intermediate, and short axes. See PARTICLE SHAPE.

**Equatorial countercurrent** See EQUATORIAL CURRENT.

**Equatorial current** Oceanic current which flows in an east–west direction in the equatorial regions of all the oceans. The broad (1000–5000km) westward-flowing currents (North and South equatorial currents) are separated by a comparatively narrow westward-flowing countercurrent (the Equatorial countercurrent). The flow tends to be limited to the upper 500 m of the water column and the velocity is 0.25–1.0 m/s. The strength and position of the equatorial currents are controlled by the overlying wind system.

**equatorial orbit** An \*orbit of a satellite in which the orbital plane deviates from the equatorial plane of a planet by less than 45°. Compare POLAR ORBIT.

**equatorial plane** See PLANE OF PROJECTION.

**equatorial trough** Shallow low-pressure zone around the equator, where the trade-wind systems tend to converge and air rises. See also INTERTROPICAL CONVERGENCE ZONE; DOLDRUMS.

**Equatorial undercurrent (Cromwell current)** A shallow, subsurface, eastward-flowing current in the central \*Pacific Ocean between 1.5° S and 1.5° N. The current is 300 km wide and flows at up to 1.5 m/s at depths of 50–300 m. The surface waters in this area flow in the opposite direction.

**Equidae (horses)** (order \*Perissodactyla) Family that includes the modern horses, asses, and zebras (all of which are placed in a single genus *Equus*), and many extinct forms. Sufficient of these are believed to be ancestral to

modern equids for the evolution of the family to have been traced in considerable detail. The horses are first represented in the fossil record by *\*Hyracotherium* ('Eohippus') which diverged from a *\*condylarth* predecessor in the *\*Palaeocene*. Numerous evolutionary lines subsequently appeared from this fox-sized prototype. Most of the evolutionary advances occurred in the New World, although the Equidae were to survive the *\*Pleistocene* only in the Old World. The conquistadores reintroduced horses to the Americas in the 16th century. The domestic horse (*E. caballus*) is probably not descended from the only true living wild horse, Przewalski's horse (*E. ferus przewalskii*), but from a progenitor closely related to it.

**equifinality** The theory that the members of morphologically similar land-forms may each have been produced by a different process or sequence of processes. For example, an armchair-shaped hollow may have been produced by a *\*cirque* glacier, by a rotational landslide, by *\*nivation*, or by spring-sapping. Their present-day similarity may make the origin of such land-forms difficult to determine.

**equilibrium** In geomorphology, a steady state of balance between the processes acting on a landscape and the resisting Earth materials so that over time the geometric shape of the landscape is little changed. Individual land-forms, such as river profiles and hillslopes, may show similar balance. As it is recognized that a perfect balance cannot exist between input and output of energy, the terms 'quasi-' and 'dynamic' equilibrium are sometimes used. Quasi-equilibrium is an apparent equilibrium which is recognized for short periods of time.

**equilibrium flow** See *STEADY FLOW*.

**equilibrium line** Line on a glacier that divides the zone of *\*ablation* (net loss) from the zone of accumulation (net gain).

**equilibrium process (e-process)** In stellar evolution, the culmination of reactions following a rise in temperature after *\*silicon* 'burning'. The e-process rearranges nucleons to produce the most stable nuclei. See *NUCLEOSYNTHESIS*.

**equinoctial gale** Term derived from the popular misconception that gales are more frequent at periods close to the equinoxes.

**equipotential 1. (line)** The two-dimensional locus of all points with equal value for a potential field. **2. (surface)** The three-dimensional distribution of all points of equal value for a potential field. Equipotential lines and surfaces are perpendicular to the lines of force associated with the potential field. For example, sea level is an equipotential surface for the Earth's gravity field.

*Equisetites hemingwayi* The first known species of the family Equisetaceae, found in the \*Carboniferous of Europe, and the direct ancestor of the only extant genus *Equisetum*. See SPHENOPSIDA.

*Equisetum* See SPHENOPSIDA.

**equivalence** A problem of non-uniqueness in the interpretation of \*electrical soundings, when identical trough- or peak-shaped apparent resistivity graphs for a three-layer case can be synthesized using different models for the middle layer. Identical peak-shaped graphs can be generated if the transverse resistance,  $R_t$ , remains a constant equal to  $h\rho$ , where  $h$  is the thickness of the layer and  $\rho$  its true resistivity. Identical trough-shaped graphs can be generated if the longitudinal conductance,  $G_x$ , remains a constant equal to  $h/\rho$ . Such problems can be resolved if access is available to borehole geologic control and the range of  $h$  or  $\rho$  can be deduced.

**equivalent aperture width** The width of a rectangular \*aperture in the \*time domain with the identical peak amplitude, containing the same amount of energy.

*Equus* See EQUIDAE.

**era** First-order geologic time unit composed of several \*periods. The \*Mesozoic era, for example, is composed of the \*Triassic, \*Jurassic, and \*Cretaceous periods. When used formally, as above, the initial letter of the term is often capitalized.

**erathem** The \*chronostratigraphic unit that is equivalent to the \*geologic-time unit \*era. An erathem comprises a number of \*systems grouped together and takes its name from the corresponding era, e.g. the \*Mesozoic Erathem would refer to the rocks laid down during the Mesozoic era. At present it is a little-used term.

**Eratosthenian system** See LUNAR TIMESCALE.

**e-ray** See EXTRAORDINARY RAY.

**ERG** See EXPLORATION OF ENERGIZATION AND RADIATION IN GEOSPACE.

**erg** 1. Sand sea in a hot desert. It is a feature of the Sahara, where sand typically is accumulated in wide shallow basins as *\*alluvial* and *\*lacustrine* deposits derived from adjacent rocky desert during the *\*Cenozoic*. It is often very large: the Grand Erg Oriental covers 196 000 km<sup>2</sup> in Algeria and Tunisia. 2. Unit of energy or work in the *\*c.g.s.* system. 1 erg = 10<sup>-7</sup> J.

**ergodic hypothesis** The proposal that sampling a land-form in space is equivalent to sampling it in time. To be true, this requires that the statistical distribution of objects and events must be the same through space as through time. There are few instances in *\*geomorphology* where this is the case.

**Erian** A *\*series* (391.8–388 Ma ago) in the Middle *\*Devonian* of N. America, underlain by the *\*Ulsterian* and overlain by the *\*Senecan*.

**Eris** The most massive solar system *\*dwarf planet*, discovered in January 2005 by astronomers at the Palomar Observatory, Eris has a diameter of 2326 km and its density is 2.52 ± 0.05 g/cm<sup>3</sup>. It has about 0.27% of the Earth's mass and is 27% more massive than *\*Pluto*, but about the same size. It has one known moon, Dysnomia. Eris has a highly eccentric orbit that brings it to 37.7 *\*astronomical units (AU)* from the *\*Sun* and as far as 97.5 AU. It has an orbital period of 556.6 years. Its *\*albedo* of 0.96 makes it the second brightest object in the solar system after *\*Enceladus*. Infrared spectroscopy reveals the presence of methane ice on its surface.

**EROS** See EARTH REMOTE OBSERVATION SYSTEM.

**Eros** A *\*solar* system asteroid (No. 433), measuring 41 × 15 × 14 km; approximate mass 5 × 10<sup>15</sup> kg; rotational period 5.270 hours; orbital period 1.76 years. After transmitting many photographs, on 2 February 2001 the *\*Near Earth Asteroid* mission crashed on Eros.

**erosion** 1. The part of the overall process of *\*denudation* that includes the physical breaking down, chemical solution and transportation of material. 2. Movement of soil and rock material by agents such as running water, wind,

moving ice, and gravitational **\*creep** (or mass movement). See **BUBNOFF UNIT**.

**erosion rate** The rate at which geomorphological **\*processes** wear away land surfaces. Rates vary widely, depending on both processes and environments, as shown by the following figures: glacial **\*abrasion**, 1000 B (**\*bubnoff** units); soil **\*creep** (under a temperate, maritime climate), 1–5 B; **\*solifluction** (cold climate), 25–250 B; slope wash, 2–200 B; solutional loss, 2–100 B; sea cliff retreat (rocks of medium hardness), 4000 B; soil erosion resulting from human activity, 2000–8000 B.

**erosion surface 1. (planation surface)** Gently undulating land surface that cuts indiscriminately across underlying geologic structures and that is the end-product of a long period of **\*erosion**. Different types are produced under different environments. Among the more important are the **\*peneplain** (humid temperate), **\*etchplain** (humid tropical) and **\*pediplain** (semi-arid). Marine processes may also cut a marine erosion surface. **2.** An irregular surface cut into rock or sediment by the eroding action of flowing water, ice, or by the wind.

**erratic** Glacially transported rock whose lithology shows that it could not have been eroded from the local **\*country rock**. An example is the occurrence near Snowdonia, Wales, of granites derived from Scotland. An ‘indicator’ is an erratic whose origin can be located precisely.

**errors** Generally, the deviation of measured values from their true values. Such errors may be random or systematic. Random errors should have a **\*Gaussian** (normal) distribution about the arithmetic mean of the measurements and this should approach the true value as the number of measurements increases. Systematic errors are consistent differences between the true value and a set of measurements, such that their arithmetic mean is displaced from the true value. Thus random errors determine the precision of a set of measurements, while systematic errors limit their accuracy. The use of the **\*isochron** is a common method for minimizing dating errors.

**eruption** The release of **\*lava** and gas from the Earth’s interior on to the Earth’s surface and into the atmosphere. Where the gas component is minimal (e.g. in basaltic **\*magma**) lava is released quietly; where the gas component is large (e.g. in **\*rhyolite** magma) lava is explosively fragmented

by the expanding gas as it is released, forming high eruption columns of **\*ash** and **\*pumice** above the vent. Such contrasts in the explosivity of eruptions have been used to define several eruption styles, ranging from the quiet Hawaiian type to the violently explosive Phreatoplinian type.

**eruptive centres (volcanic centres)** On **\*Io**, the source of plumes and the deposits they leave on the surface. Centres may be hot, central, volcanic vents, lava lakes, or lava flows that cause explosive sublimation.

**ESA** See **EUROPEAN SPACE AGENCY**.

**escape tectonics** The lateral extrusion of blocks of crust that are bounded by **\*faults** as a result of compression. The northward movement of India into southern Asia is believed to have caused the lateral extrusion of parts of China and the part of southern Asia comprising Myanmar (Burma), Thailand, Laos, Cambodia, Malaysia, and Vietnam.

**escape velocity** The velocity required for atoms or molecules at high altitude to escape from a planet's gravitational field. For example:

Planet	Escape velocity (km/s)
Earth	11.2
Moon	2.4
Mercury	4.3
Venus	10.3
Mars	5.0

**escarpment** See **SCARP**.

**escutcheon** Depressed area, variable in shape, which is found to the posterior of the **\*shell** beaks of certain bivalves. Compare **LUNULE**.

**ESE** See **EARTH SCIENCE ENTERPRISE**.

**esker** Long, sinuous, steep-sided, narrow-crested ridge which consists of cross-bedded sands and gravels. It is laid down by glacial meltwater either at the retreating edge of an ice sheet, or in a subglacial, or **\*englacial** ice tunnel, or in an ice-walled tunnel.

**Eskola, Pentti Elias** (1883–1964) A Finnish geologist and mineralogist, Eskola was a professor at the University of Helsinki from 1928 to 1953. His

major work was a study of the origin of *\*metamorphic rocks*; he developed a classification of *\*metamorphic facies*, based on typical *\*minerals*.

**essential mineral** A *\*primary mineral* whose presence in an *\*igneous* rock is essential to defining the root name of that rock. For example, *\*plagioclase* and the *\*pyroxene* *\*augite* are essential to defining the root name '*\*gabbro*' and hence are essential minerals. Where a primary mineral is present but is not essential to the naming of the rock it is termed an '*\*accessory mineral*'.

**estuary** Semi-enclosed coastal body of water which has a free connection with the open sea and where fresh water, derived from land drainage, is mixed with sea water. Estuaries are often subject to tidal action and where tidal activity is large, ebb and flood tidal currents tend to avoid each other, forming separate channels. In estuaries where tidal activity is small, the invading dense sea water may flow under the lighter fresh water forming a *\*salt wedge*. A positive estuary is one in which surface salinities are lower within the estuary than in the open sea due to freshwater inflow exceeding outflow caused by evaporation. A negative estuary is one in which evaporation exceeds freshwater inflow and therefore hypersaline conditions exist in the estuary. Normally an estuary is the result of valley drowning by the post-glacial rise in sea level. The action of tidal currents on the large amount of available sediment may give rise to a range of mobile bottom forms, including ebb and flood channels, sandbanks, and *\*sand waves*.

**Etalian** A New Zealand *\*stage* (243–232 Ma ago) of the Middle *\*Triassic* *\*epoch*, preceded by the *\*Malakovian* and followed by the *\*Kaihikuan*.

**Etalon** A Russian geodetic (see *GEODESY*), passive satellite mission, dedicated entirely to *\*satellite* laser ranging. The mission comprises two identical satellites, 1.294 m diameter and 1415 kg mass. Etalon-1 was launched on 10 January 1989 and Etalon-2 on 31 May 1989, both from the Baikonur Cosmodrome, Kazakhstan.

**Etched Plains** See *MARTIAN TERRAIN UNITS*.

**etch figures (etch marks)** In *\*crystallography*, pits (etchings), regular in shape, which may develop in certain directions consistent with the orientation of the symmetry elements for the *\*crystal* as a whole when *\*crystal* faces are treated with suitable chemical reagents. These etch

figures may therefore be used to assign crystals to their appropriate **\*crystal systems**.

**etch marks** See ETCH FIGURES.

**etchplain** Plain produced in a tropical or subtropical environment as a result of a phase of deep **\*chemical weathering** during tectonic stability, followed by one of **\*erosion** in which the weathered debris is stripped away. Etchplain relief reflects differences in the resistance of the bed-rock to weathering and involves **\*double planation**.

**etesian winds** Greek name for dry, north- easterly, easterly, northerly, or north-westerly winds which blow between May and October in the Aegean Sea. The equivalent Turkish term is *meltemi*.

**Ethiopian faunal realm** Area which corresponds with sub-Saharan Africa, although it is not completely separated from the neighbouring **\*faunal realms**; generally it is taken to include the south-west corner of the Arabian peninsula.

**ethotype** An organism that is known solely, or principally, by its behaviour.

**eu-** From the Greek *eu* meaning ‘well’ or ‘easily’, a prefix meaning ‘well’, ‘good’, etc. It is used in ecology to denote, in particular, enrichment or abundance, e.g. ‘**\*eutrophic**’, nutrient-rich; ‘euphotic’, light-rich.

**Eubacteria** A former name for the **\*domain \*Bacteria**.

**Eucaryota** See EUKARYOTA.

**eucrete** A type of **\*meteorite** of basaltic composition, mainly pigeonite (low-calcium **\*pyroxene**) and **\*plagioclase** feldspar, with a little metallic iron, **\*troilite**, and one or more **\*silicates**. All eucretes seem to have crystallized at or near the surface of the parent body.

**Euechinoidea** (class **\*Echinoidea**) Subclass of sea-urchins in which the normally rigid **\*test** is composed of five **\*ambulacra** and five **\*interambulacra**, each made up of two columns of plates. The subclass includes both **\*regular** types of echinoid (e.g. *Hemicidaris*) and **\*irregular** types (e.g. *Clypeus*, *Micraster*). They first appeared in the Upper **\*Triassic**.

**eugeocline** An association of **\*calc-alkaline** volcanics, **\*greywackes**, and **\*shales**. These materials are thought to be related to **\*island arcs**. The word

is now little used, unlike the contrasting term ‘\*miogeocline’.

**eugeosyncline** The part of a \*geosyncline that is characterized by the presence of \*volcanism and \*plutonism. Geosynclinal theory has needed some reinterpretation and modification in the light of the newer, unifying theory of \*plate tectonics.

**euhaline water** See HALINITY.

**euohedral (idiomorphic)** A morphological term referring to grains in \*igneous rocks which have a regular crystallographic shape. Euhedral forms are developed when a crystal grows freely in a \*melt and is uninhibited by the presence of any surrounding crystals. The shape of the growing crystals will thus be controlled by their own natural crystallographic form. Compare ANHEDRAL.

**Eukarya** See EUKARYOTA.

**Eukaryota (Eucaryota, Eukarya)** In \*taxonomy, the group that includes all \*eukaryotes. In the five-kingdom classification the Eukarya ranks as a superkingdom, and in the more widely used three-domain classification it is a \*domain that includes the kingdoms \*Protista, \*Animalia, \*Plantae, and \*Fungi.

**eukaryote (adj. eukaryotic)** Organism with cells that have a distinct nucleus, i.e. all \*protists, fungi, plants, and animals. The first eukaryotes were almost certainly green algae (\*Chlorophyta), and what appear to be their microscopic remains appear in \*Precambrian sediments dating from a little less than 1 500 Ma ago. Compare PROKARYOTE.

**Euler angles** See EULER’S ROTATION THEOREM.

**Eulerian current measurement** A technique for measuring the direction and speed of water movement at a series of fixed points. A current-measuring device is held at a fixed point and as the water flows past its speed is measured. A series of measurements taken at different times or places may be plotted on a map as individual current vectors or streamlines. A number of devices exist for Eulerian current measurement, the most common being the propeller-type current meter.

**Euler pole** See POLE OF ROTATION.

**Euler's rotation theorem** A theorem formulated by the Swiss-German mathematician and physicist Leonhard Euler (1707–83) and proved by him in 1775, which states that any displacement of a rigid body in three-dimensional space, such that one point on the body remains fixed, is equivalent to a single rotation of the body about an axis passing through the fixed point, known as the Euler pole or **\*pole of rotation**. This also means that any rotation can be described using three rotation angles, called Euler angles.

**eulite** See ORTHOPYROXENE.

**eulysite** A **\*metamorphic rock** consisting of iron and manganese-bearing **\*silicates**, such as the **\*pyroxene** group **\*minerals** **\*hedenbergite** and iron-rich **\*hypersthene**, the **\*olivine** group minerals fayalite and manganese fayalite, and the **\*garnet** group minerals **\*almandine** and **\*spessartine**.

**EUMETSAT Advanced Retransmission Service (EARS)** A service of the European Organisation for the Exploitation of Meteorological Satellites (EUMETSAT) comprising nine separate data services, each retransmitting observations from an instrument or instrument group, that aim to provide users with regional data from meteorological satellites in **\*polar orbit** through a network of stations that receive, process, and redistribute products in near-real time.



<https://www.eumetsat.int/website/home/Data/RegionalDataServiceEARS/index.html>

- Regional Data Service/EARS.

**Euparkeria** Bipedal **\*thecodont** ('tooth-in-socket') reptile known from the Lower **\*Triassic**. The thecodonts are considered to be ancestors of the **\*dinosaurs** and, apart from a limited number of primitive features in the skull, *Euparkeria* is a likely ancestor for most archosaurian (see ARCHOSAURIA) stocks. It was a small reptile, only 60–100 cm in length.

**euphotic zone** See PRIMARY PRODUCTIVITY.

**Euramerica** The continental mass which resulted from the fusion of north-western Europe and N. America during the **\*Caledonian orogeny**. This

cratonic (see [CRATON](#)) area subsequently fused with [\\*Angara](#) and [\\*Gondwana](#) during the Variscan orogenic event to form [\\*Pangaea](#).

**Eurasian Plate** One of the present-day major lithospheric [\\*plates](#), extending from the oceanic [\\*ridges](#) in the [\\*Atlantic Ocean](#) to a currently poorly-identified margin with the [\\*Pacific Plate](#) in the east. The southern boundary is basically the Alpine–Himalayan [\\*fold belt](#), involving a collage of [\\*microplates](#), minor plates, and the [\\*African](#) and [\\*Indo-Australian](#) Plates.

**Europa (Jupiter II)** The smallest of the [\\*Galilean](#) satellites, and the smoothest object in the solar system, with no feature more than 1 km high. The surface is icy, the ice being about 10–30 km thick, with two types of terrain, one mottled, brown or grey, with small hills, the other comprising large, smooth plains criss-crossed with straight and curved tracks, some thousands of kilometres long, producing a surface resembling that of the Arctic Ocean but with [\\*penitentes](#). There are very few craters. The crust is believed to be no more than 150 km thick and is believed to include a liquid ocean beneath the surface ice. The inner core is believed to be of iron and sulphur beneath a rocky mantle. The satellite was discovered on 7 January 1610, by Galileo. Its diameter is 3130 km; mass  $4.8 \times 10^{22}$  kg; density 2990 kg/m<sup>3</sup>; visual albedo 0.64; surface gravity 0.135 (Earth = 1); mean distance from Jupiter 670 900 km; mean distance from Sun 5.203 AU; orbital period 3.551181 days; rotational period 3.551181 days.

**European Federation of Geologists** A body that represents professional geologists within the European Union. It promotes the movement of geologists within the EU by encouraging mutual recognition of qualifications and the harmonization of educational standards. It regulates degrees, diplomas, and titles, has established a common code of professional conduct, and seeks to develop common policies regarding public issues of interest to geologists. The Federation was formally launched in 1980 in Paris, during the 26th International Geological Congress.

**European Geosciences Union** A body founded in 2002 through the merger of the European Geophysical Society and the European Union of Geosciences with the aim of promoting the Earth and planetary sciences and cooperation between scientists.


**European GeoTraverse (EGT)** A geological survey of the European continent along a traverse from the North Cape, Norway, to Tunisia with the aim of producing a three-dimensional description of the evolution, nature, structure, physical properties, and dynamics of the continent. The project was proposed in 1980, involved scientists from fourteen countries, and was completed in 1992.

**European Province** See ATLANTIC PROVINCE.

**European Space Agency (ESA)** An organization, founded in May 1975 to replace the European Space Research Organization, that conducts space research and operates its own launch vehicles.

**europium anomaly** Most *\*rare-earth* elements exist in the trivalent state. Europium, however, can exist as the  $\text{Eu}^{2+}$  *\*ion* and it can proxy for calcium in *\*plagioclase* feldspar during *\*igneous* fractionation. Crystallization of calcic plagioclase thus depletes residual *\*magmas* of europium, relative to the other rare-earth elements. The europium anomaly can therefore be used as an indication of the amount of fractionation a magma has undergone. Efforts to explain the marked europium anomaly discovered in *\*lunar \*basalts* led to much debate about the geochemistry of the *\*Moon*, with suggestions that the europium might have been lost by volatilization, retained selectively in the lunar interior, or, more probably, that melting and differentiation caused it to be depleted in the basalts but retained in the highlands.

**Euryapsida** Reptiles which have a single, upper temporal opening behind the eye. The group includes the *\*plesiosaurs*, *\*nothosaurs*, and the *\*placodonts*. The *\*ichthyosaurs* had a similarly placed opening, but there were key differences in the arrangement of the bones forming its margins: for this and other reasons, the ichthyosaurs are usually placed in a different subclass from the euryapsids, the Ichthyopterygia. However, like the ichthyosaurs, the euryapsids were typically aquatic types, especially from the mid-*\*Triassic* onwards. Euryapsids first appeared in the *\*Permian* and became extinct at the end of the *\*Mesozoic*.



## Euryapsida

**Euryarchaeota** (domain *\*Archaea*) The more derived (see *APOMORPH*) of the two kingdoms of Archaea, comprising a broad range of *\*phenotypes* including *\*methanogens*, *\*halophiles*, and *\*sulphur-reducing* organisms. Members of the Euryarchaeota show less genetic similarity to those belonging to the *\*domains* *\*Eukaryota* and *\*Eubacteria* than do those of the *\*Crenarchaeota*.

**euryhaline** Able to tolerate a wide range of *\*salinity*.

**eurypterid** See *CHELICERATA*; *MEROSTOMATA*.

**eurythermal** Able to tolerate a wide range of temperature.

**eurytopic** Able to tolerate a wide range of several factors.

**eustasy** The worldwide changing of sea level caused either by tectonic movements, or by the growth or decay of glaciers (*\*glacio-eustasy*, or glacio-eustatism).

**eutaxitic structure** A planar *\*fabric* found in welded *\*ignimbrites*, defined by flattened and elongate *\*pumice* *\*clasts* (*\*fiamme*), usually 10–40 mm in length and set in a lighter matrix of flattened and sintered, *\*ash-size*, glassy shards.

**eutectic point** See *EUTECTIC SYSTEM*.

**eutectic system** Mixture of two or more minerals in definite proportions which have crystallized from a *\*melt* or solution simultaneously. The

temperature at which this occurs is the eutectic point.

**Eutelsat 172B** A communications satellite, built by Airbus Defence and Space, that relies exclusively on electric propulsion to raise its initial orbit and for all manoeuvres needed to keep it on station. It was launched on 1 June 2017, from Kourou, French Guiana.

**Eutheria** (class *\*Mammalia*, subclass Theria) The infraclass that includes all of the placental mammals, and which probably arose during the *\*Cretaceous*. The embryo is retained in the uterus, nourished by means of an allantoic placenta, and born in an advanced stage of development.

**Eutracheophytic** An epoch in the evolution of plants, proposed in 1993 by J. Gray, that lasted from the late *\*Lochkovian* (about 411 Ma ago) in the Early *\*Devonian* until the middle *\*Permian* (about 265 Ma ago). During this time the diversity of spores and plant megafossils increases greatly, with several classic assemblages, and a marked increase in the diversity of vascular plants.

**eutrophic** Applied to nutrient-rich waters with high *\*primary productivity*. Compare OLIGOTROPHIC.

**euxinia** The state of being *\*euxinic*.

**euxinic** Applied to an environment in which the circulation of water is restricted, leading to reduced oxygen levels or anaerobic conditions in the water. Such conditions may develop in swamps, *\*barred* basins, stratified lakes, and *\*fjords*. Euxinic sediments are those deposited in such conditions, and are usually black and organic-rich. Compare ANOXIC.

**EV (ExactView)** A Canadian commercial service used by marine traffic that monitors ship movements, especially through busy shipping lanes and harbours. The EV-1 *\*microsatellite* was launched on 22 July 2012, from the Baikonur Cosmodrome, Kazakhstan; the EV-9 *\*nanosatellite* was launched on 28 September 2015, from the Satish Dhawan Space Centre, India.

**evaporation** See EVAPOTRANSPIRATION.

**evaporation pan** A broad, shallow, water-filled pan of standard size. For example, the class A pan used by the US Weather Bureau is 122 cm in diameter and 25 cm deep. The amount of water within the pan is monitored to obtain an estimate of evaporation losses.

**evaporimeter** See [LYSIMETER](#).

**evaporite** General lithologic term applied to [\\*sedimentary rocks](#) which have formed by the precipitation of salts from natural brines during evaporation of marginal salt pans, [\\*lagoons](#), supratidal flats, and saline lakes. Common rock types are [\\*limestone](#), [\\*dolomite](#), [\\*anhydrite](#), [\\*gypsum](#), and [\\*rock salt](#).

**evapotranspiration** Combined term for water lost as vapour from a soil or open water surface (evaporation) and water lost from the surface of a plant, mainly via the stomata (transpiration). The combined term is often used since in practice it is very difficult to distinguish water vapour from these two sources in water-balance and atmospheric studies.

**evapotron** An instrument developed in Australia to measure the extent and direction of vertical air eddies which are involved in the vertical transfer of water vapour, and thus to provide a direct measurement of evaporation rates over short periods of time.

**event deposit** See [STORM BED](#).

**event stratigraphy** A term first proposed by D. V. Ager (1973) for the recognition, study, and [\\*correlation](#) of the effects of significant physical events (e.g. marine [\\*transgressions](#), volcanic [\\*eruptions](#), geomagnetic [\\*polarity reversals](#), climatic changes), or biological events (e.g. extinctions), on the stratigraphic record of whole continents, or even of the entire globe. It is argued that by correlating these effects, as they are evidenced in the sedimentary record, it will be possible to define truly synchronous horizons, thus leading to greater resolution and a more accurate [\\*chronostratigraphic scale](#). More recently A. Seilacher (1984) has suggested the term 'event stratigraphy' for the study of events at the level of individual [\\*beds](#).

**event stratigraphy** See [EVENT STRATIGRAPHY](#).

**evolute** Applied to coiled [\\*cephalopod](#) conchs, in which all the whorls are exposed. Compare [INVOLUTE](#).

**evolution** Change, with continuity in successive generations of organisms. The phenomenon is amply demonstrated by the fossil record, for the changes over geologic time are sufficient to recognize distinct [\\*eras](#), for the

most part with very different plants and animals. *See also* DARWIN; MACROEVOLUTION; MICROEVOLUTION; NATURAL SELECTION; PHYLETIC EVOLUTION; PHYLETIC GRADUALISM; PHYLOGENY; PUNCTUATED EQUILIBRIUM.

**evolutionary lineage** Line of descent of a \*taxon from its ancestral taxon. A lineage ultimately extends back through the various taxonomic levels, from the species to the genus, from the genus to the family, from the family to the order, etc. *See also* CLASSIFICATION.

**evolutionary rate** Amount of evolutionary change that occurs in a given unit of time. This is often difficult to determine, for several reasons. For example, should the unit of time be geologic or biological (the number of generations)? How should morphological change in unrelated groups be compared? In practice it is necessary to adopt a pragmatic approach, such as the number of new genera per million years.

**evolutionary species** *See* CHRONOSPECIES.

**evolutionary trend** Steady change in a given adaptive direction, either in an evolutionary lineage or in a particular attribute, e.g. dentition. Such trends are often apparent in unrelated \*taxa. Formerly they were attributed to \*orthogenesis; now \*orthoselection or the contending theory of species selection are invoked.

**evolutionary zone** *See* LINEAGE-ZONE.

**evolved magma** A basaltic \*magma with a low \*magnesium number (Mg#) and low content of MgO (magnesium oxide). Such magma results from crystals separating from a melt in which the liquid in equilibrium with the solid has a lower Mg# than the solid, so as \*olivine is removed the Mg# falls and since olivine at the forsterite ( $\text{Mg}_2\text{SiO}_4$ ) end of the series is rich in MgO, the level of MgO will fall also.

**Ewing, Maurice** (1906–74) An American geophysicist and oceanographer, Ewing developed offshore \*seismic reflection \*profiling for use in oil prospecting in the 1930s. In the post-war period he made extensive studies of the structure of the floor of the Atlantic ocean, using \*seismic refraction, sediment cores, etc. Ewing was instrumental in making the Lamont–Doherty Geological Observatory a leading research centre.

**ex-** From the Latin *ex* meaning ‘out of’, a prefix meaning ‘out’ or ‘not having’.

**ExactView** See *EV*.

**exaerobic** Applied to a laminated *\*biofacies* formed in a depositional environment that was *\*anaerobic*, or almost so, and that contains epibenthic (see *EPIBENTHOS*) fossils of macroinvertebrates (e.g. *\*Brachiopoda*, *\*Bivalvia*, *\*Mollusca*).

**exaptation** A characteristic that opens up a previously unavailable niche to its possessor. The characteristic may have originated as an *\*adaptation* to some other niche (e.g. it is proposed that feathers were an adaptation to thermoregulation, but opened up the possibility of flight to their possessors), or as a neutral mutation.

**excavation** Hole created in the ground by *\*drilling*, augering (see *AUGER*), boring, blasting, scraping, ripping, or digging, depending on the strength and condition of the rock requiring removal. Excavation may be on the surface, e.g. for buildings, or underground, e.g. for mines and tunnels.

**exchangeable ions** Charged ions that are adsorbed on to sites (with a charge opposite to that on the ion) on the surface of the *\*adsorption complex* of the soil (mainly *\*clay* and *\*humus \*colloids*). Exchangeable ions can replace each other on this surface, and are also available to plants as nutrients. Although *\*cations* (e.g. calcium and magnesium) are the most common, exchanging at negatively-charged sites, some complexes (e.g. sulphate and phosphate) do exchange at positively-charged sites. See also *ANION-EXCHANGE CAPACITY*; *CATION-EXCHANGE CAPACITY*; *EXCHANGE CAPACITY*.

**exchange capacity** Total ionic charge of the *\*adsorption complex* in the soil that is capable of adsorbing *\*cations* or *\*anions*.

**exchange pool** See *ACTIVE POOL*.

**exfoliation** Weakening and separation of the surface layers of rock as a result of *\*chemical* or (possibly) thermal *\*weathering*, or of pressure release due to *\*erosion*. The decomposition of *\*biotite* and hydration of *\*feldspar* in *\*granite* causes swelling that may lead to failure. Expansion and rock fracturing may also result from temperature change (although this is

questioned by many geologists), and sheet failure may result from the release of internal stress in massive rocks when an overburden is removed. *See also* THERMOCLASTIS.

**exhumed topography** An ancient land-form or landscape that had been buried beneath younger rocks or sediment and that is exposed by their subsequent \*erosion.

**exichnia** *See* TRACE FOSSIL.

**exine** Outer, decay-resistant coat of a \*pollen grain or \*spore, composed of sporopollenin, an inert polymer. The exine is characteristic for different plant families and genera, and sometimes even for different species. Hence it forms the basis for the identification and quantitative analysis of the vegetation composition of \*peats and other suitable sedimentary deposits dating back many thousands of years. *See also* PALYNOLOGY.

**exinite** *See* COAL-MACERAL GROUP.

**exitance** The radiant flux density of \*electromagnetic radiation leaving a surface.

**exobiology** The biology of outer space: a study currently limited to the seeking of evidence for the existence of life beyond the Earth, and speculation on the possible alternative forms of such life.

**ExoCube** A 3-unit \*CubeSat developed by the California Polytechnic State University and sponsored by the National Science Foundation that measures the density of hydrogen, oxygen, helium, and nitrogen in the Earth's lower \*exosphere and upper \*ionosphere. It was launched on 31 January 2015, from California.

**exocuticle** *See* EXOSKELETON.

**exogenetic processes** A blanket term for those processes which operate on or close to the surface of the Earth and which involve \*weathering, \*mass movement, \*fluvial, \*aeolian, \*glacial, \*periglacial, and \*coastal processes. The term is normally used in contrast to the \*endogenetic processes, whose origin is within the Earth.

**exogenous dome** *See* DOME (2).

**exorheic lake** A lake that has one or more outflow streams. *Compare* ENDORHEIC LAKE.

**exoskeleton** A general term applied to the hard covering of many *\*invertebrates*. It may be a shell growing at the edges only (by accretion) or a series of plates. The term is most commonly applied to the horny skeleton enclosing the body of all *\*Arthropoda* and secreted by the underlying cellular layer. The exoskeletal material (*\*cuticle*) is composed of a complex glycoprotein, is relatively impermeable to water, and has a high strength-to-weight ratio. It provides insertion sites for muscles (apodemes) and is divided into separate plates that facilitate movements. The plates are connected by thin, untanned cuticle. In trilobites (*\*Trilobita*) and crustaceans (*\*Crustacea*) the cuticle is impregnated with mineral salts (calcium carbonate and calcium phosphate) which give increased strength. The exoskeleton is moulted periodically (*\*ecdysis*) to permit body growth.

**exosphere** Outer region of the upper atmosphere extending from a base of 500–750 km altitude. The zone has a very low concentration of gases (mostly atoms of oxygen, hydrogen, and helium, of which about 1% are ionized). Gases can escape from it into space, as molecular collisions are much reduced because of the low gas density. The exosphere and much of the underlying *\*ionosphere* form part of the *\*magnetosphere*.

**exothecal** External to the *\*theca*, or *\*test*.

**exotherm** *See* POIKILOTHERM.

**exotic** Applied to geological materials derived from an extrabasinal source, e.g. exotic *\*clasts* in a *\*conglomerate* derived from some distant or extraformational source, or an exotic *\*terrane* (i.e. a structurally emplaced *\*allochthonous* unit).

**expanding Earth** A hypothesis which was strongly supported by Warren Carey (1911–2002), but which was first proposed by M. R. Mantovani in 1907, and raised again in the 1930s by Hilgenbirg and others. It holds that the diameter of the Earth has increased with time, fragmenting the continents and causing the growth of ocean basins at spreading axes (ridges). A conference to discuss the idea was hosted by Carey at Hobart, Tasmania, in 1956.

**expanding spread** In seismic refraction shooting, a method in which a \*spread of \*geophones is used at increased \*offset for repeated shots at the same location. This is equivalent to using many spreads of geophones for a single shot. In seismic-reflection surveys it is used to provide information about \*root-mean-square velocities and depths to reflectors using the  $t^2-x^2$  method. *See also* ELECTRICAL SOUNDING.

**Experimental Geodetic Satellite (EGS)** A passive satellite developed by the \*Japan Aerospace Exploration Agency (JAXA) and Kawasaki Heavy Industries that is used in long-range \*geodetic measurements aimed at rectifying Japan's geodetic triangular network. The satellite is a hollow sphere, 2.15 m diameter and 685.2 kg mass, covered in 318 mirrors and 120 laser-reflective surfaces. It was launched on 12 August 1986, from Tanegashima Space Center, Japan.

**Exploration of energization and Radiation in Geospace (ERG)** A \*minisatellite mission of the \*Japan Aerospace Exploration Agency (JAXA) to study the acceleration and loss mechanisms of relativistic particles in the inner \*magnetosphere during space storms. The satellite was launched on 20 December 2016, from Uchinoura Space Center, Japan.

**Explorer-1 PRIME-2** *See* HRBE CUBESAT MISSION.

**Explorer 59** *See* INTERNATIONAL SUN-EARTH EXPLORER-C.

**Explorer Ridge** A \*mid-ocean ridge at a \*divergent plate boundary about 241 km west of Vancouver Island, Canada. It comprises one major segment, the Southern Explorer Ridge, and includes a deep \*rift valley.

**explosive charge** *See* CHARGE.

**exposure age (sun-tan age)** The period during which a rock has been exposed at the lunar surface as measured by fission tracks induced by exposure to \*solar flare particles, which penetrate to depths of less than 0.5 cm. The average exposure age is less than three million years.

**exsiccation** Dehydration of an area by a process, e.g. drainage, in the absence of changes in precipitation levels. Draining of marshlands and deforestation are examples of processes that can lead to exsiccation. *See also* DESICCATION.

**exsolution** Unmixing. Some homogeneous **\*solid solutions** of **\*minerals** are stable only at high temperatures. On cooling these become unstable and one mineral separates from the other at a certain temperature. An intergrowth of two separate minerals may result, e.g. **\*perthite**, an intergrowth of sodium and potassium feldspar. This occurs without loss or addition to the mineral as a whole.

**extension** A measure of the change in length of a line from its initial unit length. Extension may be positive (elongation) or negative (shortening), depending on whether the length of the line increases or decreases. The simplest type of extension ( $e$ ) is calculated from:  $e = (L_f - L_0)/L_0$ , where  $L_f$  is the final length and  $L_0$  the original unit length.

**external mould** See FOSSILIZATION.

**extinction** 1. In optical mineralogy, a **\*mineral** is said to be in extinction when the **\*vibration** direction of the two rays of a doubly refracting **\*crystal** coincide with the vibration directions of the two pieces of **\*Polaroid** in a thin-section microscope that is parallel to the **\*polarizer** and **\*analyser** so that no light reaches the eye. This phenomenon occurs four times in a complete 360° rotation of the stage. See OBLIQUE EXTINCTION; STRAIGHT EXTINCTION; SYMMETRICAL EXTINCTION; UNDULOSE EXTINCTION. 2. The elimination of a **\*taxon**. This may take place in several ways. In the simplest case the taxon disappears from the record and is not replaced. Alternatively, one taxon may replace another, the earlier group consequently disappearing. Thus there is a process of either subtraction or substitution. Extinction generally takes place at particular times and places but there are recurring periods when episodes of mass extinction have taken place. Environmental catastrophe, occurring for whatever reason, removes many groups from the environment and ecosystems collapse. Eventually new forms appear and evolution resumes. It would appear that periods of mass extinction control the pattern of evolution.

**extraclast** A fragment of **\*carbonate** rock derived from the erosion of an exposed ancient **\*limestone** on land outside the depositional basin in which it is found. Compare INTRACLAST.

**extraction** See ABSTRACTION.

**extraformational** See CONGLOMERATE.

**extraordinary ray (e-ray)** In mineral optics, one of the two rays produced when light is passed through a doubly refracting **\*crystal**. The ordinary ray (o-ray) travels with uniform velocity in all directions; the velocity of the extraordinary ray varies with direction. The extraordinary ray is refracted within the **\*mineral**.

**extra-orogen basin** A **\*drainage** basin located along the edge of a mountain belt in which water drains parallel to the **\*strike**.

**extremophile** A micro-organism (domain **\*Archaea**) that thrives under extreme environmental conditions of temperature, **\*pH**, or salinity. *See also* ACIDOPHILE; ALKALIPHILE; HALOPHILE; HYPERTHERMOPHILE; PSYCHROPHILE; THERMOPHILE.


**extrusion** The emission of **\*magma** from a vent or **\*fissure** on to the Earth's surface where it forms a **\*lava** flow. Gas is released quietly, rather than explosively, from the magma at the source vent and consequently there is little associated **\*pyroclastic** activity. Gas may also be vented from a different source, or extrusion may be of lava (magma) which has been degassed previously.

**extrusive** Applied to all ejected material of volcanic origin. The sense applies to **\*lavas** and flows, rather than to **\*pyroclastic** rocks.

**'eye' of storm** Central part of a tropical cyclone with light winds, generally clear skies, and a slight, horizontal pressure gradient. The diameter of the 'eye' averages 20 km but in a large cyclone can be 40 km or more. The 'eye' is an area of some air subsidence which produces adiabatic warming.

**eyepiece (ocular)** An eye lens, a fixed diaphragm with **\*cross-wires**, and a field lens, all contained in a short tube which is inserted into the top of a microscope. The internal construction of the eyepiece may vary depending on whether the focal plane lies above (Huygenian or negative eyepiece) or below (Ramsden or positive eyepiece) the field lens. Most eyepieces are either 5× or 10× magnification.

**eyot** See BRAIDED RIVER.



**fabric (petrofabric)** The physical arrangement of particles and \*minerals in a rock, including its texture and structure, both microscopic and macroscopic.

**fabric analysis** Analysis of the elements that make up the \*fabric of a rock to determine the response of that rock to \*stress. In a rock the three-dimensional pattern comprising the distribution, shape, size, and size distribution of \*crystals or \*grains constitutes the fabric, e.g. bedding in \*sedimentary rocks, or metamorphic banding (see GNEISS). Fabric analysis is important in engineering because, for example, rock strength parallel to banding can differ markedly from that perpendicular to banding. In soils, fabric is analysed by comparing the strength of an undisturbed sample with its strength when \*remoulded; this describes the \*sensitivity of the soil.

***Fabrosaurus australis*** Described in 1964 by Ginsburg, *F. australis* is one of the earliest recorded \*ornithischian \*dinosaurs. Like other 'bird-hipped' dinosaurs, *Fabrosaurus* possessed a predentary bone at the tip of its lower jaw. Its teeth were small and pointed, and were used to grind plant material. *Fabrosaurus* was a biped with short forelimbs, and a long tail to provide good balance.

**face pole** See POLE OF A FACE.

**facial suture** See CEPHALIC SUTURE.

**facies** 1. Sum total of features that reflect the specific environmental conditions under which a given rock was formed or deposited. The features may be lithologic, sedimentological, or faunal. In a sedimentary facies, \*mineral composition, \*sedimentary structures, and bedding characteristics

are all diagnostic of a specific rock or lithofacies. 2. See [METAMORPHIC FACIES](#).

**facies association** A group of sedimentary [\\*facies](#) used to define a particular sedimentary environment. For example, all the facies found in a [\\*fluvatile](#) environment may be grouped together to define a fluvial facies association.

**facies fossils** [\\*Fossil](#) organisms that are restricted to particular [\\*lithologies](#), reflecting the original environments of deposition.

**facies sequence** A vertical succession of [\\*facies](#); such sequences often fine or coarsen upwards, and may be repeated many times in a cyclic manner due to the migration of the facies through time and space.

**facing direction (younging)** The direction (upward or downward) in which the stratigraphy throughout most of a [\\*fold](#) becomes progressively younger. [\\*Anticlines](#) face upwards, by definition; [\\*antiforms](#) may face upwards or downwards depending on the direction in which the stratigraphy is younging. It may be determined by such 'way-up' criteria as [\\*graded bedding](#), [\\*sole structures](#), and [\\*cross-bedding](#).

**facula** (*pl. faculae*) A bright area on the surface of an extraterrestrial body.

**faculae** See [FACULA](#).

**faecal pellet** A rounded, initially soft particle, usually 100–500 µm in diameter, which has been excreted by an organism. The internal structure of the pellet is usually fine grained. Worms ([\\*Annelida](#)), [\\*gastropods](#), and [\\*crustaceans](#) produce these pellets in large quantities. Faecal pellets are most likely to accumulate and be preserved in low-energy, muddy environments colonized by an abundant fauna, e.g. [\\*lagoons](#) and [\\*tidal flats](#). As a type of [\\*fossil](#) excreta, the term is commonly applied to small droppings, often of invertebrate origin, which may make up important parts of some [\\*lithologies](#). See [COPROLITE](#).

**fahlband** A layer or lens in a [\\*metamorphic rock](#) that contains metal sulphides. The word is German: *fahl* means fawn or dun-coloured.

**failed arm** See [AULACOGEN](#); [RIFT VALLEY](#).

**failed rift** See [AULACOGEN](#); [RIFT VALLEY](#).

**failure** The process by which a body under *\*stress* loses cohesion and divides into two or more parts, commonly by means of a brittle fracture.

**failure strength** See *ULTIMATE STRENGTH*.

**failure stress envelope** On a *\*Mohr* stress diagram, the curve which joins points of *\*failure* at progressive *\*stress* configurations, thus delimiting the field of stable stress configurations (within the envelope) as distinct from the failure field outside the envelope.

**fairweather wave-base** The sea depth, typically 5–15 m, above which the sea bed is affected by the action of waves that occur during calm weather. During calm weather, material eroded from the beach is deposited as a smooth layer covering the material laid down during the most recent storm. Compare *STORM WAVE-BASE*.

**falling dune** An irregular *\*sand \*dune* that forms on the downwind side of a large obstruction, in the area where the airflow is disrupted.

**falling head permeameter** See *PERMEAMETER*.

**falling stage systems tract** See *REGRESSIVE SYSTEMS TRACT*.

**falls** *\*Meteorites* that are seen to fall or are collected immediately, and whose time and locality of impact are accurately recorded. Compare *FINDS*.

**fall-stripes** See *VIRGA*.

**false body** *\*Clay* displaying *\*thixotropy*.

**false cirrus** See *SPISSATUS*.

**false colour** A term used in *\*remote* sensing techniques to describe the display of data collected in a number of different wavelengths, usually longer or shorter than those perceptible to the naked eye. Typical false-colour images include infrared data, which are often displayed as visible red. Thus green vegetation, which is highly reflective in the infrared, typically appears red on a false-colour image.

**famatinite** See *ENARGITE*.

**Famennian 1.** The final *\*age* in the *\*Devonian* period, preceded by the *\*Frasnian*, followed by the Hastarian age (*\*Carboniferous*), and dated at

374.5–359.2 Ma ago (Int. Commission on Stratigraphy, 2004). **2.** The name of the corresponding European *\*stage*, which is roughly contemporaneous with the upper *\*Hervyan* (Australia) and the *\*Chautauquan* (N. America).

**family** See CLASSIFICATION.

**FAMOUS project** The Franco-American Mid-Ocean Undersea Study, undertaken on a 50 km length of the Mid-Atlantic Ridge between latitudes 36.5 ° N and 37 ° N.

**fan cleavage** The structural arrangement in which *\*cleavage* planes form upwardly convergent or divergent fans throughout a sequence of folded layers. Generally, cleavage planes are parallel to the *\*axial plane* only in the *\*hinge* region of a *\*fold*; elsewhere the cleavage planes may deviate systematically from a parallel alignment.

**fanglomerate** Applied generally to *\*conglomerates* and *\*breccias* deposited on *\*alluvial fans*.

**fan shooting** A simple *\*seismic* refraction method used for delineating subsurface geologic features (e.g. *\*salt* domes, buried valleys, and back-filled mineshafts) by contrasting their *\*seismic* velocities with those of the surrounding materials. *\*Geophones* are set out around a segment of arc in a fan-like *\*array* centred on one or more shot locations. A base line of geophones is used in relation to one of the shot locations to provide a time–distance curve where no subsurface feature is present, thus calibrating the travel times for a given range. Using the fan arrays, travel times of refracted rays are measured to each detector; any ray encountering a zone of anomalously high or low velocity will arrive ahead of or behind the expected travel time for that shot-to-geophone range.

**FAO** The Food and Agriculture Organization of the United Nations.

**Farallon Plate** A present-day minor lithospheric *\*plate* subducting beneath the *\*North* American Plate, the Farallon Plate is the remnant of a plate that was once large. The amount of Farallon *\*crust* which has been subducted can be estimated from the area of Pacific crust which still exists, assuming symmetrical spreading from the Pacific–Farallon *\*constructive* margin. The present-day remnants are also called the *\*Gorda*, *\*Juan de Fuca*, and *\*Cocos* Plates.

**far-field barrier (geological barrier)** In the disposal of *\*radioactive* waste, a structure with geological and hydrological characteristics which make it impermeable to radionuclides and thus ensure that it will provide permanent containment. *Compare* [NEAR-FIELD BARRIER](#).

**farra** *See* [FARRUM](#).

**farrum** (*pl. farra*) On the surface of *\*Venus*, a feature shaped like a pancake or a row of pancakes.

**fasciculate** Applied to a *\*compound* coral in which the *\*corallites* are spaced far enough apart to avoid mutual interference during growth. They may be linked by connecting processes.

**fasciole** A groove on the *\*test* of spatangoid echinoids (*\*Echinoidea*) which does not bear large spines or tubercles. Tiny spines situated in these grooves are covered in cilia (*see* [CILIUM](#)) that move water and mucus so as to remove extraneous material from the surface of the animal.

**fast breeder reactor** Nuclear reactor which uses fast neutrons to convert uranium to plutonium and which creates more fuel than it uses. A chain reaction is set up in which  $^{238}\text{U}$ , with an initial charge of  $^{239}\text{Pu}$  to begin the *\*fission* process, discards fast neutrons each of which induces fission in another nucleus. Excess  $^{239}\text{Pu}$  is produced where plutonium production exceeds rate of fission. About 60% of fuel elements in fast breeder reactors is converted to useful energy compared to 0.5–1% in *\*burner* reactors. Plutonium can be recycled, but other radioactive products are waste. *See* [RADIOACTIVE WASTE](#).

**fast Fourier transform (FFT)** An algorithm (e.g. the Cooley–Tukey method) which enables the Fourier transformation of digitized wave-forms to be accomplished more rapidly by computer than would be possible using direct evaluation of the Fourier integral. FFT usually involves iterative techniques. *See also* [FOURIER ANALYSIS](#); [FOURIER TRANSFORM](#).

**Fast On-Orbit Recording of Transient Events (FORTE)** A US satellite sponsored by the Department of Energy and designed and built by the Los Alamos National Laboratory that measures electromagnetic pulses due primarily to lightning. It was launched on 29 August 1997, from California.

**fathom** Unit of water-depth measurement, originally six feet, equal to 1.83 m.

**fatty acid** Long-chained, predominantly unbranched, carboxylic acid; it may be saturated or unsaturated. Fatty acids have the general formula  $R-(CH_2)_n-COOH$ , where R represents a hydrocarbon group, e.g.  $CH_3$  or  $C_2H_5$  and  $n$  is any whole number between 1 and 16.

**fault** Approximately plane surface of fracture in a rock body, caused by brittle failure, and along which observable relative displacement has occurred between adjacent blocks. Most faults may be broadly classified according to the direction of slip of adjacent blocks into *\*dip-slip*, *\*strike-slip*, and *\*oblique-slip* varieties. The term 'dip-slip fault' comprises both *\*normal* and *\*reverse* slip faults, and the special cases of low-angle *\*lag* and *\*thrust* faults. Strike-slip faults (wrench, transform, transcurrent) result from horizontal displacement (dextral or sinistral movements), and on a regional scale may involve *\*transpression* and *\*transtension*. *See also VOIDS.*

**fault block (fault slice)** A rock mass which is bound on at least two sides by *\*fault* planes. The block may be uplifted or depressed in relation to adjacent blocks.

**fault-block mountains** Mountains or ranges that result from the upthrow of large fault blocks and that are separated from others by basins or troughs, producing an upland unit bounded by normal or reversed faults. It is classically developed in the Great Basin, Utah, USA, where a typical block is tilted and bounded by a steep fault scarp on one side and by a more gentle dip slope on the other. Usually it is dissected by erosion. *See BASIN-AND-RANGE PROVINCE; HORST.*

**fault line** *See FAULT TRACE.*

**fault-line scarp** *See SCARP.*

**fault outcrop** *See FAULT TRACE.*

**fault plane** A discrete, planar surface along which there has been appreciable relative displacement of the rock masses on either side.

**fault-plane solution** The use of the direction of first motion of seismic waves detected at seismic stations in different areas to determine the nature and orientation of the *\*stress field* involved in the initial generation of an *\*earthquake* and hence used to determine the *\*focal* mechanism.

**fault scarp** See SCARP.

**fault slice** See FAULT BLOCK.

**fault trace (fault line, fault outcrop)** A generally linear feature which marks the intersection of a *\*fault* plane with the surface of the Earth. Fault traces are sometimes marked by positive or negative topography and the emergence of *\*springs*.

**fault trap** Structure in which water, oil, or gas may be trapped on one side of a *\*fault* plane by an impervious horizon thrown above it by a *\*fault*. Compare ANTICLINAL TRAP; REEF TRAP; STRATIGRAPHIC TRAP; STRUCTURAL TRAP; UNCONFORMITY TRAP.

**fault zone** A region, from metres to kilometres in width, which is bounded by major *\*faults* within which subordinate faults may be arranged variably or systematically. Single fault zones are marked by fault *\*gouge*, *\*breccias*, or *\*mylonites*.

**fauna** (*adj.* faunal, faunistic) The animal life of a region or geologic period. Compare FLORA.

**faunal province** See FAUNAL REALM.

**faunal realm (faunal province, faunal region, zoogeographical region)** Biological division of the Earth's surface (i.e. a large geographical area) containing a fauna more or less peculiar to it. The degree of distinctiveness varies with the region concerned and reflects partly climate and partly the existence of barriers to migration. The number of realms recognized varies from one authority to another, but a minimum of six are recognized: *\*Australian*, *\*Ethiopian*, *\*Nearctic*, *\*Neotropical*, *\*Oriental*, and *\*Palaeartic*.

**faunal succession** See LAW OF FAUNAL SUCCESSION.


**faunizone** See ASSEMBLAGE ZONE.

**fayalite** See OLIVINE.


**Faye correction** See FREE-AIR CORRECTION.

**FDSN** See INTERNATIONAL FEDERATION OF DIGITAL SEISMIC NETWORKS.

**Feammox** A process by which anaerobic bacteria oxidize ammonium (NH<sub>4</sub>) that is coupled to ferric iron (Fe(III)), which is reduced, to produce dinitrogen (N<sub>2</sub>) and nitrite (NO<sub>2</sub>); the NO<sub>2</sub> may subsequently be reduced to N<sub>2</sub>, nitrate (NO<sub>3</sub>), or NH<sub>4</sub>:



or, at pH below 6.5:



The process occurs in tropical soils, where it shortcuts the nitrogen cycle, leading to the loss of nitrogen from ecosystems. See also ANAMMOX.

**feather angle** The angle subtended between a \*streamer and the track of the towing vessel when a cross current causes the cable to drift off-line. The feather angle is of considerable importance when the streamer is several kilometres long and multi-fold coverage is to be obtained, because the position of the streamer affects the \*common-depth-point coverage.

**feather ore** See JAMESONITE.

**Federov stereographic net** See STEREOGRAM.

**feldspars** The most important group of rock-forming \*silicate \*minerals, including the \*alkali feldspars KAlSi<sub>3</sub>O<sub>8</sub> to NaAlSi<sub>3</sub>O<sub>8</sub> (K-feldspar to \*albite) and the \*plagioclase feldspars NaAlSi<sub>3</sub>O<sub>8</sub> to CaAl<sub>2</sub>Si<sub>2</sub>O<sub>8</sub> (\*albite to \*anorthite). The potassium, sodium, and calcium \*end-members can be taken to represent the three apices of an equilateral triangle into which all the alkali feldspars and the plagioclase feldspars can be plotted (usually with the plagioclase feldspars along the base of the triangle and the alkali feldspars along the left-hand side).

**feldspathic greywacke** See ARKOSIC WACKE.

**feldspathic wacke** See ARKOSIC WACKE.

**feldspathoid** The name for a group of framework *\*silicate* minerals which are similar to feldspars in their structure but contain less silica per formula unit. The group includes the minerals *\*nepheline* (a sodic feldspathoid), *\*leucite* (a potassic feldspathoid), and *\*sodalite* (a sodium, chlorine-bearing feldspathoid). The mineral *\*analcime*, although a *\*zeolite*, is closely related to the feldspathoids and often described with them. Minerals within the feldspathoid group crystallize from silica-deficient (*\*silica* undersaturated) *\*melts* and can occur instead of, or with, *\*feldspar*.

**Felidae** See CARNIVORA.

**Feliformia** See CARNIVORA.

**felsenmeer** See BLOCKFIELD.

**felsic** A term applied to light-coloured *\*igneous* *\*minerals* and *\*igneous* rocks rich in these minerals. Typical felsic minerals are *\*quartz*, *\*feldspar*, *\*feldspathoids*, *\*muscovite*, and *\*corundum*. The term felsic derives from the two common minerals, *feldspar* and *silica*.

**felsite** A very light-coloured, *\*aphanitic* *\*igneous* rock, with or without *\*phenocrysts* present. The term is used in the field as an initial classification and can refer to devitrified *\*rhyolite* glass (*\*obsidian*), or primary, *\*cryptocrystalline* rhyolite.

**felsitic** An *\*igneous* texture characterized by an equigranular, *\*cryptocrystalline* aggregate of minerals, usually *\*quartz* and *\*feldspar*, in *\*rhyolites*. The texture is formed during extreme *\*undercooling* of a *\*magma*, a condition which generates numerous *\*nucleation* sites and imposes slow rates of *\*crystal* growth.


**femic** A little-used term to describe the normative *\*ferromagnesian* *\*minerals* in a rock. The *\*CIPW* normative components *\*hypersthene* (Hy), *\*diopside* (Di), fayalite (Fa), and forsterite (Fo) are all femic components of a rock.

**femtoplankton** Marine planktonic organisms, 0.02–0.2 µm in size, about which little is known.

**femtosatellite** A space satellite with a *\*wet mass* of 10–100 g.

**femur 1.** In *\*tetrapods*, the upper bone of the hind limb. **2.** In *\*Insecta*, the third and usually the largest and most robust segment in the leg.

**fence diagram** A three-dimensional depiction of an area, resembling an open area surrounded by a 'wall' or 'fence', showing the location and relationships of its sedimentary deposits. The diagram is constructed from several stratigraphic sections drawn in positions corresponding to their actual locations and their strata are joined.



### Fence diagram

**fenestrae** Irregular cavities found in muddy *\*intertidal* to *\*supratidal* *\*carbonate* sediments. They take a number of forms: *\*birdseye* fenestrae (irregular, 'birdseye'-shaped cavities, usually 1–5 mm across, formed by

gas entrapment in the sediment); laminoid fenestrae (long, thin cavities, parallel to the sediment laminae, formed particularly in algal, laminated muds, and produced by the decay of organic material); and tubular fenestrae (cylindrical, near vertical tubes, formed by burrowing organisms or plant rootlets). Fenestral cavities may become filled with sparry calcite (*\*sparite*). If they remain unfilled, the fenestrae are responsible for the development of *\*fenestral* porosity in the sediment. The term comes from *fenestra* (pl. *fenestrae*), the Latin for an opening or window.

**fenestral fabric** See BIRDSEYE FABRIC; FENESTRAE.

**fenestral porosity** *\*Porosity* developed in *\*carbonates* due to the presence of *\*fenestrae*. Rocks with fenestral porosity will not form good *\*reservoir rocks* unless the fenestrae are interconnected to permit a good *\*permeability* to be established. See CHOQUETTE AND PRAY CLASSIFICATION.

**fenestrated** Perforated with small openings or transparent areas.

**FengYun (FY)** A series of sixteen Chinese meteorological satellites (the name means ‘wind cloud’); satellites in the odd-numbered series are in low-Earth *\*polar orbits*, those in the even-numbered series are in *\*geostationary orbits*. The first satellite, FY-1A, was launched on 6 September 1988. Still operational are FY-2D (launched 8 December 2006), FY-3A (launched 27 May 2008), FY-2E (launched 23 December 2008), FY-3B (launched 4 November 2010), FY-2F (launched 13 January 2012), FY-3C (launched 23 September 2013), FY-2G (launched 31 December 2014), FY-4A (launched 10 December 2016), and FY-3D (launched 14 November 2017).

**fenite** A rock that was first described from the Fen carbonatite intrusion in southern Norway. It is sodium-rich, metasomatic, *\*orthoclase*-*\*nepheline*-*\*arfvedsonite*-*\*aegirine*-bearing, and developed around *\*carbonatite* intrusions. Alkali-rich fluids, migrating away from the crystallizing carbonatite *\*magma*, react with the surrounding rocks, converting them to an assemblage of alkali-bearing *\*minerals*. This process of alkali *\*metasomatism* is given the specific name of ‘fentization’ and produces particularly striking results when the *\*country* rocks are *\*granites*.

**fenitization** See FENITE.

**Fennoscandian Border Zone (Tornquist Line)** An ancient *\*Caledonian* lineation separating the crystalline mass of Scandinavia and Russia from the

fragmented crustal blocks of the remainder of north-western Europe. This NW–SE trending structural feature complements the major Caledonian **\*faults** of northern Scotland (Great Glen Fault, Highland Boundary Fault, etc.). The Fennoscandian Border Zone has been active since that time, and in the **\*Mesozoic** formed the boundary between the stable, crystalline **\*basement** of Sweden and the Baltic and the subsiding Danish–Polish Furrow and North Sea Basin to the south-west. It is a complex zone of faults, **\*horsts**, and half-**\*grabens** which are best seen in southern Sweden (Scania), on the Danish island of Bornholm, and in southern Poland.

**Fennoscandian uplift** The uplift of Fennoscandinavia as a consequence of the inflow of **\*mantle** in order to return to **\*isostatic** balance following unloading of the area as the **\*Pleistocene** ice sheets melted.

**Ferdinand (Uranus XXIV)** A lesser satellite of **\*Uranus** with a radius of 10 km.

**ferns** See PTEROPSIDA.

**ferralitization** See FERRALIZATION.

**ferralization (ferralitization)** Part of the **\*leaching** process found in tropical soils, by which large amounts of iron and aluminium oxides accumulate in the B **\*horizon** of such soils as krasnozems.

**ferralsols** A reference soil group in the **\*World Reference Base for Soil Resources** classification scheme. Ferralsols have a highly weathered **\*ferric** horizon with high concentrations of iron and aluminium.

**Ferrel cell** In the **\*general circulation** of the atmosphere, an indirect cell in middle latitudes in which the air circulation is driven by the **\*Hadley** and **\*polar** cells. The cell was discovered by the American climatologist William Ferrel (1817–91).

**ferric horizon** In the **\*World Reference Base for Soil Resources** classification, a **\*soil horizon** with distinctive red mottling that is more than 15 cm thick.

**ferricrete** See DURICRUST.

**ferrimagnetism** The property of a **\*ferromagnetic** substance, i.e. one in which one of the two antiparallel magnetic lattices is stronger than the

other, so that the substance has a net **\*remanent** magnetization even in the absence of an external magnetic field. **\*Magnetite** is the commonest ferrimagnetic rock mineral.

**ferro-** A prefix attached to an **\*igneous** rock name to indicate a particular abundance of iron-bearing **\*minerals**, or a high whole-rock iron content.

**ferroactinolite** See TREMOLITE.

**ferroan** Containing ferrous iron ( $\text{Fe}^{2+}$ ).

**ferroaugite** See BENMOREITE.

**ferroelectricity** A property of certain **\*dielectric** materials that they spontaneously polarize, i.e. centres of positive and negative electric charge separate; application of an appropriate electric field causes the polarity to reverse.

**ferrohastingsite** See NEPHELINE-SYENITE.

**ferromagnesian minerals** **\*Silicate** **\*minerals** in which **\*cations** of iron and magnesium form **\*essential** chemical components. The term is used to cover such minerals as the **\*olivines**, **\*pyroxenes**, **\*amphiboles**, and the **\*micas**, **\*biotite**, and **\*phlogopite**.

**ferromagnetic** In the wide sense, applied to substances in which **\*electron** spins are magnetically coupled by either exchange or super-exchange quantum-mechanical forces. Such materials can acquire a spontaneous magnetization which is much greater than either **\*diamagnetism** or **\*paramagnetism**. There are three types of ferromagnetism: **\*ferromagnetism** (in the strict sense); **\*ferrimagnetism**; and **\*antiferromagnetism**.

**ferromagnetism** In the strict sense, magnetism occurring in substances in which the **\*electron** spins are coupled by quantum-mechanical exchange forces so that, within a single volume element, all electron-spin vectors are in the same direction. Typical materials are pure iron, nickel, and iron-nickel alloys found in **\*meteorites** and lunar samples. See also FERRIMAGNETISM; ANTIFERROMAGNETISM.

**ferropericlase (magnesiowüstite)** A magnesium iron (FeII)-bearing variety of **\*periclase** ((Mg,Fe)O) that is the second-most-abundant solid phase in

the Earth's interior (after *\*bridgmanite*).

**ferrosilite** See ORTHOPYROXENE.

**fertility, soil** Condition of a soil relative to the amount and availability to plants of elements necessary for plant growth. Soil fertility is affected by physical elements, e.g. supply of moisture and oxygen, as well as by the supply of chemical plant nutrients.

**Ferungulata** (class *\*Mammalia*, infraclass *\*Eutheria*) A cohort proposed by G. G. *\*Simpson* in 1945 on palaeontological grounds but regarded by some authorities as artificial, that includes the *\*Carnivora* (all of the modern mammalian carnivores), the primitive ungulates (including the elephants and sea cows), the *\*Perissodactyla* (tapirs, rhinoceroses, horses, etc.), the *\*Artiodactyla* (pigs, camels, cattle, etc.), and the Tubulidentata (aardvarks, etc.), which are believed to have arisen from a common population in the *\*Palaeocene*.

**Festiniogian (Franconian)** A *\*stage* of the Late *\*Cambrian* epoch, 496.8–492.5 Ma ago, preceded by the *\*Maentwrogian* and followed by the *\*Dolgellian*.

**fetch** **1.** Length of water surface over which the wind blows in generating waves. Together with wind velocity and duration, this determines wave height. Many features of coastal deposition tend to become orientated normally to the direction of maximum fetch. **2.** Distance over which an air stream has travelled across sea or ocean.

**Ffestiniogian** See MAENTWROGIAN.

**FFT** See FAST FOURIER TRANSFORM.

**fiamme** Flattened, elongate, *\*pumice* *\*clasts* that are found in many welded *\*ignimbrites*. The pumice clasts, which are in a hot, plastic state when deposited from the *\*pyroclastic flow*, are compressed and flattened by the weight of the overlying ignimbrite body, producing elongated glassy clasts with ragged, flame-like ends (*fiamme* is the Italian word for 'flame'). Many *fiamme* clasts continue to vesiculate after formation, leading to the development of spherical *\*vesicles* within them.

**fiard (fjard, firth)** A coastal inlet similar to a *\*fiord*, but with lower relief.

**fibratu**s From the Latin *fibratu*s meaning ‘fibrous’, a species of separate cloud or cloud veil which has rather curved elements, but without hooks. *See also* CLOUD CLASSIFICATION.

**fibril** Cloud trail observed in \*cumulonimbus, where \*drizzle-sized droplets are large enough for their terminal velocities to allow them to depart from the main cloud body.

**fibrolite** *See* SILLIMANITE.

**fibrou**s Applied to the physical \*form of a mineral that occurs in fine, thread-like strands which may be parallel or radiating in nature, e.g. \*asbestos.

**fibula** In \*tetrapods, the post-axial bone of the lower part of the hind limb.

**Fick’s laws of diffusion** Two laws, formulated in 1855 by the German physiologist and physician Adolf Eugen Fick (1829–1901) that describe how one substance moves into another by \*diffusion. The first law states that molecules move from a region of higher concentration to a region of lower concentration at a rate proportional to the concentration gradient. The second law describes how diffusion causes the concentration gradient to weaken over time. Fick studied diffusion in fluids, but his laws are now known to apply equally to diffusion in solids.

**fiducial point** 1. Temperature at which the atmospheric-pressure scale of a particular barometer reads correctly. The temperature at which this is so in latitude 45° is called the standard temperature of that barometer. At other temperatures and other latitudes, corrections must be applied. 2. Fixed point (indicated by a pointer) that is the zero of the scale of a \*Fortin barometer.

**field capacity** The water content which can be retained by a soil after excess moisture has drained freely away. Usually it is measured as a percentage of the soil volume or of the weight of oven-dry soil. *See also* SOIL-MOISTURE CONTENT.

**field reversal** *See* GEOMAGNETIC FIELD; POLARITY REVERSAL.

**Figtree** *See* SWAZIAN.

**Filicopsida** *See* PTEROPSIDA.

**filiform** Thread-like; long and slender.

**filling** Term used in synoptic meteorology to describe an increase in pressure at the centre of a *\*depression*. *See also* DEEPENING.

**film water** *See* PELLICULAR WATER.

**filter** **1.** A device for removing unwanted components from water. Coarse material may be recovered by the use of simple mesh sieves, but finer material and certain pollutants may require the use of other filters, e.g. of activated carbon or sand. **2.** To discriminate against a portion of information entering a device (the filter), typically by removing unwanted *\*noise* or isolating specific parts of the information (e.g. separating high-frequency from low-frequency data). Filters usually operate within the frequency domain although others exist (e.g. the velocity filter). Frequency filters have the disadvantage of invariably distorting the signal pulse shape, lengthening the pulse and so causing a phase shift, and displacing peaks and troughs in time. Linear filtering is known as *\*convolution*. *See also* ALIASING; BAND FILTER; SPATIAL-FREQUENCY FILTER; WIENER FILTER.

**filter route** Term introduced by the American palaeontologist G. G. *\*Simpson* (1940), to specify a faunal migration route along which the spread of some animals is very likely but the spread of others is improbable. The route thus filters out part of the fauna, but permits the rest to pass. Deserts and mountain ranges provide examples of filter routes.

**filtrate** *See* MUD FILTRATE.

**filtration** Essentially, the removal of solid matter from liquids. Filter systems may be physical, mechanical, biological, chemical, or electrokinetic.

**fin** Appendage of fish and fish-like aquatic animals used for locomotion, steering, and balancing of the body. The skin fold forming the fin membrane is supported by cartilaginous, horny, or bony fin rays, which can be soft and flexible (soft rays) or hard and inflexible (fin spines).

**finds** *\*Meteorites* identified by their composition and structure but not seen to fall. *Compare* FALLS.

**finer** 1. Particles of material below a specified size; or fine-grained sediment which settles very slowly. 2. In *\*ore* processing, material crushed or ground too finely, or ores too powdery, for normal smelting.

**finer 10% test** A test that is similar to the *\*aggregate crushing value (ACV)* test. A sample of aggregate is subjected to varying loads and the percentage of *\*finer* calculated for each load. A graph of loading versus percentage finer is used to find the loading required for 10% finer.

**finger lake** A long, narrow, deep lake that probably formed by glacial deepening of a pre-existing river channel. The eleven Finger Lakes in upstate New York are the most famous examples.

**Fingerlakerian (Fingerlakerian)** See *SENECAN*.

**fining-upward succession** A vertical change in a *\*facies* in which the grain size decreases with height above the base. Compare *COARSENING-UPWARD SUCCESSION*.

**finite resource** See *NON-RENEWABLE RESOURCE*.

**finite rotation** In *\*plate tectonics* theory, the amount of rotation about a fixed point that would bring a plate back to some other defined position.



<http://www.serg.unicam.it/Models.html>

- Plate Tectonic Modeling: Tools and Methods; Rotation Models.

**finite strain** The total amount of *\*strain* which has accumulated incrementally over a period through the addition of many smaller strains. The investigation of present-day rock structures involves the analysis of finite strain states which are evaluated by reference to natural objects whose original shape is known.

**fiord (fjord)** Long, narrow, deep, U-shaped coastal inlet which usually represents the seaward end of a glaciated valley that has been partially submerged. The water depths often exceed 1000 m except near the mouth where a bar or sill may be present.

**fire-ball** A very bright *\*meteor*, approximating the average magnitude (brightness) of Venus. See also *BOLIDE*.

**fire damp** A flammable gas that occurs in coal mines; it explodes readily at concentrations between 4% and 16%, and explosions are most violent at a 10% concentration. The gas is 70–98% methane mixed with small and variable amounts of other gases.

**Firefly** A low-cost, 3-unit *\*CubeSat* *\*NASA* mission, funded and managed by the National Science Foundation, to study the relationship between lightning and terrestrial gamma-ray flashes. The *\*nanosatellite* was launched on 20 November 2013, from Virginia.

**fire-fountain** A continuous spray of disrupting *\*magma* through a vent to form a persistent fountain of molten magma above the vent. The fountain, which may rise to 200 m, is supported either by the hydrostatic pressure of magma in the upper levels of the main volcanic superstructure, or by expanding gas released from the magma during the *\*eruption*. Fall-out from the column produces a *\*spatter* rampart around the vent and if the accumulation rate is high the molten spatter may coagulate to form a flow of *\*lava* (a ‘clastogenic flow’).

**firn (névé)** Snow that has survived a summer melting season. It is an intermediate material in the conversion of snow to glacial ice. Normally it is granular, due to the partial melt.

**firn limit** See *FIRN LINE*.

**firn line (annual snow-line, firn limit)** A line on a *\*glacier* marking the upper limit to which winter snowfall melts during the summer *\*ablation* season. It is often clearly marked, and on many glaciers separates hard, blue ice below from snow above.

**firn wind (glacier wind)** Downhill airflow which develops over a glacier during the day, usually in summer. The greater air density over the glacier than over the surrounding surfaces causes this air to sink.

**first arrival** See *FIRST BREAK*.

**first break (first arrival)** The first wave from a discrete seismic-source impulse, naturally or artificially generated, that is recorded at a seismic detector. First breaks are used in *\*seismic* refraction surveying. See *BREAK*.

**firth** See *FIARD*.

**fish** An instrument package towed behind a ship so that the measurements are unaffected by the ship and its equipment. A fish commonly contains a **\*magnetometer** and **\*side-scan** sonar.

**Fisher, Osmond** (1817–1914) An Anglican clergyman, Fisher was the author of the first textbook on **\*geophysics**: *The Physics of the Earth's Crust* (1881). He argued that the Earth had a thin crust, and that convection currents in the fluid interior were the cause of mountain building, **\*rifts**, etc. He postulated that the **\*Moon** had been torn from the Pacific, causing the continents to be pulled apart.

**fish-hook beach** See ZETA-FORM BEACH.

**fish-tail bit** A drill **\*bit** used for cutting through soft sediments.

**fissility** Ability of rock materials to split. The term is applied to **\*shales**, flags, **\*slates**, and **\*schists**.

**fission** Splitting of a heavy atomic nucleus by collision, with the ejection of two or more neutrons, and the release of much energy.

**fission hypothesis** One of the three classical hypotheses for lunar origin. Proposed by George Darwin in 1879, it derives the **\*Moon** from the silicate **\*mantle** of the **\*Earth**, following **\*core** separation. Although this accounts for the low density of and paucity of metallic iron in the Moon, the process requires about four times the observed **\*angular momentum** of the present Earth–Moon system. Detailed compositional differences between the Moon and the terrestrial mantle, from the Apollo data, appear fatal to the hypothesis.

**fission-track dating** Charged particles, from the spontaneous fission of  $^{238}\text{U}$  in **\*minerals** and in natural and synthetic **\*glasses**, leave a trail of damage (fission tracks, radiation tracks) as they travel through a solid medium. This is the result of the transfer of energy from the particles to the atoms of the medium. These tracks, suitably enlarged by etching, can be seen in some minerals by using a petrological microscope (see **POLARIZING MICROSCOPE**). The number of tracks per unit area is a function of the age of the specimen and its uranium concentration, provided that it cooled rapidly on formation and has not been reheated at a later date. The uranium concentration can be measured by counting tracks produced by fission of

$^{235}\text{U}$  caused by irradiation of the specimen with thermal neutrons in a nuclear reactor. Fission-track dates can, by this method, be obtained for minerals such as *\*micas*, *\*apatite*, *\*sphene*, *\*epidote*, and *\*zircon*. The dates obtained are 'cooling ages' and indicate the time elapsed since the temperature dropped below the 50% track retention value; the tracks are known to fade by annealing of solids at elevated temperatures. The method can also be used to date *\*tektites*, volcanic glass, and some archaeological objects.

**fissure volcano** A linear fracture on the Earth's surface through which *\*lavas*, *\*pyroclastics*, and gas are erupted and effused. The eruptive products accumulate most thickly along the linear fracture and build up an elongate, low-angle shield or higher-angle cone topography, constituting the volcanic pile. In addition to *\*eruptions* from the main fissure, material can be erupted from secondary fissures developed locally within the growing volcanic pile and radiating from the trend of the main fissure. *See also* [CENTRAL VENT VOLCANO](#); [VOLCANO](#).

**fixation** 1. Soil process by which certain nutrient chemicals required by plants are changed from a soluble and available form into a much less soluble and almost unavailable form. 2. (**nitrogen fixation**) The biochemical process by which certain bacteria convert gaseous nitrogen into ammonia ( $\text{NH}_3$ ).

**fixed-source method** A geophysical exploration method in which the source (transmitter) is kept at a fixed position and a detector (receiver) is moved over the survey area to take measurements which are then plotted as profiles or maps. Electromagnetic methods such as *\*VLF* and the *\*TURAM* technique are examples of fixed-source methods. *Compare* [MOVING-SOURCE METHOD](#).

**fixigena** *See* [CEPHALON](#).

**fjard** *See* [FIARD](#).

**fjord** *See* [FIORD](#).

***f-k space*** A means of representing *\*frequency-domain* data in terms of the independent variables *\*frequency* ( $f$ ) and *\*wavenumber* ( $k$ ).

**Fladbury** A site in Worcestershire, England, where palaeontological and palaeozoological evidence suggests there was a barren, tundra-like landscape during a cold period following the **\*Upton** Warren Interstadial.

**flagstone** A stonemason's term to describe a hard stone, commonly a **\*sandstone** or **\*siltstone**, that splits readily into slabs that are flat on both sides and can be used to make paving, roof tiles, and field boundaries.

**flake tectonics** Where two **\*plates** are colliding, a process in which large sections, or flakes, shear off the top of one of the plates while the lower part is subducted (see **SUBDUCTION**) into the **\*mantle**.

**flakiness index (I<sub>F</sub>)** The specification for stone for bituminous surfacing, applied to **\*aggregate** coarser than 6.5 mm. It is expressed as the percentage by weight of particles (in a sample of more than 200) whose smallest dimension is less than 0.6 times the mean dimension.

**flame photometry (flame spectrometry)** A technique analogous to emission spectrometry, but using a flame to excite electrons, rather than an arc or plasma. It is a simple and straightforward analytical technique that is basically a quantitative version of a 'flame test'. A known weight of sample is dissolved in hydrofluoric acid and either perchloric or sulphuric acids, portions of the solution are added to a flame, and the strength of emission of light of a particular wavelength produced by the potassium in the flame is recorded. This is then compared with those produced by standard solutions. The final results may be affected by sodium concentrations as well as by the sulphuric acid. Perchloric acid, iron, magnesium, aluminium, and calcium also interfere with the potassium emission but their effects may be reduced by buffering and by the removal of interfering **\*ions**.

**flame spectrometry** See **FLAME PHOTOMETRY**.

**flame structure** A sedimentary structure in which wavy tongues of mud, with shapes resembling flames, project into the rock above them, which is commonly a **\*sandstone**.


**flamma** See **CLOUD CLASSIFICATION**.

**flammagenitus** See **CLOUD CLASSIFICATION**.

**Flandrian** The present \*interglacial. Evidence suggests that the so-called post-glacial period, the warm phase following the last (\*Devensian) ice advance or cold phase, is more appropriately treated as another interglacial of the \*Quaternary (or Late \*Cenozoic) Ice Age. In Europe, the warmest Flandrian stage occurred during \*Atlantic times, about 6000 BP (the Hypsithermal is the equivalent N. American climatic optimum). No consensus view exists as to when the ice advance or extreme cold conditions will prevail once again in high mid-latitudes, nor as to how quickly these conditions will arise. The Flandrian is sometimes referred to alternatively as the \*Holocene interglacial.

**flank eruption** The release of \*lava and \*pyroclastic material from a source on the slopes of a \*volcano, away from its primary \*central vent or fissure-vent area. The location of a flank eruption is often controlled by local inflation stresses. These initiate fractures within the volcanic pile which propagate outwards from the primary volcanic conduit to feed lava towards the volcanic flanks. A flank eruption occurs where the lava-filled fractures intersect the slope of the volcano.

**flaser bedding** A form of \*heterolithic bedding characterized by \*cross-laminations draped with \*silt or \*clay. Flaser beds form in environments where \*flow strengths fluctuate considerably, thus permitting the transport of \*sand in \*ripples, followed by low-energy periods when mud can drape the ripples.



**Flaser bedding**

**flaser gneiss** See FLASER ROCK.

**flaser rock (flaser gneiss)** A rock displaying ovoidal *\*megacrysts* enclosed within a fine-grained, streaky, *\*anisotropic \*groundmass*. The groundmass is typical of a *\*mylonite* fabric produced by intense deformation. The megacrysts are regions of the original rock which have survived deformation. 'Flaser' is a German dialect word meaning 'irregular vein', or 'knot' (as in wood).

**flash flood** A brief but powerful surge of water either over a surface ('sheet flood') or down a normally dry stream channel ('stream flood'). Usually it is caused by heavy convective rainfall of short duration and is typical of semi-arid and *\*desert* environments, although it also occurs in temperate regions.

**flat** That part of a staircase *\*thrust* plane trajectory which has a horizontal or subhorizontal orientation. Flats represent areas of décollement (see DÉCOLLEMENT PLANE) along weak, layer- or bedding-parallel planes. Although initially horizontal, flats may steepen during later compression. See RAMP; THRUST.

**flat bed** See PLANE BED.

**flat-iron** 1. A land-form of roughly triangular shape, and with one side (the *\*dip slope*) that is both steep and uniform. It is formed between two adjacent valleys that cut through a *\*hogback* ridge roughly at right angles to its *\*trend*. It is a common land-form along the eastern front of the Rocky Mountains, USA. 2. Applied to *\*clasts* that have been shaped by glacial *\*erosion* at the base of sliding *\*ice*. Typically, they show a distinctive 'bullet' form, with one end plucked and the other streamlined, and may vary in size from a few centimetres to many metres.

**flat spot** A characteristic, strong, flat reflection within an otherwise dipping reflection event which is produced by a strong positive reflection from a gas-water interface, typically in a *\*trap* structure. Often a flat spot occurs below a *\*bright* spot.

**flattening** A change in the shape of an object caused by the application of a *\*stress*, and which may be described by reference to the transformation of a sphere into an oblate ellipsoid as a result of *\*pure* shear. Flattening is the strain state found in S-*\*tectonites* (i.e. tectonites marked by a single, penetrative *\*foliation*).

**F-layer** The transition zone between the liquid outer *\*core* and the inner solid core of the *\*Earth*. The boundaries are poorly defined but it is at a depth of about 5100 km.

**FLEX** See FLUORESCENCE EXPLORER.

**flexural basin** See FORELAND BASIN.

**flexural isostasy** The *\*isostasy* that occurs when a load on the *\*lithosphere* is not compensated by local *\*buoyancy*. The load is supported regionally by the strength of the lithosphere, which bends (flexes) in response.

**flexural rigidity** The flexural rigidity of an elastic sheet is defined as  $ET^3/12(1 - \sigma^2)$  where  $E$  is **\*Young's modulus**,  $T$  is the thickness of the sheet, and  $\sigma$  is **\*Poisson's ratio**. In the Earth, the **\*lithosphere** is usually treated as an elastic sheet that responds to loading by **\*ice caps**, volcanic piles, etc., from which the thickness of the lithosphere and the **\*viscosity** of the **\*mantle** can be derived.

**flexural slip** Folding in which there is **\*slip** along the contacts between parallel layers. This discontinuous **\*simple** shear mechanism may occur along **\*bedding planes** or cleavage planes.

**flexure** 1. The lateral deflection from a datum line of a planar feature as it is shortened. 2. The form of a **\*monocline**, or a gentle fold whose inter-limb angles are 120–180°.

**flexus** A low, curved ridge that forms a scalloped pattern on the surface of an extraterrestrial body.

**flight** See **STRING**.

**Flinn diagram** A graphical representation of the full range of three-dimensional **\*strain** states in deformed rocks, with or without a change in their volume. The diagram plots a value for the **\*principal strain axes**  $Y/Z$  against  $X/Y$  of ellipsoids (see **STRAIN ELLIPSOID**) which result from **\*dilation** and/or distortion of a reference sphere. The data plotted on the diagram are obtained from analyses of strain using **\*strain markers**.

**flint (silex)** Variety of **\*chert**, which occurs commonly as nodules and bands in **\*chalk**. It is deposited in the porous, permeable structures of **\*sponge**, **\*diatom**, and **\*echinoid** skeletons and also in burrows.

**floating point** A style of expressing a number that avoids losing significant figures should the number be too large or too small for a given register on a calculating device. For example, 165 400 can be written as  $1.654 \times 10^5$  in floating point (although computers usually use base-2 rather than base-10). Floating-point amplifiers rather than binary amplifiers are being used increasingly in seismic amplification systems to increase the effective range of digital recordings.

**floatstone** A coarse-grained **\*limestone** with **\*matrix-supported** **\*clasts**, 10% or more of which are coarser than 2 mm in size. See **EMBRY AND**

## CLOVAN CLASSIFICATION.

**flocculation** Process in which *\*clay* and other soil particles adhere to form larger groupings or *\*aggregates*. The reverse of this process is known as *\*dispersion*.

**floccus** From the Latin *floccus* meaning 'tuft', a species of cloud with a tufted appearance, the lower parts of which are rather ragged, often with *\*virga*. The species is most associated with *\*cirrus*, *\*cirrocumulus*, *\*altocumulus* and *\*stratocumulus*. *See also* CLOUD CLASSIFICATION.

**Flock 1** A constellation of twenty-eight Earth-imaging, 3-unit *\*CubeSats*, each with a mass of about 5 kg, developed by Planet Labs, of San Francisco. The constellation was launched on 9 January 2014, from Virginia.

**flood basalt** *\*Lava* of basaltic composition which is erupted from a laterally continuous fissure to form a widely dispersed, *\*low* aspect-ratio, flow sheet. The Roza Member of the mid-*\*Miocene* Columbia River Plateau in the USA is the largest flood *\*basalt* unit in the geologic record, comprising basalt flows with a total volume of more than 1500 km<sup>3</sup> which have travelled up to 300 km from their source fissure. The largest individual lava flows in this member have volumes up to 700 km<sup>3</sup>.

**flood forecasting** A technique which uses the known characteristics of a river basin to predict the timing, discharge, and height of flood peaks resulting from a measured rainfall, usually with the objective of warning populations who may be endangered by the flood. *Compare* FLOOD PREDICTION.

**flood-peak formulae** A number of methods (including those attributed to Benson, Potter, Morisawa, and Rodda) for the prediction of flood peaks by reference to rainfall intensity and frequency, topography, and orographic, temperature, and other relevant factors.

**floodplain** The part of a river valley that is made of unconsolidated river-borne sediment, and periodically flooded. It is built up of relatively coarse debris left behind as a stream channel migrates laterally, and of relatively fine sediment deposited when *\*bankfull* discharge is exceeded.

**flood prediction** The study of rainfall patterns, catchment characteristics, and river hydrographs to predict the future average frequency of occurrence of flood events. Flood predictions seek to estimate the probable discharge that, on average, will be exceeded only once in any particular period, hence the use of such terms as '50-year flood' and '100-year flood'. *Compare* FLOOD FORECASTING.

**flood tide** Rising *\*tide*: the phase of the tide between low water and the next high tide. *Compare* EBB TIDE.

**flood zone** *See* ACME ZONE.

**floor thrust** The lowest *\*thrust* surface bounding a *\*duplex* system, which joins the *\*roof thrust* at the leading and trailing edges of the duplex. If it is the lowest regional thrust surface it may be called a *\*sole thrust*.

**flora** (*adj.* floral, floristic) All the plant species that make up the vegetation of a given area. The term is also applied to assemblages of fossil plants from a particular geologic time, or from a geographical region in a former geologic time. Examples of all three types of usage, respectively, are: British flora, *\*Carboniferous* flora, and *\*Gondwana* flora. *Compare* FAUNA.

**Florian** An Australian *\*stage* (508–506 Ma ago) of the Middle Cambrian, preceded by the *\*Templetonian* and followed by the *\*Undillian*.

**Florida current** Part of the *\*Gulf Stream* that extends from the southern tip of Florida to Cape Hatteras, N. Carolina. It is a fast-flowing (1–3 m/s), narrow (50–75 km wide), and deep current, still evident at depths of 2000 m where velocities of up to 10 cm/s have been measured. It is an example of a western *\*boundary* current.

**floristics** *See* PHYTOGEOGRAPHY.

**flotation separation** A concentration process whereby finely ground *\*ore* is dispersed in water containing a flotation reagent which causes selected minerals to become hydrophobic. Aeration and agitation of the suspension allows hydrophobic particles to float while unaffected minerals sink. The floating particles are skimmed off or overflow the flotation cell.

**flow** See BANKFULL FLOW; BASEFLOW; CRITICAL FLOW; FROUDE NUMBER; GROUNDWATER FLOW; INTERFLOW; LAMINAR FLOW; QUICKFLOW; STEADY FLOW; SUBSURFACE FLOW; SURFACE RUNOFF; TURBULENT FLOW; UNDERFLOW; UNIFORM FLOW.

**flow cleavage** A type of \*cleavage which is intermediate between a \*slaty cleavage and a \*schistosity. The term has fallen from favour and several authorities (e.g. C. McA. Powell (1979)) recommend that it should not be used in descriptions of cleavage because it implies the mode of origin of the cleavage.

**flower structure** A series of convex-upward thrust or \*reverse faults found in \*transpressional \*strike-slip zones. On seismic sections the appearance of the structure is reminiscent of the petals of a flower or the leaves of a palm tree, hence the analogy. Such structures are important in oil exploration as they indicate strike-slip movements and therefore the possible development of pull-apart basins, and also provide potential areas for oil accumulation.

**flow folding** Folding which results mainly from continuous \*simple shear or viscous flow within layers, producing deformations analogous to lamellar flow in liquids. The term 'flow fold' has also been used synonymously with \*ptygmatic fold.

**flowmeter** An instrument for measuring the flow of liquids.

**flow slide** See RETROGRESSION.

**flowstone** See DRIPSTONE.

**flow till** Sediments which flow after they have been deposited by \*ablation. See TILL.

**fluctus** 1. A feature formed by flowing liquid on the surface of \*Io, \*Titan, and \*Venus. 2. See CLOUD CLASSIFICATION.

**flud mud** \*Fluvial material that is only partly settled in the \*subtidal zone of a well-mixed \*estuary.

**fluid inclusion** Usually minute amount of liquid and/or gas trapped in a crystal during crystallization or \*recrystallization. There are two ways in

which solid phases in fluid inclusions may originate. (a) One or more mineral grains may be trapped along with the fluid phase(s) during formation of the fluid inclusion. (b) One or more solid phases may form in a fluid inclusion after its initial formation as a result either of a reaction between the fluid and the host mineral, or by precipitation from the fluid upon cooling (in which case the solids are known as daughter minerals). The temperature and composition of the fluid from which the enclosing mineral originated can be estimated from studies of such inclusions.

**fluidization** Process of passing gas through loose, fine-grained particles causing the mixture to flow like a liquid; this facilitates mixing and chemical reaction. The faster the gas flows the more the mixture expands and movement increases. A bubble phase may form which travels upwards until all the solid particles are transported by the gas. Fluidization is used in coal-fired power stations and may occur naturally in volcanic eruptions producing *\*pyroclastic flows* and surges. It is the phenomenon which enables pyroclastic flows to travel distances in excess of 100 km and to surmount topographic obstacles hundreds of metres high.

**flume 1.** A short section of artificial channel constructed in a river in order to create a constriction in which *\*critical flow* will be established, allowing the *\*discharge* to be calculated from the water depth. **2.** An experimental channel used for studying relationships between *\*sediment* movement and *\*flow* conditions. There are a number of different flume designs but most flumes are capable of carrying water at variable depths and velocities, either in a unidirectional flow or generating waves. Flume studies have been responsible for establishing the important relationships between the grain size (see *PARTICLE SIZE*) and *\*erosion* velocities and stability fields for the various sediment *\*bedforms*.

**flumen 1.** (*pl. flumina*) On the surface of *\*Titan*, a channel through which liquid might flow. **2. (inflow band)** See *CLOUD CLASSIFICATION*.

**flumina** See *FLUMEN*.

**fluorescence** Kind of luminescence, in which an atom or molecule emits radiation when electrons within it pass back from a higher to their former, lower energy state. The term is restricted to the phenomenon in cases where the interval between absorption and emission is very short (less than  $10^{-3}$

s). See also PHOSPHORESCENCE; X-RAY FLUORESCENCE; X-RAY FLUORESCENCE SPECTROMETRY.

**Fluorescence Explorer (FLEX)** An Earth Explorer mission by the \*European Space Agency that will provide maps of vegetation fluorescence, which reflects photosynthetic activity and plant health and stress. The mission is scheduled to be launched in 2022.

**fluorite (fluorspar, Blue John)** Mineral,  $\text{CaF}_2$ ; sp. gr. 3.2; \*hardness 4; \*cubic; often yellow, green, blue, or purple, but can be colourless, pink, red, or black, and often colour banded; white \*streak; vitreous \*lustre; \*crystals often cubes, but can be \*octahedra and \*rhombohedra, and a mixture of forms; \*cleavage perfect {111}; widely distributed in mineral veins alone or as a \*gangue mineral with metallic ores, and in association with \*quartz, \*barite, \*calcite, \*galena, \*cassiterite, \*sphalerite, and many other minerals; soluble in sulphuric acid with the evolution of hydrogen sulphide. It is used extensively as a flux in the smelting of iron, in the ceramic industry, and in the chemical industry. The deep-purple, banded variety, Blue John, is used as an ornamental stone.

**fluorometer** An analytical instrument used mainly in chemical analysis to measure the fluorescent radiation emitted by any particular substance. It works by exposing the substance under investigation to monochromatic radiation.

**fluorspar** See FLUORITE.

**flushed zone (invaded zone)** The zone in a sediment bordering a drill hole in which the \*groundwater has been replaced by \*mud filtrate.

**flute cast** See FLUTE MARK.

**fluted moraine** A ground \*moraine surface which shows streamlined ridges and grooves trending (see TREND) at right angles to the ice front. Individual ridges are generally less than 1 km long, and less than 10 m high. They may form as a consequence of high vertical ice pressure forcing a subglacial, plastic \*till up into the low-pressure zone down-glacier of a large boulder.

**flute mark** A tongue-shaped scour cut into mud by a turbulent flow of water. The tongue is deepest at the up-current end and the flute can thus be

used as a [\\*palaeocurrent](#) indicator. If the flute is infilled by [\\*sediment](#) a flute [\\*cast](#) will be preserved in the base of the overlying bed. Although once believed to be diagnostic of [\\*turbidite](#) deposition, flutes can form in any setting where water flows strongly over soft mud.

**fluvial** Pertaining to a river.

**fluvial processes** The set of mechanisms that operate as a result of water flow within (and at times beyond) a stream channel, bringing about the [\\*erosion](#), transfer, and deposition of [\\*sediment](#). The erosional processes include: the displacement of bed particles through drag and lift forces; [\\*corrasion](#), the wearing away of bed and banks as mobile sediment is dragged against them; and bank collapse, a consequence of hydraulic activity. Transport processes include the transfer of material in solution and suspension, and by [\\*saltation](#). Depositional processes act when the immersed weight of a particle is greater than the force driving it down-channel.

**fluviatile** Applied to [\\*sediments](#) of fluvial (river) origin.

**fluvic horizon** In the [\\*World Reference Base for Soil Resources](#) classification, a dark-coloured [\\*soil horizon](#), usually rich in [\\*pyroclastic](#) deposits.

**fluviokarst** A [\\*karst](#) landscape in which the predominant land-forms are river valleys.

**fluvisols** A reference soil group in the [\\*World Reference Base for Soil Resources](#) classification scheme. Fluvisols form on recent [\\*alluvial](#) deposits and have a [\\*fluvic](#) horizon lying about 25 cm below the surface and extending to 50 cm depth.

**fluxgate magnetometer** [\\*Magnetometer](#) based on two parallel solenoids, equally and oppositely wound on high-permeability cores, and driven by a high-frequency alternating current. A signal coil detects any bias arising from the presence of an ambient magnetic field.

**fluxoturbidite** A poorly graded [\\*sediment](#), the product of gravity-induced flow in which little turbulent mixing of particles occurs. It is transitional between a [\\*slump](#) and a turbidity flow (see [TURBIDITY CURRENT](#)).

**flyer** See [STRING](#).

**Flying Laptop** A German *\*minisatellite* mission, developed by the Institut für Raumfahrtssysteme, to demonstrate minisatellite technologies, conduct Earth-observation experiments, observe ship traffic, and detect *\*Near Earth Objects*. The satellite was launched on 14 July 2017, from the Baikonur Cosmodrome, Kazakhstan.

**flysch** Sedimentary *\*facies* term used to describe a thick succession of redeposited, deep-sea, *\*clastic* material of *\*synorogenic* character.

**focal mechanism, earthquake** Shallow *\*earthquakes* are considered to occur when prolonged tectonic stress exceeds the local yield strength of rocks, so that brittle *\*failure* occurs suddenly, with associated earthquake and stress drop. Intermediate and deep earthquakes appear to have variable focal-plane mechanisms, although they are predominantly associated with *\*slip* along fracture planes. Other mechanisms include sudden phase transitions, resulting in an implosion as the mineral density increases and its volume decreases. See [FAULT-PLANE SOLUTION](#).

**focus (of earthquake)** See [HYPOCENTRE](#).

**fodinichnia** The excavations formed by *\*deposit* feeders in search of food. The category includes radial traces (see [TRACE FOSSIL](#)), e.g. *\*Chondrites*, and U-shaped tubes, e.g. *Rhizocorallium*.

**foehn wall** See [FÖHN WALL](#).

**foehn wind** See [FÖHN WIND](#).

**fog** A type of *\*precipitation* caused by the presence of *\*stratus* cloud at surface level, reducing horizontal visibility to less than 1 km. The formation of fog is aided by any concentration of smoke particles, which act as *\*condensation nuclei*, and may cause fog at levels of humidity below saturation point. See also [ADVECTION](#); [FRONTAL FOG](#); [RADIATION FOG](#); [SMOG](#). Compare [HAZE](#) and [MIST](#).

**föhn wall (foehn wall)** Mass of cap cloud and associated precipitation over windward slopes and parts of leeward slopes on mountain barriers, resulting from the föhn effect (see [FÖHN WIND](#)).

**föhn wind (foehn wind)** Generic term for warm, dry winds in the lee of a mountain range. It was originally used in the European Alps. After cooling


on the windward ascent at the *\*saturated adiabatic lapse rate* of about 0.5 °C/100 m, with the resulting condensation and precipitation, the air subsiding on the leeward side of the mountain range is warmed through compression at the *\*dry* adiabatic lapse rate of 1°C/100 m. This produces a warming wind on the lee side, with higher temperatures than occurred in the same air on the upslope side of the mountains.

**foid** A contraction of the term *\*feldspathoid*, which is applied to any plutonic rock containing up to 60% *\*modal* feldspathoid minerals. For example, a *\*syenite* with significant *\*nepheline* present can be termed a 'foid-bearing syenite', or a 'foid-syenite'. Such terminology is used on the Streckeisen classification of *\*igneous* rocks with a *\*colour index* of less than 90. In practice, the terms 'foid-bearing' and 'foid-' are replaced with the specific feldspathoid mineral name. In the above example, the name 'nepheline-syenite' would apply. Where the modal volume of feldspathoid minerals exceeds 60%, the rock is termed a 'foidolite'.

**foidolite** See FOID.

**fold 1.** A bend in rock strata or in any planar feature. The feature (e.g. *\*bedding*, *\*cleavage*, or layering) is deflected sideways and the amount and direction of *\*dip* is altered. Four principal regimes are responsible for folding: *\*layer-parallel* or lateral compression; differential vertical subsidence; differential shearing; and *\*thrusting*. In a simple *\*anticline-syncline* fold pair, an individual fold consists of a curved *\*hinge* zone and two planar limbs. An imaginary *\*fold axis* lies parallel to the hinge zone (line) and marks the intersection of the *\*axial plane* (or surface) with this zone. This basic geometric form gives rise to many fold profiles, including *\*parallel*, *\*similar*, *\*concentric*, *\*open*, and *\*isoclinal* fold types. To define the attitude of a fold accurately, the orientation of both the hinge line and the axial plane have to be measured. Varying orientations of the hinge line and axial plane may give rise to widely differing fold attitudes, thus *\*vertical*, *\*upright* and *\*inclined* (horizontal and plunging, see PLUNGE), and *\*reclined* and *\*recumbent* forms may be described. See AMPLITUDE; WAVELENGTH. **2.** In *\*seismic* reflection sampling, the number of *\*offset* distances which sample one *\*common depth point* (CDP). For example, if one CDP is sampled at 24 offset distances it is referred to as '24-fold'

coverage. The signals recorded for the CDP on each separate trace are then summed by *\*stacking* to improve the signal-to-noise ratio.



## Fold

**fold-and-thrust belt** A linear or arcuate belt in which compression has produced a combination of *\*thrusts* and *\*folds*. The *\*dip* of the thrust planes decreases with depth, and the belt normally lies against the *\*foreland* of an *\*orogenic* belt. See also FOLD BELT.

**fold angle (inter-limb angle)** The angle between the limbs of a *\*fold*, whose tightness or openness reflects the intensity with which the structure has been deformed. The fold angle can be quantified by measuring the angle of intersection between two tangents through the *\*inflexion* points of opposite *\*fold limbs*.

**fold axis** A line which lies parallel to the *\*hinge* line and marks the intersection of the *\*axial plane* with the hinge zone.

**fold belt** A linear or arcuate region characterized by compressional tectonics, including folding. Some authors refer to '*\*fold-and-thrust belts*' in recognition of the importance of thrusting in crustal *\*shortening*.

**folding frequency** See NYQUIST FREQUENCY.

**fold limb** The generally planar region of a *\*fold* which lies between two adjacent *\*hinge* lines and is confined between the zones of maximum curvature.

**fold test** The main test in *\*palaeomagnetism* for determining the age of the *\*remanent* magnetization of rocks. Comparison of the scatter of directions of natural remanence from both sides of a *\*fold*, before and after correcting for the tectonic effects of the folding, enable the magnetization to be determined as pre-folding if the scatter is least after correction for folding.

**foliation** A continuous, sub-planar rock *\*fabric* formed by the preferred orientation of minerals with a generally *\*platy* or *\*tabular* habit. The layers are planar, and parallel but not necessarily to *\*bedding* or *\*cleavage*. The term 'foliation' is commonly applied to textures in high-grade *\*metamorphic rocks* (see METAMORPHIC GRADE).

**folivorous** Leaf-eating.

**Folk limestone classification** A widely used classification of *\*carbonates*, based on the type of particles and the nature and proportion of the *\*matrix* and/or *\*cement* present. In his original classification Robert L. Folk defined three main components to *\*limestones*. These are *\*allochems*, comprising various *\*grains* and particles; *\*micrite* (*microcrystalline \*calcite* mud matrix); and *\*sparite* (*sparry calcite* cement). The main allochems are *\*bioclasts* ('bio-', see below), *\*pellets* ('pel-'), *\*intraclasts* ('intra-'), and *\*ooids* ('oo-'). In defining a limestone by the Folk classification the rock is named according to the nature of the material filling the spaces between the particles (i.e. micrite matrix or sparite cement), prefixed by an abbreviation to denote the main allochems present: bio- for bioclasts, pel- for pellets, oo- for ooids, and intra- for intraclasts. For example, a limestone comprising pellets in a mud matrix is called a 'pelmicrite', and a limestone comprising shell fragments with a sparry calcite cement is a '*\*biosparite*'. It is possible to combine the prefixes of several allochems where appropriate, e.g. 'oobiosparite'. Limestones which lack allochems, consisting only of micrite are termed 'micrites'. *\*Dismicrite* is used for micrites with fenestral cavities (see FENESTRAE). Organically bound limestones, e.g. *\*reef* rocks, or *\*stromatolites*, are termed *\*biolithites*. Folk subsequently modified his classification to include various carbonate textures. Under his textural scheme a limestone with varying proportions of bioclasts, mud matrix and cement would be classified as follows: micrite with less than 1% shell fragments = micrite; micrite with 1–10% shell fragments = fossiliferous micrite; micrite with 10–50% shell fragments = sparse biomicrite; micrite

with over 50% shell fragments = packed biomicrite. Limestone with micrite and sparry calcite together with shell fragments = poorly washed biosparite; poorly sorted shell fragments with sparry calcite cement = unsorted biosparite; well sorted shell fragments with sparry calcite = sorted biosparite; rounded and abraded shell fragments with sparry calcite cement = rounded biosparite. Similar textural terms apply for other allochems with micrite and/or sparite. See [BIOMICRITE](#); [INTRAMICRITE](#); [INTRASPARITE](#); [PELSPARITE](#); [OOSPARITE](#); [OOMICRITE](#).

**fondathem** A rock unit produced beneath a body of water and giving a seismic reflection.

**fondoform** A flat surface beneath a body of water that gives a seismic reflection.

**fool's gold** See [PYRITE](#).

**footwall** The *\*fault* block which lies below any inclined *\*fault* surface. Compare [HANGING WALL](#).

**foram** See [FORAMINIFERIDA](#).

**foramen** A pore or opening. Applied to various openings, e.g. the *\*pedicle foramen* in *\*Brachiopoda* through which the *\*pedicle* emerges. In cephalopods (*\*Cephalopoda*) the term is applied to the opening in the *\*septum* through which the siphuncular cord passes.

**foramen magnum** Opening at the posterior end of the skull through which the spinal cord passes.

**foramina** A series of openings connecting the various chambers of a foraminiferid (see [FORAMINIFERIDA](#)) *\*test*.

**foraminifera** See [FORAMINIFERIDA](#).

**Foraminiferida** (informally **foraminifera**, **foraminiferans**, **forams** (*pl.*); **foraminiferid**, **foraminifer**, **foraminiferan**, **foram** (*sing.*) (**phylum Rhizopoda**) Order (in some classifications, subclass **Foraminifera**) Testate, amoeboid *\*protozoa* in which the cell is protected by a *\*test*, consisting of one to many chambers, whose structure and composition is of great importance in foraminifera classification. The three main types are: (a) most primitively, a test wall composed of a secreted, chitinous-like,

organic material called tectin, which also often forms an underlying layer in the other two types; (b) a test formed from agglutinated *\*sedimentary* particles, which may be cemented with an organic, calcareous, or ferric oxide *\*cement*; (c) a fully mineralized test, composed of secreted calcareous or siliceous minerals, of which the calcareous types (*\*aragonite* and *\*calcite*) are the most common. The arrangement of multiple chambers may be linear, spiral, cone-like, etc. Numerous fossil foraminifera are known, usually less than 1 mm across; though some, like the fusulinids (*\*Carboniferous* to *\*Permian*) and *\*nummulitids* (*\*Eocene* to *\*Oligocene*) were appreciably larger (some measured up to 100 mm in diameter). All species live in marine environments. Agglutinated forms predominated in the *\*Cambrian* and *\*Ordovician*, presumably derived from a tectinous ancestor, while forms with fully mineralized tests appeared in the Ordovician and diversified greatly in the *\*Devonian*. The Foraminiferida are important zonal fossils, and some *\*planktonic* varieties can be used for stratigraphic correlation on virtually a worldwide scale. See *GLOBIGERINA OOZE*.

**foramol** Applied to an association of *\*Bryozoa*, *\*Foraminiferida*, coralline red algae (*\*Rhodophyceae*), and *\*Mollusca* that inhabits seas where the temperature often falls below 15 °C and forms a characteristic sediment. Compare *CHLORALGAL* and *CHLOROZOAN*.

**forams** See *FORAMINIFERIDA*.

**forced convection** Mechanical turbulence, with the development of eddies, in air flowing over an uneven surface.

**forced regression** *\*Regression* caused by a fall in sea level.

**forced regressive systems tract** See *REGRESSIVE SYSTEMS TRACT*.

**fore-arc** See *ARC-TRENCH GAP*.

**fore-arc basin** The part of the fore-arc (*\*arc-trench* gap) adjacent to the *\*island* arc which is characterized by flat-lying sediments, in contrast to the highly deformed *\*accretionary wedge* adjacent to the oceanic *\*trench*. Fore-arc basins lie behind the topographic high-point of the wedge, which in places forms an outer (sedimentary) island arc. Fore-arc sediments lie unconformably on accretionary wedge material and show progressive

shoaling. The source of the material is the adjacent volcanoes and the erosion of uplifted **\*plutonic**–metamorphic basement.

**forebulge** An area of increased elevation, resulting from flexing of the **\*lithosphere**, that develops some distance from a shoreline that is depressed by the weight of an ice sheet.

**foredeep (toe-trough)** A basin adjacent to a **\*craton** which is filled with a thick accumulation of sediment derived from an **\*orogenic** belt during uplift. The sediments are typically non-marine to shallow-marine and commonly suffer deformation within a few million years of deposition.

**foredune** A crescent-shaped **\*sand dune** with arms that extend downwind on either side of an obstruction. It results from the accumulation of sand on the upwind side of the obstruction.

**foreland** A stable area on the edge of an **\*orogenic** belt; a foreland is usually on the margin of a **\*craton** and is underlain by **\*continental crust**. Many forelands have suffered warping during **\*orogeny** and also carry a superficial **\*fold-and-thrust** belt. The major direction of movement in an orogenic belt is towards the foreland. Where the orogenic belt lies between two stable areas, the other is called the hinterland.

**foreland basin (flexural basin)** A basin that forms by the downward flexing of the **\*lithosphere** in response to the weight of an adjacent mountain belt in a region of compressional tectonics. There are two types: **\*peripheral** foreland basins and **\*retro-arc** foreland basins.

**foreland thrust belt** A region of a **\*thrust** belt in which the thrust faults are generally clearly discernible and affect rocks that can be identified in the adjacent **\*foreland**.

**fore reef** **\*Talus** slope on the seaward side of a **\*reef**, constantly under attack by waves and currents.

**foreset** **1.** The inclined surface within a cross set (see **CROSS-STRATIFICATION**) produced by the forward movement of the slip-face of a **\*ripple**, **\*dune**, **\*sand wave** or bar. **2.** The slip-face of a **\*Gilbert-type** delta.

**foreshock** A small **\*earthquake**, sometimes occurring in swarms, that precedes a major earthquake (or volcanic eruption).

**foreshore** Lower shore zone that lies between the normal high- and low-water marks. The foreshore may either be a plane slope dipping seawards at a low angle, or be marked by the development of longshore bars (ridge-and-runnel topography), depending on the nature of the wave attack.

**foreshortening** In *\*radar* terminology, the distortion of a radar image caused by shallow surface angles relative to the radar wavefront angle, such that the base of a hill is actually closer to the radar than the top. The resulting image gives the appearance of a shorter slope. *Compare* LAYOVER.

**forked lightning** Lightning discharge in which luminous branches from the main channel are seen. *See also* SHEET LIGHTNING.

**form** The overall shape of a crystal. If it is able to grow freely, a crystal develops with a regular pattern of *\*crystal* faces and interfacial angles which are characteristic of a particular *\*mineral*. The study of this regularity of crystal form, and of the internal structure to which it is related, is called '*\*crystallography*'.

**formal** The name of a *\*stratigraphic unit* or *\*taxon* is formal when it has been established according to the conventional principles of nomenclature and is being used in the sense of a proper name, e.g. *\*Barremian* stage, *\*Osteichthyes* (bony fish). In formal use the initial letter is often capitalized. *Compare* INFORMAL.

**formation** The fundamental unit used in *\*lithostratigraphy*. Specific features distinguish one rock formation from another. The thickness of the formation is unimportant in its definition, as a given formation may vary within different outcrops. Formations may be subdivided into members and together several formations constitute a group.

**formation age** Time which has elapsed since a *\*meteorite* formed, obtained by *\*radiometric* dating, assuming no loss of a gaseous daughter isotope has occurred (*see also* GAS-RETENTION AGE). Most meteorites give formation ages around 4.5 billion years, although there are exceptions (*see* SHERGOTTYTE/NAKHLITE/CHASSIGNITE METEORITES).

**formation evaluation** The detailed analysis and interpretation of borehole data, drilling results, geophysical down-hole logs, etc., to determine the physical characteristics of the rock *\*formations* through which the drill has

penetrated. This is done mainly to ascertain whether or not economic reserves of hydrocarbons are present and, if they are, to determine the most economical and efficient way to extract them. Formation evaluation is an important component of reservoir engineering design.

**formation velocity** ( $v_{\text{for}}$ ) The uniform seismic velocity of a particular homogeneous rock type. For a rock thickness  $h$  and a single-travel time  $t$ ,  $v_{\text{for}} = h/t$ . Compare [INTERVAL VELOCITY](#).

**form factor** See [DRAINAGE BASIN SHAPE INDEX](#).

**form-genus** Non-phylogenetic, artificial [\\*taxon](#) of convenience. In palaeobotany the disarticulated parts of fossil plants, whose natural affinities are unknown, have been assigned to form-genera on the basis of similarities in morphology. Later discoveries may show different form-genera to have derived from the same plant, e.g. [\\*Stigmara](#)-type roots and [\\*Lepidodendron](#)-type bark. See also [CORDAITALES](#); [CALAMITES CISTIIFORMES](#).

**FormoSat-3 (COSMIC; Constellation Observing System for Meteorology, Ionosphere and Climate)** An international project between the National Space Program Office of Taiwan and the University Corporation for Atmospheric Research of the USA to operate a constellation of six identical [\\*microsatellites](#) that collect atmospheric remote-sensing data for operational weather prediction and research into climate, space weather, and [\\*geodesy](#). The microsatellites were launched together on 15 April 2006, from California.

**form roughness** See [BED ROUGHNESS](#).

**form set** See [CROSS-STRATIFICATION](#).

**forsterite** See [OLIVINE](#).

**FORTE** See [FAST ON-ORBIT RECORDING OF TRANSIENT EVENTS](#).

**Fortin barometer** Portable mercury barometer that requires the accurate setting of the mercury level at a fixed point (scale zero, see [FIDUCIAL POINT](#)). The reading of the mercury height is then taken by adjusting a vernier scale to the top of the mercury column. See also [KEW BAROMETER](#).

**Förtsch discontinuity** An irregular seismic discontinuity within the upper **\*continental crust**, at 8–11 km depth, usually interpreted as a change from upper granitic rocks to deeper dioritic composition. *See also* **CONRAD DISCONTINUITY**; **MOHOROVIČIĆ DISCONTINUITY**.

**forward problem (direct problem, normal problem)** The problem of calculating what should be observed for a particular model, e.g. calculating the gravity anomaly that would be observed for a given model of a salt **\*dome**. *Compare* **INVERSE PROBLEM**.

**fossa** (*pl.* **fossae**) A linear depression on a planetary surface, closely analogous to a terrestrial fault-bounded depression or **\*graben**. The radial fractures with mainly **\*normal faults** (e.g. Tempe, Tantalus Menonia, Claritas fossae, etc.) which surround the Tharsis bulge on Mars, form type examples.

**fossil** **1.** Generally, anything ancient, especially if it is discovered buried below ground (e.g. **\*fossil fuel**, fossil soil). **2.** The remains of a once-living organism. Formerly, only the preserved remains of an organism that lived prior to the end of the last glacial period was called a fossil, i.e. fossils were older than 10 000 years, and remains younger than 10 000 years were called subfossils. That limitation has now been abandoned and any ancient remains of a living organism can be called a fossil, regardless of its age. The term includes skeletons, tracks, impressions, trails, borings, and casts. Fossils are usually found in consolidated rock, but not always (e.g. woolly mammoths living 20 000 years ago were recovered from the frozen tundra of Siberia). In its original sense, fossil meant anything dug up from the earth, including ores, precious stones, etc. The modern use of the word dates from the late 17th century. *See also* **LIVING FOSSIL**; **TRACE FOSSIL**.

**fossil fuel** All deposits of organic material capable of being burnt for fuel; chiefly coal, oil, and gas. These are formed under pressure by alteration or decomposition of plant or animal remains.

**fossiliferous micrite** *See* **FOLK LIMESTONE CLASSIFICATION**.

**fossilization** The process by which a **\*fossil** is formed. It is unusual for organisms to be preserved complete and unaltered; generally, the soft parts decay and the hard parts undergo various degrees of change. **\*Solution** and other chemical action may reduce the tissues to a thin film of carbon; this

process is called 'carbonization'. The organism may be flattened by the compaction of sediments to form compressions. Porous structures, e.g. bones and shells, may be made more dense by the deposition of mineral matter by *\*groundwater*; this process is called 'permineralization' or 'petrification'. The internal physical structures of some shells may be changed as a result of solution and reprecipitation; in this process ('recrystallization') the original structure may be blurred or lost. Many shells which were originally composed of *\*aragonite* are recrystallized into the more stable mineral *\*calcite*. The solution of an original shell and the simultaneous deposition of another mineral material constitutes 'replacement'; this may occur molecule by molecule, in which case the microstructure is preserved, or *en masse*, where it is not. Common replacement minerals include *\*silica* or iron sulphide, but there are many others. The impression of skeletal remains in surrounding sediments constitutes a 'mould'. Where the external structures are preserved it is called an 'external mould' and where the internal features are preserved it is called an 'internal mould' or 'steinkern'. Filling of a mould cavity by mineral matter may produce a 'natural cast'. Tracks, trails, burrows, and other evidence of organic activity may also be preserved. These are called 'ichnofossils' or *\*trace fossils*.

**fossil-Lagerstätte** See LAGERSTÄTTE.

**fossula** A gap or depression in the floor or *\*calyx* of a rugose coral (*\*Rugosa*). It may be formed by the absence of septa (see SEPTUM) in a particular part of the calyx.

**foundation** Lowest part of a structure, below ground surface and in contact with natural earth materials, which transmits *\*load* to the soil or rock. In a dam the foundation may include the valley floor and abutments.

**founder effect (peripatric speciation)** The derivation of a new population (e.g. on an oceanic island) from a single individual or a limited number of immigrants. The founder(s) represent a very small sample of the *\*gene pool* to which it or they formerly belonged. *\*Natural selection* operating on this more restricted genetic variety yields gene combinations different from those found in the ancestral population.

**founder lineage** In *\*phylogenetics*, an ancestral lineage, often still extant, from which other lineages have risen. The term is usually applied to

intraspecific studies of populations and used to describe operational taxonomic units that occur at *\*internal nodes* of a *\*phylogenetic tree*.

**fourchite** An *\*intrusive \*igneous* rock consisting of essential titanaugite (titanium-rich *\*augite*) and *\*kaersutite* (with or without *\*biotite*) set in a light-coloured base of *\*analcime* or *\*glass*. This rock type is a member of the alkali *\*lamprophyre* group of rocks, which includes *\*camptonites*, *\*monchiquites*, and alnoites.

**Fourier analysis** The method whereby any periodic function can be broken down into a convergent trigonometric series of the form  $f(x) = a_0/2 + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx)$ , where  $a_n$  and  $b_n$  are constant coefficients. Fourier analysis is the process of determining the *\*frequency-domain* function from a time function (e.g. a seismic-trace wave-form). *See also* **FOURIER TRANSFORM**.

**Fourier synthesis** The superimposition of sinusoidal waves of known frequency, amplitude, and phase to represent an observed wave-form. It is the process of determining the *\*time-domain* function from a frequency function. *See also* **FOURIER TRANSFORM**.

**Fourier transform** The mathematical formulae by which a time function (e.g. a seismic trace) is converted into a *\*frequency-domain* function and vice versa. *See also* **FOURIER ANALYSIS**; **FOURIER SYNTHESIS**.

**fracking** *See* **HYDROFRACTURING**.

**fractal** A geometric entity which has a basic pattern that is repeated at ever decreasing sizes. Fractal patterns are not able to fill spaces and are hence described as having fractal dimension. Fractals occur frequently in nature, such as in forked lightning or in chaotic systems (*see* **CHAOS**).

**fractional crystallization** The removal of early formed *\*crystals* from an originally homogeneous *\*magma* (for example, by gravity settling) so that these crystals are prevented from further reaction with the residual *\*melt*. The composition of the remaining melt becomes relatively depleted in some components and enriched in others, resulting in the precipitation of a sequence of different minerals. Fractional crystallization is one of the main processes of *\*magmatic* differentiation.

**fractionation** See MAGMATIC DIFFERENTIATION.

**fracto-** See FRACTUS.

**fractography** The study of fracture surfaces.

**fracture** 1. General term applied to any break in a material, but commonly applied to more or less clean breaks in rocks or minerals that are not due to \*cleavage or \*foliation. 2. See VOIDS.

**fracture cleavage** A \*cleavage (defined originally by Leith in 1905) which resembles very closely spaced, parallel \*joints or fractures, but where in fact the 'fractures' are due to a loss of material resulting from \*pressure solution. In modern terminology the term would be replaced by 'spaced cleavage'.

**fracture porosity** A form of \*secondary porosity generated by tectonic fracturing (see TECTONISM) of the rock. Such porosity can develop in any rock, allowing the development of productive \*reservoirs in rocks such as \*granites and \*gneisses. See POROSITY; CHOQUETTE AND PRAY CLASSIFICATION; RESERVOIR ROCK.

**fracture spacing index ( $I_f$ )** The number of fractures in a one-metre length of drill \*core.

**fracture zone** A linear feature on the deep-sea floor across which the \*lithosphere changes abruptly in both age and water depth. Most fracture zones transect oceanic \*ridges and are small circles whose radii of curvature depend on the distance from the \*pole of rotation of the two lithospheric \*plates which are diverging at the ridge. Many fracture zones also contain deep-sea basins.

**fractus (fracto-)** From the Latin *fractus* meaning 'broken', a species of cloud that has an irregular or ragged form. The term is applied to \*cumulus and \*stratus, i.e. fractocumulus, fractostratus. Fractus is either torn from the parent cloud or remains as fragments after the parent cloud has dissipated. See also CLOUD CLASSIFICATION.

**fragipan** Subsoil \*horizon, found deep in a \*soil profile and having a high bulk density. It is a dense, brittle, and compact layer, apparently with little or no cementation horizon, associated with \*acid soil conditions.

**fragmental** See CLASTIC.

**framestone** An *\*autochthonous*, organically bound *\*limestone*, where the organisms, e.g. corals (*\*Anthozoa*), form a rigid framework during deposition. See EMBRY AND CLOVAN CLASSIFICATION.

**framework grains** *\*Detrital* fragments, 0.065–2.00 mm in size, that make up the bulk of a *\*sandstone*.

**framework porosity** See CHOQUETTE AND PRAY CLASSIFICATION.

**Francisco (Uranus XXII)** A lesser satellite of *\*Uranus* with a radius of 11 km.

**Franconian** See FESTINOPIAN.

**franklinite** A member of the *\*spinel* group of minerals, and an *\*end-member* of the *\*magnetite* series, with composition  $ZnFe^{3+}_2O_4$ , although appreciable amounts of  $Mn^{2+}$  and  $Fe^{2+}$  are also present; sp. gr. 5.0; *\*hardness* 6; black; *\*metallic lustre*; *\*crystals cubic* but normally found as octahedra or granular aggregates; occurs at Franklin, New Jersey, in association with other zinc minerals in a metamorphosed *\*limestone*.

**Frasien** See FRASNIAN.

**Frasnian (Frasien)** A *\*stage* in the Late *\*Devonian* epoch, preceded by the *\*Givetian* (Middle Devonian), followed by the *\*Famennian*, and dated at 385.3–374.5 Ma ago (Int. Commission on Stratigraphy, 2004). The stage is noted for the culmination of the first major radiation of the single-celled *\*Foraminiferida* and for the origin of important groups of *\*goniatites*, prolecanitids, goniatitids, and clymeniids. It is roughly contemporaneous with the *\*Senecan* (N. America) and part of the *\*Condobolinian* and *\*Hervyan* (Australia).

**frazil ice** Flowing water ice that forms platelets rather than continuous sheets; often observed in Canadian rivers. The name is derived from the French *fraisil*, meaning cinder.

**free-air anomaly** The *\*gravitational* acceleration remaining after correction of a measurement for the theoretical gravity, usually the *\*International Gravity Reference Field*, and by the *\*free-air correction*. No correction has

been made for the gravitational attraction of rocks between the observing station and the reference datum, which is usually sea level.

**free-air correction (Faye correction)** The correction applied to a measurement of gravity which allows for the variation of gravity with height above a reference level, usually sea level. This correction assumes there is only air between the station and the reference level, and it is 0.3086 mgal/m. *See also* [BOUGUER CORRECTION](#).

**free atmosphere** The atmosphere above the [\\*planetary boundary layer](#) (commonly taken as above about 500 m).

**freeboard** The distance between the maximum permitted water level in the reservoir behind a dam and the top of the dam wall. *See also* [CONTINENTAL FREEBOARD](#).

**free dune** A [\\*sand \\*dune](#) that is shaped entirely by the action of the wind.

**free face** *See* [SLOPE PROFILE](#).

**free oscillations** The harmonics at which any body, e.g. the Earth, tends to vibrate most freely, i.e. resonates. There are two fundamental types: torsional (vibration with motions perpendicular to the Earth's radius); and spheroidal (vibrations that are both radial and tangential to the Earth's surface). The study of such resonances, e.g. those induced by major earthquakes, provides information on the internal nature of the Earth. A major earthquake can make the entire globe vibrate or ring like a bell, and some earthquakes have been so large that sensitive [\\*seismometers](#) have continued to record the oscillations for weeks after the event. The decay of the vibrations gives valuable information about the elastic layering of the Earth, and especially of the [\\*low-velocity zone](#). [\\*Moonquakes](#) produce similar phenomena.

**freestone** A building stone; poorly jointed [\\*sandstone](#) or [\\*limestone](#) which can be worked easily in any direction. *Compare* [DIMENSION STONE](#).

**freezing nuclei** Any nuclei, commonly ice crystals but sometimes the suitably shaped crystals of other substances, that when present in clouds at temperatures below 0 °C will cause any supercooled droplets with which they collide to change to ice (in the form of a crystalline growth upon the nucleus). *See also* [NUCLEUS](#). *Compare* [ICE NUCLEUS](#).

**freibergite** See TETRAHEDRITE.

**frequency (f)** The number of complete \*wavelengths which pass a given point in a specified time; units are hertz (Hz; one hertz is one cycle per second). The frequency of a periodic wave-form is given by  $f = 1/T$ , where  $T$  is the \*period; and by  $f = v/\lambda$ , where  $v$  is the velocity and  $\lambda$  the wavelength.

**frequency domain** A reference framework in which measurements are related to frequency rather than to time (as in the \*time domain).

**fresh water** Water containing little or no chloride ion. According to the Venice system, which classifies brackish waters by their percentage chloride content, fresh water contains 0.03% or less of chloride. Compare HALINITY.

**freta** See FRETUM.

**Fretted and Chaotic Hummocky Terrain** See MARTIAN TERRAIN UNITS.

**fretum** (*pl. freta*) On the surface of \*Titan, a narrow area of liquid that connects two larger areas of liquid.

**friable** Applied to the consistency or handling properties of soil, meaning that the soil crumbles easily.

**friction** A force that resists the movement of one body in relation to another body with which it is in contact. Where one body rests upon another there is a contact force ( $R$ ) between them acting at right angles to the contact surface. If a force  $B$  acting parallel to the contact surface is applied to one of the bodies, a force  $F$  will resist it. If  $B$  increases until the bodies move against each other,  $F$  will also increase until it reaches a maximum ( $F_L$ ).  $F_L = \mu_g R$ , where  $\mu_g$  is the coefficient of static friction. Once the two bodies are moving against each other at a constant velocity,  $F$  will fall to a value  $F_k = \mu_k R$ , where  $\mu_k$  is the coefficient of kinetic friction.

**frictional angle** See ANGLE OF INTERNAL FRICTION.

**frictional resistance, angle of** See ANGLE OF SHEARING RESISTANCE.

**frigid** See PERGELIC.

**frigid zone** In the climate classification devised by Aristotle, the region in either hemisphere between the pole and the Arctic or Antarctic Circles. In his view, this zone was uninhabitable for humans. *Compare* TEMPERATE ZONE; TORRID ZONE.

**fringing reef** *See* REEF.

**Fronian** A \*stage (437.5–436 Ma ago) of the Early \*Silurian, underlain by the \*Idwian and overlain by the \*Telychian.

**front** Boundary or boundary region separating \*air masses of different origins and characteristics. Temperature gradients in any horizontal surface are large through the front. Different types of front are distinguished according to the nature of the \*air masses separated by the front, the direction of the front's advance, and the stage of development. The term was first devised during the First World War by the Norwegian school of meteorologists (headed by Professor V. \*Bjerknes). *See also* ANAFRONT; COLD FRONT; WARM FRONT; KATAFRONT; OCCLUDED FRONT; POLAR FRONT.

**frontal arc** At a \*convergent margin, the region of the overriding plate that is nearest to the \*trench and ahead of the \*volcanic arc. Islands situated along the frontal arc experience rapid uplift, in some cases (e.g. Nias on the Sunda frontal arc) of up to 1 mm per year.

**frontal fog (precipitation fog)** \*Fog that forms along a \*front, where warm air is being lifted above cooler air. \*Stratus cloud forms in the rising warm air and produces precipitation that falls through the front into the dry air below, where it evaporates. The air is also cold, however, so the water vapour condenses into droplets, producing the fog.

**frontal wave** Wave-like deformation of the line of a \*front between two \*air masses. The wave develops from the northward incursion of warm air and usually travels along the front, with colder air ahead and to the rear. Typically, frontal waves occur in sequences, or 'families', of several waves, and develop into \*depressions or storm centres travelling more or less eastward as 'secondaries' along the extended cold front to the rear of the original low. The secondaries tend to catch up and merge with the original depression as it slows up in its fully developed stage.

**frontal zone** Transition zone, sometimes amounting to a discontinuity, that separates adjacent **\*air** masses, and where the temperature gradient is strongest. Some turbulent mixing takes place. The sloping zone separating a cold wedge under warm air typically extends about 1 km vertically and about 100–200 km horizontally.

**frontogenesis** Development and intensification of frontal boundaries between adjacent **\*air** masses.

**frontolysis** Processes of dissolution or dissipation of a **\*front**. Frontal decay results when different **\*air** masses stagnate together, or move together or in succession along the same track at the same speed, or incorporate air of the same temperature.

**frost** Condition in which the prevailing temperature is below the freezing point of water (0 °C). This may lead to a deposit of ice crystals on objects, e.g. grass or trees. Such deposits result from condensation when the **\*dew-point** temperature is below freezing. *See also* **BLACK ICE**.

**frost boil** *See* **INVOLUTION**.

**frost heave (frost heaving)** Upward movement of the ground surface or of individual particles, due to the formation of lenses of ice up to 30 mm thick in the **\*regolith**. It reaches its maximum in silt-dominated material, in which the greatest volume of ice may develop (more than 68% ice by volume). When the total uplift of the surface is measured, it is found to be approximately equal to the sum of the thicknesses of the layers of ice. Surface stones may be heaved by the development of needle-ice columns ('pipkrakes').

**frost heave test** A laboratory test in which **\*aggregate** or soil is frozen under controlled conditions. A cylinder containing rock aggregate, 150 mm high and 100 mm in diameter is placed in freezing conditions with its base in running water for 250 hours. The **\*frost** heaving must be less than 12 mm.

**frost heaving** *See* **FROST HEAVE**.

**frost hollow** Area (e.g. a valley bottom or a smaller hollow) that is very liable to severe and frequent **\*frosts** as a result of dense, cold air moving

downslope (katabatic flow) and collecting there under conditions of radiation cooling (e.g. at night).

**frost pull and frost push** \*Periglacial processes that bring about the upward migration of \*clasts through the \*regolith. Frost push takes place when an ice lens forms beneath a clast and so pushes it upwards. Frost pull occurs when a clast adheres to ice within a freezing regolith and so is drawn upwards as the ground heaves.

**frost-shattering** See FROST WEDGING.

**frost smoke** See ARCTIC SEA SMOKE.

**frost table** See TJAELE.

**frost wedging (conglifraction, frost-shattering, gelifraction, gelivation)** Fracturing of rock by the expansionary pressure associated with the freezing of water in planes of weakness or in pore spaces.

**Froude number (Fr)** A dimensionless number equal to the ratio of water velocity to the speed of a gravity wave, used to assess whether flow in an open channel is critical, tranquil, or shooting. If the Froude number is less than 1, flow is said to be subcritical or slow; if  $Fr = 1$ , flow is critical; and if  $Fr$  is greater than 1, flow is fast or supercritical.

**frustule** Silica wall of a diatom. See BACILLARIOPHYCEAE.

**Fs** Orthoferrosilite. See ENSTATITE.

**fugacity (f)** A measure of the tendency of a gas to escape or expand, used in calculations of chemical equilibrium. Fugacity ( $f_i$ ) is the pressure value needed at a given temperature to make the properties of a non-ideal gas satisfy the equation for an ideal gas, i.e.  $f_i = \gamma_i P_i$ , where  $\gamma_i$  is the fugacity coefficient, and  $P_i$  is the partial pressure for the component  $i$  of the gas. For an ideal gas,  $\gamma_i = 1$ .

**fugichnia** The so-called 'escape structures' that mark the response of animals to changes in the rate of deposition or erosion or to predation. The original trace (see TRACE FOSSIL) may be regarded as a permanent dwelling structure, but the presence of fine, crescentic laminae (\*spreiten) indicates

that the organism made efforts to escape burial or exposure. Thus the presence of spreiten implies fugichnia.

**Fujita Tornado Intensity Scale** A standard six-point scale for reporting the intensity of a \*tornado by inferring its wind force from the type and extent of the damage it caused. The scale was introduced in 1971 by Tetsuya Theodore Fujita and Allen Pearson. It has since been replaced by the \*Enhanced Fujita scale. See WIND STRENGTH; APPENDIX C.

**fulgurite** An elongated piece of \*glass produced when energy released by a lightning strike melts rock which then fuses and solidifies as it cools. Fulgurites can form in almost any type of rock, but the largest occur in loose \*sand. They can be thought of as fossilized lightning strikes.

**fuller's earth** 1. A clay consisting mainly of expanding \*smectites such as \*montmorillonite used industrially for its absorptive properties. 2. Capitalized, Fuller's Earth is the stratigraphic name of a \*Jurassic clay formation outcropping in southern Britain.

**fulvic acid** Mixture of uncoloured organic acids that remains soluble in weak acid, alcohol, or water after its extraction from soil.

**fumarole** Vent in a volcanically active area that emits steam, gas (SO<sub>2</sub>, CO<sub>2</sub>, etc.), or other \*volatile constituents at high temperatures (from 100 °C to 1000 °C). The fumarolic condition has been thought to indicate a late stage in volcanic activity, but may actually precede volcanic \*eruptions, e.g. Mt St Helens, Cascade Range, Washington, in 1980.

**fumarolic stage** See FUMAROLE.

**functional morphology** The attempted interpretation of the functions of particular organs or structures that occur in various \*fossils. In some cases it is difficult to interpret function in groups that have no living examples, but it is often possible to analyse growth and form in invertebrate groups and to relate this growth both to the biology of the organism and to the environment.

**fundamental form** See PARAMETRAL PLANE.

**fundamental strength** The maximum \*stress a material can sustain indefinitely at a given temperature and \*confining pressure. The

fundamental strength is always less than the breaking strength and **\*ultimate** strength.

**Fungi** One of the three multicellular kingdoms, along with the **\*Plantae** (plants) and **\*Animalia** (animals). Although resembling plants, Fungi feed by ingesting organic matter, whereas plants are autotrophic and require only inorganic substances as nutrients. As fungi generally lack hard parts they are rarely found as fossils, but thread-like representatives have been found in **\*Precambrian** rocks. Fungi probably left the sea about 400 million years ago, when the first plants colonized the land.

**funnel cloud** Cloud produced in a low-pressure vortex in the centre of a spiral storm. The funnel may extend below the cloud in which it originates, becoming visible because the low pressure inside it causes water vapour to condense. If the funnel touches the surface it becomes a tornado or waterspout.

**funnelling** The constraining of an airflow by valleys, leading to higher wind speed, convergence, and uplift. Similar effects occur in the air between an advancing front and the face of a mountain barrier.

**furious fifties** Popular maritime term for the prevailing westerly winds which are commonly strong over the oceans in temperate latitudes of the southern hemisphere, particularly between about 50° S and 60° S. *See also* ROARING FORTIES; SHRIEKING SIXTIES.

**fusain** Fossil charcoal, a **\*coal** lithotype sometimes called 'mother-of-coal', produced by the burning of plant material under airless conditions. This converts the material to almost pure carbon and can preserve small plant parts and cellular structures.

**fusiform** Spindle-shaped; elongated with tapering ends.

**fusinite** *See* COAL MACERAL.

**fusion** **1.** Generally, the melting of a solid substance by heat. **2.** In nuclear fusion, the combining of two light atomic nuclei to form a heavier nucleus with the sudden release of energy, e.g. in the hydrogen bomb.

**fusuline** (*adj.* **fusulinid**) One of the so-called 'larger foraminifera' (**\*Foraminiferida**) which usually have a **\*fusiform** or discoid shape. Many genera are differentiated by rapid development and evolution, making them

important *\*index fossils*, particularly in *\*Carboniferous* and *\*Permian* rocks.

**Fuvelian** A regional *\*stage* (77–70.6 Ma ago) of the Late *\*Cretaceous*, preceded by the *\*Valdonian* and followed by the *\*Maastrichtian*.

**FY** See FENGYUN.

# G

**G** See GAUSS.

**G** See GRAVITATIONAL CONSTANT.

**g** See GRAVITATIONAL ACCELERATION.

**gabbro** A coarse-grained, *\*basic \*igneous* rock, consisting of *\*essential* calcium-rich *\*plagioclase* feldspar (approximately 60%), *\*clinopyroxene* (*\*augite* or titanaugite), and *\*orthopyroxene* (*\*hypersthene* or *\*bronzite*), plus or minus *\*olivine* with *\*accessory \*magnetite* or *\*ilmenite*. Gabbros result from the slow crystallization of *\*magmas* of basaltic composition, and like the *\*basalts* they can be divided into tholeiitic and alkali types (*compare* ALKALI BASALT; THOLEIITE). Tholeiitic gabbros are characterized by the presence of two pyroxene types (augite and hypersthene) and *\*interstitial* silica-rich *\*glass*, whereas alkali gabbros are characterized by one calcium/titanium-rich pyroxene (titanaugite) and scattered interstitial *\*feldspathoid* minerals. Many large gabbroic *\*intrusions* display mineral layering, testifying to the complex processes taking place within basic *\*magma chambers*. Gabbros are commonly found intruded as ring complexes (e.g. Ardnamurchan and Skye in Scotland), large *\*lopoliths* (Bushveld complex, S. Africa), or layered complexes (Skaergaard in eastern Greenland being the most famous).

**gaging station** See GAUGING STATION.

**gahnite** See SPINEL.

**Gaian hypothesis** Hypothesis, formulated by James E. Lovelock and Lynn Margulis, that the presence of living organisms on a planet leads to major modifications of the physical and chemical conditions obtaining on the

planet, and that subsequent to the establishment of life the climate and major **\*biogeochemical** cycles are mediated by the living organisms themselves.

**gaining stream** A stream that receives water emerging from a submerged spring or other groundwater seepage which adds to its overall flow.

**gal** The unit, named after Galileo, for measuring **\*gravitational** acceleration. 1 gal = 1 cm/s<sup>2</sup>. The gal has been largely replaced by the **\*gravity unit**.

**galactic cosmic rays** See COSMIC RADIATION.

**Galápagos Rift** A spreading **\*mid-ocean ridge** between the **\*Cocos** and **\*Nazca Plates**, that forms a **\*rift valley** between latitudes 86 °W and 89 °W, where there is a large area of **\*hydrothermal vents**.

**Galápagos Rise** The oceanic **\*ridge** between the **\*Cocos** and **\*Nazca Plates**.

**Galassia** A 2-unit **\*CubeSat** mission, expected to last 6 months, developed by students at the National University of Singapore, to measure the total electron count in the **\*ionosphere** and to acquire data related to quantum-based communication. The **\*nanosatellite** was launched on 16 December 2015, from Satish Dhawan Space Centre, India.

**Galatea (Neptune VI)** A satellite of **\*Neptune**, with a diameter of 158 km; visual albedo 0.06.

**galaxite** See SPINEL.

**gale** Wind blowing at more than 30 knots (17 m/s).

**galena** Mineral, PbS; sp. gr. 7.4–7.6; **\*hardness** 2.5; **\*cubic**; lead-grey; lead-grey **\*streak**; **\*metallic \*lustre**; **\*crystals** cubes or octahedra, and often octahedral twins; **\*cleavage** perfect cubic {100}; widely distributed in **\*hydrothermal** veins and **\*syngenetic** exhalative deposits and as a replacement in **\*limestones** and dolomitic rocks, associated in veins with **\*sphalerite**, **\*pyrite**, **\*chalcopyrite**, **\*barite**, **\*quartz**, **\*fluorite**, and **\*calcite**.

**Galilean satellites** The four classical **\*satellites** **\*Io**, **\*Europa**, **\*Ganymede**, and **\*Callisto** (in order outwards from Jupiter) which were discovered in 1610 by Galileo. The observation that they orbited a body other than the

Earth was fatal to the Ptolemaic **\*cosmology**. Ganymede (radius 2638 km) is the largest satellite in the **\*solar** system, larger than both Mercury and Pluto. Europa (radius 1536 km), the smallest of the four, is a little smaller than the Moon (radius 1738 km). They occupy equatorial orbits. There is a regular decrease in density from Io ( $3550 \text{ kg/m}^3$ ) to Callisto ( $1830 \text{ kg/m}^3$ ), Callisto preserving one of the most heavily cratered surfaces of any **\*satellite**.

**Galileo** A **\*NASA** spacecraft, launched on 15 October 1989 from the shuttle *Atlantis* on an international deep-space mission, that passed and photographed Venus on 9 February 1990, for a gravity assist. On its journey through the asteroid belt, Galileo encountered **\*Gaspra** on 29 October 1991, and **\*Ida** on 28 August 1993, where it discovered the satellite Dactyl. It reached **\*Jupiter** in December 1995, then entered an orbit that brought it into repeated encounters with the **\*Galilean** satellites. The spacecraft comprised an orbiter and a probe, released in July 1995, as Galileo was still approaching Jupiter, which penetrated the atmosphere and returned data for 61.4 minutes, by which time it had descended to a level where the pressure was 24 bars, 140 km below the 1 bar pressure altitude. The Galileo mission ended on 21 September 2003, when the spacecraft plunged into the jovian atmosphere.



<http://solarsystem.nasa.gov/galileo/>

- A NASA mission to study Jupiter and its moons.

**Gallic** The middle **\*epoch** of the **\*Cretaceous** period, dated at 130–89.3 Ma ago (Int. Commission on Stratigraphy, 2004), preceded by the **\*Neocomian** and followed by the **\*Senonian**.

**gamma** A unit of magnetic field strength.  $1 \text{ gamma} = 10^{-5} \text{ gauss}$ . It has now been replaced by SI units:  $1 \text{ gamma} = 10^{-9} \text{ tesla}$  (i.e. 1 nano tesla, nT).

**gamma–gamma sonde** A **\*well-logging** instrument package in which a source of gamma radiation, usually  $^{27}_{60}\text{Co}$  or  $^{137}\text{Cs}$ , bombards the wall of the **\*borehole**, and the backscatter of **\*gamma** rays, together with natural radiation, is recorded some 45 cm above the source. The record is known as a **\*density log**, as the backscatter is an exponential function of electron

density in the rocks. Mostly the instrument responds to the nearest 10 cm of rock wall. Sometimes two detectors are used at different locations to distinguish between **\*mud cake** and rock.

**gamma-ray log** See [GAMMA-RAY SONDE](#).

**gamma rays** Electromagnetic radiation, about  $10^{-10}$  to  $10^{-14}$  m in wavelength, similar to, but of shorter wavelength than X-rays, emitted by radioactive substances.

**gamma-ray sonde** The **\*well-logging** instrument, comprising a scintillometer, used to measure the natural radioactivity of the rocks through which the drill hole passes. Potassium ( $^{40}\text{K}$ ) is the most abundant radioactive element and occurs in **\*clays** (especially the mineral **\*illite** and **\*micas**). The record is a gamma-ray log, expressed in API units (a gravity scale devised by the American Petroleum Institute and applied mainly to measurements of crude oil). The log is particularly useful for delineating the alternation of **\*clay-rich** and clay-poor **\*lithologies**, e.g. **\*claystones** interbedded with **\*limestones** or **\*sandstones**. Conventionally, claystone horizons yield API values of more than 75. High gamma-ray log values will also be recorded from organic-rich **\*shales** which also concentrate other radioactive elements, e.g. uranium and thorium. In addition, **\*glauconitic** sands, volcaniclastic sands, **\*zircon-rich** sands, and clay-matrix-rich sands produce gamma-active sediments. See also [PHOTON LOG](#).

**gamma-ray spectrometer** See [SCINTILLATION COUNTER](#).

**gamma-ray spectrometry** Analytical method used in some branches of chemistry and physics for the measurement of the intensities and energies of gamma radiation. Scintillation or semi-conductor radiation detectors, coupled to various types of electronic circuitry, enable a spectrum to be accumulated. This may be used to identify the gamma-emitting radioisotopes, and their energy intensities can be used to determine concentrations of the corresponding elements. The technique is also used in **\*remote** sensing to determine the abundance of some elements in distant objects, e.g. the surface of the **\*Moon**.


**gangue** That portion of an ore deposit which is of no commercial value but which cannot be avoided during mining; it is removed during processing as

waste. Common gangue minerals are *\*quartz*, *\*calcite*, and *\*fluorite*, see [ORE MINERAL](#); [OREBODY](#); [ORE GRADE](#).

**ganister (gannister)** A fine-grained, *\*arenaceous* rock that underlies certain coal measures. It is used for its refractory qualities, e.g. to make furnace hearths.

**gannister** See [GANISTER](#).

**ganoid scale** Type of fish scale with a rhomboid shape, found in some fossil as well as extant *\*bony* fish (e.g. Polypterus and Lepisosteus). The scale consists of a superficial layer of enamel-like ganoine, a middle layer of *\*dentine*, and a basal layer of vascular bony tissue.



**Ganoid scale**

**Ganymede (Jupiter III)** One of the *\*Galilean* satellites, and the largest jovian satellite; it is bigger than *\*Mercury* and *\*Pluto*. It is believed to have a rock and metal core surrounded by a large mantle of water or water ice, 800–900 km thick, and the surface is of ice, with two types of terrain, one very cratered and dark, the other rather lighter, with many grooves and ridges. These terrains result from tectonic activity, but the details are not known. Both terrains are extensively cratered, the craters being flat, with no ring mountains and central depressions, and suggest the surface is about 3–3.5 Ma old. Ganymede has a magnetic field, embedded within that of Jupiter. Ganymede was discovered in 1610 by Simon Marius and Galileo.

Its diameter is 5268 km; mass  $1.48 \times 10^{23}$  kg; mean density 1940 kg/m<sup>3</sup>; visual albedo 0.42; mean distance from Jupiter  $1.07 \times 10^6$  km.

**Gaofen (GF)** A series of Chinese high-resolution (the name means high resolution) Earth-observation satellites; GF-1 was launched on 26 April 2013, from the Jinqian Satellite Launch Center, and GF-2 on 19 August 2014, from the Taiyuan Satellite Launch Center.

**GaoJing (SuperView Earth Observation Constellation)** A commercial, Chinese constellation of four remote-sensing *\*minisatellites* that provide imagery with 0.5-m panchromatic resolution and 2-m multispectral resolution along a 12-km swathe. The first pair of satellites was launched on 28 December 2016, and the second pair on 9 January 2018, all from Taiyuan Satellite Launch Center, into Sun-synchronous orbits at an altitude of 530 km.

**gap** Transverse valley that cuts through a ridge. It is termed a water gap when occupied by a stream; otherwise it is a wind gap. It may be a relic of an early stage in the development of a drainage pattern.

**Gardar rifting** An episode of rifting (see *RIFT*) that occurred about 1400–1000 Ma ago, after the *\*Ketilidian* orogeny but probably not causally connected to it.

**garnet** An important rock-forming *\*mineral* group, with the general formula  $X_3Y_2Si_3O_{12}$ , where X may be Ca, Mg, Fe<sup>2+</sup>, or Mn and Y may be Al, Fe<sup>3+</sup>, or Cr<sup>3+</sup>; the main minerals are *\*grossular* (X = Mg, Y = Al), *\*pyrope* (X = Mg, Y = Al), *\*almandine* (X = Fe<sup>2+</sup>, Y = Al), *\*spessartine* (X = Mn, Y = Al), *\*andradite* (X = Ca, Y = Fe<sup>3+</sup>), and *\*uvarovite* (X = Ca, Y = Cr<sup>3+</sup>), and there is continuous chemical variation in the group; an unusual variety called hydrogrossular  $Ca_3Al_2[SiO_4]_2[SiO_4]_{1-m}(OH)_{4m}$  has hydroxyl ions in the structure and is found in the rare rock type *\*rodingite*; sp. gr. 3.6–4.3; *\*hardness* 7.0–7.5; colour very variable depending on its chemical composition, and can vary from shades of deep red-brown to almost black, green, white, yellow, and brown; usually *\*vitreous* *\*lustre*; *\*crystals* *\*cubic*, with the most common form being *\*dodecahedra*; no *\*cleavage*; found in high-grade *\*metamorphic* and *\*igneous* rocks, in beach *\*sands*, and *\*alluvial* *\*placers*. Transparent pyrope crystals may be used as *\*gemstones*, but garnet is more generally used as an abrasive.

**GARP** See GLOBAL ATMOSPHERIC RESEARCH PROGRAMME.

**gas** See NATURAL GAS.

**gas chimney** A leakage of \*natural gas from a subsurface \*reservoir.

**gas chromatography** Analytical technique in which the components of a sample are separated by partitioning between either a mobile gas and a thin layer of non-volatile liquid held on a solid support (gas–liquid chromatography), or between the gas and a solid absorbent as the stationary phase (gas–solid chromatography). Partitioning occurs repeatedly throughout the column, and, as each solute travels at its own rate, a band corresponding to each solute will form. Solute bands are eluted (washed out) in increasing order of partition ratio, and enter a detector attached to the column exit. The time of emergence of a peak on the display identifies a component, and the area under each peak is proportional to the component's concentration. Gas chromatography is used mainly in the analysis of \*volatile organic compounds.

**Gasconadian** See TREMADOCIAN.

**gas–liquid chromatography** See GAS CHROMATOGRAPHY.


**Gaspra** A \*solar system asteroid (No. 951), measuring  $19 \times 12 \times 11$  km; approximate mass  $10^{16}$  kg; rotational period 7.042 hours; orbital period 3.29 years. It was imaged by \*Galileo in October 1991.

**gas-retention age** Measurement of the age of \*meteorites using the amount of radiogenic argon from the decay of potassium-40, or of helium from uranium and thorium, to indicate the time since the meteorite was last at a temperature at which gas leakage could occur. This is based on the principle that at high temperatures gases diffuse easily through \*silicate \*lattices, e.g.  $^{40}\text{Ar}$  from  $^{40}\text{K}$  diffuses out and is lost at temperatures above about 300 °C. If a meteorite has not suffered reheating its gas-retention age is the same as its \*formation age.

**gas–solid chromatography** See GAS CHROMATOGRAPHY.

**gastrolith** A stone swallowed (e.g. by some reptiles and birds) to break up food and so assist digestion. Such stones acquire a rounding and polish.

**Gastropoda (gastropod)** (phylum *\*Mollusca*) The class of Mollusca that includes snails and slugs. They have a true head, an unsegmented body, and a broad, flat foot. They appear in the *\*Cambrian* and occur in *\*sedimentary rocks* of all ages, occupying a range of aquatic and terrestrial environments. The majority of modern gastropods and all the *\*fossil* forms possess a coiled shell, which is all that is left to the palaeontologist for determining identification. The classification of living forms is based largely upon soft parts, so that similarly shaped shells developed by unrelated groups cause problems of nomenclature.



**Gastropoda**

**gauging station (gaging station)** A point at which river flow or groundwater levels are measured.

**Gault Clay** Glutinous marine deposit found in south-eastern England and in France, containing abundant fossil *\*bivalves*, *\*gastropods*, *\*ammonites*, and vertebrates. It is Lower *\*Cretaceous* (*\*Albian*) in age.

**Gauss, Karl Friedrich** (1777–1855) A German mathematician, Gauss developed the study of spherical geometry, which is essential to the theory of *\*plate tectonics*. His major work, *Theoria Motus Corporum Coelestium* ('Theory of the Motion of Heavenly Bodies') (1809), was on planetary

movements. He also helped to initiate the international geomagnetic survey of 1834.

**Gauss** A normal *\*polarity chron* in the late *\*Pliocene*, preceded by the *\*Gilbert* and followed by the *\*Matuyama* reversed *\*chrons*, and *\*radiometrically dated* at 3.58–2.60 Ma ago (Harland et al., 1982). The Gauss contains at least two reversed *\*polarity subchrons*: *\*Mammoth* and *\*Kaena*.

**gauss (G)** The c.g.s. unit of measurement of (a) magnetic field and (b) magnetic moment per unit volume. It has now been replaced by the *\*SI* units weber/m<sup>2</sup> (Wb/m<sup>2</sup>) and tesla (T).  $1 \text{ G} = 10^{-4} \text{ T} = 10^4 \text{ Wb/m}^2$ .

**Gaussian distribution** See NORMAL DISTRIBUTION.

**Gaussian process regression** See KRIGING.

**Gazzi-Dickinson method** A *\*point counting* technique that is used to measure statistically the components of a *\*sedimentary rock*, usually *\*sandstone*, using a randomly selected *\*thin* section and a device that randomly selects a minimum 300 and preferably 500 points on the slide. For each point the investigator must then determine the type of material at that point. The resulting counts are converted to percentages that can be used in comparisons between samples.

**GCM** See GENERAL CIRCULATION MODEL.

**GCOM** See GLOBAL CHANGE OBSERVATION MISSION-WATER.

**GCOM-C1** See GLOBAL CHANGE OBSERVATION MISSION-CLIMATE 1.

**GEDI** See GLOBAL ECOSYSTEM DYNAMICS INVESTIGATION.

**Gedinnian** See LOCHKOVIAN.

**gedrite** See AMPHIBOLES; ANTHOPHYLLITE.

**gehlenite** See MELILITE.

**Geiger counter** See GEIGER–MÜLLER COUNTER.

**Geiger–Müller counter (Geiger counter)** An instrument for detecting ionizing radiation which is used in general geologic prospecting. It consists

of a cylindrical metal cathode with a wire anode along its axis, the whole being enclosed in a thin-walled tube filled with low-pressure inert gas. In operation the cathode carries a charge of about 1000 volts, which is just short of that level needed to produce an electrical discharge across the cathode–anode space. A charged particle or **\*gamma** ray traversing this space collides with atoms of the inert gas, producing positive **\*ions** and negative **\*electrons**. Under the high voltage these are rapidly accelerated towards the cathode and anode, colliding on the way with other gas atoms and producing many more charged particles in a chain reaction. This avalanche arriving at the anode and cathode is registered as a pulse which is amplified to produce a click in a headphone set, or a succession of such pulses which can be expressed as a meter reading in milliroentgens per hour or counts per second. For more accurate surveys (especially from the air) a **\*scintillation** counter is required, which is a more sensitive instrument.

**Geikie, Archibald** (1835–1924) Director of the British Geological Survey from 1881 to 1901, Geikie made studies of glacial and fluvial **\*erosion**, and attempted to calculate the age of the Earth from rates of denudation. This led to conflict with **\*Kelvin**. Geikie was also one of the first historians of geology, stressing in his *Founders of Geology* (1897, 1905) the importance of the work of his fellow Scotsman **\*Hutton**.

**gel** Translucent to transparent, jelly-like material formed by the coagulation of a **\*colloid**; a non-homogeneous gelatinous precipitate; or a liquefied mud. A gel is more solid than a **\*sol** and is able to withstand some **\*shear** stress. **\*Bentonite** slurry is used as a gel in **\*diaphragm** walls.

**Gelasian** A **\*stage** (2.588–1.806 Ma ago) of the **\*Pliocene** **\*epoch**, preceded by the **\*Piacenzian** and followed by the Early **\*Pleistocene**.

**gel-filtration** Column-chromatography technique, normally employing polymeric carbohydrate-gel beads of controlled size and porosity as a stationary phase. Mixture components are separated on the basis of their sizes and rates of diffusion into the beads. Smaller molecules tend to diffuse more rapidly into the beads, thereby leaving the mainstream of solvent and so becoming retarded with respect to larger molecules. This method can also be used to determine the molecular weight of an unknown substance.

**gelifluction (congelifluction)** Flow of water-saturated sheets of rock debris over perennially frozen ground, and on slopes as low as 1°. It is the cold-

climate variety of **\*solifluction** and occurs only in the active layer (to a depth of 3 m).

**gelifraction** See FROST WEDGING.

**geliturbate** See GELITURBATION.

**geliturbation (congeliturbation, cryoturbation)** General term for all frost-based movements of the **\*regolith**, including **\*frost** heaving and **\*gelifluction**. The material disturbed by such movements is called 'geliturbate'.

**gelivation** See FROST WEDGING.

**gemstone** Naturally occurring mineral that has been artificially polished, faceted, and shaped for decorative purposes. It is normally classified as precious (e.g. **\*diamond**, ruby, and emerald) or semi-precious (e.g. **\*garnet**, **\*zircon**, and **\*topaz**). Many gemstones are hard, clear, and free from natural imperfections.

**genal angle** The angle where the lateral and posterior margins of a trilobite (see TRILOBITA) **\*cephalon** meet.

**genal spine** A spine that occurs in some species of trilobites (**\*Trilobita**) at the **\*genal angle**.

**gendarme** See ARÊTE.

**gene** Fundamental physical unit of heredity. It occupies a fixed chromosomal locus, and when transcribed has a specific effect upon the **\*phenotype**. It may mutate, and so yield various **\*allelic** forms. A gene comprises a segment of DNA (in some viruses it is RNA) coding for one function or several related functions. The DNA is usually situated in thread-like **\*chromosomes**, together with protein, within the nucleus; in bacteria and viruses, though, the chromosomes comprise simply a long thread of DNA.

**gene flow** Movement of **\*genes** within an interbreeding group that results from mating and gene exchange with immigrant individuals. Such an exchange of genes may occur in one direction or both.

**gene pool** The total number of \*genes, or the amount of genetic information, that is possessed by all the reproductive members of a population of sexually reproducing organisms.

**genera** The plural of genus. See CLASSIFICATION.

**general adaptation** Adaptation that fits an organism for life in some broad environmental zone, as opposed to 'special adaptations' which are specializations for a particular way of life. Thus the wing of a bird is a general adaptation, while a particular kind of bill is a special adaptation.

**general circulation** The term generally used to describe the large-scale circulation of the atmosphere over the globe, or over one hemisphere, with its more or less persistent features (which may be brought to prominence by considering long-term, or even shorter-term, averages), and all the transient features on various scales. Although in its nature it is a matter of the winds, the general circulation may be studied by means of barometric pressure maps because of the intimate relation between pressure and wind.

**general circulation model (GCM)** A computer simulation of the \*general circulation of the atmosphere for purposes of climatological studies. The atmosphere is contained within a notional three-dimensional grid and its condition described by the gas laws calculated at each point of intersection within the grid. Climatic development is then simulated by repeated recalculations of the effect of changes at one intersection on the intersections surrounding it. Present GCM studies are limited by the coarseness of the grids used, which cannot reflect events, such as cloud formation, occurring on a smaller scale, and by ignorance of the detail of certain important processes, such as those affecting energy transfer between the oceans and atmosphere.

**generalized reciprocal method (GRM, Palmer method)** A method of interpreting seismic refraction profiles over irregular layers, using forward and reverse shooting to obtain matched \*time–distance graphs. It is comparable to the \*plus–minus method, but less restricted in its application.

**generating curve** See COILING.

**Genesis** A \*NASA project to collect charged particles from the \*solar wind and return them to Earth. It was launched in July 2001 and returned in September 2004.



[http://www.nasa.gov/mission\\_pages/genesis/main/index.html](http://www.nasa.gov/mission_pages/genesis/main/index.html)

- A NASA mission to study the origins of the Sun and inner planets.

**Genesis Inflatable Space Complex** A space habitat that inflates once it reaches orbit, developed by Bigelow Aerospace, of Las Vegas. Genesis I was launched on 12 July 2006 and Genesis II on 28 June 2007, each with a volume of 11.5 m<sup>3</sup>, from Dombarovsky Cosmodrome, Russia.

**genetic drift** The random fluctuations of *\*gene* frequencies in a population, such that the genes amongst offspring are not a perfectly representative sampling of the parental genes. Although drift occurs in all populations, its effects are most marked in very small isolated populations, in which it gives rise to the random fixation of alternative *\*alleles* so that the variation originally present within single (ancestral) populations comes to appear as variation between reproductively isolated populations.

**genetic stratigraphic sequence model** One of the two schools of *\*sequence* stratigraphy, proposed in 1989 by W. E. Galloway ('Genetic stratigraphic sequences in basin analysis. 1: Architecture and genesis of flooding-surface bounded depositional deposits', *Bull. Am. Ass. Petrol. Geol.*, 73, 125–42), based on studies of the coast of the Gulf of Mexico. It uses marine flooding surfaces as the boundaries of strata, because these reorganize the sedimentary system and are easy to identify and correlate. Compare [DEPOSITIONAL SEQUENCE MODEL](#).

**genetic stratigraphic unit** A large-scale set of *\*facies sequences*, which were formed by slow deposition and are bounded by breaks in sedimentation, which can be related to one another for purposes of analysis and interpretation.

**geniculate twin (knee twin)** A special kind of twinned *\*crystal* in which the *\*twin-plane* has markedly changed the shape of the crystal, rather like a knee or elbow joint. The twin-plane is a reflection plane.

**genitus** Growth of a new cloud from a mother-cloud, where only a limited part of the mother-cloud is affected by the change. See also [CLOUD CLASSIFICATION](#); [MUTATUS](#).

**genotype** Genetic constitution of an organism, as opposed to its physical appearance (**\*phenotype**).

**genus** See CLASSIFICATION.

**genus-zone** See TAXON RANGE ZONE.

**GEO** See GEOSTATIONARY ORBIT.

**geo-** From the Greek *ge* meaning 'Earth' (a version of 'Gaia'), a prefix meaning 'pertaining to the Earth'.

**geo** Narrow inlet of a cliffed coastline, which has developed along a major near-vertical **\*joint** or **\*fault**.

**geobarometer** **\*Mineral** or group of minerals whose existence, coexistence, or element distribution, is stable between known pressure bounds at given temperatures and which is therefore useful as an indicator of the pressure under which a rock equilibrated. For example, the FeS content of the mineral **\*sphalerite**, in equilibrium with **\*pyrite** and **\*pyrrhotite**, is unaffected by temperatures in the range of 300–550 °C, but has been shown to vary according to pressure, and has been used as a geobarometer in regionally metamorphosed terrains.

**geobotanical anomaly** The marked local concentration, above background levels, of one or more elements in an ecological assemblage or specific plant which may indicate the presence of an ore deposit or a concentration of hydrocarbons.

**geobotanical exploration (biogeochemical exploration)** Traditionally, the use of indicator plant species or assemblages to detect the possible presence of metal-rich deposits. It is based on the limits-of-tolerance principle, i.e. it assumes that only specialized species can withstand metal-contaminated soils. In practice, plant response may be confusingly more complex (e.g. plants may respond to the low availability of essential nutrients rather than to a high concentration of toxic minerals) which makes such indicators unreliable. In modern use the concept includes the collection and chemical analysis of plant materials or soil layers, especially humus, in which metal ions may accumulate. It is a supplementary rather than a primary prospecting method.

**geochemical affinity** Affinity of an element for a particular environment.

**geochemical anomaly** 1. Abnormal concentration of elements in earth materials compared with background levels. 2. Increase of hydrocarbons in soils.

**geochemical cycle** A continuous cycle of elements passing through and between the Earth's *\*lithosphere*, *\*biosphere*, *\*hydrosphere*, and *\*atmosphere*. For example, sodium is released from rocks (lithosphere) by *\*weathering* and is transported in solution or suspension to the sea (hydrosphere). Sediments formed in the oceans take up sodium and may be compacted to join the geologic cycle, becoming *\*sedimentary*, *\*metamorphic*, and perhaps ultimately new *\*igneous* rock. See BIOGEOCHEMICAL CYCLE.

**geochemical differentiation** See PRIMARY GEOCHEMICAL DIFFERENTIATION; SECONDARY GEOCHEMICAL DIFFERENTIATION.

**geochemical soil survey** The process of collecting and analysing unconsolidated soil sediments in order to locate geochemical anomalies in the underlying rock and to use these to find *\*orebodies*. Where the loose material or soil is stable there is a geochemical balance between the original rock, *\*weathering*, pore water, and biological activity. The soil profile is usually layered, and in a geochemical soil survey the horizon giving the sharpest contrast between background and anomaly is chosen for analysis.

**geochemistry** Branch of geology concerned with the abundance and distribution of the chemical elements and their *\*isotopes* within the Earth or within solid bodies in the solar system, their circulation in natural systems (the *\*atmosphere*, *\*hydrosphere*, *\*biosphere*, and *\*lithosphere*), and the laws governing this distribution and its evolution.

**geochronologic unit** See GEOLOGIC-TIME UNIT.

**geochronology** Determination of time intervals on a geologic scale, through either absolute or relative dating methods. Absolute dating methods involve the use of radioactive elements and knowledge of their rates of decay: this yields an actual age in years for a given rock or *\*fossil*. Relative dating involves the use of fossils or sediments to place events and rock sequences in order, and does not provide absolute dates. See also DATING METHODS;

ABSOLUTE AGE; RELATIVE AGE; PLANKTONIC GEOCHRONOLOGY;  
GEOCHRONOMETRY.

**geochronometric scale (chronometric scale)** A timescale based on years BP (conventionally before 1950). Subdivisions on the scale are defined by particular units of duration (e.g.  $10^6$  years,  $10^9$  years) rather than reference points in actual rock successions. An example of such a subdivision is the placing of the boundary between the *\*Archaean* and the *\*Proterozoic* at 2500 Ma ago (i.e.  $2500 \times 10^6$  years) ago.

**geochronometry** The determination of the length of time intervals. Geochronometric resolutions for zonations based on different organisms may be calculated by dividing the time-span of a series by the number of *\*zones* and the intervals between zones. However, this will give only an approximate measure of time. *See also* DATING METHODS; GEOCHRONOLOGY.

**geode** A hollow, rounded body, which has a lining of *\*mineral \*crystals* pointing inward, e.g. *\*quartz* or *\*calcite*. The crystals grow into the cavity unimpeded and form perfect crystals which are frequently collected and valued for their beauty.

**geodesy** The science of measurement of the shape or figure of the Earth and its gravitational field. This science has expanded from topographic and astronomic surveying with the advent of satellite positioning systems, e.g. *\*GPS* and *\*SPS*.

**geodetic latitude** The latitude as defined by the vertical, relative to the geodetic reference ellipsoid (i.e. the shape that most closely matches mathematically the figure calculated by geodesy). The centre of the reference system is not geocentric.

**geodetic measurement** Any measurement concerned with the shape figure of the Earth, often involving the *\*geoid*.

**geodynamics** The study of geological change and the processes responsible for it. It includes *\*tectonics*, *\*erosion*, the formation of *\*sedimentary rock*, and *\*volcanicity*, but is concerned only with phenomena that occur on a macroscopic scale.

**geocology (landscape ecology)** The study of landscapes, including the manner of their formation and history, and their *\*ecology*.

**geo-electric section** A diagrammatic section of stratified layers that is deduced from electrical (resistivity) depth probing or drilling, where layers are identified by their *\*apparent resistivities*. Such sections are useful in detecting *\*water-table* levels and determining whether water is saline or fresh at the water-table.

**GeoEye-1 (OrbView-5)** A high-resolution imaging mission of GeoEye Inc., of Virginia, USA; since 31 January 2013, it has formed part of the DigitalGlobe commercial constellation. The satellite was launched on 6 September 2008, from California, into a Sun-synchronous circular orbit at an altitude of 681 km.

**geognosy** See WERNER, ABRAHAM GOTTLÖB.

**geographic information system (GIS)** A computer system that gathers, stores, and analyses geographic information, and displays it on demand.

**Geographos** A *\*solar* system asteroid (No. 1620), diameter 2 km; approximate mass  $4 \times 10^{12}$  kg; rotational period 5.222 hours; orbital period 1.39 years. It was to have been visited by the Clementine spacecraft, but a computer malfunction ended the mission.

**geoid** The gravitational equipotential surface corresponding to mean sea level, including the level at which the sea would stand in a continental area if it were able to do so.

**GEO-KOMPSAT-1** See COMMUNICATION, OCEAN AND METEOROLOGICAL SATELLITE.

**geological barrier** See FAR-FIELD BARRIER.

**Geological Long Range Inclined Asdic (GLORIA)** A sonar device, towed behind a ship, that transmits one sound beam to port and another to starboard and is used for studying the sea bed.

**Geological Society of America (GSA)** A society founded in 1888 that now has approximately 24 000 members in 115 countries. It aims to promote the professional development of Earth scientists and to facilitate the exchange of information.



<http://www.geosociety.org/>

- Home page of the GSA.

**Geological Society of London (GSL)** Founded in 1807, the GSL is the United Kingdom's national society for Earth sciences. It is a learned and professional body that exists to promote the Earth sciences and the professional interests of Earth scientists. With more than 11 500 members, the GSL is the largest geoscience society in Europe.



<http://www.geolsoc.org.uk/>

- Home page of the Geological Society of London.

**geologic cross-section** An interpretation of a vertical section through the Earth's surface, most usefully a profile, for which evidence was obtained by geologic and geophysical techniques or from a **\*geologic** map.

**geologic map** A map which shows the surface distribution of rock types, including their ages and relationships, and also structural features.

**geologic map symbols** Symbols used on a geologic map to provide a reduced or condensed pictorial representation of data, so that space is conserved.

**geologic timescale** A two-fold scale that subdivides all the time since the Earth first came into being into named units of abstract time, and subdivides all the rocks formed since the Earth came into being, into the successions of rock formed during each particular interval of time. The branch of geology that deals with the age relations of rocks is known as **\*chronostratigraphy**. The concept of a geologic timescale has been evolving for the last century and a half, commencing with a relative timescale (mainly achieved through **\*biostratigraphy**), to which it has gradually become possible to assign dates (see **DATING METHODS**) which are, nonetheless, subject to constant revision and refinement. Since the first International Geological Congress in Paris in 1878, one of the main objectives of stratigraphers has been the production of a complete and globally accepted **\*stratigraphic scale** to provide a historical framework into which all rocks, anywhere in the world, can be

fitted (see STANDARD STRATIGRAPHIC SCALE; CHRONOSTRATIGRAPHIC SCALE; UNIFIED STRATIGRAPHIC SCALE). Such a standard scale is still a long way off, but the names for *\*geologic-time* units and *\*chronostratigraphic units* down to the rank of *\*period/\*system* are in common use; many *\*epoch/\*series* and *\*age/\*stage* names are still regionally variable. Appendix B gives an outline geologic timescale employing currently common names and dates (though they are not necessarily universally accepted).



<https://engineering.purdue.edu/Stratigraphy/resources/geowhendiscl.html>

- A detailed geological timescale including stages.

**geologic-time unit (geochronologic unit)** A subdivision of geologic time, based on the rock record of the corresponding *\*chronostratigraphic unit*. Each time unit coincides with a particular chronostratigraphic unit and, like them, time units are ranked in order of decreasing duration, each unit comprising a number of units of shorter time interval (e.g. two or more *\*chrons* comprise an *\*age*, two or more ages comprise an *\*epoch*, etc.). Geologic-time units generally bear the same name as their chronostratigraphic counterparts, but the terms ‘lower’, ‘middle’, and ‘upper’ are changed to ‘early’, ‘middle’, and ‘late’.

<i>Chronostratigraphic Unit</i>	<i>Geologic-time Unit</i>
<i>*eonothem</i>	<i>*eon (longest)</i>
<i>*erathem</i>	<i>*era</i>
<i>*system</i>	<i>*period</i>
<i>*series</i>	<i>*epoch</i>
<i>*stage</i>	<i>*age</i>
<i>*chronozone</i>	<i>*chron</i>

**geology** Literally, the study (Greek *logia*) of the Earth (Greek *geo*). The medieval Latin word *geologia* may have been coined in the 8th century by St Bede (The Venerable Bede, 672 or 673–735) to distinguish earthly matters from godly matters (i.e. theology). It acquired its modern meaning in the 18th century. Geology is the scientific study of the composition, structure, and history of the Earth. More recently, many branches of geology have become established as disciplines in their own right (e.g.

\*geochemistry, \*geomorphology, \*geophysics, \*mineralogy, and \*palaeontology), and 'geology' has tended to decline in use.

**geomagnetic dipole field** The best mathematical fit to the observed \*geomagnetic field using a single dipole. If not stated, this is usually the geocentric dipole field, which is axial and inclined at about 11.3° relative to the Earth's axis of rotation. The axial geocentric dipole field is the best-fitting dipole, located at the centre of the Earth and aligned along the Earth's axis of rotation, and is usually considered to represent the Earth's magnetic field after averaging out the geomagnetic \*secular variation. See also NON-DIPOLE FIELD.

**geomagnetic equator** A line joining points on the Earth's surface where the \*inclination of the \*geomagnetic field is zero. To avoid \*magnetic-anomaly distortions it is usually calculated from harmonics of the geomagnetic field less than 12.

**geomagnetic field** The \*Earth's magnetic field shows variability on all timescales, ranging from nanoseconds to millions of years. Most \*transient variations are of external origin, reflecting interactions between the \*solar wind and the Earth's \*atmosphere. Longer-term changes, called \*secular variations, are of internal origin. The average annual field has been closely defined by satellite observations. The intensity of the field varies from about 30 μT near the equator to 60 μT near the observed \*geomagnetic poles at 73° N 100° W and 68° S 143° E. Most of the field (80%) can be accounted for mathematically by a single microgeocentric dipole inclined at 11.3° to the Earth's rotation pole, known as the geomagnetic dipole, with a magnetic moment of  $8.01 \times 10^{22}$  A/m<sup>2</sup>. The remaining field, the micro-non-dipole field, forms 12 main areas, varying by ±1.5 μT. An improved mathematical fit to the observed field was obtained using an inclined dipole offset from the geocentre by 340 km. At present the pattern of the field shows a tendency to drift westwards at about 0.2°/year, but this is not considered to be persistent on archaeological timescales. The field shows an ability to reverse polarity on a geologic timescale, with three polarity changes per million years during the last 60 million years, but long periods, of about 50 million years, of constant polarity are also known. The geomagnetic field is generally attributed to fluid motions within the outer \*core; these carry magnetic lines of force with them

(\*magnetohydrodynamics), creating a coupled \*self-exciting dynamo that allows polarity reversals.

**geomagnetic polarity interval** See CHRON.

**geomagnetic pole** The location on the Earth's surface of the axes of the calculated inclined \*geomagnetic dipole field. These are not \*dip poles.

**geomagnetic reversal timescale** See MAGNETO-STRATIGRAPHIC TIMESCALE.

**geomatics** The collection, management, analysis, and interpretation of spatial data pertaining to the Earth.

**geometric distribution** See LOG-NORMAL.

**geometric factor ( $K_g$ )** A numerical multiplier defined by the geometrical spacings between electrodes, which is used in conjunction with the voltage-to-current ( $R$ ) ratio measured in \*electrical resistivity surveys to give an \*apparent resistivity ( $\rho_a$ ) such that  $\rho_a = K_g \times R$ . The generalized formula for calculating  $K_g$  for a four-electrode configuration is:  $K_g = 2\pi(1/C_1P_1 - 1/C_1P_2 - 1/C_2P_1 + 1/C_2P_2)^1$  where  $C_1$  and  $C_2$ , and  $P_1$  and  $P_2$  are the current and potential electrode positions respectively. The geometric factor for the three main electrode configurations are: dipole–dipole,  $K_g = \pi n(n + 1)(n + 2)p$ ; Schlumberger,  $K_g = (\pi p^2/q)(1 - q^2/4p^2)$ ; Wenner,  $K_g = 2\pi p$ , where  $p$  and  $q$  are defined for each case.

**geomicrobiology** The scientific study of micro-organisms and their role in causing geochemical change.

**geomorphic sequence** A sequence of strata (see STRATUM) formed during a full cycle of rising sea level and its subsequent fall.

**geomorphology** The scientific study of the land-forms on the Earth's surface and of the processes that have fashioned them. Recently an extraterrestrial aspect has developed, resulting from studies of lunar and planetary surfaces.

**geopetal structure** A sedimentary \*fabric which records the way up at the time of deposition. Geopetal structures are commonly found in cavity fills

within [\\*limestones](#), where the lower part of the cavity has been filled with [\\*sediment](#) and the upper part filled later with [\\*cement](#).

**geophone (seismometer, pickup, 'jug')** A rugged device used to detect the arrival of seismic waves by transforming the ground motion into an electrical voltage. *Compare* [HYDROPHONE](#).

**geophysics** The science concerned with all aspects of the physical properties and processes of the Earth and planetary bodies and their interpretation, including, for example, [\\*seismology](#), gravity, magnetism, [\\*heat flow](#), and [\\*geochronology](#).

**geophysiology** A term coined by the English chemist James E. Lovelock to describe the mechanisms by which the conditions prevailing on Earth are regulated to ensure they remain suitable for living organisms. This includes the biological response to perturbations in those conditions. *See* [GAIAN HYPOTHESIS](#).

**Georgian epoch** *See* [EARLY CAMBRIAN EPOCH](#).

**geosphere** The upper part of the [\\*continental crust](#), together with the soil and organic matter found on or close to the surface. It is the part of the solid Earth that supports life and interacts directly with the [\\*biosphere](#), [\\*atmosphere](#), and [\\*hydrosphere](#).

**geostatic stress** *See* [LITHOSTATIC STRESS](#).

**Geostationary Meteorological Satellite (GMS)** A series of five weather and environmental satellites in [\\*geostationary orbit](#) at 140°E, and Japan's first such national series, operated by the Japanese Meteorological Agency (JMO) and Japan Aerospace Exploration Agency (JAXA). The satellites were launched on 14 July 1977 (removed to orbit on 26 January 2001), 11 August 1981, 3 August 1984, 6 September 1989, and 18 March 1995, from the Tanegashima Space Center, Japan.

**Geostationary Operational Environmental Satellite (GOES)** A series of meteorological satellites placed in [\\*geostationary orbit](#) by [\\*NASA](#). The first generation comprised eight satellites, designated GOES-A to GOES-H, launched in 1975, 1977, 1978, 1980, 1981, 1983, 1986 (launch failed), and 1987; all are now out of commission. The second generation comprised GOES-I to GOES-M, launched in 1994, 1995, 1997, 2000, and 2001; all are

now out of commission. GOES-N (launched 24 May 2006, at 75°W), GOES-O (launched 27 June 2009, at 105°W), and GOES-P (launched 4 March 2010, at 135°W) are second-generation satellites still in operation. GOES-Q was cancelled. GOES-R, launched 19 November 2016, is the first in the third generation.



<https://www.nasa.gov/content/goes>

- GOES Satellite Network.

**Geostationary Operational Meteorological Satellite-2** See [ELEKTRO-L](#).

**geostationary orbit (GEO, Clarke orbit)** A satellite orbit in which the satellite travels on the equatorial plane in the same direction as the rotation of the Earth at a height of about 36 000 km (more than 5 Earth radii) above the equator. Its orbital period is exactly one *\*sidereal* day and therefore the satellite remains vertically above a fixed spot on the surface of the Earth. At this height it has a view of almost the whole of one hemisphere. The possibility of such an orbit was first suggested by Arthur C. Clarke, for whom the orbit is sometimes named. *Compare* [GEOSYNCHRONOUS ORBIT](#); [LOW EARTH ORBIT](#); [MEDIUM EARTH ORBIT](#).

**geostrophic current** An ocean current that is the product of a balance between *\*pressure-gradient* forces and the *\*Coriolis* effect. This produces a current flow along the pressure gradient. Such a current does not flow directly from a region of higher pressure to one of lower pressure (i.e. 'down the slope' of the sea surface) but flows parallel to the gradient. All the major currents in the oceans, such as the *\*Gulf Stream*, are very nearly true geostrophic currents. The Gulf Stream, for example, can be likened to a river that does not run down a hill but around the hill.

**geostrophic wind** The wind blowing above the *\*planetary boundary layer* that, by its strength and direction, represents the balance between the *\*pressure-gradient* force, acting directly from the region of higher pressure towards the region of lower pressure, and the *\*Coriolis* effect (CorF), deflecting moving air to the right (in the northern hemisphere and to the left in the southern hemisphere). When these are in balance the wind flows parallel to the isobars (see [BUYS BALLOT'S LAW](#)). Air is also subject to a

centrifugal force, owing to the curvature of the air's path around a centre of low or high pressure. In the boundary layer, air experiences friction with the surface, slowing it, thereby reducing the CorF and causing it to flow across the isobars, at an angle of 10–20° over the sea and 25–35° over land (where friction is greater). *See also* [GRADIENT WIND](#).

**geosynchronous orbit** A satellite orbit around the Earth that has a 24-hour periodicity, so it follows the same path over the Earth's surface every day. *Compare* [GEOSTATIONARY ORBIT](#); [SUN-SYNCHRONOUS ORBIT](#).

**geosyncline** Large, downward structure generally of considerable extent, which may develop along a [\\*continental margin](#). The term was introduced originally by James Hall in 1859, and later an elaborate terminology was developed to describe and interpret the component parts. Many of these have fallen into disuse since the replacement of geosynclinal theory by more unifying theory of [\\*plate tectonics](#); the terms 'eugeocline' and 'miogeocline' are used today to denote particular rock associations.

**GEOTAIL** A collaborative mission between the [\\*Japan Aerospace Exploration Agency \(JAXA\)](#) and [\\*NASA](#) to study the structure and dynamics of the Earth's geomagnetic tail. The satellite was launched on 24 July 1992, from Cape Canaveral, into a highly elliptical orbit.

**geotechnical map** Map recording existing geology, and estimating likely conditions in terms helpful to the selection of construction techniques and ground treatment, and to the prediction of the reaction between ground and structure. Included in geotechnical maps are analytical maps, comprehensive maps, interpretive geologic maps, and slope category maps.

**geotherm** The curve representing the [\\*geothermal gradient](#) on a diagram showing the change of temperature with depth through the [\\*lithosphere](#) and upper [\\*mantle](#).

**geothermal brine** Hot, concentrated, saline solution that has circulated through crustal rocks in an area of anomalously high [\\*heat flow](#) and become enriched in substances leached from those rocks (e.g. chlorides of Na, K, and Ca); it often contains dissolved metals, in which case it forms an important intermediary in the deposition of [\\*ore](#) deposits. One of the best-documented examples is the brine from [\\*boreholes](#) in the Salton Sea [\\*geothermal field](#) in southern California, discovered in the early 1960s,

with temperatures between 300 and 325 °C, density 1021 kg/m<sup>3</sup>, and containing appreciable concentrations of Cu, Pb, Zn, and Ag. Hot brine solutions may also form through sea water–rock reactions in hydrothermal systems at oceanic ridges, e.g. in the median valley of the Red Sea.

**geothermal field** An area of the Earth characterized by a relatively high **\*heat flow**. The anomalously high rate of heat flow may be due to present, or fairly recent, **\*orogenic** or magmatic activity, or to the **\*radioactive** decay of **\*isotopes** of K, Th, and U where these occur at very high concentrations in crustal **\*granites** (**\*hot dry rocks**). In **\*sedimentary basins**, low thermal-conductivity values for the rocks are balanced by high thermal gradients, thus maintaining a constant heat flow. The high thermal gradients raise the temperature of deep, permeating water. Extraction of the water up deep **\*boreholes** provides surface water at temperatures useful for space heating. **\*Hot springs** and **\*fumaroles** can be important surface manifestations of a geothermal field.

**geothermal gradient** The increase of temperature with depth. It usually refers to depths below 200 m. In the continents, the gradient is usually between 20 and 40 °C/km, although it can well exceed this in volcanic regions. In the oceans, the depth of penetration of most core barrels is so short that the gradient can be determined over only a few metres and varies considerably. The average geothermal gradient at the surface of the Earth is about 24 °C/km, but it is assumed to decrease with depth as widespread **\*mantle** melting would otherwise occur. The observed gradients are therefore modified to result in an estimated temperature of about 1200 °C at the top of the seismic **\*low-velocity zone** in the upper mantle. Within the mantle, the increase of temperature with depth is considered to be less than 0.1 °C/km greater than the adiabatic increase of 0.33 °C/km. *See also* **HEAT FLOW**.

**geothermic survey** A survey measuring the **\*heat-flow** variations within a region.

**geothermometer** An indicator of the temperature, or range of temperatures, at which a geologic event (e.g. the crystallization of a **\*magma** or the metamorphism of pre-existing rocks) occurred. Apart from the presence or absence of **\*minerals** or mineral assemblages known to be stable within certain temperature ranges, among the most widely used indicators are: (a)

**\*stable-isotope** distribution, e.g. the ratios of  $^{18}\text{O}$  to  $^{16}\text{O}$  between different mineral pairs varies according to temperature (*see also* **OXYGEN-ISOTOPE ANALYSIS**); (b) mineral transformations or inversions known to be temperature dependent, e.g. the transition of  $\alpha$  quartz to  $\beta$  quartz at 573 °C; (c) liquid–vapour homogenization points in **\*fluid** inclusions (subject to certain assumptions, the temperature of formation of a crystal is indicated by the temperature at which the vapour bubble coexisting in the inclusion disappears upon heating); (d) unmixing or **\*exsolution** lamellae of mineral pairs below a particular temperature, e.g. chalcopyrite–bornite at 500 °C; (e) temperature-dependent element distribution between coexisting minerals, e.g. iron–titanium oxide distribution between the coexisting mineral pairs **\*magnetite**–ulvöspinel and **\*ilmenite**–**\*hematite** (in this instance subject also to oxygen **\*fugacity**).

**germanate system** Series of compounds which have identical structures to the **\*silicates**, except that **\*phase** transitions take place at lower pressures. Used by early workers to study possible phase transitions in the Earth's **\*mantle** before apparatus capable of attaining realistic geologic pressures was available.

**geyser** A small opening on the Earth's surface which periodically spouts a fountain of boiling water into the air. The largest fountain height recorded was 500 m, from a now extinct geyser in New Zealand. Water beneath the mouth of a geyser is heated by conduction from surrounding hot rocks, water at the base of the column boiling before that higher in the column. Expanding vapour bubbles rise in the column of water, expelling water at the top and lowering the pressure at the base. This allows the onset of further boiling, the system being self-sustaining, until the entire column of water is blown out of the system as a water spout. The water involved carries a large load of dissolved minerals which precipitate around the mouth of the geyser as **\*siliceous sinter**. The name is from Geysir, about 45 km from the active volcano Hekla, Iceland, and was first used as a technical term in 1847 by the German chemist R. W. von Bunsen, who spelled it 'geysir'. This spelling is still sometimes used.

**GF** *See* **GAOFEN**.

**GHGSat-D** *See* **GREENHOUSE GAS SATELLITE – DEMONSTRATOR**.

**GHOST** See GLOBAL HORIZONTAL SOUNDING TECHNIQUE.

**ghost** A spurious seismic **\*reflection** which occurs when energy is transmitted upwards from a subsurface **\*shot** and then reflected downwards from the surface of the ground or sea. A ghost wave train may interfere with other downward-moving waves, thus modifying their wave-form, and may add a reverberation tail. This type of ghost reflection is one kind of **\*multiple** event. A seismic event recorded from a reflector located outside the plane of the seismic section is called an 'off-section ghost' and may be a problem in an area where the three-dimensional subsurface topography is pronounced but only a two-dimensional seismic survey is being conducted.

**ghost stratigraphy** The alignment of **\*country** rock **\*xenoliths** within large **\*granite** bodies such that they reflect a continuation of the **\*stratigraphy** and structure of the country rocks surrounding the **\*intrusion**. An old idea, ghost stratigraphy was regarded as evidence favouring the view that granite may form by the replacement of country rocks.

**Ghyben–Herzberg relationship** Beneath oceanic islands, the thickness ( $d$ ) of a lens of fresh **\*groundwater** (density  $\rho_w$ ) overlying sea water (density  $\rho_m$ ) can be determined if the height above sea level ( $h$ ) of the top layer of the lens is known and the conditions are static. The relationship is  $d = ah = \rho_w/(\rho_m - \rho_w)$  where  $a$  is typically about 38.

**Giacobini–Zinner** A **\*comet** with an orbital period of 6.52 years; **\*perihelion** date 21 November 1998; perihelion distance 0.996 AU.

**gibber** See GIBBER PLAIN.

**gibber plain** A term used in Australia to describe an extensive plain (normally a **\*pediplain**) that is mantled by loose rock fragments (gibber). These fragments are typically the rubble left from the destruction of a **\*silcrete** **\*duricrust** or from the breakdown of resistant **\*conglomerates** (in which case the gibber consists of **\*quartz** pebbles).

**Gibbs free energy** See GIBBS FUNCTION.

**Gibbs function (Gibbs free energy; symbol  $G$  or  $F$ )** Generally defined in terms of changes in free energy:  $\Delta F = \Delta H - T\Delta S$ , where  $H$  = enthalpy,  $T$  = absolute temperature,  $S$  = entropy. In a mixture of reactants, if  $\Delta F$  is less than 0 a reaction may take place spontaneously, whereas if  $\Delta F$  is positive

energy must be supplied to the system for a reaction to occur at all. In geochemistry, the sign and magnitude of  $\Delta F$  are important. The sign indicates whether a given reaction can occur spontaneously, and the magnitude indicates how far the reaction can go before equilibrium is attained.

**gibbsite** Mineral,  $\text{Al}(\text{OH})_3$ , that is a constituent of **\*bauxite**; sp. gr. 2.4; **\*hardness** 3; greyish-white; occurs as an alteration product of aluminium silicates in **\*laterite** and bauxite deposits.

**giga-** From the Greek *gigas* meaning 'giant', a prefix (symbol G) attached to SI units, meaning the unit  $\times 10^9$  (e.g. 2 Gm = 2 gigametres =  $2 \times 10^9$  m).

**gigantic jet** A **\*transient luminous** event that resembles a conical **\*red sprite** but that propagates upward from the core of an oceanic thunderstorm and is not linked to cloud-to-surface lightning.

**Gigantoproductus giganteus** See PRODUCTUS GIGANTEUS.

**Gilbert** A reversed **\*polarity chron** in the mid **\*Pliocene**, which is followed by the **\*Gauss** normal polarity chron. It started about 5.70 Ma ago and lasted until 3.58 Ma ago. The Gilbert contains at least four normal **\*polarity subchrons**: **\*Thvera**; **\*Sidufjall**; **\*Nunivak**; and **\*Cochiti**.

**Gilbert, Grove Karl** (1843–1918) An officer on the US Geological Survey, Gilbert made studies of Meteor Crater, Arizona, suggesting that it was the product of a collision with an **\*asteroid**, and also studied lunar **\*craters**. He investigated crustal movements around Lake Bonneville, developing a theory of crustal isostatic compensation (see ISOSTASY). He also distinguished between folded and block-faulted mountains.

**Gilbert, William** (Gilberd, William) (1540–1613) A natural philosopher and physician to Elizabeth I, Gilbert studied terrestrial magnetism. In his book *De Magnete Magneticisque Corporibus, et de Magna Magnete Tellure* (1600) he made a firm distinction between magnetism and electricity. He explained the Earth's magnetic field by likening the Earth to a vast spherical magnet.

**Gilbert-type delta** A type of river **\*delta** which consists of a wedge-shaped body of **\*sediment**, comprising relatively thin, flat-lying, **\*topset** sediments, long, steeply dipping **\*foresets** which prograde (see PROGRADATION) from

the river mouth, and thinner, flat-lying, [\\*bottomset](#) or [\\*toeset](#) deposits. Gilbert-type deltas are often developed in lakes, where river water and lake waters are of the same density. It was first described by [Grove Karl \\*Gilbert](#).

**gilgai** Undulating micro-relief of soils which contain large amounts of clay minerals (e.g. [\\*montmorillonite](#)) that swell and shrink considerably on wetting and drying, to an extent that may be sufficient to fracture pipelines or move telegraph or fence poles from the vertical. The word is derived from a settlement in Queensland, Australia, where the soils of swelling clays are especially common. *See also* [PATTERNED GROUND](#).

**gills** The respiratory organs of aquatic animals, consisting of an outgrowth from the body surface or from an internal layer of modified gut. This provides a large surface area, well supplied with blood vessels. As water passes across the gills, oxygen and carbon dioxide are exchanged across the walls of the blood vessels.

**gilsonite (asphaltite)** Solid form of [\\*hydrocarbon](#) which occurs as pure, natural [\\*bitumen](#); sp. gr. 1:05–1:10; [\\*hardness](#) 2; black; occurs in [\\*veins](#), [\\*lodes](#), and sedimentary rocks. On heating it softens rapidly and flows like a liquid. It is used in waterproof coatings, wire insulations, and lacquer.

**Ginkgoales (maidenhair tree)** An order of the [\\*Coniferopsida](#), one of the major [\\*gymnosperm](#) groups. Only one species of Ginkgo survives, *Ginkgo biloba*. The first undoubted maidenhairs occur in [\\*Triassic](#) rocks, and in the subsequent [\\*Jurassic](#) period their distribution was practically world-wide. The surviving species is restricted (in the wild) to China, and its fan-shaped leaves, with open [\\*dichotomous](#) venation are strikingly similar to fossil Ginkgo leaves. Many fossil species are known. The earliest representative of the genus, *G. digitata*, is first recorded from the Middle Jurassic. The restricted geographical range, the unchanged appearance of the leaves, and the motile male sperms (otherwise known only in living seed plants in the [\\*Cycadales](#)) have together led to the maidenhair being referred to as a [\\*living](#) fossil. Other classifications have placed ginkgos in the class Ginkgoopsida.

**Giotto** An [\\*ESA](#) mission, launched on 2 July 1985, to the comets [\\*Halley](#) and [\\*Grigg–Skjellerup](#). The mission ended on 23 July 1992.



<http://sci.esa.int/science-e/www/area/index.cfm?fareaid=15>

- An ESA mission launched in 1985 that studied Halley's Comet.

**gipfelflur** A plain or apparent surface which is made up of the summit levels in a mountainous region. *Gipfelflur* is a German word whose literal meaning is 'peak-plain'.

**GIS** See GEOGRAPHIC INFORMATION SYSTEM.

**Gisbornian** A *\*stage* (460.9–456 Ma ago) of the Middle *\*Ordovician* of Australia, underlain by the *\*Darriwilian* and overlain by the *\*Eastonian*.

**GISP (Greenland Ice Sheet Project)** A US drilling project that extracts ice cores from the Greenland ice sheet and uses them to obtain atmospheric and palaeoclimatological information. The first core reached bedrock at a depth of about 3000 m and in 1993 a second hole penetrated about 1.55 m into bedrock at a depth of 3053.44 m, about 30 km from *\*GRIP*.

**Givetian (Taghanican, Tioughniogan)** A *\*stage* in the Middle *\*Devonian* epoch preceded by the *\*Eifelian*, followed by the *\*Frasnian*, and dated at 391.8–385.3 Ma ago (Int. Commission on Stratigraphy, 2004). The corresponding European *\*stage* is zoned on goniatites and spiriferid brachiopods. It is roughly contemporaneous with the lower *\*Condobolinian* (Australia) and *\*Erian* (N. America).

**glabella** See CEPHALON.

**glaci- (glacio-)** Dominated by glacial ice. The prefix is followed by a term indicating the environment or process that is so dominated, e.g. glaciaquatic (of water derived from a *\*glacier*), glacioeustasy (the theory that changes in sea level result from the growth and decay of *\*ice* sheets), glacifluvial (of sediments or land-forms produced by meltwater streams escaping from a glacier), glacioisostasy (the theory that local flexing of the Earth's *\*crust* occurs as a result of the loading and unloading that takes place as large ice sheets wax and wane), glacialacustrine (pertaining to a lake adjacent to a glacier), and glaciomarine (of sediments laid down in a sea environment near a glacier). See also GLACIOTECTONICS.

**glacial breach** A glacially eroded trough that cuts through a ridge and so breaches a former *\*watershed*. It is formed when the outflow of a *\*glacier* (or *\*ice* sheet) is impeded, its thickness consequently increases, and ultimately a new escape route (the breach) is exploited. This process is called 'glacial diffluence' when a single glacier spills out of its valley, and 'glacial transfluence' when several breaches are formed due to the accumulation of a large ice sheet. The many breaches through the western Highlands of Scotland are due to the accumulation of transfluent ice east of the main watershed.

**glacial diffluence** See [GLACIAL BREACH](#).

**glacial diversion** Displacement of a pre-glacial stream by the action of a *\*glacier*. In upland areas the new drainage often follows a *\*glacial* breach. In lowlands, glacial drift may block an existing valley and lead to stream diversion.

**glacial drainage channel (meltwater channel)** Channel cut by the action of glacial meltwater or by water from an ice-dammed lake. Various types may be recognized, classified by the position of the channel with reference to the *\*glacier*, e.g. ice-marginal, *\*englacial*, or subglacial. Usually these channels are steep-sided and flat-floored, and are unrelated to the present drainage pattern.

**glacial horn** See [HORN](#).

**glacial lake outburst flood (GLOF)** See [OUTBURST FLOOD](#).

**glacial limit** A line marking the furthest extent of a former glacial advance. It may be identified on the ground through the recognition of features associated with glacial margins, including lateral and terminal *\*moraines*, outwash spreads (*\*sandur*), marginal *\*meltwater* channels, and *\*proglacial* lakes. See also [TRIM LINE](#).

**glacial morphoclimatic zone** A *\*morphoclimatic* region in which glacial action is at a maximum. The rate of *\*mechanical* weathering is high, especially that due to *\*frost* action, but the rate of *\*chemical weathering* is low. There is significant wind action. *\*Fluvial* action occurs during the summer melt. There is little *\*mass-wasting*, except locally.

**glacial period** A general term used to describe either a glacial *\*stage* (e.g. *\*Devensian*), or an indeterminate period of *\*glaciation*.

**glacial plucking (quarrying)** The removal of relatively large fragments of bedrock by direct glacial action. The process involves several mechanisms, including the incorporation of rock fragments into the base of the *\*glacier* when it freezes to weakened bedrock, and the removal of bedrock material when fragments already included in the ice are dragged over it.

**glacial stairway** Long profile of a *\*glacial* trough: it is characterized by alternating rock bars (*\*riegels*) and rock basins, giving the impression of a stairway. The structure is attributed to variations in the erosive power of ice, or to the influence of rock jointing.

**glacial theory** The theory, developed in the late 1830s and 1840s by Venetz, *\*de Charpentier*, and *\*Agassiz*, that most of Northern Europe, N. America and the north of Asia, had been covered by *\*ice* sheets during a period later termed the *\*Pleistocene*. The hypothesis was used to explain *\*erosion*, and the subsequent deposition of *\*till* or *\*boulder* clay, and the extinction of species such as the *\*mammoth*. Since that time, glacial theory has been developed to include multiple glaciation, and evidence of much older ice ages.

**glacial transfluence** See [GLACIAL BREACH](#).

**glacial trough** A relatively straight, steep-sided, U-shaped valley that results from glacial *\*erosion*. Its cross profile approximates to a parabola, while its long profile is often irregular, with rock bars (*\*riegel*) and over-deepened rock basins being typical features. The world's largest glacial trough is that of the Lambert Glacier, Antarctica which is 50 km wide and about 3.4 km deep.


**glaciaquatic** See [GLACI-](#).

**glaciated rock knob** See [ROCHE MOUTONNÉE](#).

**glaciation** The covering of a landscape or larger region by ice; an ice age.

**glacier** A large mass of ice, resting on or adjacent to a land surface, and typically showing movement. Glaciers may be classified in several ways. The most useful division (as it relates to work done) is based on temperature, and three categories are recognized. In temperate (or warm)

glaciers (e.g. those of the Alps) the ice is at **\*pressure melting** point throughout, except during winter when the top few metres may be well below 0 °C. Movement is largely by basal slip. Polar (or cold) glaciers (e.g. parts of the Antarctic sheet) have temperatures well below the pressure melting point and movement, which is slow, is largely by internal deformation. Subpolar glaciers (e.g. those of Spitzbergen) have temperate interiors and cold margins and so are composite. The morphological classification is based largely on the size, shape, and position of the ice mass and **\*cirque glaciers**, **\*valley glaciers**, and **\*piedmont glaciers** are among the types recognized.



## Glacier

**glacier creep** The deformation of **\*glacier** ice in response to **\*stress**, by a process involving slippage within and between ice crystals. The rate of creep is dependent on both stress and temperature. When the **\*shear** stress is doubled, the **\*strain rate** increases eight times, and a rise in temperature from -22 °C to 0 °C involves a ten-fold increase in strain rate.

**glacier ice** See ICE.

**glacier power** A general term for the ability of a **\*glacier** to erode its bed. A distinction is made between total glacier power (basal **\*shear** stress multiplied by average velocity) and effective power (determined by the contribution of **\*basal sliding** to the total velocity). It follows that temperate

glaciers may be roughly ten times more powerful than polar and subpolar types.

**glacier surge** A relatively rapid movement of a valley *\*glacier*, or of an individual *\*ice* stream within a major *\*ice* sheet. The movement may build up over a period ranging from a few months to several years and may be a hundred times faster than the 'normal' velocity. Surging may result from an increase in ice thickness or from excessive basal water.

**glacier wind** See FIRN WIND.

**glacifluvial** See GLACI-.

**glacilacustrine** See GLACI-.

**glacio-** See GLACI-.

**glacioeustasy (glacioeustatism)** The theory that sea levels rise and fall in response to the melting of ice during *\*interglacials* and the accumulation of ice during *\*glaciations*. A major glacioeustatic oscillation has an amplitude of some 100 m. See GLACI-.

**glacioisostasy** The adjustment of the *\*lithosphere* following the melting of an *\*ice* sheet. The gradual rise of the land enables an estimation to be made of the flexural rigidity of the lithosphere and the viscosity of the *\*mantle*. Adjustments need to be made to gravity readings in previously glaciated areas to allow for this effect, which occurs on a regional scale. Some 1000 m of depression may have occurred in Scandinavia during the last *\*ice* age, where 520 m of recovery has been recorded. The process gives rise to warped shorelines. See GLACI-.

**glaciology** The scientific study of *\*ice* in all its forms. It therefore includes the study of ice in the atmosphere, in lakes, rivers, and oceans, and on and beneath the ground. Commonly, however, it is the study of *\*glaciers*.

**glaciomarine** See GLACI-.

**glaciomarine sediment** High latitude, deep-ocean sediment, which originated in glaciated land areas and has been transported to the oceans by *\*glaciers* or icebergs. Such sediments may contain large dropstones, transported by and dropped from icebergs, in the midst of fine-grained sediments.

**glaciotectonics** 1. The study of structures within a *\*glacier*. These may be identified by contorted layers of rock debris. They are most common when a glacier is frozen to its bed and can move only by thrusting and folding. 2. The study of structures imposed on bedrock by glacial movement and on *\*drift* by both movement and the loss of support that occurs on melting.

**glaeboles** See CALICHE.

**glance** A mining term, from the German *glanz* (lustre), that describes any mineral with a lustre and colour indicating it is metal compound. *\*Argentite* is known as silver glance; specularite (see HEMATITE) is called iron glance.

**glass** An amorphous, *\*metastable* solid with the atomic structure of a *\*silicate* liquid. Glass can be formed by *\*quenching* a silicate *\*melt*, the short timescale for cooling or pressure reduction preventing the reorganization of the random liquid structure into an ordered crystalline structure. Since cohesion between atoms in the liquid silicate increases with increasing silica content, melts with high silicate contents are most likely to form glasses. Natural *\*igneous* glasses of *\*rhyolite* composition (70% SiO<sub>2</sub>) are termed 'obsidians'. A wide variety of glasses, formed by meteoritic impact into the lunar *\*regolith*, exist on the lunar surface. Shapes include spheres averaging 100 µm in diameter, tear-drops, dumb-bells, etc., typical of rotational shapes assumed by splashed liquids. They do not resemble meteoritic *\*chondrules*. Volcanic glasses, formed by *\*fire* fountains during eruption of *\*mare* basalts, also occur locally on the Moon. Their compositions match those of local surface rocks, soils, or minerals. No *\*tektite* compositions are found.

**glass-plate reflector** See REFLECTOR.

**glass shards** Angular, glassy particles, less than 2 mm in size, formed either by the explosive magmatic fragmentation of *\*pumice* *\*vesicle* walls, or by the chilling and brittle fragmentation of *\*magma* when it comes into contact with groundwater or surface water. Magmatically formed shards commonly have 'Y' or cusped shapes and may deform plastically and weld together if they are hot enough when deposited and the overburden load is sufficient. Shards formed by *\*hydrovolcanic* processes have a variety of shapes, ranging from those with curvilinear surfaces and low vesicularity to those with smooth, fluid-form surfaces and moderate vesicularity.

**glass sponges** See HEXACTINELLIDA.

**glassy** Applied to **\*igneous** rocks (e.g. **\*obsidian**) that cooled too rapidly for crystals to form and that, consequently, have no internal structure and resemble glass.

**glauconite** Although sometimes considered as **\*clay mineral**, glauconite is more accurately a member of the **\*mica** group, with the composition  $(K, Ca, Na)_2(Fe^{3+}, Al, Mg, Fe^{2+})_4[(Si, Al)_4O_{10}]_2(OH)_4$ ; sp. gr. 2.4–3.0; **\*hardness** 2; **\*monoclinic**; olive green, yellowish, or blackish green; dull **\*lustre**; granular; occurs in marine sediments as aggregates up to 1 mm in diameter. It is being formed on many modern **\*continental shelves**, at depths from a few tens to hundreds of metres, where the sedimentation rate is low and decaying organic matter is present in a generally oxidizing environment. In many **\*sandstones** glauconite can impart a green colour when abundant, as in the **\*Cretaceous Greensands** of Britain and the eastern USA.

**glaucony** A green-coloured, marine sedimentary **\*facies** characterized by the presence of **\*grains** of the **\*glauconite** mineral family, which develop on **\*continental margins** and on ocean highs. Glaucony is of variable mineralogy due to the replacement of mineralogically different initial substrates by **\*authigenic** minerals of the glauconite family.

**glaucophane** An important alkali **\*amphibole** of composition  $Na_2(Mg_3Al_2)[Si_8O_{22}](OH)_2$ , and **\*end-member** of the glaucophane–riebeckite  $(Na_2(Fe^{2+}_3Fe^{3+}_2)[Si_8O_{22}](OH)_2)$  series; sp. gr. 3.0; **\*hardness** 6.0; blue, bluish-black; **\*prismatic** crystals, **\*fibrous** or **\*granular** habit; it is an **\*essential** mineral of **\*blueschists** under conditions of low temperature and high pressure in metamorphosed sediments at **\*destructive** plate margins.

**glaucophane-schist facies** A set of metamorphic **\*mineral** assemblages produced by **\*metamorphism** of a wide range of starting rock types under the same high-pressure/low-temperature metamorphic conditions and typically characterized by the development of the mineral assemblage **\*glaucophane–lawsonite–\*quartz** in rocks of **\*basic \*igneous** composition. Other rocks of contrasting composition, for instance **\*shales** or **\*limestones**, would each develop their own specific mineral assemblage, even though they are all being metamorphosed under the same conditions. The variation

of mineral assemblage with starting rock composition reflects a particular range of pressure, temperature and  $P(\text{H}_2\text{O})$  conditions. Experimental studies of mineral  $P$ – $T$  stability fields indicate that the facies represents high-pressure/low-temperature conditions which can be met during **\*subduction** of **\*oceanic** crust under **\*continental crust**. Because rocks of basic igneous composition within this facies are characterized by the blue-coloured mineral glaucophane, the **\*facies** is sometimes referred to as the ‘blueschist facies’.

**G-layer** The inner, solid **\*core** of the **\*Earth**, extending from a depth of about 5150 km to the Earth’s centre at 6371 km depth.

**glaze** Clear-ice deposit on objects, produced by the freezing of supercooled water droplets on to surfaces at temperatures below 0 °C. See **CLEAR ICE**.

**Gleedonian** A **\*stage** (425.4–422.9 Ma ago) of the **\*Silurian**, underlain by the **\*Whitwellian** and overlain by the **\*Gorstian**.

**Glen’s power flow law** A model of the way ice deforms based on experiments on blocks of ice conducted in the early 1950s by J. W. Glen and adapted by J. F. Nye to apply to **\*glaciers**. Glen found that the **\*strain rate** in a block of ice subjected to constant **\*stress** reaches a steady value. In glaciers, the relationship between strain rate ( $e$ ) and effective **\*shear** stress ( $\tau$ ) is given by  $e = A\tau^n$ , where  $A$  is a constant related to temperature and  $n$  is an exponent with a mean value of approximately 3.

**gley** The product of waterlogged soil conditions, and hence an **\*anaerobic** environment; it encourages the reduction of iron compounds by micro-organisms and often causes mottling of soil into a patchwork of grey and rust colours. The process is known as gleying, or gleyzation (US usage).

**gleying** See **GLEYSOIL**.

**gley soil** In the soil classification developed by the Soil Survey for England and Wales, a soil that is subject to long periods of waterlogging. **\*Soil horizons** in a gley soil may be grey in colour, with mottling.

**gleysols** A reference soil group in the **\*World Reference Base for Soil Resources** classification scheme. Gleysols show evidence of gleying (see **GLEYSOIL**) within 50 cm of the surface.

**gleyzation** See GLEY.

**gliding tectonics** See GRAVITY TECTONICS.

**gliding twins** See DEFORMATION TWINNING.

**glimmerite** An \*ultrabasic \*igneous rock, consisting almost wholly of \*essential dark \*mica, either \*phlogopite or \*biotite. These rocks are rather rare, being found among \*ultramafic \*xenoliths in \*kimberlite pipes and within old \*basement \*gneisses. These occurrences testify to the deep-seated origin of glimmerites, which might be considered as metamorphic (see METAMORPHISM) rather than igneous.

**Glinka, Konstantin Dimitrievich** (1867–1927) A Russian soil scientist who worked at the University of St Petersburg, Glinka was a student of \*Dokuchayev and developed and organized his work. He was responsible for the soil surveys of most of European Russia and of Siberia. His book *Soil Science* was first published in 1908.

**Global Atmospheric Research Programme (GARP)** International project that aims to provide a comprehensive knowledge of atmospheric structure as an aid to prediction. The complex system is based on land and marine weather stations, together with upper-air soundings, satellite remote-sensing, and sensors carried by balloons.

**Global Change Observation Mission-Climate 1 (GCOM-C1)** A satellite mission by the \*Japan Aerospace Exploration Agency (JAXA) to provide long-term observations of the Earth's environment on a global scale. The satellite was launched on 23 December 2017, from Tanegashima Space Center, Japan, into Sun-synchronous orbit at an altitude of 798 km.

**Global Change Observation Mission-Water (GCOM)** A constellation of two satellites by the \*Japan Aerospace Exploration Agency (JAXA) to observe the sea surface and contribute to measurements related to the carbon cycle and radiation budget, with reference to climate change. The satellites were launched on 17 May 2012, from Tanegashima Space Center, Japan, into Sun-synchronous orbit at an altitude of 699.6 km.

**Global Ecosystem Dynamics Investigation (GEDI)** A laser-based instrument system that observes the structure of forest canopy. It is scheduled for launch to the International Space Station in 2019

**Global Heritage Stone Resource** A proposed designation that seeks international recognition for natural stone resources that have been widely used in human culture. It is proposed by the Heritage Stone Task Group of the [\\*International Union of Geological Sciences](#) and the [\\*International Association for Engineering Geology and the Environment](#).



<http://globalheritagestone.com>

- IIGS Subcommittee: Heritage Stones.

**Global Horizontal Sounding Technique (GHOST)** Programme for the direct sensing of the atmosphere, using balloons designed to float at various constant-density levels. These are tracked and their various measurement readings are monitored by polar orbiting satellites. Sensors on the balloons record temperature, humidity, and pressure; tracking gives mean winds for each balloon. The project forms part of the [\\*World Weather Watch](#).

**Global Positioning System (GPS)** A total of 18 satellites, in different orbits, so that at least four are visible at any time from all points on the Earth's surface. They give a locational accuracy of less than 2 cm on baselines of 1000 km. It is planned to use them for measuring the motions of small blocks of the Earth's [\\*crust](#), e.g. opposite sides of a fault system, [\\*microplates](#), etc. (The satellites also have obvious military applications.)

**Global Precipitation Measurement (GPM)** A collaborative mission between [\\*NASA](#) and the [\\*Japan Aerospace Exploration Agency \(JAXA\)](#) to provide near-global measurement of precipitation, improve the accuracy of measurements of rain rates and latent heating, and predict floods and freshwater resources. The satellite was launched on 27 February 2014, from the Tanegashima Space Center, Japan, into a Sun-synchronous circular orbit at an altitude of 407 km.

**Global-scale Observations of the Limb and Disk (GOLD)** An imaging mission within [\\*NASA's](#) Explorer Program to investigate how geomagnetic storms, ultraviolet variability during solar extremes, and atmospheric waves and tides affect the [\\*thermosphere](#), and how the night-time equatorial [\\*ionosphere](#) influences irregularities in [\\*plasma](#) density. The satellite was launched on 25 January 2018, from Kourou, French Guiana, into [\\*geostationary orbit](#).

**Global Seismographic Network (GSN)** A network of permanent seismographic stations (see **SEISMOGRAPH**) deployed evenly over the Earth's surface. The GSN comprises more than 150 stations, continuously recording seismic data at a rate of 20 samples each second, with real-time access for users. The GSN is a cooperative partnership between the Incorporated Research Institutions for Seismology (IRIS) and the United States Geological Survey.



<http://earthquake.usgs.gov/monitoring/gsn/>

- The Global Seismographic Network (GSN).

**global tectonics** The study of the relative movements of large parts of the Earth. Implicit in the phrase 'global tectonics' are the ideas that: (a) the movements may not have involved lithospheric **\*plates** (indeed, the earliest supposed plate movements are in the early **\*Proterozoic**); (b) energy changes in one part of the Earth have repercussions in other parts; and (c) large-scale tectonics have effects on many other systems, e.g. global weather patterns, evolutionary change, and the formation of natural resources.

**global warming potential (GWP)** A ranking of the absorptive capacity of the principal **\*greenhouse** gases expressed as the atmospheric warming effect of each compared with that of carbon dioxide, which is given a value of 1. The values take account of the wavelengths at which each gas absorbs radiation and its atmospheric residence time, and values are given over a specified number of years. On this scale, the GWP of methane is 56 (20 years), 21 (100 years), and 6.5 (500 years), nitrous oxide is 280 (20 years), 310 (100 years), and 170 (500 years), HFC-23 is 9100 (20 years), 11 700 (100 years), and 9800 (500 years), and sulphur hexafluoride is 16 300 (20 years), 23 900 (100 years), and 34 900 (500 years).

**global water budget** See **WATER BUDGET, GLOBAL**.

**Globigerina ooze** Deep-sea ooze in which at least 30% of the sediment consists of **\*planktonic** foraminifera (**\*Foraminiferida**) including chiefly *Globigerina*. It is the most widespread **\*pelagic** deposit, covering almost 50% of the deep-sea floor, and it covers most of the floor of the western **\*Indian Ocean**, the mid-**\*Atlantic Ocean**, and the equatorial and S. **\*Pacific**.

Species occurring in this deposit have been used to establish climatological and temperature criteria. *Globorotalia menardii* is supposed to indicate warmer conditions and *Globigerina pachyderma* to indicate colder temperatures. Another foraminiferan, *Globorotalia truncatulinoides* can coil in either a left- or right-handed manner and it is suggested that right **\*coiling** indicates warmer conditions and left coiling colder.

**Gloger's rule** Individuals of many species of insects, birds, and mammals are darkly pigmented in humid climates and lightly coloured in dry ones. This may well be a camouflage adaptation (moist habitats are usually well vegetated and tend to lack pale colours). There are many exceptions to this so-called rule. *See also ALLEN'S RULE; BERGMANN'S RULE.*

**glomeroporphyritic** Applied to those **\*phenocrysts** which have attached together to form clusters set within a finer-grained **\*groundmass**. The phenocryst clusters can form by the aggregation of separate **\*crystals** suspended in a **\*melt** or by partial disaggregation of crystal accumulations on the walls of the **\*magma chamber**.

**GLORIA** *See GEOLOGICAL LONG RANGE INCLINED ASDIC.*

**Glory** *See A-TRAIN; PARASOL.*

**Glossifungites** An assemblage of **\*trace fossils** that includes separate vertical, U-shaped, or sparsely branched **\*burrows**. These may have been temporary or permanent dwelling places, depending on the feeding habits of the animals concerned, i.e. whether they were foragers or suspension feeders. The Glossifungites assemblage is characteristic of marine **\*littoral** and shallow **\*sublittoral** environments.

**Glossopteris flora** The **\*Permian** glacial deposits of S. Africa, Australia, S. America, and Antarctica are succeeded by beds containing a **\*flora** very different from that of N. America and Europe. The flora of the south grew in a cold, wet climate, while that of the north existed under warm conditions. Plants with elongate, tongue-shaped leaves dominated the southern flora, with the genera *Glossopteris* and *Gangamopteris* being among the best known. Of these two, the genus *Glossopteris* gives its name to the flora. *Glossopteris* is characterized by a leaf with a fairly well-defined midrib and a reticulate (net-like) venation. *G. indica* is the last species

referred to the genus and the order Glossopteridales. It is known from the \*Trias of India.

**gloup** A blow-hole. See COASTAL PROCESSES.

**glow curve** See THERMOLUMINESCENCE.

**GMES** See COPERNICUS.

**GMS** See GEOSTATIONARY METEOROLOGICAL SATELLITE.

**gnamma** See PANHOLE.

**Gnathostomata** 1. Superclass comprising all vertebrates with true jaws. 2. (phylum \*Echinodermata, class \*Echinoidea) Superorder of echinoids, in which the \*periproct is outside the apical system, there are no compound \*ambulacral plates, a lantern (see ARISTOTLE'S LANTERN) and girdle are present, and the teeth are keeled. It includes the orders Holoctypoida (\*Jurassic–Recent), and Clypeasteroida (sand dollars, \*Palaeocene–Recent).

**gneiss** (*adj.* gneissose) General petrological term applied to coarse-grained, banded rocks that formed during high-grade \*regional metamorphism. The banding (gneissose banding, or gneissosity) is a result of the separation of dark minerals (e.g. \*biotite, \*hornblende, and \*pyroxenes) and the light-coloured quartzofeldspathic minerals. A gneissose rock may be described more strictly by adding a qualifying prefix, e.g. biotite gneiss, hornblende gneiss, or \*pelitic gneiss.

**gneissose banding** See GNEISS.

**gneissosity** See GNEISS.

**Gnetales** Plants, with a fossil record from the early \*Cretaceous, of particular interest because they show features of both conifers and \*angiosperms. It has been suggested that the Gnetales are the closest living relatives of angiosperms. Although commonly classed as \*gymnosperms, they possess flower-like structures and are sometimes grouped with the angiosperms in the subdivision anthophytes. See also BENNETTITALES.

**gnome** A \*transient luminous event that is believed to be a compact variety of \*blue starter.

**gnomon** A device that is erected beside a rock or structure of interest to provide a reference for the vertical and colour. It was used during the manned exploration of the Moon.

**goaf** **1.** Waste material. **2.** An area from which **\*coal** has been removed.

**gobi** An Asian name for **\*desert pavement**.

**GOCE** See GRAVITY FIELD AND STEADY-STATE OCEAN CIRCULATION EXPLORER.

**GOES** See GEOSTATIONARY OPERATIONAL ENVIRONMENTAL SATELLITE.

**goethite** A hydrous iron oxide,  $\alpha$ -FeO(OH), which forms a series with lepidocrocite,  $\gamma$ -FeO(OH); sp. gr. 4.0; **\*hardness** 5; reddish-brown; **\*massive** or earthy; occurs as an alteration product of iron-bearing minerals in association with **\*limonite** or **\*hematite**. It is named after J. W. von Goethe (1749–1832).

**Göktürk** An Earth-observation satellite mission of the Turkish Ministry of National Defence that provides high-resolution imagery for a wide range of applications. Göktürk-1 was launched on 5 December 2016, from Kourou, French Guiana, into a Sun-synchronous orbit at an altitude of 695 km; its operational status is unknown. Göktürk-2, for the Undersecretariat for the Defence Industry, was launched on 18 December 2012, from the Jiuquan Satellite Launch Center, China, into a Sun-synchronous orbit at an altitude of 684 km.

**GOLD** See GLOBAL-SCALE OBSERVATIONS OF THE LIMB AND DISK.

**gold** Malleable metallic element, Au; sp. gr. 19; **\*hardness** 2.5; bright yellow, but yellowish-white if silver is also present; occurs native as grains, threads, wire, sponge, and in many other forms, and may be combined with a variety of other metals including silver, copper, palladium, rhodium, or bismuth; occurs in **\*placer** deposits, in residual soils, and in veins associated with a variety of **\*igneous** and **\*sedimentary rocks**.

**golden algae** See CHRYSOPHYCEAE.

**golden-brown algae** See CHRYSOPHYCEAE.

**golden spike** See BOUNDARY STRATOTYPE. Compare SILVER SPIKE.

**Goldschmidt, Victor Moritz** (1888–1947) A Norwegian geochemist, Goldschmidt developed some of the basic techniques of physics for use in *\*geochemistry*, including *\*X-ray* diffraction and spectroscopy. He did important work on *\*metamorphism* and *\*trace* elements, described in his textbook *Geochemistry*, published posthumously in 1954.

**Goldschmidt's rules** In a system where the two variables, temperature and pressure, are controlled externally, the number of *\*phases* will not usually exceed the number of components. In geology the maximum number of naturally occurring minerals in a rock is equal to the number of components where a given mineral assemblage is stable over a range of temperatures and pressures.

**GomSpace Express (GOMX)** A commercially available 2-unit *\*CubeSat* kit provided by GomSpace Express, Aalborg, Denmark. GOMX-1 receives signals from aircraft over the ocean and also carries a camera for colour Earth observation. It was launched on 21 November 2013, from Yasný Cosmodrome, Russia. GOMX-3 is a collaboration between the *\*European Space Agency* and GomSpace, to demonstrate *\*nanosatellite* capabilities in attitude control, radio-frequency sensing, and high-speed data downlinking. It was launched on 19 August 2015, from the Tanegashima Launch Center, Japan. GOMX-4A and -4B are 6-unit CubeSats that demonstrate key technologies in handling satellite formations. They were launched on 4 February 2018, from the Jiuquan Satellite Launch Center, Japan.

**Gomphotheriidae** (order *\*Proboscidea*) Extinct family of mastodons, characterized by the development of multiple accessory tooth cusps and believed to be ancestral to the later mastodons, though not to modern elephants. These long-jawed mastodons or gomphotheres were one of three distinct proboscidean lines established by the *\*Miocene*. They in turn diverged into a variety of descendant lines, several of which had highly specialized lower jaws. The family persisted into the *\*Pleistocene* in both Old and New Worlds. Long snouts appeared in the earliest forms and in the Miocene Gomphotherium the lower jaw and premaxilla (front bone of the upper jaw) were very long, both bearing tusks, which may have been used for digging, and probably permitting only a short trunk. In later forms the face became shorter and the trunk presumably longer. There were many genera and species. The evolutionary importance of the gomphotheres lies

in the probable parallel between their development and that of the line which led to the true elephants.

**GOMS-2** See [ELECTRO-L](#).

**GOMX** See [GOMESPACE EXPRESS](#).

**gonatoparian suture** See [CEPHALIC SUTURE](#).

**Gondwana** Former supercontinent of the southern hemisphere from which S. America, Africa, Madagascar, India, Sri Lanka, Australia, New Zealand, and Antarctica are derived. Their earlier connection explains why related groups of plants and animals are found in more than one of the now widely separated southern land masses. Examples include: the [\\*Dipnoi](#) (lung-fish) common to S. America, Africa, and Australia; [\\*marsupial](#) mammals found today in Australia and for most of the [\\*Cenozoic](#) in S. America; and the monkey-puzzle tree (*Araucaria*) common to S. America and Australia.

**Gondwanaland** The name Eduard [\\*Suess](#) originally gave to the former supercontinent now more commonly known as [\\*Gondwana](#).

**goniatites** See [AMMONOIDEA](#).

**goniometer** See [GONIOMETRY](#).

**goniometry** The technique for measuring the [\\*interfacial](#) angles of [\\*crystals](#). The instrument used, a goniometer (or contact goniometer) resembles a 180° protractor with a pivoted straight edge at the point of origin. The angle between the normals to [\\*crystal](#) faces is then read directly from the graduated scale. More accurate measurements on very small crystals may be made using an optical system and a telescope (optical goniometry), with the crystal mounted on a rotating [\\*stage](#) fitted with a vernier scale. These instruments are known as 'one-circle' or 'two-circle' reflecting goniometers. The crystal faces are arranged with the crystal edges parallel to the axis of rotation of the goniometer head.

**Gorda Plate** A minor lithospheric [\\*plate](#) which is subducting under the [\\*North American Plate](#) to the west of Oregon. The [\\*subduction zone](#) is not marked by an oceanic [\\*trench](#), but inland there is the andesitic volcanic chain of the Cascade Mountains. The Gorda Plate is separated from the

**\*Juan de Fuca Plate** by a **\*transform fault**; both are remnants of the once-large **\*Farallon Plate**.

**Gore** The basal **\*series** in the New Zealand **\*Triassic**. It is overlain by the **\*Balfour**, and comprises the Malakovian, Etalian, and Kaihikuan **\*stages**.

**Gorstian** A **\*stage** of the Late **\*Silurian**, underlain by the **\*Homerian** and overlain by the **\*Ludfordian** and dated at 422.9–421.3 Ma ago (Int. Commission on Stratigraphy, 2004).

**GOSAT** See GREENHOUSE GASES OBSERVING SATELLITE.

**gossan** Near-surface, iron oxide-rich zone overlying a sulphide-bearing **\*ore** deposit, caused by the **\*oxidation** and **\*leaching** of sulphides. Useful in mineral exploration as a visible guide to sulphide mineralization by its yellow or red colour.

**Gothenburg** See BRUNHES.

**Gothian** A **\*stage** of the Lower **\*Proterozoic**, from about 2100–1600 Ma ago, of the Baltic Shield region, underlain by the **\*Svecofennian** and overlain by the **\*Jotnian**. According to Van Eysinga, 1975, Gothian is synonymous with Karelian.

**Gothian orogeny (Gothic orogeny)** See DALSLANDIAN OROGENY; GRENVILLIAN OROGENY.

**gouge** **\*Clay** filling in a mineral vein, or clay material between **\*fault** planes produced by movement along the fault.

**GovSat-1** A communication satellite operated as a joint venture between the Luxembourg government and Société Européenne des Satellites. It was launched on 31 January 2018, from Cape Canaveral, into a **\*geostationary orbit** at 21.5°E.


**GPM** See GLOBAL PRECIPITATION MEASUREMENT.

**GPS** See GLOBAL POSITIONING SYSTEM.

**Grabau, Amadeus William** (1870–1946) An American geologist and palaeontologist, Grabau was a professor at Columbia University, New York, and later worked on the geologic survey of China. In 1940 he developed a

theory of rhythms in the growth of the Earth's \*crust, and of repetitions in mountain building.

**graben** A downthrown, linear, crustal block, bordered lengthways by normal \*faults. The upstanding blocks on either side may have been lowered by erosion. It is a structural feature, which may be of considerable length, caused by the relative lowering of a block between two faults or \*fault zones. The faults are commonly high-angle, normal faults which are parallel in their \*strike direction. Half-graben are bounded on only one side by one or more faults and are associated mainly with \*tilt-block tectonics. Graben with a regional extent and topographic expression can also be called \*rift valleys. Compare HORST.



**Graben (includes half-graben)**

**grab sampling** See SAMPLING METHODS.

**GRACE** See GRAVITY RECOVERY AND CLIMATE EXPERIMENT.

**grade** 1. Group of things all of which have the same value. 2. A balanced condition, especially of a river (river or stream grade) when it has just sufficient energy to transport the load supplied from the drainage basin; a balance between erosion and deposition. The concept has also been applied to hillslopes ('graded slopes') that are stable dynamically and so maintain themselves in the most economical configuration. The term is no longer used widely as it oversimplifies the issues involved. 3. The fraction of a \*sediment falling within a particular size limit, e.g. sand grade, silt grade, and boulder grade. See PARTICLE SIZE. 4. The quality of a mineral ore. 5.

Classification of an ore by the quantity or purity of the mineable metal in an orebody. **6.** In civil engineering, the gradient of a road. **7.** Distinctive functional or structural level of complexity in the organization of an organism. Thus fish, *\*amphibians*, *\*reptiles*, and *\*mammals* represent successive *\*vertebrate* grades. Grades may occur within a single lineage; or the same grade may be achieved independently in different ones (e.g. warm-bloodedness evolved independently in birds and mammals). **8.** See METAMORPHIC GRADE.

**graded bedding** *\*Sedimentary structure* in which there is an upward gradation from coarser to finer material, caused by the deposition of a heterogeneous suspension of particles. The feature may be used to establish the 'way up' or natural succession of *\*strata*.

**graded reach** A length (reach) of stream channel whose gradient and cross-sectional form have become adjusted to carry just the discharge and sediment load that are normally supplied from upstream. Such a reach is said to be in equilibrium. Early definitions emphasized the smooth long profile of such a reach; it is now realized that a profile may be irregular and the reach still be graded.

**graded sediment** **1.** A sedimentary deposit which is sorted (see SORTING) with the coarsest grain size at the base and the finest at the top is termed 'normally graded'; a *\*sediment* which is sorted with the finest at the base and coarsest at the top is termed 'inverse-graded'. **2.** A well-sorted sediment.

**graded slope** See GRADE.

**gradient** The rate of change of a function (e.g. elevation pressure or temperature), at right angles to the isolines.

**gradient wind** A non-*\*geostrophic wind* that blows parallel to the *\*isobars*. It may be expressed by  $V$  in the equation:  $G = 2D\omega V \sin \phi \pm DV^2/r + F$ , where  $G$  is the pressure gradient,  $D$  the air's density,  $\omega$  the angular velocity of the Earth's rotation about its axis,  $\phi$  the latitude,  $r$  the radius of curvature of the air's path,  $F$  the friction, and  $V$  the air's velocity when the forces are in balance. The sign before the second term is + when the motion is around a cyclonic (low-pressure) centre, - when the motion is around an anticyclonic (high-pressure) centre. See also GEOSTROPHIC WIND.

**grading curve (soil grading curve)** Graph of grain size (horizontal logarithmic scale) against percentage distribution (vertical arithmetic scale); a point on the curve indicates the percentage by weight of particles smaller in size than the grain size at the given point.

**gradiometer** Any instrument that measures the gradient of a potential field rather than its absolute value. See [GRAVITY METER](#); [MAGNETOMETER](#).

**gradualism** See [PHYLETIC GRADUALISM](#).

**grain** 1. A [\\*detrital](#) mineral or rock fragment (particle), of [\\*sand](#) size. 2. Quarrying term, for the parting fabric of a rock (i.e. the direction in which a rock is most easily split by a quarry worker).

**grain boundary sliding** See [CREEP MECHANISMS](#).

**grain flow** The movement of [\\*sediment](#) under gravity where the sediment is supported by direct grain-to-grain contact. This differs from turbidity flow, where the sediment moves under the influence of gravity in a turbulent flow of water.

**grain roughness** See [BED ROUGHNESS](#).

**grain shape** See [PARTICLE SHAPE](#).

**grain size, igneous rocks** Arbitrarily defined limits for the sizes of [\\*crystals](#) within an [\\*igneous](#) rock. Commonly accepted grain-size ranges are:

Very coarse	>3 cm
Coarse	5 mm to 3 cm
Medium	1–5 mm
Fine	<1 mm
Glassy	no grains present

For sedimentary rocks, see [PARTICLE SIZE](#).

**grainstone** In the Dunham classification, a grainstone is defined as a [\\*limestone](#) consisting of [\\*grain-supported](#) particles without any mud [\\*matrix](#).

**grain-support** A sedimentary [\\*fabric](#) in which the particles are in contact with one another and form the mechanical framework of the rock or

\*sediment.

**granite** A light-coloured, coarse-grained, \*igneous rock, consisting of \*essential \*quartz (at least 20%), \*alkali feldspar, \*mica (\*biotite and/or \*muscovite), with or more commonly without \*amphibole, and \*accessory \*apatite, \*magnetite, and \*sphene. \*Hypersolvus granites are characterized by one type of alkali feldspar, usually \*microperthite, whereas subsolvus granites are characterized by two types of alkali feldspar: \*microperthite and \*albite. Granite can be formed by \*partial melting of old \*continental crust, on a local scale by *in situ* replacement of continental crust (\*granitization), by \*fractional crystallization of \*basalt \*magma, or by a combination of these processes.

**granite minimum** In the \*ternary system \*quartz–\*albite–\*orthoclase, the common low-temperature point marking either the last of a \*melt to crystallize after \*fractional crystallization has occurred, or, conversely, the first fraction of any solid mixture of \*alkali feldspars and quartz to melt upon heating. As this liquid has a very similar composition to many \*granites, the temperature point at which it forms is known as the granite minimum.

**granitic layer** The upper 10–12 km of the \*continental crust that overlies the \*Conrad discontinuity. The name is based on the common occurrence of \*granites near to the surface, and average \*seismic velocities and densities are consistent with a \*granodioritic composition.

**granitization** The conversion of crustal rocks to a granitic \*mineral assemblage by the action of metasomatic fluids (see METASOMATISM) without going through the magmatic stage. The \*essential chemical components of \*granite are introduced into the solid parent rock, and those elements not required are removed from the solid rock by the percolation of metasomatic fluids along grain margins and the \*diffusion of ions through \*crystals. ‘Granitization’ was proposed as a process for generating granites in order to get around the space problem associated with injecting large volumes of granitic \*magma into the crust; the idea is no longer much in vogue.

**granitoid** A term used to encompass the granitic rock types: \*alkali-feldspar granite, \*granite, \*granodiorite, and \*tonalite as defined in the

classification of rocks by the International Union of Geological Sciences (IUGS).

**granoblastic** A *textural* term referring to a mosaic of equidimensional *anhedral* grains in *metamorphic rocks*. If inequant grains such as *micas* are present, they are randomly oriented.

**granodiorite** A coarse-grained *igneous* rock consisting of *essential quartz*, *plagioclase feldspar*, *alkali feldspar*, *biotite*, and *hornblende*, with *accessory sphene*, *apatite*, and *magnetite*. Plagioclase is the dominant feldspar in the rock, equal to or greater than two-thirds of the total feldspar present. Where alkali feldspar is completely, or almost completely suppressed, the rock is known as a 'trondjemite' or, more generally, 'plagiogranite'. Granodiorites are commonly found as *intrusions* into the *crust* above *subduction zones*, and are the *plutonic* equivalent of *extrusive dacites*.

**granofels** A little-used name for a massive *granoblastic metamorphic rock*, consisting of *equant* grains of *quartz* and *feldspar*, hence lacking a *foliation*. Minor *mafic minerals* consist of *pyroxene* and *garnet*. The anhydrous mineral assemblages of such rocks are produced under the high-grade, anhydrous *granulite* facies conditions found at the base of the *crust*.

**granophyre** A light-coloured, medium-grained, *igneous* rock with a *granite* mineralogy but characterized by the development of a *granophyric* texture.

**granophyric (micrographic)** Applied to a fine-scale intergrowth of *quartz* and either *alkali feldspar* or *plagioclase*, found as *interstitial*, late-stage products in the *groundmass* of *granites*. The texture is formed by simultaneous and rapid crystallization of the two *phases* from the late-stage liquid trapped between the earlier formed crystals.

**granular** 1. Applied to the *texture* of equigranular *igneous* rocks with a *grain* size ranging from 0.05 to 10 mm. 2. Applied to the *form* of a *mineral aggregate* when it is composed of grains. Evenly granular mineral aggregates have a saccharoidal *texture*.

**granular iron formation** An iron-rich *formation* developed from well-sorted chemical sands and comprising discontinuous layers probably

resulting from the action of waves, currents, and storms. Layers are rare and seldom more than a few metres thick. Three types are often recognized: *\*Algoma*, *\*Rapitan*, and *\*Superior*. Compare BANDED IRON FORMATION.

**granulation** The comminution, during deformation, of crystals into smaller grains of equal size which are then able to rotate in relation to one another. The process is a common feature of *\*plastic* deformation in *\*granitoids*.

**granule** A particle between 2 and 4 mm in size. See PARTICLE SIZE.

**granulestone** A *\*siliciclastic* rock consisting of *\*granules*.

**granulite** A coarse-grained, equigranular *\*metamorphic rock*, consisting of *\*quartz*, *\*feldspar*, and the anhydrous *\*ferromagnesian minerals* *\*pyroxene* and *\*garnet*. There is some confusion over the use of the term granulite, different authors using the name in different ways. Consequently, *\*basic* granulites, rich in the ferromagnesian minerals *\*orthopyroxene* and *\*clinopyroxene*, are better termed ‘pyroxene gneisses’, whilst *\*acid* granulites, rich in quartz and feldspar, are better termed ‘charnockitic gneisses’. These rock types are thought to be formed by *\*metamorphism* of deep crustal rocks which have suffered earlier dehydration by the removal of a wet *\*granite* *\*melt*.

**granulite facies** A set of metamorphic *\*mineral* assemblages produced by *\*metamorphism* of a wide range of starting rock types under the same metamorphic conditions and typically characterized by the development, in rocks of *\*basic* *\*igneous* composition, of the mineral assemblage *\*clinopyroxene–\*plagioclase–\*orthopyroxene–\*quartz*. Other rocks of contrasting composition, for instance *\*shales* or *\*limestones*, would each develop their own specific mineral assemblage, even though they were all being metamorphosed under the same conditions. The variation of mineral assemblage with starting rock composition reflects a particular range of pressure, temperature, and  $P(\text{H}_2\text{O})$  conditions. Experimental studies of mineral  $P$ – $T$  stability fields indicate that the facies represents high-pressure/high-temperature conditions which can be met near the base of the *\*continental crust*.

**granulometry** The measurement of grain sizes.

**grapestone** A composite *\*carbonate* *\*grain*, consisting of an aggregate of carbonate *\*peloids* or other particles, bound together by *\*algae* or micritic

cement (see MICRITE). The aggregate has an appearance resembling a bunch of grapes, and forms in low-energy, \*subtidal areas.

**graphic** A \*texture seen in \*granites and \*pegmatites and characterized by a coarse intergrowth of \*quartz and \*alkali feldspar on a scale of tens of microns. The texture can be very coarse when developed in certain pegmatites, the intergrowths developing on a scale of several millimetres, and is easily seen in hand specimen, looking rather like cuneiform (i.e. wedge-shaped) writing. When developed on a microscopic scale, graphic texture is termed \*granophyric texture.

**graphic log** An outline description of a particular sequence of rocks, made in the field and set out in columnar form. It is usual to sketch in each succeeding rock unit with its appropriate \*lithologic symbol, and draw thicknesses of units to scale. Subsequent columns contain data concerning: \*particle size (shaded in according to a horizontal scale); \*sedimentary structures; \*fossils; colour; palaeocurrent directions (see PALAEOCURRENT ANALYSIS); \*weathering; \*sorting; and any other salient features gathered under the general heading of 'remarks'. See also LITHOLOGIC SYMBOL.

**graphite** Pure carbon, C; sp. gr. 2.1; \*hardness 2; greyish-black; feels soft and greasy; good basal \*cleavage; scaly, columnar, granular, or earthy; occurs in veins and may be disseminated through rocks as a result of \*metamorphism of original carbon-rich sediments. It is used as a lubricant, electrical conductor, and in the manufacture of crucibles and paint.


**graphoglyptid** A \*trace fossil comprising a system of horizontal tunnels that were used as permanent dwellings and to produce or trap food.

**graptolite** See GRAPTOLITHINA.

**Graptolithina (graptolites)** (phylum \*Hemichordata, subphylum Stomochordata) An extinct class of stick-like, colonial, marine organisms that existed from the Middle \*Cambrian to the Lower \*Carboniferous. Their fossils are used to establish a stratigraphical timescale for the Lower \*Palaeozoic. There were two principal orders, \*Dendroidea and \*Graptoloidea, and a number of minor, short-lived orders.

**Graptoloidea** (subphylum Stomochordata, class Graptolithina) An order of graptolites that existed from the Lower \*Ordovician to Lower \*Devonian.

The *\*rhabdosomes* had up to eight *\*stipes* in early forms, but two and finally one in later forms. The *\*thecae* are of only one type, equivalent to the *\*autotheca* of *\*Dendroidea*, occur on one or both sides of the stipe, and vary quite widely in morphology. In adults, there is always a *\*sicula* bearing a *\*nema*. There are three suborders: one, to which no name has been assigned, comprises a single, *\*paraphyletic* family, Anisograptidae. The other two are Dichograptina and Virgellina.



Graptoloidea (*Saetograptus chimaera*)

**grass minimum temperature** Minimum temperature recorded in open ground at night by a thermometer whose bulb is exposed over the tips of short grass.

**graupel** Soft hail, composed of particles resembling small snowballs, and formed by accretion when a snowflake falls through supercooled water droplets which freeze on contact and cover the flake.

**gravel** In the British classification for **\*particle sizes**, grains with diameters between 2 and 60 mm.

**gravimeter (gravity meter)** An instrument for measuring the **\*gravitational** acceleration. Most field instruments are relative instruments, i.e. they determine the difference in gravitational acceleration between two or more points. Most operate by determining the change in extension of a spring loaded with a constant mass. Laboratory instruments can be absolute and are based on dropping masses, vibrating strings, or oscillating pendulum systems.

**gravimetric analysis** A method of chemical analysis which involves the quantitative precipitation of an element or elements from solution in a compound whose composition is accurately known and which may be isolated in a pure form prior to weighing. When new electronic techniques of elemental analysis were introduced the term 'wet chemistry' came to be applied to the classical gravimetric techniques.

**gravimetry** The science of measuring **\*gravitational \*acceleration** at different locations.

**gravitational acceleration (g)** Following Newton's law, the force  $F$  between two masses  $m_1$  and  $m_2$ , separated by a distance  $r$ , is given by  $F = Gm_1m_2/r^2$ , where  $G$  is the **\*gravitational constant**. The gravitational acceleration,  $g$ , is then given by  $g = F/m_1 = Gm_2/r^2$  and is measured in **\*gravity units**.

**gravitational constant (G)** The constant of proportionality relating the gravitational force ( $F$ ) between two masses ( $m_1$  and  $m_2$ ) separated by a distance  $r$ .  $F = Gm_1m_2/r^2$ . The value of  $G$  is  $6.672 \times 10^{11} \text{ Nm}^2/\text{kg}^2$ .

**gravitational equipotential** A surface of equal **\*gravitational** acceleration, e.g. the **\*geoid**.

**gravitational field** Theoretically, the **\*gravitational** attraction of a unit mass extends to infinity; in practice, this field is the space within which the mass's gravity is an effective force.

**gravitational potential energy (GPE)** The energy that would be needed to move a massive body against the gravitational force exerted by the Earth's

gravitational field.

**gravitational signature** See SIGNATURE.

**gravitational water** Water that moves through soil under the influence of gravity and that must be removed before the soil can attain **\*field capacity**.

**gravity anomaly** The **\*gravitational** acceleration remaining after allowing for other factors. Common anomalies are **\*Bouguer**, **\*free-air**, and **\*isostatic**, but a gravity anomaly is any acceleration remaining after allowing for different gravitational attraction models.

**gravity assist** A technique used to accelerate or slow spacecraft without consuming fuel by transferring **\*angular momentum** from a planet to the spacecraft. The craft approaches the planet, is accelerated towards it by gravitational force, and as it leaves decelerates to its original speed relative to the planet, but not relative to the Sun. If a spacecraft approaches from behind, travelling in the same direction as the planet, it will accelerate relative to the Sun by acquiring angular momentum; if it approaches from ahead, travelling in the opposite direction to the planet, it will slow relative to the Sun by losing angular momentum to the planet.

**gravity corer** A sampling tube that penetrates into marine or lake sediments under its own weight. See also HYDRAULIC CORER.

**Gravity field and steady-state Ocean Circulation Explorer (GOCE)** A core mission of the **\*European Space Agency's** Earth Explorer Program that aims to provide new understanding of the physics of the Earth's interior, including **\*geodynamics** associated with the **\*lithosphere**, **\*mantle** composition and **\*rheology**, and uplifting and subduction processes, measurements to permit a precise estimate of the marine **\*geoid** and combined with satellite altimetry of absolute ocean circulation and transport of mass, mapping of short-wavelength features of the dynamic topography to an accuracy of 1–2 cm on a global scale, identification of features of the **\*geostrophic** current field, estimation of the thickness of polar ice sheets, and to provide a highly accurate reference system for global heights. The satellite was launched on 17 March 2009, from Plesetsk, Russia, into a Sun-synchronous circular orbit at an altitude of 250–270 km.

**gravity gliding** See GRAVITY SLIDING.

**gravity meter** See GRAVIMETER.

**Gravity Recovery and Climate Experiment (GRACE)** A satellite to satellite tracking (SST) mission, launched from Plesetsk, Russia, on 17 March 2002 and originally planned to continue for five years, in which two identical *\*minisatellites* flying about 220 km apart in a circular *\*polar orbit* at an altitude of about 500 km measure variations in the distance between them caused by changes in gravity due to the uneven distribution of the Earth's mass, providing long-term, accurate data for global, high-resolution models of the mean and time-variable components of the Earth's gravity field. This will enable a better understanding of ocean surface currents and heat transport, changes in sea-floor pressure, and the mass balance of *\*ice sheets* and *\*glaciers*, and it will also monitor changes in the storage of water and snow on the continents. The satellites also collect data that help in compiling a detailed profile of the atmosphere. GRACE is a collaborative project between NASA and the Deutsche Forschungsanstalt für Luft- und Raumfahrt (DLR) in Germany. The GRACE-FO (follow-on) mission was launched on 22 May 2018, from California.



<http://science.nasa.gov/missions/grace/>

- A joint DLR and NASA mission to accurately map variations in the Earth's gravity field.

**gravity separation** A concentration process based on differences in specific gravity between *\*ore* and *\*gangue* minerals. Separation is performed by shaking tables, Humphreys spirals, jigs, hydroclones, dense media, etc.

**gravity settling** The settling of heavy *\*minerals* and their accumulation on the floor of a *\*magma chamber*, e.g. gravity-accumulated *\*chromite* in some *\*ultramafic* rocks. See FRACTIONAL CRYSTALLIZATION.

**gravity sliding (slide, gravity gliding)** The movement of rock bodies in response to gravitational instability along particular planes in unstable regions which leads to the formation of *\*thrust*, *\*nappe*, and *\*slump* structures.

**gravity survey** A survey, usually a profile or grid, undertaken to determine the *\*gravitational acceleration* within an area.

**gravity tectonics (gliding tectonics)** Mechanism whereby large masses of rocks move down a slope under gravitational force, producing *\*folding* and *\*faulting* of varying extent and complexity.

**gravity unit** Not an official SI unit, but one commonly used to measure *\*gravitational* acceleration. One gravity unit is equal to  $10^{-6}$  m/s<sup>2</sup>, or 0.1 mgal.

**graywacke** See GREYWACKE.

**grazer** A herbivorous, vertebrate animal that feeds on grass and herbs. An aquatic invertebrate that feeds by scraping or rasping material from organic mats (e.g. *\*algal mats*) coating the surface of a sediment or rock, or that eats larger plants.

**greasy** Of *\*mineral lustre*, having the appearance of an oily coating, due to the scattering of light by a surface that is rough at the microscopic scale.

**great circle** A line on the surface of a sphere representing the circumference of a circle whose centre is coincident with the centre of that sphere. In making a *\*stereographic projection*, a horizontal (equatorial) projection of the sphere (i.e. as a plane at right angles to the N–S plane) is a primitive circle of given radius, and also a great circle. All other great circles are similar to lines of longitude. A line joining the points on the conceptual sphere at right angles to the horizontal plane will project on a *\*stereogram* as a straight line passing through the centre of the projection, with the N and S poles as its diameter, both of which plot coincidentally in the centre of the horizontal plane. See PLANE OF PROJECTION.

**Great Interglacial** See MINDEL/RISS INTERGLACIAL.

**Great Oxidation Event** The accumulation of oxygen in the atmosphere that occurred 2.5–2.45 billion years ago, when the atmospheric oxygen concentration rose from less than 0.1% of today's value (and probably much less) to about 5% of today's value. Photosynthesis by *\*cyanobacteria* is believed to have supplied the oxygen, but much of it would have been consumed by oxidizing methane as well as iron, sulphur, and other elements exposed to the air, and oxygen would have dissolved in surface water.

These processes would have delayed its accumulation, but why it accumulated relatively suddenly is uncertain.

**Great Red Spot** An anticyclonic storm system in the hydrogen–helium atmosphere of the southern hemisphere of *\*Jupiter*. It covers 10° of latitude, is about 12 000 km across (close to the diameter of the Earth), and has been observed for more than 300 years.

**green algae** See CHLOROPHYTA.

**greenalite** See CHAMOSITE; CHLORITE.

**greenhouse effect** The effect of heat retention in the lower atmosphere as a result of absorption and reradiation by clouds and gases (e.g. water vapour, carbon dioxide, methane, and chlorofluorocarbons) of long-wave (more than 4 μm) *\*terrestrial radiation*. The insulating effect is analogous to that of greenhouse glass (i.e. it is transparent to incoming short-wave radiation but partly opaque to reradiated long-wave radiation, although air in a greenhouse heats because it is unable to leave the building so the analogy is in a sense misleading) and alters the balance of incoming and outgoing radiation in the Earth's energy budget. Increases in the concentration of atmospheric carbon dioxide, generated for example by the combustion of *\*fossil fuels*, will result in a global increase of atmospheric temperatures if not offset by other (perhaps natural) changes. In the absence of any feedback effects, a doubling of atmospheric carbon dioxide will produce a rise in temperature of 1 °C. See also ATMOSPHERIC 'WINDOW'.

**greenhouse gas** A gas composed of molecules that absorb and reradiate infrared electromagnetic radiation. When present in the atmosphere, therefore, the gas contributes to the *\*greenhouse effect*. On Earth, the principal greenhouse gases are water vapour, carbon dioxide, methane, nitrous oxide, ozone, and certain halocarbon compounds. See GLOBAL WARMING POTENTIAL.

**Greenhouse Gases Observing Satellite (GOSAT)** A mission by the *\*Japan Aerospace Exploration Agency (JAXA)* to monitor sources and sinks of CO<sub>2</sub> on a sub-continental scale. GOSAT was launched on 23 January 2009, from the Tanegashima Space Center, Japan, into a Sun-

synchronous circular orbit at an altitude of 666 km. GOSAT-2, a follow-on mission, was launched in October 2018.

**Greenhouse Gas Satellite – Demonstrator (GHGSat-D)** A commercial mission by GHGSat Inc. of Montreal to monitor by remote sensing greenhouse gas and air-quality gas emissions from industrial sites. The satellite was launched on 22 June 2016, from Satish Dhawan Space Centre, India, into a Sun-synchronous circular orbit at an altitude of 512 km.

**greenhouse period** A time during which there were no glaciers on Earth. Sea levels were high, ocean waters were not well mixed and tended to be anoxic, and nutrients were recycled on *\*continental shelves*. In *\*cyclostratigraphy*, third-order cycles were predominant. *Compare ICEHOUSE PERIOD.*

**Greenlandian** The earliest *\*age/\*stage* of the *\*Holocene*, 11 700–8326 BP, dated from the NorthGRIP2 ice core (see *GRIP*).

**Greenland Ice Core Project** See *GRIP*.

**Greenland Ice Sheet Project** See *GISP*.

**greensand** Term applied to *\*glauconite*-rich *\*sandstones* and calcareous sandstones. The Lower and Upper Greensands of south-east England occur beneath and above the *\*Gault*.

**greenschist** A low-grade, *\*regional metamorphic* rock, containing abundant *\*chlorite* with *\*albite*, *\*epidote*, and *\*sericite*, and possessing a distinct *\*cleavage*. Greenschists form by *\*metamorphism* of *\*basic \*igneous* rocks, the chlorite component being derived from the *\*ferromagnesian* minerals present in the original rock. Chlorite imparts a green colour to the cleaved rock, hence the name ‘greenschist’.

**greenschist facies** A set of metamorphic *\*mineral* assemblages produced by *\*metamorphism* of a wide range of starting rock types under the same metamorphic conditions and typically characterized by the development of the mineral assemblage: *\*chlorite–\*actinolite–\*albite–\*epidote–\*quartz* in rocks of *\*basic \*igneous* composition. Other rocks of contrasting composition, e.g. *\*shales* or *\*limestones*, would each develop their own specific mineral assemblage, even though they are all being metamorphosed under the same conditions. The variation of mineral assemblage with

starting rock composition reflects a particular range of pressure, temperature, and  $P(\text{H}_2\text{O})$  conditions. Experimental studies of mineral  $P$ – $T$  stability fields indicates that the facies represents a range of moderate-pressure (about 4–7 kb), moderate-temperature (about 400–500 °C) conditions, usually developed in **\*orogenic** belts associated with continent–continent or ocean–continent plate **\*collision zones**.

**greenstone** A low-grade, **\*regional metamorphic** rock containing **\*actinolite**, **\*epidote**, and **\*albite**, and lacking a **\*cleavage**. Greenstones form by **\*metamorphism** of **\*basic \*igneous** rocks, the actinolite and epidote components being derived from the **\*ferromagnesian minerals** present in the original rock. Actinolite and epidote impart a green colour to the massive rock, hence the name ‘greenstone’.

**greenstone belt** Large geologic formation, up to 250 km across, that is largely of **\*Archaean** age. Greenstone belts are considered to represent ancient volcano–sedimentary **\*basins** bordered and intruded by granitic **\*plutons**. These formations represent an important phase of crustal evolution and currently it is commonly considered that they are remnants of **\*back-arc basins**.

**greisen** An altered, light-coloured, **\*igneous** rock consisting of white **\*mica** and **\*quartz**. The rock forms by the reaction of crystalline **\*granite** with hot fluorine-rich vapour derived from the crystallizing granite at a deeper level in the **\*intrusion**. The granite mineralogy is unstable in the presence of this vapour and reacts to give the stable mica-quartz assemblage. This process is termed **\*pneumatolysis**. Greisens are found as marginal modifications to granite adjacent to **\*mineral \*veins**, as thin veins and **\*dykes** in granite fissures, and as large bodies at the tops and sides of granite intrusions.

**Grenvillian orogeny** An episode of mountain building that ended about 1000 Ma ago and affected an area extending from what are now Columbia and Mexico, eastern N. America, eastern Greenland, and Scandinavia (where it is known as the **\*Dalslandian**, or Gothian orogeny). It is well exposed in the south-eastern part of the Canadian Shield along a line extending north-east from the northern shore of Lake Huron. It was caused by **\*plate motions** associated with the opening of the Atlantic and leading to the displacement of rocks in a north-westerly direction.

**Grenz horizon** See SUB-ATLANTIC.

**grey-brown podzolic** \*Soil-profile term describing \*eluviated, freely draining soils that have a distinctive \*clay-enriched B \*horizon. *See also ALFISOLS.*

**grey level** A calibrated sequence of grey tones, ranging from black to white. In \*remote sensing, electromagnetic radiation falling on a photosensitive receiver generates an electrical current proportional to the intensity of the radiation. The receiver is usually tuned to specific wavelength bands and the signal from each receiver is amplified and its intensity classified into different levels, usually 0 (black) to 256. These are the \*digital numbers of each of the \*pixel units that together make up a remotely sensed frame. In multispectral scanning systems, the digital numbers are classified into grey-scale bands, of which there are usually ten to twelve, but up to 256 shades of grey (brightness) may be used.

**greywacke (graywacke)** Texturally and mineralogically immature \*sandstones that contain more than 15% \*clay minerals. They may consist of angular to sub-rounded grains of \*quartz and \*feldspar, small pebbles, and a fine \*matrix of clay minerals, \*chlorite, and \*carbonate.

**grèze litée** A bedded hillslope deposit made up of alternate layers of coarse rock fragments (up to 25 mm diameter) and finer material, and laid down in a \*periglacial environment. It is probably formed by a combination of \*frost wedging, \*gelifluction, and the action of meltwater derived from the thawing of frozen ground.

**grid reference** In Great Britain, a map reference system, overprinted on all Ordnance Survey (OS) maps, which can describe uniquely the location of a point on the ground, on the smallest-scale maps to within 10 m. From a reference point to the west of the Isles of Scilly, the National Grid divides Great Britain (Ireland is excluded) into squares of 100 × 100 km, each designated by two letters. These squares are subdivided into 1 × 1 km squares whose western and southern margins are numbered, and on the smallest-scale maps 100 × 100 m squares are also used. To describe a particular location, the grid reference begins with the identifying letters of the relevant 100 km square (e.g. SH), followed by two digits (e.g. 60) to identify the western edge of the 1 km square and a third digit (e.g. 9) as an estimate of the distance of the point from the western margin of that square. This procedure is called 'easting'. 'Northing' is similar, starting with two

digits for the southern edge of the 1 km square (e.g. 54) and a third (e.g. 3) as an estimate of the distance from the southern edge. The full grid reference, consisting of two letters and six digits locates the point to within 100 m (SH 609 543 is the grid reference for the summit station of the Snowdon Mountain Railway). On smaller-scale maps, two additional digits, one added to the easting and one to the northing, locate the point to within 10 m.

**Griesbachian** The first *\*stage* (251–250.4 Ma ago) of the Early *\*Triassic* epoch, preceded by the *\*Changhsingian* and overlain by the *\*Dienerian*.

**Griffith cracks** See GRIFFITH FAILURE CRITERION.

**Griffith failure criterion** The two-dimensional relationship between *\*shear stress* and *\*normal stress* at the point of failure. The mechanism of failure is based on the formation, propagation, and joining of microscopic ‘Griffith cracks’ whose leading edges concentrate stress. Failure occurs when a critical stress is reached and the cracks propagate fully.

**Griffith–Murrell failure criterion** A three-dimensional modification of the *\*Griffith* failure criterion which includes the effect of intermediate *\*principal stress* ( $\sigma_2$ ) on failure.

**Grigg–Skjellerup** A *\*comet* with an orbital period of 5.09 years; *\*perihelion* date 22 July 1992; perihelion distance 0.989 AU.

**grike** Deep, tapering cleft, normally a solution-widened *\*joint*, cut into the surface of a near-level area of hard *\*limestone*. In north-western Yorkshire, England, it may be 15–60 cm wide at the surface, and commonly 0.5–3 m deep. The deepest grikes form where several joints intersect, and may develop into shallow caves. See also CLINT.

**GRIP (Greenland Ice Core Project)** A European drilling programme that extracts ice cores from the Greenland ice sheet, from which atmospheric and palaeoclimatological data are obtained. It operates at a site about 30 km from the second *\*GISP* site. On 17 July 2003, GRIP reached bedrock at 3084.99 m and drilling was terminated.

**GRM** See GENERALIZED RECIPROCAL METHOD; GYROREMANENT MAGNETIZATION.

**groin** See GROUYNE.

**Grooved Terrain** See MARTIAN TERRAIN UNITS.

**groove mark** A linear groove, cut in a muddy substrate by the dragging of an object through the \*sediment by flowing water. The orientation of the groove will be parallel to the current direction. Subsequent infilling of the groove by sediment will result in a groove cast being preserved on the base of the overlying bed.

**grossular** A member of the \*garnet group of minerals,  $\text{Ca}_3\text{Al}_2\text{Si}_3\text{O}_{12}$ ; sp. gr. 3.5; \*hardness 7; green to yellowish-brown; well-formed crystals or granular; occurs in metamorphosed impure \*limestones and as a \*detrital mineral, and is used as an abrasive or for gemstones.

**ground anchor** See ANCHOR.

**ground-control point** A point which can be mapped by Cartesian coordinates in two dimensions common to both an image and a topographical map. Such points allow orientation and the correction of distorted images.

**ground data (ground information, 'ground truth')** Information collected in the field, e.g. during a survey of land resources.

**ground frost** Condition in which the ground surface has a temperature below 0°C.

**ground ice** See ICE.

**ground information** See GROUND DATA.

**groundmass** The finer-grained material of \*igneous rocks in which larger crystals (\*phenocrysts) and \*xenoliths are enclosed. The groundmass of an igneous rock commonly represents that part of the \*magma system which has cooled rapidly. Rapid cooling initiates numerous crystal \*nucleation sites which limit the size to which crystals can grow. In \*lavas, the fine groundmass is formed when the magma is erupted on to the Earth's surface and is cooled rapidly by air convection over the flow. Compare MATRIX.

**ground moraine** See MORaine.

**ground range** In *\*radar* terminology, the distance between the *\*nadir* and an object. *See also* RANGE.

**ground roll** A *\*surface wave*, typically a *\*Rayleigh wave*, which propagates along the surface of the ground with a characteristically low frequency and low velocity, but with a relatively high amplitude. Such waves degrade the quality of the *\*seismic record* by masking reflections from the subsurface. In a marine environment, the same type of phenomenon is called ‘mud roll’.

**ground surge** *See* SURGE.

**ground truth** 1. In *\*remote sensing*, the verification of image interpretation by direct observation of the ground. 2. *See* GROUND DATA.

**groundwater** All the water contained in the void space within rocks. The term is generally taken to exclude vadose water (water travelling between the surface and the *\*water-table*). Most groundwater derives from surface sources (*\*meteoric water*); the remainder is either introduced by magmatic processes (*\*juvenile water*) or is *\*connate water*.

**groundwater facies** A *\*groundwater* of particular character and chemistry. The concept of hydrochemical facies is based on the assumption that the chemical composition of groundwater at any point reflects a tendency towards chemical equilibrium with the matrix rocks under the prevailing conditions. *See* CHEBOTAREV SEQUENCE.

**groundwater flow** The movement of water through interconnected voids in the *\*phreatic zone*. *See also* DARCY’S LAW.

**group** 1. A number of *\*geophones* whose output is summed to feed one *\*seismic* channel. A particularly large number of geophones used per channel may be referred to as a ‘patch’. *See also* ARRAY. 2. *See* FORMATION.

**group interval** The horizontal distance between the mid-points of adjacent *\*groups* of *\*geophones*.

**group speed (of wave)** In deep-water areas individual waves within a group move forward through the group at a faster speed than the group itself moves. The group form is produced by the interference of waves of different wavelengths (and therefore speeds) as they move forward from

their source. New waves form at the rear of the group, lose amplitude as they reach the front of the group, and disappear. The group speed is half that of the individual waves that it comprises.

**group velocity (U)** The speed with which an envelope of a wave train travels. The term is contrasted with the **\*phase velocity**.

**grouting** Injection of liquid **\*cement** or chemicals into the ground where they set thus impeding or preventing water flow by reducing **\*permeability** and improving the strength of rocks by filling **\*pores** and **\*fractures**. Primary injection holes are spaced at regular intervals with infilling of secondary holes where necessary. The type of grout and its **\*viscosity** depends on rock type because of varying size of pores and fractures and hydraulic resistance. The migration of grout is controlled by permeability.

**growan** See **SAPROLITE**.

**growth band (growth line)** A band or line, found in many organisms, that marks growth. Growth lines in **\*Brachiopoda**, for example, record the former position of the **\*commissure**; in **\*Gastropoda** growth may be intermittent and a band of rapid growth may be preserved in the shell.

**growth curve** A graph that records the changing isotopic levels of  $^{207}\text{Pb}$ ,  $^{206}\text{Pb}$ , and  $^{204}\text{Pb}$  brought about by the **\*radioactive** decay of  $^{238}\text{U}$ ,  $^{235}\text{U}$ , and  $^{232}\text{Th}$ . Very early in the Earth's cooling history there was no **\*radiogenic** lead. With the passage of time radiogenic  $^{206}\text{Pb}$ ,  $^{207}\text{Pb}$ , and  $^{208}\text{Pb}$  accumulated at the sites of uranium and thorium. The accumulation of radiogenic lead from an environment with constant ratios of U to Pb and Th to Pb would be expected to follow a simple growth curve, illustrating the increase of daughter **\*isotopes** with time. The curve would begin at the level of **\*primordial** lead and is normally plotted against axes of  $^{207}\text{Pb}:^{204}\text{Pb}$  (y), and  $^{206}\text{Pb}:^{204}\text{Pb}$  (x). The curve rises steeply at first, while more  $^{207}\text{Pb}$  than  $^{206}\text{Pb}$  is produced, but then flattens as  $^{235}\text{U}$  becomes depleted following its more rapid decay than that of  $^{238}\text{U}$ . Plotted in this way, the growth curve is the reverse of the decay curve. In reality, for any suite of samples, there would be a family of growth curves, each of which is specified by a particular value of the parameter  $^{238}\text{U}:^{204}\text{Pb}$ . Each of these curves represents the course along which lead isotope ratios would evolve in a system presently containing a given  $^{238}\text{U}:^{204}\text{Pb}$  ratio. As the equation

governing these graphs is only a function of time, systems of the same age but different  $^{238}\text{U}:^{204}\text{Pb}$  ratios will contain lead-isotope ratios lying along straight lines, known as **\*isochrons**. These isochrons will diverge from the assumed **\*common lead**-isotope ratio and will have slopes that are a function of age.

**growth fault** A **\*fault** occurring as a primary, post-depositional or syn-depositional feature commonly associated with **\*unconsolidated \*sediments** recently deposited in a **\*basin** which is undergoing active subsidence. Growth faults are typically **\*normal** and movement occurs at the same time as sedimentation, so the sediment is usually thicker on the down-faulted side. See **LISTRIC FAULT**.

**growth-fibre analysis** The analysis of the orientation of **\*quartz** or **\*calcite** **\*growth fibres**; changes in their orientation mark changes in the direction of **\*extension**.

**growth fibres** Elongate **\*crystals** which lie in a crack or **\*vein** parallel to the direction in which the walls of the fissure have opened (i.e. the direction of **\*extension**).

**growth line** See **GROWTH BAND**.

**growth twinning** A type of **\*crystal** twinning that occurs naturally during crystal growth where the energy state near the twin boundary is slightly higher than that of a single crystal. Most of these twinned crystals grow in a natural state by the addition of clusters of atoms a layer at a time, e.g. **\*aragonite** growth twins.

**groynes (groin)** A breakwater made from rock, concrete, wood, or metal, erected on a beach to inhibit the movement of sand and shingle and to protect against **\*longshore drift**.

**Grüneisen ratios (r)** A variety of parameters that have been determined to relate crystal lattice energy, pressure, and density. They are used mostly in determining the volume coefficient of thermal expansion for convecting **\*mantle** materials.

**grunerite** See **CUMMINGTONITE**.

**grus** See **SAPROLITE**.

**GSA** See GEOLOGICAL SOCIETY OF AMERICA.

**GSL** See GEOLOGICAL SOCIETY OF LONDON.

**GSN** See GLOBAL SEISMOGRAPHIC NETWORK.

**Guadalupian** An *\*epoch* (270.6–260.4 Ma ago) in the Middle *\*Permian* of N. America, underlain by the *\*Leonardian*, overlain by the *\*Ochoan*, and roughly contemporaneous with the *\*Ufimian* and lower *\*Kazanian* stages.

**guano** The leached residue of profuse accumulations of bird or bat excrement, rich in calcium phosphate. Such deposits are found particularly on arid oceanic islands and in caves. Guano is worked industrially as a phosphate resource.

**guard** See ROSTRUM.

**Guettard, Jean Étienne** (1715–86) A French geologist who in 1746 constructed the first mineralogical map of France, using symbols to indicate minerals and rock types. He produced the first mineralogical map of N. America in 1752 (although he never went there), and in 1766 began a survey of France with the aim of producing detailed geologic maps.

**guild** A group of species that have similar ecological roles, because they require the same resources and obtain them by similar means.

**Gulfian (Late Cretaceous)** An *\*epoch* (99.6–65.5 Ma ago) in the N. American Upper *\*Cretaceous*, underlain by the *\*Comanchean*, and overlain by the *\*Puercan* (*\*Palaeocene*).

**Gulf Stream** The most important ocean-current system in the northern hemisphere, which stretches from Florida to north-western Europe. It incorporates several currents: the *\*Florida Current*, the Gulf Stream itself, and an eastern extension, the *\*North Atlantic Drift*. The Florida Current is fast, deep, and narrow, but after passing Cape Hatteras the Gulf Stream becomes less effective at depth and develops a series of large meanders which form, detach, and re-form in a complicated manner. After passing the Grand Banks (off Newfoundland), the flow forms the diffuse, shallow, slower-moving North Atlantic Drift. The temperature (18–20 °C) and salinity (36?) tend to be seasonally constant, unlike neighbouring coastal water masses.

**gull** A tension fracture that develops when a competent rock that overlies incompetent material experiences extension due to gravitational sliding along the bedding, widening *\*joints*. Surface material often fills gulls. Small-scale cambering often causes gulling behind cliff tops where blocks have toppled over the edge, producing gulls that are wide at the top and narrow at the bottom.



[http://www.ubss.org.uk/resources/proceedings/vol17/UBSS\\_Proc\\_17\\_2\\_153-174.pdf](http://www.ubss.org.uk/resources/proceedings/vol17/UBSS_Proc_17_2_153-174.pdf)

- Describes two gull caves from the Wiltshire and Avon border.

**gulling** See GULL.

**gully** A feature of rain *\*erosion* that develops from the runoff of a violent torrent that bites deeply into topsoil and soft sediments. Gullies can develop on valley sides as valley-side gullies, and also along valley floors as *\*arroyos*. See also RILL-WASH.

**Gunflint Chert** See EOSPHAERA.

**Günz** The first of four glacial episodes (1.8–1.65 Ma ago) established by *\*A. Penck* and E. Bruckner in 1909. It is named after an Alpine river, and so the term is really applicable only to its type area, but it has come to be used much more widely. The Günz may correlate with the *\*Menapian* of northern Europe.

**Günz/Mindel Interglacial** An Alpine *\*interglacial \*stage*, that is perhaps the equivalent of the *\*Cromerian* stage of northern Europe.

**gust** Sharp, brief increase in wind strength close to the ground, caused by mechanical disturbance in an air flow. Gusts may also be generated by temperature lapse rates and by wind shear, e.g. in *\*clear-air* turbulence.

**Gutenberg, Beno** (1889–1960) A German seismologist who emigrated to the USA in the 1930s, in 1913 Gutenberg used seismic data to calculate the diameter of the *\*core*. In 1926, he established the existence of the low-velocity layer (see LOW-VELOCITY ZONE), noting that seismic waves from foci with depths of 50–250 km took longer to arrive than expected. At the

California Institute of Technology he continued his seismic studies in collaboration with C. F. *Richter*. *See also* GUTENBERG DISCONTINUITY.

**Gutenberg discontinuity** The seismic-velocity *discontinuity* between the Earth's *mantle* and *core*. The boundary is at a depth of about 2600 km and is thought to have surface irregularities of a few kilometres.

**gutter cast** An elongate *cast* found on the base of a bed and formed by the infilling of a gutter structure. The gutter is a linear to sinuous, U-shaped depression, up to 10 cm wide and of similar depth to its width, formed by fluid scour from helical vortices travelling parallel to the flow direction.

**guyot** Flat-topped submarine mountain or *seamount*, the summit of which lies 1000–2000 m below the ocean surface. The flat top may be a result of marine and/or subaerial erosion. It is named for the Swiss-American geologist and geographer Arnold Guyot (1807–84).

**G-wave** *See* LOVE WAVE.

**GWP** *See* GLOBAL WARMING POTENTIAL.

**Gymnodiniales** *See* DINOPHYCEAE.

**gymnosperm** A seed plant in which the ovules are carried naked on the cone scales, in contrast to the *angiosperms*, in which they are enclosed by an ovary. Gymnosperms date from the *Carboniferous* and subsequently dominated the floras of the world until the *Cretaceous*, since when they have been progressively displaced by the angiosperms (flowering plants).

**Gymnospermae (\*gymnosperms)** Formerly a subdivision of seed plants (*Spermatophyta*) that are now considered *polyphyletic*, so the name is no longer used formally. Traditionally the subdivision contained the extinct *Pteridospermales* (seed ferns), ginkgos, cycads, conifers, and gnetophytes. These are now ranked separately. *See also* GINKGOALES; CYCADOPSIDA; CONIFEROPSIDA; GNETALES.

**gypcrete** A gypsiferous (*see* GYPSUM) soil profile developed in arid regions. Gypcretes are formed by the precipitation of CaSO<sub>4</sub> from saline waters drawn to the surface by capillary action.

**gypsic** Applied to a **\*soil horizon** (a gypsic horizon) where secondary **\*gypsum** ( $\text{CaSO}_4$ ) has accumulated through more than 150 mm of soil, so that this horizon contains at least 5% more gypsum than the underlying horizon.

**gypsisols** A reference soil group in the World Reference Base for Soil Resources classification scheme. Gypsisols have a **\*gypsic** horizon within 100 cm of the surface, or more than 15% **\*gypsum** in the uppermost 100 cm.

**gypsum** **\*Evaporite** mineral,  $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ ; sp. gr. 2.3; **\*hardness** 1.5–2; **\*monoclinic**; clear white, but sometimes shades of yellow, grey, red, and brown; white **\*streak**; vitreous **\*lustre**; **\*crystals** usually **\*tabular**, often with curved faces, but can occur **\*massive** or granular; **\*cleavage** perfect {010}, good {100}, {011}; occurs in bedded deposits in association with **\*halite** and **\*anhydrite**. It is very insoluble and therefore the first mineral to precipitate from evaporating sea water; usually succeeded by anhydrite and halite. Occasionally it results from the reaction of sulphuric acid on **\*limestone** in volcanic areas. It can also result from the secondary hydration of anhydrite. Selenite is the colourless, transparent form; satin spar is the fibrous variety; alabaster is the fine-grained variety and can be carved.

**gyre (ocean gyre)** Circular or spiral motion of water, the term usually being applied to a semiclosed current system. A major gyre exists in each of the main ocean basins, centred at about  $30^\circ$  from the equator and displaced towards the western sides of the ocean ('western intensification'). Gyres are generated mainly by surface winds and move clockwise in the northern hemisphere and anticlockwise in the southern hemisphere.

**gyrocompass** The axis of a spinning gyroscope, within a set of gimbals, and with an unbalanced mass distribution, will precess about true North. Such an instrument can be used to make very precise determination of the direction of true North.

**gyroconic** Applied to the conch of a **\*cephalopod** when it coils relatively loosely.

**gyrogonite** **\*Fossil** cast of the female reproductive structure (nucule) of charophytes (stoneworts, see **CHAROPHYCEAE**). Charophyte nucules are small (around 400–600  $\mu\text{m}$ ), nut-like objects that have a fossil record

commencing in the [\\*Silurian](#). The various characteristic spiral ornamentations of gyrogonites make them useful [\\*index fossils](#) in some freshwater deposits. *See also* [PSEUDOSYCIDIUM](#).

**gyroremanent magnetization** A magnetic remanence acquired by magnetically anisotropic materials that are rotated within an alternating magnetic field. *See also* [ROTATIONAL REMANENCE](#).

**gyttja (nekron mud)** Rapidly accumulating, organic, muddy deposit, characteristic of [\\*eutrophic](#) lakes. The precise nature of gyttja varies with the producer organisms involved, which include small [\\*algae](#) or larger aquatic plants.

**Gzelian** *See* [GZHELIAN](#).

**Gzhelian (Gzelian)** A [\\*stage](#) in the [\\*Pennsylvanian \\*epoch](#) (303.9–299 Ma ago), preceded by the [\\*Kasimovian](#) and followed by the [\\*Asselian](#).

# H

**H** The horizontal component of the \*geomagnetic field.

**H<sub>0</sub>** See HUBBLE PARAMETER.

**haar** Especially in coastal areas of eastern Scotland and north-eastern England, common term for a sea fog: these are frequent in early summer.

**habit** The development of an individual crystal, or aggregate of crystals, to produce a particular external shape, with development depending on the conditions obtaining during formation. Individual crystals may possess habits such as acicular (needle-like), tabular (broad and flat), fibrous (hair-like), or prismatic (elongated in one direction). Aggregates of crystals may possess habits such as \*botryoidal, \*dendritic, or reniform (kidney-shaped).

**haboob** From the Arabic *habb* meaning 'to blow', a local term for a severe dust storm in northern Sudan. Typically, the storm is experienced late in the day during summer.

**hackly fracture** A sharp or jagged \*fracture typical of brittle minerals, e.g. in metals such as silver or platinum.


**HAD** See HELIUM ABUNDANCE INTERFEROMETER.

**hadal zone** The part of the ocean that lies in very deep trenches below the general level of the deep-ocean floor (the \*abyssal zone).

**hade** The angle between a \*fault plane, mineral \*vein, or \*lode and the vertical.

**Hadean (Priscoan)** An informal name for the \*eon of Earth history from the formation of the Earth 4567.17 Ma ago until the undefined start of the \*Archaean 3800 Ma ago (Int. Commission on Stratigraphy, 2004).

**Hadley cell** One of the most fundamental divisions of the global wind circulation, comprising the net ascent of air over the lowest latitudes due to convection, the compensatory downward motion in the subtropical anticyclone belt, and the resulting trade winds which blow toward the meteorological equator (intertropical convergence). It is named after George Hadley, who in 1735, seeking to account for the trade winds, proposed a single, large-scale convective cell representing a thermally driven, low-latitude atmospheric circulation.



**Hadley cell**

**Hadrosauridae (hadrosaurs)** Duck-billed *\*ornithischian \*dinosaurs*, of possibly amphibious habits, which averaged about 9m in body length. Several mummies of hadrosaur dinosaurs have been found, from which it was possible to establish that they were unarmoured and that the digits of

the forelimbs were connected by a web of skin. Hadrosaurs flourished in the Upper \*Cretaceous.

**hadrosaurs** See HADROSAURIDAE.

**Hadrynian** A \*stage (850–452 Ma ago) of the Upper \*Proterozoic of the Canadian Shield region, underlain by the \*Helikian and followed by the Early \*Cambrian.

**haematite** See HEMATITE.

**haemocyanin** (**hemocyanin**) A blue, copper-containing respiratory pigment that is found free in the blood plasma of malacostracans (\*Malacostraca) and some insects (\*Insecta) including the embryos of cockroaches and grasshoppers.

**Hagedoorn method** See PLUS–MINUS METHOD.

**hail** Form of precipitation comprising ice in the shape of balls or irregular particles (hailstones), whose concentric structure indicates a growth by \*coalescence and freezing of supercooled water drops. Hail is associated with \*cumulonimbus cloud and the larger the cloud the bigger the hailstones that fall from it.

**hair hygrometer** Instrument that indicates \*relative humidity, based on the expansion and contraction of a treated human hair.

**hairpin** A sudden change in a palaeomagnetic \*polar wander path, usually taken to indicate a tectonic collision between the tectonic plate concerned and some other plate.

**Haiyang-1B** (**HY-1B**; **Ocean-1B**) An ocean-observation satellite sponsored by the Chinese State Ocean Administration that monitors ocean colour and sea-surface temperature. It was launched on 11 April 2007, from the Taiyuan Satellite Launch Center, into a Sun-synchronous near-circular orbit at an altitude of 798 km.

**haldane** A unit of evolutionary morphological change, equal to one standard deviation per generation, and named for the Scottish physiologist and geneticist J. B. S. Haldane. See also DARWIN.

**Hale–Bopp** A \*comet with an orbital period of more than 4000 years; \*perihelion date 31 March 1997; perihelion distance 0.914 AU.

**Hale cycle** The 22-year cycle in sunspot activity that was discovered in 1925 by the US astronomer George Ellery Hale (1868–1938). The Hale cycle may result from the coupling of 11-year sunspot cycles with opposite magnetic polarization.

**half-field prism** See REFLECTOR.

**half-graben** See GRABEN.

**half-life** See DECAY CONSTANT; RADIOMETRIC DATING.

**half-shadow test (shadow test, Schroeder Van Der Kolk method)** A test used to determine the approximate *\*refractive index* of a mineral by immersion in a liquid of known refractive index. A card is inserted below the rotating *\*stage* to cut out half the light. The *\*condenser* is then adjusted to focus or blur the card. If the mineral differs in refractive index from the liquid, one side of the grain will have a distinct shadow. If the shadow is on the same side as the card the mineral has a higher refractive index than the liquid; if it is on the opposite side of the grain the mineral has a lower refractive index than the liquid.

**half-space** A mathematical model in which only one boundary exists, all others being infinitely far away. The medium under consideration is usually assumed to be perfectly homogeneous and *\*isotropic*.

**half-spreading rate** See SPREADING RATE.

**half-width ( $w_{1/2}$ )** Half the width of an anomaly measured between points with half the peak amplitude. It is used when determining depths to the centres of causative bodies in *\*gravity* and *\*magnetic profiles*.

**halides** A group of minerals which contain halogens, principally chlorine and fluorine. The group includes *\*halite* (NaCl), *\*sylvite* (KCl), and *\*fluorite* (CaF<sub>2</sub>). Halides are characterized by ionic bonding and are mainly *\*cubic* in form, soft, and generally light in weight. They frequently occur as precipitates resulting from the evaporation of saline waters.

**Halimede (Neptune IX)** A satellite of *\*Neptune* with a radius of 30 km, visual *\*albedo* of 0.16, and a *\*retrograde orbit* with a period of 1879.7 days.

**halinity** The extent to which particular water contains chloride. According to the Venice system, brackish waters (which are saline, but less so than sea

water) are classified by the chloride they contain and divided into zones. The zones, with their percentage chlorinity (mean values at limits) are: euhaline 1.65–2.2; polyhaline 1.0–1.65; mesohaline 0.3–1.0; alpha-mesohaline 0.55–1.0; beta-mesohaline 0.3–0.55; oligohaline 0.03–0.3; fresh water 0.03 or less.

***Haliomma vetustum*** (family Actinommidae) One of the earliest recorded species of *\*radiolarians*, characterized by a siliceous shell, and a member of a family whose members have a spherical or ellipsoidal shape and no internal spicule. *H. vetustum* is known from the *\*Ordovician* of Europe.

**halite (rock salt)** Mineral, NaCl; sp. gr. 2.2; *\*hardness* 2.5; *\*cubic*; perfect cubic *\*cleavage*; colourless, white, or shades of yellow, red, and blue; white *\*streak*; vitreous *\*lustre*; crystals usually cubes often with curved faces, but it can be granular and compact; widely distributed in stratified *\*evaporite* deposits, associated with other water-soluble minerals (e.g. *\*sylvite*) and with *\*gypsum* and *\*anhydrite* of various geologic ages, large masses frequently forming plugs which rise through and arch overlying *\*sedimentary rocks*, thereby forming oil traps; soluble in water; tastes salty. It is widely used as a road dressing in icy weather.

**Halley** A *\*comet* with an orbital period of 76.1 years; *\*perihelion* date 9 February 1986; perihelion distance 0.587 AU.

**Hall, James** (1761–1832) Scottish geologist and physicist who was the pioneer of experimental petrology. He was able to demonstrate (1800) that a *\*basalt \*melt* will crystallize if cooled slowly, thus giving important support to plutonist theory.

**Hall, James** (1811–98) American geologist, palaeontologist, and member of the US Geological Survey. A uniformitarian, he made detailed observations of the stratigraphy of the Appalachians. He proposed that the sediments were laid down in a trough, which subsided under their weight, before final uplift and erosion. He opposed catastrophist and contractionist hypotheses of mountain formation. See *CATASTROPHISM*; *CONTRACTING EARTH HYPOTHESIS*.

**Hall effect** A strip of metal or semiconductor carrying an electrical current within a strong, transverse, magnetic field develops a potential at right angles to both the current and the field. This forms the basis for some

sensitive **\*magnetometers**. The effect was discovered by Edwin Hall (1855–1938) in 1879.

**Hallian** See HOLOCENE.

**halloysite** See CLAY MINERALS.

**halmyrolysis** Early **\*diagenesis**, modification, or decomposition of sediments on the sea floor. For example, both the breakdown of **\*ferromagnesian minerals**, and the growth of **\*glauconite** aggregates in sea-floor sediments, are types of halmyrolytic process.

**halo** Rainbow-coloured, sometimes white, ring around the Sun or Moon, subtending an angle of 22° or, more rarely, 46°, due to refraction by ice crystals when thin **\*cirrus** cloud obscures the luminary. *Compare* CORONA.

**haloclasty** A type of physical **\*weathering** caused by the accumulation of salt crystals that is most marked in arid climates. Rainwater penetrates crevices in the rock and salts dissolve into it. As the water evaporates the salts crystallize and as the temperature rises, the crystals expand, widening the crevices. A later rainfall dissolves the crystals, which flow out of the rock, leaving flakes of rock detached.

**halocline** Water layer with a large vertical change in salinity. The halocline is usually well developed in coastal regions where there is much freshwater input from rivers producing surface waters of low **\*salinity**, a zone where salinity increases rapidly with depth (the halocline), and a deeper zone of more saline, denser waters.

**halokinesis** The study of salt tectonics, which includes the mobilization and flow of subsurface salt, and the subsequent emplacement and resulting structure of salt bodies (e.g. the **\*Mesozoic** salt domes of the North Sea basin). *See also* HALOTECTONISM.

**halophile** An **\*extremophile** (domain **\*Archaea**) that thrives in extremely saline environments.

**halotectonism** Movements of subsurface salt that involves tangential compressive stress.

**hamada (hammada)** A rocky **\*desert**, or desert region, which does not have surficial materials and which consists mainly of boulders and exposed

bedrock. Two basic types occur: stony hamada, jaggedly developed across crystalline rocks; and pebbly hamada, cut across sedimentary material and mantled with bedrock fragments.

**hammad** See [HAMADA](#).

**hammer chart** A circular graticule divided by radial and concentric lines to provide a template with many compartments. It is used when calculating the [\\*terrain correction](#) in gravity surveying.

**hammer source** In [\\*seismic refraction](#) surveys, a heavy sledge-hammer is swung on to a plate on the ground to provide a seismic impulse and trigger an appropriate recording system. It is used extensively with signal-enhancement [\\*seismographs](#), when repeat hammer blows can improve the signal-to-noise ratio. It is mainly a source of [\\*P-waves](#), but can be modified to become a source of [\\*S-waves](#).

**hand lens** A hand-held, steel- or plastic-mounted, optical magnifying system usually consisting of two or more optical elements, which is used in the field to provide an enlarged image of rocks, minerals, and fossils, ranging from 5× to 20×. Hand lenses vary in diameter from 0.5 to 1.5 cm, can be carried conveniently in a pocket, and form an essential tool in a geologist's field kit.

**hanging valley** Tributary valley whose floor is well above that of the adjacent main valley and where there is therefore often a waterfall. It is typical of glaciated uplands, where it may result from glacial widening and/or deepening of the main valley.

**hanging wall** The [\\*fault](#) block which lies above any inclined fault surface. See [FOOTWALL](#).

**hardground** A term first introduced into geologic literature in 1897, drawn from an oceanographic source but used in a narrower sense to describe a specific [\\*horizon](#) that had initially been lithified a short distance beneath the sea floor. [\\*Lithification](#) resulted from [\\*calcite](#) precipitation around local foci, giving rise to [\\*concretions](#). The [\\*sediment](#) above and surrounding the concretions remained soft and was consequently bioturbated (see [BIOTURBATION](#)). The initial [\\*cementation](#) of the hardground may have been triggered by the chemical influence of organisms that dwelt within the sediment. Progressive encroachment of the cemented areas on to the soft

sediment gradually resulted in total lithification. 'Incipient' hardgrounds represent periods of arrestation in the cementation process, with the result that nodular and soft-sediment units alternate one with the other. The cemented hardground may itself support a fauna of fixed or cemented organisms and may be bored extensively by bivalves (\**Bivalvia*). \*Pyrite films may line \*burrows, which may also be filled by \*silica.

**hardness** 1. A measure of the ability of water to form a carbonate scale when boiled, or to prevent the sudsing of soap. Permanent hardness is due mainly to dissolved calcium and magnesium sulphate or chloride, the bicarbonate ion causes temporary hardness. Dissolved carbon dioxide and the weathering of carbonate rocks are the main sources of hardness in water. 2. A physical property of minerals and one of the most useful tests for mineral identification. \*Mohs's scale of hardness (H), which ranks minerals by their hardness and thus makes possible a diagnostic test in which one mineral is used to scratch another, was introduced in 1822 and is still the standard used today. Useful tools for determining hardness are the finger nail (H about 2.5) and a penknife (H about 5.5). With a little practice, the hardness of a mineral may be determined by means of a scratch to within one or two points on Mohs's scale. *See also* VICKERS HARDNESS NUMBER.

**hardpan** Hardened \*soil horizon, usually found in the middle or lower parts of the profile, that may be \*indurated or \*cemented by a variety of possible cementing materials. *See also* CALICHE; DURICRUST; PAN.

**HARI** *See* HIGH-ASPECT-RATIO IGNIMBRITE.

**Harker diagram (variation diagram)** A diagram that shows the amount of each of the chemical constituents of a rock as a proportion of the main ingredient (usually silica). It is the earliest of the variation diagrams, having been devised in 1909 by the English petrologist Alfred Harker (1859–1939). *See also* AFM DIAGRAM; A'KF DIAGRAM.

**harmattan wind (the doctor)** Dry, dusty, north-easterly or easterly wind which occurs in W. Africa north of the equator. Its effect extends from just north of the equator in January, almost to the northern tropic in July. In W. Africa it is known as 'the doctor' because of its invigorating dryness compared with humid tropical air. The harmattan wind stream occasionally

extends south of the equator during the northern winter as an upper air wind over the south-westerly monsoon.

**harmonic** A *\*frequency* which is a whole multiple of the fundamental frequency; thus the second harmonic has twice the fundamental frequency, the third harmonic three times, etc.

**harmonic fold** A *\*fold* which maintains its geometric form, integral wavelength, and symmetry throughout a sequence of layers. Such folds form where the competent layers comprising the sequence are of similar thickness and evenly spaced, and the contrast in *\*competence* is constant between each layer and the next.

**harmonic tremor** A type of *\*seismicity* that occurs only as a result of *\*volcanicity*. The seismic vibrations have a single frequency, usually about 1–5Hz, and continue for hours and sometimes for days.

**Harnagian** A *\*stage* (459–458 Ma ago) of the Late *\*Ordovician* in the Lower *\*Caradoc*, underlain by the *\*Costonian* and overlain by the *\*Soudleyan*.

**HARP** See HYPER-ANGULAR RAINBOW POLARIMETER.

**Hastarian** A *\*stage* (359.2–348 Ma ago) of the *\*Mississippian* epoch (Int. Commission on Stratigraphy, 2004), preceded by the *\*Famennian* and followed by the *\*Ivorian*.

**hastingsite** See HORNBLLENDE.

**Haumurian** See MATA.

**Hauterivian** A *\*stage* in the European Early *\*Cretaceous* dated at 136.4–130 Ma ago (Int. Commission on Stratigraphy, 2004), preceded by the *\*Valanginian* and followed by the *\*Barremian*. See also NEOCOMIAN.

**Haüy, Abbé René-Just** (1743–1822) One of the founders of modern crystallography, Abbé Haüy was Professor of Mineralogy at the Sorbonne. He applied mathematics to mineralogy in order to determine the structure of crystals, proposing a ‘law’ of rational intercepts. His work, *Traité de cristallographie* (1822), was for a time the standard book on the subject.

**haiüyne** A *\*feldspathoid* and member of the *\*sodalite* group of minerals, formula  $(\text{Na,Ca})_{4-8} (\text{Al}_6\text{Si}_6\text{O}_{24})(\text{SO}_4,\text{S})_{1-2}$ ; sp. gr. 2.5; *\*hardness* 5.5; greyish-white; granular; occurs as a constituent of *\*alkali* volcanic rocks, often with *\*pyrite*.

**Hawaiian-Emperor chain** The Earth's largest volcanic mountain range, extending from the oldest *\*seamount*, dated at 78 Ma ago, near Kamchatka to an active submarine volcano south-east of Hawaii. The chain is regarded as a plume trace (see *MANTLE PLUME*). It forms two distinct, gentle curves, with a kink dated at approximately 43 Ma ago and thought to be related to a change in the direction of movement of the *\*Pacific Plate* over a stationary *\*hot spot*.

**Hawaiian eruption** A type of volcanic *\*eruption* that produces very fluid, mobile, basaltic *\*lava*, characterized by spectacular *\*fire* fountains, with little explosive activity. The lava flows that result are often formed by the accumulation of fire-fountained lava and are termed 'spatter-fed flows'. Hawaiian-type eruptions produce a broad, low-angle cone (a shield *\*volcano*). Compare *PELÜEAN ERUPTION*; *PLINIAN ERUPTION*; *STROMBOLIAN ERUPTION*; *SURTSEYAN ERUPTION*; *VESUVIAN ERUPTION*; *VULCANIAN ERUPTION*.

**hawaiite** See *ANDESITE*; *BENMOREITE*.

**Hawera** See *HOLOCENE*.

**Hawker** See *ADELAIDEAN*.

**Hayabusa 2** A successor asteroid sample-return mission to the first Hayabusa (see *MUSES-C*), the spacecraft was launched on 3 December 2014 and reached the near-Earth asteroid Ryugu on 27 June 2018. On 21 September Hayabusa (the name means 'peregrine falcon') released two landers from a height of 60 m, both called Minerva-II, which landed safely on the asteroid and began returning images. On 3 October Hayabusa released the lander Mascot (Mobile Asteroid Surface Scout) from a height of 56 m, carrying instruments to measure Ryugu's mineral composition, temperature, and magnetic properties.

**haze** Atmospheric condition in which horizontal visibility is reduced, but not to less than 2 km, because of the absorption and scattering of light by very small, dry particles, e.g. fine dust.

**HDR** See **HOT DRY ROCK**.

**head 1.** In certain areas of Britain, the name given to a sheet of poorly sorted, angular rock debris, mantling a hillslope and deposited by **\*gelifluction**. Similar material is commonly found as a fossil deposit in the extraglacial areas of N. America and Europe. Coombe rock is similar, but is found on chalk. **2.** See **BERNOULLI EQUATION**; **DARCY'S LAW**; **ELEVATION POTENTIAL ENERGY**; **HYDRAULIC HEAD**; **PRESSURE HEAD**.

**head cut** See **KNICK POINT**.

**heading blasting** An old method of quarry blasting in which explosive is confined in small tunnels at suitable intervals in the quarry face. For large blasts, several tunnels may be used.

**head shield 1.** The term sometimes applied to the **\*cephalon** in trilobites (**\*Trilobita**). **2.** The dorsal head covering of solid bone which occurs in many ostracoderms (**\*Ostracodermi**) (agnathan fish, see **AGNATHA**).

**headwall, glacial** The steep rock slope at the head of a **\*cirque** or **\*valley glacier**. It is a site of active **\*erosion**, perhaps by **\*frost wedging**.

**head wave** A **\*refracted** wave which enters and leaves a high-velocity medium at the **\*critical angle**. Usually the term refers to the refracted wave which arrives to give a refraction **\*first break**.

**heat capacity** The energy required to raise the temperature of a given mass of material by one degree Celsius.

**heat flow (heat flux)** A measure of the heat being conducted through the surface rocks of the Earth. It is usually determined by taking two or more temperature readings at different depths down a **\*borehole** or sampling tube, and measuring or estimating the thermal conductivities of the rocks in between. On land, measurements are usually taken at depths greater than 200 m to avoid temperature changes associated with climatic oscillations. The largest uncertainties are due to local or regional heat transference by convective circulation. In oceanic sediments the observed heat flow

decreases in proportion to  $t^{0.5}$ , where  $t$  is the age of the \*oceanic crust. Near the oceanic ridge crests the heat flow averages  $100 \text{ mW/m}^2$ , decreasing to less than  $50 \text{ mW/m}^2$  for crust older than 100 Ma. In the continents, the older crustal areas appear to have a lower heat flow, averaging about  $38 \text{ mW/m}^2$ , than the younger crust; in \*Mesozoic and \*Tertiary orogenic areas it is 60–75  $\text{mW/m}^2$ . The heat flow in the cratonic shields (see CRATON) is almost entirely accounted for by \*radiogenic heat production within the surface rocks themselves, implying depletion of radiogenic heat-producing elements in the \*crust and upper \*mantle beneath these areas. The annual global heat loss by the Earth is somewhat uncertain because of heat loss by convection, but is about  $(4.1\text{--}4.3) \times 10^{13} \text{ W}$  ( $1.3 \times 10^{21} \text{ J/year}$ ), about 75% of which is lost through the oceanic crust. Internal heat sources within the Earth are: (a) radiogenic heat-producing elements; (b) gravitational energy released by core formation; (c) exothermic chemical reactions; and (d) heat remaining from the initial accretion of the Earth, comprising the original heat of the materials, kinetic heat from early meteoritic and \*planetismal bombardment, and exothermic chemical reactions. The proportion of each source is not clear, although radiogenic heat is probably the major component; gravitational energy release may have been dominant during core formation.

**heat-flow anomaly** An area characterized by higher or lower \*heat flow than the average for the area.

**heat-flow unit (HFU)** A unit of  $1 \text{ cal/cm}^2/\text{s}$ , now replaced by SI units:  $1 \text{ HFU} = 41.8 \text{ mW/m}^2$ .

**heat flux** See HEAT FLOW.

**heat of formation** The amount of heat evolved or absorbed during the formation of one \*mole of a substance from its constituent elements.

**heave** **1.** The amount of horizontal (lateral) displacement between two sides of a \*dip-slip fault, measured at right angles to the \*strike of the fault. **2.** Lifting of earth due to frost (see FROST HEAVE TEST), overloading, swelling clay, etc. **3.** Upward in the floor of a mine due to the floor being too weak to resist the forces resulting from the weight of the overlying rock on the adjacent, supporting pillars.

**heavy liquids** See DENSITY DETERMINATION.

**heavy-medium separation** See DENSE-MEDIUM SEPARATION.

**hedenbergite** A member of the *clinopyroxene* group of *minerals* and the iron-rich *end-member* of the *diopside* series, with formula  $\text{CaFeSi}_2\text{O}_6$ ; sp. gr. 3.7; *hardness* 6; black; forms short, *prismatic crystals* or lamellar masses; occurs in metamorphosed iron-rich sediments, *skarns* and *eulysites*, and in some *igneous* ferrogabbros and *granophyres* along with fayalitic *olivine*.

**Heezen, Bruce Charles** (1924–77) An American oceanographer from Lamont-Doherty Observatory, Heezen is perhaps best known for his maps of the ocean floors, produced in collaboration with Marie Tharp. He also made important studies of the axial ridge systems of the oceans, and in 1952 found the first evidence for *turbidity* currents during his studies of the Grand Banks earthquake (1929).

**Heim, Albert** (1849–1937) A Swiss alpine geologist, Heim did important work concerned with the theory of mountain building, and on the formation of *overthrusts* and *nappes* in the Alps. He supported the idea of a *contracting* Earth. He also studied the mechanics of rock deformation, proposing that rocks can deform plastically under pressure and that the same pressure causes *metamorphism*.

**Heinrich events** The release into the ocean of large amounts of ice, as icebergs, at intervals of 8000–10 000 years, with major consequences for the circulation of ocean water and the global climate. The phenomenon was first described in 1988 by Hartmut Heinrich.

**Heinrich layers** North Atlantic sediments that have a high ratio of debris carried by ice to *Foraminiferida* shells and thus record episodes of major iceberg release or surges in the *Laurentide* ice sheet.

**Helderbergian** See ULSTERIAN.

**Helene (Saturn XIII)** One of the lesser satellites of *Saturn*, discovered in 1980 by *Voyager* 1, with a radius of 16 km; visual albedo 0.7.

**helicitic structure** Sigmoidal trails of mineral inclusions in *porphyroblasts* formed during metamorphic *recrystallization*. The sigmoidal inclusion

trails may represent remnants of a pre-existing fold-fabric preserved as the growing *\*crystal* enclosed the *\*fabric*, leaving untouched those minerals it did not require to provide its chemical components. Sigmoidal inclusion trails may also represent relative movement between the growing porphyroblast and the *\*groundmass* or external *\*fabric*. Thus, if the porphyroblast grows under the influence of *\*shear stress*, it rotates during growth, and may include relicts of any parallel external fabric as an S-shaped internal inclusion trail.

**helictite** See DRIPSTONE.

**Helikian** A *\*stage* (1600–850 Ma ago) of the Middle *\*Proterozoic* of the Canadian Shield region, underlain by the *\*Aphebian* and overlain by the *\*Hadrynian*.

**Heliopora** See OCTOCORALLIA.

**helium abundance interferometer (HAD)** An instrument used in *\*remote sensing* to determine very precisely the relative abundances of helium and hydrogen in planetary atmospheres by means of an *\*interferometer*.

**helium ‘burning’** The fusion of helium in the contracted core of a red giant star at extremely high temperatures, hotter than those reached in the *\*Sun*. It is particularly important because the reaction produces carbon and oxygen, so helping to explain their *\*cosmic* abundance. See CARBON ‘BURNING’; HYDROGEN ‘BURNING’; SILICON ‘BURNING’; NUCLEOSYNTHESIS.

**helium clock** A method for measuring the passage of geologic time based on the accumulation of helium in the Earth’s atmosphere. As one of the rare or ‘noble’ gases, helium is essentially inert chemically. There is very little of it on Earth and this is attributed to the fact that its low atomic weight allows it to escape from the Earth’s gravitational field. The mean *\*residence* time in the atmosphere is only a few million years. No helium in the atmosphere can therefore be residual since the formation of the Earth, but must be the product of *\*alpha* decay in the Earth’s crust. Thus a state of equilibrium must exist between the continual loss of helium into outer space and the supply of new *\*radiogenic* helium.

**Hellenic Arc** The mostly inactive east–west volcanic chain on the \*Hellenic Plate which is associated with the \*subduction of the \*African Plate. Crete forms part of the outer, non-volcanic arc and extinct \*volcanoes, including Santorini, form the volcanic arc, approximately 150 km to the north.

**Hellenic Plate** A \*microplate in the eastern Mediterranean which lies in the \*collision zone between the \*African Plate and the \*Eurasian Plate. The southern boundary of the Hellenic Plate is the \*Hellenic Trench, north of which there is a rising, non-volcanic arc. The Aegean Sea, further to the north, is undergoing widespread extension and subsidence.

**Hellenic Trench** A linear depression between the Ionian Basin and Crete which forms the boundary between the \*Hellenic Plate and the \*African Plate. To the east the boundary is a \*transform fault along the Pliny-Strabo Trench. Several hundred kilometres of \*oceanic crust are thought to have been subducted under the Hellenic Plate, and the Hellenic Trench may now be the junction between the African \*continental margin and the Hellenic Plate.

**Helminthoida** A two-dimensional grazing trace represented by tightly packed meanders or spiral \*trails. The \*trace fossil is related to the feeding behaviour of an efficient \*sediment-eating organism.

**helm wind** Local name for a type of lee wave frequent in winter and spring on the western (lee) slope of the Crossfell range in Cumbria, England. Associated with this cold, gusty, north-easterly wind, the helm itself is a thick bank of cloud along the mountain range, together with an outlier of narrow, almost motionless cloud away from the range.

**Helvetian** See SERRAVALLIAN.

**hematite (haematite, iron glance, kidney ore, red iron ore, specularite)** Iron \*mineral,  $\text{Fe}_2\text{O}_3$ ; one of the main ores for iron; sp. gr. 4.9–5.3; \*hardness 5–6; \*trigonal; steel-grey to black, often iridescent, compact varieties dull to bright red; red to reddish-brown \*streak; \*metallic \*lustre; crystals \*tabular or \*rhombohedral with curved, striated faces, also occurs as columnar, \*mammillated, and \*botryoidal masses; no \*cleavage; widely distributed as an \*accessory mineral in \*igneous rocks, \*hydrothermal \*veins, as a rock-forming mineral in \*sedimentary rocks, as a \*primary

**mineral**, as concretions or a cementing agent, and as a replacement for other minerals. Bedded ores of hematite form huge deposits in the **\*Precambrian** of N. America and elsewhere. Specularite is a grey or black variety of hematite with a brilliant metallic lustre that occurs in foliated masses or aggregates of tabular crystals. See **BANDED IRON FORMATION (BIF)**.

**hemera** A period of geological time determined by the maximum development of a **\*fossil** plant or animal.

**hemi-** From the Greek *hemi* meaning 'half', a prefix meaning 'half' or 'affecting one half'.

**Hemichordata (acorn worms)** Phylum first encountered in the Middle **\*Cambrian** **\*Burgess Shale**, British Columbia, Canada. The acorn worms are related to the **\*chordates**, for although they lack a **\*notochord**, the gill slits are very similar to those of primitive **\*vertebrates**.

**hemimorphism** The property of certain crystals in which there is no element of symmetry present to cause the repetition of upper-hemisphere faces in the lower hemisphere. A good example is **\*tourmaline**, where there are no horizontal **\*axes of symmetry** or **\*centre of symmetry**, but only one vertical axis of three-fold symmetry and three vertical planes of symmetry.

**hemimorphite** A **\*silicate** mineral  $Zn_4[Si_2O_7](OH)_2.H_2O$  which accompanies zinc, iron, and lead sulphides and is associated with **\*smithsonite**  $ZnCO_3$ , a principal **\*ore** of zinc; sp. gr. 3.4–3.5; **\*hardness** 4–5; **\*orthorhombic**; when **\*massive**, white, grey, yellow, brown, green, or light blue, sometimes colourless; **\*vitreous** **\*lustre**; **\*crystals** can be small, **\*tabular** grains, but normally form aggregates of radial or earthy masses; **\*cleavage** perfect prismatic {110}; occurs in the oxidation zone of lead-sulphide/zinc-sulphide deposits, in **\*hydrothermal** veins near the surface, and as a **\*primary mineral** in hydrothermal veins, associated with sulphides and **\*fluorite**.

**hemipelagic sediment (hemipelagite)** A deep-sea, muddy **\*sediment** formed close to continental margins by the settling of fine particles, in which biogenic material comprises 5–75% of the total volume and more than 40% of the **\*terrigenous** material is **\*silt**. Compare **\*pelagic sediment**.

**hemipelagite** See **HEMPELAGIC SEDIMENT**.

**Hengelo (Hoboken)** A Middle *\*Devensian* *\*interstadial* dating from about 40 000 years BP, when estimated July temperatures in the Netherlands were about 10°C (today the average is 17°C). Together with the *\*Moershoofd* and *\*Denekamp* interstadials, the Hengelo is equivalent to the *\*Upton Warren* Interstadial of the British Isles.

**Hennig, Willi** (1913–76) A German zoologist who originated *\*phylogenetics*. He was awarded his Ph.D. by the University of Leipzig in 1947 and conducted research on *Drosophila* larvae. His book, *Grundzüge einer Theorie der phylogenetischen Systematik* (1950), made no immediate impact, but when an English translation appeared in 1966, as *Phylogenetic Systematics* (with a second edition in 1979), it suddenly achieved authoritative status (comparable, perhaps, to *The Origin of Species!*). Hennig argued that, as *\*taxonomy* aims to depict relationships and the only objective meaning of ‘related’ means sharing a common ancestor, taxonomy must be based on *\*phylogeny*. This struck an immediate chord. Hennig coined such terms as *\*apomorphic*, *\*plesiomorphic*, and *\*sister groups*, and offered a redefinition of monophyly (see *MONOPHYLETIC*), which he insisted must be paramount in taxonomy.

**Hennig’s dilemma** Phylogenetic trees constructed from an examination of two or more characters (e.g. two different genes) may result in two contradictory *\*phylogenies*. In such cases no single tree can be constructed using all characters compatibly. (*\*Cladistics*, in which the dilemma may arise, was founded by the German entomologist W. *\*Hennig*.)

**Herangi** A *\*series* in the Lower *\*Jurassic* of New Zealand, underlain by the *\*Balfour* (*\*Triassic*), overlain by the *\*Kawhia*, and comprising the Aratauran and Ururoan *\*stages*. It is roughly contemporaneous with the *\*Lias* and *\*Aalenian*.

**Herbert Smith refractometer** See *REFRACTOMETER*.

**Hercynian orogeny** A major and prolonged episode of mountain building that began in the Late *\*Devonian* and continued throughout the *\*Carboniferous*, affecting a broad belt along an approximately WSW to ENE line from what are now south-western England and north-western Europe to southern Europe and the Iberian Peninsula, and eastern N. America and the Andes. Probably it was caused by northward thrusting and *\*plutonism* along a discontinuous front.

**hercynite** See SPINEL.

**Heretaungan** A \*stage (50–47 Ma ago) in the \*Eocene of New Zealand, underlain by the \*Mangaorapan, overlain by the \*Porangan, and roughly contemporaneous with the upper \*Ypresian and lower \*Lutetian stages.

**hermatypic** Applied to \*corals that contain zooxanthellae and are reef forming. Modern \*scleractinian hermatypic corals are characterized by the presence of vast numbers of symbiotic zooxanthellae (unicellular \*dinoflagellates) in their endodermal tissue. They live in waters of normal marine salinity, at depths of up to 90 m, in temperatures above 18 °C, and grow vigorously in strong sunlight.

**herringbone cross-bedding** A form of cross-bedding (see CROSS-STRATIFICATION) in which the \*foresets in successive sets are directed in opposite directions, so producing a structure which somewhat resembles the bones of a fish. The bipolar orientation of foresets seen in herringbone cross-bedding is commonly generated by the reversing currents developed in many tidal environments.

**Hervyan** The uppermost \*stage in the \*Devonian of Australia, underlain by the \*Condobolinian stage, overlain by the \*Carboniferous, and roughly contemporaneous with the upper \*Frasnian and \*Famennian of Europe.

**Hesperian** A division of \*areological time, lasting from 3.50 Gy to 1.80 Gy ago in the Hartmann–Tanaka Model and 3.80 Gy to 3.55 Gy ago in the Neukum–Wise Model, and divided into two epochs: Lower Hesperian (3.50–3.10 or 3.80–3.70 Gy ago); and Upper Hesperian (3.10–1.80 or 3.70–3.55 Gy ago).

**Hess, Harry Hammond** (1906–69) An American geophysicist from Princeton University, Hess made important contributions to the theory of \*plate tectonics. He devised the concept of \*sea-floor spreading (see also dietz, robert sinclair), and discovered and named \*guyots. His *Essay In Geopoetry* (1960, 1962) was an attempt to link the features of the sea floor in a common hypothesis, in which he proposed that the continents move passively on rafts or rigid plates.

**hessenbergite** See BERTRANDITE.

**Heterian** A New Zealand *\*stage* (156–152 Ma ago) of the Late *\*Jurassic* *\*epoch*. followed by the *\*Ohauan* and overlapping with the *\*Oxfordian* and Kimmeridgian stages.

**hetero-** From the Greek *heteros* meaning ‘other’, a prefix meaning ‘different from’.

**heterocercal tail** In fish, a tail in which the tip of the vertebral column turns upward, extending into the dorsal lobe of the tail fin; the dorsal lobe is often larger than the ventral lobe. The heterocercal tail is present in many fossil fish, in the sharks (*\*Chondrichthyes*), and in the more primitive bony fish, e.g. the families Acipenseridae and Polyodontidae. In the later, ray-finned fish (*\*Actinopterygii*) the homocercal tail developed, in which the vertebral column stops short of the tail fin, which is supported only by bony rays, giving rise to an apparently symmetrical type. *Compare* DIPHYCERCAL TAIL.

**heterochrony** Dissociation, during development, of factors of shape, size, and maturity, so that organisms mature in these respects at earlier or later growth stages. This leads to either *\*paedomorphosis* or *\*recapitulation*.

**heterocoelus** Applied to the condition in which the articulate surface of the *\*vertebra* centrum is saddle shaped, as is the case in birds. *Compare* AMPHICOELUS.

**Heterocorallia** (subclass *\*Zoantharia*) Small order of corals, known only from *\*Carboniferous* rocks in Europe and Asia, and represented by two genera, Hexophyllia and Heterophyllia. It is included in the *\*Zoantharia* because its members possess *\*septa*, and because of the microstructure of the calcareous skeleton. The septa are arranged differently from those in other orders. The four original septa are conjoined axially, and new septa are formed attached to these at their axial ends.

**heterodont** 1. (**invertebrate**) Applied to a *\*hinge* dentition occurring in the *\*Bivalvia*, where teeth of differing sizes occur in the hinge plate. The teeth are differentiated into *\*cardinal* teeth, which occur beneath the *\*umbo*, and lateral teeth, which occur posteriorly and anteriorly to the umbo. 2. (**vertebrate**) Possessing teeth that are differentiated into several forms, e.g. incisors, canines, premolars, and molars (as in mammals). This contrasts with the homodont condition, in which the teeth are all of the same form.

**Heterodonta** (phylum *\*Mollusca*, class Pelecypoda) Subclass of bivalves (*\*Bivalvia*), which have *\*heterodont dentition* consisting of a *\*hinge* plate with distinct *\*cardinal* teeth below the *\*umbo*, and lateral teeth posterior to the cardinals. Some forms may be *\*desmodont*. The ligament is *\*opisthodetic*. Shell shape varies according to the mode of life of the organism, most having a crossed-lamellar, *\*aragonitic* shell structure. The *\*pallial* line is entire, or with a sinus. Heterodonta are mainly *\*infaunal* siphon-feeders, the *\*siphons* being well developed. They first appeared in the Middle *\*Ordovician*.

**Heterodontosauridae** Ornithopod *\*ornithischian dinosaurs*, known only from the Upper *\*Triassic* of S. Africa. These 'different-toothed lizards' had tusk-like canines.

**heterogeneity** See INHOMOGENEITY.

**heterogeneous accretion** A model for the accretion of the planetary bodies from the primitive solar nebula (PSN), in which the rate of accretion of solid particles into the planets is slow relative to the rate at which the PSN cools. The consequence is that the surface layer of each body at any one time is in equilibrium with the pressure and temperature conditions prevailing in the nebula, and thus each planet accretes successive 'onion-skin' layers of material with different compositions. According to this model, the layered structure of planets may be partly of primary origin. Compare HOMOGENEOUS ACCRETION.

**heterogeneous simple shear** A *\*fold* mechanism in which folding is due to changes in the amount and orientation of a *\*simple* shear displacement. Layers arranged obliquely to planes of simple shear are rotated passively, producing similar folds of ideal form.

**heterolithic bedding** A closely interbedded deposit of *\*sand* and mud, generated in environments where current flow varies considerably. The three main types of heterolithic bedding are called *\*flaser*, *\*wavy*, and *\*lenticular*. Flaser bedding is characterized by cross-laminated sands with thin mud drapes over *\*foresets*. Wavy bedding consists of rippled sands with continuous mud drapes over the ripples. Lenticular bedding consists of isolated lenses and *\*ripples* of sand set in a mud *\*matrix*. Heterolithic sediments can be deposited in storm-wave influenced shallow marine environments, river *\*floodplains*, *\*tidal flats*, or *\*delta-front* settings where

fluctuating currents or sediment supply permit the deposition of both sand and mud.

**heterolithic unconformity** See NON-CONFORMITY.

**heteropygous** See PYGIDIUM.

**heterosphere** The layers of the atmosphere, above 80 km altitude, through which the chemical composition of the air changes markedly with height, principally as a result of oxygen dissociation. Compare HOMOSPHERE.

**heterospory** See SPORE.

**Heterostraci (Pteraspida; heterostracans)** (superclass \*Agnatha) Oldest known vertebrate order of jawless, heavily armoured fish-like forms, ranging from the Upper \*Cambrian to the \*Devonian. The \*dermal plates of the body lacked true \*bone cells and no internal skeleton has been preserved, so this presumably was \*cartilaginous. The anterior part of the body was covered by large dorsal and ventral plates, with smaller plates to the side. There was a single lateral gill opening, eyes at the sides of the head, and the impression of paired nasal sacs on the inner surface of the rostral shield (which protruded in front of the mouth). The rest of the body was covered by scales; the tail was hypocercal (tilted downwards). Heterostracans were usually only a few centimetres long, although some species reached 1.5 m. The body in typical forms, such as the Devonian Pteraspis, was rounded, but the group included dorso-ventrally flattened, bottom-dwelling forms such as Drepanaspis.

**heterotopy** An evolutionary change in the site at which a particular development occurs. The term was coined by E. H. Haeckel in 1866 to complement \*heterochrony.

**Hettangian** A \*stage in the European Early \*Jurassic (199.6–196.5 Ma ago, Int. Commission on Stratigraphy, 2004), preceded by the \*Rhaetian, followed by the \*Sinemurian, and roughly contemporaneous with the lower Aratauran (New Zealand). See also LIAS.


**Hexacorallia** Alternative name for the order \*Scleractinia (stony corals).

**Hexactinellida (Hyalospongea; glass sponges)** (phylum \*Porifera) A group of marine sponges that first appeared in the \*Cambrian. They are of

normal sponge shape and have skeletal *\*spicules* consisting of opaline *\*silica*. Large spicules (megascleres) and small spicules (microscleres) are connected in various patterns, and this forms the basis for the classification of the group. They were common in the *\*Palaeozoic* but many forms disappeared in the *\*Permian*. They were again important in the *\*Jurassic* and *\*Cretaceous*, but are now confined to depths of 200–2000 m.

**hexad** See CRYSTAL SYMMETRY.

**hexagonal** A *\*crystal* system characterized by four *\*crystallographic* axes of which three are horizontal and equal in length with an angle of  $120^\circ$  between them, and a fourth axis which is vertical. The vertical axis is one of six-fold symmetry. The hexagonal system differs from the *\*trigonal* system, which has a vertical axis of three-fold symmetry.



**Hexagonal**

**Hexapoda** (phylum *\*Arthropoda*) Subphylum of arthropods that includes the orders Collembola (springtails), Protura, and Diplura of wingless forms found in soil and moist litter, and the *\*Insecta* (insects). More than 950 000 species of hexapods have been described.

**HFS** See HIGH-FIELD-STRENGTH ELEMENTS.

**HFU** See HEAT-FLOW UNIT.

**hiatus** **1.** A gap in a sedimentary succession. **2.** The geologic-time interval represented by such a gap.

**hidden layer** A layer whose presence remains undetected by *\*seismic refraction* methods because (a) it is too thin and has insufficient velocity contrast to provide a discernible arrival; or (b) it is of lower velocity than the overlying medium. The term is analogous to the *\*suppressed* layer in the interpretation of *\*electrical* resistivity soundings.

**hierarchical method** **1.** The classification of organisms or units into a graded succession or hierarchy. **2.** In stratigraphy, the practice of defining *\*stratigraphic units* in terms of their *\*rank* within a hierarchy and the units of lower rank which they comprise. For example, the geologic *\*systems* are composed of *\*series*, which are themselves composed of *\*stages*. Compare TYPOLOGICAL METHOD.

**high-alumina basalt** An *\*aphyric*, *\*extrusive* *\*igneous* rock of basaltic composition containing a high proportion of *\*modal* *\*plagioclase*, and an alumina content of more than 17 wt. %. Tholeiitic and calc-alkali *\*basalts* (see THOLEIITE; CALC-ALKALINE) can have alumina contents of more than 17 wt. %, so the term does not refer to a specific basalt type. Rather, the variety of basalt should be specified, e.g. 'high-alumina basalt of calc-alkali type'.

**high-aspect-ratio ignimbrite (HARI)** An *\*ignimbrite* sheet displaying a value for the ratio of its average thickness ( $V$ ) to its horizontal extent ( $H$ ) of  $10^{-2}$  to  $10^{-3}$ , where  $H$  is taken as the diameter of a circle with a surface area equal to that of the flow. Compare LOW-ASPECT-RATIO IGNIMBRITE.

**high-field-strength elements (HFS)** Elements of high *\*valency* (greater than 2), e.g. Sn, W, and U, which are not readily incorporated into the *\*lattices* of common rock-forming *\*silicate* minerals. During crystallization of *\*igneous* rocks they are generally incorporated into accessory phases (see ACCESSORY MINERAL), e.g. *\*zircon* and *\*monazite*, or else continually concentrated into residual *\*pegmatitic* or *\*hydrothermal* fluids, from which they may be precipitated in economic concentrations.

**high-level waste** See RADIOACTIVE WASTE.


**high-pass filter** In *\*remote sensing*, a type of *\*spatial-frequency* filter which selectively removes low *\*spatial-frequency* data from an image and increases contrast between high spatial-frequency data. The result is that the lines which make up the edges of objects are more clearly defined. High-pass filters are used to achieve *\*edge enhancement*.

**high-plateau crustal type** A type of *\*crust* that results from uplift, but without folding (see *FOLD*) or *\*thrust* faulting. The elevation is high and the average crustal thickness is probably about 35 km.

**high-potassium basalt** An *\*igneous* rock of basaltic composition characterized by the appearance of *\*sanidine* as well as *\*labradorite* in the *\*groundmass*, and by  $K_2O/Na_2O > 1$ . Such *\*basalts* are commonly found well inland on the continental side of a *\*plate* *\*subduction zone*. The  $K_2O$  content of basalts in these above-subduction-zone environments increases inland as the depth to the subducted oceanic slab increases.

**highstand** A time during which sea levels are at their highest. Compare *LOWSTAND*.

**highstand systems tract (HST)** In the *\*genetic stratigraphic sequence model* used in *\*sequence* stratigraphy, a bounding surface formed by maximum marine flooding, where the sea level was high, and stable or falling slowly. See *SYSTEMS TRACT*. Compare *LOWSTAND SYSTEMS TRACT*; *TRANSGRESSIVE SYSTEMS TRACT*.



Highstand systems tract

**high symmetry** See *CRYSTAL SYMMETRY*.

**Hiller borer** See PEAT-BORER.

**Hiller peat-borer** See PEAT-BORER.


**hill fog** General term denoting low cloud that covers high ground.

**Himalayan orogenic belt** An *\*orogenic* belt interpreted as being the result of the convergence and collision of India with Asia. Because of variable uplift, great relief, high rates of *\*erosion* and upthrusts of mid-crust, most depths within the *\*orogen* are exposed, down to the mid-crust, and the belt is thought to illustrate the later stages of the *\*Wilson* cycle. To the north of the Indus *\*suture* the *\*Tibetan Plateau* has *\*crust* up to 80km thick: a few authorities interpret this as being due to *\*A-subduction* of the crust of the *\*Indo-Australian Plate* after collision in the *\*Eocene*, but most attribute the thickness to internal thrusting since the early *\*Mesozoic*.

**Himalia (Jupiter VI)** One of the lesser satellites of *\*Jupiter*, with a diameter of 170 km ( $\pm 20$  km).


**Himawari-8 and -9** Two meteorological satellites in *\*geostationary orbit* operated by the Japan Meteorological Agency. Himawari-8 was launched on 7 October 2014 and Himawari-9 on 2 November 2016, both from the Tanegashima Space Center, Japan.

**hinge** **1.** The surface region of a *\*fold* about a *\*hinge line*, which occupies the area of maximum curvature. **2.** In *\*Brachiopoda* and lamellibranchs, the area of the *\*commissure* where the two valves of the shell are permanently in contact. In those articulate brachiopod (*\*Articulata*) shells where the whole width of the posterior part of the commissure remains in contact during opening, the shell is said to be 'strophic'. In those brachiopods where the hinge axis passes through the teeth and sockets, and these represent the fulcra, the shell is said to be 'non-strophic'.



## Hinge

**hinge line** A straight or curved line which joins the points of maximum curvature along the \*hinge of a \*fold.



## Hinge line

**hinterland** See FORELAND.

**Hippocamp** (**Neptune XIV**) The smallest satellite of **\*Neptune**, discovered on 1 July 2013. It orbits Neptune in 0.9362 days, has a mean radius of 8–10 km, and has a dark surface.

**Hirnantian** A **\*stage** (445.6–443.7 Ma ago) of the Late **\*Ordovician** **\*epoch** in the Upper **\*Ashgill**, underlain by the **\*Rawtheyan** and overlain by the **\*Rhuddanian** (**\*Silurian**).

**Hiscock Radiation Belt Explorer** See HRBE CUBESAT MISSION.


**histic epipedon** Surface **\*soil horizon**, not less than one metre in depth, high in organic carbon, and saturated with water for some part of the year. See also HUMUS (2).

**histogram** A frequency graph in which the data range is divided into **\*bins**.

**histosols** An order of soils in the **\*soil taxonomy** that are composed of organic materials. Histosols must have a thickness of more than 40 cm when overlying unconsolidated mineral soil, but may be of any thickness when overlying rock.

**HJ-1** See HUAN JING-1.

**Hjulström effect** The contrast between the flow velocity at which a fine-grained cohesive **\*sediment** may be deposited and that at which it will be eroded. F. Hjulström published an important diagram which plots the relationship between grain size and velocity for **\*erosion**, transportation, deposition, and settling velocity. One of the many important consequences of these relationships is that although fine-grained cohesive sediments (fine **\*silt** and **\*clay**) will be deposited only if flow velocities are very low, a very high velocity is required to erode the same sediment, once deposited. This is because of the cohesive nature of the sediment, which makes silt and clay more difficult to erode than pebbly sediment.



### Hjulström effect


**hlaup** A burst of water through an *\*ice cap*, triggered by volcanism.

**HMS** See DENSE-MEDIUM SEPARATION.

**hoar frost** Deposit of patterned ('feathers', 'needles', 'spines', etc.) ice crystals on surfaces chilled by radiation cooling. The feature is seen particularly well on vegetation. The ice is derived from the *\*sublimation* of water vapour on surfaces, as well as from frozen dew.

**Hoboken** See HENGELO.

**hogback (hog's back)** Narrow, symmetric ridge, underlain and controlled by a resistant bed dipping at some 40° or more. It is a limiting case of a *\*cuesta*.



**Hogback**

**hog's back** See HOGBACK.

**Hohmann transfer orbit** An elliptical orbit that carries a spacecraft from one circular orbit to a different circular orbit when both orbits lie in the same plane. In transferring from a low orbit to a higher one, the spacecraft engines fire to accelerate the craft into an elliptical orbit that intersects the higher orbit. The spacecraft travels through half the ellipse. As it arrives at the higher orbit, the engines fire again to accelerate it into the higher orbit. The same manoeuvre can also be used in reverse to bring a spacecraft into a lower orbit. In practice, the two orbits may not be perfectly circular and therefore are not coplanar, in which case the transfer orbit is slightly less than  $180^\circ$ , when it is known as a Type I Hohmann transfer, or slightly more than  $180^\circ$ , when it is known as a Type II Hohmann transfer. The manoeuvre was devised by the German engineer Walter Hohmann (1880–1945).

**Holkerian** A substage (339–337.5 Ma ago) of the *\*Visean* *\*stage*, underlain by the *\*Arundian* and overlain by the *\*Asbian*.

**Holmes, Arthur** (1890–1965) A British geologist of the Universities of Durham and Edinburgh, Holmes did important work on *\*radiometric* dating and developed the concept of convection currents in the *\*mantle*, caused by heat from radioactive decay. This latter work persuaded him to become an early supporter of *\*continental drift* theory. His *Principles of Physical Geology* (1944, 1965, 1993) has been an important textbook for more than 50 years.

**holo-** From the Greek *holos* meaning ‘complete’, a prefix meaning ‘whole’ or ‘complete’.

**Holocene (Hallian, Hawera)** \*Epoch that covers the last 11 000 years. It is often referred to as Recent or Post-Glacial. The Holocene is divided into three \*ages/\*stages: \*Greenlandian, \*Northgrippian, and \*Meghalayan. See also FLANDRIAN.

**holochroal** See TRILOBITE EYE.

**holocrystalline** A \*textural term referring to \*igneous rocks or parts of igneous rocks which are composed entirely of crystals.

**holohyaline** A rarely used \*textural term referring to \*igneous rocks or parts of igneous rocks which are composed entirely of \*glass.

**holokarst** A region where almost all the land-forms result from karstic activity (see KARST). There is little or no surface runoff and there are few streams. The entire area is underlain by thick \*carbonate rocks.

**hololeucocratic** A rarely used term applied to an \*igneous rock with a \*colour index less than 5.

**holometabolous** Applied to a type of development in which distinct larval and adult forms occur.

**holomictic** Applied to lakes in which all of the water circulates, from the surface to the bed, when the lake cools in winter. Compare \*meromictic.

**holophyletic** Applied to a \*taxon that includes all descendants of the common ancestor. The term is a special case of \*monophyletic.

**Holostei** Group of marine and freshwater \*bony fish including many fossil species, e.g. Lepidotes (\*Triassic–\*Cretaceous), and Dapedius (\*Jurassic). Holosteans arose at the end of the \*Permian from \*palaeoniscid ancestors and are particularly abundant in marine Jurassic deposits. The main characteristics shown by the group relate to improvements in swimming and feeding: the use of the air sac to control buoyancy; a reduction in the bony fin rays, now unjointed; the development of a shorter and more mobile jaw; the gradual reduction in scale thickness; and the development of an almost symmetrical tail. Holosteans are represented today only by the garpike (*Lepisosteus*), and the bowfin (*Amia*), both freshwater.

**holostomatous** See APERTURE.

**holostratotype** The original **\*stratotype** chosen and described to act as the standard reference section, or reference point within a section, when a stratigraphic unit or boundary is first established. Compare LECTOSTRATOTYPE; NEOSTRATOTYPE. See also PARASTRATOTYPE; HYPOSTRATOTYPE.

**holosymmetric** Within each **\*crystal system**, applied to the class with the highest order of symmetry, where the **\*crystal** has the same symmetry as the lattice.

**holothuroid (sea cucumber)** See HOLOTHUROIDEA.

**Holothuroidea (holothuroids; sea cucumbers)** (subphylum Echinozoa) Class of worm-like echinoderms which may be free-living or attached. The mouth, surrounded by variably branched tentacles, is at one end of the elongate body, the anus at the other. The calcitic skeleton is not rigid, being reduced to small sclerites (or **\*spicules**) of very variable shape: hooks, anchors, rings, and plates. Fossil spicules of sea cucumbers are first encountered in rocks of **\*Ordovician** age.

**holotype (type specimen)** Individual plant or animal chosen by taxonomists to serve as the basis for naming and describing a new species or variety. Compare LECTOTYPE; NEOTYPE; PARATYPE; SYNTYPE.

**Holsteinian** A north European **\*interglacial** period, dating from about 0.3 Ma ago to 0.25 Ma ago, that is probably the equivalent of the **\*Mindel/Riss Interglacial** of the Alpine areas and may also be equivalent to the **\*Hoxnian** of East Anglia.

**Homalozoa ('carpoids')** (phylum **\*Echinodermata**) Extinct subphylum whose members have no trace of **\*radial symmetry**. The **\*theca** is depressed and asymmetrical. Homalozoa are known from Middle **\*Cambrian** to Middle **\*Devonian** rocks. There are three classes: Homostelea, Homoiostelea, and Stylophora. The affinities of the group have generated some controversy. The similarities between the larvae of some echinoderms and some chordates (**\*Chordata**) has led to the suggestion that the phyla are in some way related. Most recently, the Stylophora have been reassigned by

R. P. S. Jeffries on morphological grounds to a new subphylum of primitive chordates, the Calcichordata. The matter remains unresolved.

**homeomorph** An organism which, as a result of *\*convergent* evolution, comes to resemble another to which it may not be closely related.

**homeostasis** See HOMOEOSTASIS.

**homeotherm** See HOMOIOOTHERM.

**Homerian (Lockportian)** A *\*stage* (426.2–422.9 Ma ago) of the *\*Niagaran* (Early *\*Silurian*) of N. America, underlain by the *\*Sheinwoodian* and overlain by the *\*Gorstian*.

**Hominidae** (order Primates, suborder *\*Anthropoidea*, superfamily *\*Hominoidea*) The mammalian family that originally included humans and immediately ancestral forms now extinct. Hominids were distinguished from the apes (Pongidae) (*a*) by the possession of a much larger brain, in which the frontal and occipital lobes are especially well developed, allowing more complex behaviour including communication by speech; (*b*) by a fully erect posture facilitated by the positioning of the *\*foramen magnum* beneath the skull so that the head is held upright; (*c*) by a bipedal gait; and (*d*) by the slow rate of post-natal growth and development, which favours complex social organization and the emergence of distinct cultures. However, the Pongidae is no longer considered to be a *\*monophyletic* group and most modern authorities include the chimpanzee (*Pan troglodytes*), bonobo (*P. paniscus*), gorilla (*Gorilla gorilla*), and probably the orang-utans (*Pongo pygmaeus* and *P. abellii*) in the Hominidae, on the ground that they are close to humans phylogenetically (see PHYLOGENY). The Hominidae also includes the genera *Paranthropus* and *Australopithecus* (but see *\*australopithecines*) as well as *Homo*. *\*Ramapithecus*, once considered a possible human ancestor, is now known to be more closely related to the orang-utans.

**Hominoidea** (order Primates, suborder *\*Anthropoidea*) Superfamily comprising the Hylobatidae (gibbons) and *\*Hominidae* (humans, apes, and their immediate ancestors). The hominids are believed to be descended from a common stock of ‘great apes’ which diverged to form distinct Asian and African lines, the African line dividing again 4–6 million years ago into the African apes and the ancestors of modern humans. The hominoids lack

tails and cheek pouches; have opposable thumbs (reduced in some species); and differ from the Cercopithecidae (Old World monkeys) in having less-specialized dentition, larger heads, longer limbs, and wider chests which some authorities believe they inherited from ancestral brachiating forms. Today only the Hylobatidae (gibbons) are specialized brachiators (i.e. swing from branches hand over hand).

**homo-** From the Greek *homos* meaning 'the same', a prefix meaning 'the same' or 'alike'.

**homocercal tail** See HETEROCERCAL TAIL.

**homodont** See HETERODONT.

**homoeostasis (homeostasis)** The condition of a system that possesses a capacity for self-regulation whereby it resists change.

**homogeneous accretion** A model for the accretion of planetary bodies from the primitive solar nebula (PSN) in which the rate of accretion is fast relative to the rate at which the PSN cools. The consequence is that each body forms very quickly and consists entirely of material that was in equilibrium with the physical conditions of the PSN over only a very short period of time. According to this model, the layered structures of the planets are entirely of secondary origin. Compare HETEROGENEOUS ACCRETION.

**homogeneous non-rotational strain** See PURE SHEAR.

**homogeneous nucleation** Spontaneous condensation or freezing of water in the atmosphere in the absence of substances to act as nuclei. This is most likely in supercooled air below  $-40^{\circ}\text{C}$ .

**homogeneous rotational strain** See SIMPLE SHEAR.

**homogeneous strain** The \*strain, distributed evenly throughout a body, which is produced when straight lines and parallel lines in an undeformed body remain straight and parallel as the body is deformed. There are three types of homogeneous strain, varying smoothly from one to the next: axially symmetrical extension; axially symmetrical shortening; and plane strain. See PURE SHEAR; SIMPLE SHEAR.

**homogenitus** See CLOUD CLASSIFICATION.

**homogenization temperature** 1. Temperature at which an *\*exsolution* pair of minerals, e.g. *\*chalcopyrite*–*\*sphalerite*, homogenize to a single crystalline phase. 2. Temperature at which the liquid and vapour phase in a *\*fluid* inclusion homogenize to monophasic liquid or vapour. This gives an indication of the minimum trapping temperature of the fluid inclusion. See also GEOTHERMOMETER.

**homoiotherm (homeotherm)** An organism whose body temperature varies only within narrow limits. It may be regulated by internal mechanisms (i.e. in an endotherm) or by behavioural means (i.e. in an ectotherm), or by some combination of both (e.g. in humans, who light fires and wear thick clothes to keep warm in cold weather and wear light clothing to keep cool in warm weather, but who are also endothermic).

**homologous** Applied to similar structures in different animals when both are thought to have the same evolutionary origin, although their functions may differ widely, e.g. forelimbs, wings, and flippers in *\*vertebrates*.

**homomutatus** See CLOUD CLASSIFICATION.

**homoplasy** In the course of *\*evolution*, the appearance of similar structures in different lineages (i.e. not by inheritance from a common ancestor). The term includes *\*convergent evolution* and *\*parallel evolution*. See also REVERSAL.

**homopycnal flow** At a river mouth, a flow of river water that is of the same density as the water in the basin receiving it. This results in intense local mixing of waters, with considerable sedimentation, and is typical of *\*Gilbert-type* deltas. Compare HYPERPYCNAL FLOW; HYPOPYCNAL FLOW.

**homosphere** Atmospheric layer from the Earth's surface to approximately 80 km altitude, where the relative proportions of the various gaseous constituents, excluding water vapour, remain almost constant. See also HETEROSPHERE.

**homospory** See SPORE.

**homotaxis** Literally, 'the same arrangement' (from the Greek *homos* and *taxis*). The term was proposed by T. H. Huxley (1825–95), in an address to

the **\*Geological Society of London**, to describe **\*strata** from different areas that contain similar lithologic or **\*fossil** successions but are not necessarily of the same age.

**Honda–Mrkos–Pajdusakova** A **\*comet** with an orbital period of 5.29 years; **\*perihelion** date 17 January 1996; perihelion distance 0.581 AU.

**hoodoo (earth pillar)** An erosional feature that is seen in **\*badlands**. It may result from severe **\*gully** erosion, but a hoodoo often develops where a resistant **\*cap rock** protects softer sedimentary strata. Rain erodes the softer rocks, leaving a pinnacle, which is the hoodoo. Hoodoos also develop where a boulder protects **\*till**. Hoodoos are often conical in shape beneath the cap rock and they sometimes occur in groups. Their curious shapes, which from a distance may bear a faint resemblance to a robed human figure, account for their name.

**Hooke, Robert** (1635–1703) An experimental philosopher, Hooke investigated a very wide range of topics and published extensively. He is responsible for **\*Hooke's law**, which laid the foundations for the scientific study of elasticity. In the field of Earth sciences, he studied gravity and planetary movements, earthquakes, geomagnetism, and the nature of fossils and crystallography.

**Hooke's law** The law, named after **Robert \*Hooke**, which describes the behaviour of perfectly elastic materials in terms of a straight-line relationship between **\*stress** and **\*strain** in such materials. Strain is directly proportional to the applied stress provided the medium remains elastic (i.e. the **\*elastic limit** is not exceeded); stress is equal to the strain multiplied by a constant of elasticity (**\*Young's modulus**). The law also states that stress divided by strain is a constant. *See also* ELASTIC CONSTANTS.

**hopeful monster** An individual possessing a macromutation which is of no benefit to the individual carrying it (i.e. the individual is a monster), but that may prove beneficial to one of its descendants if the mutation undergoes some further but minor change (i.e. the monster is hopeful).

**'hopper' crystals** Cubic crystals of **\*halite** or **\*pseudomorphed** (replaced) halite, characterized by sunken depressions on the crystal faces.

**horizon 1.** An **\*informal** term used in **\*stratigraphy** to denote a plane within a body of **\*strata**. This may be at a boundary of lithological change, or

commonly the term may refer to a thin, distinctive **\*bed** within a lithological unit. See also **BIOHORIZON**. **2.** An interface separating two media with different geophysical properties. **3.** In soil, a horizontal layer that can be distinguished from the layers below and (except for the surface layer) above it. Identified by a coding system using a capital letter, sometimes followed by a subscript, such layers are used to diagnose soil types. See **SOIL HORIZON**.

**horizontal drilling** The drilling of an oil or natural gas well at an angle to the vertical, so the well runs parallel to the formation containing the oil or gas. Production from the resulting well, known as a horizontal hole, is often three to five times greater than that from a vertically drilled well.

**horizontal hole** See **HORIZONTAL DRILLING**.

**horizontal stack** See **COMMON-DEPTH-POINT STACK**.

**horn (glacial horn, pyramidal peak)** A steep, horn-shaped mountain summit that is the result of glacial action when two adjacent **\*cirques** erode into the mountainside and form steep **\*arêtes**. The Matterhorn, in the Swiss Alps, is a famous example.



<https://sites.google.com/site/mtwashingtonmountains/formation-of-presidentials/glaciation>

- Describes glacial horns.

**hornblende** An important rock-forming **\*silicate** mineral and member of the **\*amphiboles**, particularly the calcium-rich **\*monoclinic** amphiboles; it is the general name given to a group of amphiboles (the hornblende series) with similar properties, which includes hastingsite  $\text{Ca}_2(\text{Mg}_4\text{Al})[\text{Si}_7\text{AlO}_{22}](\text{OH},\text{F})_2$ , tschermakite  $\text{Ca}_2(\text{Mg}_3\text{Al}_2)[\text{Si}_6\text{Al}_2\text{O}_{22}](\text{OH},\text{F})_2$ , edenite  $\text{NaCa}_2\text{Mg}_5[\text{Si}_7\text{AlO}_{22}](\text{OH},\text{F})_2$ , and pargasite  $\text{NaCa}_2(\text{Mg}_4\text{Al})[\text{Si}_6\text{Al}_2\text{O}_{22}](\text{OH},\text{F})_2$ ; sp. gr. 3.0–3.5; **\*hardness** 5.0–6.0; black or greenish-black; white **\*streak** with a greenish tint; **\*vitreous** **\*lustre**; **\*crystals** normally **\*prismatic**, **\*columnar**, and occasionally isometric; **\*cleavages** in two directions set at  $124^\circ$ , prismatic {110}; occurs as a common constituent of medium **\*basic**, **\*igneous** rocks (e.g. **\*syenites**, **\*diorites**, and

\*granodiorites), also in contact \*metasomatic zones, and extensively in \*metamorphic rocks (e.g. \*schists, \*gneisses, and \*amphibolites).

**hornblende–hornfels facies** A set of metamorphic mineral assemblages produced by \*contact metamorphism of a wide range of starting rock types under the same metamorphic conditions and typically characterized by the development of the mineral assemblage \*hornblende–\*plagioclase in rocks of \*basic \*igneous composition. Other rocks of contrasting composition, e.g. \*shales and \*limestones, would each develop their own specific mineral assemblage, even though they are all being metamorphosed under the same conditions. The variation of mineral assemblage with starting rock composition reflects a particular range of pressure, temperature, and  $P(H_2O)$  conditions. Experimental studies of mineral  $P$ – $T$  stability fields indicate that the facies represents the low-pressure, moderate-temperature conditions met close to \*igneous \*intrusions.

**hornblendite** An \*ultrabasic \*igneous rock composed of more than 90% \*hornblende. Where other \*ferromagnesian minerals constitute 10–50% of the rock, the rock can be named according to the other types of ferromagnesian minerals, for example, pyroxene hornblendite, olivine–pyroxene hornblendite.

**hornfels** (*pl.* hornfelses) A massive, fine-grained, \*granoblastic contact metamorphic rock (see CONTACT METAMORPHISM), commonly displaying \*conchoidal fracture and splintery debris. Heat from an \*igneous \*intrusion initiates \*recrystallization of surrounding rocks, thereby producing a contact metamorphic hornfels which is usually finer-grained than the \*country rocks. Hornfelses formed by recrystallization of \*shales, and occurring away from the contact with the igneous body, quite often develop prominent \*graphite-rich spots formed by the aggregation of organic material in the original shales.

**hornitos** Small cones or chimneys of \*lava \*spatter, often several metres high, found on the surface of \*pahoehoe lava flows. Lava spatter is thrown out from the flow interior by the explosive release of primary magmatic gas or by the explosive conversion to steam of trapped pockets of \*groundwater beneath the flow.

**horotely** (*adj.* horotelic) Normal or average rate of \*evolution per million years, of genera within a given \*taxonomic group. Thus slowly or rapidly

evolving lines may be horotelic during certain episodes in their history.

**horse** A lenticular or sigmoidal mass of rock which is completely bounded by two or more *\*thrust* faults which rejoin along the *\*strike* and up-*\*dip*. The term may also be used for the analogous structure in strike-slip terrains (see *STRIKE-SLIP FAULT*).

**horse latitudes** Subtropical latitudes coinciding with a major anticyclonic belt; they are characterized by generally settled weather and light or moderate winds. When sailing ships carrying cargoes of horses were becalmed in these latitudes, horses sometimes died from thirst and were thrown overboard.

**horsetails** See *SPHENOPSISIDA*.

**horst** Upthrown block lying between two steep-angled *\*fault* blocks. Compare *GRABEN*. See also *RIFT*.

**hortic horizon** In the *\*World Reference Base for Soil Resources* classification, a *\*soil horizon* that has been strongly influenced by cultivation.

**Horton analysis** See *DRAINAGE MORPHOMETRY*.

**Hortonian flow** See *SURFACE RUNOFF*.

**hot dry rock (HDR)** Rocks, usually *\*granite*, which have abnormally high heat production as a result of the decay of *\*radiogenic* elements rather than merely residual heat. Potentially these are a source of geothermal energy. One method of exploiting the heat generated is to fracture the rocks at depth using small, down-hole, explosive charges, and then initiate a water circulation system from the surface. When cold water is pumped down it returns considerably warmer, and this energy can be extracted by heat exchangers. See also *GEOTHERMAL FIELD*; *GEOTHERMAL GRADIENT*.

**hot spot** An area of high volcanic activity. Some hot spots, e.g. Iceland, are located on *\*constructive margins*. Others occur within lithospheric *\*plates*, often lying at the end of a chain of progressively older volcanoes, e.g. the *\*Hawaiian-Emperor Chain*. The hot spot is thought to be stationary, or nearly so, and to produce volcanoes intermittently as the plate moves over it. It has been suggested that *\*mantle plumes* lie beneath hot spots.

**hot spring** A continuous flow of hot water through a small opening on to the Earth's surface. The water is usually \*groundwater heated at depth by hot rocks and recycled to the surface by convection. Hot spring waters are rich in dissolved minerals which are often precipitated around the spring mouth.

**hot working (annealing, polygonization)** A \*strain-recovery process in which new, unstrained, polygonal grains develop from and replace highly strained grains at high temperatures.

**Howard, Luke** (1772–1864) An English manufacturing chemist and meteorologist who was the first person to devise a successful classification system for clouds, which he published in 1803 as a paper, 'On the Modifications of Clouds'. Howard defined four main and several secondary cloud types, calling the main types 'stratus', 'cumulus', 'cirrus', and 'nimbus'. He became very well known and his classification system forms the basis of the one used today. Howard lived in London.

**Hoxnian** 1. An \*interglacial period. 2. A series of temperate-climate deposits, named after Hoxne, Suffolk, England, with a characteristic vegetational sequence that occurs in \*tills of the earliest glacial \*stages, sometimes filling deep channels. They may be equivalent to the \*Holsteinian deposits of continental Europe. Sometimes during the Hoxnian Interglacial the sea rose to well above its present level. There is a correlation between this stage and the Boyn Hill terraces of the Thames Valley, and also with \*raised beaches (30m ordnance datum) found on the Sussex coast.

**HRBE CubeSat Mission (Hiscock Radiation Belt Explorer; E1P-2; Explorer-1 PRIME-2)** A 1-unit \*CubeSat mission by Montana State University, sponsored by \*NASA, to detect the intensity and variability of energetic electrons in the \*Van Allen belts. The CubeSat, with 10-cm sides weighing less than 1 kg, was launched on 28 October 2011, from California. It is named for William A. Hiscock, founder and former director of the Montana Space Grant Consortium.

**HREE** Abbreviation for 'heavy \*rare-earth element'. See MID-OCEAN-RIDGE BASALT.

**HSR** See HIGHSTAND SYSTEMS TRACT.

**Huan Jing-1 (HJ-1)** A constellation of Earth-observing radar *\*minisatellites* to monitor disasters and for environmental protection that form a national program by the National Committee for Disaster Reduction and State Environmental Protection Administration of China (*Huan Jing* means ‘environment’). The mission comprises three minisatellites. Two were launched on 5 September 2008, into a Sun-synchronous *\*polar orbit* at an altitude of 649 km, and the third on 19 November 2012, at an altitude of 499 km, all from the Taiyuan Satellite Launch Center.

**Hubble constant** See HUBBLE PARAMETER.

**Hubble parameter ( $H_0$ )** A measure of the rate at which the recessional speed of galaxies varies with distance, calculated from *\*Hubble’s law*;  $H_0 = 50\text{--}100$  km/s/Mpc (megaparsec). This was formerly known as the Hubble constant.

**Hubble’s law** As formulated in 1936 by the American astronomer Edwin Powell Hubble (1889–1953) and his colleague Milton Lasell Humason (1891–1972), the distance to a galaxy in light-years is equal to one-hundredth of its red-shift velocity in miles per second.

**Hubble Space Telescope** The first major optical telescope to be placed in space, the Hubble Space Telescope was launched on 24 April 1990 and deployed on 25 April into a *\*low Earth orbit* at an altitude of 547 km. It is 13.2 m long, has a 4.2 m maximum diameter, and completes one Earth orbit in about 95 minutes.



[https://www.nasa.gov/mission\\_pages/hubble/story/index.html](https://www.nasa.gov/mission_pages/hubble/story/index.html)

- About the Hubble Space Telescope.

**Hudsonian orogeny** A *\*Proterozoic* phase of mountain building, which ended about 1750–1800 Ma ago, and that affected the shield area in what is now Canada. It was preceded by the *\*Kenoran* and succeeded by the *\*Grenvillian* orogenies.

**hue** A measure of the relative amounts of the *\*additive* primary colours which contribute to the colour of an object. See MUNSELL COLOUR.

**Hugoniot** The relation between pressure and density within the *\*Earth* as derived from *\*seismic velocities*. It is named for the French engineer Pierre Henri Hugoniot (1851–87).

**Humberian orogeny** *See* TACONIC OROGENY.

**Humboldt, Friedrich Heinrich Alexander von** (1769–1859) A Prussian naturalist and physicist, and great explorer, Humboldt made contributions to the study of volcanoes, vegetation and its relation to climate, ocean currents, and mountain ranges. He made a particular study of *\*geophysics*, aiming to collect and correlate all relevant data, and played an important role in the international geomagnetic survey of 1834, which determined declinations and inclinations worldwide.

**Humboldt current** *See* PERU CURRENT.

**humerus** In *\*tetrapods*, the upper bone of the forelimb.

**humic acid** Mixture of dark-brown organic substances, which can be extracted from soil with dilute alkali and precipitated by acidification to *\*pH* 1.0–2.0 (in contrast with fulvic acid, which remains soluble in acid solution).

**humic coal** *See* COAL.

**humidity** Expression of the moisture content of the atmosphere. Measures of humidity include statements of the total mass of water in 1 m<sup>3</sup> of air (absolute humidity), the mass of water vapour in a given mass of air (specific humidity), *\*relative humidity*, vapour pressure, and the *\*mixing ratio*.

**humid mid-latitude morphoclimatic zone** A *\*morphoclimatic* region in which the rate of *\*chemical weathering* is moderate, and high in low latitudes. *\*Mechanical* weathering is moderate, with *\*frost* action being important at high latitudes. *\*Fluvial activity* is moderate, and the rate of *\*mass-wasting* is moderate to high. Wind action is significant only near coasts.

**humid tropical morphoclimatic zone** A *\*morphoclimatic* region in which the rate of *\*chemical weathering* is high, but that of *\*mechanical* weathering fairly low. *\*Mass-wasting* occurs sporadically. Locally there are high rates of transport of *\*dissolved load* and *\*suspended load*.

**humification** The sequence of reactions by which decaying organic material is converted to *\*humus*.

**humilis** From the Latin *humilis* meaning ‘low’, a species of shallow *\*cumulus* cloud which typically has a flattened appearance. See also CLOUD CLASSIFICATION.

**humite** See CHONDRODITE.

**hummocky cross-bedding** A form of cross-bedding (see CROSS-STRATIFICATION) characterized by cross-laminations which have both concave and convex-upwards forms. The cross sets cut across one another with concave and convex-upwards surfaces, and the cross-beds have an external form of convex-upwards hummocks with wavelengths of about 1–5 m. The structure has been recognized only in ancient sediments and is thought to be the product of storm waves. See SWALEY CROSS-BEDDING.

**hummocky moraine** A strongly undulating surface of ground *\*moraine*, with a relative relief of up to 100 m, and showing steep slopes and deep, enclosed depressions. It results from the downwasting (i.e. thinning) of ice which is usually stagnant. Blocks of ice may squeeze debris released from the ice into *\*crevasses* between the blocks.

**humus** 1. Decomposed organic matter in soils that are *\*aerobic* for part of the year. It is dark brown and amorphous, having lost all trace of the structure and composition of the vegetable and animal matter from which it was derived. 2. Surface organic *\*soil horizon* that may be divided into types: *\*mor* (acid and layered) or *\*mull* (alkaline and decomposed). It is now known as a ‘*\*histic epipedon*’.

**Huronian** 1. A *\*system* of the *\*Proterozoic*, extending from about 2500 to 1400 Ma ago (Int. Commission on Stratigraphy, 2004) and containing three glacial cycles. 2. A Lower–Middle Proterozoic *\*stage*, extending from about 2600 to 1500 Ma ago, preceded by the *\*Neoproterozoic* era, and followed by the *\*Riphean* stage.

**hurricane** 1. Strictly, the name given to a *\*tropical cyclone* that develops over the N. Atlantic and Caribbean. Hurricanes move westward, then swing north, on tracks that often carry them across inhabited islands and coastal areas of Mexico and the USA. 2. A widely used alternative name for a

tropical cyclone, regardless of where it develops. **3.** A wind blowing at more than 120 km/h (75 mph), which is Force 12 on the [\\*Beaufort scale](#).

**Hutchinsonian** A New Zealand [\\*stage](#) (21–20 Ma ago) of the Early [\\*Miocene](#), preceded by the [\\*Otaian](#) and followed by the [\\*Awamoan](#).

**Hutton, James** (1726–97) Scottish natural philosopher, prominent advocate of Plutonist and uniformitarian theories, who published *The Theory of the Earth* (1795). His ideas were based on the study of crystalline rocks, but he paid little attention to stratigraphy and organic remains. He recognized that unconformities implied that earth movements must have occurred. His work was popularized by [\\*Playfair](#). See [PLUTONISM](#); [UNIFORMITARIANISM](#).

**Huygenian eyepiece** See [EYEPIECE](#).

**Huygens** A mission by [\\*NASA](#) and [\\*ESA](#), carried by [\\*Cassini](#) and launched in 1997, to [\\*Titan](#). Huygens reached Titan in 2004.



<http://www.esa.int/esaMI/Cassini-Huygens/>

- A joint ESA and NASA probe launched from the Cassini orbiter that landed on Titan, moon of Saturn.

**Huygens' principle** Each point of an advancing wavefront can be thought of as a new source of secondary wavelets, so that the envelope tangent to all these wavelets forms a new wavefront.

**HY-1B** See [HAIYANG-1B](#).

**Hyaenidae** See [CARNIVORA](#).

**Hyakutake** A [\\*comet](#) with an orbital period of more than 65 000 years; [\\*perihelion](#) date 1 May 1996; perihelion distance 0.230 AU.

**hyaline** Translucent or transparent; glass-like.

**hyaloclastite** An aggregate of fine, glassy debris formed by the sudden contact of hot, coherent [\\*magma](#) and either cold water or water-saturated sediment. Rapid heat loss from the magma to the cold water sets up tensile thermal stress in the magma carapace as it cools, chills, and contracts, causing the glassy, chilled zone to fragment and form a quench-fragmented debris. Thick hyaloclastite deposits form over [\\*basalt](#) flows when they

erupt beneath the sea or enter the sea. If the deposit remains in contact with water after its formation, the glassy debris can easily be hydrated to form **\*palagonite**.

**hyalopilitic** A **\*texture** found in **\*extrusive \*igneous** rocks and characterized by a felt of crystals set in a background of **\*glass**. The felts of crystals represent an arrested stage of early crystal growth from a liquid now represented by the background glass.

**Hyalospongea** See **HEXACTINELLIDA**.

**hydration** Chemical combination of water with another substance, e.g. the addition of water to a mineral (such as **\*anhydrite**) to produce a hydrous phase (in this case, **\*gypsum**). This usually involves expansion. Hydration may be important in the mechanical **\*weathering** of rocks to produce **\*clays** and economically important minerals such as **\*kaolin**, **\*talc**, and **\*goethite**.

**hydraulic boundary** Within a **\*groundwater** system, the interface between regions of different hydraulic characteristics such as **\*porosity**, storativity (see **STORAGE COEFFICIENT**), conductivity (see **PERMEABILITY**), or **\*transmissivity**. For example, when modelling groundwater systems or carrying out pumping tests, workers must make judgements regarding the homogeneity of those systems. Distinct and marked changes in the hydraulic characteristics (e.g. where an **\*aquifer** abuts an **\*aquiclude**) may require marking as hydraulic boundaries.

**hydraulic conductivity** See **PERMEABILITY**.

**hydraulic corer (piston corer)** A sampling tube that penetrates marine or lake sediments by hydraulic methods, instead of being either a **\*gravity corer** or a **\*drilling corer**. One version is the **\*Mackereth corer**.

**hydraulic equivalent** **\*Grains** of different density will settle through a fluid at different rates. The concept of hydraulic equivalence is used to relate the size of a mineral grain to the size of a **\*quartz** grain with the same settling velocity. Thus a **\*magnetite** grain (sp. gr. 5.18) of 0.2 mm diameter is the hydraulic equivalent grain size to a 0.5 mm diameter **\*quartz** particle (sp. gr. 2.65).

**hydraulic fracture** An extensional fracture and/or crack-seal vein produced by the expulsion of fluid from a **\*sediment** that is subjected to a rapid

increase in **\*load** pressure and therefore high fluid pressure.

**hydraulic fracturing** See **HYDROFRACTURING**.

**hydraulic geometry** A description of the adjustments made by a stream in response to changes in discharge, both at a cross-section and in the downstream direction. Adjustments are made in width, mean depth, mean velocity, slope, frictional resistance, suspended-sediment load, and water-surface gradient. The relationship between discharge and adjustment is expressed as a power function:  $y = aQ^b$ , where  $y$  is the adjusting variable,  $Q$  is discharge, and  $a$  and  $b$  are coefficients.

**hydraulic gradient** A measure of the change in **\*groundwater** head over a given distance. Maximum flow will normally be in the direction of the maximum fall in head per unit of horizontal distance, i.e. in the direction of the maximum hydraulic gradient. See **HYDRAULIC HEAD**.

**hydraulic head** In general, the elevation of a water body above a particular datum level. Specifically, the energy possessed by a unit weight of water at any particular point, and measured by the level of water in a **\*manometer** at the laboratory scale, or by water level in a well, **\*borehole**, or **\*piezometer** in the field. The hydraulic head consists of three parts: the elevation head (see **ELEVATION POTENTIAL ENERGY**), defined with reference to a standard level or datum; the **\*pressure head**, defined with reference to atmospheric pressure; and the velocity head. Water invariably flows from points of larger hydraulic head to points of lower head, down the **\*hydraulic gradient**.

**hydraulic radius** Ratio between the cross-sectional area of a stream channel and the length of the water-channel contact at that cross-section (the wetted perimeter). It is a measure of channel efficiency: the higher the ratio, the more efficient is the channel in transmitting water.

**hydric** See **MESIC**.

**hydrocarbon** Naturally occurring organic compound containing carbon and hydrogen. Hydrocarbons may be gaseous, solid, or liquid, and include **\*natural gas**, **\*bitumen**, and **\*petroleum**.

**hydrochemistry** The study and representation of the chemical composition of waters, normally those of natural occurrence.

**hydroclast** A *\*clast* produced by a reaction between *\*magma* and water. Compare ALLOCLAST; AUTOCLAST; EPICLAST.

**hydrocollapsibility** A condition of some soils (e.g. *\*loess*) that are susceptible to sudden mechanical collapse when the accumulation of water makes the ground beneath them close to saturation. Hydrocollapsibility is due to the high *\*porosity* of the material combined with weak *\*cementation*.

**hydrofracturing (fracking, hydraulic fracturing)** Process of breaking up rocks under pressure by introducing water or other fluids; usually done to increase *\*permeability* in oil, gas, and geothermal reservoirs. The pressure opens joints, cracks, and bedding planes which can be kept open by the introduction of sand, glass beads, or aluminium balls. Hydrofracturing may occur naturally as a result of internal hydraulic overpressures, e.g. in the formation of porphyry deposits, and is often used to increase recovery from oil and gas wells. See GEOTHERMAL FIELD; PORPHYRY COPPER.

**hydrogen ‘burning’** A thermonuclear process that produces energy by the combination of hydrogen nuclei, and referred to in stellar evolution as the ‘main-stage sequence’. The stage ends when the inner core of hydrogen is exhausted, causing the contraction of the inner part of the star and the expansion of the outer part to produce a red giant. See CARBON ‘BURNING’; HELIUM ‘BURNING’; SILICON ‘BURNING’; NUCLEOSYNTHESIS; THERMONUCLEAR REACTIONS.

**hydrogeologic map** A geologic map that is specially prepared to emphasize features of hydrogeologic importance. For example, rocks are shown not just according to their age or lithology but also as *\*aquifers* or *\*aquicludes*, and details may be included of *\*groundwater* levels, *\*springs*, and water sources. Unlike many geologic maps, they provide a high degree of interpretation for the user.

**hydrogeology** The scientific study of the occurrence and flow of *\*groundwater* and its effects on earth materials.

**hydrograph** A graph showing the plot of water flow in a water course, or the elevation of *\*groundwater* in a *\*borehole* above a particular datum point, against time. The ‘unit hydrograph’ is the name given to a method of calculation which allows rainfall to be converted to stream flow and so

facilitates the prediction of how particular river basins will respond to changing precipitation patterns. The discharge hydrograph shows the flow rate of water against time for a discharging water body. The stage hydrograph shows water level against time.

**hydroids** See [HYDROZOA](#).

**hydro-isostasy** Uplift that occurs in [\\*oceanic](#) crust during [\\*ice](#) ages. The accumulation of ice to form [\\*ice](#) sheets on the continents removes water from the oceans, thus reducing the weight of water loading the ocean floor.

**hydrologic cycle** Representation of the flow of water in various states through the terrestrial and atmospheric environments. Storage points (stages) involve [\\*groundwater](#) and surface water, ice caps, oceans, and the atmosphere. Exchanges between stages involve evaporation and transpiration from the Earth's surface, condensation to form clouds, and precipitation followed by runoff. See also [RESIDENCE TIME](#).

**hydrologic modelling (hydrologic simulation)** The use of small-scale physical models, mathematical analogues, and computer simulations to characterize the likely behaviour of real hydrologic features and systems.

**hydrologic network** An integrated array of meteorological, [\\*groundwater](#)-level, and stream-flow measuring stations which in combination give a complete measurement of the [\\*hydrologic](#) cycle for a particular area. In modern practice the various measuring and [\\*gauging](#) stations are sometimes linked by telemetry to a central monitoring unit for use in flood forecasting. See [FLOOD PREDICTION](#).

**hydrologic regions** The smaller units, with fixed boundaries, into which large tracts of country are divided for the purpose of collecting hydrologic data. The establishment of hydrologic regions allows data to be collected on the same basis from year to year and so facilitates historical analysis. Well-identified regions have a common climate and geologic and topographical structure, so that the [\\*hydrologic](#) cycle operates fairly uniformly within each region.

**hydrologic simulation** See [HYDROLOGIC MODELLING](#).

**hydrology** The study of the [\\*hydrologic](#) (water) cycle. While it involves aspects of geology, oceanography, and meteorology, hydrology emphasizes

the study of bodies of surface water on land and how they change with time.  
See [HYDROGEOLOGY](#).

**Hydrology and Water Resources Programme** A project by the [\\*World Meteorological Organization](#) to promote international collaboration in the evaluation of water resources and the development of hydrological networks and services.



[http://www.wmo.int/pages/prog/hwrp/index\\_en.php](http://www.wmo.int/pages/prog/hwrp/index_en.php)

- The World Meteorological Organization's hydrology and water resources programme.

**hydrolysate** Sediment consisting of undecomposed, finely ground rock and insoluble material derived from weathered primary rocks; typical of [\\*clays](#), [\\*shales](#), and [\\*bauxites](#), with Al, Si, K, and Na as major components.

**hydrolysis** 1. Reaction between a substance and water in which the substance is split into two or more products. At the points of cleavage the products react with the hydrogen or hydroxyl ions derived from water. 2. Process of enriching the soil [\\*adsorption complex](#) with hydrogen after [\\*exchangeable](#) metallic ions have been replaced by hydrogen ions.  
*Compare* [WEATHERING](#).

**hydromuscovite** See [ILLITE](#).

**hydrophone** A microphone used to detect acoustic (including seismic) waves under water. A number of hydrophones are linked together to form a [\\*streamer](#).

**hydrosphere** The whole of that body of water which exists on or close to the surface of the Earth. The hydrosphere formed as the Earth cooled and atmospheric water condensed.

**hydrospire** See [BLASTOIDEA](#).

**hydrostatic stress** The component of [\\*confining](#) pressure derived from the weight of pore water in the column of rock above a specified level. All [\\*principal](#) stresses are equal and changes in hydrostatic pressure produce changes only in the volume and density of the material. It can be simulated

experimentally by enclosing material in a jacket and pumping in liquids to produce equal pressures throughout.

**hydrostratigraphic units** Sections of a geological formation that exhibit similar hydraulic properties (see [HYDROGEOLOGY](#)), regardless of their composition. These units may be very extensive. The region of Alberta containing the Athabasca [\\*tar sands](#), for example, covers approximately 67 260 km<sup>2</sup> and comprises three hydrostratigraphic units. These consist of [\\*Cretaceous](#) and [\\*Holocene](#) sediments with alternately vertical and horizontal water flow, Upper [\\*Devonian](#) strata with predominantly horizontal flow, and formations lying beneath [\\*evaporites](#) where water flows horizontally towards a [\\*fault zone](#). The concept of hydrostatic units is important in planning, when the movement of [\\*groundwater](#) must be predicted, and in tracing the source of groundwater pollutants.

**hydrothermal activity** Any process associated with [\\*igneous](#) activity involving the action of very hot waters. The waters involved can be derived directly from an igneous [\\*intrusion](#) (i.e. [\\*juvenile water](#)) as a residual fluid formed during the late stages of crystallization of the body, or can be external [\\*groundwater](#) heated during crystallization of the intrusion. The hydrothermal fluids can react with and alter the rocks through which they pass, or can deposit minerals from solution. Hydrothermal reactions include [\\*serpentinization](#), chloritization (see [CHLORITE](#)), saussuritization (see [SAUSSURITE](#)), uralitization (see [URALITE](#)), and [\\*propylitization](#); whilst hydrothermal vein and replacement mineral deposits include Cu, Pb, and Zn sulphides. Hydrothermal activity should not be confused with geothermal activity which involves the convection and movement of hot waters but is not necessarily connected with an igneous intrusion (see [GEOTHERMAL FIELD](#); [GEOTHERMAL GRADIENT](#)).

**hydrothermal metamorphism** [\\*Metamorphism](#) that occurs when hot waters, laden with minerals and chemically active, intersect surrounding [\\*country](#) rock. It may occur at low temperature and pressure, typically yielding [\\*pegmatites](#), [\\*serpentinite](#), and soapstone (see [TALC](#)), often as an after-effect of [\\*igneous](#) activity. Oceanic hydrothermal metamorphism occurs at [\\*rift](#) centres, where [\\*magma](#) oozes on to the ocean floor to form [\\*pillow](#) lava.

**hydrothermal mineral** A mineral formed by precipitation from a very hot hydrothermal fluid (see [HYDROTHERMAL ACTIVITY](#)) as it passes down a temperature or pressure gradient. Common hydrothermal minerals precipitated in [\\*veins](#) and cavities are [\\*quartz](#), [\\*fluorite](#), [\\*galena](#), and [\\*sphalerite](#).

**hydrothermal vent** A place on the ocean floor, on or adjacent to a mid-ocean [\\*ridge](#), from which there issues water that has been heated by contact with molten rock, commonly to about 300 °C. The vent water often contains dissolved sulphides. These are oxidized by chemosynthetic bacteria, which fix carbon dioxide and synthesize organic compounds. Near the vents, at temperatures up to 40 °C, there are highly productive communities comprising animals that utilize the organic compounds or live symbiotically with the chemosynthetic bacteria; these organisms support carnivores and detritivores. These communities include beard worms (phylum [\\*Pogonophora](#)) that completely lack a digestive tract, Munidopsis crabs (superfamily Galatheaidea), giant clams (e.g. *Calyptogena magnifica*), mussels, acorn worms (class Enteropneusta), and many more. Vent fluids containing high concentrations of iron, manganese, and copper tend to be hot (about 350 °C) and black. They are known as 'black smokers'. 'White smokers' flow more slowly, are cooler, and contain high concentrations of arsenic and zinc. See [HYDROTHERMAL ACTIVITY](#), [HYDROTHERMAL MINERAL](#). See also [COLD SEEP](#).

**hydrovolcanic processes** A sequence of events initiated by the interaction of a body of [\\*magma](#) and water external to the magma system. For example, when a rising body of magma encounters external [\\*groundwater](#) it converts the groundwater to steam which expands rapidly, fragmenting the surrounding rocks and magma to generate a phreatomagmatic eruption (see also [PHREATIC ACTIVITY](#)). When lava is erupted under water, the water removes heat rapidly from the outer margin of the flow, which is chilled to a glass, and fractured and fragmented during contraction. The fragmented glassy margin of the flow forms a [\\*hyaloclastite](#) envelope around the flow. See [PILLOW LAVA](#).

**hydroxides** Applied to [\\*minerals](#) whose chemical composition includes the OH radical, and often to those with water molecules, whose presence gives hydrated oxides. Among the important hydroxide minerals are: [\\*brucite](#),

Mg(OH)<sub>2</sub>; [\\*gibbsite](#), Al(OH)<sub>3</sub>; [\\*diaspore](#), α-AlO(OH); [\\*boehmite](#), γ-AlO(OH); [\\*limonite](#), FeO(OH).nH<sub>2</sub>O; and [\\*goethite](#), α-FeO(OH).

**hydroxyapatite** A hydrated calcium phosphate mineral, which also contains fluoride, chloride, and carbonate calcium salts. It is often formed as a consequence of [\\*biomineralization](#), producing hard structures, such as [\\*bone](#).

**Hydrozoa (hydroids)** (phylum [\\*Cnidaria](#)) Class of multicellular, mainly marine animals with cells arranged in two layers, the epidermis and the gastrodermis (endodermis), separated by a gelatinous mesogloea. These enclose a continuous digestive cavity (coelenteron), which communicates directly with the exterior by a single aperture (mouth) and is lined by a gastrodermis. The gastrodermis lacks nematocysts. Hydrozoa are Lower [\\*Cambrian](#) to Recent. *See also* [MILLEPORINA](#); [STYLASTERINA](#).

**hyetograph** A chart that shows the amount and duration of rainfall at a particular place. The instrument producing the chart is also called a hyetograph. It comprises a reservoir in which rainwater collects, with a float connected mechanically to a pen, and a rotating drum bearing the graph paper. As the water level rises and falls in the reservoir the movement is transmitted to the pen, which traces a line on the graph paper.

**hygromagnetophile elements** *See* [INCOMPATIBLE ELEMENTS](#).

**hygrometer** Instrument for measuring atmospheric humidity. Types include the wet-bulb–dry-bulb, dew-point, and [\\*hair](#) hygrometers, and there is one type based on electrical resistance. *See also* [PSYCHROMETER](#).

**hygroscopic nucleus** Microscopic particle (e.g. of sulphur dioxide, salt, dust, or smoke) in the free air, on which water vapour may condense to form droplets. [\\*Aerosols](#) that are soluble in water (e.g. salt or sulphuric acid) can induce condensation in unsaturated air, e.g. salt nuclei can induce it at a [\\*relative humidity](#) of less than 80%. The size of nuclei may be from 0.001 μm to more than 10 μm (i.e. ‘giant’ nuclei such as particles of sea-salt). *See also* [AITKEN NUCLEI](#); [NUCLEUS](#).

**hygroscopic water** Water absorbed from the atmosphere and held very tightly by the soil particles, so that it is unavailable to plants in amounts sufficient for them to survive. *Compare* [CAPILLARY MOISTURE](#).

**hygrothermograph (thermo-hygrograph)** An instrument used for the continuous recording of both the temperature and humidity of the air, on separate traces.

***Hylonomus lyelli*** Ancient, probably the oldest-known, **\*reptile**. *Hylonomus* is a stem reptile and a member of the **\*Captorhinomorpha**. It was first found in fossilized tree stumps in the **\*Coal Measures** of Nova Scotia. The skull roof of this species was fully ossified and there were no openings behind the eye sockets. *Hylonomus* measured some 25 cm and possessed a long tail.

**hyoid** A bone or bones developed from the second visceral arch: it or they support the tongue.

**Hyalolithida** (phylum **\*Mollusca**, class **\*Calypptomatida**) Order of pyramid-shaped calypptomatids which have conical embryonic chambers. The **\*operculum** possesses one or two pairs of **\*muscle** scars which are bilaterally symmetrical. The shell is externally undifferentiated, and internally it is non-septate. The order ranges in age from Lower **\*Cambrian** to Middle **\*Permian**. The main genus, *Hyalolithes*, had worldwide distribution during the Cambrian, with over 300 species.

**hyp-** See **HYPO-**.

**hypabyssal** Applied to medium-grained, **\*intrusive \*igneous** rocks which have crystallized at shallow depth below the Earth's surface. There is, however, no sharply defined or agreed depth limit to the term 'shallow'.

**hyper-** From the Greek *hyper* meaning 'beyond' or 'over', a prefix meaning 'exceeding' or 'greater than normal'.

**Hyper-Angular Rainbow Polarimeter (HARP)** A joint 3-unit **\*CubeSat** mission between the University of Maryland Baltimore County and Utah State University Space Dynamics Laboratory to measure the microphysical properties of cloud water and ice particles. The **\*nanosatellite** was launched on 18 April 2016, from Cape Canaveral, Florida, into a near-circular **\*polar orbit** at an altitude of about 400 km.

**Hyperion (Saturn VII)** One of the major satellites of **\*Saturn**, with a radius measuring  $185 \times 140 \times 113$  km; visual albedo 0.19–0.25. It was discovered in 1848 by W. Bond.

**hypermorphosis** Acceleration of development so that the organism reaches its adult size and form well before the attainment of sexual maturity, and continues to develop into a 'super-adult'.

**hyperpycnal flow** At a river mouth, a flow of river water that is denser than the water in the basin receiving it. This occurs during floods. The denser water flows beneath the basin water, as a density current, carrying sediment beyond the shore and inhibiting the progradation of a delta. *Compare* HOMOPYCNAL FLOW; HYPOPYCNAL FLOW.

**hypersolvus granite** A \*granite which has crystallized above the solvus temperature and hence contains only one \*alkali feldspar type. The solvus temperature is that temperature below which two alkali feldspars of contrasting composition are in equilibrium. Granites which crystallize below the solvus temperature (subsolvus granites) immediately crystallize two alkali feldspar types, one K-rich, the other Na-rich. The feldspars from either hyper- or subsolvus granites may suffer sub-\*solidus \*exsolution if cooling is slow enough. This is an equilibrium process which takes place when the rock is entirely solid and involves splitting of the feldspars into K- and Na-rich lamellae by internal \*diffusion of \*ions, to give perthitic (see PERTHITE) or antiperthitic feldspars.

**hypersolvus syenite** See SYENITE.

**Hyperspectral Precursor and Application Mission (PRISMA; PRecursore IperSpettrale della Missione Applicativa)** A \*minisatellite mission by the Italian Space Agency to provide medium-resolution hyperspectral imaging to monitor natural resources and atmospheric characteristics. It was launched on 22 March 2019.

**hypersthene** A member of the \*orthorhombic \*pyroxenes with approximately equal amounts of Fe and Mg in its composition  $MgFeSi_2O_6$ , it is a \*mineral occurring within the series \*enstatite to (ortho)\*ferrosilite, although the name hypersthene has fallen into disuse (with the more general term orthopyroxene followed by the exact composition being preferred); sp. gr. 3.5; \*hardness 5–6; \*orthorhombic; green to greenish, or brownish-black; vitreous \*lustre; \*crystals can be \*prismatic or \*tabular, but usually irregular grains; \*cleavage good, prismatic {110}; occurs in iron-rich, bafic, \*igneous rocks, e.g. noritic \*gabbro, \*trachytes, and \*andesites.

**hyperthermic** See PERGELIC.

**hyperthermophile** An *\*extremophile* (domain *\*Archaea*) that thrives in environments where the temperature is extremely high, in some cases preferring a temperature of about 105 °C, tolerating 113 °C, and failing to multiply below 90 °C. Compare THERMOPHILE.

**hypichnia** See TRACE FOSSIL.

**hypidiomorphic fabric** See HYPIDIOTOPIC FABRIC.

**hypidiotopic fabric** A rock *\*texture* characterized by the presence of minerals some of which show their *\*crystal* form (i.e. some *\*euhedral* habits). ‘Hypidiotopic fabric’ refers to *\*sedimentary rocks*; for *\*igneous* or *\*metamorphic rocks*, ‘hypidiomorphic fabric’ is used.

**hypo-** (**hyp-**) From the Greek *hupo* meaning ‘under’, a prefix meaning ‘below’, ‘slightly’, or ‘lower than normal’. ‘Hypo-’ is generally used before words beginning with a consonant, ‘hyp-’ before words beginning with a vowel.

**hypocentre (focus)** The actual location, usually within the Earth, of the first motion of an *\*earthquake*, i.e. the location of the focus of the earthquake. The point at the Earth’s surface overlying the hypocentre is the *\*epicentre*.

**hypocrystalline (merocrystalline)** A *\*textural* term referring to *\*igneous* rocks, or parts of igneous rocks, which contain both *\*crystals* and *\*glass*.

**hypogene** *\*Mineral* deposit formed by generally ascending solutions in or from below the Earth’s *\*crust*; or processes such as *\*volcanicity* operating within the crust.

**hypolimnion** The lower, cooler, non-circulating water in a thermally stratified lake in summer. If, as often occurs, the *\*thermocline* is below the compensation level, the dissolved oxygen supply of the hypolimnion depletes gradually: replenishment by *\*photosynthesis* and by contact with the atmosphere are prevented. Re-oxygenation is possible only when the thermal stratification breaks down in autumn.

**hyponome** A tube or funnel occurring in cephalopods (*\*Cephalopoda*), through which water is expelled from the *\*mantle cavity*. It is used for jet

propulsion and its presence may cause an embayment or slit in the ventral margin of a cephalopod shell, called the 'hyponomic sinus'.

**hyponomic sinus** See [HYPONOME](#).

**hypopycnal flow** At a river mouth, a flow of river water that is less dense than the water in the basin receiving it. The river water is buoyant and flows above the basin water. This is typical of the situation where a river enters the sea, because of the difference in density between fresh and salt water.

*Compare* [HOMOPYCNAL FLOW](#); [HYPERPYCNAL FLOW](#).

**hypostomal suture** See [CEPHALIC SUTURE](#).

**hypostratotype** (**reference section, auxiliary reference section**) An additional, subordinate *\*stratotype*, selected after the establishment of a stratigraphic unit, and in another region, to supplement the information contained in the original stratotype (the *\*holostratotype*). *Compare* [PARASTRATOTYPE](#). *See also* [LECTOSTRATOTYPE](#); [NEOSTRATOTYPE](#).

**hypotheca** See [DINOPHYCEAE](#).

**hypothermal** Applied to a mineral deposit originating at great depth and at temperatures between 300 and 500 °C, below the *\*mesothermal* zone.

**hypothesis** An idea or concept that provides a basis for arguments or explanations which can be tested by experimentation. In inductive or inferential statistics, the hypothesis is usually stated as the converse of the expected results, i.e. as a null hypothesis ( $H_0$ ). This helps workers to avoid reaching a wrong conclusion, since the original hypothesis  $H_1$  will be accepted only if the experimental data depart significantly from the values predicted by the null hypothesis. Working in this negative way carries the risk of rejecting a valid research hypothesis even though it is true (a problem with small data samples); but this is generally considered preferable to the acceptance of a false hypothesis, which would tend to be favoured by working in the positive way.


**hypothesis testing** In statistics, the comparison of a data set with a particular theory, called a null hypothesis, in order to see if it deviates significantly from this theory, in which case an alternative theory may more appropriately explain the data. A statistic is considered significantly deviant

from the null hypothesis when the statistic falls outside the **\*confidence** interval predicted by the null hypothesis for a defined confidence level. An example is the **\*chi-squared test**.

**Hypsilophodontidae** Family of bipedal (ornithopod), **\*ornithischian** **\*dinosaurs**, ranging from the Upper **\*Triassic** to the Upper **\*Cretaceous**. Hypsilophodon itself, from the **\*Wealden** of the Lower Cretaceous, is the most primitive of the ornithopod dinosaurs known to date, despite its relatively late appearance in the **\*Mesozoic** fossil record.

**Hypsithermal** See FLANDRIAN.

**hypsographic curve** A graphic representation of the elevation and depth of points on the surface of a planet with reference to a datum; on Earth sea level is used as the datum. This shows that on Earth, on average land projects about 850 m above sea level and the average depth of ocean basins is about 3730 m below sea level. Compare **HYPSONOMETRIC CURVE**.




**Hypsographic curve**

**hypsonometric curve** A graphic representation of the proportion of the surface of a planet that is elevated or depressed in relation to a datum. Compare **HYPSONOGRAPHIC CURVE**.

***Hyracotherium*** Known formerly as Eohippus (the ‘dawn horse’), the earliest known **\*perissodactyl**, which is placed in the family **\*Equidae**. It was only 27 cm high, the size of a fox terrier: it was short-faced with low-crowned cheek teeth, and had four toes on the forefeet, and three on the hind. Abundant in the Early **\*Eocene** of N. America and Europe, it was recently discovered in **\*Palaeocene** deposits in Mongolia. It was a browser dwelling in forest glades, and the likely ancestor for all the horses. Because of its small size, when the fossils were first found, in Europe, they were mistakenly associated with the African hyraxes, hence the name. In America the fossils were identified correctly and the name ‘Eohippus’ was adopted, but according to the rules of **\*taxonomic** nomenclature ‘Hyracotherium’ takes precedence.

**hysteresis loop** The curve formed when the intensity of magnetization acquired by a **\*ferromagnetic** material is plotted against the magnitude of a direct magnetic field applied in one direction and then in the opposite direction. This loop is used to define the **\*coercivity**, **\*saturation magnetization**, and **\*susceptibility** of such materials.

**hystricospheres** Cysts in which the fertilized eggs (zygotes) of dinoflagellates (see **DINOPHYCEAE**) remain dormant during times when conditions are unfavourable. Fossilized hystricospheres, comprising a cellulose skeleton bearing spines or processes by which genera can be identified, are important in **\*biostratigraphy**.



**IAEG** See INTERNATIONAL ASSOCIATION FOR ENGINEERING GEOLOGY AND THE ENVIRONMENT.

**-ian** The preferred suffix that is added to the geographical location of a *\*type* section or *\*type area* to form the name of a *\*stage* or *\*age*, e.g. ‘*\*Dolgellian*’, ‘*\*Frasnian*’, ‘*\*Barremian*’.

**Iapetus (Saturn VIII)** One of the major satellites of *\*Saturn*, with a radius of 718 km; mass  $15.9 \times 10^{20}$  kg; mean density  $1020 \text{ kg/m}^3$ ; visual albedo 0.05–0.5. It was discovered in 1671 by G. D. Cassini.

**Iapetus Ocean (proto-Atlantic)** The late *\*Precambrian* and early *\*Palaeozoic* ocean that lay between *\*Baltica* and *\*Laurentia*. The *\*oceanic* crust and upper *\*mantle* of the Iapetus Ocean floor is presumed to have been subducted during the early Palaeozoic, and the ocean to have disappeared completely by the latest *\*Silurian*–early *\*Devonian* (about 400 Ma ago). The ancient *\*suture* is thought to extend from WSW to ENE across the Solway Firth and Borders Region, Scotland. The *\*Caledonian* *\*orogenic* belt, which removed all trace of the Iapetus Ocean, extends all along the border between the two ancient *\*cratons*, affecting the areas now known as Norway, eastern Greenland, Scotland, northern England, Wales, Ireland, eastern Canada, and the eastern USA.

***Iberomesornis*** The best-known of several Early *\*Cretaceous* birds, from Las Hoyas, central Spain. It is the first bird to show evidence of a perching foot.

**IBEX** See INTERSTELLAR BOUNDARY EXPLORER.

**Ibexian** A N. American *\*stage* (491–471.8 Ma ago) of the *\*Palaeozoic* *\*era*, preceded by the Sunwaptan and followed by the *\*Whiterockian*.

**Icarus** A \*solar system asteroid (No. 1566), diameter 1.4 km; approximate mass  $10^{12}$  kg; rotational period 2.273 hours; orbital period 1.12 years. Its highly eccentric orbit crosses the orbit of Earth.

**ICE** See INTERNATIONAL SUN—EARTH EXPLORER.

**ice** **1.** Water which has frozen into a crystal lattice. Pure water freezes at  $0^{\circ}\text{C}$  at 10 MPa (megapascals) pressure. The presence of salts in solution depresses the freezing point of water. Liquid water has its maximum density at  $4^{\circ}\text{C}$ , in consequence of which ice floats on water. With increasing pressure, a series of denser polymorphs of ice forms, each designated by a Roman numeral, ordinary ice being ice I. **2.** Several properties and varieties of ice are important in geomorphological \*processes. Expansion on freezing (9.05% in specific volume) generates very high pressures. In an enclosed space in the laboratory the pressure reaches 216 MPa at  $-22^{\circ}\text{C}$  but reaches only about 10% of this when unenclosed, as in nature. Such ice I converts into the denser ice III at lower temperatures, but the pressure exerted by it changes little. ‘Ground ice’ forms when \*interstitial water freezes, and this may bring about heaving as well as frost wedging. ‘Glacier ice’ is a relatively opaque mass of interlocking crystals, and has a density of 0.85–0.91  $\text{g}/\text{cm}^3$ . \*Regelation ice’ is relatively clear and is formed by the freezing of meltwater beneath a temperate \*glacier. **3.** In planetary geology other ices are important. Water ice condenses at 160 K at \*solar nebular pressures and appears in abundance forming the surfaces of the \*Galilean satellites Europa, Ganymede, and Callisto. The \*satellites of the \*jovian planets are mostly mixtures of water ice and rock. Water ice will exist in high-pressure polymorphs (e.g. ice VIII, density  $1670 \text{ kg}/\text{m}^3$ ) above about 1.5–2.0 MPa in satellite interiors. Other possible ices important in satellites (e.g. Titan) include  $\text{NH}_3 \cdot \text{H}_2\text{O}$ ,  $\text{CH}_4 \cdot n\text{H}_2\text{O}$ , and  $\text{H}_2\text{O} \cdot \text{CO}_2$ .

**ice ages** Periods when ice has accumulated at the poles and the continents have been glaciated repeatedly. Exactly why glaciation occurred is not clear. There are suggestions of a middle \*Precambrian glaciation about 2300 Ma in N. America, S. Africa, and Australia. More information exists to suggest that the Earth was glaciated between 950 and 615 Ma ago, and there are at least two glacial \*horizons in Africa, Australia, and Europe. There is good evidence for a glaciation at the end of the \*Ordovician in N. Africa, but glacial deposits described from elsewhere at this period are problematical,

so the extent of the glaciation is not known. The Permo–Carboniferous glaciation of S. America, S. Africa, India, and Australia was widespread and is well documented. There is no evidence for further glaciation until the **\*Quaternary**. Suggestions have been made for other ice ages during the **\*Palaeozoic** but evidence for them is sparse. The **\*Pleistocene** ice age is the best documented, but there is undoubted evidence of earlier glaciations in the geologic record.

**iceberg calving** See CALVING.

**ice blink** Whitish appearance seen above the horizon, caused by light reflection over an ice or pack-ice surface.

**ice cap** Ice mass less than 50 000 km<sup>2</sup> in area, but large enough to submerge the underlying topography and whose flow behaviour is a consequence of its size and shape. It consists of a central **\*ice** dome together with outlet **\*glaciers** radiating from the periphery.

**Ice, Cloud, and land Elevation Satellite (ICESat)** A NASA satellite mission that ran from 2003 to 2009, measuring ice sheet mass balance, cloud and **\*aerosol** heights, surface topography, and vegetation characteristics. The satellite was decommissioned on 17 August 2010.



<http://icesat.gsfc.nasa.gov/icesat/index.php>

- A NASA mission to measure ice sheet mass balance, cloud and aerosol heights.

**ice core** A long cylinder of ice that is obtained by drilling vertically downwards through an **\*ice** sheet. The seasonal accumulation of ice produces bands by which the age of the core can be determined and sections of it dated. Analysis of ice from particular sections reveals information regarding past climate (see **OXYGEN-ISOTOPE ANALYSIS**). Dust trapped in the ice indicates arid conditions (and low temperatures) and bubbles of air record the chemical composition of the atmosphere. Cores taken from the Greenland ice sheet (the Greenland Ice Core Project (**\*GRIP**) and Greenland Ice Sheet Project (**\*GISP**) have obtained samples up to 20 000 years old. Cores drilled near the Vostok Station, Antarctica, have reached ages of more than 40 000 years.

**ice crystal** Frozen water composed of crystalline structures, e.g. needles, dendrites, hexagonal columns, and platelets. Ice clouds are composed entirely of ice crystals.

**IceCube** A 3-unit **\*CubeSat** mission by **\*NASA**'s Goddard Space Flight Center to measure cloud ice. It was launched on 18 April 2017 to the International Space Station and released later into a near-circular orbit at an altitude of about 400 km.

**ice dome** A main component (with the **\*outlet** glacier) of an **\*ice** sheet or **\*ice** cap. It has a convex surface form, of parabolic shape, and tends to develop symmetrically over a land mass. Its thickness often exceeds 3000 m.

**ice field** A nearly level field of **\*ice**, whose area may range from about 5 km<sup>2</sup> to near-continental size, formed when the land surface is sufficiently high or uniform for ice to accumulate. It differs from an **\*ice** cap in that it lacks a domed form and its flow is controlled by the underlying relief.

**icehouse period** A time during which glaciers reached their maximum extent, ocean waters were well mixed and oxygenated, and sea levels were low. In **\*cyclostratigraphy**, fourth- and fifth-order cycles were predominant. *Compare* GREENHOUSE PERIOD.

**Iceland low** The region of the N. Atlantic in which the average value of atmospheric pressure is low, owing to the frequency of low-pressure systems (cyclones or **\*depressions**) moving into and occupying the region. Any one of these systems, when present on an individual day, may be called 'an Iceland low', but the term is used mostly to describe the statistical or climatic feature.

**ice mound** A low mound, up to about 2 m high, 20 m long, and 10 m wide, that is found in **\*tundra** environments. It consists of a core of clear ice covered by a layer 0.3–1.0 m thick of sand, gravel, and soil, with tundra plants growing on it. *See also* PINGO.


**ice nucleus** Crystalline, microscopic particle, either of ice or some other substance, which can induce the growth of ice crystals upon it from saturated air in cloud at temperatures of about –25 °C or below. Deposition of water in this way—directly from vapour to form ice, without forming liquid in between—is called 'sublimation'. During growth, splintering of

the crystals (e.g. as a result of updraughts) causes large numbers of new nuclei to be formed. Compare [FREEZING NUCLEI](#).

**ice-rafted detritus** \*Terrigenous material that is transported within the ice matrix of a \*glacier and deposited in marine or \*lacustrine sediments when the ice melts.

**ICESat 1.** See [ICE, CLOUD, AND LAND ELEVATION SATELLITE](#). **2.** ICESat-2 is a follow-up mission launched on 15 September 2018, from California, into a near-polar, low Earth orbit at an altitude of 496 km.

**ice sheet** A large ice mass, with an area usually greater than about 50 000 km<sup>2</sup>, made up of \*ice domes and \*outlet glaciers. The Antarctic ice sheet is the world's largest, with an area of about  $13.9 \times 10^6$  km<sup>2</sup> and a mean thickness of about 2100 m.



**Ice sheet (cross section)**

**ice shelf** The outer part of an \*ice cap or \*ice sheet that extends into and over the sea. It typically ends in a cliff that may be 30 m high, and the total ice thickness may be 200 m. Ice wastage is by \*calving (the breaking away of ice blocks) and by bottom melt.

**ice stream** See [OUTLET GLACIER](#).

**ice wedge** Tapering, vertically layered mass of ice, about a metre wide at the top and extending downward for some 3–7 m. It results from contraction cracking of the ground during extreme cold, followed by water penetration from the \*active layer and subsequent freezing. It is

characteristic of uniform sediment, such as river **\*alluvium**, under **\*periglacial** conditions.

**ice-wedge polygon** See **PATTERNED GROUND**.

**ichnoclast** A **\*trace fossil** that has been reworked by a later organism.

**ichnocoenosis** An assembly of **\*trace fossils** made by members of a single community.

**ichnofabric** The structure and texture of a sediment that is produced by the activity of living organisms.

**ichnofacies** A rock or sequence of rocks characterized by its **\*lithology**, inorganic **\*sedimentary structures**, and specific **\*trace fossils**. The traces represent the behaviour of **\*fossil** organisms under particular environmental conditions. The relative abundance of different trace fossil **\*ethotypes** is critical to the determination of the correct ichnofacies.

**ichnofossil** See **FOSSILIZATION**; **TRACE FOSSIL**.

**ichnogenus** A group of **\*trace fossils** that is given a name because the similarity of the traces suggests they were made by closely related species of organisms. Ichnological taxonomy, which applies the principles of biological nomenclature to non-biological material, is governed by the *International Code of Zoological Nomenclature*. Above the level of genus, the Code indicates names should be used formally only to the family level; at higher levels all names are informal. Names of ichnogenera are conventionally written italicized and with a capital initial; ichnogenus is abbreviated as igen. See **DIPLOCRATERION**; **RHIZOCORALLIUM**; **ZOOPHYCUS**.

**ichnoguild** A group of **\*ichnospecies** that shared particular resources or behaviour in the manner of a **\*guild**.

**ichnology** The study of the tracks, burrows, and other traces made by living organisms on and within a substrate. If the traces are recent and made by organisms that are still living, the study is called neoichnology. If the organisms are long disappeared and evidence of their presence is preserved as **\*trace fossils**, the study is called palaeoichnology.

**ichnospecies** A species (trivial) name assigned to **\*trace fossils** within an **\*ichnogenus** and conventionally written italicized and with a lower case

initial; ichnospecies is abbreviated as isp.

**ichnotaxobase** Any morphological feature of a *\*trace fossil* that can be used in classification.

**ichnotaxonomy** The formal classification of *\*trace fossils*.

***Ichthyornis*** See AVES.

**Ichthyosauria** (class *\*Reptilia*, subclass Ichthyopterygia) Order of so-called 'fish lizards', and the only order in the subclass, the first members of the group date from the *\*Triassic* and were primitive in type. The typical, shark-like form, *Ichthyosaurus*, appeared in the *\*Jurassic*, when the group in general was especially common. Ichthyosaurs disappeared before the end of the *\*Cretaceous*.

***Ichthyostega*** *\*Amphibians* in the form of *Ichthyostega* and *Ichthyostegopsis* first appear in the Upper *\*Devonian*. *Ichthyostega* shows refinements of the skull, and the development of strong limbs. It retains a long, rather fish-like tail, and links with the *\*crossopterygian* fish are confirmed by the presence of teeth with a labyrinthine infolding of the enamel. *Ichthyostega* grew to just under 1 m in length. An unexpected finding about *Ichthyostega* and its relatives, from recently discovered, more complete specimens, is that they had more than the five digits per limb that characterize modern tetrapods: *Ichthyostega* had seven on the hind feet (the forefeet are still unknown), and *Acanthostega* had eight on both fore- and hind feet.

***Ichthyostegopsis*** See ICHTHYOSTEGA.

**ICP** See INDUCTIVELY COUPLED PLASMA EMISSION SPECTROMETRY.

**-id** See -IDE.

**Ida** A *\*solar* system asteroid (No. 243), measuring 58 × 23 km; approximate mass 10<sup>17</sup> kg; rotational period 4.633 hours; orbital period 4.84 years. Images of Ida from *\*Galileo*, taken on 23 August 1993, showed it has a small satellite, later named Dactyl.

**Idamean** A *\*stage* (497–494 Ma ago) of the Upper *\*Cambrian* of Australia, underlain by the *\*Mindyallan* and overlain by the Post-Idamean.

**-ide (-id, -ides)** From the Greek *ides* meaning ‘son of’, a suffix attached to the name of an element which is a member of a series (e.g. actinide, halide), or to the more electronegative element or radical in a binary compound (e.g. sodium chloride).

**-ides** See **-IDE**.

**idioblastic** A **\*textural** term referring to **\*metamorphic rocks** in which the grains display fully developed **\*crystal** forms.

**idiomorphic** See **EUHEDRAL**.

**idiomorphic fabric** See **IDIOTOPIC FABRIC**.

**idiotopic fabric** The **\*fabric** developed in a crystalline **\*sedimentary rock** in which most of the **\*crystals** are **\*euhedral**. In an **\*igneous** or **\*metamorphic rock** the term ‘idiomorphic fabric’ is used.

**idocrase (vesuvianite)** A mineral, with the formula  $\text{Ca}_{10}(\text{Mg,Fe})_2\text{Al}_4\text{Si}_9\text{O}_{34}(\text{OH,F})_4$ , closely related to the **\*garnet** group; sp. gr. 3.4; **\*hardness** 6.5; green; normally **\*massive** or granular; occurs in contact-metamorphosed (see **THERMAL METAMORPHISM**) **\*limestones** along with **\*grossular**, **\*scapolite**, and **\*wollastonite**.

**Idwian** A **\*stage** (439–437.5 Ma ago) of the Lower **\*Silurian**, underlain by the **\*Rhuddanian** and overlain by the **\*Fronian**.

**I<sub>f</sub>** See **FRACTURE SPACING INDEX**.

**IfSAR** See **INTERFEROMETRIC SYNTHETIC APERTURE RADAR**.

**IGC** International Geological Congress.

**igen.** Abbreviation for **\*ichnogenus**.

**igneous** Applied to one of the three main groups of rock types (igneous, **\*metamorphic**, and **\*sedimentary**), to describe those rocks that have crystallized from a **\*magma**.

**igneous rock classification** Any scheme for categorizing **\*igneous** rocks. The simplest classifications define rocks by their chemical and **\*mineral** composition, as **\*felsic**, **\*intermediate**, **\*mafic**, and **\*ultramafic**, and by their **\*texture**, as **\*intrusive** and coarse-grained, **\*extrusive** and fine-grained,

or glass. In coarse-grained rocks the **\*crystals** are visible to the naked eye; in fine-grained rocks they are microscopic; and in glass there are no crystals (and therefore no minerals). The table shows a simple classification with examples. *See also* [QAPF CLASSIFICATION](#).

<u>Occurrence</u>	<u>Composition</u>			
	Felsic	Intermediate	Mafic	Ultramafic
Intrusive	Granite	Diorite	Gabbro	Peridotite
Extrusive	Rhyolite	Andesite	Basalt	Komatiite

**ignimbrite** A poorly sorted, **\*pyroclastic** rock body formed by deposition from a pumiceous pyroclastic flow. The passage of one pyroclastic flow deposits one ignimbrite flow unit with a number of layers directly related to the components of the flow. Ignimbrite flow units deposited by the passage of successive flows can accumulate into a loosely consolidated, composite sheet. Where the rock-body temperature and the accumulating overburden load are both sufficiently high, sintering and flattening of **\*groundmass** shards and **\*pumice \*clasts** can occur in the lower part of the body, producing a welded zone within the ignimbrite body which is then characterized by the development of a **\*eutaxitic** texture. Hot gases escaping through the ignimbrite deposit can precipitate minerals between the loosely packed shards and pumice in the upper parts of the flow, creating a lithified **\*sillar** horizon in which the glass of the pumice and shards is devitrified. Ignimbrites are found on all scales, from a few hundred metres to more than 100 km long, and from one metre to tens of metres thick. The geometry of the ignimbrite sheet is described using the aspect ratio. *See* [HIGH-ASPECT-RATIO IGNIMBRITE](#); [LOW-ASPECT-RATIO IGNIMBRITE](#).

**IGRF** *See* [INTERNATIONAL GEOMAGNETIC REFERENCE FIELD](#).

**IGTS** *See* [ISOBARIC GAS-TIGHT SAMPLER](#).

**Iguanodontidae** **\*Jurassic–\*Cretaceous** family of bipedal (ornithopod), **\*ornithischian \*dinosaurs**, whose best-known representative is Iguanodon, remains of which have been found in Europe, Asia, and Africa.

**ijolite** A medium- to coarse-grained, ultra-alkaline \*plutonic \*igneous rock, consisting of \*essential \*nepheline, \*aegirine (or aegirine-\*augite, or sodic \*diopside) with or without melanite \*garnet, and \*accessory \*apatite, \*sphene, and \*cancrinite. The rock can be considered an undersaturated alkali \*syenite (see SILICA SATURATION).

**ilium** In \*tetrapods, the dorsal section of the pelvis, which articulates with one or more sacral \*vertebrae.

**Illinoian** The third (0.55–0.4 Ma ago) of four glacial \*stages recognized in N. America. It is represented by deposits from ice moving from the north-east, and in formerly glaciated areas to the west pollen evidence suggests that mean annual temperatures were 2–3 °C cooler than they are now. At the end of this period the climate became warmer and drier. The Illinoian is approximately equivalent to the \*Mindel and \*Riss glacials of the Alps.

**illite (hydromuscovite)** Common \*clay mineral and important member of the 2:1 group of \*phyllosilicates (sheet silicates) with the formula  $K_{1-1.5} Al_4[Si_{7-6.5} Al_{1-1.5} O_{20}](OH)_4$ , which possesses an overall negative charge due to incomplete charge balance; sp. gr. 2.6–2.9; \*hardness 1–2; \*monoclinic; \*crystals form tiny flakes; formed by the \*weathering decomposition or \*hydrothermal alteration of \*muscovite or \*feldspar.

**illuminator** The light source used in a transmitted- or \*reflected-light microscope. For most routine work, an incandescent, tungsten filament lamp with a variable rheostat is used. The colour temperature is normally about 2800 K and a pale blue filter may be used to provide a more daylight-like colour temperature. The standard illuminating system may contain additional lenses, diaphragms, and a \*polarizer.

**illuviation** Process of deposition (inwashing) of soil materials, either from suspension or solution, and usually into a lower \*soil horizon, after removal from above or from a lateral source.

**ilmeneite** Mineral,  $FeTiO_3$ ; sp. gr. 4.5–5.0; \*hardness 5–6; \*trigonal; black; black to brownish-red \*streak; \*sub-metallic \*lustre; crystals normally thick and \*tabular, but often \*massive and compact; no \*cleavage; magnetic; occurs as an \*accessory mineral in \*igneous rocks, e.g. \*gabbro and \*diorite, in \*quartz veins and \*pegmatites, in \*gneisses, in association with \*hematite and \*chalcopyrite, and because it is resistant to \*weathering

it occurs extensively in *\*alluvial* deposits with *\*magnetite*, *\*monazite*, and *\*rutile*. It is used as a source of iron and titanium. The name is derived from that of the Ilmen Mountains, Russia.

**IMA** International Mineralogical Association. See COMMISSION ON NEW MINERALS, NOMENCLATURE, AND CLASSIFICATION.

**IMAGE** See IMAGER FOR MAGNETOPAUSE-TO-AURORA EXPLORATION.

**image intensifying** See IMAGING.

**imaginary component (quadrature, out-of-phase component)** An electromagnetic field induces a secondary field in a conductor and the resultant vector of the two fields can be resolved into two components, one of which is the imaginary component, lagging  $\pi/2$  behind the *\*in-phase* component.

**Imager for Magnetopause-to-Aurora Exploration (IMAGE)** A mission of *\*NASA*'s Goddard Space Flight Center to study the global response of the Earth's *\*magnetosphere* to changes in the *\*solar wind*. The satellite was launched on 25 March 2000, from California, into a highly elliptical *\*polar orbit*.

**imaging** The instrumental recording and interpreting of various portions of the electromagnetic spectrum from planetary or *\*satellite* surfaces. This information is obtained mostly in digital form, either from spacecraft or telescopically. A common form is a photographic image in the visible portion of the spectrum. 'Multispectral imaging' consists of simultaneous recording, typically of four or more spectral bands ranging from the visible to infrared wavelengths. 'Image intensifying' is the enhancement of the image by computer processing. See REMOTE SENSING.

**imaging radar** A type of *\*radar* that constructs an image from the echoes reflected back to its antennae.

**imaging spectrometer** A *\*remote* sensing instrument which records an image in a large number of spectral *\*channels*.

**imbibition** The displacement of one fluid by another *\*immiscible* fluid in a porous medium. There are several biological examples and imbibition is

also important in oil recovery from *\*water-drive reservoirs* because a porous rock is able to imbibe both water and oil.

**Imbrian** 1. A *\*period* of the *\*Archaean*, dated at about 3850–3800 Ma ago (Harland et al., 1989). The name is no longer used. 2. Two epochs in the *\*lunar timescale*, the Lower Imbrian (3850–3800 Ma ago) and the Upper Imbrian (3800–3200 Ma ago).

**imbricate** With parts overlapping one another like tiles.

**imbricate structure (schuppen structure)** Fabric resulting from the stacking of rock fragments, particles, or tectonic units. Some pebble beds show an imbricate structure, with the pebbles leaning in the direction of the current. *See also BAR.*

**immersion objective method (oil immersion)** A technique commonly used in *\*reflected-light* microscopy, especially when high magnification and high resolution are required. A drop of immersion oil or water, with a *\*refractive index* of about 1.51, is placed on the polished surface of the mineral and the *\*objective* is lowered carefully into the liquid. The advantages of the method include better observations of colour differences, *\*birefractance*, and *\*anisotropy*.

**immiscible** *See MISCIBILITY.*

**immobilization** Conversion of a chemical compound from an inorganic to an organic form as a result of biological activity; the compound is thereby removed from the reservoir of compounds available to plant roots.

**IMP-8** *See INTERNATIONAL MONITORING PLATFORM.*

**impact cratering** The modification of the surface of a planet or satellite by collision with a *\*bolide*. On Earth, *\*tectonics* and *\*erosion* obliterate most signs of such events, but they are a major process in fashioning the surface of other solar-system bodies, most notably the Moon. An impacting bolide travels at up to 150 000 km/h. As it penetrates the surface its kinetic energy is instantly converted to heat, causing an explosion that vaporizes the object and the material it strikes and producing a crater much larger than the size of the bolide might suggest. An impacting *\*meteorite* about 3 m across and weighing 50 t will make a lunar crater 150 m in diameter.

**impactite** A rock type produced during the impact of a *\*meteorite* on to a planetary or *\*satellite* surface. *See also* SUEVITE.

**impactogen** *See* RHINE GRABEN.

**impeded dune** A *\*sand \*dune* with a shape that is strongly influenced by the effect of vegetation, a topographic barrier, or a highly localized source of sand.

**imperforate** Sometimes applied to those gastropod (*\*Gastropoda*) shells which do not possess an *\*umbilicus*. Although the term is much used, 'anomphalous' or 'non-umbilicate' are to be preferred since an umbilicus is not a perforation.

**impervious rock** Rock which will not permit oil, water, or gas to flow through it. *See also* PERMEABILITY.

**implosion** A sudden collapse into a zone of very low pressure. In marine *\*seismic* surveying, the discharge of air into water causes water to implode into the bubble space and the impact of the water colliding with itself creates a seismic signal, the *\*bubble* pulse.

**impulse response function** The specification of the effect of a *\*filter*. For example, if a spike (*\*Dirac* function) is put through a linear filter the output function will appear modified and spread out, this characteristic filter effect being the impulse response function. It is important in *\*convolution* and *\*deconvolution* in data processing.

**impunctate** *See* PUNCTATE.

**Inarticulata** (phylum *\*Brachiopoda*) A class of brachiopods, existing from the Lower *\*Cambrian* to the present day, in which the shell is calcareous, but its valves are not hinged by teeth and sockets and the *\*pedicle* is much reduced or absent. There are three orders.

**Inarticulate brachiopods** *See* INARTICULATA.

**inceptisols** An order of mineral soils in the US *\*soil taxonomy* that have one or more *\*soil horizons* in which mineral materials have been weathered or removed. Inceptisols are in the early stages of forming visible horizons, and are only beginning the development of a distinctive *\*soil profile*. The term embraces *\*brown earths*.

**incident angle** See ANGLE OF INCIDENCE.

**incipient hardground** See HARDGROUND.

**incised meander** See MEANDER.

**inclination** The angle between the horizontal and a magnetic vector. Conventionally, a vector with a magnetic north pole dipping below the horizontal is considered positive, and an upward vector is negative.

**inclined extinction** See OBLIQUE EXTINCTION.

**inclined fold** A *fold* in which the angle of *dip* of the *axial plane* is between 10° and 80°, and the highest and lowest points on the fold surfaces do not necessarily coincide with the *hinge* points. Compare OVERFOLD; UPRIGHT FOLD; RECUMBENT FOLD.

**inclinometer, geomagnetic** An instrument to measure the *inclination* of the *geomagnetic* field. It is usually a dip circle, but inclination may also be determined from separate measurements of the horizontal and vertical components of the geomagnetic field.

**included fragments** See PRINCIPLE OF INCLUDED FRAGMENTS.

**incompatible elements (hygromagmatophile elements)** Elements that, owing to their size, charge, or *valency* requirements, are difficult to substitute into the crystal structure of a rock-forming mineral (e.g. the boron *ion* is very small and the tungsten ion may have a +6 charge). This results in their being preferentially introduced into a *magma* on *partial* melting and less likely to crystallize out of it. During the crystallization of *igneous* rocks, incompatible elements (e.g. Sn, Li, Rb, Sr, and *rare-earth* elements) are often concentrated into *pegmatitic* or *hydrothermal* fluids. During the formation of the *Earth's* crust the incompatible elements have been transferred through magmatic processes to the crust from the *mantle*, which has consequently become depleted in these elements.

**incompetent** A relative rheological (see RHEOLOGY) term, referring to the ease with which a rock or layer of rock may be deformed. It is applied to materials which are less rigid than competent materials (see COMPETENCE) and tend to flow rather than fracture when deformed. Incompetence reflects

the inability of a material to transmit **\*compressive** stresses over large distances.

**incompressibility modulus** See BULK MODULUS.

**incongruent dissolution** Dissolution of a mineral with decomposition or reaction in the presence of a liquid, converting one solid **\*phase** into another, e.g. the conversion of orthoclase (see ALKALI FELDSPAR) to **\*kaolinite**:  $2\text{KAlSi}_3\text{O}_8 + 11\text{H}_2\text{O} \rightarrow \text{Al}_2\text{Si}_2\text{O}_5(\text{OH})_4 + 4\text{Si}(\text{OH})_4 + 2\text{K}^+ + 2\text{OH}^-$ .

**incongruent melting** Melting of a mineral with decomposition or reaction, such that it is replaced by a **\*melt** and a solid **\*phase** of different composition. For example, **\*orthoclase** melts incongruently to give **\*leucite** and a silica-rich liquid.

**inconsequent drainage (insequent drainage)** A **\*drainage pattern** which bears no apparent relationship to the underlying rock type or structure.

**incumbent replacement** An evolutionary mechanism, proposed in 1991 by M. L. Rosenzweig and R. D McCord, according to which a well-adapted species (the incumbent) becomes extinct, due to a chance combination of adverse factors, and its vacated niche is occupied by an invading species population.

**incus** From the Latin *incus* meaning 'anvil', a supplementary cloud feature comprising the flattened, anvil-like shape of the top of **\*cumulonimbus** cloud. See also CLOUD CLASSIFICATION.

**INDEX** See REIMI.

**index ellipsoid** See INDICATRIX.

**index fossil (index species, zone fossil)** **\*Fossil** whose presence is chosen to denote the **\*zone** in which it occurs and after which the zone is named. Index fossils are selected for their distinctiveness and/or abundance. To be of use in **\*biostratigraphy**, ideally an index fossil should have a narrow range in time (i.e. to have undergone rapid evolutionary change) but have had a wide geographical distribution. **\*Trilobites** (**\*Cambrian**); **\*graptolites** (**\*Ordovician** and **\*Silurian**); **\*ammonites** (**\*Jurassic**); and **\*foraminifera** (**\*Cretaceous** and **\*Cenozoic**) are among the most notable index fossils.

**index mineral** A diagnostic \*mineral in a \*regional metamorphic terrain whose first appearance, going in the direction of increasing \*metamorphic grade through the metamorphic sequence, marks the outer limit of a \*metamorphic zone. A line marking the first appearance of an index mineral is termed an 'isograd', and represents a line of constant metamorphic grade. To indicate the regional distribution of metamorphic grade the first appearance of index minerals in rocks of \*pelitic composition (i.e. \*shales) are usually mapped out in the field, as this type of rock is extremely sensitive mineralogically to changes in metamorphic grade.

**index species** See INDEX FOSSIL.

**Indian jade** See AVENTURINE

**Indian National Satellite-3 (INSAT-3)** A series of seven multipurpose communications and meteorological spacecraft operated by the Indian Space Research Organization (ISRO). INSAT-3B was launched on 21 March 2000, -3C on 3 February 2002, Kalpana-1 (MetSat-1) on 12 September 2002, -3A on 9 April 2003, -3E on 28 September 2003, and -3D on 25 July 2013, all from Kourou, French Guiana. INSAT-3DR, a repeat mission similar to -3D, was launched on 8 September 2016, from Satish Dhawan Space Centre, India.

**Indian Ocean** One of the world's major oceans, lying between Africa, India, and Australia. It has a surface area of 77 million km<sup>2</sup> and an average depth of 3872 m. The ocean receives a great deal of sediment from three of the world's major rivers (the Ganga, Indus, and Brahmaputra).

**Indian Remote Sensing (IRS)** A programme by the Indian Space Research Organization (ISRO) that began with the launch of satellite IRS-1A from Baikonur, Kazakhstan, on 17 March 1988, into a Sun-synchronous \*polar orbit. The IRS series comprises satellites that help map, monitor, and manage natural resources. There is now a constellation of eleven IRS operational satellites, the largest constellation for civilian use.



<https://www.isro.gov.in/saga-of-indian-remote-sensing-satellite-system>

- The Saga of Indian Remote Sensing Satellite System.

**Indian summer** See SINGULARITY.

**indicated reserve** See RESERVE.

**indicator** See ERRATIC.

**indicatrix (optical indicatrix, index ellipsoid)** An ellipsoid which represents geometrically the different \*vibration directions in a \*mineral and illustrates conceptually the optical features of a \*crystal. The origin is a point lying at the centre of the ellipsoid and the axes of the ellipsoid are proportional in length to the \*refractive indices of beams of light vibrating at right angles along them. These axes are commonly termed X, Y, and Z, or  $n_\alpha$ ,  $n_\beta$ , and  $n_\gamma$ , for \*orthorhombic, \*monoclinic, or \*triclinic minerals. \*Tetragonal, \*hexagonal, and \*trigonal minerals are represented by an indicatrix with one principal section circular; and the \*cubic minerals are represented by an indicatrix (the isotropic indicatrix) which is a sphere with all axes equal. Measurements of the elliptical plane sections give optical properties which aid specific identification of a mineral. See OPTIC AXIS.

**Indo-Australian Plate** One of the present-day major lithospheric \*plates, which is having new material added to its south and south-west along the \*Carlsberg Ridge and the south-east Indian Rise, but its other margins are the \*collision zone of the Himalayan \*orogenic belt, \*subduction zones (e.g. in the E. Indies), or \*transform faults (e.g. the Alpine Fault in New Zealand). It is thought this plate may now be breaking into two separate plates along the line of the 90° E Ridge.

**Induan** A \*stage of the \*Triassic period from 251–249.7 Ma ago, preceded by the \*Changhsingian and followed by the \*Olenekian.

**induced polarization (IP, induced potential, over-voltage, interfacial polarization)** An exploration method which uses either the decay of an excitation voltage (\*time-domain method) or variations in the Earth's \*resistivity at two different but low frequencies (\*frequency-domain method). A variety of different \*electrode configurations can be used. See also CHARGEABILITY.

**induced potential** See INDUCED POLARIZATION.

**induced pulsed transient** See INPUT.

**induction** The creation of a voltage by changing the magnetic flux such that the amount of voltage induced is directly proportional to the rate of change of the magnetic flux according to Faraday's or Neumann's law (*see also LENZ'S LAW*). In applied geophysics, induction is a fundamental process in *\*electromagnetic* (EM) prospecting; a primary EM field is used to induce a secondary field in any subsurface conductors and the resultant of the two fields is measured. The strength of the secondary field, which is a direct function of the electrical *\*conductivity* of the ground, can then be determined.

**induction log** A *\*well-logging* system which uses *\*electromagnetic* *\*induction* methods to measure electrical *\*conductivity* and *\*resistivity* in the adjacent *\*formations*.

**induction sonde** A variety of *\*electromagnetic* (EM) instrument for producing and measuring eddy currents within the rocks surrounding a *\*borehole*. The strength of the eddy currents is proportional to the conductivity (1/resistivity) of the surrounding rocks. Penetration is usually greater than by *\*laterolog* sondes. The main application of the log in hydrocarbon exploration and production is in the estimation of water saturation. Interpretation is affected by the *\*Delaware* effect.

**inductively coupled plasma emission spectrometry** Technique for the chemical analysis of *\*trace* elements in a wide variety of materials, using a source that depends on the interaction between the magnetic field of an oscillating radio-frequency current and the charged species present in the *\*plasma*. Argon gas is passed through the field producing a toroidal (doughnut-shaped) plasma at temperatures of over 5000 °C. The sample is introduced as an *\*aerosol* into the gas and is vaporized and atomized in the plasma. The source has a *\*monochromator* (a means of selecting a single frequency) and a readout system. It has low limits of detection for most elements, and is a very rapid technique.

**induration** Process of forming a *\*soil horizon* or *\*hardpan* that has a high bulk density and is hard or brittle. *\*Cementing* materials may be present and responsible for the induration.

**industrial mineral** Any earth material of economic importance, excluding metal *\*ores* and fuels; e.g. *\*barite*, *\*fluorite*, and china clay (*\*kaolin*).

**inertial reference frame** In *\*geodesy*, a reference frame that allows for the rotation of celestial bodies, but assumes uniform motion. It is used to express positional coordinates derived from astronomical or satellite measurements.

**inertinite** See COAL-MACERAL GROUP.

**infaunal** Applied to benthic organisms that dig into the sea bed or construct tubes or burrows. They are most common in the subtidal and deeper zones (i.e. the area seaward of the low-water mark). Compare EPIFAUNAL.

**inferred reserve** See RESERVE.

**inferred tree** A *\*phylogenetic tree* based on empirical data pertaining to extant taxa.

**infiltration** Downward entry of water into soil.

**infiltration capacity** The maximum rate at which soils and rocks can absorb rainfall. The infiltration capacity tends to decrease as the soil-moisture content of the surface layers increases. It also depends upon such factors as grain size and vegetation cover.

**infinitesimal strain** The extensions and deflections that are required to deform an infinitesimally small reference cube into a *\*strain parallelepiped*. The concept is of use when considering the amount of strain absorbed by a material during each instant of its progressive deformation.

**inflexion point** 1. The point of no curvature in a *\*fold*. 2. The point at which the curvature of a fold changes sense between one closure and the next.

**inflow band (flumen)** See CLOUD CLASSIFICATION.

**influent stream** See GAINING STREAM.

**influx** An inflowing of *\*sediment*, fluid, mineralizing solution, or other material.

**informal** The informal naming of a *\*stratigraphic unit* occurs when (a) the unit-term is referred to as an ordinary noun, and not in the context of a proper name, e.g. 'the geologic periods'; and (b) when a unit-term (such as *\*zone*, or *\*formation*) is referred to without having been specifically

established in classification, e.g. 'a sandstone formation', 'a mineralized zone'. Terms used informally are not capitalized. *Compare* [FORMAL](#). *See* [STRATIGRAPHIC NOMENCLATURE](#).

**infrared radiation** \*Electromagnetic radiation which has a \*wavelength between 0.7  $\mu\text{m}$  and 100  $\mu\text{m}$ . *See also* [NEAR-INFRARED](#); [MID-INFRARED](#); [REFLECTED INFRARED](#); [THERMAL INFRARED](#).

**infrared remote sensing** Method of distinguishing various types of vegetation, rocks, etc. formerly using either monochrome or coloured infrared film which can be used in conventional cameras, but nowadays using an infrared sensor. Potentially, \*aerial photography using an infrared camera may be of benefit in preparing \*geologic maps. Longer-wavelength infrared can discriminate most rocks, shorter wavelengths can reveal iron oxide, etc.; alteration effects around certain mineral deposits, e.g. \*porphyry coppers with their attendant \*clay minerals, can also be distinguished, and the technique allows a consideration of plant species affected by the nature of the \*soil and rock substrate.

**ingrown meander** *See* [MEANDER](#).

**inhomogeneity (heterogeneity)** Irregular spatial variability in physical properties. *Compare* [ANISOTROPY](#).

**inhomogeneous strain** \*Strain, distributed unevenly throughout a deformed body, in which straight lines and parallel lines in the undeformed material become curved and non-parallel on deformation. The mathematical theory describing the geometry of inhomogeneous strain is very complex and it is usual to divide this strain into smaller components of \*homogeneous strain.

**initial levée** *See* [LAVA LEVÉE](#).

**initial strontium ratio (common strontium)** Strontium has four naturally occurring \*isotopes:  $^{88}\text{Sr}$ ;  $^{87}\text{Sr}$ ;  $^{86}\text{Sr}$ ; and  $^{84}\text{Sr}$ . Of these,  $^{87}\text{Sr}$  is perhaps the most important because it is formed by the natural radioactive decay of the rubidium isotope  $^{87}\text{Rb}$ ; this decay provides the basis for one of the most important geochronological methods. At its simplest, the initial strontium ratio (common strontium) of a rock is the ratio between the radioactively produced isotope  $^{87}\text{Sr}$  and the 'ordinary', non-radiogenic isotope  $^{86}\text{Sr}$  at the

time when the rock crystallized. In a hypothetical rock containing no rubidium this ratio would remain unchanged for ever, but most rocks contain some rubidium; thus radioactive decay constantly increases the amount of  $^{87}\text{Sr}$ , and the ratio of  $^{87}\text{Sr}$  to  $^{86}\text{Sr}$  constantly increases, at a rate proportional to the amount of rubidium in the rock. The initial strontium ratio of a rock is determined by measuring the present-day  $^{87}\text{Sr}:$  $^{86}\text{Sr}$  ratio in several of its constituent minerals. At the time when it first crystallized each mineral in the rock would have had the same  $^{87}\text{Sr}:$  $^{86}\text{Sr}$  ratio, but each mineral contains a different amount of rubidium, so after any given time its  $^{87}\text{Sr}:$  $^{86}\text{Sr}$  ratio will have increased away from the initial value by an amount exactly determined by the relative proportions of rubidium and strontium in it. Therefore, both the initial strontium ratio and the age of the rock can easily be found by plotting the measured present-day ratios of the constituent minerals on an **\*isochron** diagram. In **\*petrology**, initial strontium ratios are important because they provide information that otherwise would not be available on the chemical composition and ages of the source regions of **\*igneous** rocks. For example, an igneous rock that has a very young radiometric age but a very high initial strontium ratio must have been derived from a source rich in  $^{87}\text{Sr}$ . Such a source must have been rich in rubidium and old enough for the  $^{87}\text{Sr}$  to have accumulated by the radioactive decay of  $^{87}\text{Rb}$ . Young **\*granites** in continental **\*collision zones** (e.g. the Alps) have extremely high (up to 0.8) initial strontium ratios and formed by the melting of old crustal **\*gneisses**. Granites formed by melting of the same rocks at different times can be distinguished immediately by their initial strontium ratios since the source rocks would have accumulated different amounts of  $^{87}\text{Sr}$  when melting occurred. Granites formed in **\*island** arcs are derived from young **\*mantle** materials and have strikingly lower (0.704–0.706) initial strontium ratios. Similar arguments can be applied to the source regions of **\*basalts** in the mantle, though the differences between present and initial  $^{87}\text{Sr}:$  $^{86}\text{Sr}$  ratios are very much smaller because the mantle is poor in rubidium; typical ocean **\*ridge** basalts have initial strontium ratios close to 0.703. At the time of its formation, 4.6 billion years BP, the bulk Earth ratio is thought to have been 0.699. *See also* **ISOCHRON**.

**inland sea** Extensive body of water that is largely or wholly surrounded by land. Any connection to the open ocean is restricted to one or a few narrow

sea passages. Examples of such areas are the Baltic and the Mediterranean Seas.

**inlier** Structure where older rocks are surrounded completely by newer rocks. It may result from *\*faulting* or *\*folding* followed by *\*erosion*.

**inner planet** See TERRESTRIAL PLANET.

**Innovative Technology Demonstration Experiment** See REIMI.

**inosilicate (chain silicate, band silicate)** Applied to the structure of silicate *\*minerals* where  $\text{SiO}_4$  *\*tetrahedra* are linked together into chains by sharing oxygens. Two important groups of rock-forming minerals are included: *\*pyroxenes*, where two of the four oxygens are shared to give a 'single chain' structure with the ratio Si:O = 1:3; and *\*amphiboles*, where half the  $\text{SiO}_4$  tetrahedra share two oxygens and the other half share three oxygens, to give a 'double chain' or 'band' structure and a ratio Si:O = 4:11.

**in-phase component (real component)** An electromagnetic field induces a secondary field in a conductor and the resultant vector of the two fields can be resolved into two components. One of these is the in-phase (real) component, in phase with the primary field, the other is the *\*imaginary* component (quadrature).

**INPUT (INDuced PULsed Transient)** An airborne electromagnetic surveying system that comprises a large coil transmitter looped around the aircraft beneath the nose and tail and around each wing-tip. A *\*bird* is trailed behind the aircraft to detect the decaying secondary field at times when the primary field is between source pulses.

**InSAR** See INTERFEROMETRIC SYNTHETIC APERTURE RADAR.

**INSAT-3** See INDIAN NATIONAL SATELLITE-3.

**Insecta (subphylum Hexapoda, insects)** (phylum *\*Arthropoda*) Class of arthropods that have three pairs of legs and, usually, two pairs of wings borne on the thorax. Typically, there is a single pair of antennae and one pair of compound eyes. Gas exchange takes place through a tracheal system and the gonoducts open at the posterior end of the body. The oldest fossil insects occur in *\*Devonian* rocks, and the first winged representatives are known from *\*Carboniferous* rocks. Dragonflies and beetles were

established before the end of the **\*Palaeozoic**; social varieties such as ants and wasps are present in **\*Cretaceous** sediments. The evolution of the flowering plants had a marked influence on insect development, so that many new forms appeared in the Cretaceous. More than 750 000 extant species of insects have been described. Insects are now believed to be descended from a crustacean ancestor, and the **\*Hexapoda** and **\*Crustacea** are grouped in the **\*clade** Pancrustacea.

**inselberg** Steep-sided, isolated hill that stands above adjacent nearly flat plains. It may have a narrow **\*pediment** at its base. Locally, flared or steepened margins occur. It is best developed under a savannah climate. *See also* BORNHARDT.

**insequent drainage** *See* INCONSEQUENT DRAINAGE.

**InSight (Interior Exploration using Seismic Investigations, Geodesy and Heat Transport)** A **\*NASA** Discovery Program mission to Mars that on 26 November 2018 placed a lander on the surface with a **\*seismometer** to monitor Marsquakes, a thermal probe 5 m long to measure the subsurface thermal gradient, and the spacecraft's communication system to measure the planet's rotation. The overall aim was to study the internal structure of Mars to provide information on the way rocky planets form.



[https://www.nasa.gov/mission\\_pages/insight/main/index.html](https://www.nasa.gov/mission_pages/insight/main/index.html)

- InSight Mars Lander.

**insolation** The amount of incoming solar radiation that is received over a unit area of the Earth's surface. Solar energy received over the planet's surface varies according to season, latitude, transparency of the atmosphere, and aspect or ground slope. On average, equatorial areas receive approximately 2.4 times as much insolation as polar areas.

**insolation weathering** *See* THERMOCLASTIC.

**instability** Atmospheric condition in which displaced air continues to rise once it has begun to do so. This occurs when rising air cools at an adiabatic lapse rate that is less than the **\*environmental lapse rate**, so a rising air parcel remains warmer than the surrounding air at every height, thus

retaining positive buoyancy. See also [CONDITIONAL INSTABILITY](#); [POTENTIAL INSTABILITY](#); [STABILITY](#).

**instantaneous field of view** In *\*remote* sensing, the angle through which a detector is receiving *\*electromagnetic* radiation. It is often expressed as a function of the ground area visible at any one time, which is dependent on height and the angle of radiation reception.

**instantaneous rotation** The relative rotation of a body about its *\*pole of rotation* at a given instant.

**instantaneous strength (short-term strength)** The amount of *\*stress* at which a rock fails in a testing machine when the stress continues for a short time. If the rock is exposed to a lower stress for a longer period it will undergo *\*creep* and fail at a stress much lower than the instantaneous strength.

**Institute of Space and Astronautical Science (ISAS)** The Japanese national research institute dedicated to space and astronautical science under the auspices of the Ministry of Education, Science, and Culture. It was founded in 1981 and in 2003 was merged with other agencies to form the *\*Japan Aerospace Exploration Agency*.

**insula** (*pl.* insulae) On the surface of *\*Titan*, an isolated area of land (i.e. an island) or group of such areas surrounded by liquid.

**intact rock strength** The strength of a rock excluding weaknesses due to *\*fractures* and *\*joints*. It is measured by the rebound value in a *\*Schmidt hammer test* and by its response to a blow from an ordinary hammer. A very strong rock such as *\*quartzite*, *\*dolerite*, or *\*basalt* will break only under severe blows from the hammer and has a rebound value of 100–60. A hammer blow will break a hand-held specimen of a strong rock (60–50) such as *\*marble*, *\*granite*, or *\*gneiss*. A firm blow will make a shallow indentation on the surface of a moderately strong (50–40) rock such as *\*sandstone* or *\*slate* and a deep indentation on a weak (40–35) rock such as *\*coal* or *\*schist*. A very weak (35–10) rock such as *\*chalk* or *\*halite* can be cut with a knife and a sharp hammer blow will make it crumble.

**Integrated Ocean Drilling Programme** See [INTERNATIONAL PROGRAMME OF OCEAN DRILLING](#).

**intensification** In meteorology, the increase of pressure gradient around a pressure system. *See also* WEAKENING.

**intensity** In *\*remote* sensing, the energy reflected or emitted by a surface.


**intensity (earthquake)** *See* EARTHQUAKE MAGNITUDE; JAPANESE METEOROLOGICAL AGENCY SEISMIC INTENSITY SCALE; MERCALLI SCALE; MOMENT MAGNITUDE SCALE; RICHTER SCALE.

**intensity-hue-saturation processing** In *\*remote* sensing, a form of *\*contrast stretching* in which the visibility of *\*pixel colour* is enhanced. Usually the saturation values of pixels are stretched to fill parameter space.

**inter-** From the Latin *inter* meaning 'between', a prefix meaning 'between'.

**interambulacral** *See* INTERAMBULACRUM.

**interambulacrum (interamb, adj. interambulacral)** In *\*Echinodermata*, that area of the body surface lying between ambulacra. *See* AMBULACRUM.



Interambulacrum

**inter-arc basin** A type of *\*back-arc* basin which is floored by *\*oceanic* crust. The main sediments are turbiditic volcanoclastics derived from the volcanic arc. *See also* INTER-ARC TROUGH.

**inter-arc trough** A *\*fore-arc* basin developed between an outer, non-volcanic arc and a volcanic arc. *Compare* INTER-ARC BASIN.

**interarea** In some brachiopods (*\*Brachiopoda*), a weight-bearing shelf of shell material, typically triangular, that stabilizes the *\*valves*.

**interbiohorizon zone** See INTERVAL ZONE.

**interception** 1. The capture of rainwater by vegetation from which the water evaporates and is thus prevented from reaching the *\*water-table* and contributing to *\*surface runoff*, *\*soil moisture*, or *\*groundwater* recharge. 2. The abstraction of groundwater part of the way along its flow path, where otherwise the water might be lost, e.g. as coastal spring discharges.

**intercept ratio** See AXIAL RATIO; PARAMETER.

**intercept time** The arrival time of a seismic wave, determined from the intercept of the extrapolation of the refracted straight-line segment of a *\*time–distance* graph at zero *\*offset*.

**intercrystalline boundary** See VOIDS.

**intercrystalline porosity** See CHOQUETTE AND PRAY CLASSIFICATION.

**interdigitating (interfingering, interlocking)** Applied to a *\*facies* boundary where the line of lithological change between one *\*sedimentary rock* type and the next laterally adjacent rock type is itself broken down into a series of wedge-shaped zig-zags or tongues. Where such interfingering occurs it is a record of the fluctuations in the local depositional environment, implying that both types of *\*sediment* were being laid down at the same time.

**interface** 1. A device which connects two machines and allows them to communicate with one another. The term is used most commonly in respect of equipment linked to a computer. 2. The boundary between two substances that have different properties. See also HORIZON.

**interface-controlled growth** *\*Crystal* growth in a *\*melt* or *\*solution* where the rate of crystal growth is controlled by the transport of material across the crystal–liquid interface. This type of growth tends to characterize large *\*supercooling* regimes in one-component systems.

**interfacial angle** In *\*crystallography*, the angle subtended by the normals to two *\*crystal faces*. It is not the external angle observed or the internal

angle between them; it is, however, 180° minus the internal angle. A *\*goniometer* is used to measure interfacial angles.

**interfacial polarization** See *INDUCED POLARIZATION*.

**interference** The combination of waves. Constructive interference occurs when peaks add to peaks; destructive interference occurs when a peak coincides with and cancels out a trough.

**interference colour chart** See *MICHEL–LÉVY CHART*.

**interference colours (polarization colours)** In *\*mineral* optics, the colours produced when the *\*analyser* is inserted on a thin-section microscope. They are produced as a result of *\*birefringence* (double refraction) whereby one ray of light is retarded relative to the other. The different degrees of retardation give different interference colours. These colours are used in a number of ways as an aid to identification.

**interference figure** In *\*crossed-polar* mineral optics, the faintly coloured rings and dark curves produced as a result of retardation when convergent polarized light passes through *\*anisotropic* (i.e. *\*double refracting*) *\*minerals*. The black curves or crosses are called ‘isogyres’. There are two kinds of interference figures: uniaxial and biaxial. When centred, the uniaxial interference figure resembles a black cross with coloured concentric circles; the biaxial interference figure is more complex, with two curved isogyres and coloured elliptical bands. The coloured bands represent zones of equal retardation and tests using accessory plates give optical properties which are characteristic of individual minerals.

**interference pattern** A two-dimensional *\*outcrop* pattern resulting from the superimposition of two or more sets of *\*folds* of different generations. The form of the patterns seen depends on the relative *\*attitudes* of the superimposed folds; J. G. Ramsay (1967) recognized four basic types: redundant superposition (in which later folding has not altered the original pattern); dome and basin (egg box); dome–crescent–mushroom (see *CRESCENT-AND-MUSHROOM*); and convergent–divergent (double zig-zag).

**interferogram** A diagram produced by an *\*interferometer*. It can reveal earth movements due to tectonic deformation, volcanic activity, subsidence, ice flow, etc. and also produce digital elevation maps (DEM).

**interferometer** An instrument used in *\*remote* sensing that forms and uses interference patterns in radiation to measure the wavelengths of that radiation.

**interferometric synthetic aperture radar (InSAR, IfSAR)** A *\*radar* technique in which two or more *\*synthetic-aperture radar* images are used to generate maps showing surface deformation. The radar equipment is carried on an orbiting satellite or aircraft, and the technique is used in *\*geodesy* and *\*remote sensing*.

**interfingering** See INTERDIGITATING.

**interflow (throughflow)** The lateral movement of water through the upper *\*soil horizons*, normally during or following significant precipitation events. Shallow *\*groundwater* or interflow may emerge at the surface at the bottom of slopes and flow across the ground surface for a time. This is known as 'return flow'.

**interfluve** The elevated part of the landscape that extends between two adjacent valleys. It is normally seen as lying above the steeper slopes of each valley side.

**interglacial** Period of warmer climate that separates two *\*glacial* periods. Mid-latitude interglacials show a characteristic sequence of vegetation change. *\*Pollen* of heathy *\*tundra* is replaced in the pollen record by abundant herbaceous pollen, which in turn is replaced by that of *\*boreal* and subsequently deciduous forest, including pollen of thermophilous (warmth-loving) species, e.g. Tilia (lime). From this peak the sequence reverses as the trend to colder conditions predominates.

**intergrade** Soil or *\*soil horizon* that has the properties of two genetically different soils or horizons. An intergrade can be regarded as transitional between two distinctive soils or horizons.

**intergranular** Applied to an *\*igneous \*texture*, especially well developed in *\*basalts*, in which the wedge-shaped spaces between a meshwork of lath-shaped crystals, such as *\*plagioclase*, are filled with granules of other minerals.

**intergranular displacement** The displacement of individual grains within a rock that is undergoing *\*plastic* deformation. Such grain movements

induce permanent **\*strain** in the rock.

**intergranular pores** See VOIDS.

**Interior Exploration using Seismic Investigations, Geodesy and Heat Transport** See INSIGHT.

**inter-limb angle** See FOLD ANGLE.

**interlocking** See INTERDIGITATING.

**intermediate rock** **\*Igneous** rock whose chemical composition lies between those of **\*basic** and **\*acidic rocks**, e.g. **\*andesite**. The limits are not fixed rigidly and a number of schemes exist that are based on modal mineralogy and the whole rock chemistry (see MODAL ANALYSIS). Compare ACID ROCK; BASIC ROCK. See also ALKALINE ROCK.

**intermittent stream** A stream which ceases to flow in very dry periods. Such streams tend to have permeable beds and during periods of flow water leaking through their beds is added to the local **\*groundwater**. The chalk bournes of southern England provide typical examples of intermittent streams. See also LOSING STREAM.

**intermontane** 1. Between mountains or mountain ranges. 2. Applied to basins which are being infilled by sediment eroded from surrounding mountains.

**internal angle of friction** See ANGLE OF SHEARING RESISTANCE.

**internal mould** See FOSSILIZATION.

**internal node** Within a **\*phylogenetic tree**, a point where two branches join, representing an ancestral species or gene.

**internal reflection (IR)** The reflection of light off **\*cleavage** or **\*fracture** planes just below the surface of certain slightly translucent **\*ore minerals** (e.g. **\*cassiterite** and **\*sphalerite**) and seen as a faint glow when the mineral is viewed under **\*crossed** polars in reflected-light or ore microscopy.

**internal standard** In many instrumental analytical techniques, the mixing of accurately known amounts of a convenient element or known compound with the sample being considered. For example, in emission spectrometry,

an internal standard is employed in order to relate the intensities of the line spectra to the concentration.

**internal wave** Wave that forms within a water mass at the boundary of two water layers that have different densities. The boundary may be abrupt or gradual, and the slow-moving waves can be detected only by instrumental observations of temperature or *\*salinity*, and by acoustic scattering.

**International Association for Engineering Geology and the Environment (IAEG)** A scientific society founded in 1964 that aims to advance and encourage engineering geology. It is affiliated to the *\*International Union of Geological Sciences* and has more than 5200 members in fifty-nine national groups.



<https://uia.org/s/or/en/1100003651>

- IAEG Homepage.

**International Cometary Explorer** See INTERNATIONAL SUN—EARTH EXPLORER.

**International Federation of Digital Seismic Networks (FDSN)** A worldwide organization that coordinates the work of groups that install and maintain *\*seismographs* either globally or within their own territories. Membership is open to any group operating more than one broadband station and the FDSN issues network codes that ensure data streams are unique.



<http://www.fdsn.org/>

- The International Federation of Digital Seismic Networks.

**International Geomagnetic Reference Field (IGRF)** The best mathematical fit to the observed *\*geomagnetic* field at any specific time. It is usually evaluated annually.

**International Gravity Formula** Formula used to determine the *\*gravitational* acceleration at a given latitude ( $g_\varphi$ ) for a model of the Earth that comprises a rotating, oblate spheroid.  $g_\varphi = g_0(1 + \alpha \sin^2 \varphi + \beta \sin^2 \varphi)$ ,

where  $g_0$  is the value at the equator, of 978.0318 gals,  $\varphi$  is the latitude, and the constants  $\alpha$  and  $\beta$  are 0.0053024 and  $-0.0000058$ .

**International Gravity Standardization Network** A network of stations where the absolute \*gravitational acceleration has been established. This enables \*gravimeters that are differential instruments to be calibrated, so that they provide absolute values. Such stations are commonly at or near airports.

**International Mineralogical Association** See COMMISSION ON NEW MINERALS, NOMENCLATURE, AND CLASSIFICATION.

**International Monitoring Platform (IMP-8)** The last in a series of ten \*NASA-Goddard Space Flight Center missions to monitor the \*solar wind and measure the \*plasma field environment of the Earth's \*magnetosheath and the magnetotail. It was launched on 26 October 1973, from Cape Canaveral, into a geocentric elliptical orbit. Decommissioned in 2001, it was the longest-lived geocentric space physics mission NASA had ever flown.

**International Polar Year (IPY)** A programme of 220 science and outreach projects aimed at determining the environmental status of polar regions, improving projections of future changes affecting them, and to advance understanding of them. The IPY also aimed to educate and involve the public and to train the next generation of engineers and scientists. The IPY was conducted under the auspices of the International Council for Science (ICSU) and the \*World Meteorological Organization (WMO) and began on 1 March 2007.



<http://www.ipy.org/>

- Explains and describes the IPY.

**International Programme of Ocean Drilling (IPOD)** An international project that evolved from the \*Deep Sea Drilling Programme, and was supported financially by the USA, USSR, West Germany, and France. The project has involved scientists from many countries, and the drilling and analysis of many deep \*boreholes in the deep sea and \*continental shelf

areas. In 2004 the Programme was renamed the Integrated Ocean Drilling Programme.

**International Research Institutions for Seismology (IRIS)** A US university research consortium that is dedicated to exploring the interior of the Earth by collecting and distributing seismographic data (*see* SEISMOGRAPH). IRIS is supported by the National Science Foundation, other federal agencies, universities, and private foundations.

**International Sun–Earth Explorer (ISEE)** A group of three spacecraft comprising a \*NASA and \*ESA mission to study the relationship between Earth and the Sun, the \*solar wind, and \*cosmic rays. ISEE-1 and ISEE-2 were launched on 22 October 1977 and ISEE-3 on 12 August 1978. The US contributed ISEE-1 and ISEE-3, and ISEE-2 was built and managed by ESA. In 1982, ISEE-3 was removed from its orbit about one of the \*Lagrangian libration points, eventually into a heliocentric orbit ahead of Earth that would intersect the comet \*Giacobini–Zinner, at which point the spacecraft was renamed International Cometary Explorer (ICE). It traversed the plasma tail of the comet in September 1985. ISEE-1 and ISEE-2 reentered the Earth’s atmosphere on 26 September 1987.



<http://heasarc.gsfc.nasa.gov/docs/heasarc/missions/isee3.html>

- A joint ESA and NASA mission to study the Earth’s magnetosphere.

**International Union of Geological Sciences (IUGS)** An organization that encourages international cooperation in geological sciences. It was founded in 1961, is based in Paris, and currently represents 121 Adhering Organisations, each from a different country or region.

**interparticle porosity** *See* CHOQUETTE AND PRAY CLASSIFICATION.

**interpenetrant twin (penetration twin)** A twinned \*crystal where parts of the twin appear to intertwine to give an irregular or indefinable contact surface. It is a special kind of \*contact twin, e.g. \*quartz Dauphiné twins and interpenetrant twins of \*fluorite.

**inter-record gap** *See* SEISMIC GAP.

**intersection cleavage** A *\*cleavage* which crosses another planar feature and consequently forms an intersection *\*lineation* with that feature.

**intersection lineation** See DIRECTIONAL FABRIC; INTERSECTION CLEAVAGE; LINEATION.

**interseismic period** See SUBDUCTION EARTHQUAKE CYCLE.

**intersertal** Applied to an *\*igneous \*texture*, especially well developed in *\*basalts*, in which the wedge-shaped spaces between a meshwork of lath-shaped crystals, such as *\*plagioclase*, are filled with *\*glass*.

**interstade (interstadial)** Phase of warmer climate within a glacial period, but of shorter duration (and thought to be less warm) than an *\*interglacial*. Warmth-demanding (thermophilous) species, e.g. *Tilia* (lime), are not represented in the *\*pollen* record, which shows *\*Boreal* affinities. The absence of thermophilous species may, however, be as much a consequence of the shorter time-span of an interstade as of the lack of warmth.

**interstadial** See INTERSTADE.

**Interstellar Boundary Explorer (IBEX)** A *\*NASA \*microsatellite* mission to investigate the global interaction between the *\*solar wind* and the interstellar medium. It was launched on 19 October 2008, from Kwajalein Atoll, Marshall Islands, into a highly elliptical orbit.

**interstellar clouds** Unusually dense patches of gas and dust, usually about 10 000 times more massive than the *\*Sun*, from which stars are believed to form. The clouds break up into many smaller, rotating fragments, which may become stars.


**interstellar medium** Substance, predominantly hydrogen (with some calcium, sodium, potassium, *\*hydrocarbons*, and cyanogen) found in the space between the stars, mainly in the plane of the Milky Way.

**interstitial** Pertaining to the spaces (interstices) between sedimentary particles.

**intertidal zone** Area between mean high-water level and mean low-water level in a coastal region. See LITTORAL ZONE.

**intertropical confluence** Alternative term for the **\*intertropical convergence zone** (ITCZ), preferred by some purists because of the discontinuous occurrence of convergence within it.

**intertropical convergence zone (ITCZ)** Low-latitude zone of convergence at the boundary between north-easterly and south-easterly **\*trade winds**. Convergence causes air to rise (see **HADLEY CELL**), producing a region of low pressure, the **\*equatorial trough**, and resulting in strong convection and heavy precipitation, mainly in the ocean sectors. The ITCZ moves latitudinally with the seasons. See **INTERTROPICAL CONFLUENCE**.



**Intertropical convergence zone**

**intertropical front** The name formerly given to the **\*intertropical convergence zone**.

**interval** The time elapsing between two geologic events. See also **POLARITY INTERVAL**.

**intervallum** The space between the inner and outer walls of **\*Archaeocyatha**.

**interval time** The difference in **\*two-way** travel time between two reflection events on a **\*seismic section**.

**interval velocity ( $V_{\text{int}}$ )** Seismic velocity over a depth interval  $z$ . If the rock type is uniform through that depth interval, then  $V_{\text{int}}$  is equal to the **\*formation velocity**. If the depth interval covers a number of rock beds, then the interval is equal to the **\*average** velocity ( $V$ ) calculated over the distance  $z$ . If  $z_i$  is the thickness of the  $i$ th interval and  $t_i$  is the one-way travel time through it, then  $V_{\text{int}} = z_i/t_i$ . A specific form of  $V_{\text{int}}$  is given by the **\*Dix**

formula, where the interval is defined in terms of the two-way travel time rather than by a discrete difference in depth.

**interval zone (interbiohorizon zone, biostratigraphic interval-zone)** The unit of *\*strata* lying between the top boundary of a distinctive lower *\*biostratigraphic unit* and the base of an equally distinctive, but different, upper biostratigraphic unit. An interval-zone may or may not contain *\*fossils*.

**intortus** Twisted or entangled, the name of a variety of *\*cirrus* cloud. See CLOUD CLASSIFICATION.

**intra-** From the Latin *intra* meaning 'inside', a prefix meaning 'within' or 'on the inside'. 1. Prefix used in the *\*Folk* classification to specify a *\*limestone* dominated by *\*intraclasts*. 2. General prefix for a process or object found or developed within the setting in question, e.g. 'intraformational conglomerate' is a conglomerate derived from within the *\*formation* of deposition.

**intraclast** A *\*carbonate* fragment of lithified, or partly lithified *\*sediment*, derived from the erosion of nearby sediment and redeposited within the same area (*compare* EXTRACLAST). Such reworked fragments are often amorphous and structureless. Carbonate lumps, formed by the complete micritization (*see* MICRITE) of shell fragments are indistinguishable from true intraclasts. They are, of necessity, classified with intraclasts, although clearly they are not true intraclasts.

**intrafolial fold** A thinned and modified, tight to *\*isoclinal* fold (*see also* FOLD) which commonly shows a fold *\*hinge* with only vestiges of detached limbs (*see* FOLD LIMB). Such folds are the result of intense deformation which causes *\*bedding* to be transposed along zones of shear and *\*solution*. *See also* SHEAR STRESS.

**intraformational** *See* CONGLOMERATE.

**intramicrite** A *\*limestone* consisting of *\*intraclasts* set in a *\*micrite* *\*matrix*. *See* FOLK LIMESTONE CLASSIFICATION.

**intra-orogen basin** A *\*drainage* basin located within a mountain belt, with the drainage network (*see* DRAINAGE PATTERN) oriented parallel to the

\*strike.

**intraparticle porosity** See CHOQUETTE AND PRAY CLASSIFICATION.

**intrasparite** A \*limestone consisting of \*intraclasts cemented together with sparry \*calcite (\*sparite). See FOLK LIMESTONE CLASSIFICATION.

**intrinsic permeability** See PERMEABILITY.

**intrusion** A body of rock, usually \*igneous, that is emplaced within pre-existing rocks. Intrusions are classified according to their size, their shape, and their geometrical relationship to the enclosing rocks.

**intrusive** Applied to an \*intrusion.

**intrusive phonolite** See TINGUAITE.

**invaded zone** See FLUSHED ZONE.

**Inverian** A sub-\*stage of the \*Lewisian, from about 2300–1600 Ma ago (Van Eysinga, 1975). Other authors place this stage within the \*Scourian as a metamorphic event.

**inverse problem** The problem of determining the nature of a physical feature by examining the effects it has on, for example, potential fields, as in \*gravity surveying when an anomaly is interpreted to produce a geologic model. Compare FORWARD PROBLEM.

**inversion** A reversal of a particular trend. **1.** A rock sequence in which the younger sediments are at the bottom. Inversion can be caused by \*overfolding or \*thrusting. It is a major feature of an \*accretionary wedge in which progressively younger oceanic and \*trench \*sediment is \*underthrust, so that while each thrust slice is not inverted, each new thrust affects younger sediment, so producing the inversion. In a positive inversion, \*normal faults on \*passive margins become thrusts on collision. In a negative inversion thrusts become normal faults at the end of an \*orogeny. The inversion of sediments takes place during \*strike-slip faulting. **2.** During an \*orogeny, the uplift that follows subsidence as a reversal of vertical direction. **3.** (velocity inversion) Seismic velocity usually increases with depth, but occasionally a zone of anomalously low velocity occurs between layers of higher velocities giving rise to a velocity

inversion. 4. See POLARITY REVERSAL, GEOMAGNETIC. 5. See TEMPERATURE INVERSION.

**inversion axis** In \*crystallography, an \*axis of symmetry which can be inverted through 180° about its centre in order to achieve a higher degree of symmetry for the crystal. Thus, an axis of two-fold symmetry on inversion every 90° becomes an axis of four-fold symmetry. An inversion axis also obviates the need for a \*centre of symmetry. The notation used to indicate an inversion axis is a 'bar' above the maximum number of repeat positions, e.g. 2<sup>-</sup> and 4<sup>-</sup> for a two-fold and four-fold inversion axis respectively.

**inversion tectonics** A process by which a \*normal fault is reactivated in compression or extension. Compression produces positive \*inversion, generating uplift, and extension produces negative inversion, resulting in subsidence and faulting.

**invertebrate** An animal without a backbone; invertebrates make up about 95% of all animal species and are found in every available habitat on Earth.

**inverted metamorphism** The condition in which higher-grade metamorphic rocks (see METAMORPHIC GRADE) lie on top of lower-grade rocks.

**inverted relief** Inverse relationship between a land-form and the underlying geologic structure, as when a hill is developed in a \*syncline and a valley in an \*anticline. It is a stage beyond the 'normal' or Jura-type relief which is characterized by anticlinal hills and synclinal valleys. The term also denotes the more general case in which, through \*erosion, a hill becomes a valley or vice versa.

**inviscid** Applied to a fluid flow that is assumed to have no \*viscosity.

**involute** Having edges that roll under or inwards. Applied to those coiled cephalopods (\*Cephalopoda) where the final \*whorl of the shell envelops earlier ones. The opposite of involute is evolute, where all the previous whorls are exposed. The term 'serpenticone' is sometimes applied to those evolute cephalopods where the shell resembles a coiled snake. Very laterally compressed shells (discus-shaped) are called 'oxycones'; very flat and inflated types are called 'cadicones'; those which are subspherical are called 'sphaericones'.

**involution (festooning)** Contorted bedding in the near-surface zone of unconsolidated earth material or bedrock. Deformation may be regular, producing festoon-like features (e.g. 'frost boils'), or highly irregular, showing pronounced distortion and twisting. Involution is characteristic of past and present **\*periglacial** zones, and is due basically to ground freezing.

**Io (Jupiter I)** One of the **\*Galilean** satellites and the most geologically (and especially volcanically) active body in the solar system, its volcanoes being due to heat generated by tidal heating and eruption temperatures reaching more than 1000 K. Io radiates more heat than it receives from the Sun. The **\*Voyager** spacecraft observed nine volcanic eruptions. It has a metallic core, rock mantle, and a rocky surface covered with sulphur and sulphur compounds, including sulphur dioxide frost. Io was discovered in 1610 by Simon Marius and Galileo. Its equatorial radius is 1821.3 km; mass  $8.93 \times 10^{22}$  kg; mean density 3530 kg/m<sup>3</sup>; visual albedo 0.61; mean distance from Jupiter 421 600 km; orbital period 1.769138 days; rotational period 1.769138 days; surface temperature about -143 °C, but with one volcanic feature with a temperature measured as 17°C.

**ion** Atom that has acquired an electric charge by the loss (cation; positive charge) or gain (anion; negative charge) of one or more electrons.

**ion drive** See ION ENGINE.

**ion engine (ion drive, ion thruster)** A propulsion device for spacecraft that strips electrons from the atoms of a neutral gas, creating a cloud of positive **\*ions**, which it accelerates along an electric field to generate thrust. Compare PLASMA ENGINE.

**ion exchange (IX)** Reversible exchange of **\*ions** in a crystal for other ions in solution, without disturbance of the crystal lattice or its electrical neutrality. This occurs by **\*diffusion**, particularly in crystals where weakly bonded ions form one- or two-dimensional channelways. Natural **\*zeolites** are used to capture **\*anions** and **\*cations** from solution. Artificial ion-exchange resins with three-dimensional **\*hydrocarbon** networks are commonly used (e.g. in water softeners; for separating **\*isotopes**; in **\*desalination**; and in the chemical extraction of elements from ores).

**ionic bond** Bond formed when an **\*electron** is transferred from one atom to another. The atom that loses the electron becomes a positively charged **\*ion**

and the atom that gains the electron becomes a negatively charged ion. A strong electrostatic force then bonds the two ions together. The bonding in a sodium-chloride crystal (NaCl) is ionic, the crystal lattice containing  $\text{Na}^+$  ions and  $\text{Cl}^-$  ions. *Compare* COVALENT BOND.

**ionic charge** The electrical charge of an *\*ion*, created by the gain (negative charge) or loss (positive charge) of one or more *\*electrons* from an atom or group of atoms. Ionic charge is important in determining the strength of bonding in *\*minerals* (e.g.  $\text{Fe}^{3+}$  makes a stronger bond than  $\text{Fe}^{2+}$  with similar ions), and also which elements can substitute for each other within a crystal *\*lattice*. *See also* IONIC BOND; IONIC RADIUS; VALENCY.

**ionic potential** Measure of the strength of attraction of *\*ions*, expressed as the ratio of *\*ionic charge* ( $Z$ ) to *\*ionic radius* ( $r$ ):  $Z/r$ .

**ionic radius** The distance between the nucleus of an *\*ion* and the electron in its outermost shell, usually reported in *\*ångstroms* or picometres (pm). Although no precise measurement can be made of the size of individual *\*ions*, in practice various techniques (e.g. *\*X-ray* diffraction) can be used to estimate ionic radii in particular crystal structures. Generally, it is found that: (a) within the same group of the periodic table ionic radius increases with increasing atomic number; (b) for elements of the same period (i.e. same horizontal row) that form positive ions, ionic radius decreases with increasing positive charge (reflecting the greater nuclear attraction on the same number of extranuclear *\*electrons*), for example,  $\text{Na}^+ = 1.02\text{Å}$ ,  $\text{Mg}^{2+} = 0.72\text{Å}$ ,  $\text{Al}^{3+} = 0.53\text{Å}$ ,  $\text{Si}^{4+} = 0.40\text{Å}$ ; (c) for the same reasons, if an element can exist in different valence states, the higher the positive charge the smaller the ion, for example,  $\text{Mn}^{2+} = 0.82\text{Å}$ ,  $\text{Mn}^{3+} = 0.65\text{Å}$ ; (d) for elements of the same period forming negative ions, the ionic radius increases with increasing negative charge (due to electronic repulsion).

**ionic substitution (proxy, diadochy)** Replacement or replaceability of one or more kinds of *\*ion* in a *\*crystal* lattice by other kinds of ions of similar size and charge (e.g. in the *\*olivine* series  $\text{Fe}^{2+}$  and  $\text{Mg}^{2+}$  substitute for each other).

**ionization potential** Energy needed to drive an *\*electron* from an atom or molecule without imparting kinetic energy to the electron; this leaves a positive *\*ion* (*\*cation*).

**ionosphere** The part of the atmosphere that lies above about 80km altitude, with the highest concentrations of *\*ions* and free *\*electrons*. The most intense concentration is at 100–300 km altitude. Long-distance radio communications use waves that are reflected by certain regions of the ionosphere where there are particular concentrations of ions and free electrons. This allows radio waves to be transmitted around the curved surface of the Earth. Communications satellites make it easier to transmit higher-frequency waves (e.g. television transmissions) around the Earth, but reflection from the ionosphere continues to be used for radio transmissions, being cheaper.

**ion pair** A positive *\*ion* and a negative ion produced by the transference of an *\*electron* from one atom or molecule to another.

**ion thrust** See ION ENGINE.

**IPOD** See INTERNATIONAL PROGRAMME OF OCEAN DRILLING.

**Ipswichian** **1.** The temperate, last *\*Pleistocene* *\*interglacial*, named after Ipswich, Suffolk, England. **2.** Late Pleistocene deposits that occur in river valleys, often associated with *\*terraces*. Pollen diagrams compiled from them indicate that the climate was not much different from that of the present day. The deposits can perhaps be correlated with those of the *\*Eemian* of north-western Europe.

**IPY** See INTERNATIONAL POLAR YEAR.

**IR** See INTERNAL REFLECTION.

**IRAZÚ** A 1-unit *\*CubeSat* mission by the Asociación Centroamericana de Aeronáutica y del Espacio (ACAE) to demonstrate ACAE's capability to develop and operate such a mission and to allow Costa Rican scientists to collect data on the country's rainforests. The satellite was launched on 2 April 2018, from Cape Canaveral, into a near-circular *\*polar orbit* at an altitude of about 400 km.

**iridescence** Physical phenomenon in which fine colours are produced on a surface by the interference of light that is reflected from both the front and back of a thin film.

**iridescent clouds** High clouds, usually observed within 30° of the Sun, with red, green, yellow, blue, or violet tints along their edges. Tiny cloud particles cause diffraction of light rays, producing the coloured effect.

**iridium anomaly** The anomalously high (typically 50 ppb) concentrations of iridium, relative to typical crustal abundances of less than 1 ppb, observed world-wide in sediments straddling the **\*Cretaceous–\*Tertiary** boundary. The anomaly is attributable to fall-out resulting from a massive **\*asteroidal** or **\*cometary** impact, that is thought by many to have been responsible for the mass **\*extinctions** that define the Cretaceous–Tertiary boundary.

**Iridium NEXT** A constellation of sixty-six operational satellites developed by Iridium Satellite, deployed between 2015 and 2017 in low-Earth **\*polar orbit** at an altitude of 780 km. They host government and scientific payloads, providing real-time visibility 24/7 over the entire Earth's surface and atmosphere. This was augmented in 2013 by Iridium PRIME, which offers space on additional copies of the Iridium NEXT satellites, the first of which were launched in 2017.

**IRIS** See **INTERNATIONAL RESEARCH INSTITUTIONS FOR SEISMOLOGY**.

**Irish elk** (*Megaloceras giganteus*) See **ALLOMETRY**.

**IRM** See **ISOTHERMAL REMANENCE**.

**iron, native** Native occurrences of the metal, Fe, may be of terrestrial or meteoritic origin; sp. gr. 7.5; hardness 4.5; grey; **\*massive** or granular; malleable; if meteoric may be alloyed with nickel and consists of small fragments that have fallen from the outer atmosphere; terrestrial iron is fairly rare but occurs in **\*igneous** rocks in Greenland and in carbonaceous sedimentary deposits.

**iron formation** Iron-rich **\*sedimentary rocks**, mostly of **\*Precambrian** age, containing at least 15% iron. The iron occurs as an **\*oxide**, **\*silicate**, **\*carbonate**, or **\*sulphide**, deposited as laminated, deep-water, shelf-sea, and lagoonal **\*sediments**, often associated with **\*cherts** (see also **BANDED IRON FORMATION**). Other iron formations contain iron-rich **\*ooids**, **\*pellets**, and **\*intraclasts**, representing deposits comparable to shallow marine **\*limestones**. The source of the iron in iron formations is the subject of

considerable debate; origins from volcanic sources, biochemical precipitation, and the diagenetic (see [DIAGENESIS](#)) replacement of [\\*limestones](#) are among the suggestions that have been made.

**iron glance** See [HEMATITE](#).

**iron hypothesis** The proposal that during [\\*ice](#) ages the more arid global climate increased the amount of airborne dust falling on to the ocean surface. The dust particles contain iron that stimulates the [\\*primary productivity](#) of marine phytoplankton. The increased rate of [\\*photosynthesis](#) draws down carbon dioxide from the atmosphere and fixes it in the tissues, shells, and [\\*tests](#) of marine organisms. The overall mechanism would therefore exert a negative [\\*greenhouse](#) effect.

**iron meteorite (siderite)** [\\*Meteorite](#) composed of iron and nickel (4–30% Ni) with only a small proportion of [\\*silicate](#) minerals. Although very common in museum collections, iron meteorites form only a few per cent of observed meteorite [\\*falls](#).

**iron pan** [\\*Indurated](#) [\\*soil horizon](#), found usually at the top of the B horizon, in which iron oxide is the main [\\*cementing](#) material.

**ironstone** Iron-rich [\\*sedimentary rock](#). The source of the iron is primary and/or diagenetic (see [DIAGENESIS](#)), developed by a number of possible processes. These include: (a) replacement of [\\*carbonate](#) particles by [\\*hematite](#), [\\*siderite](#), or [\\*chamosite](#); (b) diagenetic development of nodules, and more continuous horizons, of [\\*siderite](#) within [\\*claystone](#) sequences; (c) primary deposition of [\\*oooids](#) from Fe/Al-rich gels, and subsequent conversion to [\\*chamosite](#) during shallow burial. See [PLINTHITE](#). See also [MINETTE](#); [CLINTON IRONSTONE](#).

**irradiance** The flux density of [\\*electromagnetic](#) radiation falling onto a surface.

**irragic horizon** In the [\\*World Reference Base for Soil Resources](#) classification, a [\\*soil horizon](#) that has been strongly affected by irrigation.

**irregular echinoids** Echinoids ([\\*Echinoidea](#)) in which a [\\*bilateral](#) symmetry is superimposed on the [\\*radial symmetry](#) and where the apical system no longer contains the [\\*periproct](#). In some groups, e.g. the spatangoids, the [\\*aboral](#) portions of the ambulacra (see [AMBULACRUM](#))

terminate above the **\*ambitus** and are situated in recessed, leaf-like ‘petals’, the mouth is situated far forward, and behind the mouth there is a flattened area (the ‘plastron’) that bears flat spines and is formed from the posterior interambulacra (see **INTERAMBULACRUM**).

**Irregulares (Anthocyathea)** (phylum **\*Archaeocyatha**) Class of solitary, rarely colonial invertebrate animals found in Lower, Middle, and Upper **\*Cambrian** rocks. The conical cup is from cylindrical to discoid in outer form, often with an irregular outline, having one, or more usually two, porous walls. The **\*intervallum** contains rods and bars, or **\*septa**, always with **\*dissepiments** and commonly with **\*tabulae**. Compare **REGULARES**.

**irrigation** The process of artificially augmenting the amount of water available to crops. The water may be sprayed directly on to the plants or made available to their root systems through a series of surface channels or ditches.

**irrotational wave** See **P-WAVE**.

**$I_s$**  See **POINT LOAD INDEX**.


**IRS** See **INDIAN REMOTE SENSING**.

**ISAS** See **INSTITUTE OF SPACE AND ASTRONAUTICAL SCIENCE**.

**ischium** In **\*tetrapods**, the part of the **\*pelvis** that projects backward on the ventral side. In Primates, it bears the weight of the sitting animal.

**ISEE** See **INTERNATIONAL SUN—EARTH EXPLORER**.

**island arc** Series of **\*volcanoes** that lies on the continental side of an oceanic **\*trench** of a lithospheric **\*plate**. The **\*volcanicity**, whose products are mainly of intermediate composition, results from the **\*subduction** process; typically it occurs approximately 100 km above the down-going oceanic plate. Island arcs are the sites of strong seismic activity, and have distinctive thermal and magnetic properties. See also **BENIOFF ZONE**; **PLATE MOTIONS**; **PLATE TECTONICS**.



Island arc

**island-arc crustal type** A type of very unstable \*crust with an average thickness of 30 km that comprises fragments of \*continental crust together with volcanic peaks. There is much \*volcanicity and seismic activity.

**iso-** From the Greek *isos* meaning 'equal', a prefix meaning 'equal'.

**isobar** Line on a weather map connecting points at the same atmospheric pressure. On surface charts the values are 'reduced' to sea level. Such isolines are drawn at a given interval in millibars. Contours of \*isobaric surfaces may be drawn to represent surfaces in the upper atmosphere composed of points at the same pressure.

**isobaric gas-tight sampler (IGTS)** An instrument made from titanium that is used to collect samples of hot, corrosive, high-pressure, hydrothermal fluids from a deep submergence vehicle. Being gas-tight, the sample cannot degas as it is carried to the surface and the pressure decreases.

**isobaric surface** A surface on which any point experiences the same atmospheric pressure. *See also* [ISOBAR](#).

**isobase** A line drawn on a map linking points at a particular height above sea level (i.e. a contour line) but that lie on surfaces formed at sea level at a specified time in the past.

**isobath** A line drawn on a map linking points at the same depth below the water surface.

**isochore** A line on a map which joins points of equal vertical interval between two datum planes.

**isochron** A line joining points of equal time intervals or ages. In **\*geochronology** the slope of the isochron may be used to determine the age of a suite of rocks. For example, if it is assumed that all the rocks formed from one **\*magma** had the same initial  $^{87}\text{Sr}:^{86}\text{Sr}$  ratio (see **INITIAL STRONTIUM RATIO**), then there is a simple equation to describe the growth of **\*radiogenic**  $^{87}\text{Sr}$ :  $^{87}\text{Sr} = ^{87}\text{Sr}_0 + ^{87}\text{Rb}(e^{\lambda t} - 1)$  where  $^{87}\text{Sr}_0$  is the number of atoms of the  $^{87}\text{Sr}$  **\*isotopes** incorporated into the rock at the time of formation,  $^{87}\text{Sr}$  and  $^{87}\text{Rb}$  are the numbers of these isotopes after time  $t$ , and  $\lambda$  is the **\*decay** constant. Because the number of  $^{86}\text{Sr}$  is constant, we can derive an equation:  $^{87}\text{Sr}:^{86}\text{Sr} = ^{87}\text{Sr}_0: ^{86}\text{Sr} + (^{87}\text{Rb}:^{86}\text{Sr})(e^{\lambda t} - 1)$  which would give a family of straight lines when plotted on a graph of  $^{87}\text{Sr}:^{86}\text{Sr}$  (on the  $y$  axis) against  $^{87}\text{Rb}:^{86}\text{Sr}$  (on the  $x$  axis). All rock specimens belonging to a co-magmatic suite will plot as points on a straight line called an 'isochron' because all points on the line represent systems having the same age ( $t$ ) and the same initial  $^{87}\text{Sr}:^{86}\text{Sr}$  ratio. In order to date co-magmatic **\*igneous** rocks by the **\*whole-rock** isochron method, a suite of rocks must be collected which span as wide a range of Rb/Sr ratios as possible, so that the slope of the isochron will be well defined. The age of the suite of rocks is obtained from the slope ( $m$ ) using the equation  $m = e^{\lambda t} - 1$ . Isochrons can also be determined in **\*lead-lead dating** by plotting a series of **\*growth curves**. See also **ISOCHRON MAP**.

**isochron map** **1.** A map showing the variation in the difference of the **\*two-way** travel times between two **\*seismic** reflection events. **2.** A map of two-way travel times to the same reflector event marked by **\*isochrons** (lines of equal travel time).

**isochronous** Applied to events or lithological units that either (a) occupy the same time interval (i.e. are of the same duration); or (b) occur simultaneously or are of the same age. In the latter sense 'isochronous' is a synonym for 'synchronous'. Compare **DIACHRONOUS**.

**isoclinal fold** A **\*fold** in which the two limbs are parallel.

**isoclinic chart** A map showing lines of equal *\*inclination* of the *\*geomagnetic* field.

**isoconductivity map** A map whose contours link points of equal electrical or thermal *\*conductivity*.

**isodynamic line** A line on a map joining places of equal magnetic intensity.

**isofrigid** See PERGELIC.

**isogal** Line on a map joining places of equal *\*gravitational* acceleration.

**isogon** A line joining points of equal *\*dip* on the inner and outer bounding surfaces of a folded layer. See DIP-ISOGON METHOD.

**isogonic** Applied to lines on a map joining places of equal angle, commonly geomagnetic *\*declination*.

**isograd** See INDEX MINERAL.

**isogyre** See INTERFERENCE FIGURE.

**isohel** Line connecting points of equal average sunshine duration.

**isohyet** Line on a climate map connecting points of equal average rainfall.

**isohyperthermic** See PERGELIC.

**isomagnetic chart** A map of equal magnetic elements, usually geomagnetic intensity or direction.

**isomesic** See PERGELIC.

**isometric** See CUBIC.

**isomorphous** Applied to two compounds having the same, or nearly the same, crystal *\*form* and containing *\*ions* of approximately the same size or relative size. Isomorphous compounds may show *\*solid solution*. Compare ISOTYPIC.

**isomyarian** See MUSCLE SCAR.

**isoneph** Line on a map joining points of uniform cloud cover.

**isopach** A contour line joining points of equal thickness in a rock layer. The isopach should represent the true thickness, i.e. corrected for dip effects, but

this is rare. Isopachs are essential for estimating the volumes and dispersal of volcanic ashes (i.e. the thickness in centimetres or metres of an air-fall ash around a volcanic vent).

**isopach map** A subsurface geologic map showing *\*isopachs* in plan view throughout a particular geographic area. Isopach maps are constructed, for example, to enable an estimate to be made of the size and shape of a *\*petroleum \*reservoir*, or of the approximate topographic relief of an underlying, older land surface.

**isopycnal** A line joining points of equal density within a water mass. A three-dimensional surface of equal density is called an isopycnal surface.

**isopygous** See PYGIDIUM.

**iso-resistivity map** A map whose contours link points of equal electrical *\*apparent resistivity*.

**isoseismal map** A map with lines of either equal seismic intensity or equal frequency of *\*earthquakes*.

**isospore** See SPORE.

**isostasy** A model for the upper region of the Earth in which differences in elevation are compensated by either low-density roots or lower-density surface rocks. The rigidity of the tectonic *\*plate* allows some departure from this model. See ISOSTATIC ANOMALY; AIRY MODEL; PRATT MODEL.

**isostatic anomaly** A *\*gravity anomaly* on a scale of more than 100 km that is associated with areas previously loaded, e.g. by ice (see GLACIOISOSTASY), lakes, etc., or where recent tectonic activity has loaded the *\*crust*, e.g. mountain formation, volcanic loading, etc. It is generally removed from *\*Bouguer anomalies* as part of the regional gradient, but it can also be calculated. See AIRY MODEL; PRATT MODEL.

**isostatic compensation** The flexural adjustment of the *\*lithosphere*, increase in topography, or presence of low-density roots that is introduced into a model to account for *\*isostatic anomalies*. The actual compensation depends on the model used for the Earth's lithospheric structure. See AIRY MODEL; PRATT MODEL.

**isostructural** See ISOTYPIC.

**isotherm** Line on a climate map connecting points of equal average temperature.

**isothermal remanence (IRM)** The remanence acquired by \*ferromagnetic substances when they are placed in direct magnetic fields at room temperature. See COERCIVITY; SATURATION REMANENCE.

**isothermic** See PERGELIC.

**isotope** One of two or more varieties of a chemical element whose atoms have a common number of protons and electrons (i.e. their atomic number is the same) but which vary in the number of neutrons in their nucleus (i.e. their atomic weight, signified by their mass number, is different). For example, hydrogen exists in the forms  $^1_1\text{H}$  (one proton, no neutron),  $^2_1\text{H}$  deuterium: one proton, one neutron), and  $^3_1\text{H}$  (tritium: one proton, two neutrons). Water in which  $^2_1\text{H}$  replaces the more common  $^1_1\text{H}$  is known as 'heavy water'. There are 300 naturally occurring isotopes, but only 92 naturally occurring elements, and in nature elements often occur as a mixture of isotopes, with one form being the most common. Isotopes may be produced by various nuclear reactions and the products are frequently radioactive. There are three different ways of specifying an isotope; for example  $^{235}\text{U}$ , U-235, and uranium 235 all indicate the isotope of uranium with a mass number of 235. See also ISOTOPIC DATING.

**isotope dilution** An analytical technique used to determine the concentration of an element in a sample by means of a mass spectrometer. The method is based on the determination of the isotopic composition of the element in a mixture. A known quantity of a compound containing an unknown quantity of a particular element is mixed with a \*spike (a known weight of a radioactive \*isotope of the element). The specific activity (disintegrations per second per kilogram) of the spike is known precisely, so the isotopic composition of the mixture can be used to calculate the amount of the element in the sample. A small amount of the mixture is isolated from the sample, weighed, and its specific activity measured. The concentration of the inactive element in the sample may be estimated by the dilution of the radiotracer. Isotope dilution analysis can be applied to all

elements that have two or more naturally occurring isotopes (about 80% of all elements), provided that a spike enriched in one of the isotopes of that element is available. As a technique it has several advantages over other analytical methods. It is free of interference from other elements present and its accuracy is governed by the calibration of the spike solution.

**isotope fractionation** The separation of *\*isotopes* of an element during naturally occurring processes as a result of the mass differences between their nuclei. Although the chemical properties of the *\*isotopes* of an element are the same, there are differences in their physical properties (e.g. density, vapour pressure, boiling point, and melting point) due to the greater vibrational energy of the lighter isotope. Separation (fractionation) of isotopes will occur during such processes as *\*evaporation* or condensation, melting or crystallization, *\*diffusion* through crystals, and isotopic exchange reactions between water in a *\*melt* and *\*minerals* or mineral pairs. The extent of fractionation is dependent on temperature and is more pronounced the greater the mass difference between isotopes in relation to their individual isotopic mass. Significant fractionation occurs naturally with carbon, oxygen, sulphur, and hydrogen-deuterium. Fractionation ratios and isotopic ratios are useful in determining palaeotemperatures, geologic processes, and the modes of formation of rocks and minerals. See [D/H RATIO](#); [OXYGEN-ISOTOPE RATIO](#); [OXYGEN-ISOTOPE ANALYSIS](#); [STABLE-ISOTOPE STUDIES](#); [ISOTOPE GEOCHEMISTRY](#).

**isotope geochemistry** The study of the abundance ratios of *\*isotopes* (both *\*stable* and *\*radioactive*) of major and *\*trace* elements in rocks (e.g. Rb/Sr, Pb/U, etc.), to elucidate a number of geologic problems and processes. These include the age relationships of rocks, and the age of the Earth itself (see [GEOCHRONOLOGY](#); [ISOTOPIC DATING](#); [RADIOMETRIC DATING](#)); palaeotemperatures and geothermometry (see [OXYGEN-ISOTOPE ANALYSIS](#); [GEOTHERMOMETER](#)); and the provenance of natural waters, ore-forming fluids, and *\*magmas* (see [ISOTOPE FRACTIONATION](#); [SMOW](#); [D/H RATIO](#); [OXYGEN-ISOTOPE RATIO](#); [STABLE-ISOTOPE STUDIES](#)).

**isotope hydrology** The use of naturally occurring and introduced *\*isotopes* to date and identify water bodies. Among the most commonly used isotopes are tritium, deuterium, carbon-13, carbon-14, chlorine-36, and oxygen-18.

**isotope tracer** Radioactive isotope, whose movement can be monitored, that is used to trace the pathways by which individual substances move through an organism, a living system, the *\*abiotic* environment, etc. Non-radioactive chemical analogues of certain substances may be used for the same purpose if their movement can be monitored (e.g. caesium, which can be substituted for potassium).

**isotopic dating** Means of determining the age of certain materials by reference to the relative abundances of the parent *\*isotope* (which is radioactive) and the daughter isotope (which may or may not be radioactive). If the decay constant (the *\*half-life* or disintegration rate of the parent isotope) and the concentration of the daughter isotope are known, it is possible to calculate an age. *See also* DATING METHODS; RADIOACTIVE DECAY; RADIOCARBON DATING; RADIOMETRIC DATING.

**isotropic** *See* BELLEROPHONTIFORM.

**isotropic** Applied to substances whose optical or other physical properties are the same from whatever direction they are observed. In thin-section and polished-section microscopy, an isotropic mineral has only one *\*refractive index* or one *\*reflectance* value respectively, whatever the orientation of the mineral as indicated in *\*crossed* polars. The beam of light is not split into two vibration directions, therefore, but passes through or is reflected off the mineral with no change in its optical characteristics.

**isotropic indicatrix** *See* INDICATRIX.

**isotropy** The uniformity of physical characteristics of a medium irrespective of the direction in which they are measured. *Compare* ANISOTROPY; INHOMOGENEITY.

**isotypic (isostructural)** Applied to a pair of *\*isomorphous* compounds in which the relative sizes of the *\*ions* to each other are the same in each pair, but their absolute sizes are different. *\*Solid solution* is impossible.

**isovelocity plot** *See* VELOCITY SURVEY.

**isovol** A line drawn on a map of a coalfield joining points at which coals have similar proportions of *\*volatiles*.

**isp** Abbreviation for *\*ichnospecies*.

**ISSC** International Subcommittee on Stratigraphic Classification.

**Isuan** A *\*system* of the Lower *\*Archaean*, extending from about 3800 Ma to about 3500 Ma ago, and named from the Isuan supercrustal rocks of western Greenland that include the Amitsôq *\*gneisses*.

**itabirite** The name given in Brazil to *\*banded* iron formations.

**ITCZ** See **INTERTROPICAL CONVERGENCE ZONE**.

**iterative evolution** Repeated *\*evolution* of similar or parallel structures in the development of the same main line. There are many examples of iterative evolution in the fossil record, spanning a wide range of groups. This evolutionary conservatism probably is due to the overriding morphogenetic control exerted by certain regulatory *\*genes*.

**Itokawa** 25143 Itokawa is an *\*asteroid* made from rubble, named for the Japanese aerospace engineer Hideo Itokawa (1912–99). The asteroid was discovered on 26 September 1998, and in 2005 the Japanese spacecraft Hayabusa (‘peregrine falcon’) touched down on it twice, collecting samples that it delivered to Earth on 13 June 2010. These were the first samples ever captured and returned from an asteroid.



[http://global.jaxa.jp/article/special/hayabusa\\_sp3/index\\_e.html](http://global.jaxa.jp/article/special/hayabusa_sp3/index_e.html)

- Close-Up on the Asteroid Itokawa.

**IUGS** International Union of Geological Sciences.

**IVD** Abbreviation for *\*ignimbrite* veneer deposit. See **MANTLE BEDDING**.

**Ivorian** A regional *\*stage* of the *\*Mississippian* *\*epoch*, dated at 348–345.3 Ma ago (Int. Commission on Stratigraphy, 2004), preceded by the *\*Hastarian* and followed by the *\*Visean*.

**I-wave** See **SEISMIC-WAVE MODES**.

**IX** See **ION EXCHANGE**.

# J

**Jaccard's index (Jaccard's coefficient)** In **\*biogeography**, an index of faunal resemblance between two regions. It is calculated as  $C/N_1 + N_2 - C$ , where  $C$  is the number of taxa shared between a pair of regions and  $N_1$  and  $N_2$  are the number of species in each of the two regions.

**jacobsite** See MAGNETITE.

**Jacob's staff** A calibrated rod used in the field to measure the thickness of **\*strata**. It is usually marked off in 10 cm and 1 m intervals.

**jade** See JADEITE.

**jadeite** Isolated, rare **\*mineral** of the **\*clinopyroxenes** with the composition  $\text{NaAlSi}_2\text{O}_6$ ; sp. gr. 3.2–3.4; **\*hardness** 6; **\*granular** crystals or **\*massive**; jade is the precious variety of jadeite, although the term includes the **\*amphibole** mineral nephrite; it occurs with **\*albite** in **\*blueschists** (**\*glaucophane** schists) under conditions of low temperature and high pressure at **\*destructive** plate margins.

**Jameson, Robert** (1774–1854) Professor of Natural History at the University of Edinburgh, who studied with **\*Werner** and actively promoted Wernerian theories to his own students and in the *Edinburgh Philosophical Journal*. He translated Cuvier's *Discours préliminaire* into English (1813), adding Wernerian interpretations.

**jamesonite** **\*Sulphide**,  $\text{Pb}_4\text{FeSb}_6\text{S}_{14}$ ; sp. gr. 5.6; **\*hardness** 2.5; greyish-black; **\*metallic** **\*lustre**; **\*massive**, fibrous, or as **\*acicular** crystals with a feather-like form ('feather ore'); occurs in vein deposits associated with other antimony sulphides.

**Janjukian** A *stage* (30–27.5 Ma ago) in the *Oligocene* of south-eastern Australia, underlain by the *Aldingan*, overlain by the *Longfordian* (*Miocene*), and roughly contemporaneous with the *Rupelian* and *Chattian* (Early Oligocene) and lowermost *Aquitanian* (Miocene) stages.

**Janus (Saturn X)** One of the lesser satellites of *Saturn*, with a radius measuring  $99.3 \times 95.6 \times 75.6$  km; mass  $0.0198 \times 10^{20}$  kg; mean density  $650 \text{ kg/m}^3$ ; visual albedo 0.8. It was discovered in 1966 by A. Dolfus, Gerard Kuiper, J. Fountain, and S. Larsen.

**Japan Aerospace Exploration Agency (JAXA)** An institution formed on 1 October 2003 from the merger of the *Institute of Space and Astronautical Science (ISAS)*, the National Aerospace Laboratory of Japan (NAL), and the National Space Development Agency of Japan (NASDA). JAXA performs the activities of all of these agencies relating to research and development in aerospace.

**Japanese Meteorological Agency seismic intensity scale (JMA scale)** A scale for reporting earthquake intensity that is used in Japan and Taiwan. It is based on units of *shindo*, or seismic intensity, on an eight-point scale. For each shindo value it describes the effects on people, indoor situations, outdoor situations, wooden houses, buildings of reinforced concrete, lifelines (utilities, e.g. gas, water, electricity), ground and slopes, and peak acceleration.



<http://www.jma.go.jp/jma/en/Activities/inttable.html>

- Tables explaining the JMA seismic intensity scale.

**Japan Trench** The oceanic *trench* which lies between the northern Japanese islands (an *island arc*) bordering the *Eurasian Plate* and the *Pacific Plate*.

**Japan-type margin** See ACTIVE MARGIN.

**Jaramillo** A normal *polarity subchron* which occurs within the *Matuyama* reversed *chron* and is dated between 0.98 and 1.05 Ma ago (Bowen, 1978).

**jarosite** Member of the *\*alunite* group of *\*minerals*,  $\text{KFe}_3(\text{SO}_4)_2(\text{OH})_6$ ; sp. gr. 3; *\*hardness* 3; yellowish-brown; *\*resinous \*lustre*; massive coatings or fine crystals; occurs as a *\*secondary* mineral, frequently as an alteration product of iron-rich mineral deposits.

**Jason** See COPERNICUS.

**Jason-1** A five-year joint satellite mission between NASA and the Centre National d'Études Spatiales (CNES) of France, launched on 7 December 2001, that monitors the global ocean circulation and events such as El Niño and ocean eddies, and collects data on the ties between atmospheric and ocean circulation. It passes over every point on the surface at intervals of ten days at an altitude of about 1336 km, and it flew in tandem with *\*TOPEX/Poseidon* until that craft ceased operation in 2005. Jason-1 now continues the work of the earlier mission.



<http://www.nasa.gov/centers/jpl/missions/jason.html>

- A joint CNES and NASA mission to study global ocean circulation and its link with atmosphere and climate.

**Jason-2** A joint satellite mission between NASA and the Centre National d'Études Spatiales (CNES) of France, launched on 20 June 2008 into the same *\*polar orbit* as *\*Jason-1*, inclined at  $66^\circ$  to the equator. It complements Jason-1, continuing the work started by *\*TOPEX/Poseidon*.



[http://www.nasa.gov/mission\\_pages/ostm/main/index.html](http://www.nasa.gov/mission_pages/ostm/main/index.html)

- A joint CNES and NASA mission to study ocean current direction and to measure variations in sea surface height.

**jasper** A variety of chalcedonic (see CHALCEDONY) silica ( $\text{SiO}_2$ ) which is reddish-brown, opaque, and *\*cryptocrystalline*. It may be banded to give 'Egyptian jasper' ('ribbon jasper'). A contact metamorphosed (see THERMAL METAMORPHISM) *\*shale* may also be baked to give 'porcelain jasper' which may be similar in appearance but has a different composition.

**jaspillite** \*Jasper interbedded with \*hematite, the name given in Australia to \*banded iron formations.

**Java Trench** The oceanic \*trench which marks the outer, deep edge of the E. Indies \*subduction zone, where the \*Indo-Australian Plate is subducting beneath the \*Eurasian Plate. The trench is about 6 km deep along Java, but becomes shallower to the north-west because of progressive infilling by Bengal Fan \*turbidites. There is an outer, non-volcanic arc, formed of an \*accretionary wedge, and a \*fore-arc basin.

**jawless fish** *see* AGNATHA.

**JAXA** *See* JAPAN AEROSPACE EXPLORATION AGENCY.

**Jeffreys, Sir Harold** (1891–1989) A British geophysicist, astronomer, and mathematician who studied \*seismic waves and developed a model for the interior structure of the Earth. His tables of \*earthquake travel times, calculated in collaboration with K. E. Bullen (1906–76), are still in use. He proposed the \*tidal theory for the origin of the solar system.

**Jeffreys–Bullen curves** The travel times for seismic waves passing through the Earth, including direct, reflected, and refracted waves. They are fundamental to the determination of the density structure of the Earth's interior.

**jet** A hard, black, and lustrous form of \*lignite, found in isolated masses within organic-rich \*shales. Jet is thought to form from the waterlogged debris of driftwood.

**jet stream** Concentrated, high-speed air flow, generally in a broadly westerly (i.e. west to east) direction. The principal global jets are the polar front and the subtropical jets, at heights of about 10–12 and 12–15 km respectively, and the polar-night or winter jet stream in the upper \*stratosphere or \*mesosphere at about 50–80 km. The intensity of the jets in narrow bands (the maximum velocity is commonly about 50–100 m/s, but greater speeds are sometimes observed) results from a large poleward increase in pressure gradient with altitude. This is a product of the pole–equator temperature gradient in the air beneath the jet, i.e. it is a \*thermal wind. As pressure decreases more rapidly with height the lower the temperature, so pressure in the colder, polar air masses decreases more rapidly with height than over the regions with warmer air masses. Above

the jet, wind speed diminishes as the pressure gradient declines with increasing height, owing to the effects of a different heating pattern in the stratosphere. *See also* POLAR-FRONT JET STREAM; SUBTROPICAL JET STREAM.

**JMA scale** *See* JAPANESE METEOROLOGICAL AGENCY SEISMIC INTENSITY SCALE.

**Johannian** A \*stage (52–36 Ma ago) in the \*Eocene of south-eastern Australia, underlain by the \*Wangerripian, overlain by the \*Aldingan, and roughly contemporaneous with the \*Ypresian and \*Lutetian stages.

**JOIDES** *See* JOINT OCEANOGRAPHIC INSTITUTIONS FOR DEEP EARTH SAMPLING.

**joint 1.** A discrete brittle fracture in a rock along which there has been little or no movement parallel to the plane of fracture, but slight movement normal to it. Fracture may be caused by shrinkage, due to cooling or desiccation, or to the unloading of superincumbent rocks by \*erosion or \*tectonism. A group of joints of common origin constitutes a ‘joint set’ and the joints are usually planar and parallel or subparallel in orientation. ‘Joint systems’ comprise two or more joint sets, which are usually arranged systematically with respect to the \*principal stress axes of regional deformation. Cooling joints (shrinkage joints), such as those which split a rock into long prisms or columns to form ‘columnar joints’, most commonly found in \*lavas, are due to differential volume changes in cooling and contracting \*magmas. Unloading joints result from erosional unloading of the \*crust and form flat-lying, sheet-like joint sets, e.g. those found in granitic rocks. **2.** *See* VOIDS, TYPES OF.

**Joint Oceanographic Institutions for Deep Earth Sampling (JOIDES)**  
The original deep-sea drilling programme. This evolved into the \*Deep Sea Drilling Programme (DSDP) and then the \*International Programme of Ocean Drilling (IPOD).



<http://iodp.tamu.edu/index.html>

- JOIDES Resolution Science Operator.

**Joint Polar Satellite System-1 (JPSS-1; NOAA-20)** The first of four satellites forming an operational environmental system operated in partnership between \*NASA and the National Oceanic and Atmospheric Administration (NOAA) that images fires, volcanoes, oil spills, and other incidents and monitors atmospheric temperature, moisture, and pressure. JPSS-1 was launched on 18 November 2017, from California, into a Sun-synchronous \*polar orbit at an altitude of 824 km. Once in orbit, the mission name changed to NOAA-20.

**joint set** See JOINT.

**joint system** See JOINT.

**jöklar** See JÖKULL

**jökulhlaup** Sudden, violent, but short-lived increase in the discharge of a meltwater stream issuing from a \*glacier or \*ice cap, sometimes due to volcanic activity beneath. A lake may develop above the heat source; this is subsequently breached to produce a torrent of meltwater, e.g. Lake Grimsvotn on Vatnajökull, Iceland. Flow velocity may reach 7–8 m/s and the discharge may attain 100 000 m<sup>3</sup>/s (e.g. the Katlahlaups from Myrdalsjökull, Iceland), comparable to rates of flow of the Amazon river.

**jökull** (*pl.* jöklar) The Icelandic word for glacier.

**Jolly balance** See DENSITY DETERMINATION.

**Joly, John** (1857–1933) An Irish physicist and geologist, Joly studied radioactivity in the Earth, showing that it was a source of internal heat and could give rise to convection currents in the interior. He never accepted that radioactivity could be used to determine the age of the Earth, believing that his own calculations, based on the \*salinity of the oceans, were more reliable.

**Jotnian** A \*stage of the Middle \*Riphean, from about 1400 to 1250 Ma ago, of the Baltic Shield region, underlain by the \*Gothian and overlain by the \*Varangian.

**Jotnian orogeny** An episode of mountain building that involved the \*Proterozoic \*sedimentary rocks of the Baltic Shield in what is now Scandinavia.

**jovian** Adjective derived from Jupiter, also known as Jove, and used in descriptions of the planet. In planetary geology it is not usual to spell the adjective with an initial capital letter.

**jovian planet (outer planet)** A general term used to refer to the four outer, giant, gaseous planets of the *\*solar* system: *\*Jupiter*, *\*Saturn*, *\*Uranus*, and *\*Neptune*; it contrasts them to the small, rocky, inner or *\*terrestrial* planets: *\*Mercury*, *\*Venus*, *\*Earth*, and *\*Mars*.

**jovian satellites** See *ADRASTEIA (JUPITER XV)*; *AMALTHEA (JUPITER V)*; *ANANKE (JUPITER XII)*; *CALLISTO (JUPITER IV)*; *CARME (JUPITER XI)*; *ELARA (JUPITER VII)*; *EUROPA (JUPITER II)*; *GANYMEDE (JUPITER III)*; *HIMALIA (JUPITER VI)*; *IO (JUPITER I)*; *LEDA (JUPITER XIII)*; *LYSITHEA (JUPITER X)*; *METIS (JUPITER XVI)*; *PASIPHAE (JUPITER VIII)*; *SINOPE (JUPITER IX)*; *THEBE (JUPITER XIV)*.

**JPSS-1** See *JOINT POLAR SATELLITE SYSTEM-1*.

**Juan de Fuca Plate** The small lithospheric *\*plate* in the north-east *\*Pacific Ocean* which is being subducted slowly under the *\*North* American Plate, giving rise to the generally andesitic volcanic chain from northern California to southern British Columbia. The Juan de Fuca Plate is a remnant of the *\*Farallon* Plate. The Juan de Fuca Ridge is offset from the East Pacific Rise by the San Andreas Fault.

**Juan de Fuca Ridge** A centre of mid-ocean spreading at the *\*divergent margin* between the *\*Pacific* and *\*Juan de Fuca Plates*.

**jug** The colloquial name for a *\*geophone*.

**Juliet (Uranus XI)** One of the lesser satellites of *\*Uranus*, with a diameter of 42 km. It was discovered in 1986.

**Juno** A *\*solar* system asteroid (No. 3), diameter 268 km; approximate mass  $2 \times 10^{19}$  kg; rotational period 7.21 hours; orbital period 4.36 years. It was discovered in 1804 by K. Harding.

**Jupiter** The fifth and largest planet in the *\*solar* system, distant 5.203 AU from the *\*Sun*. It has a radius of 71 900 km, and a mass 318 times and a volume 1403 times that of the *\*Earth*. Its density is 1310 kg/m<sup>3</sup> and it is comprised mainly of hydrogen and helium. The atmosphere is 0.9H–0.1He

(with traces of H<sub>2</sub>O, CH<sub>4</sub>, and NH<sub>3</sub>) which grades down into a liquid shell, overlying a zone of metallic hydrogen. In the centre is a small rock–ice core of about ten Earth masses. Jupiter has at least 61 *\*satellites*, including the four *\*Galilean* satellites.

**Jurassic** One of the three *\*Mesozoic \*periods*: it lasted from 199.6 to 145.5 Ma ago, following the *\*Triassic* and preceding the *\*Cretaceous*. The Jurassic period is subdivided into eleven *\*stages* (*\*Hettangian*, *\*Sinemurian*, *\*Pliensbachian*, *\*Toarcian*, *\*Aalenian*, *\*Bajocian*, *\*Bathonian*, *\*Callovian*, *\*Oxfordian*, *\*Kimmeridgian*, and *\*Tithonian*), with *\*clays*, calcareous *\*sandstones*, and *\*limestones* being the most common rock types. *\*Brachiopods*, *\*bivalves*, and *\*ammonites* are abundant fossils, along with many other invertebrate stocks. *\*Reptiles* flourished on land and in the sea, but mammals were relatively insignificant and presumed to have been predominantly nocturnal. The first birds, including *\*Archaeopteryx*, appeared in the Late Jurassic.

**Jura-type relief** See *INVERTED RELIEF*.

**Juvavic** See *NORIAN*.

**juvenile** 1. Applied to volcanic material derived directly from a *\*magma*. Material derived from the surrounding wall rocks is termed ‘foreign’. 2. See *JUVENILE WATER*.

**juvenile water** Original water, formed as a result of magmatic processes. Juvenile water has never been in the atmosphere. Magmatic water can form in very large quantities. A *\*magma* body with a density of 2.5, an assumed water content of 5% by weight, a thickness of 1 km, and an area of 10 km<sup>2</sup> contains some  $1.25 \times 10^9$  m<sup>3</sup> of water. See also *GROUNDWATER*.

**J-wave** See *SEISMIC-WAVE MODES*.

# K

**Kaena** A reversed **\*polarity subchron** which occurs at about  $2.87 \pm 0.03$  Ma ago within the **\*Gauss normal \*polarity chron**.

**kaersutite** A member of the alkali **\*amphiboles** with the composition  $(\text{Na,K})\text{Ca}_2 (\text{Mg,Fe})_4\text{Ti}[\text{Si}_6\text{Al}_2\text{O}_{22}](\text{OH})_2$  in a group which includes katophorite  $\text{Na}(\text{Na,Ca})(\text{Mg,Fe}^{2+})_4\text{Fe}^{3+} [\text{Si}_7\text{AlO}_{22}] (\text{OH})_2$  and oxyhornblende  $\text{NaCa}_2(\text{Mg,Fe,Fe}^{3+},\text{Al,Ti})_5[\text{Si}_6\text{Al}_2\text{O}_{22}](\text{OH,O})_2$ ; sp. gr. 3.2–3.5; **\*hardness** 5.0–6.0; small **\*euhedral \*crystals**; dark brown to black; this rare group of amphiboles occurs in **\*intermediate alkali \*igneous** rocks.

**KAGUYA** See SELENE.

**Kaiatan** A **\*stage** in the Lower **\*Tertiary** of New Zealand, underlain by the **\*Bortonian**, overlain by the **\*Runangan**, and roughly contemporaneous with the **\*Bartonian** stage.

**Kaihikuan** A New Zealand **\*stage** (232–225 Ma ago) of the **\*Triassic**, preceded by the **\*Etalian** and followed by the **\*Oretian**.

**kainite** **\*Evaporite** mineral,  $\text{MgSO}_4 \cdot \text{KCl} \cdot 3\text{H}_2\text{O}$ ; sp. gr. 2.1; **\*hardness** 2.5–3.0; **\*monoclinic**; variable in colour, from white through yellow to reddish; vitreous **\*lustre**; crystals rare, usually forms granular masses; occurs widely in salt deposits in association with **\*halite**, **\*carnallite**, etc. It is used as a fertilizer and as a source of potassium salts.

**Kainozoic** See CENOZOIC.

**Kalb light line** In optical **\*mineralogy**, a test made on **\*polishing** relief, which is visually similar to the **\*Becke line test**, but different in origin. The microscope is focused clearly on the boundary between the two mineral grains. The microscope tube is raised (or the **\*stage** lowered) and as the

specimen begins to go out of focus a 'line of light' will move towards the softer mineral.

**Kalimnan** A *\*stage* (4.3–3.4 Ma ago) in the *\*Pliocene* of south-eastern Australia, underlain by the *\*Cheltenhamian*, overlain by the *\*Yatalan*, and roughly contemporaneous with the upper *\*Zanclean* (Tabianian) and lower *\*Piacenzian* stages.

**kaliophilite** A potassium-rich *\*feldspathoidal* mineral  $\text{KAlSiO}_4$  that is closely related to *\*kalsilite*.

**Kalpana-1** (**MetSat-1; Meteorological Satellite-1**) An Indian meteorological satellite that was launched on 12 September 2002, from the Satish Dhawan Space Centre, India, into *\*geostationary orbit* at  $74.1^\circ\text{E}$ .

**kalsilite** A *\*feldspathoidal* mineral  $\text{KAlSiO}_4$  and *\*end-member* of a series with *\*nepheline*  $\text{NaAlSiO}_4$  with which it forms limited *\*solid solution*; sp. gr. 2.6; *\*hardness* 5.5; has been reported from the *\*groundmass* of potassium-rich *\*lava* flows.

**Kama** See *KAZANIAN*.

**kame** Steep-sided mound composed of bedded sand and gravel which often shows signs of marginal slumping. It is a land-form of glacial deposition, associated with stagnant ice whose removal by melting causes the collapse.

**kame delta** Flat-topped mound of stratified sand and gravel laid down in standing water at an ice margin. Subsequent ice melt leading to loss of support brings about collapse at the ice-contact margin.

**kamenitza** See *PANHOLE*.

**kame terrace** Continuous valley-side land-form consisting of stratified sand and gravel whose outer edge typically shows collapse features. It is laid down by meltwater at the junction between an ice mass and the valley wall.

**kamikaze grains** Particles found on Mars that shatter into fine fragments when they smash into solid rock surface. The particles form through the aggregation by electrostatic bonding of silt- to clay-sized particles. The very low surface atmospheric pressure on Mars means that martian winds drive them at much higher speeds than similar particles could attain on Earth.

When they smash into solid rock surfaces, the aggregates break into fragments about 20µm in size.

**kandite** See ALLOPHANE; CLAY MINERALS.

**Kanopus-V1** See KANOPUS-VULKAN N1.

**Kanopus-V-1K 1** See KANOPUS-VULKAN-INFRA-KRASNY-1.

**Kanopus-Vulkan-Infra-Krasny-1 (Kanopus-V-1K 1)** A Russian cluster of remote-sensing *\*minisatellites* for Earth observation that has an infrared capability allowing it to detect fires as small as 5 × 5 m on a 2000-km swathe. They were launched on 14 July 2017, from Baikonur, Kazakhstan, into three different *\*polar orbits* at about 500 km, 600 km, and 475 km.

**Kanopus-Vulkan N1 (Kanopus-V1)** A Russian Earth-observation *\*minisatellite* mission to provide data for mapping, to detect forest fires and large-scale emissions of pollutants, and to monitor agriculture and resources. It was launched on 22 July 2012, from Baikonur, Kazakhstan, into a Sun-synchronous near-circular *\*polar orbit* at an altitude of about 510 km.

**Kansan I and II** The second (0.9–0.7 Ma ago) of four glacial *\*stages* occurring in N. America, during which *\*isotope* evidence suggests that the climate was less extreme than during the *\*Nebraskan* episode, although the Kansan glaciation extended further south. The Kansan is approximately equivalent to the *\*Günz* glacial of the Alps.

**kaolin (china clay)** See CHINASTONE; KAOLINITE; KAOLINITIZATION.

**kaolinite (dickite, nacrite, China clay, kaolin)** A very important group of *\*clay minerals* belonging to the 1:1 group of *\*phyllosilicates* (sheet silicates), and with the general formula  $Al_4[Si_4O_{10}](OH)_8$ , kaolinite represents the final product from the *\*chemical weathering* of feldspars to give clays; sp. gr. 2.6–2.7; *\*hardness* 2.0–2.5; *\*monoclinic*; white, greyish, or stained a variety of colours; dull earthy *\*lustre*; on a microscopic scale crystals are hexagonal plates, but it is usually *\*massive*; *\*cleavage* {001}; occurs as a *\*secondary* mineral produced by the alteration of aluminosilicates, especially *\*alkali* feldspars; is widely distributed in *\*igneous* rocks, *\*gneisses*, and *\*pegmatites*, and in *\*sedimentary rocks* due to the action of acidic solutions on a wide variety of rocks by processes of

weathering and low-temperature **\*hydrothermal** reactions. It is distinguished by its plastic feel, but normally it has to be identified by optical and physical tests. It is extensively used when pure as a cheap, general-purpose filler and coating material for paper, in ceramics, and also in chemicals and paints. It is distinctive in soils for its physical stability during wetting and drying and for its small **\*cation-exchange** capacity. Kaolinite dominates the clay minerals present in certain acid and very old soils, but not in **\*oxisols**, some **\*latosols**, or soils formed in arid climates. The name is derived from the Chinese *kau-ling* ('high ridge'), referring to the hill from which the first samples were taken.

**kaolinitization (kaolinization)** High-temperature hydrothermal alteration (see **HYDROTHERMAL ACTIVITY**) and replacement of **\*feldspars**, to varying degrees, to form a fine-grained aggregate of the mineral **\*kaolinite**. Kaolinitization in **\*granites** can be so complete that the rock is reduced to a rotten, friable condition, **\*quartz** being the only mineral to survive the process unscathed. In this condition, the altered granite can be easily broken down by a high-pressure water jet and the kaolinite settled out from suspension in settling pools to produce china clay (kaolin) concentrates. See also **CHINASTONE**.

**kaolinization** See **KAOLINITIZATION**.

**Kapitean** A **\*stage** (6–4.8 Ma ago) in the **\*Neogene** of New Zealand, underlain by the **\*Tongaporutuan**, overlain by the **\*Opoitian**, and roughly contemporaneous with the uppermost **\*Messinian** and lower **\*Zanclean** (Tabianian) stages.

**Karatau** A **\*period** of the **\*Riphean** sub-era, following the **\*Yurmatian**, succeeded by the **\*Sturtian**, and dated at about 1100–800 Ma ago (Int. Commission on Stratigraphy, 2004).

**Karatavian** A **\*stage** of the Upper **\*Proterozoic**, from about 1000 to 700 Ma ago, underlain by the **\*Yurmatian** and overlain by the **\*Vendian** **\*series** (Van Eysinga, 1975).

**Karelian** See **GOTHIAN**.

**Karelian orogeny** An **\*Archaean** episode of mountain building affecting the Baltic Shield along a NW–SE line in what is now central Finland. It

began about 2000 Ma ago and ended about 1900 Ma ago.

**Karginsky** See VALDAYAN/ZYRYANKA.

**von Karman–Prandtl equation** An equation describing the logarithmic variation of water velocity within a channel from zero flow at the stream bed to a maximum velocity at the water surface. Originally developed in aerodynamics, the equation also describes the profile of wind velocity above the ground. See EDDY VISCOSITY.

**K–Ar method** See POTASSIUM–ARGON method.

**Karnian** See CARNIAN.

**karren** German term describing the group of solutional features developed at the surface of an outcrop of hard \*limestone. Forms range from runnels (shallow troughs) a few millimetres deep to fissures (\*grikes) extending several metres into the limestone. The word is sometimes used as a synonym of \*lapiés.

**karst** 1. Area of the Dinaric Alps of Yugoslavia underlain by \*limestone. 2. Any region underlain by limestone and characterized by a set of land-forms resulting largely from the action of \*carbonation. 3. By extension, regions in which other processes produce similar types of land-forms, e.g. \*thermokarst in the \*periglacial environment.

**karstic aquifer** An \*aquifer within a \*karst limestone rock matrix. Such aquifers are normally characterized by large void spaces, relatively high values for hydraulic conductivity (see PERMEABILITY), flat \*water-tables, and extensive networks of solution channels within which \*Darcy's law is not obeyed and flow may be \*turbulent.

**Kashirskian** A \*stage (309.2–308 Ma ago) in the \*Pennsylvanian \*epoch, preceded by the \*Vereiskian and followed by the \*Podolskian.

**Kasimovian** 1. An \*epoch in the \*Pennsylvanian, dated at 306.5–303.9 Ma ago (Int. Commission on Stratigraphy, 2004) comprising the Krevyakinskian, Chamovnicheskian, and Dorogomilovskian ages (these are also \*stage names in eastern European stratigraphy). The Kasimovian is preceded by the \*Moscovian and followed by the \*Gzhelian epochs. 2. The name of the corresponding eastern European \*series, which is roughly

contemporaneous with part of the *\*Stephanian* (western Europe), and the *\*Missourian* and lower *\*Virgilian* (N. America).

**kastanozems** A reference soil group in the *\*World Reference Base for Soil Resources* classification scheme. Kastanozems have a *\*mollic* horizon more than 20 cm below the surface and concentrations of calcium compounds within 100 cm of the surface.

**katabatic wind (drainage wind, mountain breeze)** Generic term for the wind that occurs when cold, dense air, chilled by radiation cooling, usually at night, moves downslope gravitationally beneath warmer, less dense air. The occurrence is frequent and widespread in, for example, the fjords of Norway, and as an outblowing wind over ice-covered surfaces in Antarctica and Greenland, where the wind may be extremely strong near the coasts and less severe in many mountain regions. *Compare ANABATIC WIND.*

**katafront** A weak frontal condition in which warm-sector air sinks relative to colder air. The term was coined by T. Bergeron. *Compare ANAFRONT.*

**katatectic layer** A solution residue, usually of *\*anhydrite* or *\*gypsum*, that is found in a *\*cap rock*.

**katazone (catazone)** The deepest region of *\*metamorphism*, where temperatures are 500–700° C, hydrostatic pressure is mainly strong, and there is little or no shearing *\*stress*. Rocks formed in the katazone include high-grade *\*schists*, *\*gneisses*, *\*granulites*, *\*eclogites*, and *\*amphibolites*. The term is little used nowadays because it is the physical conditions of metamorphism that are considered important, rather than the depth at which they occur.

**katophorite** *See KAERSUTITE.*

**Kawhia** A *\*series* in the *\*Jurassic* of New Zealand, underlain by the *\*Herangi*, overlain by the *\*Oteke*, and comprising the Temaikan, Heterian, and most of the Ohauan *\*stages*. It is roughly contemporaneous with the *\*Bajocian*, *\*Bathonian*, *\*Callovian*, *\*Oxfordian*, and *\*Kimmeridgian*.

**Kazanian 1.** A *\*substage* of the Middle *\*Permian*, which is preceded by the *\*Kungurian*, followed by the *\*Wuchiapingian*, and dated at 270.6–260.4 Ma ago. Some authors extend the Kazanian to incorporate the Ufimian age, referring to this longer time-span as the Kazanian (or Kama) age. **2.** The

name of the corresponding eastern European *\*stage*, which is roughly contemporaneous with the upper *\*Zechstein* (western Europe), upper *\*Guadalupian* and lower *\*Ochoan* (N. America), and the upper Basleoan and lower Amarassian (New Zealand).

**KazEOSat-1** The first satellite in a two-satellite Earth-observation mission developed on behalf of Kazcosmos (Kazakhstan National Space Agency). It provides high-resolution images; the second satellite will provide medium-resolution images. It was launched on 30 April 2014, from French Guiana, into a Sun-synchronous near-circular *\*polar orbit* at an altitude of 759 km.

**kb** See KILOBAR.

**K-band** A range of radar frequencies, 10–36 GHz, commonly used in Doppler radar systems and for scanning vegetation cover.

**keel (carina)** **1.** An external, longitudinal ridge situated on the *\*venter* of an ammonoid (*\*Ammonoidea*). **2.** In flying birds, a ridge projecting forward from the sternum and serving for the attachment of the enlarged pectoral muscles.

**Keewatinian** A *\*stage* of the Upper *\*Archaean* of New Zealand, overlain by the *\*Laurentian*.

**KEI** See KEY EVOLUTIONARY INNOVATION.

**Keiloran** A *\*stage* (443.7–433 Ma ago) of the Early *\*Silurian* of southeastern Australia, underlain by the *\*Bolindian* (*\*Ordovician*) and overlain by the *\*Eildonian*.

**kelly** The driving chuck of a *\*rotary* drill system through which the rotation is transmitted to the drill stem.

**kelp** Brown seaweeds that grow below the low-tide level. Large brown *\*algae*, e.g. Laminaria species, which anchor themselves firmly to the sea bed, are typical. Kelp accumulations are important sources of iodine and potash.

**Kelvin, Lord** See THOMSON, WILLIAM.

**kelvin** See KELVIN SCALE.

**Kelvin scale** A scale of temperature proposed by Lord Kelvin (**William \*Thomson**), which does not include negative values. The unit of the scale is the kelvin (K),  $1 \text{ K} = 1 \text{ }^\circ\text{C}$ ,  $0 \text{ K} = -273.16 \text{ }^\circ\text{C}$ . The base of the scale, absolute zero, is the lowest possible temperature for all substances at which no molecule possesses any heat energy. The **\*triple point** of water is given as 273.16 K.

**Kelvin–Voigt model (Voigt model)** A description of the **\*viscoelastic behaviour** of a medium. The model has two components, an elastic spring and viscous damper, which are connected in parallel, and yields equations that resolve the **\*stress** and **\*strain** and their rates of change over time. The model was devised by **William \*Thomson** (Lord Kelvin) and the German physicist Woldemar Voigt (1850–1919). *Compare* **BURGERS MODEL**; **MAXWELL MODEL**.

**kelyphitic rim (kelyphite)** A **\*mineral layer** that surrounds a mineral of different composition. The term is most often applied to minerals coating **\*garnets**, the kelyphitic rim containing **\*orthopyroxene**, **\*spinel**, **\*clinopyroxene**, and **\*amphibole**, often as two concentric layers, the outer one poor in spinel and the inner one rich in spinel. Several explanations have been proposed for the formation of kelyphitic rims; the most widely accepted is that they result from **\*exsolution** from orthopyroxene and metamorphic reaction between **\*olivine** and garnet during **\*retrograde metamorphism**.

**Kenoran orogeny** The name, no longer much in use, given to a **\*Proterozoic** phase of mountain building affecting the shield area in what is now the Lake Superior region of Canada; it was followed by the **\*Hudsonian** orogeny.

**Kenorland** A **\*supercontinent** that formed 2700–2400 Ma ago, during the **\*Neoarchean \*era**, and comprised what are now N. America, Greenland, Scandinavia, the Baltic region of Eurasia, W. Australia, and part of Africa.

**Kent Ridge 1 (KR1)** An Earth-observing **\*microsatellite** developed at the National University of Singapore (Kent Ridge is the location of the main campus) in collaboration with Berlin Space Technology. It was launched on 16 December 2015, from the Satish Dhawan Space Centre, India, into a near-equatorial orbit at an altitude of 550 km.

**kenyte** A type of \*mafic \*phonolite characterized by abundant large \*phenocrysts of \*anorthoclase, smaller phenocrysts of \*nepheline, with or without \*olivine, and with alkali \*pyroxene, set in a glassy \*groundmass (see GLASS). The type locality is Mount Kenya in E. Africa.

**Kepler's laws of planetary motion** (1) The orbits of the planets are ellipses with the Sun at a common focus. (2) The line joining a planet to the Sun sweeps out equal areas in equal times. (3) The squares of the periodic times are proportional to the cubes of the mean distances from the Sun. These laws were formulated by the German astronomer Johannes Kepler (1571–1630) and published during the period 1609–19.

**keratophyre** A fine-grained, \*igneous rock consisting of \*albite or \*oligoclase as the major component, accompanied by \*chlorite, \*epidote, or \*calcite. \*Phenocrysts of unaltered \*augite are sometimes present. The \*mineral assemblage is almost entirely secondary in origin, indicating that the original rock was a \*tholeiitic \*andesite, the volcanic equivalent of plagiogranite. Keratophyres are found on the ocean floor, and in \*ophiolite complexes where they are associated with \*spilites. Like that of the spilites, the mineral assemblage of keratophyres may result from low-grade, ocean-floor \*metamorphism.

**kernal function (K)** A mathematical function for \*resistivity and depth, from which electrical layering can be computed from \*apparent resistivity data.

**kerogen** Solid, bituminous substance formed of fossil organic material in \*oil shales, which can yield oil by destructive distillation.

**kersantite** A type of \*lamprophyre, characterized by essential \*biotite and \*plagioclase feldspar. If \*augite is present the rock is termed an 'augite-kersanite'.

**Ketilidian orogeny** A \*Proterozoic phase of mountain building that occurred about 1800–1600 Ma ago, affecting what is now Greenland. See also GARDAR RIFTING.

**kettle hole (kettle)** Depression in the surface of glacial drift (especially ablation or kettle moraine), resulting from the melting of an included ice mass. It may be filled with water to form a small lake ('kettle lake').

**kettle lake** See KETTLE HOLE.

**Kew barometer** Barometer in which the scale markings are adjusted to take account of the changes of the mercury level in the cistern, so eliminating the need for adjustment of the cistern to the **\*fiducial** point as is required in the **\*Fortin** barometer.

**Keweenawan** 1. A **\*stage** of the Upper **\*Proterozoic** of N. America, underlain by the **\*Huronian** and followed by the **\*Cambrian** (Van Eysinga, 1975). 2. A Lower-Middle **\*Proterozoic** **\*system**.

**key bed** See MARKER BED.

**key evolutionary innovation (KEI)** A newly evolved feature that preadapts the **\*clade** possessing it for new ecological opportunities.


**khamsin** Hot, dry, dusty wind originating in N. Africa and blowing across Egypt and Sudan, usually between April and June, ahead of **\*depressions** which move eastward or north-eastward in the Mediterranean Sea or across N. Africa, with high pressure to the east. The term is also applied to very strong southerly or south-westerly winds in the Red Sea.

**Kibalian orogeny** See BUGANDA-TORO-KIBALIAN OROGENY.

**kidney ore** See HEMATITE.

**kieselguhr** See DIATOMACEOUS EARTH.

**kill curve** A graph, devised by D. M. Raup, on which the number of taxa becoming extinct is plotted against the time intervals between extinction events. It shows, for example, that an extinction of 30% of species occurs, on average, about every 10 Ma.



Source: Skelton, P. *Evolution: A biological and palaeontological approach*. Addison-Wesley & OU, p. 797.

### Kill curve

**kilobar (kb)** Unit of pressure, equal to 1000 bars (986.923 atmospheres,  $10^8$  N/m<sup>2</sup>, or  $10^8$  Pa).

***Kimberella quadrata*** A fossil of \*Precambrian age, known from Australia and the coast of the White Sea, Russia, that is believed to have been a bilaterally symmetrical animal, possibly resembling a mollusc.

**kimberlite** A brecciated (see BRECCIA), potassic, \*ultrabasic \*igneous rock, consisting of \*megacrysts of \*olivine, \*enstatite, Cr-rich \*diopside, \*phlogopite, pyrope-almandine \*garnet, and Mg-rich \*ilmenite, in a fine-

grained \*groundmass of \*serpentine, phlogopite, \*carbonates, \*perovskite, and \*chlorite, in varying proportions. Kimberlites contain abundant \*xenoliths derived from the \*mantle and, as such, they are extremely useful for investigating mantle mineralogy and chemistry. Many of the mantle-derived xenoliths have been brought up from such great depth that they contain high-pressure minerals, the best known being \*diamond, the high-pressure form of carbon. Many of the megacrysts of enstatite, Cr-diopside and garnet are likewise thought to be derived from the upper mantle through which the kimberlite ascended, and as such should be termed ‘\*xenocrysts’. Near the surface, kimberlite is found in clusters of pipe-like bodies called \*diatremes. Mine workings show that they coalesce at depth and link up with \*dykes of non-fragmental kimberlite.

**Kimmeridgian** A \*stage in the European Upper \*Jurassic (155.7–150.8 Ma ago), preceded by the \*Oxfordian, and succeeded (depending on \*biostratigraphic criteria) by the \*Tithonian (Tethyan) or \*Volgian (Boreal) stages. *See also* MALM.

**Kinderhookian** A \*stage (359.2–348 Ma ago) in the \*Mississippian \*epoch of N. America, underlain by the \*Famennian, overlain by the \*Osagean, and containing the \*Hastarian regional stage.

**Kinderscoutian** A \*stage (318.1–317 Ma ago) in the \*Pennsylvanian \*epoch, preceded by the \*Serpukhovian and overlain by the \*Marsdenian.

**K-index** In geomagnetism, the maximum values for the horizontal intensity ( $H$ ) and the \*declination ( $\tan^{-1} X/Y$ ), each on a range 0 to 9, for a three-hourly interval.

**kinematics** The study of the movement of rocks.

**kinematic viscosity** In a flowing fluid, the ratio of \*dynamic viscosity to density, an important factor in determining the amount of turbulence in the flow.

**kinematic wave** A mechanism by which an increase in the mass of ice at a particular point in a \*glacier is transmitted down the glacier. If ice accumulates, the glacier will be thicker at that point and the locally increased mass will slide 2–5 times faster than the ice on its downslope side. The resulting compression will raise a bulge on the downslope side,

which will press against the ice on its downslope side. Thus the local increase in mass will be transmitted down the glacier as a wave.

**king crab** See CHELICERATA.

**kink band** An asymmetric, linear zone of deformation characterized by short **\*fold limbs** and very small **\*hinge** zones. Kink bands commonly occur as **\*conjugate** sets (see CONJUGATE FOLD).

**kipuka** A Hawaiian term describing an 'island' of land completely surrounded and cut off by **\*lava**. Kipukas are also known by the term 'dagalas' in Italy.

**Kirkfield** See SOUDLEYAN.

**Kirkwood Gaps** See RESONANCE.

**Klazminskian** A Russian **\*stage** (303.9–300.5 Ma ago) in the **\*Pennsylvanian \*epoch**, preceded by the **\*Dorogomilovskian (\*Kasimovian epoch)** and followed by the **\*Noginskian**.

**klippe** In geology, a **\*tectonic \*outlier** produced by the **\*erosion** or **\*gravity sliding** of one or more **\*nappes**. The front portions of the nappes become detached to produce the klippe structure.

**knee twin** See GENICULATE TWIN.

**knick point (headcut)** Abrupt change of gradient in the generally smooth long-profile of a stream, and typically separating two concave-up segments. It is often attributed to a fall in base level: this, it is said, initiates a knick point which then slowly travels upstream. It may alternatively be due to a change in rock type or load size, or to tributary entry.

**knob and kettle (sag and swell topography)** The landscape sometimes found on a recent terminal **\*moraine** complex and consisting of a hummocky mound (the 'knob') alternating with a depression (the 'kettle'). The 'kettle' results from the melting of a block of ice enclosed in the **\*drift**.

**Knobby Terrain** See MARTIAN TERRAIN UNITS.

**knock and lochan** Term descriptive of a glaciated landscape of low relief which is made up of ice-moulded hillocks and intervening lochans (small lakes) eroded along zones of rock weakness. It is especially well developed

in the Lewisian **\*gneiss** area of the coastal lowlands of north-western Scotland.

**knot** Unit of speed equal to one nautical mile per hour (0.515 m/s). It is still used in many countries as a measure of wind speed and current velocity (as well as for the speed of ships and aircraft).

**Knott, Cargill Gilston** (1856–1922) A Scottish seismologist and mathematician whose equations (**\*Knott's equations**) formed the basis of modern reflection seismology. Knott was born in Penicuik and studied at the University of Edinburgh, where he obtained his PhD in 1879. He taught at Edinburgh University until 1883 when he moved to Tokyo Imperial University, where he collaborated with others in establishing a network of seismometers throughout the Japanese Empire. He also undertook the first geomagnetic survey of Japan. Knott returned to Edinburgh University in 1891 as reader in applied mathematics. He was elected a Fellow of the Royal Society, a Fellow of the Royal Society of Edinburgh, and was president of the Scottish Meteorological Society and founder of the Edinburgh Mathematical Society. On leaving Japan the Emperor Meiji awarded Knott the Order of the Rising Sun. Knott died in Edinburgh.

**Knott's equations** A set of equations that were the first to describe the partition of energy between reflected and refracted **\*seismic waves** that reach the interface between two different types of rock at a non-normal incidence. The equations were formulated in 1899 by Cargill Gilston **\*Knott**. *See also* ZOEPPRITZ'S EQUATIONS.

**Kohoutek** A **\*comet** with an orbital period of 6.24 years; **\*perihelion** date 28 December 1973; perihelion distance 1.571 AU.

**Kolmogorov Wiener prediction** *See* KRIGING.

**komatiite** An **\*extrusive** **\*igneous** rock of **\*peridotite** composition, dominated by **\*essential** magnesium **\*olivine** accompanied by lesser amounts of aluminous **\*clinopyroxene** and **\*chromite**, and found as **\*lava** flows and shallow **\*sills** within **\*Archaean** and **\*Proterozoic** rock successions. Many komatiite flows were erupted above their **\*liquidus** temperature and may have had temperatures in excess of 1600°C. Because their melting temperature is much higher than that of **\*basalts**, **\*sediments**, etc., they were quite capable of melting the rocks over which they flowed.

Flows commonly show an upper zone containing magnesium *\*olivine* or aluminous *\*clinopyroxene* crystals with *\*spinifex* texture, indicating extreme *\*undercooling* of the upper part of the flow by contact with the atmosphere on extrusion. Komatiites represent the only known examples of peridotitic *\*magma*.

**KOMPSAT** See KOREA MULTI-PURPOSE SATELLITE.

**kona storm** Storm conditions associated with very strong southerly winds over the Hawaiian islands, bringing heavy rainfall. The winds blow in conjunction with the passage of a *\*depression* to the north of the islands.

**Königsberger ratio** ( $Q$ ,  $Q_n$ ) Originally the ratio ( $Q$ ) of the intensity of *\*natural remanent magnetization* (NRM) of an *\*igneous* rock to the intensity of magnetization induced at room temperature in a magnetic field the same as that in which the original NRM was acquired. Now it usually means the ratio ( $Q_n$ ) of the intensity of NRM to that induced in a magnetic field of 50  $\mu\text{T}$  at room temperature.

**kopje** See KOPPIE.

**Köppen, Wladimir Peter** (1846–1940) A meteorologist who developed the climate classification system that bears his name. He was born in St Petersburg, Russia, of German parents, studied at the universities of Heidelberg and Leipzig, and during 1872–3 worked in the Russian meteorological service. In 1875 he moved to Hamburg, Germany, where he headed a new division of the Deutsche Seewarte, formed to issue weather forecasts for the land and sea areas of northern Germany. He was able to devote himself entirely to research from 1879. He died in Graz, Austria.

**Köppen climate classification** System devised in 1918 by Wladimir Peter *\*Köppen* (1846–1940), with modifications that were completed in 1936, by which climates are divided into six broad groups according to the major vegetation types associated with them, broadly determined by critical temperature and the seasonality of precipitation. For example, a summer temperature of 10 °C defines the poleward limit of tree growth; a winter temperature of 18 °C is critical for certain tropical plants; a temperature of –3 °C indicates some period of regular snow cover. The groups are: (A) tropical rainy climates with temperatures in the coldest month higher than 18 °C; (B) arid climates; (C) warm, temperate, rainy climates in which

temperatures in the coldest month are between 3 °C and 18 °C, and in the warmest month higher than 10 °C; (D) rainy climates typical of boreal forest, in which temperatures in the coldest month are lower than –3 °C (in US usage modified to 0 °C), and in the warmest month higher than 10 °C; (E) tundra, in which temperatures in the warmest month are 0–10 °C; (F) permanent frost and *\*ice* caps, in which temperatures in the warmest month are below 0 °C. Subsets of the main classes (written as capital and lower case letters, e.g. Cs) are: absence of a dry season (f); a dry summer season (s); a dry winter season (w); a monsoon climate, with a dry season and rains at other times (m). Arid climates (B) are subdivided into semi-arid steppe-type (S), and arid desert (W). The temperatures within class (B) are indicated as: mean annual temperature higher than 18 °C (h); mean annual temperature lower than 18 °C with the warmest month higher than 18 °C (k); mean annual temperature and warmest month both lower than 18 °C (k'). Some criticism of the system centres on its arbitrary criteria of temperature associated with fixed boundaries, and the failure of data on temperature and precipitation to account fully for the effectiveness of the precipitation. *See also* STRAHLER CLIMATE CLASSIFICATION; THORNTHWAITE CLIMATE CLASSIFICATION.

**koppie (kopje)** An Afrikaans word for 'hill', and applied to a land-form widely described in Africa. It is similar to a *\*tor*, being a steep-sided, isolated land-form, about the size of a house, and best developed on granitic *\*outcrops*. It often shows an angular, castellated outline, when it is called a 'castle koppie'. A koppie may be a late stage in the destruction of a *\*bornhardt*.

**Korangan** A New Zealand *\*stage* (119–110 Ma ago), preceded by the *\*Mokoian* and followed by the *\*Urutawan*.

**Korea Multi-Purpose Satellite (KOMPSAT)** A South Korean programme to develop a satellite system to provide high-resolution imagery of the Korean Peninsula, including coastal zones and adjacent ocean. KOMPSAT-1 was launched on 20 December 1999; contact was lost on 30 December 2007. KOMPSAT-2 was launched on 28 July 2006, from the Plesetsk Cosmodrome, Russia, into a Sun-synchronous circular orbit at an altitude of 685 km. KOMPSAT-3 was launched on 17 May 2012, from the Tanegashima Space Center, Japan, into a Sun-synchronous orbit at 685.1

km. KOMPSAT-3A was launched on 25 March 2015, from the Jasny Dombrovsky site, Russia, into a Sun-synchronous orbit at 528 km. KOMPSAT-5 was launched from the same site on 22 August 2013, into a Sun-synchronous near-circular orbit at a mean altitude of 550 km. KOMPSAT-6 is scheduled for launch in 2020.

**kosava** A strong, usually cold, easterly or south-easterly wind that blows through the Iron Gate gorge in the Danube valley and continues through the Carpathian Mountains to the Hungarian Plain. The term is also used in a generic sense to refer to a type of *\*ravine* wind.

**Kotlassia prima** Amongst the *\*amphibians*, a few genera referred to the Seymouriamorpha show many developments that suggest they were evolving towards the reptilian condition. Some palaeontologists believe that these animals have attained the 'half-way' stage in the evolution of the first *\*reptiles*. Others, however, including A. L. Panchen, claim that *Kotlassia prima* and its relatives retain a lateral-line system and a tadpole stage in the early part of their life cycle. *Kotlassia prima*, known from the Upper *\*Permian*, is one of the last survivors of this important stock. See also SEYMOURIA.

**KR1** See KENT RIDGE 1.

**KREEP volcanism** The term for possible pre-mare volcanic activity in the lunar crust, producing 'KREEP basalts' with high concentrations of Th, U, K, REE, P, and other incompatible elements. They are elusive as a rock type and only a few lunar specimens have been identified so far. The Apennine Bench formation, in the Imbrium Basin, has been proposed to be composed of such material. An alternative explanation is that both it and KREEP basalts are derived from impact-generated melts in basin-forming collisions.

**Krevyakinian** A Russian *\*stage* (306.5–306 Ma ago) in the *\*Pennsylvanian \*epoch*, preceded by the *\*Myachkovskian* (*\*Moscovian* stage) and followed by the *\*Chamovnicheskian*.

**kriging** (**Gaussian process regression, Kolmogorov Wiener prediction**) In statistics, a method for estimating values that can be interpolated where data is missing, using adjacent data. The method is most often used in spatial studies (e.g. in estimating surface temperatures in areas with few

monitoring stations) and in computer modelling. The method was devised by the South African mining engineer Danie G. Krige (1919–2013).

**krotovina (crotovina)** Animal burrow that has been filled with organic or mineral material from another **\*horizon**.

**K-selection** **\*Natural** selection of those organisms that breed in such a way as to maximize their competitive ability, the strategy of equilibrium species. Most typically it is a response to stable environmental resources. This implies selection for low birth rates, high survival rates among offspring, and prolonged development. *K* represents the carrying capacity of the environment for species populations showing an S-shaped population-growth curve. *Compare* **R-SELECTION**.

**K/T boundary event** The impact of an asteroid, about 10–11 km in diameter, that struck the Earth about 65 Ma ago near Chicxulub, on the coast of northern Yucatán, Mexico. Initial evidence for the impact was an iridium-enriched layer of clays, found in many parts of the world. The Chicxulub crater, now filled with sediments, is about 180 km in diameter. Many scientists have concluded that the impact caused the mass extinction at the end of the **\*Cretaceous**.

**Kuehneosaurus latus** First of the eolacertilians, flying lizards which flourished during the Late **\*Triassic**. Essentially these were of ancestral stock, but the outward extension of the ribs indicate a specialized adaptation towards gliding.

**Kuenen, Philip Henry** (1902–72) A Dutch geologist and one of the founders of modern sedimentology, Kuenen's participation in the Snellius deep-sea expedition to the Moluccas (1929–30) directed his attention to marine geology. One of the very few experimental geologists, he demonstrated the efficacy of high-density **\*turbidity** currents in transporting sand to deep-sea environments. In 1950 Kuenen, together with the Italian geologist C. I. Migliorini, interpreted the graded **\*sandstone** beds of ancient **\*flysch** sequences as **\*turbidites**. The study of their **\*sedimentary structures** has contributed much to **\*palaeogeography**. His most widely known book in the English language is the early classic on *Marine Geology* (1950).

**Kuiper, Gerard Peter** (Gerald Kuiper) (1905–73) A Dutch-born American astronomer who discovered **\*Miranda** and **\*Nereid**, was a member of the

team which discovered **\*Janus**, and who made many important studies of planetary and lunar features and the origin of the **\*solar** system. He also predicted, in 1948, that carbon dioxide is one of the major constituents of the martian atmosphere and discovered that **\*Titan** has a methane atmosphere. He suggested that material left over from the formation of the solar system might form a belt, now called the **\*Kuiper belt**, beyond the orbit of **\*Pluto**. Kuiper was born at Harenskarspel and educated at Leiden. He migrated to the USA in 1933 and became a US citizen in 1937. He worked at the Yerkes Observatory, University of Chicago, and was its director from 1947 to 1949 and again from 1957 to 1960. From 1960 until his death he was director of the Lunar and Planetary Laboratory, University of Arizona. The International Astronomical Union named a crater and geologic period on Mercury for him.

**Kuiper belt** A region of bodies, made mainly from ice, that lies just beyond the orbit of **\*Pluto** and supplies the **\*solar** system with short-period comets, e.g. **\*Halley**. At its outer margins the Kuiper belt extends into the **\*Oort** cloud. The existence of the Kuiper belt was proposed by **Gerard P. \*Kuiper**, in 1988 its role as the source of short-period comets was suggested, and in the early 1990s the first objects belonging to the Kuiper belt were discovered.

**Kuiperian** The fifth and most recent period of the **\*Mercurial geologic timescale**, lasting from 1000 Ma ago until the present and named for **Gerard P. \*Kuiper**. Craters formed during the Kuiperian are surrounded by **\*rays**; some craters are so deeply and permanently shaded that ice, probably derived from **\*comets**, survives in them.

**Kula Plate** A formerly large lithospheric **\*plate** in the north-western Pacific that met the **\*Pacific** and **\*Farallon** plates at a **\*triple junction**. It was subducted during the **\*Jurassic**, **\*Cretaceous**, and **\*Cenozoic**, only fragments now remaining as **\*terranes** in the North American **\*cordillera**.

**Kungurian** The final **\*stage** in the Early **\*Permian \*epoch**, preceded by the **\*Artinskian**, followed by the Roadian (Middle Permian), and dated at 275.6–270.6 Ma ago (Int. Commission on Stratigraphy, 2004). It is roughly contemporaneous with the Weissliegende (western Europe), upper **\*Leonardian** (N. America), and Tae Weian (New Zealand). Boundary and

correlation uncertainties have led to some rocks being assigned to either the Artinskian or Ufimian stages.

**kunkar** See CALCIFICATION.

**Kuril Trench** The oceanic *\*trench* which marks part of the *\*destructive* plate margin between the *\*Pacific* and *\*N. American Plates*. The Kuril Trench is backed by the Kuril *\*Island* Arc, running from Kamchatka to northernmost Japan.

**Kuroshio current** Oceanic surface current that flows northwards from the Philippines, along the Japanese coast, and then out into the N. Pacific. It is an example of a western *\*boundary* current: fast-flowing (up to 3 m/s), narrow (less than 80 km), and relatively deep. It is second in strength only to the Gulf Stream. The warm water transports heat polewards. The volume transport is variable but is normally about  $4 \times 10^7 \text{ m}^3/\text{s}$ .

**kurtosis** The degree of peakedness in a distribution curve. If kurtosis ( $K$ ) is 1, the curve is said to be mesokurtic; if  $K$  is greater than 1 it is leptokurtic; if  $K$  is less than 1 it is platykurtic.

**Kutorginida** (phylum *\*Brachiopoda*) Order of brachiopods with biconvex, calcareous shells, and cardinal areas in both valves. No teeth or sockets are found. In the past this group of Lower *\*Cambrian* fossils has been assigned to both the *\*Inarticulata* and *\*Articulata* classes, but it is suspected to have been derived independently of these classes and in the future may come to be regarded as a distinct class.

**K-wave** See SEISMIC-WAVE MODES.

**kyanite** An important metamorphic *\*index mineral* and one of the three  $\text{Al}_2\text{SiO}_5$  mineral *\*polymorphs*, the other two being *\*sillimanite* and *\*andalusite*; sp. gr. 3.5–3.6; *\*hardness* 4–7; *\*triclinic*; light to darker blue, sometimes green or yellow, occasionally black; *\*vitreous* *\*lustre*; *\*crystals* usually long, columnar, often flattened and bladed, with occasional stellate intergrowths; *\*cleavage* pinacoidal, good {100}, {010}, parting {001}; occurs in regional *\*metamorphic rocks* that have been subjected to high pressures and low to moderate temperatures, and may be found in *\*schists* or *\*gneisses* associated with *\*staurolite* and *\*garnet*.

**Kyushu Satellite for Earth Observation System Demonstration (QSat-EOS)** A \*[microsatellite](#) mission by Kyushu University, Japan, to monitor natural disasters, observe the \*[geomagnetic field](#), detect micro debris, and observe water vapour in the upper atmosphere. It was launched on 6 November 2014, from the Yasný Cosmodrome, Russia, into a Sun-synchronous near-circular orbit at an altitude of 504 km.



**labes** On the surface of *\*Mars*, chaotic surface areas caused by debris resulting from a landslide.

**Labrador current** Oceanic current that brings cold arctic waters southwards into the N. Atlantic along the western margin of Greenland. Frequently it carries icebergs southwards, concentrating them in the area to the east of the Grand Banks in late spring to early summer. Fog banks often occur off the coast of Newfoundland where the Labrador current meets the *\*Gulf Stream*.

**labradorite** See *PLAGIOCLASE FELDSPAR*.

**Labyrinthodontia** Subclass of the *\*Amphibia*, primitive in character, and ranging in length from a few centimetres to several metres. They were the *\*Palaeozoic* and *\*Triassic* amphibians. There are no known connections with other amphibians, but they show affinities with *\*Crossopterygian* fish and with *\*reptiles*.

**labyrinthus** An area of intersecting ridges and valleys on the surface of an extraterrestrial body.

**laccolith** *\*Concordant*, lenticular *\*pluton*, which is circular or elliptical in plan. The floor essentially is flat, but the roof is distinctly domed. Compare *LOPOLITH*.

**lacuna** (*pl.* lacunae) On the surface of *\*Titan*, an irregularly shaped depression resembling a dried lake bed.

**lacunosus** From the Latin *lacunosus* meaning ‘with holes’, a variety of cloud usually associated with the genera *\*altocumulus* and *\*cirrocumulus*.

Thin layers or sheets of cloud display a fairly regular set of holes with frilled edges, so forming a net. *See also* CLOUD CLASSIFICATION.

**lacus** From the Latin *lacus*, meaning ‘lake’, a term that has been used to name small, isolated patches of \*mare basalt on the lunar surface (e.g. Lacus Somniorum, the ‘Lake of Dreams’, north of Mare Serenitatis). On \*Titan the term describes a small, dark plain with clearly defined boundaries.

**lacustrine** Pertaining to a lake.

**LAD** *See* LAST-APPEARANCE DATUM.

**Ladinian** A Middle \*Triassic \*stage, preceded by the \*Anisian, followed by the \*Carnian, and dated at 237–228 Ma ago (Int. Commission on Stratigraphy, 2004). It is roughly contemporaneous with the Fa Lang (China) and Kaihikuan (New Zealand). It should not be confused with ‘Landenian’, a Lower \*Eocene stage.

**lag** A \*normal, \*dip-slip fault, dipping less than 45°, on which there has been a relative downward displacement of the \*hanging wall. *See also* FAULT.

**lag breccia (co-ignimbrite breccia)** A coarse, lithic-rich deposit accumulated synchronously with the formation of \*ignimbrite, but confined to a proximal, near-vent location. As the outer margins of a \*pyroclastic \*eruption column collapse, forming a pyroclastic flow moving under gravity away from the \*vent area, \*clasts which are too large to be supported by the column or flow accumulate as a lag \*breccia around the vent. Because they are deposited as the ignimbrite forms, they are now called ‘co-ignimbrite’ breccias to distinguish them from other types of breccias.

**lag deposit** A coarse-grained residue left behind after finer particles have been transported away, due to the inability of the transporting medium to move the coarser particles.

**LAGEOS** *See* LASER GEODYNAMICS SATELLITE.


**Lagerstätte** (*pl.* Lagerstätten) A sedimentary deposit that is of value because of the \*fossils it contains. A fossil-Lagerstätte contains an

abundance of fossils; a conservation-Lagerstätte is a stratum in which fossils are exceptionally well preserved; a concentration-Lagerstätte is a large accumulation of fossils of particular species, such as a shell bed. The term was introduced by Adolf Seilacher, from the German Lager, 'stratum', and Stätte, 'place'.

**lagoon** Coastal body of shallow water, characterized by a restricted connection with the sea. The water body is retained behind a **\*reef** or islands.

**Lagrangian current measurement** A technique for measuring water movements by tracing the path of a water particle over a long time interval. A device is released into the water and allowed to drift passively with it. Measurements can be made, e.g. by following and plotting the progressive position of a **\*neutrally** buoyant float for subsurface currents, or a drift pole or buoy for surface-water movements.

**Lagrangian points** The locations where the gravitational attraction of two orbiting bodies is equal. Objects located at such points remain fixed so that they are favoured positions for future space stations. There are five such points in the Earth-Moon system; two of them,  $L^4$  and  $L^5$ , lie  $60^\circ$  ahead of and behind the lunar orbit. The Trojan family of **\*asteroids** occupy positions close to two of the jovian Lagrangian points. Lagrangian points are named after the French astronomer Comte J. L. Lagrange (1736–1813).



### Lagrangian points

**lahar (mudflow)** Catastrophic mudflow on the flank of a **\*volcano**. It is a notable feature of the volcanic areas of Indonesia, especially Java. Lahars are the cause of most volcanic fatalities. They may reach more than 100 km from the source volcano, when confined along pre-existing valleys.

**Lake Mungo** See BRUNHES.

**Lamarckism** The theory of evolution propounded by Lamarck.

**Lamarck, Jean Baptiste Pierre Antoine de Monet, Chevalier de** (1744–1829) A French naturalist who, in 1809, proposed the first formal theory of evolution. He advanced the theory that evolutionary change may occur by the inheritance of characteristics acquired during the lifetime of the individual. For example, fossil evidence suggests that the ancestors of the giraffe had short necks: Lamarck proposed that competition for food encouraged them to stretch upward in order to browse among higher vegetation, causing their necks to lengthen, and that this lengthening was passed on to their descendants. Over millions of years the minute increases from each generation to the next culminated in the long-necked form. It is interesting to note that the theory of the inheritance of acquired characteristics did not hold a central position in Lamarck's own writings.

His cardinal point was that evolution is a directional, creative process in which life climbs a ladder from simple to complex organisms. He believed the inheritance of acquired characteristics provided a mechanism for this evolution. Lamarck explained that this progress of life up the ladder of complexity is complicated by organisms being diverted by the requirements of local environments; thus cacti have reduced leaves (and giraffes have long necks). *Compare* [DARWIN, CHARLES ROBERT](#).

**Lamb, Hubert Horace** (1913–97) A British climatologist, who was among the first scientists to draw attention to the natural variability of climates and the social and economic effects of climate change, which he described in his detailed studies of climate history. He studied geography at Cambridge University and in 1936 joined the staff of the Meteorological Office. After refusing to work on the meteorology of spraying poisonous gas, he was transferred to the Irish Meteorological Office, but resigned in 1945, after a disagreement with the director, and returned to the UK Meteorological Office in 1946. In the following years he travelled to Antarctica and in 1954 was transferred to the climatology department of the Meteorological Office. He left this post in 1971 to establish the Climatic Research Unit at the University of East Anglia and was its director until his retirement in 1977.

***Lambeophyllum profundum*** One of the earliest recorded rugose corals from the Upper [\\*Ordovician](#) of N. America. The majority of early [\\*solitary](#) corals were rather small and horn-shaped. They belonged to the subclass [\\*Rugosa](#) and were noted for the presence of well-defined radiating partitions, the [\\*septa](#).

**Lambertian reflector** A perfectly [\\*diffuse](#) reflector which reflects [\\*electromagnetic radiation](#) equally in all directions.

**Lamb's dust-veil index** Index, devised by H. H. [\\*Lamb](#), of the amount of finely divided material suspended in the atmosphere after great volcanic eruptions, and of the duration of an effective veil intercepting the Sun's radiation. It can be calculated from estimates of the amount of solid matter thrown up, or from the reduction of intensity of the solar beam, or from the reduction of temperatures prevailing at the surface of the Earth. The latitude of the volcano affects the index value also, because the maximum extent of the veil over the Earth varies, being greatest after eruptions in low latitudes. (The great eruption of Krakatoa in Indonesia in 1883, which ejected about

17 km<sup>3</sup> of particulate matter into the atmosphere, where it remained for three years, gave an index value of 1000.)

**Lamé constant ( $\lambda$ )** 1. An *\*elastic* constant equal to the *\*bulk modulus* minus two-thirds of the shear modulus. *See also* POISSON'S RATIO. 2. *See* SHEAR MODULUS.

**lamellar** In *\*crystallography*, applied to a special kind of *\*crystal* twinning (also called 'multiple' or 'repeated' twinning) where there are many twin individuals, each one having a tabular or plate-like appearance. Lamellar twinning is common within the *\*plagioclase* feldspars, in places where two adjoining twin slabs or lamellae are mutually reversed with respect to each other and every alternate twin 'plate' or 'slab' has an identical atomic structure.

**Lamellibranchia** *See* BIVALVIA.

**lamina** The finest sedimentary layer, less than 1 cm in thickness. *\*Strata* more than 1 cm thick are termed 'beds' (*see* BEDDING).

**laminar flow** Type of flow (normally in water) characterized by the movement of fluid particles parallel to each other, with no transverse movement or mixing. Velocity increases steadily away from the bed. Laminar flow in channels is found only at low velocities and adjacent to smooth surfaces. It is almost ubiquitous in soil moisture and *\*groundwater* (except in *\*karstic* aquifers).

**lamine** Comprising layers (laminae, sing. lamina) of material.

**lamination** Very fine stratification, composed of discrete layers (laminae, sing. lamina) of *\*sediment*, a millimetre or so in thickness. Lamination is a smaller-scale feature than *\*bedding*.

**laminite** A finely *\*laminated* *\*sediment* or *\*sedimentary rock*.

**laminoid fenestrae** *See* FENESTRAE.

**lamprophyre** A dark-coloured, strongly *\*porphyritic*, *\*intrusive* *\*igneous* rock, containing abundant *\*euhedral* *\*phenocrysts* of *\*biotite* and/or *\*amphibole* which can be accompanied by phenocrysts of *\*olivine*, *\*diopside*, *\*apatite*, or opaque oxides, set in a *\*mafic*, *\*felsic*, or glassy *\*groundmass*. There are no felsic phenocrysts present in this rock type. The

lamprophyres are subdivided into a number of types on the basis of their most abundant mafic component and the presence and type of feldspar in the groundmass. Lamprophyre types include *\*minette*, kerstantite, *\*alnöite*, *\*vogesite*, *\*camptonite*, and *\*monchiquite*, and are found intruded as *\*dykes* and *\*sills*.

**lamp shells** See BRACHIOPODA.

**Lancefieldian** A *\*stage* (482–475 Ma ago) of the Early *\*Ordovician* of Australia, underlain by the *\*Warendian* and overlain by the *\*Bendigonian*.

**land and sea breezes** Circulations of air common along coasts, caused by a low-level pressure gradient due to the differential heating of land and sea. On summer days, the land surface heats more quickly than the adjacent sea, air rises over the land, moves over the sea at height, then subsides, producing a shallow convection cell and reducing the surface pressure over land and increasing it over the sea. The resulting pressure gradient from sea to land results in a gentle, cooling, landward ‘sea breeze’ whose maximum strength is usually developed by late afternoon. Land then cools more rapidly than the sea, and at night and in early morning the cooler land and relatively warmer sea produce a reverse-flow convection cell, with a seaward ‘land breeze’. The horizontal extent of well-developed land and sea breezes is typically limited to about 40 km from the coast, but associated air movements can often be detected over a much wider coastal belt.

**land bridge** Connection between two land masses, especially continents, e.g. the Bering land bridge that once linked Alaska and Siberia across the Bering Strait, that allows migration of plants and animals from one land mass to the other. Before the widespread acceptance of *\*continental drift*, the existence of former land bridges was often invoked to explain faunal and floral similarities between continents now widely separated. On a smaller scale, the term may be applied to land connections that have now been removed by recent tectonics or the *\*Flandrian \*transgression* (e.g. between northern France and south-eastern England).

**landfill** See MADE GROUND.

**Landsat** A series of *\*satellites*, mainly carrying *\*multispectral scanners*, and more recently also carrying *\*thematic mappers*. They are primarily designed for scanning the vegetation cover of the land surface of the Earth

and evaluating the effectiveness of satellite-based scanning systems for routine monitoring. The recordings are available on magnetic tape for computer analysis or in photographic form (usually known as Landsat images). Landsat is complemented by Seasat. Landsat-7 is still operational and is due to be replaced in December 2020 by Landsat-9; Landsat-8 (formerly called the Landsat Data Continuity Mission, LDCM) is also operational.



<http://landsat.gsfc.nasa.gov/>

- A NASA programme of satellites from 1972 to collect visual information about the Earth.

**landscape ecology** See GEOECOLOGY.

**landslide** See MASS-WASTING.

**Langhian** A *\*stage* in the *\*Miocene* epoch, dated at 15.97–13.65 Ma ago (Int. Commission on Stratigraphy, 2004) which is preceded by the *\*Burdigalian* and followed by the *\*Serravallian* ages. Most authors subdivide the Langhian into the Early Langhian age (Early Miocene) and Late Langhian age (Middle Miocene). It is roughly contemporaneous with the *\*Relizian* and lower *\*Luisian* (N. America), *\*Cliffdenian* (New Zealand), and *\*Balcombian* (Australia). The *\*stratotype* is found between Cessole and Case dei Rossi, northern Italy.

**La Niña** A strengthening of the south-easterly *\*trade winds* and the *\*Equatorial* current in the South Pacific, linked to the *\*southern oscillation*. It is the opposite of *\*El Niño*. See also ENSO EVENT.

**lanthanide** See RARE-EARTH ELEMENT.

**Laomedeia (Neptune XII)** A satellite of *\*Neptune* with a radius of 20 km, visual *\*albedo* of 0.16, and an orbital period of 3167.9 days.

**LAPAN-A2** An Indonesian *\*microsatellite* (Lembaga Penerbangan dan Antariksa Nasional) that provides Earth observation with a resolution of 5 m along a 3.5-km swathe. It was launched on 28 September 2015, from the Satish Dhawan Space Centre, India, into a near-circular, near-equatorial orbit at an altitude of 650 km.

**LAPAN-A3 (LiSat)** A cooperative, remote-sensing *\*microsatellite* project between LAPAN (Lembaga Penerbangan dan Antariksa Nasional) of Indonesia and the Bogor Agricultural University to monitor food resources and environmental change. The microsatellite was launched on 22 June 2016, from the Satish Dhawan Space Centre, India, into a Sun-synchronous circular orbit at an altitude of 515 km.

**lapiés** Variety of *\*karren*, consisting of shallow, straight grooves incised by solution into a sloping surface of *\*limestone*. It may constitute a dense, subparallel network which develops rapidly upon exposure of the limestone.

**lapilli** *\*Pyroclastic* fragments or *\*tephra* ranging in size from 2 to 64 mm. Lapilli may be composed of primary magmatic material (e.g. *\*pumice*), accessory lithic material, accidental lithic material, or accretions of wet *\*ash*-size material (*\*accretionary lapilli*). The size-term 'lapilli' is usually applied to the lithological *\*clast* type to give descriptive terms, e.g. 'pumice lapilli' or 'accessory lithic lapilli'.

**lapilli-tuff** See TUFF.

**lapis lazuli** See LAZURITE.

**Laplace, Pierre Simon, Marquis de** (1749–1827) A French mathematician and physicist, Laplace is best known for his nebular hypothesis of the formation of the solar system, published in *Exposition du système du monde* (1796). He also worked on planetary motions, the theory of tides, etc.

**Laplace's equation** The sum of the rates of change of a potential field gradient in three perpendicular directions is equal to zero. For a potential function  $U(x,y,z)$ , Laplace's equation states:  $\nabla^2 U = \delta^2 U / \delta x^2 + \delta^2 U / \delta y^2 + \delta^2 U / \delta z^2 = 0$ , where  $\nabla$  is the Laplace operator.

**lapout** The lateral termination of a stratum at the limit of its original deposition. There are two types: *\*baselap* and *\*toplap*, baselap being further divided into *\*onlap* and *\*downlap*.

**lappets** See APERTURE.

**lapse rate** Rate of decrease of temperature per unit height in the atmosphere. In the *\*troposphere* the average rate is approximately 6.5 °C

per 1000 m. *See also* [ADIABATIC](#); [DRY ADIABATIC LAPSE RATE](#); [ENVIRONMENTAL LAPSE RATE](#); [SATURATED ADIABATIC LAPSE RATE](#).

**lapse time** Estimate of the time interval between the ending of [\\*nucleosynthesis](#) and the consolidation of [\\*meteorite](#) parent bodies. Lapse times may be calculated from the detection in meteoritic material of the daughter [\\*isotopes](#) (*see* [DAUGHTER](#) nuclide) of short-lived radionuclides which are now extinct, but which survived until meteoritic condensation. Times of about  $10^8$  years have been deduced.

**Lapworth, Charles** (1842–1920) An English geologist, Lapworth was responsible for ending the controversy between [\\*Murchison](#) and [\\*Sedgwick](#) over the Lower [\\*Palaeozoic](#), by proposing in 1879 that the area in dispute should form a third system, the [\\*Ordovician](#). He also published a classification of [\\*graptolites](#) (1873), and worked on the structure of the Scottish Highlands, identifying the Moine thrust.

**Laramide-Columbian orogeny** A late [\\*Cretaceous](#) to early [\\*Eocene](#) mountain-building episode, affecting an area extending from what are now the south-western USA to northern S. America, caused by thrusting associated with the [\\*subduction](#) of the [\\*Farallon](#) Plate beneath the [\\*North American](#) Plate.

**Laramide Province** A North American region of north-trending mountains and basins located in Wyoming and extending into Colorado, Montana, Utah, and South Dakota, where [\\*Precambrian](#) [\\*basement](#) has been uplifted due to shortening along basement faults. *See* [LARAMIDE–COLUMBIAN OROGENY](#).

**LARES** *See* [LASER RELATIVITY SATELLITE](#).

**large-aperture seismic array (LASA)** An [\\*array](#) of [\\*geophones](#) in Montana, USA, built in 1965, which is used to detect seismic events and to distinguish between those caused by nuclear explosions and those caused by [\\*earthquakes](#).

**large-ion lithophile (LIL)** Element of large [\\*ionic radius](#) and with a [\\*valency](#) of 1 or 2 (e.g.  $\text{Rb}^{2+}$ ,  $\text{Pb}^{2+}$ , and  $\text{Ba}^{2+}$ ), which during [\\*igneous](#) fractionation tends to be concentrated in silicic [\\*melts](#), and from which LILs are incorporated mainly into potassium [\\*silicates](#) such as the [\\*alkali](#)

feldspars and \*micas. See also FRACTIONAL CRYSTALLIZATION; INCOMPATIBLE ELEMENTS.

**large low shear velocity province (LLSVP)** One of two regions at the base of the \*mantle, one beneath Africa and the other beneath the Pacific, extending upward for about 1000 km, through which \*S-waves travel more slowly than they do through surrounding material. \*Earth tides are sensitive to the density, hence \*buoyancy, of the LLSVPs.

**LARI** See LOW-ASPECT-RATIO IGNIMBRITE.

**Larissa (Neptune VII)** A satellite of \*Neptune, measuring  $208 \times 178$  km; visual albedo 0.06.

**larvikite** A coarse-grained, \*intrusive \*igneous rock that consists of essential potash \*feldspar, \*oligoclase feldspar, titaniferous \*augite, and sodic \*amphibole, together with accessory \*biotite, \*quartz, or \*nepheline (depending on the \*silica saturation), \*magnetite, \*zircon, and \*apatite. The two feldspar types occur in equal proportions, making the rock a type of \*monzonite. Larvikite was first described by \*Brøgger in 1890 from the Larvik district of southern Norway. Larvikite can be an impressive ornamental stone.

**LASA** See LARGE-APERTURE SEISMIC ARRAY.

**Laschamp** See BRUNHES.

**laser** An acronym for *light amplification by stimulated emission of radiation*, a device for emitting a single, intense beam of coherent, monochromatic light (i.e. light at a single wavelength). See also MASER.

**laser altimeter** An instrument that measures the distance from an orbiting spacecraft to a planetary or \*satellite surface below by measuring the time of travel of a reflected \*laser beam. Accuracy is within a few metres. Since the position of the spacecraft is known, elevations on the surface can be obtained.

**Laser Geodynamics Satellite (LAGEOS)** A \*NASA passive satellite used for geodynamic measurements by laser ranging from its orbit and perturbations in the orbit, allowing its position to be measured to within 10 cm. LAGEOS-I was launched on 4 May 1976, from California, into a near-

circular **\*polar orbit** at an altitude of 5858–5958 km. LAGEOS-II was launched in 1992 into a complementary orbit.

**laser ranging** The establishment of precise Earth-Moon distances by aiming **\*laser** beams telescopically, from the Earth at reflecting mirrors placed on the lunar surface by the Apollo 11–15 missions, and measuring the return time to the receipt of the reflections. By making such measurements from different terrestrial sites relative **\*plate motions** can be determined.

**LAsER RELativity Satellite (LARES)** An Italian geodynamic satellite mission to measure frame-dragging due to Earth's angular momentum and to test the gravitomagnetic field. The satellite was launched on 13 February 2012, from Kourou, French Guiana, into a circular orbit at an altitude of 1450 km.

**last-appearance datum (LAD)** The last recorded occurrence of a key taxon in biological history.

**Late-Devensian Interstadial** See **WINDERMERE INTERSTADIAL**.

**late-glacial** Term usually applied to the time between the first rise of the temperature curve after the last minimum of the **\*Devensian** glacial stage and the very rapid rise of temperature that marks the beginning of the post-glacial, or **\*Flandrian** period. The late Devensian extends in Europe from about 15 000 to 10 000 BP and shows a characteristic three-fold climatic and hence depositional sequence, from cold, Older **\*Dryas** deposits, to a warmer **\*Bølling-Allerød** interstadial, to colder, Younger Dryas.

**late heavy bombardment (LHB)** A period from about 4.1-3.8 billion years ago when many **\*asteroids** collided with the Earth. The bombardment is late in the formation of the Earth and the other inner planets by accretion. The evidence for the LHB comes from the **\*radiometric dating** of impact-melted lunar rocks collected during the **\*Apollo** programme.

**latent heat of transition** The heat required to activate a **\*phase** change from a solid to a liquid or from a liquid to a gas, i.e. to a higher energy state (e.g. latent heat of melting), measured in joules per mole (J/mol). Latent heat is released by reactions in the reverse direction (e.g. latent heat of crystallization).

**lateral** To or on the side.

**lateral accretion deposit** Inclined layers of *\*sediment*, deposited laterally rather than in horizontal *\*strata*, particularly by the lateral outbuilding *\*sediment* on the surface of a river *\*point* bar. The inclined surfaces thus record the progressive migration of the point bar. The *\*dip* of the lateral accretion deposit can be used to determine the size of the point bar and *\*channel* geometry. *See also* EPSILON CROSS-BEDDING.

**lateral dispersion** *See* DISPERSION.

**lateral moraine** *See* MORAINE.

**laterite** Weathering product of rock, composed mainly of hydrated iron and aluminium oxides and hydroxides, and *\*clay* minerals, but also containing some silica. It is related to *\*bauxites* and is formed in humid, tropical settings by the *\*weathering* of such rocks as *\*basalts*. *See also* PLINTHITE.

**laterolog sonde** An electrical sonde for measuring the electrical resistivity of rocks within a *\*borehole* (resistivity logging). Short sondes (40 cm or less) normally measure the resistivity of the *\*mud cake* and *\*invaded* (flushed) zone. Longer laterologs measure the resistivity of the outer edges of the invaded zone and of the uninvaded rocks. Major differences in the resistivity values measured by shallow and deep sondes will occur in *\*reservoir rocks* containing appreciable quantities of hydrocarbons, so permitting the estimation of the thickness of hydrocarbon reservoirs. Interpretation of long laterologs is affected by the *\*Delaware* effect. *See also* MICROLOG.

**latite** A *\*porphyritic*, *\*extrusive* *\*igneous* rock containing *\*euhedral* *\*phenocrysts* of calcium-rich *\*plagioclase* feldspar, accompanied by lesser amounts of *\*biotite*, magnesium *\*clinopyroxene*, magnesium *\*orthopyroxene*, iron-titanium oxides, and *\*olivine* phenocrysts set in a *\*trachytic* *\*groundmass* dominated by *\*alkali* feldspar. Plagioclase accounts for 40–60% of the total feldspar content of the rock. Latites are the volcanic equivalents of monzonites and, as such, they are members of the *\*calc-alkaline* *\*magma* series. The term is sometimes used to describe potassium-rich *\*andesites*.

**latite-andesite** A \*porphyritic, \*extrusive \*igneous rock containing \*phenocrysts of zoned \*plagioclase feldspar, \*augite (\*clinopyroxene), \*orthopyroxene, and some \*olivine set in a \*groundmass of \*andesine plagioclase and \*alkali feldspar, \*pyroxene, and iron oxides. Plagioclase accounts for 60–90% of the total feldspar content. This rock is transitional to a true \*andesite, which is characterized by plagioclase forming more than 90% of the total feldspar content of the rock.

**latitude correction** The correction made to \*gravity readings to allow for the difference in \*gravitational acceleration as a function of distance from the equator. See INTERNATIONAL GRAVITY FORMULA.

**latosol** A red soil composed of fine, loose, mineral grains.

**lattice** A regular, three-dimensional framework which indicates the ordered arrangement of atoms in \*crystals. The smallest complete lattice is known as the ‘unit cell’ and it may be repeated many times to form a complete crystal. The shape of the unit cell varies according to the arrangement of the points of the lattice in space, i.e. a ‘space lattice’.

**lattice energy** The work required to decompose a crystal lattice of one \*mole of a substance into elementary structural units, and displace them so there is a large distance between them.

**lattice gliding** See DEFORMATION TWINNING.

**laumontite** Member of the \*zeolite group of minerals,  $\text{CaAl}_2\text{Si}_4\text{O}_{12}\cdot 4\text{H}_2\text{O}$ ; sp. gr. 2.3; \*hardness 3.5; whitish; square \*prismatic crystals; occurs as a \*secondary mineral in veins and cavities within volcanic rocks and some associated mineral deposits, and may also occur within the zeolite \*facies of \*metamorphic rocks.

**Laurasia** The northern continental mass produced in the early \*Mesozoic by the initial rifting of \*Pangaea along the line of the N. Atlantic Ocean and \*Tethys. Laurasia included what was to become N. America, Greenland, Europe, and Asia, while the large, southern, continental mass (called \*Gondwana) was later to divide into S. America, Africa, India, Australia, and Antarctica.

**Laurentia (Laurentian Shield)** The \*Precambrian shield (see CRATON) of central eastern Canada. The name is derived from the St Lawrence River

and has been applied to a series of *\*granites*, *\*gneisses*, and *\*metasediments* that are older than 2500 Ma. The Laurentian Shield forms the ancient 'core' of Canada, around which younger *\*orogenic* belts have been accreted.

**Laurentian** A *\*stage* of the uppermost *\*Archaean* of New Zealand, underlain by the *\*Keewatinian*.

**Laurentian Shield** See LAURENTIA.

**Laurentide ice sheet** An area of continental ice that lay over the eastern part of Canada during the *\*Pleistocene* glaciations. The centre of the ice mass may have originated in or near northern Quebec, Labrador, and Newfoundland, and spread out to the south and west. At its maximum spread it may have covered an area of  $13 \times 10^6$  km<sup>2</sup>.

**lava** Molten rock, normally a *\*silicate*, erupted by a *\*volcano*. It may be *\*vesicular*, glassy, or *\*porphyritic* in texture, and varies between *\*acidic* and *\*basic* in composition. Its behaviour on extrusion and its relief-forming capacity depend largely on its *\*viscosity*, which is affected by silica content, temperature, and amount of dissolved gases and solids. Generally, the less viscous the lava the faster the flow, and the more viscous the lava the greater the tendency towards explosive eruption. Two varieties of basaltic lava surface are recognized: 'aa', a jagged, stony clinker, bristling with sharp points; and 'pahoehoe', characterized by a smooth, ropy appearance. Andesitic and rhyolitic lavas tend to have 'blocky' surfaces, characterized by smooth-faceted blocks, 1–5 m in diameter.

**lava blister** The surficial swelling of a plastic *\*lava* flow crust in response to the puffing up of gas or vapour from beneath the flow. Blisters may also form through hydrostatic or *\*artesian* forces in the lava. They are usually 1–150 m in diameter, with a maximum height of 30 m, and are hollow. Compare TUMULUS.

**lava channel** The depression between two parallel and closely spaced retaining walls (*\*lava* levées) in which a flow of *\*lava* is confined. The retaining walls are composed of cooled blocks of lava from the flow itself. As the lava flows, the height of the flow surface pulsates and lava may overflow its retaining walls; this adds a coating of chilled lava to the walls,

thereby increasing their height and the effectiveness with which they constrain the flow.

**lava haze (laze)** White cloud that forms when hot *\*lava* reacts with sea water. The lava cools rapidly, forming glass that shatters to produce fragments of all sizes, including tiny shards, and chemical reactions between the lava and salt ions in the water produce hydrochloric acid. Lava haze consists of the resulting mixture of steam, HCl, and glass shards.

**lava lake** Molten *\*lava*, usually basaltic in composition, held in a depression (e.g. a large *\*caldera* or *\*crater*) over a magmatic vent. A lava pond may form where a lava flow fills a topographic depression. Large lava lakes found in lava-flooded calderas or craters often display surface features indicating the presence of numerous thermal convection cells within the lake. Lava lakes may be sustained for many years (e.g. Kilauea Crater, Hawaii, which has been an important tourist attraction, and at present there is a continuous lava pond at Pu'u O'o, Hawaii).

**lava levée** One of the scoriaceous retaining walls on either side of a *\*lava channel*. Four types are recognized: initial levées form when the yield strength of the lava permits the margins of a flow to remain stationary and crystallize while the central part of the flow continues to move; accretionary levées are formed near *\*boccas* where piles of surface clinker are accreted to the flow margins; rubble levées form when the sides of a flow expand outwards by the avalanching of surface and marginal crystalline clinker; and overflow levées form when lava repeatedly overflows existing rubble levées.

**lava tube** A hollow passage beneath the surface of a solidified *\*lava* flow, formed by the withdrawal of molten lava from a former distributory tunnel, which fed lava below the stationary surface of the flow to the advancing flow margin, sometimes transporting the lava over long distances. Tubes range in width from less than one metre to more than 30 m and in height up to 15 m. They can be tens of kilometres long, e.g. in Victoria, Australia. Most tubes are developed in flows displaying *\*pahoehoe* surface morphology, although they can develop in flows with *\*aa* surface morphology. Mount Etna, Sicily, has examples of both.

**law of constancy of interfacial angles** In all *\*crystals* of the same substance, the angles between corresponding faces have the same value

when measured at the same temperature. This concept was first proposed by *\*Steno* in 1669 and was formulated as a law by the French mineralogist Jean-Baptiste Louis Romé de l'Isle (1736–90) in 1772.

**law of constant proportions** Pure substances always contain the same elements in the same proportion by weight.

**law of correlation of facies** See WALTHER'S LAW.

**law of cross-cutting relationships** An *\*igneous* rock, *\*fault*, or other geologic feature must be younger than any rock across which it cuts.

**law of faunal succession** The principle, first recognized at the beginning of the 19th century by William *\*Smith*, that different *\*strata* each contain particular assemblages of *\*fossils* by which the rocks may be identified and *\*correlated* over long distances; and that these fossil forms succeed one another in a definite and habitual order. This law, together with the *\*law of superposition of strata*, enables the relative age of a rock to be deduced from its content of fossil faunas and floras.

**law of Haüy** See LAW OF RATIONAL RATIOS OF INTERCEPTS.

**law of original horizontality** *\*Sedimentary rocks* are always deposited as horizontal, or nearly horizontal, *\*strata*, although these may be disturbed by later earth movements. The law was first proposed in the mid-17th century by Nicolaus *\*Steno*.

**law of rational indices** See LAW OF RATIONAL RATIOS OF INTERCEPTS.

**law of rational ratios of intercepts (law of rational indices, law of Haüy)** *\*Crystal faces* cut the *\*crystallographic axes* in simple, whole-number ratios. These ratios may be measured with reference to a unit plane having a ratio of 1:1:1 for three crystallographic axes and 1:1:1:1 for four crystallographic axes.

**law of superposition of strata (principle of superposition)** *\*Strata* are deposited sequentially, so that in an undisturbed sedimentary succession each layer of rock is younger than the layer beneath it. Subsequent earth movements may overturn and invert this sequence. The law was first proposed in the 17th century by Nicolaus *\*Steno*.

**law of the wall** In a *\*turbulent flow*, the average velocity increases logarithmically with distance from a wall or the bed of a river channel.

**laze** See LAVA HAZE.

**Laxfordian** A sub-*\*stage* of the *\*Lewisian*, from about 1600–1100 Ma ago (Van Eysinga, 1975), named from Loch Laxford, north-western Scotland.

**Laxfordian orogeny** A mountain-building episode that occurred about 1800–1600 Ma ago, and which is recorded in north-west trending *\*folds* in the Lewisian *\*gneisses* of what is now the extreme north-west of Scotland. It is a continuation of the *\*Proterozoic \*Ketilidian* and *\*Nagssugtoqidian* orogenies of Greenland, and may be a continuation of the *\*Scourian orogeny* which immediately preceded it.

**layer cloud** One of the principal forms of cloud, with flattened, sheet-like appearance and of limited vertical extent. Common types of cloud exhibiting this form are: (a) low-level layer clouds, e.g. *\*fog* and *\*stratus*; and (b) multi-layered clouds, e.g. *\*altostratus*, *\*cirrostratus*, and *\*nimbostratus*. See also CLOUD CLASSIFICATION.

**Layered Deposits** See MARTIAN TERRAIN UNITS.

**layered silicate** See PHYLLOSILICATE.

**layer-parallel shortening** The *\*homogeneous strain* which is developed in a layered rock when shortening occurs parallel to the layering. Buckling instabilities, which would normally form *\*folds*, are restricted in such a case, and the layer shortens and thickens as a result.

**layover** In *\*radar* terminology, the distortion of a radar image caused by steep surface angles relative to the radar wavefront angle, such that the top of a hill is actually closer to the radar than the base. The resulting image gives the appearance of hills or mountains leaning toward the radar. Compare FORESHORTENING.

**Lazarus taxon** A taxon that disappears from the fossil record close to an extinction horizon, but reappears again much later in the sequence.

**lazurite** *\*Mineral*  $\text{Na}_8[(\text{Si},\text{Al})_6\text{O}_{24}](\text{S},\text{SO}_4)$  belonging to the *\*sodalite* group of minerals and with a similar composition to *\*häüyne* and *\*nosean*; sp. gr. 2.3–2.4; *\*hardness* 5.5; *\*cubic*; intense, deep, azure-blue, violet, light

blue, or greenish-blue; **\*vitreous lustre**; usually compact, **\*massive**; **\*cleavage** imperfect, rhombododecahedral; occurs in the contact zone of **\*alkaline**, **\*igneous** rocks, in association with **\*carbonate** rocks, and in alkaline **\*lavas**. The variety lapis lazuli is used to make ornaments.

**L-band** Radar frequency band between 390 and 1550  $\mu\text{Hz}$ , used for radar scanning of the Earth's surface.

**leachate** Solution formed when water percolates through a permeable medium. In some cases the leachate may be toxic or carry bacteria when derived from solid waste. In mining, leaching of waste tips can produce a mineral-rich leachate which is collected for further processing, as in heap leaching of **\*porphyry** copper and gold deposits.

**leaching** Removal of soil materials in solution.

**lead–lead dating** A dating method based on the comparison of **\*isotopes** of lead (Pb). Throughout geologic time the isotopic composition of **\*common lead** in the Earth has evolved from that of **\*primordial** lead by the addition of **\*radiogenic** leads ( $^{206}\text{Pb}$ ,  $^{207}\text{Pb}$ , and  $^{208}\text{Pb}$ ) derived from uranium and thorium decay (see **DECAY SERIES**).  $^{204}\text{Pb}$  is not derived from any radioactive parent and appears to be a standard against which the other lead values can be compared. It is normally assumed that in any small part of the Earth's **\*crust** and underlying **\*mantle** which, at the time of formation, contained primordial lead together with uranium and thorium, no radiogenic lead could have been present. With the passage of time atoms of radiogenic  $^{206}\text{Pb}$ ,  $^{207}\text{Pb}$ , and  $^{208}\text{Pb}$  gradually replaced uranium and thorium atoms. If, at one instant in time, all the lead in the area under discussion was removed in solution and deposited as a lead ore, then this would preserve a record of the isotopic balance of lead at the time. Given that this ore mineral would not contain any uranium or thorium, it would be preserved as a unique point on the lead growth curve. Using the Holmes–Houtermans model, and plotting  $^{207}\text{Pb}:^{204}\text{Pb}$  against  $^{206}\text{Pb}:^{204}\text{Pb}$ , a series of **\*growth curves** would be obtained based on the different isotopic ratios. These curves can then be used to plot **\*isochrons**, the slope of which determines the age of the particular lead assemblage.

**lead loss** Loss of daughter lead **\*nuclides** during the **\*radioactive** decay of uranium to lead, in which  $^{238}\text{U}$  decays to  $^{206}\text{Pb}$ , and  $^{235}\text{U}$  decays to  $^{207}\text{Pb}$

(see [DECAY SERIES](#)). These lead [\\*isotopes](#), together with that produced by the  $^{232}\text{Th}$  to  $^{208}\text{Pb}$  series, are moderated by the presence of non-[\\*radiogenic](#)  $^{204}\text{Pb}$  in any calculations of age. After correcting for the original lead in any analysis, if the mineral being investigated has remained a closed system, the  $^{235}\text{U}:^{207}\text{Pb}$  and the  $^{238}\text{U}:^{206}\text{Pb}$  ages should be concordant. On a graph of  $^{235}\text{U}:^{207}\text{Pb}$  against  $^{238}\text{U}:^{206}\text{Pb}$  the loci of all [\\*concordant ages](#) define a curve called Concordia (see [CONCORDIA DIAGRAM](#)). If they do not agree, they are discordant and the ratios do not fall on the Concordia curve. Because daughter atoms tend to escape from the system, especially when heated or otherwise disturbed, [\\*discordant ages](#) are generally on the young side. Because  $^{207}\text{Pb}$  and  $^{206}\text{Pb}$  are chemically identical, they are not fractionated by natural processes. As a result of this any lead loss from a mineral is in the same isotopic proportion as that in which it occurs in the mineral. The lead loss would be the same for all parts of the rock body, and the plots of  $^{238}\text{U}:^{206}\text{Pb}$  against  $^{235}\text{U}:^{207}\text{Pb}$  fall on a straight line below the Concordia curve. The two points where this straight line intersects the Concordia curve give the age of the rock (higher value) and the time of the lead loss (lower value). In some systems there is not a single time of lead loss but a more continuous process of [\\*diffusion](#). The lower of the two intersects described above may then be further depressed by this continuous diffusion and the otherwise straight-line relationship may be lost especially at the lower value end.

**Leaf Margin Analysis (LMA)** A technique for estimating mean annual temperatures in the past from the proportion of dicotyledonous plant species present that had untoothed leaf margins, based on the observation of a strong positive relationship between warmth and plants with smooth-edged leaves. In addition to this univariate approach there is also a multivariate technique. See [CLIMATE—LEAF ANALYSIS MULTIVARIATE PROGRAM](#).

**leaky transform fault** A type of [\\*transform fault](#) in which there is production of basaltic [\\*magma](#) along the [\\*fault](#) plane, indicating some separation of the blocks on each side of the fault. Such transform faults do not exactly follow arcs of small circles about the [\\*pole of rotation](#) of the relevant plate pair. An example is the Azores Fracture Zone, which forms the boundary between the [\\*Eurasian](#) and [\\*African](#) Plates from the [\\*Mid-](#)

Atlantic Ridge into the Mediterranean. Leaky transform faults are one kind of *\*combined* plate margin.

**least-work principle** The theory that geomorphological *\*processes* always operate in such a way as to achieve the work that has to be done with a minimum expenditure of energy (and maximum *\*entropy*). This is typically achieved by the adoption of a certain profile or shape (e.g. a river *\*meander* may be that shape best suited for carrying the discharge and sediment with the least loss of energy). See *LEAST-WORK PROFILE*.

**least-work profile** That profile whose gradient is just sufficient for the associated geomorphological *\*process* to occur with the minimum possible expenditure of energy. An example is a long river profile, whose concave-up form is the shape best suited for the transfer of increasing quantities of water and sediment in accordance with the *\*least-work* principle. Such a profile expresses a state of high *\*entropy*.

**lebensspuren** Biologically formed, *\*sedimentary structures* found in *\*sediments*, including *\*tracks*, *\*trails*, *\*burrows*, *\*borings*, faecal casts, and *\*coprolites*.

**lechatelierite** A white, pale yellow, or colourless *\*mineraloid* consisting of fused *\*silica* that solidifies at temperatures above 1750 °C. It forms through *\*meteorite* impact, volcanism, and lightning strikes and occurs in association with shocked *\*quartz*, volcanic rocks, and sandy soil. Hardness 6.5, sp. gr. 2.5–2.7, lustre vitreous, cleavage absent, fracture conchoidal, streak white.

**lectostratotype** A *\*stratotype*, chosen after the original establishment of a *\*stratigraphic unit*, that is designed to serve as the standard in the absence of a satisfactory original stratotype (i.e. a *\*holostratotype*). A lectostratotype may be selected from outside the *\*type area*. Compare *NEOSTRATOTYPE*. See also *PARASTRATOTYPE*; *HYPOSTRATOTYPE*.

**lectotype** One of a collection of *\*syntypes* which, subsequent to publication of the original description, is chosen and designated through published papers to serve as the ‘type specimen’. Compare *HOLOTYPE*; *NEOTYPE*; *PARATYPE*.

**Leda (Jupiter XIII)** One of the lesser satellites of **\*Jupiter**, with a diameter of 10 km.

**lee depression** Non-frontal **\*depression** that develops on the lee side of an upland barrier across the airflow as a result of contraction leading to cyclonic curvature. Dynamic processes are responsible for the low-pressure system rather than wave development along a **\*front**. Such depressions are common, for example in winter on the southern lee side of the Alps.

**lee dune** An elongated **\*sand** **\*dune** that forms on the downwind side of an obstruction.

**lee waves** Air waves in the lee of a mountain barrier, where a stable layer of air, after displacement by movement over the barrier, returns to its original level. This process results in a series of stationary ('standing') waves extending downwind on the lee side of the barrier. Clouds often form along the wave crests in lenticular form: they may appear stationary, due to condensation of water vapour at the upward side, caused by the upward air movement, and evaporation on the downward side of the wave. The wavelength can be up to 40 km and the wave amplitude is most pronounced in the intermediate levels of the airstream. Circular air motion (in the vertical plane) beneath the wave crests may reverse wind direction locally within the general air flow. This phenomenon is termed a 'rotor'. In addition to stable air at an intermediate level, lee-wave formation requires a constant wind of at least 8 m/s. Well-known wave clouds on the lee side of barriers include the Sierra wave of the Sierra Nevada, California, the helm wave of Cumbria, England, and the moazagotl of Silesia.

**left lateral fault** See **SINISTRAL FAULT**.

**Lehmann, Inge** (1888–1993) A Danish geophysicist, Lehmann was Director of the Seismological Department of the Royal Danish Geodetic Institute from 1928 to 1953. In 1936 she was able to demonstrate, from studies of the refraction of **\*P-waves**, that the Earth has a solid inner **\*core**.

**Lehmann, Johann Gottlob** (1719–67) A German mining engineer who moved to Russia, Lehmann worked on the **\*stratigraphy** of mountain rocks, and distinguished between Primary (non-fossiliferous and metal-bearing) and Secondary sediments. His ideas were afterwards extended and developed by **\*Werner**.

**Leighton–Pendexter classification** A classification for *\*limestones* and *\*dolomites* proposed in 1962 by W. M. Leighton and C. Pendexter, now used infrequently. The classification defines *\*carbonate* rocks by the percentage of *\*grains*, *\*micrite*, *\*cement*, and voids present. Limestones with more than 50% grains are named as ‘limestones’, prefixed by the name of the main grain type (e.g. ‘skeletal limestone’, ‘pellet limestone’); limestones with 10–50% grains are termed ‘micritic limestones’, prefixed by the name of the dominant grain type (e.g. ‘skeletal micritic limestone’); limestones with less than 10% grains are termed micritic limestones; limestones built by organic frame-builders are termed ‘coralline limestones’ or ‘algal limestones’ according to the type of frame-builder. Leighton and Pendexter also define carbonates with variable amounts of dolomite, *\*calcite*, and impurities on a triangular classification.

**Leitz–Jelley refractometer** See REFRACTOMETER.


**Lemur-2** A commercial constellation of 150 *\*nanosatellites* that provide ship tracking and weather monitoring over 97% of the Earth. The first of the *\*CubeSats* were launched in 2015. Some of the satellites are deployed from the International Space Station (ISS), others from vehicles visiting the ISS.

**Lenian** A *\*substage* of the Early Cambrian epoch, underlain by the *\*Atdabanian* and overlain by the Middle Cambrian and dated at 524–513 Ma ago (Int. Commission on Stratigraphy, 2004).

**lenticula** (*pl.* lenticulae) A small, dark area on the surface of *\*Europa*.

**lenticulae** See LENTICULA.

**lenticular bedding** A form of *\*heterolithic* *\*sediment* characterized by the presence of isolated *\*sand* *\*ripples* and lenses, set in a mud *\*matrix*. The ripples may be symmetrical or asymmetrical. Such sediments form in low-energy, muddy environments which suffer episodic higher flows, e.g. areas of *\*continental shelf* lying in water depths affected by storm *\*waves*, or the low-energy zones on *\*intertidal* flats.



**Lenticular bedding**

**lenticularis** From the Latin *lenticularis* meaning ‘biconvex’ or ‘lens-shaped’, a form of cloud consisting of clearly defined, elongated lenses. The form is typical of *\*lee-wave* clouds and may affect such clouds as *\*stratocumulus*, *\*altocumulus*, and *\*cirrocumulus*. See also CLOUD CLASSIFICATION.

**Lenz’s law** When a magnetic field and an electrical circuit are moved in relation to one another, an electric current is induced in the circuit such that it forms a magnetic field opposing the motion.

**LEO** See LOW EARTH ORBIT.

**Leonardian** A *\*stage* (280–270.6 Ma ago) in the Early *\*Permian* of N. America, underlain by the *\*Wolfcampian*, overlain by the Middle Permian, overlapping the *\*Artinskian*. and containing the *\*Kungurian* stage. In some areas it is zoned by the use of *\*fusulinid* foraminiferids. It is the N. American equivalent of the *\*Rotliegende*. *\*Red-bed* localities of

**\*Wolfcampian**—Leonardian age in Texas and Colorado have yielded many vertebrate fossils.

**Le Pichon, Xavier** (1937– ) A French marine geologist formerly of the Brittany Oceanographic Centre at Brest, who worked in the USA in the late 1960s and is now a professor at the Collège de France, Le Pichon studied the mechanisms of **\*sea-floor** spreading. He worked out the geometry of **\*plate motions** on a sphere, and was the first to recognize that there are six major tectonic plates.

**lepidoblastic** Applied to a **\*metamorphic rock** that has a flaky texture due to **\*schistosity**.

**lepidocrocite** See GOETHITE.

**Lepidodendron selaginoides** Important species of **\*Palaeozoic** plant, characterized by a **\*dichotomous** branching, by diamond-shaped leaf scars, and by large cones. During the Upper **\*Carboniferous**, *Lepidodendron* species flourished on several continents and the trunks grew up to 30 m before branching. See also LYCOPSIDA.

**lepidolite** Lithium-bearing **\*mica** with composition  $K_2(Li,Al)_{5-6} [Si_{6-7}Al_{2-1}O_{20}] (OH,F)_4$ ; sp. gr. 2.8–2.9; **\*hardness** 2.5–4.0; **\*monoclinic**; **\*crystals** as small flakes; perfect basal {001} **\*cleavage**; lilac to greyish; occurs in late-stage **\*pegmatites** associated with other lithium minerals, such as **\*spodumene**, and minerals of a **\*pneumatolytic** origin, such as **\*tourmaline** and **\*topaz**. Lepidolite is an important **\*ore mineral** of lithium.

**lepidomelane** An iron-rich variety of **\*biotite**.

**Lepidophloios kilpatrickense** The earliest known representative of the important **\*Palaeozoic** family of plants, the Lepidodendraceae. Closely related to *Lepidodendron*, *Lepidophloios kilpatrickense* is distinguished by its internal anatomy. It is recorded from the Upper **\*Carboniferous** of Scotland. See also LYCOPSIDA.

**Lepidosauria** See DIAPSID TYPE.

**lepisphere** A **\*microcrystalline**, blade-shaped crystal of a metastable variety of **\*quartz**, composed of **\*cristobalite** with interlayered lattices of

tridymite, aggregates of which often occur during the transformation of \*opal into quartz \*chert.

**Lepospondyli** Group of small \*Palaeozoic \*amphibians, distinguished by the possession of \*vertebrae with spool-shaped centra below the neural arches. Each centrum (the massive part of the vertebra) was perforated lengthwise, providing a channel for the \*notochord. The lepospondylous, or 'husk', vertebra also occurs in living amphibians, but its evolutionary derivation is unclear.

**leptograptid** A type of Middle to Upper \*Ordovician graptoloid (\*Graptoloidea, suborder Didymograptina) characterized by \*biramous, slender, flexuous \*stipes which are either horizontal or reflexed.

**leptokurtic** See KURTOSIS.

**leptosols** A reference soil group in the \*World Reference Base for Soil Resources classification. Leptosols are weakly developed soils that either have hard rock within 25 cm of the surface, or overlie material with more than 40% calcium carbonate within 25 cm of the surface, or contain less than 10% fine earth to a depth of 75 cm.

**lessivage** The movement of insoluble particles, especially clay particles, down a \*soil profile suspended in a soil \*colloid.

**leste** Regional easterly or south-easterly wind affecting N. Africa and Madeira, which blows ahead of a low-pressure area and brings hot, dry conditions.

**leucite** An important \*feldspathoid mineral  $\text{KAlSi}_2\text{O}_6$ ; sp. gr. 2.5; \*hardness 5.5–6.0; \*tetragonal; normally white or grey; \*vitreous \*lustre; \*crystals form very characteristic tetragonal trioctohedra; \*cleavage imperfect {110}; occurs in potassium-rich rocks which may be silica deficient, such as leucite-basanites, leucite tephrites, lamproites, and leucitophyres or \*leucitites.

**leucite–basanite** See BASANITE.

**leucitite** An \*extrusive, undersaturated (see SILICA SATURATION), \*mafic \*igneous rock consisting of \*essential \*leucite and \*augite, with \*accessory iron oxide and \*apatite. When \*olivine or \*nepheline are also present, the

rock is termed an olivine-leucitite or a nepheline-leucitite respectively. Leucitites are allied to **\*phonolites**, with which they are commonly associated in the field.

**leuco-** A prefix attached to a rock name to indicate a lighter than usual colour for the particular rock type. For example, a **\*gabbro** lighter in colour than a normal gabbro owing to the presence of a larger amount of feldspar than usual would be called a 'leucogabbro'.

**leucocratic** Applied to a rock with a **\*colour index** between 5 and 30.

**leucosome** A coarse-grained, quartzofeldspathic vein, varying in thickness from a few centimetres to a metre or two, and found as a high-grade metamorphic product (see **METAMORPHISM**) in rocks of pelitic (see **PELITE**) to psammitic (see **PSAMMITE**) composition. The finer material between the leucosomes is enriched in **\*mafic** and aluminous minerals. Leucosomes may represent a low-melting-point liquid segregated from the **\*metasediment** during high-grade metamorphism to give rise to what is, in effect, a **\*migmatite fabric**.

**levanter** Local wind from the east experienced in the Straits of Gibraltar and associated with standing waves in the lee of the Rock of Gibraltar. The wind is especially prevalent in late summer and autumn and brings high humidity.

**leveche** Local wind affecting south-eastern Spain, especially in summer, and similar to such other hot, dry, dusty winds of tropical continental origin as the **\*scirocco** and **\*khamsin**, which blow in the Mediterranean region.

**levée** **1.** Raised embankment of a river, showing a gentle slope away from the channel. It results from periodic overbank flooding, when coarser sediment is immediately deposited due to a reduction in velocity. This may lead to a situation in which the river flows well above the level of its outer **\*floodplain**. **2.** See **LAVA LEVE'E**.

**level of compensation** See **PRATT MODEL**.

**Lewisian** A **\*stage** of the **\*Proterozoic** of north-western Scotland, from about 2600–1100 Ma ago, named from the Outer Hebridean Isle of Lewis.

**LHB** See **LATE HEAVY BOMBARDMENT**.

**lherzolite** A two-*\*pyroxene-* and *\*olivine-*bearing, coarse-grained, *\*ultrabasic* rock consisting of essential magnesium-rich olivine, chromium-magnesium *\*clinopyroxene* (Cr-diopside), magnesium *\*orthopyroxene* (*\*enstatite*), and *\*garnet* or *\*spinel*, which is found as *\*xenoliths* in *\*basalts* and as a component of Alpine *\*ultramafic* bodies. A *\*mantle* origin for many lherzolites is attested to by their metamorphic (see *METAMORPHISM*) texture, their high-pressure *\*mineral* assemblage, and their ultrabasic composition. Using evidence from experimental *\*petrology*, lherzolites are interpreted as examples of undepleted mantle (i.e. mantle which has not undergone partial melting to generate basalt *\*magma*). The name is taken from the type locality, at Lherz in the French Pyrenees.

**Lias** Former name for what is now called the Early *\*Jurassic \*epoch*, dated at 199.6–175.6 Ma ago (Int. Commission on Stratigraphy, 2004), and comprises the *\*Hettangian*, *\*Sinemurian*, Lotharingian, *\*Pliensbachian*, and *\*Toarcian \*stages*. Blue-grey *\*shales* and muddy *\*limestones* are typical of the Lias, with Dactyloceras and Gryphaea among the more important *\*fossils* found in the extensive outcrops of England and France.

**Libby, Willard Frank** (1908–80) Professor of chemistry at the University of California and director of the Institute of Geophysics, Libby developed a method for dating *\*sediments* and artefacts using *\*carbon-14*. The method is described in his book *Radiocarbon Dating*, first published in 1952.

**libeccio** Local south-westerly wind which brings stormy conditions, especially in winter, to the central Mediterranean.

**libration** The slow oscillation of a satellite, as seen from the planet around which it orbits. One libration is due to a parallax effect: e.g., because of the rotation of the *\*Earth*, more of the eastern limb of the *\*Moon* is visible at moonrise, while more of the western limb can be seen at moonset. In the Earth–Moon system, a second libration is that of lunar longitude, with a monthly period, since the Moon’s orbit around the Earth sometimes exceeds and sometimes lags behind its axial rotation, and the third lunar libration is of latitude, due to the 6° inclination of the lunar equator to its orbital plane, so that more of the polar regions become visible when the Moon is north or south of the *\*ecliptic*.

**librigena** See *CEPHALON*.

**lichenometry** A technique for dating rock surfaces from measurements of the diameter of lichens growing on them. It is especially useful in environments where trees cannot survive. The first step is to find lichens growing on rock surfaces of a known age (e.g. gravestones or *\*made ground*) and to determine the mean diameter of the five largest lichens growing on them. That provides a lichenometric dating curve against which other exposed surfaces can be dated. An alternative method, useful where there are few surfaces of known age, involves measuring the diameters of selected lichens at intervals over several years to determine their rates of growth. Using very large samples of lichens makes it possible to define confidence limits for the calculated ages.

**Lichida** An order of *\*Trilobita* that lived from the Middle *\*Cambrian* to Middle *\*Devonian*. They were medium-sized to very large, usually spiny, and in many the *\*pygidium* is larger than the *\*cephalon*. There were three suborders.

**life assemblage (biocoenosis)** A *\*fossil* community that is interpreted as representing a former living community. Most assemblages interpreted as life assemblages represent only a small fraction of a former community.

**lifting condensation level** Level at which air becomes saturated when forced to rise. (The level can be determined on a *\*tephigram*, where a dry-*\*adiabat* line intersects a line of saturation mixing ratio through the *\*dew-point* temperature.)

**ligand** Atom, *\*ion*, or molecule that acts as the electron-donor partner in one or more coordination bonds. A heterocyclic ring is formed if the ligand is an organic compound, and the product is termed a chelate. *See also* CHELATION; COORDINATION NUMBER.

**light absorption** Most of the light entering the oceans is absorbed within the first 100 m of water. The depth of light penetration is governed by the amount of material in the water that absorbs and scatters light, e.g. suspended organic material. The absorption of light varies with differing wavelengths of light, blue light penetrating more deeply than red light.

**Lightning Imaging Sensor** *See* LIS.

**lignite** Poor-quality, brown *\*coal* with a visibly woody structure, relatively unaltered, between *\*peat* and *\*bituminous* coal in *\*rank*. It is non-coking,

burns with a long, smoky flame, and contains up to 40% moisture when mined, although this can be reduced to 10–15% by drying in air. It is found in *\*Palaeogene* basins in Britain, and also in continental Europe where it is economically important.

**Likhvin** A series of sediments in European Russia, perhaps dating from the *\*Mindel/Riss* Interglacial, and the equivalent of the *\*Holsteinian* or the *\*Hoxnian*.

**LIL** See *LARGE-ION LITHOPHILE*.

**Lillburnian** A *\*stage* (15–11.5 Ma ago) in the *\*Neogene* of New Zealand, underlain by the *\*Cliffdenian*, overlain by the *\*Waiauan*, and roughly contemporaneous with the *\*Serravallian* stage.

**lime** Compounds, mostly of calcium carbonates but also other basic (alkaline) substances, used to correct soil acidity and occasionally as a fertilizer to supply magnesium.

**lime boundstone** See *DUNHAM CLASSIFICATION*.

**lime grainstone** See *DUNHAM CLASSIFICATION*.

**lime mud** A general term for *\*carbonate \*sediment* composed of particles up to 62 µm in size. Lime muds are found in a wide range of depositional settings, ranging from *\*pelagic* to *\*intertidal*. The mud comes from various sources; it may be derived from microfauna, calcareous *\*algae*, and from the mechanical or biological breakdown of coarser particles or, more problematically, it may be formed by chemical/biochemical precipitation (this is as yet unobserved in modern marine settings).

**lime mudstone** See *DUNHAM CLASSIFICATION*.

**lime packstone** See *DUNHAM CLASSIFICATION*.

**limestone** Sedimentary type of rock composed mainly of *\*calcite* and/or *\*dolomite*, which is often of organic, chemical, or *\*detrital* origin. See *FOLK LIMESTONE CLASSIFICATION*; *DUNHAM CLASSIFICATION*; *LEIGHTON—PENDEXTER CLASSIFICATION*.

**limestone pavement** See *CLINT*.

**lime wackestone** See DUNHAM CLASSIFICATION.

**limit-equilibrium analysis** In rock and soil mechanics, the study of the point at which a material has reached the limit of its stability using the concept of yield criteria and the associated flow rule in the stress–strain relationship, e.g. where toppling occurs in a steeply jointed rock slope, where a soil becomes plastic, etc.

**limnic** Pertaining to fresh water.

**limnic eruption (lake overturn)** A rare event in which a large amount of gas (usually carbon dioxide but also possibly methane) erupts from deep below the surface of a lake. Being heavier than air, the CO<sub>2</sub> may then spread at ground level across the land surrounding the lake, causing widespread poisoning. Two limnic eruptions have been recorded, at Lake Monoun, Cameroon, in 1984, killing 37 persons, and at Lake Nyos, Cameroon, in 1986, when the release of more than 80 million m<sup>3</sup> of CO<sub>2</sub> killed 1700–1800 people. For a limnic eruption to occur the lake water must be saturated with gas. The deep water is held under greater pressure than the overlying water and is also cooler. Any event (e.g. landslide or earthquake) that displaces some of the deep water, reducing the pressure, or raising its temperature, reducing the solubility of CO<sub>2</sub> or CH<sub>4</sub>, may trigger a release of gas.

**limnology** The study of all aspects of inland waters.



<http://www.limnology.org/>

- The International Society of Limnology.

**limonite** Mineral, FeO(OH).nH<sub>2</sub>O; sp. gr. 2.7–4.3; \*hardness 4.0–5.3; yellowish-brown to reddish-brown; normally \*earthy \*lustre; usually amorphous; occurs as a \*secondary mineral from the weathering of iron in rocks and mineral deposits, and may accumulate to give iron-rich mineral deposits.

**limpet** See ARCHAEOGASTROPODA; ‘PATELLA’ BEACH.

**Limulus** See CHELICERATA.

**Lindgren, Waldemar** (1860–1936) A Swedish-born officer of the US Geological Survey, and later professor at the Massachusetts Institute of Technology, Lindgren is best known for his classification of ore deposits according to their genesis (1913). He was the first to propose the hydrothermal theory of ore deposition.

**linea** (*pl.* lineae) A linear feature, which may be dark or light, straight or curved, on the surface of an extraterrestrial body.

**lineae** See **LINEA**.

**lineage zone (phylozone, evolutionary zone, morphogenetic zone)** A unit of *\*strata* containing a clearly defined portion of an evolutionary lineage, marked above and below by some distinct and specified change in form.

**lineament** A linear landscape feature that is distinctly different from other features nearby and that reflects an underlying geological structure.

**linear regression** In statistics, a comparison of two sets of data to see if there is a linear relationship.

**linear sand ridge** Submarine sand mound typical of shallow sea and wide *\*continental-shelf* areas, 3–10 m high, 1–2 km wide, which may extend for tens of kilometres across the shelf and have an average spacing of about 3 km. Such ridges have been described from the North Sea (off Norfolk, England) and the eastern seaboard of the USA. They are the product of storm and tidal action.

**lineation** 1. Any linear feature that appears on the surface of a rock. Lineation may be formed during deformation by the parallel alignment of minerals, fossils, or pebbles; by parallel crenulation *\*cleavages*; or by striations and grooves resulting from the movement of a rock over a plane, e.g. a *\*fault* surface (see **SLICKENSIDE**), or *\*flexural slip* during folding. An intersection lineation is caused by the crossing of any two planes, e.g. cleavage and *\*bedding*. 2. Lineations are a series of parallel lines on a rock surface, formed by tectonic processes, by the transportation and deposition of sand under upper-flow-regime plane-bed conditions, or by the movement of glacial ice over the rock surface.


**line scanner** In *\*radar* terminology, an imaging device which uses photoelectronic detectors combined with a rotating mirror which sweeps

across the ground surface taking in image data a line at a time in order to produce a *\*raster*.

**line spectrum** See SPECTRUM.

**line squall** An individual *\*squall* that occurs on a *\*squall line*. The squall is caused by a large, vigorous *\*cumulonimbus* cloud; as one cloud dissipates, cold air emerging at the base from downdraughts inside the cloud moves beneath adjacent warm, moist air, lifting it and thus triggering the development of another storm cloud.

**Lingulata** (phylum *\*Brachiopoda*) A class of brachiopods, existing from the Lower *\*Cambrian* to the present day, in which the shell is chitinophosphatic, the valves are not hinged by teeth and sockets, and the *\*pedicle* is thick.



Lingulata

**Lingulella viridis** One of the earliest recorded *\*inarticulate* brachiopods, and very similar to extant species of the genus *Lingula*, *L. viridis* is described from the lower *\*Cambrian* of England. It is one of several species referred to the cosmopolitan genus *Lingulella*. *Lingula* has been cited as an example of *\*bradytelic* evolution.

**Lingulida** (class *\*Inarticulata*) Order of *\*Brachiopoda*, with *\*valves* usually of calcium phosphate with some layers of organic material. The valves may be finely *\*punctate* or *\*impunctate*. The *\*pedicle* emerges between the valves posteriorly. Lingulida are usually marine, but some are tolerant of reduced salinity. They appeared first in the Lower *\*Cambrian*. There are two superfamilies: the Lingulaceae (which includes *\*Lingulella* and the extant *Lingula*); and the Trimerellaceae, ranging from the *\*Ordovician* to the *\*Silurian*, with a biconvex, probably aragonitic shell, with internal muscle platforms (e.g. *Trimerella*, *Dinobolus*).

**linguoid** Applied to tongue-shaped asymmetrical *\*ripples* which have a highly sinuous crest and a strong three-dimensional form. *See also* AKLÜ.

**liptinite** *See* COAL-MACERAL GROUP.

**liquation** A form of *\*magmatic* differentiation in which a sulphide liquid separates from a mixed *\*sulphide*–silicate *\*magma* and sulphide droplets sink to the bottom of the *\*intrusion* or *\*lava* flow, where they accumulate. The sulphide liquids have a strong affinity for nickel, copper, iron, and platinum-group elements. Consequently, liquation produces valuable *\*orebodies*.

**liquefaction** The process of becoming or making a liquid by heating, cooling, or a change in pressure. In soils, the temporary transformation of material to a fluid state due to a sudden decrease in shearing resistance caused by a collapse of the structure associated with a temporary increase in *\*pore* fluid pressure.


**liquid limit** *See* ATTERBERG LIMITS.

**liquidus** In a *\*temperature*–*composition* diagram, the liquidus is the locus of points marking the boundary above which the *\*phases* are all liquid. In a *\*binary* system it is represented by a line, in a *\*ternary* system by a curved

surface, and in a *\*quaternary system* by a volume. Between the liquidus and the *\*solidus*, both liquid and solid phases are present. See PHASE DIAGRAM.

**LIS (Lightning Imaging Sensor)** An instrument that observes lightning. It was launched on 19 February 2017 to the International Space Station.

**listric fault (lystric fault)** A curved extensional *\*fault* which characteristically flattens at depth into a *\*décollement* horizon. The *\*hanging wall* rotates down along the concave-up fault surface, in some cases forming a *\*rollover* anticlinal structure. Listric faults are commonly found in extensional regimes (see EXTENSION).



**Listric fault**

**lithic arenite** A *\*sandstone* containing less than 15% mud *\*matrix*, and with a *\*grain* composition comprising more than 25% rock fragments, and more rock fragments than *\*feldspar* present. See DOTT CLASSIFICATION.

**lithic fragment** The dense or crystalline components of a *\*pyroclastic* deposit. Three types are recognized. Cognate lithics are fragments of non-vesiculated, juvenile, magmatic material, e.g. dense, angular, glass fragments. Accessory lithics are fragments of *\*country* rocks ejected explosively during *\*eruption*. Accidental lithics are *\*clasts* picked up locally by pyroclastic flows and surges. Lithics range from blocks to *\*ash*-size fragments and are usually angular, but may be rounded by in-vent abrasion during eruption.

**lithic greywacke (lithic wacke)** A \*sandstone containing more than 15% but less than 75%, mud \*matrix, and with a \*grain composition comprising more than 5% rock fragments and with more rock fragments than \*feldspar present. See DOTT CLASSIFICATION.

**lithic wacke** See LITHIC GREYWACKE.

**lithification** The process of changing unconsolidated \*sediment into rock. This involves cementation (see CEMENT) of the \*grains, but not necessarily burial alteration or \*compaction.

**litho-** From the Greek *lithos* meaning ‘stone’, a prefix meaning ‘pertaining to rock or stone’.

**lithoclast** A sedimentary \*clast composed of a pre-existing rock type.

**lithofacies** Rock noted for a distinctive group of characteristics, e.g. composition and grain size.

**lithofacies map** A map which shows the distribution of different \*lithofacies for a particular stratigraphic interval or horizon (see STRATIGRAPHIC COLUMN). Such maps may be interpreted to yield a \*palaeogeography or environmental interpretation for the mapped horizon.

**lithologic symbol** A mark, or set of marks, representing a particular textural feature or rock type. For example, conventionally a sandstone is represented by stipple, limestone by a ‘brick’ emblem. However, as there is not a complete range of generally accepted lithologic symbols, where they are used (e.g. in \*columnar sections) it is customary to give a key-panel explaining the meaning of the symbols. Symbols may be idiosyncratic to particular authors but they are often standardized, especially within large oil companies.

**lithologic trap** See STRATIGRAPHIC TRAP.

**lithology** The description of the macroscopic features of a rock, e.g. its \*texture or \*petrology.

**lithology classification** The identification of rocks in the field, based on their texture, structure, composition, colour, and grain size, and their allocation into types.



<http://civilblog.org/2015/02/11/how-to-classify-rocks-on-site-geological-or-lithological-classification/>

- How to classify rocks on site? (Geological or lithological classification).

**lithomarge** A soft \*saprolite rich in \*kaolinite.

**lithomorph soil** A major \*soil group in the soil classification developed by the Soil Survey for England and Wales. A lithomorph soil has a shallow \*soil profile, with organic \*soil horizons directly overlying bedrock.

**lithophile** Applied to elements with a strong affinity for oxygen and which concentrate in the Earth's \*crust in \*silicate rather than metal or \*sulphide minerals; or to elements with greater free energy of oxidation per gram of oxygen than iron. They occur as oxides, and especially in the \*silicate minerals which make up 99% of the crust. Examples of lithophile elements are Al, Ti, Ba, Na, K, Mn, Fe, Ca, and Mg. Compare **ATMOPHILE**; **BIOPHILE**; **CHALCOPHILE**; **SIDEROPHILE**.

**lithophysae** Concentric shells of \*aphanitic material enclosing a hollow core, which form rounded masses a few centimetres in diameter within glassy or aphanitic, \*felsic \*igneous rocks. The hollow, lithophysal core is sometimes filled by secondary \*silica to form a \*geode.

**lithosequence** A sequence of soils where the changing character of the soil is related to, or caused by, the changing \*lithology of \*rocks and superficial mineral deposits.

**lithosome** A mass of rock of approximately uniform character penetrated by tongues of rock from adjacent masses with a different \*lithology.

**lithosphere** The upper (oceanic and continental) layer of the solid \*Earth, comprising all crustal rocks and the brittle part of the uppermost \*mantle. It is generally considered to deform by brittle fracture and if subjected to stresses of the order of 100 MPa. It comprises numerous blocks, known as tectonic \*plates, which have differential motions giving rise to \*plate tectonics. The concept was originally based on the requirement for a rigid upper layer to account for \*isostasy. Its rigidity is variable, but much

greater than  $10^{21}$  P, which corresponds with the underlying **\*asthenosphere**. Its thickness is variable, from 1–2 km at mid-oceanic **\*ridge** crests, but generally increasing from 60 km near the ridge to 120–140 km beneath older **\*oceanic** crust. The thickness beneath **\*continental crust** is uncertain, probably some 300 km beneath the cratonic (see **CRATON**) parts of the continental crust, but the absence of the asthenosphere in these regions makes definition difficult. Compare **ATMOSPHERE**; **HYDROSPHERE**.

**lithospheric plate** See **PLATE**.

**lithostatic stress (geostatic stress)** The component of **\*confining** pressure derived from the weight of the column of rock above a specified level.

**lithostratigraphic unit (rock unit, rock-stratigraphic unit)** A body of rock forming a discrete and recognizable unit, of reasonable homogeneity, defined solely on the basis of its lithological characteristics (see **LITHOLOGY**). A lithological unit may be **\*sedimentary**, **\*igneous**, **\*metamorphic**, or a combination of these. As with other **\*stratigraphic units**, lithostratigraphic units are defined according to **\*type** sections. Their boundaries are placed at surfaces of lithologic change, usually sudden but sometimes gradational. As the physical nature of the units reflects depositional environments rather than time spans, the boundaries of lithological units may be **\*diachronous**. Lithostratigraphic units are comparatively local in extent when compared to the worldwide compass of **\*chronostratigraphic units**. They are ranked in decreasing order of magnitude in **\*supergroups**, **\*groups**, **\*formations**, **\*members**, and **\*beds**. A diverse, but distinctive and interrelated body of rock that cannot be subdivided into any other lithostratigraphic unit is termed a ‘complex’. See also **BIOSTRATIGRAPHIC UNIT**; **CHRONOSTRATIGRAPHIC UNIT**; **STRATIGRAPHIC UNIT**.

**lithostratigraphy** Branch of stratigraphy concerned with the description of rock units in terms of their lithological features. It deals with the spatial relations of such rock units, but does not take into consideration (a) the evolution of the organisms contained within the units (**\*biostratigraphy**), or (b) geologic time (**\*chronostratigraphy**).

**Lithothamnion** See **RHODOPHYCEAE**.

**litter (L-layer)** Accumulation of dead plant remains on the soil surface.

**Little Ice Age** A period between about 1350 and 1860 during which the global climate became generally harsher and there was a worldwide expansion of *\*glaciers*. The effects have been recorded in the Alps, Norway, and Iceland, where farm land and buildings were destroyed, and in many other parts of both hemispheres. There were times of especial severity during this period, e.g. the early 1600s when glaciers were particularly active in the Chamonix valley, in the French Alps.

**littoral drift** See BEACH DRIFT; LONGSHORE DRIFT.

**littoral zone** 1. The area in shallow fresh water and around lake shores where light penetration extends to the bottom sediments, giving a zone colonized by rooted plants. 2. In marine ecosystems the shore-area or intertidal zone where periodic exposure and submersion by tides is normal. Since the precise physical limits of tidal range vary constantly a biological definition of this zone, which essentially reflects typical physical conditions rather than rarely experienced events, is generally more useful. Thus in Britain the littoral zone is defined as the region between the upper limit of species of the seaweed *Laminaria*, and the upper limit of *Littorina* species (periwinkles), or of the lichen *Verrucaria*.

**living fossil** An organism that has persisted, essentially unchanged, since its first appearance. For example, *Lingula* (a *\*brachiopod*) has remained much the same since *\*Ordovician* times and *Sphenodon* (tuatara) since the early *\*Mesozoic*.

**lixisols** A reference soil group in the *\*World Reference Base for Soil Resources* classification. A lixosol is any soil that has an *\*argic B* *\*soil horizon* within 100–200 cm of the surface, other than *\*albeluvisols*, *\*alisols*, *\*acrisols*, and *\*luvisols*.

**lizardite** See SERPENTINE.

**Llandeilo** A *\*stage* (464–460.9 Ma ago) of the Middle *\*Ordovician*, underlain by the *\*Llanvirn* and overlain by the *\*Caradoc*.

**Llandovery** A sub-*\*epoch* (443.7–428.2 Ma ago) of the Early *\*Silurian*, underlain by the *\*Ashgill* *\*stage* (Late *\*Ordovician*) and overlain by the *\*Wenlock* sub-epoch.

**Llanvirn** A *\*stage* (471.8–464 Ma ago) of the Middle *\*Ordovician*, underlain by the *\*Arenig* and overlain by the *\*Llandeilo*.

**L-layer** See LITTER.

**LLSVP** See LARGE LOW SHEAR VELOCITY PROVINCE.

**LMA** See LEAF MARGIN ANALYSIS.

**load** Total amount of material carried by a stream or river, or the mass of rock overlying a geologic structure.

**load cast** A bulbous depression formed on the base of a bed of *\*sediment*. Load casts develop by the differential sinking of the sediment, while still soft, into less dense sediment below. Load ‘casts’ are not strictly casts, as they do not infill an existing depression as in the case of flute casts (see FLUTE MARK).

**loadstone** See MAGNETITE.

**loam** Class of soil texture composed of *\*sand*, *\*silt*, and *\*clay*, which produces a physical property intermediate between the extremes of the three components.

**lobe** See SUTURE.

**lobe fins** Paired fins, characteristic of *\*Crossopterygii*, that are borne on fleshy lobes containing an axial skeleton.

**lobsters** See MALACOSTRACA.

**local range zone** See TEILZONE.

**local wind** Air movements, generally of limited geographical range, characteristic of a particular region and/or area of particular land and atmospheric configuration. Many local winds are especially associated with orographic peculiarities or with particular high- or low-pressure systems, and local names may be applied to winds of a broad, general type. Examples include the *\*föhn winds*, of orographic origin, and the many regional winds of the Mediterranean area, whose origins lie in the tropical continental *\*air* masses of the Sahara.

**lochan** See KNOCK AND LOCHAN.

**Lochkovian (Gedinnian)** The earliest *\*stage* in the *\*Devonian* period, preceded by the Late *\*Silurian*, followed by the *\*Praghian*, and dated at 416–411.2 Ma ago (Int. Commission on Stratigraphy, 2004).

**Loch Lomond stadial** Relatively cold period that occurred towards the end of the last (*\*Devensian*) glacial stage in Scotland. The event took place about 11 000–10 000 radiocarbon years BP. It is characterized by the development of small *\*ice* caps and *\*cirque glaciers* in the Highlands.

**Lockportian** See HOMERIAN.

**lode** Mineral vein or system of veins; in Cornwall (UK) especially, refers to productive veins only.

**lodestone** See MAGNETITE.

**lodgement till** See TILL.

**loess** *\*Unconsolidated*, wind-deposited sediment composed largely of *\*silt*-sized *\*quartz* particles (0.015–0.05 mm diameter) and showing little or no stratification. It occurs widely in the central USA, northern Europe, Russia, China, and Argentina. It can give rise to a rugged topography with steep slopes (up to 70°).

**logging** See WELL LOGGING.

**log-normal distribution (geometric distribution)** A distribution in which the logarithms of the values have a *\*Gaussian* (normal) distribution. This distribution is common in geologic contexts, e.g. grain sizes.

**logs** Sedimentological logs are vertically measured records of sedimentary successions, illustrating with symbols the vertical sequence of *\*lithology*, grain size, *\*sedimentary structure*, and *\*fossil* content.

**Longfordian** A *\*stage* (27.5–16.5 Ma ago) in the *\*Cenozoic \*era* of southeastern Australia, underlain by the *\*Janjukian (\*Oligocene)*, overlain by the *\*Batesfordian*, and containing the *\*Aquitanian* stage.

**longiconic** Applied to a cephalopod (*\*Cephalopoda*) shell that is very elongate.

***Longisquama insignis*** One of the first *\*archosaurs* known to have the ability to glide or parachute. Described from Soviet Kirgizstan in the 1970s,

Longisquama is noted for the development of elongate paired scales along its back; its Latin name means 'remarkable long scale'. It is recorded from sediments of *\*Triassic* age.

**longitudinal conductance (S)** The sum of all the thickness/resistivity ratios of  $n - 1$  layers which overlie a semi-infinite substratum of resistivity  $\rho_n$ , such that  $S \times h_1/\rho_1 + h_2/\rho_2 + h_3/\rho_3 + \dots + h_{n-1}/\rho_{n-1}$  (mho), where  $h_1, h_2$ , etc. are the depths and  $\rho_1, \rho_2$ , etc. the resistivities, of successive layers. A knowledge of  $h_i/\rho_i$  for the  $i$ th layer when it is sandwiched between two layers of much higher resistivity is of importance in resolving the problem of *\*equivalence*.

**longitudinal dispersion** See DISPERSION.

**longitudinal-type coast** See PACIFIC-TYPE COAST.

**Longmyndian** A *\*stage* of the Upper *\*Proterozoic* of the Welsh borders, underlain by the *\*Charnian* and overlain by the *\*Moinian*.

**long-range forecasting** Weather forecasting for periods beyond a few days.

**longshore bar** Linear ridge of sand whose long axis is parallel to the shore and that is in, or immediately seaward of, the intertidal zone. See also RIDGE AND RUNNEL.

**longshore current** Current that flows parallel to the shore within the zone of breaking waves: it is generated by the oblique approach of waves.

**longshore drift (littoral drift)** Movement of sand and shingle along the shore. It takes place in two zones. *\*Beach drift* occurs at the upper limit of wave activity, and results from the combined effect of *\*swash* and *\*backwash* when waves approach at an angle. Movement also occurs in the *\*breaker* zone, where currents transport material thrown into suspension.

**Longtanian** See WUCHIAPINGIAN.

**Longvillian** A *\*stage* (457–455 Ma ago) of the *\*Ordovician* in the Middle *\*Caradoc*, underlain by the *\*Soudleyan* and overlain by the *\*Marshbrookian*.

**Longwangmiaoan** A Chinese *\*stage* (518–513 Ma ago) of the Early *\*Cambrian* *\*epoch*, preceded by the *\*Changlangpuan* and followed by the

**\*Maozhangian.**

**long wave** Meander in the flow of the circumpolar upper westerly winds, commonly with a wavelength of about 2000 km and sometimes large amplitude. *See also* [DISH-PAN EXPERIMENT](#); [ROSSBY WAVES](#).

**Lonsdale, William** (1794–1871) A curator and librarian of the [\\*Geological Society of London](#), who studied fossils, and especially [\\*corals](#) and [\\*oolitic limestone](#). Lonsdale is credited with using his knowledge of fossils to demonstrate that sediments in Devon were of the same age as the [\\*Old Red Sandstone](#) (1837), and thus helping to establish the [\\*Devonian](#) system.


**look angle** In [\\*radar](#) terminology, the angle between the vertical plane passing through the radar antenna and the line between the antenna and object. *Compare* [DEPRESSION ANGLE](#).

**looping** A geophysical field method in which a [\\*base station](#) is visited regularly because the survey is designed in loops focused through it. This allows instrumental [\\*drift](#) to be measured, especially in gravity and [\\*geomagnetic](#) investigations, and misties and misclosures to be checked for in seismic work.

**lophodont** Applied to cheek teeth ([\\*molars](#)), found in some [\\*Mammalia](#), in which the cusps are fused to form transverse ridges (lophs) that aid the mastication of plant material. *See also* [BILOPHODONTY](#).

**lophophore** *See* [BRACHIOPODA](#); [BRYOZOA](#); [PTEROBRANCHIA](#).

**lopolith** [\\*Concordant](#) [\\*igneous](#) [\\*intrusion](#) that has a sagging, saucer-like form. The shape of small lopoliths may be controlled by their emplacement in folds, but there is no such obvious control for great lopolithic intrusions. *Compare* [LACCOLITH](#).



## Lopolith

**Lopstedt** See BRØRUP.

**Los Angeles abrasion test** A method for measuring abrasion resistance in which the sample and a set of steel spheres are tumbled inside a closed, hollow, steel cylinder, about 700 mm diameter and 500 mm long, which is rotated on a horizontal axis.

**losing stream (influent stream, disappearing stream)** A stream with a permeable bed through which water can seep to the *\*water-table*. See also INTERMITTENT STREAM.

**Lost City Hydrothermal Field** A field of alkaline *\*hydrothermal vents*, more than 120 000 years old, located on a dropped-down bench 70 m below the southern wall of the summit of the *\*Atlantis* Massif and comprising about thirty carbonate chimneys 30–60 m tall. Hydrothermal activity occurs at Poseidon, a chimney 60 m high and 100 m across, where fluid is vented at up to 91°C and pH 9–11. The field supports a variety of small invertebrates. The field was discovered in 2000.



<https://agupubs.onlinelibrary.wiley.com/doi/full/10.1002/2015GC005869>

- Geological Evolution of the Lost City Hydrothermal Field.

**Love, Augustus Edward Hugh** (1853–1940) An applied mathematician and physicist, Love worked at the University of Oxford. His studies of the

theory of elasticity led him to suggest that two types of surface *\*earthquake* waves should exist, one of which has been named for him. *See* [LOVE WAVE](#).

**Love wave (SH-wave, Q-wave, L-wave, L<sub>Q</sub>-wave, G-wave)** A type of *\*surface wave* which occurs when the shear-body-wave velocity in the surface medium is lower than that in the underlying strata. Love waves are characterized by horizontal motion normal to the direction of travel, with no vertical motion. In effect a Love wave is a polarized *\*shear* wave and travels slightly faster than a *\*Rayleigh wave*.

**low** Common term for a low-pressure system, i.e. a depression.

**low-angle fault** A fault which *\*dips* less than 45°. If the sense of displacement is normal (*see* [NORMAL FAULT](#)) it is called a *\*thrust*, if reverse (*see* [REVERSE FAULT](#)) it is called a *\*lag*.

**low-aspect-ratio ignimbrite (LARI)** An *\*ignimbrite* sheet that displays a value for the ratio of its average thickness ( $V$ ) to its horizontal extent ( $H$ ) of  $10^{-4}$  to  $10^{-5}$ , where  $H$  is taken as the diameter of a circle whose area is equal to that of the flow. *Compare* [HIGH-ASPECT-RATIO IGNIMBRITE](#).


**low Earth orbit (LEO)** An orbit at an altitude of 2000 km or lower. *Compare* [GEOSTATIONARY ORBIT](#); [MEDIUM EARTH ORBIT](#).

**low-level waste** *See* [RADIOACTIVE WASTE](#).

**low-potassium tholeiite** *See* [MID-OCEAN-RIDGE BASALT](#).

**lowstand** A time during which sea levels are at their lowest. *Compare* [HIGHSTAND](#).

**lowstand systems tract (LST)** In the *\*genetic* stratigraphic sequence model used in *\*sequence* stratigraphy, a bounding surface formed either from *\*allochthonous* deposits eroded from the platform margin and slope, or from *\*autochthonous* wedges deposited on the upper slope. *See* [SYSTEMS TRACT](#). *Compare* [HIGHSTAND SYSTEMS TRACT](#); [TRANSGRESSIVE SYSTEMS TRACT](#). *See also* [REGRESSIVE SYSTEMS TRACT](#).



**Lowstand systems tract**

**low symmetry** See CRYSTAL SYMMETRY.

**low-velocity layer** See ELEVATION CORRECTION.

**low-velocity zone (LVZ)** The zone within the upper *\*mantle* beneath the oceans within which seismic *\*P-waves* are slowed and *\*S-waves* are slowed and partially absorbed. The top of the zone is some 40–60 km deep near the oceanic spreading *\*ridges*, and this depth increases to 120–160 km beneath the older *\*oceanic* crust. The bottom of the zone is poorly defined, but in the region of 250–300 km in depth. Beneath the continents, a restricted low-velocity zone occurs beneath *\*crust* areas subjected to *\*orogenesis* during the last 600 million years or so, but is not found beneath cratonic (see CRATON) areas. It is attributed to the presence of a 0.1% fluid phase and commonly ascribed to the partial melting of mantle rocks at these depths. It is often considered coincident with the *\*asthenosphere*, but probably this is valid only for oceanic areas.

**Lq** See DIURNAL VARIATION (2).

**L<sub>Q</sub>-wave** See LOVE WAVE.

**LREE** Abbreviation for ‘light *\*rare-earth* element’. See MID-OCEAN-RIDGE BASALT.

**LST** See LOWSTAND SYSTEMS TRACT.

**L–S-tectonite** See SHAPE FABRIC.

**L-tectonite** See SHAPE FABRIC.

**lucinoid** Applied to one of the two types of **\*hinge-line** dentition found in **\*heterodont** bivalves (**\*Bivalvia**). The lucinoid type has two **\*cardinal** teeth in each valve and may have more lateral teeth in the right valve than in the left. Compare CYRENOID.

**Ludfordian** A **\*stage** (421.3–418.7 Ma ago) of the Late **\*Silurian** **\*epoch**, underlain by the **\*Gorstian** and followed by the **\*Pridoli**.

**Ludhamian** The Lower **\*Pleistocene**, lowermost, temperate-climate, shelly, sand deposit of the tripartite division of the deposits in a **\*borehole** at Ludham, in eastern England. See also ANTIAN; BAVENTIAN; PASTONIAN; THURNIAN.

**Ludlow** A sub-**\*epoch** (422.9–418.7 Ma ago) of the Late **\*Silurian**, underlain by the **\*Wenlock** and overlain by the **\*Pridoli**.

**Luisian** A **\*stage** (15.5–13.5 Ma ago) in the **\*Miocene** of California, underlain by the **\*Relizian**, overlain by the **\*Mohnian**, and roughly contemporaneous with the upper **\*Langhian** and lower **\*Serravallian** stages.

**lumbar vertebra** See VERTEBRA.

**luminance** A measure of the **\*intensity** of **\*electromagnetic** radiation emitted by a source in a specified direction.

**luminous night clouds** See NOCTILUCENT CLOUDS.

**lumps, carbonate** See INTRACLAST.

**Luna** A series of Soviet lunar missions that ran from 1959 to 1976.



<http://www.lpi.usra.edu/lunar/missions/luna/>

- An RSA mission to collect information about the Moon.

**lunar** From the Latin *luna* (Moon), of, pertaining to, affecting, caused by, or involving the Moon.

**Lunar-A** A Japanese **\*ISAS** mission, to be launched in 2004, comprising an orbiter and an instrument package to be delivered to the surface. After

many delays, the mission was cancelled in January 2007.

**Lunar Highlands** See TERRA.

**lunar magnetic variation** See DIURNAL VARIATION (2).

**Lunar Orbiter** A series of \*NASA lunar mapping missions that ran from 1966–1967.



<http://nssdc.gsfc.nasa.gov/planetary/lunar/lunarorb.html>

- A NASA mission to map the Moon prior to the Apollo manned landing in 1969.

**lunar timescale** See TIMESCALES.

**lunate** Half-moon shaped.

**lunette (clay dune)** An accumulation of \*aeolian sediment, consisting of clay pellets the size of sand grains, found around the margins of some salt lakes.

**lunule** 1. In \*Bivalvia, a depressed plane or curved area along the \*hinge line, anterior to the \*umbo. Compare ESCUTCHEON. 2. ‘Keyhole-like’ perforations in the \*tests of many flat clypeasteroid (sand dollar) \*Echinoidea.

**lustre** Of \*minerals, the ability to reflect light. The quality and nature of the reflection may be a useful diagnostic aid. The term ‘lustre’ is often used in conjunction with the qualifying terms ‘\*adamantine’, ‘\*metallic’, ‘\*resinous’, ‘\*waxy’, ‘\*pearly’, ‘\*silky’, ‘\*greasy’, and ‘\*vitreous’. The intensity of the reflection is qualified by descriptions such as ‘shiny’, ‘dull’, or ‘\*splendent’.

**lustre mottling** A spotted, shimmering appearance seen on the surface of some calcareous \*sandstones, caused by the reflection of light from coarse mineral \*cement crystals which enclose \*detrital \*quartz grains.

**Lutetian** A \*stage (48.6–40.4 Ma ago) in the Middle \*Eocene sub-epoch, preceded by the \*Ypresian (Early Eocene), followed by the \*Bartonian (Int. Commission on Stratigraphy, 2004). It is roughly contemporaneous with the upper \*Ulatian and lower \*Narizian (N. America), the upper

\*Heretaungan and lower \*Bortonian (New Zealand), and the upper \*Johannian and lower \*Aldingan (Australia).

**lutite** See ARGILLITE.

**luvisols** A reference soil group in the \*World Reference Base for Soil Resources classification. Luvisols have an argic B horizon (see ARGIC HORIZON) with a \*cation-exchange capacity of more than 24 cmol<sub>c</sub>/kg, with illuvial (see ILLUVIATION) accumulations of \*clay.

**luxullianite** See TOURMALINE; TOURMALINIZATION.

**LVL** See ELEVATION CORRECTION.

**LVZ** See LOW-VELOCITY ZONE.

**L-wave** See LOVE WAVE.

**lycopods** See LYCOPSIDA.

**Lycopsidea (lycopods)** Class of the \*Pteridophytina, represented today by relatively small plants, the club mosses, although in the \*Carboniferous they included large trees which dominated the coal-swamp vegetation of the time. The oldest lycopods date from the \*Devonian and derived from the most primitive of vascular plants, the \*psilophytes. Lycopods are characterized by \*dichotomizing branches; spores borne in spore cases (sporangia) on the upper side of the fertile leaves (sporophylls), which are sometimes organized into cones; and 'microphyllous' leaves, i.e. with a single strand of vascular tissue proceeding unbrokenly (without a 'leaf gap') from the vascular tissue of the stem. The leaves of some lycopods were long and grass like. When they were shed, the leaves left distinctive, spirally arranged leaf scars on the trunk of the lycopod tree. Fragments of the bark are common as fossils in coal balls and the roof shales of Carboniferous coal seams. Prominent genera included \*Lepidodendron, \*Lepidophloios, and Sigillaria (in which the spiral leaf-scar pattern fell into discrete vertical rows).

**Lyell, Charles** (1797–1875) Author of the influential *Principles of Geology*, 12 editions of which were published between 1830 and 1875. He was an extreme uniformitarian, emphasizing a great age for the Earth. He

thus influenced **\*Darwin** by allowing the time needed for evolutionary changes to occur. See **UNIFORMITARIANISM**; **DRIFT**.

**lysimeter (evaporimeter)** Device for the direct estimation of **\*evapotranspiration**. Typically it comprises a vegetated block of soil of volume 0.5–1 m<sup>3</sup>, to which the amount of water added is known, and from which the amount lost as runoff or percolation may be measured. Recording the changing weight of the soil-vegetation system (keeping vegetation change due to growth static or monitored) reveals the amount of water retained by the system, and thus, by difference, the amount lost as evapotranspiration. For geographic comparisons an easily standardized, short, grass vegetation cover is used. For water-budget experiments, vegetation cover may be varied to simulate different crop types or semi-natural communities.

**Lysithea (Jupiter X)** One of the lesser satellites of **\*Jupiter**, with a diameter of 24 km.

**lysocline** The ocean depth at which the rate at which calcium carbonate dissolves increases markedly; this may mark the upper boundary of the bottom water.

**lystric fault** See **LISTRIC FAULT**.

# M

**m-** See **MILLI-**.

**M-** See **MEGA-**.

**M** See **MAGNETIC MOMENT**.

**Ma** Abbreviation meaning ‘million years’.

**maar** \*Crater, often occupied by a shallow lake, produced by an explosive volcanic \*eruption. Normally it is surrounded by a low rampart or ring of ejected material. Typically, maars are formed by the explosive interaction of volcanic \*magma with \*groundwater. The ejected material is a mixture of \*country rock and highly fragmented \*ash.

**Maastrichtian (Maestrichtian)** Final \*stage (70.6–65.5 Ma ago) of the \*Cretaceous in Europe, preceded by the \*Campanian and followed by the \*Danian. The \*stratotype is described from the Maastricht area of the Netherlands. Throughout western Europe, the Maastrichtian is characterized by chalk \*limestones. *See also* **SENONIAN**.

**Mab (Uranus I)** A lesser satellite of \*Uranus with a radius of 5 km.

**McAuliffe** A \*solar system asteroid (No. 3352); diameter 2–5 km; orbital period 2.57 years. Its orbit crosses that of \*Mars. A plan for the \*New Millennium Deep Space-1 spacecraft to visit McAuliffe in 1999 was cancelled.

**mackerel sky** Pattern of wavy \*cirrocumulus (or \*altocumulus) cloud with holes which produces an overall resemblance to the body markings of mackerel. *See also* **VERTEBRATUS**.

**Mackereth corer** A form of **\*hydraulic** corer commonly used to obtain lake-sediment cores.

**macroclimate** The climatic character of a large region.

**macroevolution** **\*Evolution** above the species level, i.e. the development of new species, genera, families, orders, etc. There is no agreement as to whether macroevolution results from the accumulation of small changes due to **\*microevolution**, or whether macroevolution is uncoupled from microevolution.

**macropygous** See PYGIDIUM.

**macrotidal** Applied to coastal areas where the **\*tidal range** is in excess of 4 m. Tidal currents dominate the processes active in macrotidal areas, e.g. the coast of the British Isles.

**macula** (*pl.* maculae) A dark spot on the surface of an extraterrestrial body.

**maculae** See MACULA.

**maculose** Applied to the texture of contact-metamorphic (see **CONTACT METAMORPHISM**) rocks that display spots, each spot representing a fine-grained aggregate of **\*minerals**.

**Madagascar** A large island lying in the **\*Indian Ocean** off the eastern coast of Africa. It became separated from the main mass of Africa at a time when India was migrating northwards across the Indian Ocean, about 95 Ma ago. Madagascar is famous for the occurrence of primitive prosimians (lemurs) which survive there because of a lack of mammalian predators. Elsewhere, prosimians have been greatly reduced, probably as a result of competition with true monkeys and apes following their appearance in the **\*Oligocene**.

**made ground (made land)** Area of dry land that has been constructed by people, generally through the reclamation of marshes, lakes, or shorelines. An artificial fill (landfill) is used, consisting of natural materials, refuse, etc.

**Madelung constant** The sum of the mutual potential coulombic attraction energies of the **\*ions** in a **\*crystal**. The constant is used in the equation for **\*lattice energy**.

**Madreporaria** See SCLERACTINIA.

**madreporite** In *\*Echinodermata*, a porous, button-shaped process on the *\*aboral* surface of the body; through its perforations the water-vascular system is connected to the water outside.

**Maentwrogian (Dresbachian)** A *\*stage* of the Upper *\*Cambrian*, underlain by the Late Cambrian, preceded by the Middle Cambrian, followed by the Festiniogian, and dated at 501–496.8 Ma ago (Int. Commission on Stratigraphy, 2004).

**maerl** The Breton name for a mixture of carbonate-rich (skeletal) sand and seaweed used as an agricultural dressing. *See also* **MARL**.

**Maestrichtian** *See* **MAASTRICHTIAN**.

**maestro** Local north-westerly wind of the Adriatic Sea which affects the western coasts, especially in summer. The term is also applied to north-westerly winds in the Ionian Sea and to winds off the coasts of Corsica and Sardinia.

**mafic** Applied to any *\*igneous* rock which has a high proportion of *\*pyroxene* and *\*olivine*, such that its *\*colour index* is between 50 and 90 (i.e. it is dark coloured).

**Magellan** A radar mapper, launched by *\*NASA* in 1989 to explore *\*Venus*.



<http://nssdc.gsfc.nasa.gov/planetary/magellan.html>

- A NASA mission to study and map the surface of Venus.

**Magic Mountain** A large *\*hydrothermal vent* field located about 150 km west of Vancouver Island, Canada, close to but outside the *\*Explorer Ridge*.

**magma** A hot, *\*silicate*, *\*carbonate*, or *\*sulphide \*melt* containing dissolved *\*volatiles* and suspended *\*crystals*, which is generated by partial melting of the Earth's *\*crust* or *\*mantle* and is the raw material for all *\*igneous* processes. The melt component of silicate magmas, by far the most common magma type, comprises a disordered mixture of single Si–O tetrahedra and chains, branching chains, and rings of Si–O tetrahedra, between which are located randomly positioned *\*cations* (e.g. Ca<sup>2+</sup>, Mg<sup>2+</sup>, Fe<sup>2+</sup>, and Na<sup>+</sup>) and *\*anions* (e.g. OH<sup>-</sup>, F<sup>-</sup>, Cl<sup>-</sup>, and S<sup>-</sup>) loosely coordinated with the oxygens in the silicate tetrahedra. The greater the silica content of

the magma, the more chains and rings of silicate tetrahedra there are to impede each other and hence increase the **\*viscosity** of the magma. The pressure regime and composition of the magma control which minerals nucleate and crystallize from a magma when it cools.

**magma chamber** A region, postulated to exist below the Earth's surface, in which **\*magma** is received from a source region in the deep **\*crust** or upper **\*mantle**, stored, and from which it moves to the Earth's surface at the site of a **\*volcano**. The existence of 'magma chambers' is often invoked by geochemists to provide a location for processes such as **\*fractional crystallization**, because the chemistry of **\*lavas** is explicable only in terms of such processes. When magma moves rapidly from the chamber, as in a **\*pyroclastic flow \*eruption**, the unsupported chamber roof may collapse to produce a **\*caldera** at the surface. The diameter of the caldera can be used to estimate the diameter of the underlying magma chamber; diameters of up to 40 km are not uncommon for terrestrial subvolcanic magma chambers.

**magnaphile elements** Elements that are partitioned into silicate melts during partial melting of magma. They include calcium (Ca), aluminium (Al), iron (Fe), sodium (Na), potassium (K), rubidium (Rb), caesium (Cs), barium (Ba), strontium (Sr), lanthanum (La), lead (Pb), thorium (Th), and uranium (U).

**magmatic arc** Where an oceanic plate is sinking beneath a less dense continental plate at a **\*subduction zone**, the region of raised elevation between the **\*fore-arc** and **\*back-arc basins**, where **\*magma** is rising.

**magmatic differentiation (magmatic fractionation)** Formation of a variety of rock types from an initial single parental **\*magma**. No specific mechanism is implied by the term. *See also* FRACTIONAL CRYSTALLIZATION.

**magmatic fractionation** *See* MAGMATIC DIFFERENTIATION.

**magmatic-segregation deposit** Concentration of particular minerals in different parts of a **\*magma chamber** during consolidation, by **\*gravity settling**, filter pressing, flow, **\*fractional crystallization**, liquid immiscibility, or gas transference; for example, the accumulation of heavy minerals such as **\*chromite** and **\*magnetite** by gravity settling.

**magnesiochromite** *See* CHROMITE; SPINEL.

**magnesioferrite** See MAGNETITE.

**magnesiowüstite** See FERROPERICLASE.

**magnesite** \*Carbonate,  $\text{MgCO}_3$  and \*end-member of a \*solid solution series with \*siderite,  $\text{FeCO}_3$ ; sp. gr. 3.0; \*hardness 4; whitish; \*earthy \*lustre; compact or granular; occurs as an alteration product of \*serpentines, \*dolomites, or \*limestones and may form as a chemical precipitate. It is mined commercially and used in the production of magnesium compounds, refractory products, and special cements.

**magnesium number (Mg#)** The ratio of magnesium to iron in an \*igneous rock, which is related to the chemical composition of the mantle material. As the magma cools, lighter minerals solidify first, forming a scum on the surface of the molten rock. If the composition of the mantle is known, the magnesium number can be predicted, and the magnesium number provides information on the composition of the mantle rock. The magnesium number also provides information on the thermal history of the rock. One task for the SMART-1 (Small Mission for Advanced Research in Technology) mission to the Moon, which was deliberately crashed into the lunar surface on 3 September 2006, was to determine the magnesium number of lunar surface rocks. If, as expected, the lunar magnesium number is lower than that of rocks on Earth it will support the view that the Moon was blasted from the Earth by an impact and the surface rocks on both bodies are derived from the same mantle material. The lower lunar value would reflect the different thermal histories of the Earth and Moon. If the lunar magnesium number is the same as that on Earth, however, it would prove that the two bodies could not have originated from the same source.

**magnesium-sulphate soundness test** A test identical to the \*sodium-sulphate soundness test, but in which magnesium sulphate is used instead of sodium sulphate.

**magnetic age** See MAGNETIC ANOMALY PATTERN; CHRON; MAGNETIC DATING.

**magnetic anomaly** Any magnetic field remaining after allowance for some particular model. Normally this is the field remaining after allowance for the \*International Geomagnetic Reference Field, but sometimes it is the

field remaining after further removal of either *\*regional* or near-surface anomalies.

**magnetic anomaly pattern (magnetic age)** Phenomenon, discovered originally in the north-eastern Pacific Ocean, of linear *\*magnetic anomalies* which lie parallel to oceanic *\*ridges* where spreading has occurred. The magnetic striping, which is determined by instrumental measurements, results from repeated *\*reversals* of the Earth's magnetic field (see *GEOMAGNETIC FIELD*): it exists as corresponding patterns on either side of an oceanic spreading centre (see *MID-OCEAN RIDGE*). These can be correlated with the *\*magnetostratigraphic* timescale (polarity timescale), so assigning a magnetic age to some of the individual magnetic anomalies. See also *SEA-FLOOR SPREADING*.

**magnetic dating** The use of magnetic properties for the age assessment of archaeological and geologic materials. It uses the *\*natural remanent magnetization* acquired at a specific time. Its direction can be compared with timescales of geomagnetic *\*secular* variations, *\*polarity reversal* sequences, or *\*polar wander paths*. Its intensity can be converted to the strength of the field in which it was acquired and so compared with the timescales of intensity changes of the *\*geomagnetic* field. *\*Viscous* remanent magnetization may sometimes be used to establish how long the sample has been in a specific position.

**magnetic domain** A volume of a *\*ferromagnetic* material in which all *\*electron* spins are aligned in the same direction. In most naturally occurring minerals a magnetic domain is about 1µm in diameter.

**magnetic equator (aclinic line)** The line that joins points on the Earth's surface at which the magnetic *\*dip* is zero. It lies close to the geographic equator.

**magnetic fabric determination** The determination of the *\*petrofabric* using magnetic methods. Generally it is the determination of the *\*magnetic susceptibility* ellipsoid, as this is dependent on the shape of *\*magnetite* grains or the crystalline alignment of *\*hematite* grains. The magnetic fabric may also be due to *\*paramagnetic* and diamagnetic materials. Some instruments may be sensitive to the *\*anisotropy* of electrical conductivity in the absence of *\*ferromagnetic* materials.

**magnetic field** See GEOMAGNETIC FIELD.

**magnetic flux** The magnetic induction, perpendicular to the surface area of a nearby body, multiplied by the area of the body.

**magnetic gradiometry** A geophysical technique, sometimes used in archaeology, that measures the magnetic properties of subsoil materials.

**magnetic induction ( $B$ )** The vector,  $B$ , which is equal to the magnetic permeability of a material ( $\mu$ ) multiplied by the magnetic field intensity vector ( $H$ ) applied to it.

**magnetic moment ( $M$ )** The measured total intensity of magnetization of a sample, irrespective of the volume or weight of the sample itself. It is measured in units of  $A \cdot m^2$ .

**magnetic orientation** The sensing by certain organisms of the direction of the \*geomagnetic field and their use of it for purposes of orientation. See BIOMAGNETISM. See also MAGNETIC SAMPLING.

**magnetic permeability** See PERMEABILITY.

**magnetic profile** A series of determinations of the intensity of the \*geomagnetic field along a traverse, usually oriented north–south.

**magnetic quiet zone** Any area showing few or no \*magnetic anomalies. Magnetic quiet zones are commonly oceanic areas where magnetic anomalies appear to be absent because either the \*geomagnetic field did not change \*polarity during the formation of the \*oceanic crust or the original magnetization has been destroyed by the effects of thermal blanketing by later sediments.

**magnetic sampling** The collection of samples in order to study their magnetic properties in the laboratory. The samples may be unoriented if they are only for rock-magnetic properties, e.g. \*magnetic susceptibility, magnetic intensity. For most \*archaeomagnetic and \*palaeomagnetic purposes samples are collected using a small portable drill and oriented using Sun, magnetic, or surveying methods, or \*gyrocompasses, and often utilizing special orientation devices to ensure precision.

**magnetic separator** A concentrator in which \*ore particles pass through a magnetic field or series of fields. The magnets may be permanent or

electromagnets, and may work wet or dry.

**magnetic signature** See SIGNATURE.

**magnetic storm** Major disturbance of the Earth's magnetic field resulting from the passage of high-speed charged solar particles, following a solar flare. The particles are deflected towards the regions of the Earth's magnetic poles.

**magnetic survey** A survey along a profile or grid using a \*magnetometer to determine the strength of the \*geomagnetic field at particular points.

**magnetic susceptibility** The constant of proportionality ( $\kappa$ ) between an applied field ( $H$ ) and the \*magnetic moment induced ( $J$ ), i.e.  $J = \kappa H$ ;  $\kappa$  is measured in dimensionless SI units. See also SUSCEPTIBILITY METER.

**magnetic variations** Periodic and irregular changes in the \*geomagnetic field, usually subdivided into brief \*transient variations and long-term \*secular variations.

**magnetite (loadstone, lodestone)** An important rock-forming iron \*oxide  $\text{Fe}^{2+}\text{Fe}^{3+}_2\text{O}_4$  ( $\text{FeO}\cdot\text{Fe}_2\text{O}_3$ ) and member of the \*spinel group of \*minerals; belongs to the magnetite series along with magnesioferrite ( $\text{MgFe}^{3+}_2\text{O}_4$ ), \*franklinite, jacobsonite ( $\text{MnFe}^{3+}_2\text{O}_4$ ), and trevorite ( $\text{NiFe}^{3+}_2\text{O}_4$ ); sp. gr. 5.2 (to 4.6); \*hardness 5.5–6.5; \*cubic; normally iron-black with a blue tarnish; black streak; \*sub-metallic \*lustre; crystals often octahedral and \*rhombohedronal, but it is also \*massive and granular; \*cleavage poor, octahedral; strongly magnetic; found extensively in magmatic rocks associated with \*basic rocks, in \*pegmatites, in contact \*metasomatic zones (especially in \*limestones in \*regionally metamorphosed rocks due to the dehydration of iron hydroxides), and also in black sands associated with gold gravels.

**magnetochemistry** Geochronological system that is based on geomagnetic \*polarity reversals. As no theory yet specifies the existence and duration of separate geomagnetic/polarity intervals, a true magnetochemistry is said not to exist. See also MAGNETOSTRATIGRAPHIC TIMESCALE.

**magnetogram** A recording of the time variations of the *\*geomagnetic field*, usually in analogue form. See *MAGNETOGRAPH*.

**magnetograph** An instrument for producing a record of changes of geomagnetic elements as a function of time. See *MAGNETOGRAM*.


**magneto hydrodynamics** The science of relating magnetic fields, mainly mathematically, within a moving conducting medium. It is mainly applicable to the Earth's *\*core*, where the *\*geomagnetic field* is generated by the motion of magnetic lines of force that are 'frozen' within a moving, electrically conducting medium, but it is also applicable to all systems involving the fluid motion of electrically conducting materials within a magnetic field.

**magnetometer** An instrument for measuring either the direction or the intensity of a magnetic field. Common field instruments include the *\*fluxgate* type and *\*nuclear-precession* types. The fluxgate magnetometer can also be used as a gradiometer and it is therefore most sensitive to shallow magnetic sources. Observatory instruments include the dip-circle and astatic types.

**magnetopause** The boundary within the *\*magnetosheath*, between the *\*solar* wind and the region closer to the planet which is shielded from the *\*solar* wind.

**magnetosheath** The region between the *\*bow shock*, where *\*solar wind* particles are slowed from supersonic to subsonic velocities, and the *\*magnetopause*.

**magnetosphere** The space around a planet in which ionized particles are affected by the planet's magnetic field. The Earth's magnetosphere reaches far beyond the *\*atmosphere*. In the magnetosphere, charged particles are concentrated at altitudes of about 3000 km and 16 000 km. The charged particles oscillate between the northern and southern hemispheres. The outer boundary of the magnetosphere is sharp and well defined, extending to about 10 Earth radii on the sunlit side of the Earth and to perhaps 40 Earth radii on the dark side; but the boundary changes its position in response to solar activity, being depressed by the *\*solar wind*. See also *EXOSPHERE*; *IONOSPHERE*.



## Magnetosphere

**magnetostratigraphic timescale (polarity timescale, geomagnetic reversal timescale, reversal timescale)** A timescale based on the periodic **\*polarity reversals** in the Earth's **\*geomagnetic field**. Magnetic minerals within a rock retain an orientation induced by the field at the time the rock was formed (see **NATURAL REMANENT MAGNETISM**). Provided they include suitable minerals, **\*strata** from all over the world thus contain a record of the **\*normal** (as at present) or reversed state of the geomagnetic field at the time of their formation. This reversal pattern has been correlated between different successions of rocks to produce a sequence that, when combined with a **\*dating** method such as **\*potassium–argon dating**, has given a timescale measured in units of normal or reversed polarity. The scale was first established in detail for the last 4.5 Ma using data from terrestrial, mainly **\*extrusive**, rocks; it has now been extended back to the Upper **\*Jurassic** by means of the **\*magnetic anomaly** patterns in **\*oceanic** crust. The terms proposed by the **\*ISSC** for geochronologic units of the magnetostratigraphic timescale are **\*polarity superchron**, **\*polarity chron** (replacing the earlier term 'epoch'), and **\*polarity subchron** (replacing the earlier term 'event'). The corresponding terms proposed for rocks deposited during those **\*polarity intervals** are polarity superchronozone, polarity chronozone, and polarity subchronozone.

**magnetostratigraphy** Branch of stratigraphy based on geomagnetic \*polarity reversals.

**magnetotelluric sounding** The use of changing components of the \*geomagnetic field to study the electrical conductivity of the rocks within the Earth. The geomagnetic-field changes induce eddy currents within the Earth that are in proportion to the electrical conductivity of the rocks concerned, but they can readily be separated from the inducing currents because of their phase difference. It is generally applied to studies of the lower \*crust and upper \*mantle.

**magnetozone** A unit of rock of the same magnetic polarity (character). See also CHRON.

**magnitude, earthquake** See EARTHQUAKE MAGNITUDE; MERCALLI SCALE; MOMENT MAGNITUDE SCALE; RICHTER SCALE.

**Magnolia** One of the most ancient of flowering plants, noted for the relatively primitive structure of its flower. *Magnolia* leaves are first recorded from Dakota, USA, in the mid-\*Cretaceous. *Magnolia paepetala* is a reconstruction, by E. E. Leppik, of a whole flower based on the earliest \*angiosperm petal ever found (also from the Dakota fossil flora).

**magnon** In a magnetic material, a quantum of spin-wave energy.

**main-stage sequence** See HYDROGEN 'BURNING'; NUCLEOSYNTHESIS.

**majanna** A dry, level plain with a surface encrusted with salt.

**malachite** Mineral,  $\text{Cu}_2\text{CO}_3(\text{OH})_2$ ; sp. gr. 3.9–4.0; \*hardness 3.5–4.0; \*monoclinic; bright green; pale green \*streak; in fibrous condition with a silky \*lustre, crystals with \*adamantine or \*vitreous lustre; crystals very rare, and it is usually found as \*botryoidal, encrusting masses with bands of varying colour, and frequently in a fibrous, radiating \*habit; common \*secondary mineral in the oxidized zone of copper deposits, associated with \*azurite, native \*copper, and \*cuprite; soluble with effervescence in dilute hydrochloric acid. It is often used for ornaments and pigments, and as an \*ore mineral for copper. See SPESSARTITE.

**Malacostraca** (phylum \*Arthropoda, subphylum \*Crustacea) One of eight classes of Crustacea, the malacostracans appear in the \*Cambrian. There

are usually eight pairs of **\*biramous**, thoracic limbs, some forms have pincer-like appendages, and the limbs may be modified for swimming, feeding, or other purposes. The group is diverse and includes crabs, lobsters, and the shrimps, and it has quite a good **\*fossil** record.

**Malagasy aardvark** See **BIBYMALAGASIA**.

**Malakovian** A New Zealand **\*stage** (248–243 Ma ago) of the **\*Triassic**, followed by the **\*Etalian**.

**malleolus** A small prominence on the **\*distal** end of the **\*tibia** and of the **\*fibula**; in humans they form the ‘knobs’ of the ankle.

**Mallet, Robert** (1810–81) A civil engineer from Dublin, Mallet became interested in **\*earthquakes** and **\*volcanoes**, making a detailed study of the Neapolitan earthquake of 1857. He investigated the velocity of **\*seismic waves**, and compiled an earthquake catalogue from which he was able to make a seismic map of the world.

**malleus** In **\*Mammalia**, the outer **\*ossicle** of the middle ear, which is derived from the articular bone of ancestral vertebrates.

**Malm** An alternative name for the Late **\*Jurassic \*epoch**, preceded by the Middle Jurassic, followed by the **\*Berriasian** (**\*Cretaceous**), and dated at 161.2–145.5 Ma ago (Int. Commission on Stratigraphy, 2004). It comprises the **\*Oxfordian**, **\*Kimmeridgian**, and **\*Tithonian** (see also **VOLGIAN**) **\*ages**. It is roughly contemporaneous with the upper **\*Kawhia** and **\*Oteke** (New Zealand).

**Malvernian** A **\*stage** of the Upper **\*Proterozoic** of the Welsh borders and English Midlands, underlain by the **\*Monian** and overlain by the **\*Uriconian**.

**mamma (mammatus)** From the Latin *mamma* meaning ‘udder’, a cloud feature consisting of projections from the basal surface of the **\*anvil** of a large **\*cumulonimbus** storm cloud. It indicates violent turbulence in the cloud and may precede a tornado.

**Mammalia (mammals)** (phylum **\*Chordata**) Class of **\*homoiothermic** animals in which the head is supported by a flexible neck, typically with seven **\*vertebrae**, articulating through two **\*occipital condyles**. The lower jaw is composed of one bone only, the dentary, which articulates directly

with the skull; the middle ear contains three small bones, two of which are derived from bones in the lower jaw of *\*reptiles*. Typically, mammalian teeth are of differing forms (*\*heterodont*), they are set in sockets (thecodont), and milk teeth are shed and replaced by a second set. A hard *\*palate* separates the nasal cavity from the mouth. Except in *\*monotremes*, the egg is small and develops in the uterus; and the young are fed milk secreted by mammae (which give the class its name). The skin has at least a few hairs. Many mammalian features were present in *\*therapsids* (mammal-like reptiles) during the *\*Triassic*. Mammals are believed to have appeared first toward the end of the Triassic and to have diversified rapidly from the end of the *\*Mesozoic*, 100 Ma later, following the mass *\*extinction* which marks the *\*Mesozoic–\*Tertiary* boundary 65 Ma ago.

**mammal-like reptiles** See SYNAPSIDA.

**mammatus** See MAMMA.

**mammillary (mammillated)** Applied to the physical *\*habit* of a mineral which has grown from radiating crystals to give curved or rounded surfaces.

**mammillated** See MAMMILLARY.

**mammillated topography** Hill relief with a streamlined, rounded, and smoothed appearance, normally resulting from the scouring action of an *\*ice* sheet, as in the Adirondack Mountains, eastern USA. However, some crystalline rocks may support a pseudo-mammillated surface formed by non-glacial processes, as on Ben Lomond, Tasmania.

**Mammoth** A reversed *\*polarity subchron* which occurs within the *\*Gauss* normal *\*polarity chron*.

**mammoth** See MAMMUTHUS.

**Mammuthus (mammoth)** Line of *\*Pleistocene* elephants adapted to steppe and *\*tundra* habitats. The tusks were elongated and strongly curved, and the skull was shorter and higher than that of other elephants. The woolly mammoth was adapted to arctic environments. The largest mammoth, indeed the largest *\*proboscidean* of all time, was *Mammuthus armeniacus* of Eurasia, which stood about 4.5 m at the shoulder.

**Mammutidae** Extinct family of mastodons, comprising the one genus Mammut (*Mastodon*) of elephant-like animals, which diverged from the evolutionary line leading to the modern elephants. It was long-lived, extending from the Lower **\*Miocene** to the Recent, and it survived in Africa, N. America, and Eurasia at least until the end of the **\*Pleistocene**. Mastodons were shorter and heavier in build than elephants. Mammut (or *Mastodon*) species had short, high skulls, longer jaws than elephants, and usually tusks in both upper and lower jaws, the upper tusks often being large and curving outward and upward. There were never more than two teeth in use at a time, never more than a vestige of the lower incisors, and the molars were low-crowned, simple, and lacked cement. It is believed that the evolution of the mastodons paralleled that of the gomphotheres (**\*Gomphotheriidae**).

**mandible** In **\*Crustacea**, **\*Insecta**, and Myriapoda (centipedes, millipedes, etc.), one of the pair of mouthparts most commonly used for seizing and cutting food. In birds, specifically the lower jaw but the term is also used to denote the two parts of the bill of a bird, as upper and lower mandibles.

**manganese nodule** Concretion of iron and manganese oxides which also contains copper, nickel, and cobalt. Nodules are variable in size, shape, and composition, and are layered internally. The average composition is: manganese 30%; iron 24%; nickel 1%; copper 0.5%; cobalt 0.5%. Manganese nodules are widely distributed on the sea floor of every ocean and in some temperate lakes. They are found in areas of negligible sedimentation and/or strong bottom currents, e.g. on the N. Pacific **\*abyssal plain** at depths of 3500–4500 m, and on the shallow, current-swept Blake Plateau off the east coast of the USA. Submarine mining of these deposits is now considered viable to recover copper and other scarce metals. Growth of manganese nodules apparently reached a peak in the early **\*Palaeogene** period.

**manganite** Mineral,  $\text{MnO}_2\text{Mn}(\text{OH})_2$ ; sp. gr. 4.2–4.4; **\*hardness** 3–4; **\*monoclinic**; dark steel-grey to black; reddish-brown to black **\*streak**; **\*sub-metallic** **\*lustre**; crystals **\*prismatic** with striated faces, often as bundles or radiating aggregates; **\*cleavage** perfect, pinacoidal; occurs in deposits precipitated in sedimentary conditions in which there is an oxygen deficiency, and in low-temperature **\*hydrothermal** veins in association with

**\*barite** and **\*goethite**; soluble in hydrochloric acid with the evolution of chlorine. It is often used in steel-making.

**Mangaorapan** A **\*stage** (52–50 Ma ago) in the **\*Eocene** of New Zealand, underlain by the **\*Waipawan**, overlain by the **\*Heretaungan**, and roughly contemporaneous with the upper **\*Ypresian** stage.

**Mangaotanean** A New Zealand **\*stage** (93–87 Ma ago) of the Late **\*Cretaceous** **\*epoch**, preceded by the **\*Arowhanan** and followed by the **\*Teratan**.

**Mangapanian** A **\*stage** (2.6–1.96 Ma ago) in the Late **\*Pliocene** of New Zealand, underlain by the **\*Waipipian**, overlain by the **\*Nukumaruan**, and roughly contemporaneous with part of the **\*Piacenzian** stage.

**mangrove swamp** Characteristic vegetation of tropical, muddy coasts, and typically associated with river mouths where the water is shallow and the load of suspended sediment is high. The aerial roots of the mangrove trees trap the sediment, favouring the gradual seaward extension of the land area.

**man-made soil** A major **\*soil** group in the soil classification developed by the Soil Survey for England and Wales. Man-made soils result from such operations as land restoration following mining or quarrying.

**Manning equation** An equation that relates the velocity ( $V$ ) of water flowing through a stream to the **\*hydraulic** radius of the stream ( $R$ ), its slope ( $s$ ), and an estimate of its **\*bed roughness** ( $n$ ).  $V = (R^{2/3}s^{1/2})/n$ . The equation was developed by the Irish engineer Robert Manning (1816–97) and its first version appeared in 1891 in Manning's paper 'On the flow of water in open channels and pipes' (Transactions of the Institution of Civil Engineers of Ireland).



<http://manning.sdsu.edu/>

- An account of Manning's life and work.

**manometer** An instrument that can be used for the direct measurement of vacuum, positive pressure, and differential pressure. It usually consists of two interconnected tubes filled with fluid, e.g. water or mercury. A difference in fluid levels in the two tubes records the pressure. It can also be modified to measure flow.

**Mansurian** The fourth period of the *\*Mercurial geologic timescale*, lasting 3000–1000 Ma ago, during which the contraction of the planet *\*Mercury* came to an end, crater formation lessened, and the planet cooled, leading to a thickening of its lithosphere, and there was a decrease in geologic activity. Craters formed during the Mansurian lacked *\*rays*.

**Mantell, Gideon Algernon** (1790–1852) A Sussex surgeon, whose hobby was the study of fossils from the Chalk and Weald, on which he became an expert. Mantell discovered *\*iguanodon* and other fossil *\*dinosaurs* which, with the remainder of his collection, were donated to the British Museum (Natural History), now the Natural History Museum. He also published a number of popular works on geology.

**mantle** **1.** Zone lying between the Earth's *\*crust* and *\*core*, approximately 2300 km thick. The mantle is probably similar in composition to *\*garnet* *\*peridotite* and it represents about 84% of the Earth's volume and 68% of its mass. A mantle is present in most *\*terrestrial* planets and the *\*Moon*, but is of a different composition in each case. **2. (pallium)** In *\*Brachiopoda* and *\*Mollusca*, a layer of tissue that covers the body and is responsible for the secretion of the shell.

**mantle array** A term used in isotope geochemistry to describe a graphical plot of  $^{144}\text{Nd}:^{143}\text{Nd}$  against  $^{87}\text{Sr}:^{86}\text{Sr}$  for *\*igneous* rocks. Rocks which have been derived from the *\*mantle* tend to plot on a straight line; those that show evidence of crustal contamination tend to fall off the line, often in systematic ways.

**mantle bedding** The maintenance of a uniform thickness of *\*pyroclastic* material over all but the steepest topographic features. Fall deposits are the main group of pyroclastic rocks which commonly display mantle bedding, although *\*ignimbrite* veneer deposits may also display it.

**mantle cavity** In *\*Mollusca* and some *\*Brachiopoda*, the space containing the *\*gills* that formed by the folding of the tissues comprising the *\*mantle*.

**mantle convection** The transfer of heat by movement of material within the *\*mantle*. Material derived from within the mantle is added to the *\*lithosphere* at *\*constructive* margins and cool lithosphere descends at *\*subduction zones*, thus some return flow of material must take place at depth. The return flow may involve only material at shallow depth below

the lithosphere, or material through the whole mantle, or possibly two or more layers of convection cells transferring heat between the layers by conduction. **\*Hot spots** may overlie isolated plumes of material rising from the **\*core–mantle interface**.

**mantle creep** The mechanism by which **\*mantle** material deforms. Dislocation creep (see **CREEP MECHANISMS**) is believed to predominate in the upper mantle in the vicinity of **\*subduction zones**, and diffusion creep predominates in the lower mantle.

**mantled gneiss dome** A **\*dome** of granitic **\*migmatites** and **\*gneisses** surrounded by a ‘mantle’ of metasediments which characteristically show outward-dipping parallel **\*fabrics** and **\*foliations**. The feature is thought to be produced by gravitational instabilities due to a density inversion in which dense rocks overlie a less dense, **\*granite–\*gneiss** core.

**mantle lobe** See **BRACHIOPODA**.

**mantle plume** Localized, hot, buoyant material which is hypothesized to be rising through the **\*mantle**. Mantle plumes are thought by some geologists to rise beneath **\*hot spots**, causing **\*domal** uplift. Plumes may originate near the mantle–core boundary, and it has been suggested that they generally have the form of a cylinder with a radius of about 150 km.

**manto** A horizontal, bedded **\*ore** deposit; the term may describe a sedimentary **\*stratum**, or a replacement, strata-bound deposit.

**Maozhangian** A Chinese **\*stage** (513–509 Ma) of the Middle **\*Cambrian**, preceded by the **\*Longwangmiaoan** and followed by the **\*Zuzhuangian**.

**map projection** The representation on a plane surface of part or all of the surface of the Earth or a celestial body. Various projections have been used in the Earth sciences because of their particular suitability for certain work, e.g. equal-area projections for the global distributions of different sedimentary deposits, polar stereographic projection for the relationships within **\*Gondwana**, Mercator’s projection for the absolute movements of the lithospheric **\*plates**, and conical projections for the Himalayan–Alpine **\*fold belt**. Projections based on Mercator’s are most frequently used because of their general familiarity, unless the information portrayed is peculiarly distorted.

**marble** Non-foliated, metamorphosed **\*limestone** which is produced by **\*recrystallization** and is hard enough to take a polish. The hardest and most attractive marbles have been used in statuary and for building since antiquity and are still quarried, e.g. from the Carrara quarry which supplied Michelangelo. The statue of Abraham Lincoln at the Lincoln Memorial, Washington, DC, is made from marble quarried in Georgia, USA. Marbles may be variously coloured or banded, depending on their chemical and mineralogical composition (mostly **\*calcite**), e.g. Carrara marble is pure white, but Siena marble, quarried in Tuscany, has red mottling.

**marcasite** Mineral,  $\text{FeS}_2$ ; sp. gr. 4.8–4.9; **\*hardness** 6.0–6.5; **\*orthorhombic**; pale bronze-yellow; greyish-black **\*streak**; **\*metallic \*lustre**; crystals **\*tabular** and spear-like, but it is also **\*massive**, stalactitic, and radiating; **\*cleavage** {101}; occurs in low-temperature mineral **\*veins** together with zinc and lead ores, and in **\*sedimentary rocks**, especially **\*limestone**, **\*chalk**, and **\*clay**. It has been used in the production of sulphuric acid. *Compare* PYRITE.

**mare** (*pl.* maria) The Latin word for ‘sea’, originally used by Galileo in 1610 to denote the smooth, grey areas on the **\*Moon**, visible to the naked eye, that are now known to be plains of basaltic lava filling circular basins excavated by **\*planetismal** impact. **\*Basalt** lavas, mainly in maria, cover 17% of the surface of the Moon, mostly on the near side. The term is also applied to low-lying areas on **\*Mars**.

**mare ridge** See WRINKLE RIDGE.

**mares’ tails** Popular name for tufted **\*cirrus** clouds with **\*virga** (precipitation trails) seen below each cloud.

**Margaret (Uranus XXII)** A lesser satellite of **\*Uranus** with a radius of 10 km.

**margarite** A dioctahedral **\*mica** with the general composition  $\text{Ca}_2\text{Al}_4(\text{Si}_4\text{Al}_4\text{O}_{20})(\text{OH},\text{F})_4$ , found in fine-grained, low-grade, regional **\*metamorphic rocks** (see METAMORPHIC GRADE). Calcium is the main **\*cation** found between the individual **\*silicate** sheets of the mica structure. Margarite has the typical appearance of mica, but is much harder than common mica types. **\*Cleavage** sheets are also less elastic than those of the common micas and hence margarite is a member of the ‘brittle mica’ group.

At one time this mineral was thought to be a rare component of metamorphic rocks but recent *\*X-ray diffraction* work has demonstrated that it can be abundant in low-grade metapelites (see *META-*). It occurs with *\*corundum* in *\*emery* deposits and in *\*mica* schists along with *\*tourmaline* and *\*staurolite*.

**marginal basin** 1. A synonym for *\*back-arc basin*. 2. A basin developed along a *\*passive margin*.

**marginal sea** 1. A semi-enclosed body of water adjacent to, and widely open to, the ocean (e.g. the Gulf of Mexico, Caribbean Sea, and Gulf of California). 2. See *PERICONTINENTAL SEA*.

**marginal suture** See *CEPHALIC SUTURE*.

**marialite** See *SCAPOLITE*.

**Marianas Trench** The oceanic *\*trench* which marks the *\*destructive* margin between the *\*Pacific* and *\*Philippine Plates*. The trench is about 11 km deep, and there is evidence of little or no *\*accretionary wedge*. The *\*subduction zone* steepens at depth, becoming almost vertical, with *\*hypocentres* down to approximately 700 km.

**marine platform** See *SHORE PLATFORM*.

**Mariner** A series of *\*NASA* planetary missions exploring *\*Venus* (Mariners 2 (1962), 5 (1967), and 10 (1973) and *\*Mars* (Mariners 4 (1964), 6 (1969), 7 (1969), 9 (1971)); Mariner 10 also travelled to *\*Mercury*. There was no Mariner 1.



<http://nssdc.gsfc.nasa.gov/planetary/mars/mariner.html>

- A NASA programme of missions that studied Mars, Venus, and Mercury.

**maritime air** *\*Air* mass with properties of temperature and humidity derived either from a source region in, or from long passage over, the oceans.

**maritime climate** General term applied to a climate much modified by oceanic influences. Typical characteristics include relatively small diurnal

and seasonal temperature variation, and increased precipitation due to more moist air.

**marker bed (key bed, chronohorizon, chronostratigraphic horizon, datum)** A thin bed of a distinctive character, often widely distributed and capable of being recognized and traced over a large geographical area. Marker beds are taken to be the results of either (geologically speaking) very short episodes of deposition (e.g. a thin *\*coal* seam), or of almost instantaneous events (e.g. turbidity deposits (see [TURBIDITY CURRENT](#)), or *\*bentonite* beds derived from volcanic *\*ash* falls). Such beds are thus very valuable as time markers in stratigraphic *\*correlation*.

**Markov chain** In statistics, a set of sequential observations in which the probability of one member of the sequence occurring conditional on all the preceding members occurring is equal to the probability of that member occurring conditional only on the immediately preceding member occurring.

**marl** A *\*pelagic* or *\*hemipelagic* sediment (an *\*arl*), typically found interbedded with purer oozes in beds up to 1.5 m thick, with a composition intermediate between a non-biogenic sediment and a calcareous or siliceous ooze. It is 30% clay and 70% microfossils, at least 15% of its volume being siliceous microfossils. *Compare* [SARL](#); [SMARL](#).

**marlstone** A semi-lithified *\*marl*. A fully lithified marl is termed an argillaceous *\*limestone*.

**Mars** (*adj.* martian) The fourth planet in the *\*solar system*, 1.524 AU from the *\*Sun*. Its radius is 3390 km, its density 3940 kg/m<sup>3</sup>, and the inclination of equator to orbit is 25.1°. It has a small *\*atmosphere* (700 Pa) of CO<sub>2</sub>. The polar caps are of water ice with seasonal solid CO<sub>2</sub>. The northern-hemisphere crust is mainly basaltic plains and *\*volcanoes*; the southern an ancient cratered terrain (see [CRATER](#)). The Tharsis Bulge is an uplifted or volcanic plateau. Large canyons exist and there is evidence of former water *\*erosion*. Some *\*basaltic* meteorites are derived from Mars. It has two small *\*satellites*, Phobos and Deimos, probably captured *\*asteroids*.

**Mars 96** A Russian mission to *\*Mars*, launched in 1996, that was intended to place a vehicle in orbit and land instruments on the surface. The launch failed, the rocket and its payload falling back to Earth.



<http://www.iki.rssi.ru/mars96/mars96hp.html>

- An RSA mission launched in 1996 to investigate Mars.

**Marsdenian** A \*stage (317–315.5 Ma ago) of the \*Pennsylvanian epoch, preceded by the \*Kinderscoutian and followed by the Yeadonian.

**Mars Global Surveyor** A \*NASA mission to \*Mars, launched in 1996, which placed a vehicle in orbit about the planet.



[http://www.nasa.gov/mission\\_pages/mgs/index.html](http://www.nasa.gov/mission_pages/mgs/index.html)

- A NASA mission launched in 1996 to survey and map Mars.

**Marshbrookian** A \*stage (455–454 Ma ago) of the Late \*Ordovician in the Upper \*Caradoc, underlain by the \*Longvillian and overlain by the \*Actonian.

**Mars Observer** A \*NASA mission to \*Mars that was launched in 1992. It should have orbited the planet, surveying and mapping, but communication with the spacecraft was lost before it could be manoeuvred into orbit and the mission was abandoned.

**Mars Pathfinder** A \*NASA environmental survey mission to \*Mars that was launched in 1996 and soft-landed successfully in July 1997. After landing Pathfinder was renamed the Sagan Memorial Station, in honour of Carl Sagan. Sojourner, a small rover, travelled short distances from the main lander, exploring rocks selected by controllers on Earth from images transmitted by the lander.



<http://nssdc.gsfc.nasa.gov/planetary/mesur.html>

- A NASA mission launched in 1996 to study the feasibility of landings on, and explorations of, Mars.

**marsquake** The martian equivalent of an \*earthquake or \*moonquake. None were positively detected by the Viking Lander \*seismometer.

**Mars Surveyor** Several \*NASA missions to \*Mars, some consisting of two launches. Mars Surveyor '98, carrying an orbiter and lander, was lost during insertion into Mars orbit. Mars Surveyor 2001, planned for launch in 2001, was cancelled due to cost overruns. \*Mars Global Surveyor, launched in 2003, reached Mars in 2004; its two rovers, Spirit and Opportunity, were deployed successfully and found evidence of past water.



[http://www.windows2universe.org/mars/exploring/MSP\\_overview.html](http://www.windows2universe.org/mars/exploring/MSP_overview.html)

- NASA missions to map and study the climate of Mars.

**Marsupialia** (subclass Theria, infraclass Metatheria) An order that comprises some 250 species of living marsupials and many extinct forms. It is sometimes divided into three suborders (Polyprotodonta, which includes the opossum-like insectivorous, carnivorous, and omnivorous forms; Diprotodontia, containing the phalangers, kangaroos, and other forms evolved from an opossum-like stock, but differing structurally from the polyprotodonts; and Caenolestoidea (classed by others as a superfamily), containing a small group of 'opossum rats'), but nowadays it is usual to divide the marsupials into several orders, often allocated to two cohorts: Ameridelphia and Australidelphia. In this scheme, the name Marsupialia would cease to be used formally. Marsupials are characterized principally by their method of reproduction. The egg is yolky and has a thin shell protecting it from maternal antigens. Placental development is usually very limited and except in the Peramelemorpha the allantois serves no nutritional function, but uterine milk may be taken up by the yolk sac. Within 10–12 days of the breaking of the shell, the embryo (whose fore limbs and associated neural development, mouth, and olfactory system have developed precociously) is born. It crawls into the pouch (marsupium) and attaches itself to a teat, its lips growing around the teat, which injects milk without choking the embryo. In the later stages of its development an offspring may receive high-fat, low-protein milk from one teat while a newer embryo receives high-protein, low-fat milk from another. Marsupials also differ from placentals in their dentition, in the possession of an inflected angular process to the jaw, and in the presence of two marsupial bones which articulate with the pubes. Marsupials and placental mammals apparently diverged from a common ancestor in the \*Cretaceous. The first

marsupials were similar in general form to the opossums of America. In Australia the marsupials radiated to produce a wide array of adaptive types, while in S. America they filled the insectivorous and carnivorous niches for much of the \*Cenozoic, while placentals occupied the herbivorous niches.

**marsupials** See MARSUPIALIA.

**martian** Of, or pertaining to, or about, the planet Mars.

**martian canals** ('canali') Optical illusions, produced by telescopic viewing of Mars with a resolution of poorer than about 100 km, first reported by Schiaparelli ('canali' is the Italian for 'channels') and especially championed by Percival Lowell (1855–1916). These observers produced maps of the martian surface showing interconnected networks of canals, implying the presence of intelligent life on Mars. The intelligence which devised the canals was, however, on the terrestrial side of the telescope.

**martian terrain units** Polar units include Layered Deposits and Etched Plains; Volcanic units include Volcanic Constructs and Volcanic Plains; Cratered units include Ancient Cratered Terrain and Cratered Plains; Modified units include Knobby Terrain, Fretted and Chaotic Hummocky Terrain, Channel Deposits, and Grooved Terrain.

**mascon** Abbreviation for mass concentration. There are about ten large-scale positive \*gravity anomalies on the \*Moon, associated with the large circular maria (see MARE). The \*basalt filling the multi-ringed impact basins only partly accounts for the excess mascons per unit area, and central subsurface uplift of denser \*mantle material into the \*feldspathic \*lunar highland crust during the impact is required to provide the observed values of up to  $+2.2 \times 10^{-3} \text{ m/s}^2$  (+220 mgal) observed in the Mare Imbrium.

**maser** An acronym for *microwave amplification by stimulated emission of radiation*, a device resembling a \*laser but emitting radiation at microwave frequencies.

**mass balance** 1. Generally, a term used in comparisons of the inputs and outputs of processes. 2. The balance of elements in ocean water, which is assumed to be constant, i.e. influx and removal of elements occurs at the same rate. Influxes include river water, elements released from sediment pore fluid, and melting ice. Reaction between sea water and rocks is probably insignificant. The removal of elements from sea water occurs by

precipitation of chemical sediments, *\*ion exchange*, and the burial of pore fluids. **3.** The relation between the input and output of a *\*glacier*. Input (accumulation) is dominated by snow precipitation and output (*\*ablation*) by surface melting. The difference between accumulation and ablation for a glacier over a year is the net (mass) balance. A positive net balance implies that a glacier is growing; a negative net balance that it is shrinking; and a zero net balance that it is stable.

**mass extinction** See *EXTINCTION*.

**mass flow** A downslope slide of *\*sediment* which moves under the force of gravity. Mass flows include rockfalls (accumulations of *\*scree*), slumps and slides (where masses of sediment move downslope along discrete *\*shear* planes), debris flows (in which ill-sorted masses of sediment move downslope due to the loss of internal strength of the sediment mass), liquefied sediment flows, *\*grain flows*, and turbidity flows.

**massif** A very large topographic or structural feature, usually of greater rigidity than the surrounding rock. The term 'massif' is used of older crystalline blocks in an *\*orogenic* belt.

**massive** Lacking in any form or structure, e.g. massive beds are those without internal grading (see *GRADE*), and lacking *\*sedimentary structures*. Applied to a compact *\*mineral* which has no distinguishing *\*crystal* form.

**massive sulphide deposit** Rich mass of metallic *\*sulphide* minerals, with little *\*gangue*, which are relatively easy to mine. Typical examples are copper–nickel sulphides and copper–zinc–lead sulphides.

**mass movement** See *MASS-WASTING*.

**mass number** The sum of the protons and neutrons in the *\*nucleus* of an atom.

**mass spectrometry** Technique that allows the measurement of atomic and molecular masses. Material is vaporized in a vacuum, ionized, and then passed first through a strongly accelerating electric potential and then through a powerful magnetic field. This serves to separate the ions in order of their charge:mass ratio; the ions are detected, commonly by means of an electrometer which measures the force between charges and hence the

electric potential. Mass spectrometry is used in the *\*radiometric dating* of rocks, and in *\*isotope* geochemistry.

**mass-wasting (mass movement)** General term for the transfer of Earth material down hillslopes. It includes four main categories: flow, slide, fall (see *ROCK FALL*), and *\*creep*. Of these, creep is the most important if least spectacular. It is the result of gravity acting on material that has lost cohesion, typically as a result of an increase in water content. An avalanche is a rapid and often destructive flow of rock or snow. A slide (or landslide) is a comparatively rapid displacement of Earth material over one or more failure surfaces which may be curved or planar. Failure on an arcuate surface is typical of clays, and gives rise to rotational slides such as those of Folkestone Warren, England.

**master curve** One of a set of theoretical curves, calculated for known models, against which a field curve can be matched. If the two fit very closely the model is considered to apply reasonably well to the field situation and the curve is known as the master curve. Master curves were used extensively in *\*electrical* resistivity depth sounding, but are being replaced by microcomputer curve-matching which is much more accurate and more sensitive to real-life situations.

**mastodon** See *GOMPHOTHERIIDAE*; *MAMMUTIDAE*.

**Mata** A *\*series* in the Upper *\*Cretaceous* of New Zealand, underlain by the *\*Raukumara* series and overlain by the *\*Teurian* *\*stage*. It comprises the Piripauan and Haumurian stages, which are roughly contemporaneous with the upper *\*Santonian*, *\*Campanian*, and *\*Maastrichtian*.

**Mathilde** A *\*solar system* asteroid (No. 253), measuring 50 × 53 × 57 km; approximate mass 10<sup>17</sup> kg; density 1.3 g/cm<sup>3</sup>; rotational period 417.7 hours; orbital period 4.32 years. It was imaged by the *\*Near Earth Asteroid Rendezvous* mission in June 1997, revealing five craters more than 20 km across, the largest being 30 km wide and 6 km deep.

**matrix** Lithologic or petrographic term denoting the interstitial material lying between larger crystals, fragments, or particles. It is the background material of small particles in which larger particles and fragments occur. The term is applied to *\*sedimentary rocks*; the *\*igneous* equivalent is *\*groundmass*, although 'matrix' is also commonly used of igneous rocks.

**matrix-support** See MUD-SUPPORT.

**maturation** The evolution of **\*hydrocarbons** through the increasing pressure and temperature associated with burial. In the immature stage the product is gas; as it matures the development of heavy oils is succeeded by medium, and finally light oils. If the temperature exceeds approximately 100 °C, however, dry gas is the sole product and the stage of incipient **\*metamorphism** is reached. See also NATURAL GAS; PETROLEUM.

**Matuyama, Motonori** (1884–1958) Professor of theoretical geology at Kyoto Imperial University, Matuyama worked on gravity surveys, and magnetism in **\*basalts**. In 1929 he showed that the polarity of **\*remanent** magnetism in some recent basalts depended on the age of the rock, concluding that the Earth's magnetic field must undergo periodic reversals.

**Matuyama** A reversed **\*polarity chron** at the end of the **\*Pliocene** and the beginning of the **\*Pleistocene**. It is preceded by the **\*Gauss** and followed by the **\*Brunhes** normal-polarity chrons, and is **\*radiometrically dated** as occurring between 2.60 and 0.78 Ma ago. The Matuyama contains at least three normal **\*polarity subchrons**: **\*Réunion**, **\*Olduvai**, and **\*Jaramillo**.

**Maury, Matthew Fontaine** (1806–73) An American oceanographer and naval officer, Maury studied the winds and currents of the N. **\*Atlantic** and **\*Indian** Oceans, and produced the first bathymetric chart of the N. Atlantic. His *Physical Geography of the Sea* (1855) is said to be the first oceanographic textbook.

**maximum-likelihood classification** A **\*remote** sensing **\*classification** system in which unknown **\*pixels** are assigned to classes using contours of probability around **\*training** areas using the maximum-likelihood statistic. See also BOX CLASSIFICATION; MINIMUM-DISTANCE-TO-MEANS CLASSIFICATION.

**maximum-likelihood tree** In **\*phylogenetics**, a tree-building method that uses the maximum-likelihood statistical estimator to calculate the **\*topology** with the highest probability of being correct under assumed rates of **\*character** change.

**maximum-parsimony tree** The method for selecting a **\*phylogenetic tree** from all possible tree **\*topologies** that requires the smallest number of

substitutions.

**maximum thermometer** Thermometer that records the highest temperature to which it has been exposed (e.g. by allowing a rise of liquid past a restriction in the tube, but preventing the liquid's return on contraction). It is most commonly used to record maximum daily temperatures.

**Maxwell mantle** A region of the upper *\*mantle* in which the material exhibits *\*elastoviscous behavior*.

**Maxwell model** A description of a substance that exhibits *\*elastoviscous behavior*. The model has two components, an elastic spring and viscous damper, which are connected in series. *Compare* BURGERS MODEL; KELVIN–VOIGT MODEL.

**Mayan (Menevian)** A Russian-Kazakhstanian *\*stage* (502–501 Ma ago) of the Middle Cambrian, preceded by the Amgan and followed by the Ayusokkanian.

**Mayvillian** A *\*stage* (453–447.5 Ma ago) of the Late *\*Ordovician* in the Middle *\*Cincinnatian \*series* of N. America, preceded by the *\*Actonian* and followed by the *\*Onnian*.

**mean** In statistics, the *\*average* as calculated by the sum of each data point divided by the total number of data points. *See also* VARIANCE.

**meander** The sinuous trace of a stream channel whose length is normally equal to or greater than 1.5 times the down-valley (or straight-line) distance. It is best developed in cohesive *\*floodplain \*alluvium*. The relationships between its geometric properties vary little with size: e.g. meander wavelength (the straight-line distance between two points at similar positions (e.g. outward extremities of curve) on two successive curves along the trace) is normally 10–14 times the channel width, irrespective of size. Meander origin is uncertain, but the sinuous curve may be that shape best fitted for the transfer of channelled flow in accordance with the *\*least work principle*. Over time, a meander may move laterally and/or vertically. The process of sideways movement is known as 'meander migration'; it involves the deposition of point *\*bars* on the inner sides of bends and erosion on the outer, and is limited to a tract of floodplain called the 'meander belt'. The migration of two adjacent, concave bands may narrow the floodplain between them, and the restriction is a 'meander neck'. This

widens out to form a bulbous feature, the 'meander core', around which the river swings. The surface of a core may show 'meander scrolls', which are low, curved ridges of relatively coarse material lying parallel to the main channel and deposited by the stream. An 'incised meander' results from downcutting, and two types are found. (a) If incision is fairly slow, and sideways movement occurs, the result is an 'ingrown meander'. The slope down which the stream has migrated during incision is called a 'slip-off slope'. (b) When incision is rapid, with mainly vertical erosion, the consequence is an 'entrenched meander'.

**meander belt** See MEANDER.

**meander core** See MEANDER.

**meander migration** See MEANDER.

**meander neck** See MEANDER.

**meander scroll** See MEANDER.

**meander wavelength** See MEANDER.

**mean sea level** The average height of the surface of the sea, for all stages of the tide, over a long (usually 19-year) period, being determined from hourly readings of tidal height. A tidal gauge measures the sea level in relation to the adjacent coast; since the land may also be moving vertically, this does not represent an absolute measure of sea level. Radar altimetry measures the absolute sea level as the distance between the sea surface and the satellite housing the radar transmitter and receiver.

**mean square** See VARIANCE.

**measure** A \*lithostratigraphic term that (in the past) has been used both formally (e.g. Coal Measures, Culm Measures, see FORMAL) and informally (see INFORMAL) to denote a succession of coal-bearing \*strata.

**mechanical layering** See EARTH RHEOLOGICAL LAYERING.

**mechanical weathering** The *in situ* breakdown of \*rocks and minerals by a set of disintegration processes that do not involve any chemical alteration. The chief mechanisms are: crystal growth, including \*gelifraction and \*salt

weathering; *\*hydration* shattering; insolation weathering (see THERMOCLASTIC); and *\*pressure* release.

**medial moraine** See MORAINE.

**median** In statistics, the *\*average* as defined by the central value in a data set when the data set is ordered by value.

**median filter** A spatial filter which reduces noise in an image by substituting the digital number value of each *\*pixel* in an image for the median value calculated from surrounding classes using contours of probability around *\*training* areas using the maximum-likelihood statistic. See also BOX CLASSIFICATION and SPATIAL-FREQUENCY FILTER.

**median network** A specialist type of *\*maximum-parsimony tree* method for phylogenetic reconstruction in which all the most parsimonious trees are represented in a reticulate grid in a three-dimensional perspective. Median networks are applied to data where the degree of *\*homoplasy* is very high and the number of informative sites very low (e.g. with data derived from populations).

**median suture** See CEPHALIC SUTURE.

**median valley (axial rift, axial trough)** The valley which lies along the axis of some oceanic *\*ridges*. From observations from submarines and the interpretation of *\*seismic* records of *\*earthquakes* in the median valley, the inner floor has been found to have scattered volcanoes and *\*black* smokers, with *\*normal* faults dipping towards the inner floor on each side. These faults raise slices of old floor to the *\*ridge* crest, from where it is lowered by further normal faults which *\*dip* away from the ridge axis. Median valleys develop on slower-spreading ridges (e.g. *\*Mid-Atlantic* and *\*Carlsberg*) and are up to 3 km deep.

**Medinian** A N. American *\*stage* (443.5–438 Ma ago) of the Early *\*Silurian* *\*epoch*, preceded by the *\*Cincinnatian* and followed by the *\*Niagaran*.

**mediocris** From the Latin mediocris meaning ‘medium’, a species of *\*cumulus* with limited vertical development and characterized by very slight projections on the upper surface. See also CLOUD CLASSIFICATION.

**Mediterranean climate** Distinctive climatic type, which occurs around latitude 35° N and 35° S, and which is associated with warm-temperate west coasts. Summers generally are hot and dry, winters mild to cool and rainy. The climate is strongly influenced by westerly air streams in winter, and subtropical high pressure in summer. In the type area, the Mediterranean basin, there is a variety of climatic regimes owing to the complex configuration of seas and mountainous peninsulas in the 3000 km incursion into Eurasia. Annual rainfall is broadly 500–900 mm, but less in more continental locations. Other areas of this type include the coasts of Chile in corresponding latitudes, southern California, south-western Africa, and south-western Australia. Cool ocean currents offshore bring lower temperatures to parts of the Chilean and Californian coasts, but rainfall and temperatures are much affected by the differences of slope and elevation inland.

**Mediterranean-type margin** See [ACTIVE MARGIN](#).

**Mediterranean water** A water mass, formed in the arid eastern Mediterranean, that flows westward, sinking in the Algero-Ligurian and Alboran basins to a depth of approximately 500 m due to its high *\*salinity* (36.5–39.1‰). This dense water flows into the *\*Atlantic Ocean* through the relatively shallow Straits of Gibraltar at a depth below 150 m, while above it lighter, Atlantic water flows eastward into the Mediterranean Sea. The Mediterranean water in the Atlantic then sinks to about 1000 m, where it forms a clearly identifiable water mass.

**medium Earth orbit (MEO)** An orbit at an altitude between 2000 km and 35 000 km. Compare [GEOSTATIONARY ORBIT](#); [LOW EARTH ORBIT](#).

**Medusina mawsoni** An early jellyfish. During the uppermost *\*Precambrian*, beach deposits laid down in the Ediacara Hills of southern Australia were covered by a layer of fine muddy sediments. The result was the unique preservation of numerous soft-bodied organisms including jellyfish such as *Medusina mawsoni* and *Medusinites*. See [EDIACARAN FOSSILS](#).

**Medusinites** See [MEDUSINA MAWSONI](#); [EDIACARAN FOSSILS](#).

**mega- (M-)** From the Greek *megas* meaning ‘great’, a prefix meaning ‘very large’. Attached to SI units, it denotes the unit  $\times 10^6$ .

**megabreccia** A **\*breccia** in which individual **\*clasts** may be more than 1 km in their longest dimension.

**megacryst** Applied to the texture of any **\*igneous** or **\*metamorphic rock** which contains large, usually **\*euhedral \*crystals** set in a finer-grained **\*groundmass**. The term has no genetic connotation, unlike 'phenocryst' which implies crystallization from a **\*magma**, and 'porphyroblast' which implies solid-state **\*recrystallization** during **\*metamorphism**.

**Meganeura (giant dragonfly)** (order Odonata, suborder Meganisoptera, family Meganeuridae) Genus of gigantic fossil dragonfly-like insects, from the Upper **\*Carboniferous**. They were probably the largest insects ever to have lived. *M. gracilipes* had a wing-span of 70 cm, and *M. monyi* a wing-span of 60–70 cm.

**megaphyll** See ENATION THEORY.

**megaregolith** The fractured and **\*brecciated** zone of rock formed on the **\*lunar** highland crust due to the intense early bombardment during the period from 4.4 billion years ago, when the crust was forming, to 3.85 billion years ago, when the Imbrium and Orientale collisions marked the close of the massive basin-forming events. Estimates of the depth of the megaregolith extend from 1 km to 25 km, the latter estimate coinciding with an observed seismic boundary.

**megaripple** See DUNE BEDFORM.

**megasclere** See HEXACTINELLIDA.

**megasequence** A stratigraphic sequence of rocks formed by sedimentation in a tectonically produced extensional basin, commonly at a **\*constructive margin**.

**megasplay fault** A large, landward-dipping **\*fault** that cuts through an **\*accretionary wedge**. The fault begins near the top of the subducting plate and ends at the landward end of the wedge.

**megasporangia** See SPORE.

**megaspore** See SPORE.

**megathermal** Applied to the seasonal patterns of river flow in certain equatorial regions, where the main characteristics are high temperatures and evaporation rates all year round, so that river flow maxima reflect the months of maximum precipitation.

**megathermal climate** High-temperature climate type, known more commonly in Europe (e.g. in the **\*Köppen classification**) as humid subtropical or tropical, with the coldest monthly mean temperature above 18 °C. The term is also defined in the **\*Thornthwaite** classification by potential evapotranspiration and moisture-budget criteria.

**Meghalayan** The most recent **\*age**/**\*stage** of the **\*Holocene**, 4250 BP to the present, dated from a **\*dripstone** in Mawmluh Cave, Meghalaya, India; Global Auxiliary **\*Stratotype**, Mount Logan Ice Core, Canada. The beginning of this age coincided with the abrupt onset of a 200-year drought and climatic cooling that affected agricultural societies throughout the world, linked to the collapse of civilizations and major migrations in Egypt, Greece, Syria, Palestine, Mesopotamia, the Indus Valley, and the Yangtze River Valley.

**Megha-Tropiques** See METEOROLOGICAL LEO OBSERVATIONS IN THE INTERTROPICAL ZONE.

**meionite** See SCAPOLITE.

**Meishuchuan** A Chinese **\*stage** (542–532 Ma ago) of the Early Cambrian **\*epoch**, preceded by the **\*Sinian** and followed by the **\*Qungzusian**.

**mela-** A prefix attached to the name of an **\*igneous** rock when the rock has a darker colour than is usual. For example, a **\*syenite** which is darker than normal is called a ‘melasyenite’.

**mélange** A mappable body of rock composed of broken rock fragments, of all sizes and many origins, in a sheared **\*matrix**. A **mélange** with a chaotic nature, initially of sedimentary origin, is called an ‘**\*olistostrome**’. Tectonic **mélanges** are thought to form in **\*subduction** zones at shallow depth.

**melanocratic** Applied to **\*igneous** rocks whose **\*colour index** is between 60 and 90. The high colour index of such rocks is due to the presence of a high proportion of **\*ferromagnesian minerals**.

**melanosome** Dark-coloured bands rich in \*mafic and aluminous minerals, e.g. \*biotite, \*sillimanite, and \*garnet, which are found between multiple coarse-grained quartzofeldspathic veins in regionally metamorphosed (see REGIONAL METAMORPHISM) \*pelites and \*psammites. The melanosomes represent planar regions which have undergone extreme shortening by dissolution (or melting), and where removal of the \*quartz and \*feldspar components of the original rock along non-penetrative \*cleavage planes developed as a result of high \*shear stresses imposed during the regional metamorphic event. The mafic and aluminous components of the melanosome represent the undissolved or unmelted mineral residue left over from the original rock.

**Melbournian** A \*stage (428.2–416 Ma ago) of the Upper \*Silurian of south-eastern Australia, underlain by the \*Eildonian.

**Melekesskian** A Russian \*stage (313.4–311.7 Ma ago) in the Early \*Pennsylvanian sub-epoch, preceded by the \*Cheremshanskian and followed by the \*Vereiskian (\*Moscovian stage).

**melilite** A member of the melilite group of \*minerals, that has the general formula  $(\text{Mg,Al}) (\text{Ca,Na})_2 (\text{Si,Al})_2 \text{O}_7$ . There is a \*solid solution series between gehlenite (aluminium-rich) and åkermanite (magnesium rich); melilite occupies an intermediate position, with some substitution of sodium for calcium, and of aluminium for silicon or magnesium. Sp. gr. 2.9; \*hardness 5.5; white or greenish-white; granular or in \*tabular \*crystals; occurs in \*basic lavas, metamorphosed \*limestones, and furnace slags.

**melilitite** A dark-coloured, \*extrusive, \*ultrabasic, \*alkaline \*igneous rock containing \*essential \*melilite, \*olivine, and \*nepheline. The active alkaline \*volcano Nyiragongo in the Democratic Republic of Congo is famous for its melilitite \*lavas which are found associated with other ultra-alkaline lavas, e.g. \*leucitites and \*nephelinites.

**melt** Liquid part of \*silicate \*magma. Magma may consist solely of melt, or of melt and suspended crystals. ‘Wet’ melt contains dissolved \*volatile constituents, mainly water. Conversely, anhydrous melts, without volatiles, are referred to as ‘dry’ melts.

**melteigite** An undersaturated, \*intrusive \*igneous rock containing essential \*nepheline and alkali \*pyroxene (\*aegirine or aegerine-\*augite) and

displaying a *\*colour index* of 70–90. With decreasing *\*mafic* *\*mineral* content and increasing nepheline content melteigites pass into *\*ijolites* and then into *\*urtites*. All three rock types are essentially *\*feldspar*-free, undersaturated *\*syenites* (see *SILICA SATURATION*) and are well displayed in the Fen intrusive complex of southern Norway.

**meltemi** See *ETESIAN WINDS*.

**meltout till** See *TILL*.

**meltwater channel** See *GLACIAL DRAINAGE CHANNEL*.

**MEM** See *MICROEROSION METER*.

**member** See *FORMATION*.

**membrane stress** The stress on a tectonic *\*plate* resulting from its motion over an *\*Earth* with a non-spherical shape. It is largely restricted to plates with a latitudinal motion, i.e. moving away from, or towards, the equatorial bulge and therefore adjusting to the new curvature.

**Menapian** A glacial *\*stage* (0.9–0.8 Ma ago) of the northern European plain, perhaps equivalent to the *\*Günz* glacial of the Alps and the *\*Kansan* of N. America.

**mendip** Hill or ridge that has been buried by younger rocks and subsequently exposed by *\*erosion*. The type example is provided by the Mendip Hills, England, described by W. M. Davis in 1912, but today the term is used only very rarely.

**Menevian** See *MAYAN*.

**mensa** (*pl.* mensae) A *\*mesa* on the surface of an extraterrestrial body.

**MEO** See *MEDIUM EARTH ORBIT*.

**Meramecian** A *\*stage* (340–333 Ma ago) in the Middle *\*Mississippian* of N. America, underlain by the *\*Osagean*, overlain by the *\*Chesterian*, and roughly contemporaneous with the Arundian, Holkerian, and Asbian stages of the *\*Visean* series of Europe.

**Mercalli, Giuseppe** (1815–1914) An Italian professor of natural sciences who studied the major *\*earthquake* districts of Italy, Mercalli is best known

for his scale of earthquake intensity, first formulated in 1897, and refined several times. *See also* [MERCALLI SCALE](#).

**Mercalli scale** An *\*earthquake-intensity* scale based on direct observation. It ranges from scales I (not felt) and II (felt by most people at rest), to VII (difficult for a person to remain standing), X (most structures destroyed), and XII (total devastation). It is not a reliable scale for energy release because the extent of destruction depends on the local geology, type of buildings, etc. *See also* [MOMENT MAGNITUDE SCALE](#); [RICHTER SCALE](#).

**Mercurial geologic timescale** The standardized timescale for the planet *\*Mercury*.

Period	Epoch	Dates (Ma ago)
Pretolstojan		4568–3900
Tolstojan		3900–3850
Calorian		3850–3000
	Early Calorian	3850–3800
	Late Calorian	3800–3000
Mansurian		3000–1000
Kuiperian		1000–0

**Mercury** The innermost planet of the *\*solar system*, 0.387 AU from the Sun, its distance from Earth varying from  $77.3 \times 10^6$  km to  $221.9 \times 10^6$  km. It has a tenuous atmosphere, with a surface pressure of about  $10^{-15}$  bar, composed of oxygen (42%), sodium (29%), hydrogen (22%), helium (6%), potassium (0.5%), and possibly trace amounts of argon, carbon dioxide, water, nitrogen, xenon, krypton, and neon (the proportions are approximate). Its diameter is 4880 km; volume  $6.085 \times 10^{10}$  km<sup>3</sup>; mass  $0.3302 \times 10^{24}$  kg; mean density 5427 kg/m<sup>3</sup> (making it the densest of the planets); surface gravity 3.7 (Earth = 1); visual albedo 0.11; average surface temperature 440 K (on the sunward side ranging from 590 K to 725 K). *See also* [MERCURIAL GEOLOGIC TIMESCALE](#).

**Mercury Orbiter** An *\*ISAS/\*NASA* mission to place a vehicle in orbit about Mercury, planned to launch in 2009, but cancelled.

**meridional circulation (meridional flow)** A movement of air or water in a direction at right angles to lines of latitude, i.e. parallel to the meridians

(lines of longitude). *See also* [HADLEY CELL](#); [ZONAL FLOW](#).

**meridional overturning circulation** A system of ocean currents that plays a major role in the transport of heat from low to high latitudes and, therefore, in global climates. It is driven by the convective overturning of water near the edge of the northern sea ice, where cold, saline water sinks to the ocean floor and travels south as the [\\*North Atlantic Deep Water](#), eventually to the Southern Ocean, where it joins the [\\*Antarctic Circumpolar Current](#). Part of the current turns north into the Indian Ocean, rising to become an intermediate current flowing about 1000 m below the surface. The current crosses the equator, follows a clockwise path in the N. Pacific, continues westward through the islands of Indonesia, crosses the equator once again, crosses the Indian Ocean, flows around Africa, then northward, crossing the equator for the fourth time and returning to the N. Atlantic and the edge of the northern sea ice. It is sometimes known informally as the Atlantic Conveyor.

**Merioneth** A [\\*stage](#) (501–488.3 Ma ago) of the Late [\\*Cambrian](#), underlain by the Middle Cambrian and overlain by the [\\*Tremadocian](#) (Early [\\*Ordovician](#)).

**Merionsian** A [\\*stage](#) in the Lower [\\*Devonian](#) of Australia, underlain by the [\\*Crudinian](#), overlain by the [\\*Cunninghamian](#), and roughly contemporaneous with the [\\*Siegenian](#) of Europe.


**MERLIN** *See* [METHANE REMOTE SENSING LIDAR MISSION](#).

**merocrystalline** *See* [HYPOCRYSTALLINE](#).

**meromictic** Applied to lakes whose waters are stratified permanently, usually because of some chemical difference (e.g. contrasting salinities, and hence densities) between [\\*epilimnion](#) and [\\*hypolimnion](#) waters. *Compare* [HOLOMICTIC](#).

**Merostomata** (phylum [\\*Arthropoda](#)) Class which includes the king crabs (horseshoe crabs) and the extinct eurypterids or ‘water scorpions’. Eurypterids were usually 10–20 cm in length, but some were much larger, e.g. *Pterygotus* which grew up to 2 m long. They ranged from the [\\*Ordovician](#) to the end of the [\\*Palaeozoic](#). The horseshoe crabs also appeared in the Lower Palaeozoic and have survived to the present.

**mesa** Flat-topped hill of limited extent, but wider than a **\*butte** and normally underlain by near-horizontally bedded sediments. A land-form similar to a mesa but larger is called a 'plateau'.



**Mesa**

**mesentery** In corals, a radial infolding of the fleshy body wall, forming a partition in the gut. In this central cavity the mesenteries alternate with the skeletal **\*septa**.

**mesic** Applied to an environment that is neither extremely wet (hydric) nor extremely dry (xeric). *See also* **PERGELIC**.

**Mesoarchaean** An **\*era**, 3200–2800 Ma ago, of the **\*Archaean eon**, preceded by the **\*Palaeoarchaeon** and followed by the **\*Neoarchaeon** eras. The first direct evidence of life is from this time, and the earliest (Pongola) **\*glaciation** occurred.

**mesoclimate** General term applied to the characteristics of a relatively small region, e.g. a valley or urban area.

**mesocratic** Applied to **\*igneous** rocks with a **\*colour index** between 30 and 60. They are intermediate in colour between **\*leucocratic** and **\*melanocratic** rocks.

**mesocyclone** A region of rapidly rotating air, up to 10 km in diameter, inside a large **\*cumulonimbus** cloud. Rotation commences near the middle of the cloud and extends downwards. If it continues beneath the cloud base it appears as a **\*funnel cloud**, and if it reaches to the ground it is a **\*tornado**.

**mesohaline water** *See* **HALINITY**.

**mesokurtic** *See* **KURTOSIS**.

**mesopause** In the atmosphere, the inversion at about 80 km height, which separates the *\*mesosphere* from the *\*thermosphere* above. The first 10 km of the mesopause are almost isothermal. *See also* [ATMOSPHERIC STRUCTURE](#).

**Mesoproterozoic** The middle part of the *\*Proterozoic*, lasting from about 1600–1000 Ma ago.



<http://www.palaeos.org/Mesoproterozoic>

- The Mesoproterozoic Era of the Proterozoic Eon.

**Mesosauria** Order of *\*anapsid* reptiles, with just one family, the Mesosauridae. The mesosaurs (e.g. *Mesosaurus*) are known only from late *\*Carboniferous* or early *\*Permian* rocks in S. America and S. Africa. They were adapted to life in fresh water, lightly built, and up to one metre in length.

**mesosphere** **1.** Upper-atmospheric layer above the *\*stratopause* (at 50 km) through which temperature decreases with height up to about 80 km, where temperatures reach a minimum of about  $-90\text{ }^{\circ}\text{C}$ . This level is the *\*mesopause*, an inversion above which temperatures rise again. *See also* [ATMOSPHERIC STRUCTURE](#). **2.** That part of the *\*Earth* underlying the *\*asthenosphere*. The term is no longer in current use in this sense.

**mesothem** *See* [CYCLOSTRATIGRAPHY](#).

**mesothermal** **1.** Applied to the seasonal patterns of river discharge of certain warm, subtropical, and temperate areas. **2.** Applied to a *\*mineral* deposit formed by hot, ascending solutions at about  $200\text{--}300\text{ }^{\circ}\text{C}$  at moderate depths in the Earth's *\*crust*.

**mesothermal climate** Climatic type with moderate temperatures, known most commonly in Europe, e.g. in the *\*Köppen* classification, as a warm-temperate rainy climate having a coldest month with temperatures of  $-3\text{ }^{\circ}\text{C}$  to  $+18\text{ }^{\circ}\text{C}$  and a warmest month above  $+10\text{ }^{\circ}\text{C}$ . Such climates are found typically in latitudes  $30\text{--}45^{\circ}$ . The *\*Thornthwaite* classification defines this type according to *\*evapotranspiration* and moisture budget.

**mesotidal** Applied to coastal areas where the **\*tidal range** is 2–4 m. Tidal action and wave activity both tend to be important in such areas.

**mesotype** A term applied to **\*igneous** rocks displaying a **\*colour index** between 30 and 60.

**Mesozoic** The middle of three **\*eras** that constitute the **\*Phanerozoic** period of time. The Mesozoic (literally ‘middle life’) was preceded by the **\*Palaeozoic** era and followed by the **\*Cenozoic** era. It began with the **\*Triassic** approximately 251 Ma ago and ended around 65.5 Ma ago at the start of the Cenozoic. The Mesozoic comprises the Triassic, **\*Jurassic**, and **\*Cretaceous** periods.



<http://scienceviews.com/dinosaurs/mesozoic.html>

- The Mesozoic Era.

**Messenger probe** A **\*NASA** mission to **\*Mercury** that was launched on 3 August 2004, entered Mercury orbit on 17 March 2011, and completed its mission on 30 April 2015, when it crashed into the planet’s surface at more than  $3.91 \text{ km sec}^{-1}$ .

**Messinian** The final **\*stage** (7.246–5.332 Ma ago) in the **\*Miocene \*epoch**, preceded by the **\*Tortonian**, followed by the **\*Zanclean** (Tabianian) (Int. Commission on Stratigraphy, 2004). It is roughly contemporaneous with the upper **\*Mohnian** and lower **\*Delmontian** (N. America), upper **\*Tongaporutuan** and lower **\*Kapitean** (New Zealand), and part of the **\*Mitchellian** (Australia). The **\*neostatotype** is found between Caltanissetta and Enna, Sicily. It is an important stage, marked by the presence of thick **\*evaporite** deposits in the Mediterranean. These may indicate that the Straits of Gibraltar were closed approximately 6.5–5.1 Ma ago and that the Mediterranean was reduced to a series of evaporite basins.

**meta-** From the Greek *meta* meaning ‘with’ or ‘after’, a prefix implying change and meaning ‘behind’, ‘after’, or ‘beyond’. The prefix may be attached to the name of any rock which has undergone **\*metamorphism**. For instance, a **\*basalt** which has been metamorphosed may be termed a metabasalt, and a pelitic rock (see **PELITE**) which has suffered metamorphism may be termed a metapelite.

**metabasalt** See META-.

**metacarpal** In the forelimb of a **\*tetrapod**, one of the rod-like bones that articulate proximally with the carpus (wrist) and distally with the phalanges (digits). In humans, the metacarpals occupy the palm region of the hand.

**metacryst** See PORPHYROBLAST.

**metagenesis** Following **\*catagenesis**, a phase in the formation of **\*petroleum** and **\*natural gas** during which the temperature rises above 150 °C and may exceed 200 °C, causing destruction of **\*kerogen** and release of gas (mainly methane); at the higher temperatures petroleum is destroyed, leaving only natural gas.

**metal factor (MF)** A **\*frequency-domain** measure of **\*induced polarization** where **\*apparent resistivities** are measured at two low frequencies, usually a factor of 10 apart in magnitude ( $\rho_{dc}$  and  $\rho_{ac}$ );  $MF = 2\pi 10^5 (\rho_{dc} - \rho_{ac}) / \rho_{ac}^2$ .

**metalimnion** In a thermally stratified lake, the layer of water between the **\*epilimnion** and **\*hypolimnion**.

**metallic** Of mineral **\*lustre**, having the sheen characteristic of a metal.

**metallic bond** A special type of **\*covalent** bond found in such solids as native **\*copper**. Every atom shares its **\*electrons** with each of its neighbours in turn. The sharing atoms must be able to move and so in metals they appear to be positive **\*ions** moving in a sea of electrons.

**metallogenesis** Study of the origin of **\*ore** deposits and of the interdependence in time and space of this process with other geologic processes such as **\*tectonics**.

**metallogenic province** An area of characteristic mineralizing activity, or a particular association of **\*mineral** deposits. It may contain several episodes of mineralization.

**metaluminous** Applied to **\*igneous** rocks in which there are fewer molecules of  $Al_2O_3$  than of  $(CaO + Na_2O + K_2O)$ . ‘Metaluminous’, together with the terms ‘peraluminous’ and ‘**\*peralkaline**’, describe the alumina saturation of an igneous rock, portraying variations in what is usually the second most abundant oxide component in the rock (after silica).

**metameric segmentation** The repetition of organs and tissues at intervals along the body of an animal.

**metamorphic aureole** See CONTACT AUREOLE.

**metamorphic differentiation** Separation of an originally homogeneous rock into bands of contrasting mineralogy during high-grade \*metamorphism (see METAMORPHIC GRADE). The bands of contrasting mineralogy may be produced either by solid state \*diffusion of elements during the metamorphic event, or by local \*metasomatism.

**metamorphic facies** A set of metamorphic \*mineral assemblages derived from rocks of contrasting composition which have been subjected to the same grade or conditions of \*metamorphism. For example, a metamorphosed \*shale, \*basic \*lava, and \*limestone which are found adjacent to each other in a metamorphic \*terrain must have been subjected to the same grade of metamorphism, yet each of these contrasting rock types displays a different metamorphic mineral assemblage, the individual assemblages reflecting both the starting rock composition and the grade of metamorphism. Since all three rocks have been subject to the same grade of metamorphism, the set of contrasting mineral assemblages reflects only the contrasts in rock composition, and thus constitutes a metamorphic \*facies. Any change in the mineral assemblage observed in a particular rock composition represents a mineralogical response to changing metamorphic conditions and thus would define a new facies. The metamorphic conditions represented by a particular facies can be deduced from experimental studies of the overlapping pressure–temperature stability fields of mineral assemblages in that facies. However, the definition of a facies is purely descriptive and based entirely on the mineral assemblages observed. The concept of metamorphic facies was first proposed by the Finnish petrologist P. E. \*Eskola in 1920 after he had compared the mineral assemblages in similar rock compositions from Oslo, Norway, and Orijarvi, Finland, two areas which had been subjected to contrasting metamorphic conditions.

**metamorphic grade** A measure of the relative intensity of \*metamorphism. In pelitic rocks (see PELITE), an increase of metamorphic grade is marked by a progressive dehydration of the mineral assemblages present whilst in \*limestones and impure limestones an increase in grade is marked by progressive decarbonation of the mineral assemblages present.

In general, mineral assemblages stable at progressively higher pressures and temperatures characterize an increase in metamorphic grade (e.g. [\\*garnet grade](#)).

**metamorphic rock** An aggregate of [\\*minerals](#) formed by the [\\*recrystallization](#) of pre-existing rocks in response to a change of pressure, temperature, or [\\*volatile](#) content. Metamorphic rocks can generally be divided into four types: (a) regional metamorphic rocks, formed in response to changes leading to high temperature and high pressure ([\\*shearing stress](#) and hydrostatic pressure: see [HYDROSTATIC STRESS](#)) accompanying orogenic events (see [OROGENY](#); [REGIONAL METAMORPHISM](#)); (b) contact metamorphic rocks, formed in response to changes leading to high temperature (with low pressure) around an [\\*igneous \\*intrusion](#) (see [THERMAL METAMORPHISM](#)); (c) cataclastic or dynamic metamorphic rocks, formed in response to an increase in directed pressure (shearing stress) particularly in [\\*fault](#) and [\\*thrust](#) zones (see [CATACLASITE](#)); and (d) burial metamorphism, formed in response to changes leading to high pressure (with low temperature).

**metamorphic rock classification** The arrangement of [\\*metamorphic rocks](#) into types according to the deformation they experienced, in particular whether or not they exhibit [\\*foliation](#), indicating whether they were subjected to directed pressure or [\\*shear stress](#), although some do not exhibit foliation because their [\\*minerals](#) do not show alignment. Rocks are also classified by [\\*metamorphic grade](#) as very low, low, medium, and high. The table shows a selection of parent rocks and the changes effected by [\\*metamorphism](#).

Parent rock	Very low grade 150–300 °C	Low grade 300–450 °C	Medium grade 450–550 °C	High grade Above 500 °C
Mudrock	Slate	Phyllite	Schist	Gneiss
Granite	No change	No change	No change	Granite gneiss
Basalt	Chlorite schist	Chlorite schist	Amphibolite	Amphibolite
Sandstone	No change	Little change	Quartzite	Quartzite
Limestone	Little change	Marble	Marble	Marble



<https://opentextbc.ca/geology/chapter/7-2-classification-of-metamorphic-rocks/>

- Classification of Metamorphic Rocks.

**metamorphic zone** The area between two successive *\*isograds* in a metamorphic terrain. The zone is named after the lower-grade isograd (see *METAMORPHIC GRADE*). For example, the area lying between the *\*garnet* isograd and the *\*kyanite* isograd for pelitic rocks in a *\*regional metamorphic* terrain is termed the garnet zone. The zone defines a set of pressure, temperature,  $P(\text{H}_2\text{O})$ , and  $P(\text{CO}_2)$ , over which there were no mineralogical reactions in the particular rock composition concerned. Normally, the chemical composition of *\*pelites* is used as a marker.

**metamorphism** The process of changing the characteristics of a rock in response to changes in temperature, pressure, or *\*volatile* content. Most metamorphic changes do not include bulk chemical changes, but merely the crystallization of new mineral *\*phases*. These isochemical changes cause major textural changes. Compare *METASOMATISM*. See also *BARROVIAN-TYPE METAMORPHISM*; *BARROW'S ZONES*; *BURIAL METAMORPHISM*; *DYNAMIC METAMORPHISM*; *REGIONAL METAMORPHISM*; *THERMAL METAMORPHISM*; *METAMORPHIC GRADE*.

**metapelite** See *META-*.

**Metaphyta** See *PLANTAE*.

**metaquartzite** See *QUARTZITE*.

**metasediment** See *META-*.

**metaseptum** (*pl.* metasepta) See *SEPTUM*.

**metasilicate** See *CYCLOSILICATE*.

**metasomatism** Kind of *\*metamorphism* that involves the introduction of chemical constituents into a rock, or their removal from it (or both), via a *\*volatile* phase. Complete *\*mineral* transformations may occur, but the original rock *\*texture* may remain (resulting in a *\*pseudomorph*).

**metaspecies** Ancestral species. They are not **\*monophyletic** (in the strictest sense) so do not, and cannot, conform to the canon that taxa must always be monophyletic.

**metastable** Applied to a **\*phase** which is apparently stable but is capable of reaction if disturbed, a state usually due to the slowness of a system to attain equilibrium. The term is applied, for example, to supersaturated solutions. A phase is said to be metastable if it exists in the same temperature range in which another phase with lower **\*vapour pressure** is stable. Many minerals that occur at room temperature and pressure are metastable (e.g. **\*diamond**) and most **\*metamorphic rocks** formed under high temperatures and pressures are metastable at the surface under normal temperatures and pressures. A metastable system (mineral assemblage) is in a temporary state of equilibrium and requires only a minimal disturbance to initiate a change to a state of true equilibrium. A **\*gabbro** mineral assemblage which crystallized at high temperatures and moderate pressures, and is thus in equilibrium under these conditions, would be in a metastable state when exposed by **\*erosion** to the low-temperature, low-pressure environment at the Earth's surface. Rainwater falling on the exposed mineral assemblage would act as the disturbance or catalyst initiating the reaction of the gabbro mineral assemblage to a low-temperature, low-pressure, **\*chlorite–\*clay** mineral assemblage, thus re-establishing the equilibrium.

**metatarsal** In the hind limb of a **\*tetrapod**, one of the rod-like bones that articulate proximally with the tarsus (ankle) and distally with the phalanges (toes).

**Metazoa** See **ANIMALIA**.

**metazoan** A multicellular animal. See **ANIMALIA**.

**meteor** Transient, incandescent trail of a **\*meteoroid** entering the **\*Earth's \*atmosphere**. All the material burns up before reaching the ground. It is popularly referred to as a 'shooting star'. A very bright meteor is called a fireball. Compare **METEORITES**.

**Meteor** The generic name for a family of Russian meteorological satellites in low Earth orbit. Meteor-1 was launched in 1969. The most recent, Meteor-M2, was launched on 8 July 2014.



<http://www.russianspaceweb.com/meteor.html>

- Meteor Spacecraft Family.

**Meteor-3M** A planned series of Russian spacecraft, the first of which was launched on 10 December 2001 from Baikonur Cosmodrome, Kazakhstan, into a *\*Sun-synchronous* orbit at a height of about 1020 km in a joint programme between *\*NASA* and the Russian Federal Space Agency (RSA) to measure ozone, *\*aerosols*, water vapour, and other key components of the atmosphere, as well as monitoring temperature and trace gases in the *\*stratosphere* and *\*mesosphere*. Meteor-3M carries NASA's Stratospheric Aerosol and Gas Experiment (SAGE III) as well as other instruments and is designed to last for five years, but technical problems in March 2006 left it almost completely disabled. Meteor-3M (2) and Meteor-3M (3) were cancelled.



<https://directory.eoportal.org/web/eoportal/satellite-missions/m/meteor-3m-1>

- A joint NASA and RSA mission in Sun-synchronous orbit to measure ozone, aerosols, and water vapour in the Earth's atmosphere.

**meteoric diagenesis** The changes (*\*diagenesis*) that take place in marine carbonate sediments buried at a shallow depth when they are exposed to *\*meteoric water*. These usually involve dissolution, as acid water dissolves *\*aragonite*, followed by precipitation, mainly of *\*calcite*, as the water becomes saturated with respect to  $\text{CaCO}_3$ , and the formation of soils, especially *\*caliche*.

**meteoritic abundance of elements** The relative abundances of elements contained within chondritic *\*meteorites* (see **CHONDRITE**), and in particular within C1 *\*carbonaceous* chondrites, are believed to provide the best estimate of the composition of the primitive material in the *\*solar* system from which the planets formed. Comparison of C1 carbonaceous chondrites with the *\*solar* abundances of elements (obtained by spectroscopic studies) show that, for non-*\*volatile* elements, the abundances are almost identical.

On the other hand, the composition of ordinary chondrites is believed to be similar to the **\*bulk** composition of the Earth.

Element	Average chondrite (wt. %)
O	33.24
Fe	27.24
Si	17.10
Mg	14.29
S	1.93
Ni	1.64
Ca	1.27
Al	1.22
Na	0.64
Cr	0.29
Mn	0.25
P	0.11
Co	0.09
K	0.08
Ti	0.06

**meteoric water** Water of atmospheric origin which reaches the Earth's **\*crust** from above, either as rainfall or as seepage from surface-water bodies. *See also* **GROUNDWATER**.

**meteorites** Small, extraterrestrial bodies, most of which probably originate in the asteroid belt, that enter the **\*Earth's \*atmosphere** and land on the surface. Most are only a few centimetres in size. Meteorites are classified into four main groups according to their composition and structure as: **\*chondrites; \*achondrites; \*stony-irons; and \*irons**. *See also* **ANTARCTIC METEORITE; AUBRITE; BASALTIC METEORITES; CARBONACEOUS CHONDRITE; EUCRITE; SHERGOTTITE/NAKHLITE/CHASSIGNITE METEORITES; STONY METEORITE; TEKTITE**.

**meteoroid** Small, extraterrestrial body, within the **\*solar** system, that may enter the **\*Earth's \*atmosphere** if its orbit around the **\*Sun** crosses that of the **\*Earth**. Strictly, at the point where a meteoroid enters the atmosphere it is referred to as a **\*meteor**; if it reaches the Earth's surface it is known as a **\*meteorite**.

**Meteorological LEO Observations in the Intertropical Zone (Megha-Tropiques)** A collaborative mission between the space agencies of India

and France that studies convective systems affecting the [\\*intertropical convergence zone](#), particularly between latitudes 10° and 20°. The [\\*microsatellite](#) was launched on 12 October 2011, from Sriharikota, India, into a circular orbit at an altitude of 865 km.

**Meteorological Operational Satellite Program of Europe (MetOp)** A series of three satellites to be launched over 14 years that will supply data for operational meteorology. The first, MetOp-A, was launched on 19 October 2006, MetOp-B on 17 September 2012, both from Baikonur, Kazakhstan, into a near-circular, Sun-synchronous [\\*polar orbit](#) at a mean altitude of 817 km. MetOp-C is scheduled for launch in November 2018, from Kourou, French Guiana. A fourth satellite, MetOp-SG (second generation), is planned to replace the first generation satellites between 2021 and 2023.

**Meteorological Satellite-1** See [KALPANA-1](#).

**Meteosat** A series of several experimental and four operational meteorological satellites launched by the European Space Agency ([ESA](#)) and operated by Eumetsat on behalf of the ESA. The first was launched in 1972 by [\\*NASA](#) and the fourth, Meteosat-7, in 1997 by the ESA. Four Meteosat Second Generation satellites were launched between 2002 and 2012 and will remain operational until 2020, when they will be succeeded by Meteosat Third Generation satellites. The first Third Generation satellite is scheduled for launch in 2021.



<https://www.eumetsat.int/website/home/Satellites/CurrentSatellites/Meteosat/index.html>

- [Meteosat](#).

**methane hydrates** See [NATURAL GAS HYDRATES](#).

**Methane Remote Sensing Lidar Mission (MERLIN)** A joint Franco-German [\\*minisatellite](#) mission that will obtain spatial and temporal gradients of atmospheric methane. It is scheduled for launch in 2021 or 2022.

**methanogen** A single-celled organism, belonging to domain [\\*Archaea](#), that produces [\\*methane](#) gas as a product of its metabolism.

**Metis (Jupiter XVI)** The innermost of the known jovian planets, discovered in 1979. It orbits inside the main ring of *\*Jupiter* (it is a *\*moon*); it and Adrastea may be the source of the material comprising the ring. Metis has a diameter of 40 ( $\pm 20$ ) km; mass  $9.56 \times 10^{16}$  kg; mean distance from Jupiter 128 000 km.

**metocean data** Measurements of oceanographic and meteorological conditions, including currents, waves, air temperature, surface atmospheric pressure, and wind speed and direction.

**MetOp** See METEOROLOGICAL OPERATIONAL SATELLITE PROGRAM OF EUROPE.

**MetOp-SG** See METEOROLOGICAL OPERATIONAL SATELLITE PROGRAM OF EUROPE.

**MetSat-1** See KALPANA-1.

**MF** See METAL FACTOR.

**M-fold** The ideal form of a *\*minor* fold which occurs in the *\*hinge* region of a major *\*antiform* (a *\*synform* is distinguished by W-fold forms). In practice it is difficult to recognize ideal M-folds and they are less informative than *\*Z-fold* and *\*S-fold* configurations.

**Mg#** See MAGNESIUM NUMBER.

**Miacidae** See CARNIVORA.

**miarolitic** Applied to a texture found in shallow, level, *\*plutonic*, *\*igneous* rocks and characterized by void spaces or pockets with outlines shaped by neighbouring crystals. The texture is primary representing entrapment of gas segregated by vesiculation (see **VESICLE**) of the late-stage *\*magma* during the final stages of crystallization of the *\*intrusive* body.

**micas** An important group of *\*phyllosilicates* (sheet silicates) with a 2:1 atomic structure (see **CLAY MINERALS**) and including the important *\*minerals* *\*muscovite*, *\*biotite*, and *\*phlogopite*; the group is characterized by the silicon oxygen tetrahedral layers of composition  $[\text{Si}_4\text{O}_{10}]_n$  and a general composition may be written  $(\text{K,Na})_2\text{Y}_6 [\text{Z}_4\text{O}_{10}]_2(\text{OH,F})_4$ , where Y

= Mg, Fe, Fe<sup>3+</sup>, or Al, and Z = Si or Al; the micas include the common micas already mentioned, plus *\*glaucosite*, *\*lepidolite*, and *\*zinnwaldite*, the brittle micas (see *MARGARITE*), and also the related minerals *\*talc*, *\*stilpnomelane*, and *\*pyrophyllite*.

**Michel–Lévy chart (birefringence chart, interference colour chart)** A chart of standard colours used in measuring *\*birefringence*. In crossed-polarized light, *\*thin* sections of *\*anisotropic \*minerals* with a standard thickness of 0.03 mm (30 µm) will give a series of *\*interference colours*, depending on the amount of birefringence (double refraction). A number of sections are examined in different orientations and the maximum interference colour matched with the equivalent colour on the chart. From the point representing 30 µm thickness on the appropriate colour a radiating line is followed to the edge of the chart to give a reading for birefringence. The accessory *\*quartz* wedge may also be used to determine birefringence.

**Michell, John** (?1724–93) An astronomer and experimental philosopher from Cambridge, Michell made the first scientific investigations of *\*seismic waves*. He studied *\*earthquake* phenomena, especially the Lisbon earthquake of 1755. He proposed a theory for the motion of seismic waves, estimating their velocity, and showed how to determine the *\*epicentre* of an earthquake.

***Micraster*** A genus of Upper *\*Cretaceous* echinoids (*\*Echinoidea*) that is common in France and England and occurs in much of Europe. Much has been written about evolution in this genus, starting with A. R. Rowe in 1899, and adaptations in the genus either to burrowing or to varying substrates have also been suggested.

**micrinite** See *COAL MACERAL*.

**micrite 1. (lime mud)** Microcrystalline *\*calcite*, with a grain size finer than 4 µm. This may originate from *\*biogenic* sources in the form of aragonitic calcareous *\*algae*, by bioerosion and physical erosion of coarser *\*carbonate* fragments, inorganic precipitation from carbonate saturated waters, and biochemical precipitation by algal activity. **2. \*Limestones** consisting mainly of lithified *\*lime mud*. See *FOLK LIMESTONE CLASSIFICATION*; *LEIGHTON–PENDEXTER CLASSIFICATION*.

**-micrite** See *FOLK LIMESTONE CLASSIFICATION*.

**micrite envelope** A dark-coloured, fine-grained, \*carbonate exterior found around carbonate skeletal fragments. It is produced by the alteration of the original mineralogy of the fragment to \*micrite, commonly caused by the boring into the skeletal material of certain \*algae. In some micrite envelopes the tiny, tube-like borings of the algae are preserved. Some micrite envelopes are formed by coatings of filamentous algae over the grain. If the development of the micrite envelope proceeds to an advanced stage, the primary structure of the skeletal fragment will be totally replaced by micrite, leaving an \*amorphous \*peloid, or lump (see INTRACLAST).

**micritic limestone** See LEIGHTON—PENDEXTER CLASSIFICATION.

**micritization** The formation of \*micrite by the boring into skeletal \*carbonate particles by cyanobacteria (blue-green algae), and the subsequent precipitation of micrite within the borings. Micritization can also occur if filamentous endolithic algae (i.e. those within the sediment) coat the grain. See MICRITE ENVELOPE.

**micro- (m)** **1.** From the Greek *mikros* meaning ‘small’, a prefix meaning ‘extremely small’. Attached to SI units it denotes the unit  $\times 10^{-6}$ . **2.** In Earth sciences, micro- is a prefix applied in the strict sense to very fine \*igneous textures. Individual particles are likely to be below the resolution of the naked eye so they can be resolved only with the aid of a petrological microscope (see POLARIZING MICROSCOPE). \*Porphyritic texture which can only be resolved by using such a microscope is termed ‘microporphyritic’ texture and the \*phenocrysts are ‘microphenocrysts’. **3.** The prefix may also be applied to igneous rock names. For example, a \*syenite with a \*grain size below that normally accepted for a syenite but above that expected of its fine-grained equivalent, a \*trachyte, may be termed a microsyenite.

**Microaccelerometric Measurements of Satellite Accelerations (MIMOSA)** A \*microsatellite mission of the Czech Republic that studies variations in density distributions in the upper \*ionosphere by measuring orbital perturbations due to atmospheric drag and solar radiation pressure. The satellite was launched on 30 June 2003, from Plesetsk, Russia, into a Sun-synchronous orbit at a \*perigee of 326 km.

**microbialite** A sedimentary body formed on the bed of a lake from the remains of benthic (see BENTHOS) communities of algae and

\*cyanobacteria.

**MicroCarb** See CARBON DIOXIDE MONITORING MISSION.

**microclimate** The atmospheric characteristics prevailing within a small space, usually in the layer near the ground that is affected by the ground surface. Special influences include the impact of vegetation cover on humidity (by \*evapotranspiration) and on temperature and winds. See also URBAN CLIMATES.

**microcline** See ALKALI FELDSPAR.

**microcontinent** See MICROPLATE.

**microcraters** See 'ZAP PITS'.

**microcrystalline** Applied to a crystalline texture which is so fine grained that the individual \*crystals are too small to be observed by the naked eye but may be distinguished under the microscope.

**microdiorite** A medium-grained \*igneous rock characterized by the \*mineral assemblage and chemical composition of \*diorite.

**microearthquake** See MICROSEISM.

**microerosion meter (MEM)** A device for measuring the rate at which an exposed rock surface is lowered, perhaps by \*weathering. It consists of a gauge which records the extension of a probe. For measurement purposes the unit is placed on three studs, already set in the rock surface, and which provide a reference level. The extension of the probe is then a measure of \*erosion.

**microevolution** Evolutionary change within species, which results from the differential survival of the constituent individuals in response to natural selection. The genetic variability on which selection operates arises from mutation and sexual reshuffling of \*gene combinations in each generation.

**microfabric** Structure or arrangement of \*mineral and organic particles on a microscopic scale, seen only with a microscope.

**microfossil** Any \*fossil that is best studied by means of a microscope. Material may include dissociated fragments of larger organisms, whole organisms of microscopic size, or embryonic forms of larger fossil

organisms. Various groups, e.g. *\*acritarchs*, *\*Foraminiferida*, *\*Ostracoda*, and *\*Conodontophora*, are studied and used as stratigraphic markers. See [MARKER BED](#); [MICROPALAEONTOLOGY](#).

**microfracture** See [VOIDS](#).

**microgabbro** See [DOLERITE](#).

**microgranite** A medium-grained (1–5 mm grain diameter) *\*igneous* rock characterized by the *\*mineral* assemblage and chemical composition of *\*granite*.

**microgranodiorite** A medium-grained (1–5 mm grain diameter) *\*igneous* rock characterized by the *\*mineral* assemblage and chemical composition of *\*granodiorite*.

**micrographic** See [GRANOPHYRIC](#).

**microkrystite** A *\*crystallite* found in a microtektite (see [TEKTITE](#)) formed by rapid quenching from a high temperature, from either vapour or molten material, in the ejecta produced by an air-fall impact event.

**microlite** An extremely small *\*crystal*, usually found embedded in a glassy *\*groundmass* (see [GLASS](#)) and resolvable only by using high power magnification on a petrological microscope (see [POLARIZING MICROSCOPE](#)). Such crystals represent the frozen initial stages of crystal *\*nucleation* and growth, and are usually preserved in rapidly chilled *\*lavas*.

**microlog** The record produced by a small electrical sonde, 3–5 cm between the electrodes, that is usually combined with a caliper sonde (to measure *\*borehole* diameter), the caliper maintaining the microlog against the borehole wall. The microlog mainly records the resistivity and thickness of the *\*mud cake*, and the log is used to assist interpretation of *\*laterologs*. See *also* [CALIPER LOG](#).

**MicroMAS-2** See [MICRO-SIZED MICROWAVE ATMOSPHERIC SATELLITE-2](#).

**micropalaeontology** The study of *\*microfossils*. Since the time of A. D. d'Orbigny (1802–57), who is credited with founding the discipline, many thousands of papers have been written on various microfossil groups and many thousands of species have been described. Commercial, or applied,

micropalaeontology began in 1877 when the age of strata in a well in Austria was determined by means of *\*Foraminiferida*, since when it has become a powerful tool in geologic investigations. Many oil companies either have their own laboratories devoted to micropalaeontology or employ consultants in this field.

**micropertthite** See PERTHITE.

**microphenocryst** See MICRO-.

**microphyll** See ENATION THEORY.

**micropiracy** A type of small-scale *\*river capture* that occurs on hillsides with parallel *\*rills*. Some rills are deeper or lower than others and during heavy storms when *\*surface runoff* overflows the ridges between rills water flows from the shallower or higher rill to the deeper or lower rill, eroding the ridge between them. What is then the larger rill then grows wider.

**microplate** Any small lithospheric *\*plate*. To be classed as a microplate, any present-day, small fragment of *\*lithosphere* should have identifiable *\*plate margins*, though adherence to this requirement is not strict (*see also TERRANE*). Microplates with *\*continental crust* are also ‘microcontinents’, though the reverse is not necessarily true: Japan is a microcontinent but a part of the *\*Eurasian Plate*, and the Rockall microcontinent has long ceased its independent movement. Microplates, especially those that have continental crust, are considered by many geologists to be important in the formation of several *\*orogenic belts*, e.g. the Cordillera of western N. America. Most authors suggest that the break-up of *\*Gondwana* produced many microplates whose subsequent collision with and *\*accretion* to Eurasia may explain the complexities of the Alpine–Himalayan belt, with the strain being taken up partly by the slipping, rotation, and slicing up of microplates and partly by deformation of other terranes.

**microporphyrific** See MICRO-.

**micropulsations** See PULSATIIONS.

**micropygous** See PYGIDIUM.

**microsatellite** A space satellite with a *\*wet mass* of 10–100 kg.

**Micro-Satellite à trainée pour l'Observation du Principe d'Equivalence (MicroSCOPE)** A collaborative \*minisatellite mission between the French space agency, the \*European Space Agency, and the University of Bremen to test the general theory of relativity. It was launched on 25 April 2016, from Kourou, French Guiana, into a Sun-synchronous quasi-circular orbit at an altitude of about 710 km.

**microsclere** See HEXACTINELLIDA.

**MicroSCOPE** See MICRO-SATELLITE A ` TRAINÉE POUR L'OBSERVATION DU PRINCIPE D'EQUIVALENCE.

**microseism (microearthquake)** A small \*earthquake, usually with a \*Richter magnitude less than 2.

**Micro-sized Microwave Atmospheric Satellite-2 (MicroMAS-2)** A 3-unit \*CubeSat mission by the Massachusetts Institute of Technology (MIT) to monitor atmospheric temperature and moisture; it follows the MicroMAS-1 mission, which failed. MicroMAS-2 was launched on 12 January 2018, from the Satish Dhawan Space Centre, India, into a Sun-synchronous orbit at an altitude of 550 km.

**microspar** Fine, crystalline \*calcite with crystals 4–10 µm in size, formed by the \*recrystallization of \*micrite. Microspar is not synonymous with \*microsparite, and the term is reserved solely for neomorphosed micrite. See NEOMORPHISM.

**microsparite** Sparry calcite in 5–20 µm range. Compare MICROSPAR.

**microsporangium** See SPORE.

**microspore** See SPORE.

**microstylolite** A complex, irregular, \*pressure-dissolution surface, lined with insoluble residues. It is developed particularly in \*carbonate rocks.

**microsyenite** A medium-grained \*igneous rock characterized by the \*mineral assemblage and chemical composition of \*syenite.

**microtektite** See TEKTITES.

**microthermal** Applied to the seasonal patterns of river discharge in areas where at least one month has a mean temperature below  $-30^{\circ}\text{C}$ .

**microthermal climate** Low-temperature climate of short summers, defined in the *\*Köppen classification* as having mean winter temperatures of less than  $-3^{\circ}\text{C}$ . Examples include the cold *\*boreal* forest climate types in continental interiors, and along some eastern seabords in latitudes  $40\text{--}65^{\circ}$ . The term is also applied in the *\*Thornthwaite* classification according to potential-evapotranspiration and moisture-budget criteria.

**microthermometry** A technique for determining the approximate composition of *\*fluid* inclusions by studying their behaviour on freezing and melting. A microscope with a *\*stage* that can heat and freeze the specimen is used to study thin, doubly-polished sections.

**microtidal** Applied to coastal areas in which the tidal range is less than 2 m. Wave action dominates the processes active in microtidal areas, e.g. the Mediterranean Sea and the Gulf of Mexico.

**microwave** *\*Electromagnetic* radiation which has a wavelength between 100  $\mu\text{m}$  and 30 cm and frequencies between 1 GHz and 300 GHz. Microwaves lie between *\*infrared* and radiowaves. *See also* PASSIVE MICROWAVE.

**microwave demagnetization** The use of *\*microwaves* to excite *\*magnons*, thereby causing loss of magnetization without significant heating and concomitant chemical change.

**microwave sounding unit (MSU)** An instrument carried on *\*Television and Infrared Observation Satellites* (TIROS-N) and NOAA satellites that measures *\*microwave* emissions from molecular oxygen in the *\*troposphere*. The readings are used to calculate the mid-tropospheric atmospheric temperature with an estimated accuracy of  $\pm 0.01^{\circ}\text{C}$ .

**Mid-Atlantic Ridge** The oceanic *\*ridge* which separates the *\*North* and *\*South American Plates* from the *\*African* and *\*Eurasian Plates*. It is a slow-spreading ridge, with rugged topography and a well-developed *\*median valley*.

**mid-continent crustal type** A type of very stable *\*crust* averaging 38 km in thickness that resembles a *\*craton*, but has thick layers of *\*sedimentary*

rocks covering extensive basins.

**mid-infrared** \*Infrared radiation which has a wavelength between 8  $\mu\text{m}$  and 14  $\mu\text{m}$ .

**mid-oceanic crustal type** A type of unstable \*crust, averaging 10 km in thickness, that is made from basaltic lava. There is much \*volcanicity and there are frequent shallow \*earthquakes. In some places there are rift valleys at the centres of \*mid-ocean ridges and ridges may protrude as islands above the surface.

**mid-ocean ridge** A long, linear, elevated, volcanic structure often lying along the middle of the ocean floor. Such ridges tend to occupy central positions because the oceans have formed by the symmetrical spreading of two lithospheric \*plates from the ridge sites. Ocean ridges occur in all the Earth's oceans, but may be offset from a central position, e.g. the E. Pacific ridge, where one side of the oceanic crust is being consumed along a \*subduction zone. At mid-ocean ridges, ocean floor is being formed. At the centre there is a rift valley, formed as discrete segments, bordered by high mountains on both sides. At a fast-spreading ridge (opening at up to 15 cm a year) the crust is smoother, with flat lavas flowing from fissures, than at a slow-spreading ridge (about 2 cm a year), where the median valley contains a chain of small volcanoes linked by fissure eruptions.

**mid-ocean-ridge basalt (MORB)** A type of tholeiitic \*basalt (see THOLEIITE), erupted from mid-ocean-\*ridge constructive-plate margins; it is one of the most abundant of all rocks and covers much of the Earth's surface. It is characterized by very low concentrations of  $\text{K}_2\text{O}$  and  $\text{TiO}_2$ ; low iron,  $\text{P}_2\text{O}_5$ , Ba, Rb, Sr, Pb, Th, U, and Zr; and high CaO. When the concentration of each \*rare-earth element in the basalt is divided by its mean concentration in \*chondrite meteorites (a standard for comparison), this type of basalt shows a progressive lowering of the ratios for the light rare-earth elements (LREEs), compared to the ratios for the heavy rare-earth elements (HREEs). MORBs are said to be LREE depleted, a reflection of the chemically depleted nature of the \*mantle source regions from which they are derived. Since leaving their source region in the mantle, these basalts, often termed low-potassium \*tholeiites, have not been contaminated by passing through any \*continental crust and therefore retain the chemical signature of the mantle from which they were derived.

MORBs thus provide an insight into the composition of the sub-oceanic mantle.

**mid-Pleistocene transition (MPT)** A shift in the period of the Pleistocene glacial cycle from 40 000 years to 100 000 years that occurred about 950 000 years ago.

**midstand systems tract** See REGRESSIVE SYSTEMS TRACT.

**Mie scattering** Scattering of electromagnetic radiation, mainly in a forward direction, by spherical particles. The theory was proposed by G. Mie in 1908. See also RAYLEIGH SCATTERING.

**migmatite** A coarse-grained, heterogeneous mixed rock consisting of: (a) a high-grade metamorphic component with a *\*gneissose* texture (see METAMORPHIC GRADE); and (b) an *\*igneous* component with a *\*granite* mineralogy and a foliated or unfoliated texture (see FOLIATION). Migmatites are found in high-grade metamorphic terrains where a sequence from high-grade metamorphic rocks through migmatites to granite bodies is often seen in the field. The granite component is thought to form by *\*partial melting* of the rock during extreme *\*metamorphism*. Migmatites may thus be a record of the initial stages in the generation of large bodies of granite magma and, as such, they would represent the high-temperature boundary between metamorphic and igneous rocks. Migmatites have an attractive appearance, often being marked with irregular small stripes or patches of contrasting shades ranging from almost white to dark grey, and are widely used as building stone, sometimes being polished for ornament.

**migmatization** The process whereby a rock undergoes *\*partial* melting during extreme *\*metamorphism*, producing a *\*migmatite*.

**migration** A method of reconstructing a *\*seismic section* so that dipping reflection events are repositioned to lie beneath their true surface locations and at corrected vertical two-way travel times. There are several migration methods. These include, for example, wave-equation, dip-moveout, ray-trace, finite-difference, wavefront-common-envelope, diffraction, and *\*frequency-domain*.




**migration route** Link between two biogeographical regions that permits the interchange of plants and/or animals. Various types are recognized in

the literature: for example, G. G. \*Simpson's (1940) '\*corridors', '\*filters', and '\*sweepstakes dispersal routes' are widely referred to in connection with mammalian, and more recently reptilian, migrations.

**Mikulino** A local name, in Russia, for sediments of \*Eemian age.

**Milankovitch, Milutin** (1879–1958) A Serbian mathematician and physicist from the University of Belgrade, Milankovitch made important studies of solar radiation. He concluded that there were periodic changes in the amount of radiation received on Earth, caused by eccentricities in the Earth's orbit. These led to cyclical climatic changes, causing glacial periods, etc. *See also* MILANKOVITCH CYCLES.

**Milankovitch cycles** Cyclical changes in the rotation and orbit of the Earth that \*Milankovitch correlated to climatic effects. There are three cycles: changes in the eccentricity of the Earth's orbit, altering the distance between Earth and the Sun at \*aphelion and \*perihelion, with a period of about 100 000 years; variations in the tilt of the Earth's rotational axis (obliquity of the ecliptic), with a period of about 40 000 years; and a movement (wobble) in the angle by which the axis of the Earth's rotation is tilted in respect of the orbital plane, altering the seasons at which aphelion and perihelion occur (precession of the equinoxes), with a period of about 21 000 years. Climatic changes associated with Milankovitch cycles may be recorded in \*cyclic sedimentation.



**Milankovitch cycles**

**Milazzian** See QUATERNARY.

**milioline winding** \*Coiling which occurs in the suborder Miliolina of the \*Foraminiferida. Coiling starts with a \*planispirally coiled \*proloculus and may continue in this way. Usually, however, tubular chambers are added lengthwise around the growth axis. It is common to find chambers added at 140° or 120°, leaving either three or five chambers visible on the outside of a \*test.

**Milleporina** (millepores; massive hydrocorals) (class \*Hydrozoa) Order of \*reef-forming \*Cnidaria whose members build massive, calcareous \*exoskeletons which have pores through which the polyps protrude. Milleporina are known from the Upper \*Cretaceous to \*Recent. See also STYLASTERINA.

**Miller, William Hallowes** (1801–80) A British mineralogist from the University of Cambridge, Miller developed a classification system for crystallography, based on crystal axes. His method was published in *Treatise on Crystallography* (1839). See MILLER INDICES.

**Miller–Bravais indices** In the \*hexagonal and \*trigonal \*crystal system, there are four \*crystallographic axes. M. A. Bravais adapted the \*Miller notation to express the intercepts of \*crystal faces on these four axes using the modified symbols  $h, k, i, l$ ; these symbols are the reciprocals of the ratios of the intercepts on the  $a_1, a_2, a_3,$  and  $c$  (or  $x, u, y,$  and  $z$ ) axes respectively.

**Miller indexes** See MILLER INDICES.

**Miller indices (Miller indexes)** One of the methods of notation devised to express the intercepts of \*crystal faces on \*crystallographic axes, proposed by W. H. \*Miller. The symbols  $h, k,$  and  $l$  represent whole numbers which are the reciprocals of intercepts along the  $a, b,$  and  $c$  (or  $x, y,$  and  $z$ ) crystallographic axes respectively. A face which is parallel to a crystallographic axis is indicated by the symbol  $o$ . If a crystal face intercepts the negative end of an axis, a bar negative sign) is placed above the appropriate symbol. A \*prism will have the general notation of  $100$  or  $110$ , a \*pinacoid  $001$ , and a \*pyramid  $101$  or  $111$ . When the indices describe a crystal face they are conventionally written without brackets; when they describe crystal \*form, they are enclosed in brackets.

**millerite** Mineral, NiS; sp. gr. 5.2–5.6; \*hardness 3.0–3.5; \*trigonal; brass-yellow and opaque; greenish-black \*streak; \*metallic \*lustre; crystals usually long, slender, \*acicular, and in radiating groups; \*cleavage perfect, rhombohedral; occurs as tufts of radiating fibres in cavities, and as replacement for other nickel minerals, also in veins carrying nickel minerals and other \*sulphides, and around some volcanoes as sublimation products.

It is a minor ore of nickel and named after the British mineralogist [W. H. \\*Miller](#).

**milli- (m-)** From the Latin *mille* meaning 'one thousand', a prefix meaning 'one-thousandth' (e.g. a 'milli-equivalent' is one-thousandth of an equivalent weight). Attached to SI units it denotes the unit  $\times 10^{-3}$ .

**millibar** See [BAR](#).

**milligal** One-thousandth of a gal; equivalent to 10 [\\*gravity units](#).

**millipedes** See [MYRIAPODA](#).

**Millstone Grit series** See [NAMURIAN](#).

**Milne, John** (1850–1913) A British mining engineer who became professor of geology and mining in Tokyo. Milne became interested in [\\*earthquakes](#), developing and testing [\\*seismographs](#), and recording waves produced by explosions. He was the first to show that distant earthquakes could be recorded. He also made a study of earthquake-proof buildings.

**Mimas (Saturn I)** One of the major satellites of [\\*Saturn](#), with a radius of 198.8 km; mass  $0.375 \times 10^{20}$  kg; mean density 1140 kg/m<sup>3</sup>; visual albedo 0.5. It was discovered in 1789 by Sir William Herschel.

**mimetic twins** Crystals which, on being twinned, appear to have a higher order of [\\*crystal](#) symmetry than is the case. Examples are commonly found in [\\*aragonite](#) and the [\\*zeolite](#) group of minerals.

**mimetite** See [VANADINITE](#).

**Mimistrobell** A [\\*solar system](#) asteroid (No. 3840), with an orbital period of 3.38 years.

**MIMOSA** See [MICROACCELEROMETRIC MEASUREMENTS OF SATELLITE ACCELERATIONS](#).

**Mindel** The second of four glacial episodes, taking its name from an Alpine river, and established by [A. \\*Penck](#) and E. Bruckner in 1909. It may be equivalent to the [\\*Elsterian](#) of northern Europe.

**Mindel/Riss Interglacial (Great Interglacial)** An Alpine [\\*interglacial](#) [\\*stage](#) that is possibly the equivalent of the [\\*Hoxnian](#) of East Anglia,

England, or the **\*Holsteinian** of northern Europe.

**Mindyallan** A **\*stage** (501–497 Ma ago) of the Late **\*Cambrian** of Australia, preceded by the **\*Boomerangian** and overlain by the **\*Idamean**.

**mineral** Usually inorganic substance which occurs naturally, and typically has a crystalline structure whose characteristics of **\*hardness**, **\*lustre**, colour, **\*cleavage**, fracture, and relative density can be used to identify it. Each mineral has a characteristic chemical composition. **\*Rocks** are composed of minerals. More loosely, certain organic substances obtained by mining are sometimes termed ‘minerals’.

**mineralization** Conversion of organic tissues to an inorganic state as a result of decomposition by soil micro-organisms.

**mineral layering** The concentration of one **\*mineral** type or a combination of mineral types into layers varying in thickness from a few centimetres to 2–3 m and commonly found in large **\*intrusive \*igneous** rock bodies. The sudden appearance or disappearance of a particular mineral marks the boundary between the mineral layers. Mineral layering is also known by the term ‘phase layering’. *Compare* CUMULATE.

**mineralogy** The scientific study of **\*minerals**, comprising **\*crystallography**, mineral chemistry, economic mineralogy, and determinative mineralogy (concerned mainly with physical properties).

**mineraloid** A substance resembling a mineral but lacking a crystal structure, e.g. **\*jet**, **\*obsidian**, **\*opal**.

**mineral saturation index** An index showing whether a water will tend to dissolve or precipitate a particular mineral. Its value is negative when the mineral may be dissolved, positive when it may be precipitated, and zero when the water and mineral are at chemical equilibrium. The saturation index (SI) is calculated by comparing the chemical activities of the dissolved **\*ions** of the mineral (ion activity product, IAP) with their solubility product ( $K_{sp}$ ). In equation form,  $SI = \log(IAP/K_{sp})$ .

**mineral soil** Soil composed principally of mineral matter, in which the characteristics of the soil are determined more by the mineral than by the organic content.

**minette** A type of *\*lamprophyre*, characterized by essential *\*biotite* and *\*orthoclase* feldspar (see ESSENTIAL MINERAL). If *\*augite* is present the rock is termed an ‘augite-minette’.

**minette ironstone** Iron *\*ore* from Alsace-Lorraine; *\*limonite* (iron oxide) is the main iron mineral with some *\*siderite* (iron carbonate).

**minimum-distance-to-means classification** A *\*remote* sensing *\*classification* system in which the mean point in digital parameter space is calculated for *\*pixels* of known classes, and unknown pixels are then assigned to the class which is arithmetically closest when digital number values of the different *\*bands* are plotted. See also BOX CLASSIFICATION; MAXIMUM-LIKELIHOOD CLASSIFICATION.

**minimum melting curve** A univariant *\*solidus* curve in pressure–temperature space that defines the initial melting temperature of a solid as a function of pressure under water-saturated conditions. Minimum melting temperatures are always achieved under water-saturated conditions.

**minimum temperature** The lowest temperature recorded diurnally, monthly, seasonally, or annually, or the lowest temperature of the entire record. Daily air-temperature minima are recorded by the screen minimum thermometer. See also GRASS MINIMUM TEMPERATURE.

**minimum thermometer** Thermometer that records the lowest temperature to which it has been exposed, e.g. by allowing a fall of liquid past a restriction in the tube, but preventing the liquid’s return on expansion. It is most commonly used to record minimum daily temperatures.

**minisatellite (small satellite)** A space satellite with a *\*wet mass* of 100–500 kg, or any satellite with a wet mass of less than 500 kg.

**minor fold** A *\*fold* which is generally in *\*harmonic* accordance with the geometry of a major fold system and which has a characteristic *\*M-fold*, *\*S-fold*, or *\*Z-fold* profile in specific areas of an *\*asymmetrical fold*.

**mio-** From the Greek *meion* meaning ‘less’, a prefix meaning ‘less’ (e.g. *\*Miocene*, ‘mio’ plus ‘cene’ (from the Greek *kainos*, ‘new’), meaning ‘less new’).

**Miocene** First of the two \*epochs of the \*Neogene period, extending from the end of the \*Oligocene, 23.03 Ma ago, to the beginning of the \*Pliocene, 5.332 Ma ago. Many mammals with a more modern appearance evolved during this epoch, including deer, pigs, and several elephant stocks. The Miocene comprises the \*Aquitainian, \*Burdigalian, Early and Late \*Langhian, \*Serravallian, \*Tortonian, and \*Messinian \*ages.



<https://ucmp.berkeley.edu/tertiary/miocene.php>

- The Miocene Epoch.

**miogeocline** An association chiefly of \*carbonates, \*shales, and clean \*sandstones, with an absence of volcanics. These sediments are thought to have formed in shallow water on a continental margin.

**miogeosyncline** An obsolete term for that part of a \*geosyncline characterized both by sediments deposited in shallow water and by the absence of volcanics. In the classification of geosynclines, miogeosynclines had a relatively thin sedimentary pile and were developed closer to the \*craton than the volcanic \*eugeosyncline. The geosynclinal theory of the formation of \*orogenic belts has been superseded by various \*plate tectonic models, and the term 'miogeosyncline' has been replaced by '\*miogeocline' which is a purely descriptive word having no connection with geosynclinal theory.

**mirage** Optical effect in which a major variation in temperature between layers of the lower atmosphere produces differential refraction of light, resulting for example in raised images and in gaps, which may give the appearance of a water surface.

**Miranda (Uranus V)** One of the major satellites of \*Uranus, discovered by Gerard \*Kuiper in 1948. Its radius is  $240 \times 234.2 \times 232.9$  km; mass  $0.659 \times 10^{20}$  kg; mean density  $1200 \text{ kg/m}^3$ ; albedo 0.27; gravity 0.01 (Earth = 1); distance from Uranus 129 850 km. The surface temperature is about 43 K. Miranda is composed of rocky material and water ice in approximately equal proportions. The first pictures of the surface were provided in 1986 by \*Voyager 2 as it passed close to the satellite to gain a gravity assist to take it to \*Neptune. The surface of Miranda comprises areas of rolling, cratered terrain, with grooves, valleys, and cliffs, one more than 15 km

high. Within this terrain there are three coronae. These are square, with rounded corners, up to about 260 km across, and contain dark and bright patches and sets of parallel ridges and grooves. It is now believed that the crust of Miranda was pulled apart by internal forces as the interior was evolving and the coronae formed above major upwellings of partially melted ice, and tidal distortions by Uranus caused heating.

**mirror plane** See CRYSTAL SYMMETRY.

**miscibility** A property of two or more liquids that can be mixed to form a homogeneous *\*solution*. Liquids that fail to form a solution when mixed in certain proportions are said to be immiscible. Oil and water are immiscible in all proportions, ethanol and water are miscible in all proportions, but diethyl ether and water form a solution in some proportions, but not in others, so they are immiscible. Immiscible metals will not form alloys; they will mix when molten but separate as they cool. Similarly, some *\*magmas* contain immiscible rocks that separate as the magma cools.

**miscible** See MISCIBILITY.

**mise-à-la-masse method (charged-body potential method)** A form of *\*constant-separation* traversing in which one current electrode is placed directly into a conductive *\*orebody*. The other three electrodes of the *\*array* are on the surface, as in normal *\*electrode* configurations. This method enables the subsurface extent of the conductor to be established more readily than with other electrical methods.

**misfit stream (underfit stream)** Stream that is too small to have cut the valley it currently occupies. Applies particularly to a meandering stream whose dimensions are much smaller than those of the meandering valley through which it flows.

**mispickel** See ARSENOPYRITE.

**Mississippian** 1. The Early *\*Carboniferous \*epoch* which is preceded by the Late *\*Devonian* and followed by the *\*Pennsylvanian*, and comprises the *\*Tournaisian*, *\*Visean*, and *\*Serpukhovian \*epochs*. It is dated at 359.2–318.1 Ma ago (Int. Commission on Stratigraphy, 2004). 2. The name of the corresponding N. American sub-system which comprises the *\*Kinderhookian*, *\*Osagean*, *\*Meramecian*, and *\*Chesterian \*series*. It is

roughly contemporaneous with the *\*Dinantian* plus the *\*Namurian A* of western Europe.

**Missourian A** *\*stage* (306.5–305 Ma ago) in the *\*Pennsylvanian* of N. America, underlain by the *\*Desmoinesian*, overlain by the *\*Virgilian*, and roughly contemporaneous with the Krevyakinskian and lower Chamovnicheskian stages of the *\*Kasimovian* series.

**mist** Surface-layer atmospheric condition in which visibility is reduced by very fine, suspended water droplets 0.0005–0.05 mm in diameter. In synoptic meteorology, the *\*relative humidity* in a mist condition is more than 95% and overall visibility is at least 1 km. *See also* HAZE; FOG.

**mistral** Strong, cold, northerly wind that blows offshore with great frequency along the Mediterranean coast from northern Spain to northern Italy, and that is particularly frequent in the lower Rhône valley. The wind may persist for several days, and is best developed when a *\*depression* is forming in the Gulf of Genoa to the east of a ridge of high pressure. The airstream that feeds the mistral is commonly derived from polar air of maritime origin. In the Rhône valley and similar areas of occurrence, the airflow is strengthened by *\*katabatic* and funnelling effects producing speeds of up to 75 knots (139 km/h), compared with the typical 40 knots (74 km/h) experienced along the coast.

**Mitchellian** A *\*stage* (10.5–5 Ma ago) in the *\*Neogene* of south-eastern Australia, underlain by the *\*Bairnsdalian*, overlain by the *\*Cheltenhamian*, and roughly contemporaneous with the *\*Tortonian*, *\*Messinian*, and lower *\*Zanclean* (Tabianian) stages.

**mitchellite** *See* SPINEL.

**mites** *See* CHELICERATA.

**mitochondrial-DNA (mt-DNA)** Circular DNA that is found in mitochondria. It is entirely independent of nuclear DNA and, with very few exceptions, is transmitted from females to their offspring with no contribution from the male parent.

**mitochondrial Eve** The postulated female ancestor of all modern humans, who was an archaic human living in Africa about 140 000 to 280 000 years ago, based on studies of *\*mitochondrial-DNA* from modern populations by

a team led by Allan Wilson. The data suggest there may have been a bottleneck associated with the speciation event leading to *Homo sapiens*.  
*See also* [ADAM](#).

**mixed cloud** Cloud containing unfrozen water droplets as well as ice crystals. Typically the condition is found in [\\*cumulonimbus](#), [\\*nimbostratus](#), and [\\*altostratus](#).

**mixed pixel** In [\\*remote](#) sensing terminology, a [\\*pixel](#) that has a [\\*digital](#) number which represents the average energy emitted or reflected from several different surfaces occurring within that area represented by the pixel.

**mixing condensation level** The lowest level at which condensation can occur in air that is thoroughly mixed, e.g. in the amalgamation of [\\*air masses](#) with different temperatures.

**mixing depth** The vertical extent of an atmospheric layer measured from the surface of the Earth, usually a sub-inversion layer, in which convection and turbulence lead to mixing of the air and any pollutants in it.

**mixing ratio** Ratio of the mass of a given gas (e.g. water vapour) to that of the remaining gas (e.g. dry air) in the mixture. The examples given yield the 'humidity mixing ratio', expressed most conveniently in grams per kilogram of dry air. *See also* [SPECIFIC HUMIDITY](#).

**mixolimnion** The upper layer of a [\\*meromictic](#) lake, lying above the [\\*chemocline](#), where the water is mixed by the wind, circulates freely, and has a low density. *Compare* [MONIMOLIMNION](#).

**MMS** *See* [MOMENT MAGNITUDE SCALE](#).

**moberg** The Icelandic name for a flat-topped mountain produced by the subglacial eruption of a [\\*central vent volcano](#).

**mobile belt** A synonym for [\\*orogenic](#) belt, most often used for those earlier (i.e. early [\\*Precambrian](#)) belts where [\\*plate-tectonic](#) models cannot easily be applied.

**mocha stone** *See* [AGATE](#).

**modal analysis** The determination of the *\*mode* of a *\*rock*, usually carried out by point counting using an *\*automatic point counter* attached to a *\*polarizing microscope*.

**mode** 1. The percentage by volume of each of the *\*minerals* which make up an *\*igneous* rock. Occasionally the term is also applied to *\*metamorphic rocks*. 2. In statistics, the *\*average* as defined by the most frequently occurring value in a data set.

**model** Representation of reality in which the main features of some aspect of the real world are presented in simplified terms in order to make that aspect easier to comprehend, and often to facilitate the making of predictions.

**Modified units** See MARTIAN TERRAIN UNITS.

**Moershoofd (Poperinge)** A Middle *\*Devensian* *\*interstadial* from the Netherlands during which the climate was relatively cool, with average July temperatures of 6–7 °C. This division, together with the *\*Hengelo* and *\*Denekamp* interstadials, is perhaps the equivalent of the *\*Upton Warren* Interstadial of the British Isles.

**mofette (moffette)** A *\*fumarole* that releases abundant carbon dioxide, often accompanied by methane, nitrogen, or oxygen, at a temperature below 100 °C, during the final stages of volcanic activity.

**moffette** See MOFETTE.

**Mogi doughnut hypothesis** A method for *\*earthquake prediction* proposed in 1969 by the Japanese seismologist Kiyoo Mogi (1929– ). It states that major earthquakes tend to occur in *\*seismic gaps* surrounded by regions of high seismic activity.

**mogote** A steep-sided hill up to 100 m high, or sometimes higher, that resembles a tower. Groups of mogotes, separated by wide, *\*alluvial* valley floors, occur in *\*limestone* landscapes.

**Mohawkian** A N. American *\*Ordovician* *\*stage*, 462–451 Ma ago, preceded by the *\*Whiterockian* and followed by the *\*Cincinnatian*.

**Mohnian** A *\*stage* (13.5–7.5 Ma ago) in the *\*Neogene* of California, underlain by the *\*Luisian*, overlain by the *\*Delmontian*, and roughly

contemporaneous with the upper *\*Serravallian*, *\*Tortonian*, and lower *\*Messinian* stages.

**Moho** See MOHOROVIČIĆ DISCONTINUITY.

**Mohorovičić, Andrija** (1857–1936) A Croatian seismologist, Mohorovičić discovered a seismic discontinuity since named after him (see MOHOROVIČIĆ DISCONTINUITY) at the *\*crust–\*mantle* boundary, when analysing the results of an earthquake in 1909. He found two sets of *\*P-wave* arrivals with different travel times, and thus was able to calculate the depth of the discontinuity.

**Mohorovičić discontinuity (Moho)** The seismic discontinuity that marks the boundary between the *\*Earth's \*crust* and *\*mantle*. The mantle has *\*P-wave* velocities of about 8.1 km/s, higher than that of the lower-density crustal rocks. The Moho was originally recognized at a depth of some 20 km in Europe by A. *\*Mohorovičić* on the basis of the arrival of *\*refracted* P-waves before *\*direct* P-waves. Previously considered to be very sharp, the discontinuity is now known to be more complex.

**Mohr stress diagram** A two-dimensional, graphical representation of the relationship between *\*shear* stress, *\*confining pressure*, and the angle of *\*failure* at the point at which a failure in a material occurs. Circles whose centres lie along the axis of *\*normal stress* (*\*principal stress axes* ( $\sigma_1 + \sigma_3/2$ )) and whose diameters represent the differential stress (principal stress axes  $\sigma_1 - \sigma_3$ ) represent the *\*stress* state, the data for them having been obtained experimentally. Tangents to each circle join the points of failure for different stress states and so delimit the field of stable stress states (within a Mohr stress envelope) from the failure field outside.

**Mohs, Friedrich** (1773–1839) A German mineralogist, Mohs was a student of *\*Werner* and succeeded him at Freiberg University before moving to become professor of mineralogy in Vienna. In 1812, he developed a decimal scale for the hardness of minerals which is still in use. His *Grundriss der Mineralogie* (English edition called *Outline of Mineralogy*) was published in 1825. See MOHS'S SCALE OF HARDNESS.

**Mohs's scale of hardness** Scale, devised in 1812 by the German mineralogist Friedrich *\*Mohs*, of the scratch *\*hardness* of *\*minerals*, as: 1

\*talc; 2 \*gypsum; 3 \*calcite; 4 \*fluorite; 5 \*apatite; 6 \*orthoclase; 7 \*quartz; 8 \*topaz; 9 \*corundum; and 10 \*diamond. The scale is linear up to a hardness of 9 (corundum), but then rises dramatically to 10, with diamond about 10 times harder than corundum. The scale is still used today.

**Moinian** A \*stage of the Upper \*Proterozoic of the north-western Highlands of Scotland, underlain by the \*Longmyndian and overlain by the \*Dalradian (\*Cambrian).

**moisture balance** See MOISTURE BUDGET.

**moisture budget (moisture balance, water budget)** The balance of water fluxes into and out of a defined area over a defined time period, as represented broadly by the equation: precipitation = runoff + evapotranspiration + the change in soil-moisture storage. In mid-latitudes, for example, the annual budget is balanced by a high level of potential \*evapotranspiration and utilization of \*soil moisture in summer, compensated by a water surplus and recharge of soil moisture in winter when evaporation is less and precipitation is sometimes greater.

**moisture index** A term based on the computation of an annual moisture budget by C. W. Thornthwaite (1955), and calculated from the \*aridity and \*humidity indices, as  $I_m = 100 \times (S - D)/PE$ , where  $I_m$  is the moisture index,  $S$  is the water surplus in months when precipitation exceeds \*evapotranspiration,  $D$  is the water deficit in months when evapotranspiration exceeds precipitation, and  $PE$  is the potential evaporation.

**Mokoian** A New Zealand \*stage (145.5–119 Ma ago) of the Early \*Cretaceous \*epoch, preceded by the \*Puaruan and followed by the \*Korangian.

**Mokolian** A South African \*stage (2050–900 Ma ago), preceded by the \*Vaalian and followed by the \*Namibian.

**mol** See MOLE.

**molarity** See CONCENTRATION.

**molasse** A term originally used to describe the mainly shallow-marine and non-marine \*sediments produced from the \*erosion of a mountain belt after

the final stage of uplift in an *\*orogeny*. It is now clear that much so-called molasse is not *\*post-tectonic*, but *\*syntectonic*, developed from the erosion of *\*nappes* while uplift and deformation are still progressing; some workers therefore consider that the term should be discontinued. See *FLYSCH*.

**Molasse Basin** See *NORTH ALPINE FORELAND BASIN*.

**moldic porosity (mouldic porosity)** A form of *\*secondary* porosity, developed by the preferential dissolution of shell fragments or other particles, to leave empty spaces previously occupied by the particles. See *POROSITY*; *CHOQUETTE AND PRAY CLASSIFICATION*.

**mole (mol)** The amount of a substance containing as many elementary units as there are carbon atoms in 12 g of carbon-12 (i.e. the *\*Avogadro* number). The elementary units by which the amount of substance is being measured may be atoms, molecules, *\*ions*, *\*electrons*, radicals, or any other expressly named constituent.

**molecular sieve action** Technique used in synthetic *\*zeolites* to separate molecules by trapping them in the *\*crystal* lattice. Sizes are selected to suit particular molecules.

**mole drain** Drain that can be made in soils by pulling a bullet-shaped device through the soil so that the compacted sides of the tunnel maintain that form for several years.

**mollic horizon** Surface *\*soil horizon* of mineral soil that is dark in colour, and relatively deep, and contains (dry weight) at least 1% organic matter, or 0.6% organic carbon, the determination of either being acceptable. It is the *\*diagnostic* horizon of *\*mollisols* and is associated with base-rich materials and grassland vegetation.

**mollisols** An order of mineral soils in the US *\*soil taxonomy* that are identified by a deep *\*mollic* surface *\*soil horizon* (well-decomposed and finely distributed organic matter) and base-rich *\*mineral soil* below.

**Mollusca (molluscs)** A very diverse phylum of *\*invertebrates* which have a common body plan modified in various ways. There is usually a shell, secreted by a series of tissues (called the *\*mantle*), and beneath this there is the body, which contains a space (mantle cavity) in which lie the gills. The body plan is modified to give the major classes of the phylum. One class,

the **\*Monoplacophora**, shows signs of internal segmentation. The **\*Amphineura** (chitons) have a **\*bilaterally symmetrical** shell of eight plates. Scaphopoda have tapering, curving shells which are open at both ends. **\*Bivalvia** have the soft parts enclosed between two shells. **\*Gastropoda** usually possess a coiled univalve shell. The most advanced of the molluscs are the **\*Cephalopoda**, which possess either internal or external, chambered shells. Other groups are also assigned to the Mollusca, but only cephalopods, bivalves, and gastropods have a good geologic record.

**molluscs** See MOLLUSCA.

**molybdenite** Mineral,  $\text{MoS}_2$ ; sp. gr. 4.6–4.8; **\*hardness** 1.0–1.5; **\*hexagonal**; pale bluish to lead-grey; greenish-grey **\*streak**; **\*metallic lustre**; crystals hexagonal, often **\*tabular** grains, and it also occurs as foliated and scaly masses; **\*cleavage** perfect basal {0001}; quite widespread but seldom occurs in large quantities, found as an **\*accessory mineral** in **\*granites**, **\*quartz** veins, and **\*pegmatites**, in contact **\*metamorphic zones** with **\*garnets**, **\*pyroxenes**, **\*scheelite**, **\*pyrite**, and **\*tourmaline**, and in **\*veins** with scheelite, **\*wolframite**, **\*cassiterite**, and **\*fluorite**. It is a major **\*ore mineral** for molybdenum.

**moment magnitude scale (MMS)** A scale used to measure and report the magnitude of **\*earthquakes** in terms of the amount of energy they release. The designation ( $M_W$  for mechanical work) refers to the **\*seismic moment** of the earthquake. MMS has largely replaced the **\*Richter scale**.

**moment of inertia (*I*)** The kinematic properties of a rotating body; a measure of the rotational inertia of an object around a specific axis of rotation, in units of  $\text{kg/m}^2$ . In the **\*Earth**, the principal moment of inertia lies close to the **\*axis of rotation** and passes through the centre of mass of the Earth. Changes in the mass distribution, e.g. ice sheets, seasonal atmospheric changes, etc., cause changes in the location of the moments of inertia.

**monadnock** Isolated hill or range of hills standing above the general level of a **\*peneplain** and resulting from the **\*erosion** of the surrounding terrain. It may be located on relatively resistant rock or in a **\*watershed** position where erosion is least. It is named after Mount Monadnock, New Hampshire, USA.

**monazite** Mineral (Ce,La,Y,Th)PO<sub>4</sub>; sp. gr. 4.9–5.4; \*hardness 5.0–5.5; \*monoclinic; clove-brown to reddish-brown to orange and green; off-white \*streak; resinous to waxy \*lustre; crystals small, short, \*prismatic to \*tabular grains, with larger crystals showing striated faces; \*cleavage imperfect basal {001}; found extensively as an \*accessory mineral in \*granites and \*pegmatites, in \*gneisses and \*carbonatites, and concentrated in \*alluvial sands and \*placers. It is used as a source of cerium, thorium, and other rare-earth metals and compounds.

**monchiquite** A type of \*lamprophyre, characterized by \*essential \*analcite, \*barkevikite (an alkali \*amphibole), and/or \*augite. There is no \*feldspar present in this type of lamprophyre.

**Monera** One of the five kingdoms of life in the classification originally proposed by R. H. Whitaker (1959, 1969), comprising the \*prokaryotic Cyanophyta (blue-green algae) and \*bacteria (Schizomycophyta). The bacteria were the first forms of life on Earth, dating from at least 3300 million years ago, while the first \*cyanophytes appeared about 2600 million years ago.

**Monian** A \*stage of the Middle-Upper \*Proterozoic of Anglesey and the Lleyn Peninsula, north Wales, equivalent to the Lower \*Torridonian.

**monimolimnion** The lower layer of a \*meromictic lake, lying below the \*chemocline, where the water is dense, static, and does not mix with the water above. Compare MIXOLIMNION.

**Monitor-E** A low-orbiting experimental spacecraft that was launched by the Russian Federal Space Agency (RSA) on 26 August 2005 into a \*Sun-synchronous orbit at an altitude of about 550 km, covering swathes 90–160 km wide with a resolution of 8–20 m. Its purpose was to acquire remote sensing data for environmental monitoring, emergency monitoring, and research into natural resources. The spacecraft was designed to live for five years but communication with it was lost and Monitor-E was removed from service in August 2008.



[http://eng.ntsomz.ru/ks\\_dzz/satellites/monitor\\_E](http://eng.ntsomz.ru/ks_dzz/satellites/monitor_E)


- An RSA mission to study and map the geology and environment of the Earth.

**mono-** From the Greek *monos* meaning 'alone', a prefix meaning 'single' or 'one'.

**monochromator** In *\*reflected-light microscopy*, an accessory used to observe the optical properties of minerals with incident light of a specific wavelength. Fixed monochromatic interference filters with a band width of 15–50 nm and continuous-spectrum monochromators are available. They are used primarily for *\*reflectance* measurements at specific wavelengths, normally 546 to 589 nm.

**monocline** A one-limbed flexure on either side of which the *\*strata* are horizontal or *\*dip* uniformly at low angles.

**monoclinic** Applied to a *\*crystal* system where the *\*Bravais* lattices have three sets of edges of unequal length, two sets of which are not at right angles to one another. They can therefore be referred to three unequal *\*crystallographic axes* *a*, *b*, and *c* (or *x*, *y*, and *z*). The *a* (or *x*) axis (clino) is inclined and forms an obtuse angle with the vertical *c* (or *z*) axis usually denoted by the symbol  $\beta$ . The *b* (or *y*) axis (ortho) is horizontal and at right angles to both the *a* (*x*) and *c* (*z*) axes.



**Monoclinic**

**Monocraterion** A single, tube-like, vertical **\*burrow** that has a series of stacked funnel structures that point towards the top surface of the bed. The multiples of funnel structures are a response to increased sedimentation and reflect the upward migration of the animal.

**Monocyathea** See REGULARES.

**monodactyl** Applied to the condition in which the lateral **\*metacarpals** and **\*metatarsals** are reduced, leaving only one functional digit.

**monograptid** **1.** A member of a suborder of **\*Graptoloidea**, occurring in Lower **\*Silurian** to **\*Emsian** marine rocks, characterized by the possession of a single, **\*uniseriate** **\*stipe**. **2.** Applied to the latest of four graptolite faunas occurring in Silurian and **\*Devonian** rocks.

**Monograptus** (class **\*Graptolithina**) The first recorded **\*monograptid** genus, from the Early **\*Llandoveryan** (Lower **\*Silurian**). Successive evolutionary trends within the graptolites resulted in the development of the single-branched monograptids. These had only a limited number of thecal cups (see **THECA**) on one side of the branch although the cups were large and exhibited considerable morphological variation. Compare **DENDROIDEA**; **GRAPTOLOIDEA**.

**monoete** See **SPORE**.

**monomictic** Applied to lakes in which only one seasonal period of free circulation occurs. In cold monomictic lakes, typical of polar latitudes, the seasonal overturn occurs briefly in summer; in other seasons, surface-water temperatures are below 4 °C, which induces density stratification. In warm monomictic lakes, typical of warm temperate or subtropical regions, the seasonal overturn occurs in winter; at other times thermal stratification, with the formation of a distinct **\*epilimnion**, prevents free circulation through the depth of the lake.

**monomineralic** Applied to **\*rocks** composed of one **\*mineral** type only. Examples would include the **\*igneous** rock **\*anorthosite** (composed entirely of **\*plagioclase** feldspar) and the **\*metamorphic rock** **\*marble** (composed entirely of **\*calcite**).

**monomyarian** See **MUSCLE SCAR**.

**monophyletic** Applied to members of a **\*taxon** that are descended from a common ancestor, e.g. the families within a class would be monophyletic if they were all descended from the same family or lower taxonomic unit. Under the strictest definition they would all have to be descended from a single species. *Compare* POLYPHYLETIC.

**Monoplacophora** (phylum **\*Mollusca**) Class of primitive, almost **\*bilaterally symmetrical**, univalved molluscs, whose limpet-shaped shell of calcium carbonate consists of an external periostracum (the outer, non-calcareous layer), a prismatic layer, and an internal, nacreous deposit. The internal organs are **\*metamerically segmented**, and the circular foot and radula are ventrally placed. Typically, Monoplacophora have paired **\*muscle** scars. All are marine, benthic filter-feeders. Fossil forms inhabited shallow water, but modern forms occur in deep water. They first appeared in the Lower **\*Cambrian**.

**monopodial** **1.** Type of branching in which lateral branches arise from a definite main, central stem. **2.** With a single axis, an extension growth from the apex.

**Monorhina** Older name for the group of fish, including the lampreys and related jawless fossils, which possess a single median nostril between the eyes.

**monostatic radar** *See* BISTATIC RADAR.

**monosulcate** *See* POLLEN.

**Monotremata** (class **\*Mammalia**, subclass Prototheria) An order comprising the duck-billed platypus (*Ornithorhynchus anatinus*) and the echidnas or spiny ant-eaters, *Tachyglossus* and *Zaglossus*. There were some extinct forms of which very few are known in detail, though of those some attained large sizes. The echidnas have no fossil record older than the **\*Pleistocene**, but a fossil platypus, *Obdurodon*, is known from the **\*Miocene** and in the early 1990s teeth of an undoubted platypus-like form, **\*Monotrematum sudamericanum**, were discovered in **\*Palaeocene** deposits in Patagonia. Two **\*Cretaceous** genera, **\*Steropodon** and *Kollikodon*, are also known. In view of their reptilian affinities they are thought to represent a separate and direct line of descent from the earliest **\*Mesozoic** animals, independent of the line leading to other mammals. They retain many

primitive features and are quite unlike *\*marsupials* or *\*eutherians* (placental mammals).

***Monotrematum sudamericanum*** A *\*Palaeocene* platypus from Patagonia, Argentina, known so far only by its molar teeth. Its presence demonstrates that monotremes (*\*Monotremata*), like marsupials, were part of the *\*Gondwana* fauna.

**monotremes** See MONOTREMATA.

**monotypic** In *\*classification*, applied to a group that has only a single representative. Thus a phylum may have only a single class or order and might be represented by a single genus and species.

**mons** (*pl.* montis, montes) The Latin word for ‘mountain’ and a term applied to large planetary mountains. The type example is Olympus Mons, the 26 km high volcano on Mars. Other examples include the Maxwell Montes on Venus.

**monsoon** From the Arabic *mausim* meaning ‘season’, a seasonal change of wind direction and properties associated with widespread temperature changes over land and water in the subtropics. Seasonal alternations of pressure systems together with shifting upper wind patterns and *\*jet streams* produce seasonal winds, called ‘monsoon winds’. The climate of the Indian subcontinent is especially characterized by the monsoon, where a distinct rainy season occurs in the south-westerly summer monsoon and the north-easterly winter monsoon is dry. Other major areas of monsoons are eastern and south-eastern Asia, the west African coast (latitude 5–15°N) and northern Australia.

**monster curve** In *\*fractal* geometry, an object for which the edge length approaches infinity as the ruler measuring it becomes infinitely short. The edge length of a fractal object depends on the size of the ruler that is used to measure it. This length increases as progressively shorter rulers are used, because the smaller rulers are able to take account of finer detail.

**montes** On *\*Mars*, a mountain range, e.g. Montes Alpes, Montes Cordillera.

**month degrees** The excess of mean monthly temperatures above 6 °C (43 °F), added together and used as accumulated temperature (indicative of

conditions for vegetation growth) in some climate classifications (e.g. that of A. Miller, 1951).

**Montian** See DANIAN.

**montmorillonite** An important \*clay mineral of the approximate composition  $\{Al_4[Si_3AlO_{10}]_2(OH)_4\}^{2-}.nH_2O$  with some  $K^+$ ,  $Na^+$ , or  $Ca^+$  ions also present; it belongs to the 2:1 group of \*phyllosilicates (sheet silicates) and the montmorillonite or \*smectite group includes \*bentonite; most members of this group are designated expansive clays by engineers, because they can accommodate many water molecules into their structure and they all possess an overall negative charge; sp. gr. variable but 2.0–2.7; \*hardness 2; \*monoclinic; white to grey with tints of blue, pink, pink-red, and green; dull \*lustre; usually occurs in \*massive, \*microcrystalline aggregates of very fine, scale-like crystals, but in soils as a hydrous aluminosilicate \*clay mineral with a layer-lattice structure (two sheets of \*tetrahedral silicon crystals enclosing a sheet of \*octahedral aluminium crystals) that expands when water enters between layers, thus making a soil material, typically with variable water content, subject to swelling and contraction; found very extensively, it results from the decomposition of volcanic ashes in marine basins and also occurs in the weathering crust of \*basic \*igneous rocks, e.g. \*diabases, \*basalts, \*gabbros, and \*peridotites. It is used extensively as an absorbant, for refining out suspended matter, in the textile and chemical industries.

**monzodiorite** A coarse-grained \*igneous rock consisting of \*essential \*plagioclase feldspar, \*orthoclase feldspar, \*hornblende, and \*biotite, with or without \*pyroxene. Plagioclase is the dominant feldspar making up 60–90% of the total feldspar and varying from \*oligoclase to \*andesine in composition. The presence of the orthoclase feldspar distinguishes this rock from a \*diorite.

**monzogabbro** A coarse-grained \*igneous rock consisting of essential \*plagioclase feldspar (see ESSENTIAL MINERAL), \*orthoclase feldspar, and \*pyroxene, with or without \*biotite. Plagioclase of labradorite composition is the dominant feldspar, making up 60–90% of the total feldspar. The presence of orthoclase feldspar distinguishes this rock from a \*gabbro.

**monzonite (syenodiorite)** A coarse-grained *\*igneous* rock consisting of essential *\*plagioclase* feldspar (see *ESSENTIAL MINERAL*), *\*orthoclase* feldspar, *\*pyroxene*, and *\*biotite*. The plagioclase feldspar and orthoclase feldspar are in roughly equal proportions, the plagioclase forming between 40% and 60% of the total feldspar.

**moom** A satellite that orbits inside the rings of its planet.

**Moon** The *\*Earth's* *\*satellite*, with a mass  $7.35 \times 10^{22}$  kg (about 1/81 that of the Earth), density  $3344 \text{ kg/m}^3$ , and radius 1737.5 km. The average Moon–Earth distance is 384 500 km. The Moon has no atmosphere and surface temperature extremes range from  $127 \text{ }^\circ\text{C}$  to  $-173 \text{ }^\circ\text{C}$ . A feldspathic lunar highland crust 60–120 km thick overlies a silicate mantle. Basaltic lavas cover 17% of the surface. There is probably a small iron core of 300–400 km radius (2–3% of lunar volume).

**moonquake** Seismic activity on the *\*Moon* resulting from internal sources (differential cooling or tidal distortions) or external sources, e.g. meteoritic impact. It is the lunar equivalent of an *\*earthquake*. All moonquakes are less than 2 on the *\*Richter* magnitude scale. About 3000 per year were recorded by the Apollo *\*seismometers*. The lunar seismic signals have very low attenuation and a large degree of wave scattering, consistent with travel through a brecciated crust. Most moonquakes occur at about 1000 km depth and are probably induced by tidal stresses, related to periodic changes in the Earth–Moon distance. No lunar tectonic activity is evident. Rare surface moonquakes are caused by the impacts of *\*meteorites*.

**moonstone** A variety of adularia. See *ALKALI FELDSPARS*; *PERTHITE*.

**mor** Type of surface *\*humus* *\*soil horizon* that is acid in reaction, lacking in microbial activity except that of fungi, and composed of several layers of organic matter in different degrees of decomposition.

**moraine** Term originally applied to the ridges of rock debris around Alpine *\*glaciers*. Subsequently its meaning has been widened to include all of the rock debris deposit. For example, ‘ground’ moraine may denote an irregularly undulating surface of *\*till*, glacial drift, or *\*boulder clay*; or it may describe the deposit itself. An ‘end’ or ‘terminal’ moraine is a ridge of glacially deposited material laid down at the leading edge of an active glacier. Its height is in the range 1–100 m, and it is accumulated by a

combination of glacial dumping and pushing. A 'recessional' moraine is morphologically similar and is laid down at the terminus of a glacier during a period of still-stand that interrupts a long period of retreat of the ice margin. A 'lateral' moraine is a ridge of debris at the margin of a valley glacier and largely derived from rock fall. It is a prominent feature of many contemporary Alpine glaciers. A 'medial' moraine results from the merging of lateral moraines when two glaciers converge. A 'washboard' moraine is a single ridge in a closely spaced pattern (perhaps 9–12 per kilometre) and stands some 1–3 m above the adjacent depressions. It is found in the 'end' moraine belt. A 'push' moraine is a morainic ridge made of unconsolidated rock debris and pushed up by the *\*snout* of an advancing glacier. *See also* [DE GEER MORAINE](#); [FLUTED MORAINE](#); [HUMMOCKY MORAINE](#).

**Morarian orogeny** What may have been an episode of mountain building that occurred about 1050–730 Ma ago, prior to the *\*Caledonian* orogeny, affecting the Morar area north-west of the Great Glen Fault, in what is now Scotland. Alternatively, the evidence suggesting an *\*orogeny* may be explained as a phase of *\*pegmatite* formation. If it was a genuine orogeny it may mark the start of the *\*subduction* that led to the closing of the *\*Iapetus* Ocean.

**MORB** *See* [MID-OCEAN-RIDGE BASALT](#).

**Morisawa flood-peak formula** *See* [FLOOD PEAK FORMULAE](#).

**morphoclimatic zone** Area that is characterized by a distinctive assemblage of land-forms and that coincides with a major climatic zone. It is believed that the land-forms largely result from the action of a unique combination of surface processes controlled by climate.

**morphogen** An environmental factor, such as a chemical substance, that establishes a concentration gradient affecting *\*morphogenesis* in organisms exposed to it.

**morphogenesis** The development of the form and structure of an organism by growth and differentiation.

**morphogenetic zone** *See* [LINEAGE-ZONE](#).

**morphological mapping** A method of mapping the form of a land surface. It is based on the assumption that land surfaces can be divided into a

number of components, each of which has a uniform gradient or curvature, and which are separated by abrupt or gentle changes of slope. The nature of the change of slope is shown on the map by a standard set of symbols. The angle of each component may be measured instrumentally.

**morphological system** In *\*geomorphology*, a theoretical construct consisting of the relationship between the physical properties of a natural *\*system* (geomorphological). For example, the physical dimensions of a beach (angle of slope seaward, average grain size, porosity, and moisture content) may be related to each other in an orderly manner, and so constitute a morphological system, and the geometric properties of a valley-side slope are typically correlated with certain characteristics of soil and vegetation.

**morphology** Form and structure of individual organisms or land surfaces.

**morphometric analysis** See DRAINAGE MORPHOMETRY.

**morphometrics** A technique of taxonomic analysis using measurements of the form (*\*morphology*) of organisms and typically involving multivariate statistics.

**morphometry (geomorphometry)** The quantitative analysis of the structure and form (morphology) of land-forms.

**morphospace** In *\*theoretical* morphology, all the possible forms an evolving taxon might take from a given set of initial parameters.

**morphospecies** A group of biological organisms that differs in some morphological respect from all other groups.

**morphotectonics** The study of the relationship between *\*tectonics* and the development of land-forms.

**morphotype** In taxonomy, a specimen chosen to illustrate a morphological variation within a species population.

**Morrowan** See BASHKIRIAN.

**mortar texture** Closely interlocking *\*crystals* of fine *\*grain* size, formed by the crushing and *\*recrystallization* of a rock that was originally coarser grained. It is a texture produced by shearing of the rock during *\*dynamic metamorphism*. In low-grade dynamic metamorphic rocks, deformation

occurs primarily along crystal boundaries, producing an envelope of mortar texture around each crystal.

**Mortensnes** A *\*stage* and substage of the Varangian stage, dated at 640–630 Ma ago (Int. Commission on Stratigraphy, 2004), preceded by the *\*Smålfjord* and followed by the *\*Wonokan*.

**mosaic evolution** Differential rates of development of various adaptive attributes within the same evolutionary lineage. For example, a particular *\*taxon* might show greatly different rates of change with respect to the head, body, and limbs. This is a common phenomenon and makes the reconstruction of transitional fossil types very difficult.

**mosaic heterochrony** *\*Heterochrony* in which a number of heterochronic processes occur simultaneously, so different parts of an organism develop at different rates. *Compare* ASTOGENETIC HETEROCHRONY; ONTOGENETIC HETEROCHRONY.

**mosaic texture** Interlocking *\*crystals* of fine to medium *\*grain* size, displaying triple-point contacts between grains. Such a texture can form during metamorphic *\*recrystallization* of a rock (see METAMORPHISM) in response to temperature change without any imposed directional stress.

**Mosasauridae (mosasaurs)** (order Squamata, suborder Sauria) A family of marine lizards, fossils of which have been found in Upper *\*Cretaceous* rocks throughout the world. They evolved into many types but all became extinct at the end of the Cretaceous. Mosasaurs were large, some reaching 9 m in length, with a long, slim body, short neck, and long head. The tail was used for swimming and the limbs for steering; the digits may have been webbed. The teeth were set in pits rather than being fused to the jaws. Most mosasaurs fed on fish, although some are believed to have fed on molluscs.

**mosasaurs** See MOSASAURIDAE.

**Moscovian** A *\*stage* in the *\*Pennsylvanian \*epoch* dated at 311.7–306.5 Ma ago (Int. Commission on Stratigraphy, 2004), comprising the Vereiskian, Kashirskian, Podolskian, and Myachkovskian Russian stages, preceded by the *\*Bashkirian* and followed by the *\*Kasimovian* epochs. It is roughly contemporaneous with the *\*Atokan* and *\*Desmoinesian* (N.

America) and the **\*Westphalian** B, C, and D plus the lowermost **\*Stephanian** (western Europe).

**Moskva Drift** A **\*drift** unit that overlies the deposits of the **\*Odintsovo \*Interstadial**, occurring in European Russia. It is equivalent to part of the **\*Saalian** of western Europe.

**moss agate** See AGATE.

**Mössbauer spectroscopy** An analytical spectroscopic technique based on the fact that certain **\*nuclides**, solidly built into a **\*crystal \*lattice**, and at temperatures significantly below a certain temperature, emit **\*gamma rays**. The crystal absorbs the recoil momentum, making it negligible because of the large number of atoms in the lattice, and there is virtually no loss of energy by the gamma photons.

**mossy** Applied to dendritic (branch-like) aggregates of **\*minerals** that are frequently secondary chemical precipitates, such as iron or manganese oxides (e.g. moss agate and mocha stone).

**mother cloud** Incipient cloud from which a well-defined cloud type can form and develop. See also CLOUD CLASSIFICATION.

**mother-of-coal** See FUSAIN.

**mother-of-pearl cloud** See NACREOUS CLOUD.

**mottling** Patchwork of different colours in **\*mineral soil** (usually orange or rust against a background of grey or blue) which indicates periods of anaerobic conditions.

**motu** A small island in an **\*atoll** that is made from wave-borne **\*detritus**. Some motus are protected from erosion by the inclusion of **\*beach rock** or the presence of a **\*pakakota**.

**Motuan** A New Zealand **\*stage** (104–99.6 Ma ago) of the Early **\*Cretaceous \*epoch**, preceded by the **\*Urutawan** and followed by the **\*Ngaterian**.

**mould** See FOSSILIZATION.

**moulin** A system of tunnels made by meltwater through a glacier.

**mouldic porosity** See MOLDIC POROSITY.

**Mount Agung** A stratovolcano on Bali, Indonesia, which erupted violently in March 1963, emitting large quantities of gases and ash. Sulphate particles rose to 22–23 km, spread throughout the world within about six months, remained aloft for several years, and produced spectacular twilight displays of colour. The temperature in the lower stratosphere rose by 6–7 °C shortly after the eruption and surface temperatures were depressed by 0.18–1.3 °C for several years. The volcano erupted again in November 2017, following more than 800 volcanic earthquakes in September. A phreatic eruption (see PHREATIC ACTIVITY) on 21 November sent ash to a height of about 3800 m and magmatic eruptions occurred on 25 and 26 November, after which the volcano subsided, although ash reached 2000 m on 8 December. The 2017 eruptions were rated VEI2 on the \*Volcanic Explosivity Index.

**mountain breeze** See MOUNTAIN WIND.

**mountain wind (mountain breeze)** Term commonly used to describe a \*katabatic wind or breeze and its counterpart, the upslope anabatic wind or breeze occurring on warm days over heated mountain slopes.

**mouthbar** An accumulation of \*sediment at the mouth of a river that eventually obstructs the river, causing the water flow to divide around it. A mouthbar comprises sediments that are coarse on the upstream side, becoming gradually finer downstream.

**moveout (stepout) 1.** The difference between the \*two-way travel times of reflected energy detected at two \*geophone \*offset distances. **2. (normal moveout, NMO,  $\Delta T$ )** At an offset distance  $x$ , the difference in two-way travel time between the detected reflection event at  $x$  ( $t_x$ ) and at zero offset ( $t_0$ ), such that  $\Delta T = t_x - t_0 \approx x^2/(2V^2t_0)$ , where  $V$  is the seismic velocity of the medium above the reflector. **3. (dip moveout, DMO,  $\Delta T_d$ )** In the case of a planar dipping reflector, the difference between the moveout updip and downdip, proportional to the angle of dip  $\theta$ , such that:  $\Delta T_d = 2x \sin \theta / V$ , where  $x$  is the offset distance from the mid-point.

**MPT** See MID-PLEISTOCENE TRANSITION.

**$M_{\text{sat}}$**  See SATURATION MAGNETIZATION AND MOMENT.

**M-shape** Characteristic shape shown in an analysis profile of nickel distribution in *\*Widmanstätten structure* in *\*meteorites*. The temperature-dependence of the *\*diffusion* of nickel within the metal *\*phase* away from kamacite (Ni–Fe alloy with 6% Ni) into taenite (Ni–Fe alloy with 30% Ni), and the faster rate of diffusion in kamacite, leads to disequilibrium nickel distribution and the M-shape.

**MSU** See MICROWAVE SOUNDING UNIT.

**MTSAT** See MULTIFUNCTION TRANSPORT SATELLITE.

**mud, drilling** During rotary drilling, the fluid mud that is circulated to cool the drill *\*bit*, convey rock chippings to the surface, and to seal permeable layers and fractures. The drilling mud is usually maintained under pressure to prevent the drill stem being blown out if it penetrates a pocket of pressurized gas. The muds may be oil- or water-based, and are frequently *\*thixotropic* *\*bentonites*, lime, or *\*barite*.

**mud cake (cake)** The solid clay deposit formed in a *\*borehole* on a permeable layer when the liquid *\*mud filtrate* permeates into the surrounding rocks.

**mudcracks** See DESICCATION CRACKS.

**mud drape** A layer of mud deposited over a pre-existing morphological feature, e.g. a *\*bar*, *\*ripple*, or *\*dune*.

**mud filtrate (filtrate)** The liquid part of a drilling *\*mud* that can penetrate into a permeable layer, leaving behind a solid *\*mud cake* and totally or partially replacing the *\*groundwater* in the rocks.

**mudflat** Area of a coastline where fine-grained silt or sediment and clay is accumulating. Its development is favoured by ample sediment, by sheltered conditions, and by the trapping effect of vegetation. It is an early stage in the development of a *\*salt marsh*.

**mudflow** 1. Heavily loaded ephemeral stream whose *\*viscosity* increases with evaporation as it flows over a desert fan. 2. Rapidly moving variety of *\*earthflow*. This is a typical phenomenon of areas underlain by sensitive clays which may liquefy and flow following a shock, perhaps initiated by sliding. 3. See LAHAR.

**mud log** 1. The record produced by a *\*microlog* measuring the *\*resistivity* of the *\*mud cake* in a drill hole. 2. 'Mud log' is also the term used for the record maintained by well site geologists of the cuttings recovered from the hole.

**mudlump** See MUD VOLCANO.

**mud mound** A build-up of *\*carbonate \*sediment* in the form of a bank or mound, dominated by mud. The accumulation of mud occurs by its deposition in the lee of *in situ* organisms, such as corals (*\*Anthozoa*) or *\*crinoids*; by the sweeping of mud into banks by currents; or by the entrapment and precipitation of carbonate mud by *\*algae* and other organisms acting as baffles.

**mudrock** A lithified mud (see LITHIFICATION).

**mud roll** See GROUND ROLL.

**mud/shale diapir** See MUD VOLCANO.

**mudstone** 1. *\*Argillaceous* or clay-bearing *\*sedimentary rock* which is non-plastic and has a *\*massive* or non-foliated appearance. Compare CLAYSTONE. 2. See LIME MUD; DUNHAM CLASSIFICATION; EMBRY AND CLOVAN CLASSIFICATION.

**mud-support (matrix-support)** Texture of *\*sedimentary rocks* in which there is complete separation of larger fragments or *\*clasts* from each other within the finer-grained *\*matrix*. Compare GRAIN SUPPORT.

**mud volcano (mudlump, mud/shale diapir)** A volcanic feature found in places with a high rate of sedimentation and a plentiful supply of mud. Mud volcanoes are fed from a chamber filled with mud and linked to the surface by a *\*clastic \*dyke*. They range from less than one metre to several hundred metres wide and up to 100 m high, and erupt hot mud.

**mugearite** See ANDESITE; BENMOREITE.

**mull** Type of surface *\*humus \*soil horizon* that is chemically neutral or alkaline in reaction, that is well aerated, and that provides generally favourable conditions for the decomposition of organic matter. Mull humus is well decomposed and intimately mixed with mineral matter.

**mullion structure** A *\*rodding* structure composed of elements tens of centimetres wide. The origin of mullions and rodding is not known with certainty but they are thought to be a type of *\*lineation* associated with folding at a competent–incompetent interface (see [COMPETENCE](#); [INCOMPETENCE](#)).

**multichannel seismic reflection** A *\*reflection* event which occurs across a number of different seismic channels, data from which can be used to enhance data processing and the subsequent interpretation of seismic sections.

**Multifunction Transport Satellite (MTSAT)** A Japanese *\*geostationary* satellite programme that augments air-traffic control services by improving communications and position information and also acquires meteorological data. The satellite was launched on 26 February 2005, from the Tanegashima Space Center, Japan.

**multifurcation** In a *\*phylogenetic tree*, the occurrence of a split in an ancestral branch into more than two branches at an *\*internal node*, because the order in which the progenic branches occur cannot be determined.

**multilocular** See [UNILOCULAR](#).

**multiple** Generally, re-reflected seismic energy. More specifically, multiple *\*reflection* which is detrimental to the data and should be removed by data processing. ‘Short-path’ multiples are *\*ghosts* and near-surface phenomena; ‘long-path’ multiples, which are easier to identify and remove, are due to extra reflections deep within the subsurface.

**multiple common-depth-point coverage** A method of seismic *\*profiling* in which the same part of the subsurface is sampled on different records. See [FOLD](#).

**multiple twinning** See [LAMELLAR](#).

**multiplexing (muxing)** The transmission of several channels of information over a single channel without interference and the subsequent recording of that information on a hard disc or magnetic tape. The digital data recorded are in the order: channel 1, sample 1; channel 2, sample 1; ... channel *n*, sample 1; channel 1, sample 2; channel 2, sample 2;...channel *n*, sample 2; etc., which is a very efficient way to record information on disc

or magnetic tape. **\*Demultiplexing** is the process by which the tape is made readable by a computer and its data are made available for processing.

**multi-ring basin (ringed basin)** A large basin excavated by **\*asteroidal** or **\*planetismal** impact on a planetary surface. Several concentric rings of mountains surround the basin. The ratio of the diameters of adjacent rings approximate to  $\sqrt{2}:1$ . The minimum dimension for rings to occur is about 300 km on the Moon; elsewhere examples of up to 3000 km diameter are known (Valhalla basin on Callisto). The classic example is the lunar Mare Orientale, 920 km diameter, with three mountain rings. Typical relief is 3 km. The energies involved in excavating the basins are of the order of  $10^{27}$  J.

**multispectral imaging** See **IMAGING**.

**multispectral scanner** In **\*remote** sensing, a **\*line scanner** which records data in several different **\*channels**.

**multispectral scanning systems** The simultaneous use of several different sensors to obtain images or spectral information from various portions of the **\*electromagnetic spectrum**. An example was the multispectral scanner on the Landsat Earth-orbiting satellites, which simultaneously recorded green, red, and two infrared spectral bands. Since different materials (e.g. soil, rock, vegetation) reflect differing amounts of light at different wavelengths, they can be identified by their characteristic spectral 'signatures'.

**multistate character** A **\*character** that can occur in several **\*character** states.

**multistorey sandbody** A series of **\*sandstone** beds, each deposited by the infilling of a river **\*channel**, stacked one above the other with little or no intervening **\*mudstone**. The multistorey body is formed by the repeated and rapid migration of the channel network over the **\*alluvial** plain, so allowing little chance for the preservation of fine **\*floodplain** **\*sediment**.

**Multituberculata** (class **\*Mammalia**) Extinct order of rather rodent-like mammals, which appeared first in the late **\*Jurassic** in Europe, flourished during the **\*Cretaceous** and **\*Palaeocene**, but became extinct during the **\*Eocene**. Probably they were the first herbivorous mammals, with skull and teeth analogous to those of rodents. Most were small, but some attained the

size of modern woodchucks. The limbs sprawled more widely than those of most mammals. The olfactory bulbs were large, which suggests that the animals depended heavily on their sense of smell. The skull was massive, but unlike that of other mammalian groups. The multituberculates appear to have been a side branch from the main line of mammalian evolution, and they are believed not to be related closely to other groups.

**Multi-User System for Earth Sensing** See [MUSES](#).

**multivariate analysis** In a statistical analysis, the measurement of several different attributes of each unit of observation.

**Munsell colour** Soil-colour system, devised originally in the USA by the painter and art teacher Albert Henry Munsell (1858–1918), that is based on the three variables of colour: hue, value, and chroma, and given notation such as ‘10YR 6/4’. It is now used as an international method for reporting soil colour.

**Murchison, Roderick Impey** (1792–1871) A gentleman of private means who devoted much of his life to science, Murchison turned professional when he became director of the Geological Survey in 1855. His *Silurian System* (1839) was the first stratigraphic system defined by fossil content, rather than [\\*lithology](#). He helped clarify the [\\*Devonian](#) system and, after a visit to Russia, the [\\*Permian](#).

**Murray, John** (1841–1914) A Canadian-born biologist and oceanographer, Murray did most of his work in Britain. He was leader of the [\\*Challenger](#) expedition and took part in other oceanographic investigations. He completed the publication of the scientific reports of the *Challenger* survey and produced a bathymetric chart of the oceans.

**muris (wall cloud)** See [CLOUD CLASSIFICATION](#).

**Muschelkalk** **1.** The sea-way (Muschelkalk Sea) that extended across north-western Europe during the [\\*Triassic](#), from the present-day UK in the west to northern Germany and Poland in the east. It was bordered to the south by the Bohemian [\\*Massif](#) and in the north by the Baltic Shield (see [BALTICA](#)). **2.** The mid-Triassic, when using the European tripartite divisions of the Triassic system: Bunter, Muschelkalk, and Keuper. The German word Muschelkalk means ‘shelly limestone’.

**muscle scar** A depressed or raised marking on the interior of a shell, usually of a brachiopod (*\*Brachiopoda*) or bivalve (*\*Bivalvia*), which was the point of attachment of a muscle. In articulate brachiopods (*\*Articulata*) there are commonly two pairs of muscle scars occurring in the floor of the *\*dorsal* and *\*ventral* valves, one pair formed by the adductor muscles that close the shell and the second pair by the diductors, that open it. There are additional muscles present in inarticulate brachiopods (*\*Inarticulata*) and these also leave scars. In bivalves there are commonly two muscles present (dimyarian); these are the adductor muscles that close the shell against the opening moment of the ligament. In some species the scars are equal in size (isomyarian), in others unequal (anisomyarian) and the smaller scar is always the anterior one. Sometimes the anterior muscle is lost and the remaining muscle may increase in size; in this condition the bivalve is 'monomyarian'. Connecting the two scars on the interior of the valve there is the pallial line, a linear depression marking the inner margin of the thickened *\*mantle* edges that may be marked by an inward deflection at the posterior part, the pallial sinus. This defines the space for the retraction of the *\*siphons* and the place of attachment of the siphonal retractor muscles.

**muscovite** An important rock-forming *\*mineral* and member of the mica group with composition  $K_2Al_4[Si_3AlO_{10}]_2(OH,F)_4$  (see *MICAS*); when K is replaced by Na the mineral is called paragonite and is related to the *\*phyllosilicates* (sheet silicates) *\*lepidolite* and *\*zinnwaldite* (both lithium-bearing micas); sp. gr. 2.8–2.9; *\*hardness* 2.5–3.0; *\*monoclinic*; normally colourless to very pale grey to green or light brown; vitreous to pearly *\*lustre*; crystals *\*tabular*, *\*hexagonal*, and also occur in foliated masses and disseminated flakes; *\*cleavage* basal perfect {001}; widely distributed in *\*igneous* *\*alkali* *\*granites*, in *\*pegmatites*, as a *\*secondary mineral* resulting from the decomposition of *\*feldspars*, and can be a major constituent of *\*clastic* sediments (e.g. *\*sandstone* and *\*silts*).

**MUSES (Multi-User System for Earth Sensing)** An Earth-imaging platform that hosts instruments such as high-resolution digital cameras and hyperspectral imagers for Earth viewing. It was launched on 3 June 2017, to the International Space Station.

**Muses-C (Hayabusa)** A JAXA mission to return samples from asteroid Itokawa. It was launched in 2003, after which the mission name was

changed from Muses-C to Hayabusa, and arrived at Itokawa in September 2005. Its attempts to land failed and it was unable to collect samples, but it returned large amounts of data. Hayabusa 2 reached Ryugu in June 2018.



[http://global.jaxa.jp/projects/sat/muses\\_c/](http://global.jaxa.jp/projects/sat/muses_c/)

- About Asteroid Explorer 'Hayabusa' (MUSES-C).

**mushroom rock** See PEDESTAL ROCK.

**Mustelidae** See CARNIVORA.

**mutation** 1. Process by which a *\*gene* or *\*chromosome* set undergoes a structural change. 2. Gene or chromosome set that has undergone a structural change. Mutations are the raw material for *\*evolution*: they provide the source of all variation. For mutations to affect subsequent generations, though, they must occur in gametes (reproductive cells) or in cells destined to be gametes, since only then will they be inherited. A mutation that occurs in a body-cell is called a 'somatic' mutation: it is transmitted to all cells derived, by mitosis, from that cell. Most mutations, however, are deleterious; evolution progresses through the few that are favourable.

**mutation rate** In *\*phylogenetics*, the number of *\*mutations* arising at a single nucleotide site per gene per unit time.

**mutatus** Form of cloud evolution in which the cloud changes shape rapidly because of new cloud masses developing from it.

**muxing** See MULTIPLEXING.

**Myachkovskian** A *\*stage* (307.2–306.5 Ma ago) and substage of the *\*Moscovian* stage, preceded by the *\*Podolskian* and followed by the *\*Krevyakinskian* (Late *\*Pennsylvanian* *\*epoch*).

**mylonite** Rock produced in zones of tectonic dislocation, e.g. *\*fault* and shear zones. Mechanical crushing and grinding or *\*cataclasis* produces a rock that has a *\*foliation* which is often crude but which is sometimes very well developed, and that has a much finer grain size than its precursor. Mylonite is well laminated and often hard and splintery. The name literally

means 'milled rock' from the Greek *mylon*, a grinding mill. See [DYNAMIC METAMORPHISM](#).

**myophore** A plate or rod-like structure on the inside of the shell of some bivalves ([\\*Bivalvia](#)). It occurs at the centre of the [\\*hinge](#) for the attachment of muscle.

**Myriapoda** A class of [\\*Arthropoda](#) that is no longer recognized, because relationships between its members are considered to be superficial. Members of the Myriapoda are now allotted to the classes Chilopoda (centipedes), Diplopoda (millipedes), Pauropoda (small, soft-bodied animals resembling millipedes but not closely related to them), and Symphyla (small, fast-moving animals that resemble centipedes but have mouthparts more like those of insects).

**Mytiloida** (phylum [\\*Mollusca](#), class [\\*Bivalvia](#)) Order of [\\*epifaunal](#), [\\*byssate](#) bivalves which have an equivalve shell that has a highly non-equilateral shape and a prismatic-nacreous microstructure (e.g. *Mytilus edulis*, the common mussel). They have a [\\*dysodont](#) [\\*dentition](#), the ligament is elongate and [\\*opisthodetic](#). Mytiloida have an anisomyarian musculature (i.e. the posterior adductor muscle and its scar are larger than the anterior adductor muscle). The pallial line is complete, the [\\*siphons](#) poorly developed. They first appeared in the [\\*Devonian](#). See [MUSCLE SCAR](#).



# N

**N** See NEWTON.

**n** See NANO-.

**NA** See OBJECTIVE.

**Nabarro–Herring creep** A form of *\*diffusion creep* in which atoms migrate within the *\*crystal \*lattice*. See also COBLE CREEP.

**nabkha** See SHRUB-COPPICE DUNE.

**nacreous cloud (mother-of-pearl cloud)** Type of cloud seen occasionally at great altitude (approximately 22–4 km), just before sunrise or after sunset, characterized by iridescent colouring. The cloud is fine and rather lenticular.

**nacrite** See KAOLINITE.

**nadir** In *\*remote sensing*, the point on the ground directly beneath a remote sensing system.

**NADW** See NORTH ATLANTIC DEEP WATER.

**Nafe–Drake relationship** An empirical relationship between the *\*P-wave* velocity and density of water-saturated *\*sediments* and *\*sedimentary rocks*. It is commonly used to evaluate the density of sedimentary rocks in shallow seismic surveys, but errors of the order of 100 kg/m<sup>3</sup> are involved. See also DENSITY OF ROCKS.

**Nagssugtoqidian orogeny** A late *\*Archaean* and *\*Proterozoic* phase of mountain building affecting a belt 240–300 km wide in what is now western Greenland. It comprised two major episodes, the first about 2600 Ma ago,

the second about 1900–1500 Ma ago, making it approximately contemporaneous with the *\*Hudsonian*, *\*Laxfordian*, and *\*Svecofennian orogenies*.

**Naiad (Neptune III)** A satellite of *\*Neptune*, with a diameter of 58 km; visual albedo 0.06.

**Namibian** A South African *\*stage* (900–542 Ma ago), preceded by the *\*Mokolian* and followed by the Early *\*Cambrian \*epoch*.

**Nammalian** See SMITHIAN.

**Namurian** A European *\*stage* (326.4–315 Ma ago) in the *\*Silesian \*epoch*, underlain by the *\*Dinantian*, overlain by the *\*Westphalian*, and subdivided into Namurian A, B, and C. It is dated at 326.4–315 Ma ago and is contemporaneous with the Millstone Grit series (western Europe) and the *\*Serpukhovian* and Lower *\*Bashkirian* series (eastern Europe).

**nannofossil** See NANOFOSSIL.

**nano-** (**n**) From the Greek *nanos* meaning ‘dwarf’, a prefix meaning ‘extremely small’ (e.g. *\*nannofossil*). Attached to SI units it denotes the unit  $\times 10^{-9}$ .

**nannofossil (nannofossil)** A *\*fossil* of the smallest member of the *\*plankton* (the nanoplankton). Nanofossils are of plants and include various forms, e.g. the coccolithophores (*\*coccolithophorids*), which are 5–60  $\mu\text{m}$  in size and have been used as stratigraphic indicators in addition to the larger *\*microfossils*.

**nanoplankton** Marine planktonic organisms 2.0–20  $\mu\text{m}$  in size.

**NanoRacks-QB50** A project that uses the International Space Station to deploy a constellation of twenty-eight *\*CubeSats* to study the upper regions of the atmosphere. It was launched on 18 April 2017.

**NanoSatC-Br1** See BRAZILIAN CUBESAT PROJECT-1.

**nanosatellite** A space satellite with a *\*wet mass* of 1–10 kg.

**nappe** From the French *nappe*, meaning ‘cover’, a thrust mass or folded body in which the *\*fold limbs* and axes are approximately horizontal.

**Naraoiidae** An order of *\*Trilobita* that lived in the Middle *\*Cambrian*. They were uncalcified and had no thoracic segments.

**Narizian** A Californian *\*stage* (48–35 Ma ago) in the *\*Eocene*, underlain by the *\*Ulatisian*, overlain by the *\*Refugian*, and roughly contemporaneous with the upper *\*Lutetian* and *\*Bartonian* stages.

**NASA** The National Aeronautic and Space Administration, the agency of the US federal government that was established under the National Aeronautics and Space Act 1958 to plan, direct, and conduct all US aeronautical and space activities, other than those that are primarily military.

**naticiform** Applied to a gastropod (*\*Gastropoda*) shell that is globular and umbilicate, resembling those of the genus *Natica*. See **UMBILICUS**.

**native element** An element which occurs in a free state as a *\*mineral*, e.g. *\*gold*, *\*copper*, and *\*carbon*.

**natric horizon** A mineral *\*soil horizon* that is developed in a subsurface position in the profile, that satisfies the definition of an *\*argillic horizon*, and that also has a columnar structure and more than 15% saturation of the exchangeable *\*cation* sites by sodium.

**natrolite** Member of the *\*zeolite* group of *\*minerals*, with the formula  $\text{Na}_2\text{Al}_2\text{Si}_3\text{O}_{10}\cdot 2\text{H}_2\text{O}$ ; sp. gr. 2.2; *\*hardness* 5.0; white; *\*massive*, fibrous, or crystalline; occurs as a *\*secondary mineral* infilling cavities in *\*basic* volcanic rocks and as an alteration product of *\*nepheline*.

**natron lake (soda lake)** A saline lake, rich in the sodium carbonate salt natron ( $\text{Na}_2\text{CO}_3\cdot 10\text{H}_2\text{O}$ ).

**‘natural break’** Concept that the Earth’s history has been punctuated by natural worldwide changes that should be discernible in the stratigraphic record.

**natural cast** See **FOSSILIZATION**.

**natural gas** Gaseous *\*hydrocarbons*, chiefly methane ( $\text{CH}_4$ ), ethane ( $\text{C}_2\text{H}_6$ ), propane ( $\text{C}_3\text{H}_8$ ), and butane ( $\text{C}_4\text{H}_{10}$ ), trapped in *\*pore spaces* in rocks with or without liquid *\*petroleum*. It has a high heat value and burns without smoke or soot; it provides raw material for the chemical industry

for making plastics, detergents, fertilizers, etc. Gas of this composition is also termed 'natural gas' if it occurs as a gas chimney or after production.

**natural gas hydrate** A type of *\*clathrate* in which a naturally occurring gas, mainly methane, is held within a lattice of water ice. Such hydrates form under conditions of pressure and temperature found in the upper 300–2000 m of marine sediments where the water temperature is at or below 0 °C and the pressure is greater than 4 MPa, and occur offshore or on land, often beneath *\*permafrost*. The hydrates can block gas transmission pipes and disturbing them can cause the release of methane (a *\*greenhouse gas*). Natural gas hydrates are likely to be developed as a source of recoverable methane and the amount present in sediments, measured as carbon, is estimated to be more than twice that of all other recoverable and non-recoverable fossil fuels combined.

**natural remanent magnetism (NRM)** The *\*remanent magnetization* of rocks and naturally occurring objects that has been acquired by normal processes, i.e. excluding laboratory-induced magnetizations. In *\*igneous* rocks and fired archaeological materials, NRM is normally of thermal or chemical origin; in *\*sediments* it is usually of depositional origin; and in *\*sedimentary rocks* it is usually of chemical origin.

**natural selection ('survival of the fittest')** Complex process in which the total environment determines which members of a species survive to reproduce and so pass on their *\*genes* to the next generation. This need not necessarily involve a struggle between organisms.

**natural theology (physico-theology)** A philosophy which tries to link the study of natural phenomena with the notion of divine providence, stressing that harmony and order in nature are evidence of God's design. It became important in 18th-century England, being associated with the work of John Ray, William Paley, and many others. The 19th-century Bridgewater Treatises were the last major exposition of natural theology. See [BUCKLAND](#).

**nautical mile** Length of mile used in ocean navigation, equivalent to one minute (1/60°) change in latitude. It is internationally defined as being equal to 1852 metres.

**nautilicone** See [CONVOLUTE](#).

**nautiliconic** Applied to the conch of a *\*cephalopod* when it is coiled and highly *\*involute*.

**Nautiloidea** (phylum *\*Mollusca*, class *\*Cephalopoda*) Subclass of cephalopods which possess a multichambered, external shell composed of calcium carbonate, which is siphunculate (see *SIPHUNCLE*) and may be coiled. The animal lives in the last-formed chamber, the body chamber. The gill structure is tetrabranchiate (four-gilled). Simple *\*sutures* are produced by contact between the internal *\*septa* and the shell wall. The subclass includes the oldest cephalopods, first recorded from Upper *\*Cambrian* rocks. They diversified and became common throughout the *\*Palaeozoic*, but were greatly reduced at the end of this *\*era*. Further diversification occurred in the *\*Mesozoic*, but the group dwindled again in the *\*Cenozoic*. There is only one extant genus, *Nautilus*, which dates from the *\*Oligocene*. See also *PLECTRONOCERAS CAMBRIA*.

**Nazca Plate** The lithospheric *\*plate* which is subducting under the *\*South American Plate* along the line of the *\*Peru–Chile Trench*. Its other *\*plate margins* are the *\*ridges* and *\*transform faults* which separate it from the *\*Pacific*, *\*Cocos*, and *\*Antarctic Plates*. The Nazca Plate is thought to have decreased in size; the former, more extensive plate has also been called the Phoenix Plate.

**neap tide** Tide of small range occurring every 14 days, near the times of the first and last quarter of the Moon when the Moon, Earth, and Sun are at right angles. The neap tidal range is 10–30% less than the mean tidal range.

**NEAR** See *NEAR EARTH ASTEROID RENDEZVOUS*.

**Nearctic faunal realm** The fauna of N. America, south to Mexico. At the order and family level the fauna is essentially the same as that of the Palearctic realm, reflecting their former connection via the *\*Bering \*land bridge*, but some genera and more especially species are distinctive to the Nearctic.

**Near Earth Asteroid Rendezvous (NEAR)** The first of the *\*NASA \*Discovery* missions, launched on 17 February 1996. The spacecraft was equipped with an X-ray/gamma ray spectrometer, near-infrared imaging spectrograph, multispectral camera with CCD (see *CHARGE-COUPLED DEVICE*) imaging detector, laser altimeter, and magnetometer. A radio

science experiment was also performed to estimate the gravity field of \*Eros. On 27 June 1997, the NEAR spacecraft approached to within 1200 km of \*Mathilde. The spacecraft reached and achieved orbit around Eros, returning large amounts of data, and at the end of its mission was deliberately crashed to the surface, returning more data.



<https://heasarc.gsfc.nasa.gov/docs/heasarc/missions/near.html>

- NEAR.

**near-field barrier** In the disposal of \*radioactive waste, the containers in which waste is held, the vault in which these are stored, and the immediately surrounding rocks, which prevent the migration of radionuclides, but which provide a level of containment that is expected to fail over some thousands of years. Containment will then be provided by the \*far-field barrier.

**near-infrared** \*Infrared radiation which has a wavelength between 0.7  $\mu\text{m}$  and 2.5  $\mu\text{m}$ . Near-infrared is subdivided into very-near infrared (0.7  $\mu\text{m}$ –1  $\mu\text{m}$ ) and short-wave infrared (1.0  $\mu\text{m}$ –2.5  $\mu\text{m}$ ). Photographic films respond to wavelengths between 0.7  $\mu\text{m}$  and 1.0  $\mu\text{m}$ , hence very-near infrared is also known as photographic infrared.

**near-infrared mapping spectrometer** (near-infrared mapping spectrophotometer; NIMS) A \*remote sensing instrument that makes measurements in the \*near-infrared region of the light spectrum, from which the chemical composition, structure, and temperature of planetary and satellite atmospheres can be determined, as well as the mineral composition and geochemistry of planetary and satellite surfaces.

**near-infrared mapping spectrophotometer** See NEAR-INFRARED MAPPING SPECTROMETER.

**near-shore current system** System of currents caused by wave activity within and adjacent to the \*breaker zone. The current system includes: the shoreward mass transport of water; \*longshore currents; and seaward-moving \*rip currents. This wave-induced current system often has a reversing tidal-current system superimposed on it.

**nebkha** See SHRUB-COPPICE DUNE.

**Nebraskan** The first N. American \*glaciation of the \*Pleistocene epoch, which is identified by deposits found in Nebraska. It began about 1.8 Ma ago and ended 1.65 Ma ago. It is approximately equivalent to the \*Günz glaciation of Europe.

**nebulosus** From the Latin *nebulosus* meaning ‘fog-covered’, a species of cloud that forms a rather indistinct layer or veil with no distinguishable features. *See also* CLOUD CLASSIFICATION.

**necrology** The scientific study of all the processes affecting dead animal and plant material, including decomposition and \*fossilization.

**Nectarian** A \*period in the late \*Hadean, marking the time during which intense bombardment of the Earth ceased and dated at about 3975–3900 Ma ago (Int. Commission on Stratigraphy, 2004).

**Nectarian system** *See* LUNAR TIMESCALE.

**needle ice** *See* PIPKRAKE.

**Néel, Louis Eugène Félix** (1904–2000) A French physicist, Néel became Director of the Centre for Nuclear Studies in Grenoble in 1956. Much of his work was in the field of magnetism, including terrestrial magnetism, and he was the first to explain the existence of remanent \*palaeomagnetism in rocks.

**Néel temperature** The temperature at which the magnetic coupling between two magnetic sublattices in an \*antiferromagnetic material is overcome by thermal vibrations; if the material is heated above the Néel temperature it no longer exhibits antiferromagnetic behaviour until it has cooled to or below this temperature.

**negative inversion** *See* INVERSION.

**nekton** (*adj.* nektonic) Free-swimming organisms in aquatic ecosystems. Unlike \*plankton, they are able to navigate at will. The nekton includes fish, amphibians, and large swimming insects.

**nema** In graptolites, a long, hollow, rod- or thread-like structure extending beyond the \*sicula. *See* GRAPTOLOIDEA.

**Nemakit-Daldynian** A Russian-Kazakhstan *\*stage* (542–534 Ma ago) of the Early *\*Cambrian \*epoch*, preceded by the *\*Ediacaran* and followed by the *\*Tommotian*.

**Nematoda** (nematodes, eelworms, threadworms, roundworms) *\*Phylum* of worms which vary greatly in size, from about 1 mm to 5 cm. The cuticle has flanges and may also have ridges or spines. Morphologically they are all similar, but they occur both as parasites in plants and animals and as free-living forms. They are first known from rocks of *\*Carboniferous* age (e.g. *Scorpiophagus*).

**Nemertini** (proboscis worms, ribbon worms) *Phylum* of unsegmented, non-parasitic worms which are bilaterally symmetrical (see *BILATERAL SYMMETRY*) and elongate. Adults are ciliated (see *CILIUM*) and anteriorly possess a thin, retractile tube, the proboscis. The mouth and brain are well developed, the *\*coelom* is indistinct. Most are marine, although freshwater and terrestrial forms occur. Most are not hermaphrodites. The first fossil forms occur in the Middle *\*Cambrian \*Burgess Shales* of British Columbia, Canada.

**Neoproterozoic** An *\*era* of the *\*Proterozoic* eon, 2500–541 Ma ago, preceded by the *\*Mesoproterozoic* and followed by the *\*Phanerozoic*. During this era, continental plates and *\*plate tectonics* appear, the supercontinent *\*Kenorland* formed, and there is evidence for oxygen-releasing *\*photosynthesis*.

**Neobothriocidaritis** See *BOTHRIOCIDAROIDA*.

**Neocomian** The earliest European *\*epoch* and substage of the Early *\*Cretaceous \*epoch*, preceded by the *\*Tithonian* and followed by the *\*Barremian \*stage*. It is dated at 145.5–130 Ma ago (Int. Commission on Stratigraphy, 2004) and comprises the *\*Berriasian*, *\*Valanginian*, and *\*Hauterivian* ages. Some authors restrict their use of Neocomian to just the Berriasian and Valanginian, whilst others extend it to include the Barremian and *\*Aptian*.

**Neogene** The middle of the three *\*periods* which comprise the *\*Cenozoic* era, preceded by the *\*Palaeogene* and followed by the Pleistocene, beginning 23.03 Ma ago and ending 1.81 Ma ago. The Neogene period is divided into the *\*Miocene* and *\*Pliocene* epochs.



<http://www.geologypage.com/2014/05/neogene-period.html>

- Neogene Period.

**Neohelikian** A sub-*\*stage* of the Upper *\*Helikian* of the Canadian Shield region.

**neoichnology** See *ICHTHOLOGY*.

**neomineralization** The formation of a new mineral or minerals from pre-existing minerals during *\*metamorphism*.

**neomorphism** The diagenetic replacement (see *DIAGENESIS*) of a mineral by a different crystal *\*form* of the same mineral. In *\*limestones* neomorphism often results in the replacement of *\*calcite* by coarser calcite crystals, e.g. *\*micrite* may become neomorphically replaced by *\*microspar* (4–10 µm crystals), or pseudospar (10–50 µm crystals). Neomorphic spar is characterized by irregularly shaped crystals, often patchily developed, and passing gradationally into unaltered areas.

**Neoproterozoic** The most recent part of the *\*Proterozoic*, from 1000–542 Ma ago.



<http://www.geologypage.com/2014/01/neoproterozoic-era.html>

- Neoproterozoic Era.

**neostratotype** A *\*stratotype* chosen to supersede the original stratotype (i.e. the *\*holostratotype*), if the first one has been demolished or invalidated. A neostratotype may be selected from outside the *\*type area*. Compare *LECTOSTRATOTYPE*. See also *HYPOSTRATOTYPE*; *PARASTRATOTYPE*.

**neotectonics** The study of the processes involved in the most recent *\*tectonism* and their effects. The term is usually applied to Late *\*Cenozoic* events. See also *SEISMOTECTONICS*.

**neoteny** Slowing down of bodily development, so that sexual maturity is achieved while the organism still looks like a juvenile; this leads to

**\*paedomorphosis.** Since the juvenile stages of many organisms are less specialized than the corresponding adult stages, such shifts allow the organisms concerned to switch to new evolutionary pathways. The word comes from the Greek *neos* (meaning ‘youthful’). Some features of human evolution have been ascribed to neoteny.

**Neotethys** See TETHYS.

**Neotropical faunal realm** Region which includes S. and Central America, including southern Mexico, the W. Indies, and the Galápagos Islands. Much of the area was isolated for the greater part of the **\*Palaeogene** and **\*Neogene** periods, which explains the distinctiveness of the fauna and the survival of ancient forms of mammals, e.g. the marsupials and edentates.

**neotype** Specimen chosen to act as the ‘type’ material subsequent to a published original description: this occurs in cases where the original types have been lost, or where they have been suppressed by the International Commission on Zoological Nomenclature (ICZN). Compare HOLOTYPE; LECTOTYPE; PARATYPE.

**NEP** See NEPHELOMETER.

**nephanalysis** Interpretation of cloud type and amount from satellite pictures in facsimile or digitized form.

**nepheline** An important member of the **\*feldspathoid** group of **\*silicate** minerals with composition  $\text{Na}_3(\text{Na,K})[\text{Al}_4\text{Si}_4\text{O}_{16}]$ , which approximates to  $\text{NaAlSiO}_4$  (with some K replacing Na); nepheline is an **\*end-member** of the series with **\*kalsilite** ( $\text{KAlSiO}_4$ ); sp. gr. 2.56–2.66; **\*hardness** 5.5–6.0; **\*hexagonal**; colourless, grey, yellowish, brownish, reddish, or greenish; **\*vitreous** to **\*greasy lustre**; **\*crystals** normally **\*prismatic**, short, and columnar, thick, and **\*tabular**, or coarse, granular aggregates; **\*cleavage** imperfect prismatic {1010}, poor basal {0001}; it is a primary crystallizing mineral of silica-poor **\*alkaline igneous** rocks in association with **\*aegirine** augite and alkali **\*amphiboles**, and is an **\*essential mineral** in some silica undersaturated rocks (see SILICA SATURATION) such as **\*nepheline-syenites** and **\*nephelinites**.

**nepheline-basanite** See BASANITE.

**nepheline-monzonite** An undersaturated \*monzonite (see SILICA SATURATION) containing essential \*nepheline (see ESSENTIAL MINERAL) in addition to the monzonite mineral assemblage.

**nepheline-syenite** A medium- to coarse-grained \*igneous rock consisting of essential \*alkali feldspar (see ESSENTIAL MINERAL), \*nepheline, \*pyroxene (\*aegirine-\*augite or sodic \*hedenbergite) and \*amphibole (ferrohastingsite,  $\text{NaCa}_2(\text{Fe}^{2+})_4\text{AlAl}_2\text{Di}_6\text{O}_{22}(\text{OH})_2$ , which is essentially a sodic, iron-rich \*hornblende, \*barkevikite and/or \*arfvedsonite) with accessory \*sphene, \*apatite, \*titanomagnetite, \*ilmenite, and \*zircon (see ACCESSORY MINERAL). The rock is, in effect, a \*syenite which is undersaturated in silica (see SILICA SATURATION).

**nephelinite** A fine-grained, often \*porphyritic, ultra-alkaline \*extrusive \*igneous rock consisting of essential \*titanaugite (see ESSENTIAL MINERAL), \*nepheline, and \*titanomagnetite, with accessory \*apatite, \*sphene, and \*perovskite (see ACCESSORY MINERAL). Where present, \*phenocrysts usually consist of the titanaugite, and occasionally nepheline. \*Sodalite and \*analcite may also be found in the \*groundmass. Where \*olivine is an additional component, the rock is termed an 'olivine nephelinite'. Nephelinites are essentially super-undersaturated \*basalts (see SILICA SATURATION) with nepheline taking the place of all the \*plagioclase feldspar.

**nepheloid layer** A body of water with a high concentration of suspended \*sediment which occurs near the deep ocean bottom, close to the base of the \*continental slope. Nepheloid layers are usually 1–300 m thick, and carry sediment up to 12  $\mu\text{m}$  in size, in concentrations of 0.3–0.01 mg/l. The sediment is suspended and transported by the movement of the oceanic \*thermohaline bottom currents. See CONTOUR CURRENT.

**nephelometer (NEP)** An instrument used in \*remote sensing that uses a transmitted \*laser beam to determine the scattering of light by atmospheric particles.

**nephrite** See JADEITE.

**Neptune** The eighth planet of the *\*solar system* (formerly the ninth when *\*Pluto*'s eccentric orbit carries it inside the orbit of Neptune, but Pluto is no longer ranked as a planet). Neptune is at 30.06 AU from the *\*Sun* and between  $4305.6 \times 10^6$  km and  $4687.3 \times 10^6$  km from Earth. Its equatorial radius is 24 766 km; polar radius 24 342 km; volume  $6254 \times 10^{10}$  km<sup>3</sup>; mass  $102.43 \times 10^{24}$  kg; mean density 1638 kg/m<sup>3</sup>; gravity 11 (Earth = 1); visual albedo 0.41; black-body temperature 33.2 K. Neptune has a dense atmosphere, with a surface pressure greater than 100 bars, composed of molecular hydrogen (89%), helium (11%), and traces of methane, with *\*aerosols* of ammonia ice, water ice, ammonia hydrosulphide, and possibly methane ice (similar to that of Uranus). The average surface temperature is about 58 K, and winds blow at up to 200 m/s. Neptune has 14 known satellites (see NEPTUNIAN SATELLITES).

**Neptune XIV** See HIPPOCAMP.


**neptunian dyke (sandstone dyke)** A sheet-like body of *\*sand* cutting through bedded *\*sediment* in a manner analogous to an *\*igneous \*dyke*. Neptunian dykes form by upward injection of liquefied sand through a fissure, often as a result of *\*earthquake* activity.

**neptunian satellites** See DESPINA (NEPTUNE V); GALATEA (NEPTUNE VI); HALIMEDE (NEPTUNE IX); HIPPOCAMP (NEPTUNE XIV); LAOMEDEIA (NEPTUNE XII); LARISSA (NEPTUNE VII); NAIAD (NEPTUNE III); NEREID (NEPTUNE II); NESO (NEPTUNE XIII); PROTEUS (NEPTUNE VIII); PSAMATHE (NEPTUNE X); SAO (NEPTUNE XI); THALASSA (NEPTUNE IV); TRITON (NEPTUNE I).

**neptunism** A theory of the formation of the Earth, popular at the end of the 18th and the beginning of the 19th century, and associated with *\*Werner*. Werner taught that the oldest, primitive rocks, including *\*granite* and *\*basalt*, had crystallized from a primordial ocean. They were followed by Transition and Floetz formations, also precipitated, and then by Alluvia and Volcanic rocks.

**Nereid (Neptune II)** A satellite of *\*Neptune*, with a diameter of 340 km; visual albedo 0.16, and an orbital period of 360.14 days.

**Nereites** A bilobed \*trail that winds over a \*bedding plane. The trail has a transverse ornament and was formed by a \*deposit feeder during its methodical search for food. The term 'Nereites facies' is given to deep-water deposits characterized by Nereites and depth-related genera.



**Nereites**

**Nereus** A \*solar system asteroid (No. 4660), diameter 2 km; orbital period 1.82 years. It is a near-Earth asteroid, scheduled to be visited by a sample-return mission.

**neritic province** See **NERITIC ZONE**.

**neritic zone (neritic province)** The shallow-water, or near-shore, marine zone extending from the low-tide level to a depth of 200 m. This zone covers about 8% of the total ocean floor. It is the area most populated by \*benthic organisms, due to the penetration of sunlight to these shallow depths.

**Neso (Neptune XIII)** A satellite of Neptune with a radius of 30 km and a retrograde orbit with a period of 9374 days.

**nesosilicates (orthosilicates)** A group of \*silicate minerals characterized by independent  $\text{SiO}_4$  \*tetrahedra with no shared oxygens. The structure is held together by bonding with interstitial cations. The group includes \*olivine, \*garnet, \*sphene, \*zircon, \*staurolite, \*chloritoid, \*topaz, \*chondrodite, and the  $\text{Al}_2\text{SiO}_5$  \*polymorphs.

**net flux radiometer** A \*radiometer carried by satellites that is used to measure the energy absorbed and emitted by the atmosphere of a planet.

**net slip** See SLIP.

**network-former** Aluminium (Al), silicon (Si), and often phosphorus (P) atoms, which tend to form network structures in **\*silicate \*melts**.

**network-modifier** Large **\*cations** (e.g.  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Fe}^{2+}$ ,  $\text{Na}^+$ , and  $\text{K}^+$ ) occurring as major constituents in **\*magmas**, which usually disrupt and modify the network structures.

**neural arch** See VERTEBRA.

**neural spine** See VERTEBRA.

**neutral fold** A **\*fold** which closes laterally and is therefore neither **\*antiformal** nor **\*synformal**. Where the **\*fold axis** and **\*axial plane** are inclined vertically, a neutral fold is known as a 'vertical fold'.

**neutrally buoyant float (Swallow buoy)** A device developed by J. C. Swallow in 1955 for measuring current speed and direction at depth in the oceans. It consists of two aluminium tubes, one containing batteries and sound-generating circuitry, the other containing adjustable weights to allow the device to float and remain at any desired depth. It emits a series of sound pulses which can be tracked by ship, allowing water motions to be measured at selected depths.

**neutral soil** Soil with a **\*pH** value of 6.6–7.3.

**neutron-activation analysis** The analysis of a material by bombarding a sample with fast-moving neutrons in the core of a nuclear reactor. Neutrons are added to the nuclei of stable **\*isotopes** to form new radionuclides which decay, producing particles with characteristic energies that can be measured with a **\*scintillation counter**.

**neutron–gamma sonde** An instrument package for measuring the amount of hydrogen present within the rocks surrounding a **\*borehole**. The **\*sonde** contains a neutron source and a gamma-ray detector. The gamma rays induced in the rocks by neutron bombardment are proportional to the amount of hydrogen present. The device can be used to determine the **\*porosity** of the surrounding rocks.

**neutron log** The recording of neutron bombardment by a **\*neutron–neutron sonde** being raised through a **\*borehole**. The neutron log is used to estimate

formation **\*porosity** and is presented on a scale calibrated in limestone porosity units.

**neutron moisture meter (neutron soil-moisture probe)** An instrument which uses high-energy (fast) neutrons to detect indirectly the water content of a soil. The fast neutrons are moderated to slow neutrons by the hydrogen atoms in the soil moisture. By measuring the density of the slow-neutron backscatter, an indication is obtained of how much water is present.

**neutron–neutron sonde** A radioactive instrumental package for lowering down a **\*borehole**. It comprises a neutron source and detector. The number of neutrons back-scattered to the detector is proportional to the number of hydrogen atoms within the rocks surrounding the borehole. It can be used to determine the **\*porosity** of the surrounding rocks. *See also* **NEUTRON–GAMMA SONDE**; **NEUTRON LOG**.

**neutron soil-moisture probe** *See* **NEUTRON MOISTURE METER**.

**névé** *See* **FIRN**.

**Newer Drift** The deposits marking the maximum extent of the last (**\*Devensian**) glaciation. A morphological distinction occurs between the more-weathered **\*drift** to the south and the less-weathered drift to the north, in the British Isles. The boundaries between the less-weathered Newer Drift and the **\*Older Drift** have been shown to be less distinct than was once supposed.

**New Horizons** A **\*NASA** mission to **\*Pluto** and the **\*Kuiper Belt** that launched on 19 January 2006, flew past Pluto on 14 July 2015, and by September 2018 was entering the Kuiper Belt.



<http://pluto.jhuapl.edu>

- New Horizons: NASA's Mission to Pluto and the Kuiper Belt.

**New Millennium Deep Space-1** A successful **\*NASA** mission, launched in 1998 and retired in 2001, that tested new technologies and encountered comet Borrelly.



<http://nssdc.gsfc.nasa.gov/nmc/spacecraftDisplay.do?id=1998-061A>

- A NASA probe designed to test new technology.

**New Millennium Deep Space-2** A \*NASA mission to \*Mars, launched in 1999. Communication with the probe was lost.

**New Millennium Program (NMP)** A \*NASA programme involving twelve spacecraft that would be used to test and demonstrate new technologies for space exploration. Technologies relevant to planetary missions were called Deep Space, renamed Space Technology in 2000, and those for tasks in Earth orbit were called Earth Observing. Two Deep Space, two Space Technology, and one Earth Observing satellites were launched, but Congress cancelled funding for the programme in 2009, and seven launches were abandoned.

**New Red Sandstone** Name given to the terrestrial \*red-bed deposits that overlie the \*Carboniferous. These deposits are of Permo–Triassic age.

**newton (N)** The derived SI unit of force, named after Sir Isaac Newton (1642–1727), being the force required to produce an acceleration of  $1 \text{ m/s}^2$  in a mass of 1 kg;  $1 \text{ N} = 1 \text{ J/m}$ .

**Newtonian behaviour** The manner in which a fluid flows if it exhibits ideal viscous \*strain. The application of a deforming stress induces permanent strain from which there is no recovery when the stress is removed. *Compare* ELASTIC DEFORMATION.

**Newtonian fluid** *See* BINGHAM FLUID.

**Newton's law of gravitation** *See* GRAVITATIONAL ACCELERATION.

**NEXRAD** *See* NEXT GENERATION WEATHER RADAR.

**Next Generation Weather Radar (NEXRAD)** A network of 175 \*Doppler radar installations, completed in 1996, that cover the whole of the USA. They are located at weather stations, airports, and military bases. Each is mounted on a tower and provides three-dimensional images of weather conditions with clear resolution to 200 km and poorer resolution to 320 km.

**Ngaterian** A New Zealand \*stage (99.6–95 Ma ago) of the Late \*Cretaceous \*period, preceded by the \*Motuan and followed by the \*Arowhanan.

**Niagaran** A *\*stage* (438–421.3 Ma ago) of the *\*Silurian* of N. America, preceded by the *\*Medinian* and followed by the *\*Cayugan*. It is equivalent to the Upper *\*Llandovery* to Lower *\*Ludlow* period.

**niccolite** Metallic mineral, NiAs; sp. gr. 7.8; *\*hardness* 5.0; greyish-red; usually *\*massive* or disseminated; occurs as irregular *\*aggregates* and as complex intergrowths with other *\*sulphides*, associated with *\*basic igneous* rocks.

**Nicol, William** (1768–1851) A mineralogist from Edinburgh, Nicol invented the calcite prism named after him and used in polarizing microscopes (see **NICOL PRISM**). His description of a microscope based on his prisms was published in 1829, although the instrument was in use much earlier.

**nicol prism** Two pieces of optically clear *\*calcite*, cemented by *\*Canada balsam* into the shape of a prism. Light entering the base of the prism is doubly refracted, and when both rays reach the Canada-balsam cement one ray is reflected away from the prism while the other ray continues through the prism. Thus the light emerging from the prism is *\*plane polarized*. These prisms were invented by **William Nicol**. Early *\*polarizing microscopes* were fitted with nicol prisms for the *\*analyser* and *\*polarizer*, but modern microscopes are fitted with *\*Polaroid* instead, and, although the term ‘crossed nicols’ is still in common use, this has been largely replaced by the term ‘*\*crossed polars*’.

**NigeriaSat-2** A Nigerian *\*minisatellite* mission that provides high-resolution surface imagery along a 20-km swathe. It was launched on 17 August 2011, from Yasný Cosmodrome, Russia, into a Sun-synchronous circular orbit at an altitude of 700 km.

**NigeriaSat-X** A Nigerian *\*microsatellite* mission that provides high-resolution surface imagery. It was launched on 17 August 2011, from the Yasný Cosmodrome, Russia, into a Sun-synchronous near-circular orbit at an altitude of 663 km.

**Niggli, Paul** (1888–1953) A Swiss crystallographer and petrologist at the University of Zurich, Niggli studied *\*magmas* and *\*metamorphic rocks*, and introduced the use of *\*phase diagrams* in *\*petrology*. He also made a classification of *\*sedimentary rocks* according to grain shape and, in 1928,

he and his students attempted to calculate the bulk chemical composition of the Earth. See [NIGGLI METHOD](#).

**Niggli method** The grouping of the oxide components of an *\*igneous* rock according to their chemical similarity: ( $\text{Al}_2\text{O}_3 + \text{Cr}_2\text{O}_3 + \text{*REEs}$ ); ( $\text{FeO} + \text{MnO} + \text{MgO}$ ); ( $\text{Na}_2\text{O} + \text{K}_2\text{O} + \text{Li}_2\text{O}$ ); and ( $\text{CaO} + \text{BaO} + \text{SrO}$ ). The oxide groupings are calculated according to *\*cation* percentage, the sum for the rock being normalized to 100. Each group corresponds to an individual Niggli value, each of which can be plotted against silica for a comparison of associated rocks. The method was devised by P. *\*Niggli*.

**nimbostratus** From the Latin *nimbus*, 'rain', and *stratus*, 'spread out', a dark or grey cloud that obscures the Sun, associated with more or less continuous rainfall which makes the cloud base diffuse. See also [CLOUD CLASSIFICATION](#).

**NIMS** See [NEAR-INFRARED MAPPING SPECTROMETER](#).

**niobite** See [COLUMBITE](#).

**nitisols** A reference soil group in the *\*World Reference Base for Soil Resources* classification scheme. Nitisols have a nitric B horizon (see [NITRIC HORIZON](#)) more than 30 cm below the surface, with a *\*cation-exchange capacity* of less than 36  $\text{cmol}_c/\text{kg}$ . No evidence of clay *\*lessivage* is evident in *\*thin sections* taken within 100 cm of the surface.

**nitratine** See [SODA NITRE](#).

**nitre (saltpetre)** Mineral,  $\text{KNO}_3$ ; sp. gr. 1.9–2.3; *\*hardness* 2; *\*orthorhombic*; white, grey, reddish-brown, or lemon-yellow; *\*vitreous* *\*lustre*; crystals *\*acicular*, usually forms granular crusts and uneven masses; occurs in very arid conditions where vegetation is very sparse, in association with *\*evaporite* minerals of the arid desert type (e.g. *\*soda nitre*, *\*gypsum*, *\*halite*, and occasionally iodates), and little precipitation is required to wash an encrustation of nitre into hollows where it forms *\*massive* deposits. It is used as a fertilizer.

**nitric horizon** A *\*soil horizon* that contains more than 30% *\*clay* consisting of 1:1 *\*clay minerals*.

**nitrogen fixation** See [FIXATION \(2\)](#).

**nival** Applied to the geomorphological (see [GEOMORPHOLOGY](#)) processes that result from the action of snow.

**nivation** Complex of surface erosional processes acting under a snow cover. It includes [\\*gelifraction](#), and the removal of shattered debris by [\\*solifluction](#) and the movement of melted snow. It is an initial process in [\\*cirque](#) development.

**NMO** See [MOVEOUT \(2\)](#).

**NMP** See [NEW MILLENIUM PROGRAM](#).

**NOAA-20** See [JOINT POLAR SATELLITE SYSTEM-1](#).

**NOAA-20/JPSS-1** See [JOINT POLAR SATELLITE SYSTEM-1](#).

**NOAA-class satellite** See [TELEVISION AND INFRARED OBSERVATION SATELLITE](#).

**Noachian** A division of [\\*areological](#) time, lasting from 4.60 Gy to 3.50 Gy in the Hartmann–Tanaka Model and 4.60 Gy to 3.80 Gy in the Neukum–Wise Model, and divided into three epochs: Lower Noachian (4.60–3.92 or 4.60–4.50 Gy); Middle Noachian (3.92–3.85 or 4.50–4.30 Gy); and Upper Noachian (3.85–3.50 or 4.30–3.80 Gy).

**NOAA POES Series** A fifth-generation series of [\\*polar](#) orbiting environmental satellites that provide an uninterrupted flow of global environmental information. The satellites were launched on 13 May 1998 (NOAA-15), 21 September 2000 (NOAA-16), 24 June 2002 (NOAA-127), 20 May 2005 (NOAA-18), and 6 February 2009 (NOAA-19).

**noctilucent clouds (luminous night clouds)** Type of clouds, occurring at 80–85 km altitude, near the upper limit of the [\\*stratosphere](#), characterized by a blue to yellow hue, and similar in appearance to [\\*cirrostratus](#). The clouds are seen on summer nights at latitudes between about 50° and 65° in both northern and southern hemispheres; they move rapidly, at speeds up to 300 knots (555 km/h), often in a wave formation.

**nocturnal radiation** The long-wave radiation from the surface of the Earth at night that is in excess of the incoming radiation from the atmosphere. See

also ATMOSPHERIC 'WINDOW'; RADIATION BUDGET; TERRESTRIAL RADIATION.

**node** 1. A point of zero displacement in a material transmitting *\*standing waves*, produced by destructive *\*interference* between waves propagating in opposite directions. 2. In a *\*phylogenetic tree*, a representation of an extant (terminal node) or ancestral (internal node) operational taxonomic unit.

**nodes** The points, diametrically opposite, at which the orbit of a planet intersects the plane of the *\*ecliptic*, or at which the orbit of a *\*satellite* intersects the orbital plane of a planet.

**nodular (concretionary)** Applied to the *\*habit* of a mineral when its outer surface is rounded or spherical in shape. It may be the result of progressive chemical precipitation, giving a concentric appearance. Minerals such as iron or *\*manganese nodules* form in this way.

**nodule** A spherical or oval concretion.

**Noginskian** A Russian *\*stage* (300.5–299 Ma ago) and substage of the *\*Gzhelian* stage (Late *\*Pennsylvanian* sub-epoch), preceded by the *\*Klazminskian* and followed by the *\*Asselian* (*\*Rotliegendes* epoch of the *\*Permian*).

**Noguerornis** The earliest known bird after *\*Archaeopteryx*, from the basal *\*Cretaceous* of Montsec, northern Spain. It was the size of a finch and was the first bird to have a wing that was well developed, as indicated by the elongation of the distal portions of the forelimb and the rigid interlocking of the hand bones.

**noise** A signal that conveys no useful information (e.g. background sound that makes it difficult to hear a conversation). If the useful signal comprises data that are being recorded, random (white) noise can be reduced by summing the recorded signals; incoherent noise is effectively damped out and the coherent signal is enhanced, thus improving the signal-to-noise ratio.

**nomogenesis** See ARISTOGENESIS.

**non-conformity (heterolithic unconformity)** An *\*unconformity* in which younger, *\*sedimentary rocks* rest unconformably on older, *\*igneous* or *\*metamorphic rocks*.

**non-dipole field** The field remaining after removal of the *\*dipole field* (usually the inclined dipole) from the observed *\*geomagnetic* or planetary magnetic-dipole field.

**non-frontal depression** Low-pressure system that does not develop from a frontal wave as do typical mid-latitude frontal cyclones (or *\*depressions*). Most tropical cyclones are non-frontal. Various conditions can lead to the formation of depressions without frontal characteristics. *See also* COLD LOW; LEE DEPRESSION; POLAR DEPRESSION; THERMAL LOW.

**non-metallic** Applied to the *\*lustre* of a *\*mineral* which reflects light from its surface but does not shine like a metal. 'Non-metallic' may be further qualified as 'glassy' or 'vitreous', 'silky', 'resinous', etc.

**non-polarizable electrode** An electrode whose potential is not affected by the current passing through it, e.g. the porous-pot electrode extensively used in *\*spontaneous potential* measurements which comprises a copper rod immersed in copper sulphate solution that makes ionic contact with the ground through the porous base of the electrolyte reservoir.

**non-recoverable strain** *See* PERMANENT STRAIN.

**non-renewable resource (finite resource)** Resource that is concentrated or formed at a rate very much slower than its rate of consumption and thus, for all practical purposes, is nonrenewable. *Compare* RENEWABLE RESOURCE.

**non-rotational strain** *See* PURE SHEAR.

**non-selective scattering** The *\*scattering* of all wavelengths of *\*electromagnetic radiation* equally in the atmosphere, usually caused by particles which are much larger than the energy wavelengths.

**non-sequence** A minor break in a *\*concordant* succession of *\*strata*, representing a period during which either no deposition of *\*sediment* occurred or it was subsequently removed. Such periods may be of short duration and localized, but are not necessarily so. *Compare* DIASTEM.

**non-spectral hue** A *\*hue* which is not present in the spectrum of colours produced by splitting white light with a prism. Non-spectral hues include brown and the pastel colours. *Compare* SPECTRAL HUE.

**non-steady flow** See STEADY FLOW.

**non-strophic** See HINGE.

**non-umbilicate** See IMPERFORATE.

**non-uniform flow** See UNIFORM FLOW.

**norbergite** See CHONDRODITE.

**Nor'easter** A storm that occurs along the eastern coast of N. America, most frequently and violently between September and April, bringing winds predominantly from the north-east.



<https://www.weather.gov/safety/winter-noreaster>

- What is a Nor'Easter?

**Norian (Juvavic)** A Late \*Triassic \*stage, preceded by the \*Carnian and followed by the \*Rhaetian, and dated at 216.5–203.6 Ma ago (Int. Commission on Stratigraphy, 2004). It is roughly contemporaneous with the Houbachong (China) and Warepan (New Zealand).

**norite** A coarse-grained, \*basic \*igneous rock consisting of essential \*plagioclase feldspar, \*orthopyroxene (\*hypersthene or \*bronzite), and \*clinopyroxene (\*augite), with accessory \*ilmenite (see ACCESSORY MINERAL; ESSENTIAL MINERAL). Orthopyroxene is dominant over clinopyroxene, and the plagioclase is a calcic type (labradorite or bytownite). Norites, like \*gabbros, are found as layers in many large, layered basic intrusions (see INTRUSIVE), as well as forming intrusions in their own right.

**norm** See CIPW NORM CALCULATION.

**normal distribution (Gaussian distribution)** In statistics, a continuous \*probability distribution which is asymptotic and symmetrically bell-shaped about the \*mean. The normal distribution is widely applied in statistics to model continuous variation. The two parameters of the distribution are the mean and \*variance. See also CENTRAL LIMIT THEOREM.

**normal fault** A high-angle (more than 50°), **dip-slip fault** on which displacement of the hanging wall is downwards, relative to the **footwall**. Normal faults commonly occur in **conjugate fault sets** which cause discrete block subsidence and uplift in the form of **grabens** and **horsts**.

**normal field** A magnetic field, usually geomagnetic, that has the same **polarity** as the present field, i.e. the north magnetic pole either lies in the northern hemisphere or is clearly a continuation of the north-pole **polar wander path**.

**normal incidence** The condition in which a wavefront is parallel to an interface, such that the ray path is perpendicular (normal) to the surface. The **angle of incidence** is zero. *See also* SNELL'S LAW.

**normalized vegetation index** *See* VEGETATION INDEX.

**normally consolidated clay** Clay that is compacted by exactly the amount to be expected from the pressure exerted by the **overburden**; clay which has never been overloaded. *Compare* OVER-CONSOLIDATED CLAY.

**normal moveout** *See* MOVEOUT.

**normal problem** *See* FORWARD PROBLEM.

**normal stress** ( $\sigma$ ) The **stress** which acts perpendicularly to the plane to which a force has been applied. Normal **compressive stresses** (with positive values) inhibit sliding along a plane; normal tensile stresses (with negative values) separate rocks along a plane.

**normal travel time** The **time–distance** graph compiled from **first arrivals**, but only where there is no anomalous geologic structure. It is used to calibrate travel times for a given **offset** range in **fan-shooting** refraction surveys.

**normal twins** In **crystallography**, a twinned **crystal** where the twin **axis of rotation** is at right angles to the **crystal face**.

**normal zoning** *See* CRYSTAL ZONING.

**normative constituents** *See* CIPW NORM CALCULATION.

**normative mineral** A chemical formula that represents the ideal **mineral** composition of a rock, with each constituent expressed as the percentage by

weight (wt. %). This is used in the normative classification of rocks (see [CIPW NORM CALCULATION](#)). The minerals most commonly used in normative classification are: [\\*quartz](#) (Q); [\\*corundum](#) (C); orthoclase (Or, see [ALKALI FELDSPAR](#)); albite (Ab, see [ALKALI FELDSPAR](#); [PLAGIOCLASE FELDSPAR](#)); anorthite (An, see [ALKALI FELDSPAR](#)); [\\*nepheline](#) (Ne); [\\*wollastonite](#) (Wo); [\\*enstatite](#) (En); ferrosilite (Fs, see [ORTHOPYROXENE](#)); forsterite (Fo, see [OLIVINE](#)); fayalite (Fa, see [OLIVINE](#)); [\\*ilmenite](#) (Il); [\\*magnetite](#) (Mt); and [\\*apatite](#) (Ap).

**NorSat-1 and -2** A [\\*microsatellite](#) mission by the Norwegian Space Centre to investigate solar radiation and space weather and (NorSat-2) to develop new methods for detecting and managing ship traffic. The satellites were launched on 14 July 2017, from Baikonur, Kazakhstan, into orbit at 600 km.

**norte (papagayo)** Cold, northerly, local wind affecting the coasts of the Gulf of Mexico, most commonly in winter. Sometimes it brings rainfall.

**North Alpine foreland basin (Molasse Basin)** A [\\*foreland basin](#) that extends for about 1000 km along the long axis of the Alps and to their north, from Lake Geneva to lower Austria. It formed during the [\\*Oligocene](#) and [\\*Miocene](#) through the [\\*flexure](#) of the [\\*Eurasian Plate](#) in response to the weight of the Alps during the [\\*Alpine–Himalayan orogeny](#).

**North American Plate** One of the present-day major lithospheric [\\*plates](#), the N. American Plate extends from the [\\*Mid-Atlantic Ridge](#) in the east to a complex pattern of [\\*subduction zones](#) and [\\*transform faults](#) which form its boundaries with the [\\*Pacific](#), [\\*Juan de Fuca](#), [\\*Gorda](#), [\\*Cocos](#), [\\*Caribbean](#), and [\\*South American Plates](#) in the west and south. Along the western side, the N. American Cordillera has been interpreted as a collage of accreted [\\*allochthonous](#) [\\*terranes](#).

**North Atlantic deep water (NADW)** A water mass (salinity 34.9–35.03%, temperature 1.0–2.5 °C) that was originally believed to form in an area off the southern tip of Greenland, where winter cooling of saline waters was thought to cause a body of water to sink and spread south. It is now recognized that the main source is in the Norwegian Sea, from which deep water flows over the sills between Scotland, Iceland, and Greenland and cascades into the depths of the Atlantic.

**North Atlantic Drift** Oceanic surface current in the N. Atlantic, flowing from the Grand Banks off Newfoundland eastwards to north-western Europe and forming a northerly extension of the **\*Gulf Stream**. This diffuse, shallow and relatively warm current has an ameliorating influence on the climate of the coastal regions of north-western Europe.

**norther** A strong, cold, northerly wind that blows in winter across the south-eastern USA, sometimes extending across Mexico and into the Pacific.

**'northern lights'** See **AURORA**.

**Northgrippian** The second **\*age**/**\*stage** of the **\*Holocene**, 8326–4250 BP, dated from the NorthGRIP1 ice core (see **GRIP**); Global Auxiliary **\*Stratotype**, Gruta do Padre Cave speleothem, Brazil.

**North Pacific current** Oceanic surface current in the N. **\*Pacific** that flows eastwards, as an extension of the **\*Kuroshio current**, towards California. It occupies a position in the N. Pacific similar to that in the **\*Atlantic** of the system comprising the **\*Gulf Stream** and **\*North Atlantic Drift**.

**nor'wester** **1.** Convective storm, usually from the north-west, which affects Assam and Bengal between March and May, prior to the start of the south-westerly **\*monsoon**. It is characterized by violent conditions, including **\*line squalls**. **2.** A hot wind blowing from the mountains in South Island, New Zealand. **3.** In S. Africa, a **\*depression** associated with an active weather **\*front**.

**nosean** A **\*feldspathoid** mineral and member of the **\*sodalite** group  $\text{Na}_8[\text{Al}_6\text{Si}_6\text{O}_{24}]\text{SO}_4$ , where it is closely related to **\*häüyne**; sp. gr. 2.3–2.4; **\*hardness** 5.5–6.0; **\*cubic**; grey, white, or greenish, often with light-blue tinges; **\*vitreous** **\*lustre**; **\*crystals**, when present, **\*rhombohedra**, but it is also **\*massive**; **\*cleavage** poor, rhombododecahedral; occurs in extrusive **\*alkaline** undersaturated (see **SILICA SATURATION**) **\*igneous** rocks such as nosean leucitophyres and **\*phonolites**.

**Nothosauria (nothosaurs)** (order **\*Sauropterygia**) Suborder of marine **\*reptiles** that flourished during the **\*Triassic**. They possessed long necks, and their limbs were adapted to swimming. The nothosaurs were replaced by the **\*plesiosaurs** in the Early **\*Jurassic**.

**nothosaurs** See NOTHOSAURIA.

**notochord (chorda dorsalis)** Somewhat flexible, rod-like structure, composed of disc-like, turgid cells, which extends virtually the entire length of the body of adult and/or larval members of the phylum **\*Chordata**. Lying below the nerve cord, but **\*dorsal** to the intestine, the notochord provides a form of flexible support to the body. In **\*vertebrates** the notochord is replaced wholly or partly by the vertebral column, but it is retained throughout life in **\*Cephalochordata** and **\*Agnatha**.

**notothyrium** A triangular opening in the posterior margin of the brachial valve of a brachiopod (**\*Brachiopoda**) shell, through which the **\*pedicle** may partly emerge. In some species it is open, in others it is closed by a single plate of shell material, the chilidium, or by two chilidial plates.

**nova** A new star.

**novaculite** A dense, fine-grained to **\*cryptocrystalline** **\*chert**.

**NovaSAR-S** A UK **\*minisatellite** mission to monitor floods, agricultural crops, and forests and to map land use, assist with disaster management, detect shipping, monitor oil spills, and help with maritime safety. It was launched successfully on 16 September 2018, from the Satish Dhawan Space Centre, India, into a Sun-synchronous orbit at an altitude of about 580 km.

**NRM** See NATURAL REMANENT MAGNETISM.

**nSight-1** A 2-unit **\*Cubesat** mission comprising a constellation initially of fifty CubeSats (called QB50) developed commercially in South Africa; nSight-1 carries instruments to study the lower **\*thermosphere**. The first twenty-eight satellites were deployed in batches from the International Space Station in 2017 into near-circular orbit at an altitude of about 400 km. The second batch, of eight CubeSats, will be placed in Sun-synchronous orbit at 500 km.

**nuclear-magnetic log** The combined outputs of radioactive and magnetic **\*sondes** used in **\*well logging**.

**nuclear-precession magnetometer** A **\*magnetometer** based on the ability of protons to precess about the **\*geomagnetic field** with a frequency dependent on the strength of the ambient magnetic field. A strong field is

applied which aligns the protons, and they then precess about the geomagnetic field after the applied field is removed. The instrument has a sensitivity of the order of 1 nT, and is capable of rapid repetition to produce an almost continuous signal.

**nuclear waste** See RADIOACTIVE WASTE.

**nucleation** 1. Formation of an embryonic \*crystallite from a \*melt which is followed by the growth of a nucleus to crystal dimensions. See also MICROLITE. 2. Theory according to which \*geosynclines developed on the edges of \*cratons; the ensuing \*orogenic belt then became part of the craton and the products of subsequent erosion filled a new geosynclinal trough which developed on the edge of the enlarged craton. The theory of \*plate tectonics, with the emphasis on horizontal movement, modified nucleation in many ways, not least by including oceanic sediment, scraped off the subducting plate (see SUBDUCTION), into the new craton. See ACCRETION.

**nucleic acids** Nucleotide polymers, with high relative molecular mass, produced by living cells and found both in the \*nucleus and cytoplasm. They occur in two forms, designated DNA and RNA, and may be double- or single-stranded. DNA embodies the genetic code of a cell or organelle, while various forms of RNA function in the transcriptional and translational aspects of protein synthesis.

**nucleosynthesis** The process by which elements are formed. Modern theories suggest that nucleosynthesis is intimately linked with the stages in the life-cycle of stars (stellar evolution), and that, commencing with hydrogen, heavier elements are created by nuclear fusion of lighter \*nuclides at the temperatures and pressures existing in the cores of stars. Because the lighter elements are consumed to produce energy these \*thermonuclear reactions are referred to as 'burning', although they have nothing to do with combustion. The stages of stellar evolution conform well with the overall pattern of peaks and troughs in the \*cosmic abundance of elements in order of increasing atomic number (Z). During the first and longest (main-sequence) phase, hydrogen (which is by far the most abundant element of the stellar material) is consumed to produce helium (\*hydrogen burning). Hydrogen burning is followed in turn by \*helium burning, \*carbon and \*oxygen burning, and \*silicon burning, each phase producing heavier elements from lighter ones. The heaviest elements are

formed in the last stages in the sequence: the **\*equilibrium (e) process** (coinciding with the 'iron peak' elements, Cr, Mn, Fe, Co, and Ni), followed by the '**\*slow-neutron (s) process** producing elements up to Bi (atomic number 83), and finally (in supernova events) the **\*rapid-neutron (r) process** producing elements with atomic number greater than 83. *See also* **PROTOSTAR**.

**nucleus** (*pl.* nuclei) **1.** The centre of an atom, composed of protons and neutrons and accounting for nearly all of its mass. A proton has a positive electrical charge, equal in magnitude to the negative charge of an **\*electron**; a neutron carries no electrical charge. The nucleus of the hydrogen atom contains a single proton; uranium, the heaviest naturally occurring element, has 92 protons and 142, 143, and 146 neutrons in **\*isotopes** 234, 235, and 238 respectively. **2.** A small, solid particle, e.g. of dust, salt, or smoke on to which water vapour will condense. Such particles are called 'condensation nuclei' and some of them have hygroscopic properties that encourage condensation in unsaturated air. Other nuclei of a suitable shape, e.g. some clay particles such as **\*kaolinite**, probably act as 'freezing nuclei' in the initial stage of ice-crystal formation. *See also* **AITKEN NUCLEUS**; **BERGERON THEORY**; **CONDENSATION NUCLEUS**; **ICE NUCLEUS**. **3.** The double-membrane-bound organelle containing the **\*chromosomes**, that is found in most non-dividing **\*eukaryotic** cells; it is essential to their long-term survival. It is variously shaped, although it is normally spherical or ovoid. It disappears temporarily during cell division. It is absent from viruses. The **\*chromosomes**, though probably intact, are not visible when the cell is in a resting stage (i.e. not dividing). The nucleus also contains nucleoli, small spherical dense bodies made up of ribosomal RNA and protein, which gives it its integrity. **4.** *See* **NUCLEATION**.

**nucleus number** The number of **\*crystal** nuclei formed in a unit volume of **\*magma** in unit time, during the cooling of the magma body.

**nuclide** A widely used alternative name for an atom. The composition of any nuclide can be given by means of the chemical symbol of the element, the **\*mass number** written as a superscript, and the **\*atomic number** written as a subscript. For example,  $^{14}_6\text{C}$  is an atom of carbon (C) having six protons (the atomic number) and 14 nucleons (the mass number).

**nuée ardente** A French phrase, meaning ‘glowing cloud’, first used in 1903 by Alfred Lacroix (1863–1948) to describe the complete phenomenon of a basal **\*pyroclastic flow** and its overriding **\*ash cloud** formed by the collapse of a volcanic **\*dome**. Since the pyroclastic flow and ash cloud behave differently from one another during transport and produce contrasting deposits, volcanologists now tend to avoid using the term.

**Nukumaruan** A New Zealand **\*stage** (1.96–1.1 Ma ago) of the **\*Neogene \*period**, underlain by the **\*Mangapanian**, overlain by the **\*Castlecliffian**, and roughly contemporaneous with most of the Calabrian stage.

**Nullaginian** A **\*stage** (2500–1800 Ma ago) of the Lower **\*Palaeoproterozoic** of south-eastern Australia, preceded by the **\*Neoarchaeon \*era** and overlain by the **\*Carpentarian**.

**null hypothesis** See HYPOTHESIS.

**null point 1.** In nearshore, shoaling waters, each particle-size of **\*sediment** should hypothetically have a null point, where there is no net movement of the particle landward or seaward. It is the point where there is a balance between the component of gravity acting down the slope in a seaward direction and the force on the particle resulting from the difference between the crest and trough velocities of the waves which tends to move the particle landwards. **2.** In an **\*estuary**, the point at which the residual sea water landward flow is balanced by the seaward residual river flow.

**numerical aperture** See OBJECTIVE.

**numerical taxonomy** The classification of organisms by purely mathematical means. It is based on quantifying observable characteristics of organisms and may be operated at various taxonomic levels to deal with species or higher taxa. It involves the grouping and computation of the similarity of characters; the results are usually displayed graphically, as a **\*phenogram** or dendrogram. There has been discussion as to whether classifications produced in this way are valid and reflect **\*phylogeny**. See TAXON; TAXONOMY.

**Nummulites** A genus of larger **\*Foraminiferida**, which have a flattened, discoidal or lenticular **\*test**. They lived in warm, shallow, marine waters from the top of the **\*Palaeocene** to the Upper **\*Oligocene**, and in some areas they are numerous enough to be major rock formers. Many species have

been described; they occur at different stratigraphic levels and in different **\*facies associations**. *Nummulites planulatus*, which is Cuisian in age, is often found in association with alveolinids and miliolids in a sandy facies; *N. globulus* occurs in the Ilerdian of the Pyrenees in marly and sandy facies; *N. gizahensis* is one of the largest forms and is usually found in sub-reefal **\*limestones** of the Upper **\*Lutetian** of the **\*Tethyan realm**; *N. variolarius* preferred shallow, calm environments and is found with miliolids in sandy limestones of the Middle and Upper Lutetian.

**Nuna (Columbia)** The earliest known **\*supercontinent**, which is believed to have formed about 1600 Ma ago and to have broken up about 1400 Ma ago.

**nunatak** Rocky summit or mountain range that stands above a surrounding **\*ice sheet** in an area that currently is being glaciated.


**Nunivak** A normal **\*polarity subchron** which occurs within the **\*Gilbert** reversed **\*chron**.

**Nusselt number** A dimensionless number relating to the **\*Rayleigh number**. It corresponds to 1 for initial convective motion and increases with an increased probability of convective motions occurring.

**nutaton** Irregularities in the orbital motion of the Earth (or other planets) superimposed on the **\*precession** of its axis. The dominant nutation of the Earth's axis has an 18.6-year periodicity.

**nutrient cycle** See **BIOGEOCHEMICAL CYCLE**.

**Nyquist frequency (folding frequency,  $f_N$ )** A frequency which is half of the **\*sampling frequency**. The Nyquist interval is the frequency range from zero to  $f_N$ , where  $f_N = 1/(2\Delta t)$  and  $\Delta t$  is the **\*sampling interval**. For example, if the sampling interval  $\Delta t$  is 2 ms (i.e. a sampling frequency of 500 Hz), then  $f_N = 1/(2 \times 0.002) = 250$  Hz. It is important to know the value of the Nyquist frequency, as frequencies greater than  $f_N$  will alias (or 'fold back') and appear as lower frequencies from which they cannot be distinguished.



**oasis** A fertile spot in the desert, the basis of which is a supply of water that is available throughout the year and which normally originates as **\*groundwater**.

**obduction** The lateral, subhorizontal displacement of a lithospheric **\*plate** on to a **\*continental margin** at a **\*destructive** plate boundary. It is the opposite of **\*subduction**.

**Obdurodon** (order **\*Monotremata**, family Ornithorhynchidae) A genus of early **\*Miocene** platypus, with two species, known from very well-preserved material from such sites as Riversleigh, Queensland, Australia. They possessed well-developed, functional teeth, unlike the living platypus, and had skull bones that were less fused, from which it can be seen that monotremes (Monotremata), unlike other living mammals, retained **\*septomaxillae**.

**Oberon (Uranus IV)** One of the major satellites of **\*Uranus**. Its radius is 761.4 km; mass  $30.14 \times 10^{20}$  kg; mean density 1630 kg/m<sup>3</sup>; albedo 0.24. The surface is extensively cratered, the craters being surrounded by bright ejecta. Near the centre of Oberon there is a large crater with a bright central outcrop and background of dark material.

**Obik Sea** See **URAL SEA**.

**objective** The magnifying lens which is situated at the base of a microscope tube immediately above the specimen. Objectives may be described in terms of lens quality (e.g. achromat, apochromat, or fluorite), magnification (which may be 5×, 10×, 20×, or larger), and numerical aperture (NA) which is a measure of the amount of light converging on the specimen through the lens. Its value is normally between 0.1 and 0.9.

**oblate** **1.** Applied to the ellipsoid formed by rotating an ellipse about its minor axis. **2.** Description of the shape of a **\*clast** which is tabular or disc-shaped in appearance. It is characterized by a ratio of intermediate to long diameters of more than 2:3, and a ratio of short to intermediate diameters of less than 2:3. See **PARTICLE SHAPE**.

**oblate uniaxial strain** The **\*strain** state obtained when a reference sphere is shortened along its z-axis and extended by an equal amount along its x- and y-axes and all intermediate axes lying in their plane and perpendicular to the z-axis. Such strain alters the sphere into an **\*oblate** ellipsoid.

**oblique extinction (inclined extinction)** In optical mineralogy, the **\*extinction** of a mineral (i.e. it becomes dark) in **\*crossed** polars with its cleavage traces or **\*crystal** boundaries positioned at an oblique angle to the E–W and N–S planes of vibration of the **\*polarizer** and **\*analyser**. Several different angles of extinction may be determined for the same mineral, and the maximum angle may be used for identification purposes.

**oblique-slip fault** A **\*fault** in which the displacements of the **\*strike-slip** and **\*dip-slip** components have very similar magnitudes; fault movement occurs obliquely across the fault surface.

**oblique-slip margin** A **\*convergent** or **\*divergent** margin where two **\*plates** have a significant transform motion (see **TRANSFORM FAULT**).

**obliquity of the ecliptic** See **MILANKOVITCH CYCLES**.

**obrusion** See **OBRUTION**.

**obruption (obrusion)** Sudden burial.

**obruption deposit** A fossil assemblage that has been preserved by the very rapid burial of intact organisms.

**obsequent** Applied to a land-form whose orientation is opposite to that which may have been expected. For example, an obsequent stream flows against the **\*dip** of underlying strata; and an obsequent **\*fault-line** scarp faces the opposite way to the original fault scarp.

**observation well** A well that is used to observe changes in **\*groundwater** levels over a period, or more specifically during a pumping test. Pumping

does not normally take place from observation wells which are often relatively small in diameter.

**obsidian** Volcanic \*glass of dacitic (see [DACITE](#)) or rhyolitic (see [RHYOLITE](#)) composition.

**Occam's razor (Ockham's razor)** The axiom that *pluralitas non est ponenda sine necessitate* (multiplicity ought not to be posited without necessity); i.e. when alternative hypotheses exist, the one requiring the fewest assumptions should be preferred. This was first proposed by William of Occam (William Ockham) (c. 1280–1349). See also [PARSIMONY](#).

**occipital** Pertaining to the posterior part of the \*cranium.

**occipital condyle** At the back of the skull, a bony knob which articulates with the first \*vertebra. It is absent in fish, and double in \*amphibians and \*mammals.

**occluded front** The composite front formed when a \*warm front and the sector of warm air behind it is lifted and overtaken by a \*cold front. See also [OCCLUSION](#).

**occlusion** Stage in the development of a frontal \*depression during which the warm-sector air is gradually lifted from the surface and above the colder air. The word is used to describe an \*occluded front. An occlusion may be a 'warm occlusion' such that warmer air follows the frontal system, or a 'cold occlusion', such that colder air follows in the rear. A 'warm occlusion' has characteristics similar to those of a warm front, but there is an upper cold front ahead and aloft which brings showers from \*cumulus and \*cumulonimbus cloud. A 'cold occlusion' exhibits cold-front features, but is preceded by warm-front cloud types. Occlusions formed in the later stages in the development of travelling depressions commonly cross north-western Europe, especially the 'warm' type in winter and the 'cold' type in summer.

**ocean** Salt water mass that occupies more than two-thirds of the surface of the Earth (70.8%). The oceans contain  $1370 \times 10^6$  km<sup>3</sup> of water; the average depth is 3730 m.

**Ocean-1B** See [HAIYANG-1B](#).

**ocean-basin crustal type** The type of very stable \*crust that covers vast areas of the ocean floor. It averages 11 km in thickness and is broken by long linear \*faults or \*fractures. There are \*volcanoes in some places.

**ocean-basin floor** The ocean floor in those parts of the oceans that are more than 2000 m deep. It occupies approximately one-third of the \*Atlantic and \*Indian Ocean floors, and three-quarters of the \*Pacific Ocean floor.

**ocean current** Large-scale water movement in an ocean, arising from three main causes: (a) wind stresses acting on the surface of the sea; (b) tidal motion caused by the variable attractions of the Sun and Moon; and (c) density differences in sea water, caused by differential heating and cooling, \*salinity differences, or variations in the suspended-sediment concentration of sea-water masses.

**ocean gyre** See GYRE.

**oceanic crust** The oceanic rocks which form 65% of the Earth's surface, and are the upper part of the oceanic \*lithosphere, overlying the \*Mohorovičić seismic discontinuity. The oceanic crust comprises four seismic layers, commencing at an average depth of 4.5 km below sea level. The uppermost layer (\*sediments) varies in thickness from being absent over the \*mid-oceanic ridges to 2–3 km near the \*continental shelves. The other three layers are of remarkably constant thickness and \*seismic velocity. Layer 2 is formed by \*basaltic lavas with \*P-wave velocities of 5 km/s and a \*dyke complex just under 2 km thick, overlying a 5 km thick gabbroic layer (layer 3) with a P-wave velocity of 6.7 km/s. Layer 4 is a thin (less than 0.5 km) layer with a P-wave velocity of 7.4 km/s, immediately overlying the \*mantle where the P-wave velocity is 8.1 km/s. The total thickness is about 11 km and shows little variation throughout the ocean basins, despite the change in ocean \*bathymetry from the ridges to the \*trenches. The crust is cut by \*fracture zones. It is largely aseismic away from \*spreading ridges and \*subduction zones. The presence of linear \*magnetic anomalies, mostly paralleling the present mid-oceanic ridges, allows the dating of the ocean floors, as does the dating of sediments immediately overlying the \*igneous part of the crust and obtained by drilling during the \*DSDP and \*IPOD projects. The oldest oceanic crust, within the present ocean basins, is less than 200 million years old; it is

found in the W. Pacific and the N. Atlantic, and was formed by igneous activity at spreading (accretionary) **\*plate margins** as part of the **\*sea-floor** spreading process. Layer 3 represents the cooled **\*magma chamber** that originally fed the overlying basaltic dykes and lavas.

**oceanicity** The effects of maritime influences on a climate. *See also* CONTINENTALITY.

**oceanic plateau** Extensive, topographically high area of an ocean floor that rises to within 2–3 km of the sea surface above the abyssal floor, e.g. the plateau on which Iceland stands, the Galápagos Islands platform, and the Azores platform. Many Pacific plateaux, e.g. the Magellan Rise and Ontong Java Plateau, have a thick covering of carbonate oozes overlying volcanic rocks. The origin of such plateaux is volcanic, but many are now inactive.

**oceanic trench** *See* TRENCH.

**oceanic-trench crustal type** A type of very unstable **\*crust** in which **\*earthquakes** are frequent. Ocean trenches are associated with mountain belts and **\*island** arcs.

**ocean-island basalt (OIB)** Quartz **\*tholeiites**, **\*alkali basalts**, and **\*nephelinites** found on **\*volcanoes** which build up from the ocean floor to form the ocean islands away from ocean **\*ridges**. Examples include the Cape Verde Islands, Ascension Island, Tristan da Cunha Island, and Gough Island in the Atlantic, and the **\*Hawaiian-Emperor** **\*sea-**mount chain in the Pacific. Compared to MORBs (*see* MID-OCEAN-RIDGE BASALT), OIBs are enriched in (a) **\*large-ion** lithophile (LIL) elements, (b) light **\*rare-earth** elements (LREE) relative to heavy rare-earth elements (HREE), and (c) **\*incompatible** elements such as Ti, Ga, Li, Nb, V, Zn, Zr, and Y. Radiogenic-**\*isotope** and **\*trace-element** evidence suggests that OIBs are formed by **\*partial** melting of enriched **\*mantle**.

**OceanSat-2** A spacecraft developed by the Indian Space Research Organization that studies surface winds over the ocean and ocean surface strata, monitors chlorophyll concentrations and phytoplankton blooms, and studies **\*aerosols** and suspended sediments in water. It was launched on 23 September 2009, from India, into a Sun-synchronous near-circular orbit at an altitude of about 720 km. *See also* SCATTEROMETER SATELLITE-1.

**oceanus** A very extensive, dark area on the surface of an extraterrestrial body.

**ocean wave** Disturbance of the ocean's surface, seen as an alternate rise and fall of the surface. Ocean waves are of several types: (a) wind-generated waves, e.g. sea waves (chaotic wave pattern) and swell (long-period waves); (b) catastrophic waves, e.g. \*tsunamis, landslide surges, and storm surges; and (c) internal waves (subsurface waves at the boundary between two water layers).

**Ochoan** The final \*stage (260.4–251 Ma ago) in the Late \*Permian of N. America, underlain by the \*Guadalupian, overlain by the \*Triassic (see SCYTHIAN), and roughly contemporaneous with the upper \*Kazanian and \*Changhsingian stages.

**ochre** An iron-rich sediment, used as a pigment. *Compare* UMBER.

**ochric horizon** A light-coloured, mineral \*soil horizon, usually at the soil surface, and characteristic of arid-environment soils.

**Ockham's razor** *See* OCCAM'S RAZOR.

**OCO-2** *See* ORBITING CARBON OBSERVATORY-2.

**octahedrite** *See* ANATASE.

**octahedron** A three-dimensional, eight-sided form in which all the faces are equilateral triangles (i.e. it resembles a double pyramid). In \*crystallography, a \*crystal form in which the \*crystal faces cut all three \*crystallographic axes at equal distances.

**Octocorallia (octocorals)** A subclass of the class \*Anthozoa, poorly known in the fossil record, that ranges from \*Ordovician to Recent. The colony is shaped like a flat fan of interconnecting branches from which project tubes with many zooids. *Heliopora* is an important \*reef former. The pennatulaceans (Pennatulacea) are also octocorals.

**octocorals** *See* OCTOCORALLIA.

**ocular** *See* EYEPIECE.

**OD** Ordnance datum.

**Odderade** The third *\*interstadial* within the *\*Weichselian*, which occurred between 70 000 years BP and 60 000 years BP, between the *\*Brørup* and *\*Moershoofd* Interstadials. The vegetational evidence indicates tundra conditions with some temperate boreal elements having been present. After the Odderade Interstadial polar-desert conditions prevailed until the Moershoofd Interstadial.

**Oddo–Harkins rule** Rule stating that the *\*cosmic* abundance of elements with an even atomic number is greater than that of adjacent elements with an odd atomic number. Consequently, a graph plotting relative atomic abundance against increasing atomic number (*Z*) displays a ‘toothed’ curve, rather than a smooth line. The reason for this is connected with processes such as helium burning (see **NUCLEOSYNTHESIS**).  $^4_2\text{He}$  is a basic building block, and so additions produce even numbers, e.g.  $^4_2\text{He} + ^4_2\text{He} \rightarrow ^8_4\text{Be}$ ;  $^8_4\text{Be} + ^4_2\text{He} \rightarrow ^{12}_6\text{C}$ .

**Odin** An international *\*minisatellite* mission led by Sweden, in partnership with Canada, Finland, and France, that supplies data for atmospheric research, including information on stratospheric ozone chemistry, the summer *\*mesosphere*, and the coupling of atmospheric regions. It was launched on 20 February 2001, from the Svobodny Cosmodrome, Russia, into a Sun-synchronous *\*polar orbit* at an altitude of 600 km.

**Odintsovo** An *\*interstadial* in the Saale Drift of European Russia, marked by a temperate, broadleaved-forest flora.

**ODP** Ocean Drilling Programme. See **DEEP SEA DRILLING PROGRAMME**.

**Oe** See **OERSTED**.

**oedometer** Instrument for testing *\*consolidation* of small samples, including *\*coefficient* of compressibility and *\*coefficient of consolidation*. The sample is compressed in a cylinder allowing only vertical consolidation; the resulting changes in volume over a specific time are measured.

**oersted (Oe)** A unit of magnetic field intensity, now replaced by SI units.  $1 \text{ A/m} = 4\pi \times 10^{-3} \text{ Oe}$ .

**offlap** A conformable sequence of inclined strata, deposited during a marine *\*regression*, in which each stratum is succeeded laterally by progressively younger units, marking the direction in which the sea retreated. *Compare* **OVERLAP**. *See also* **OVERSTEP**; **SEISMIC STRATIGRAPHY**.

**offset** The distance between a *\*geophone*, or the centre of a geophone *\*group*, and the *\*shot* position. In-line offset is the displacement within the line of the spread; perpendicular (normal) offset is the distance at right angles between the shot and the line of the spread; reflection offset is the horizontal displacement of a dipping reflection event to its migrated position.

**offshore bar** *See* **BAR**.

**offshore zone** Zone extending seaward from the point of low tide to the depth of wave-base level or to the outer edge of the *\*continental shelf*.

**ogive** Banded pattern on the surface of a *\*glacier*. Normally it is convex down-glacier due to relatively high velocities at the centre. The banding effect may be due to alternating bands of white ice (containing many air bubbles) and dark ice (few air bubbles), or to variations in longitudinal pressure.

**Ohauan** A New Zealand *\*stage* (152–150.8 Ma ago) of the Late *\*Jurassic* *\*epoch*, preceded by the *\*Heterian* and followed by the *\*Puaroran*.

**Ohm's law** The ratio of the voltage ( $V$ ) applied to a conductor and the electric current ( $I$ ) caused to flow through it at constant temperature is constant, and is the electrical resistance ( $R$ ) of the conductor, such that  $V/I = R$ . At high current densities the law may break down for some materials.

**OIB** *See* **OCEAN-ISLAND BASALT**.

**oikocryst** A crystal that encloses others in a poikilitic fabric.

**oil** *See* **PETROLEUM**.

**oil immersion** *See* **IMMERSION OBJECTIVE**.

**oil shale** Dark grey or black *\*shale* containing organic substances that yield liquid *\*hydrocarbons* on distillation, but that do not contain free *\*petroleum*.

**Oka/Demyanka** An equivalent in Russia of the *\*Elsterian* (*\*Mindel*), or its *\*Anglian* equivalent, which passes beneath the *\*Dnepr \*drift*. Little is known of its occurrence.

**okta** See CLOUD AMOUNT.

**Older Drift** All the older *\*drift* deposits that mark the maximum extent of ice advance during the last (*\*Devensian*) glaciation in the British Isles. Compare NEWER DRIFT.

**Older Dryas** See DRYAS.

**Oldest Dryas** See DRYAS.

**Oldham, Richard Dixon** (1858–1936) A British seismologist, Oldham worked for the Indian Geological Survey. In 1897 he showed that *\*P-* and *\*S-waves* could be distinguished from each other on *\*seismograms*, and that they travelled through the *\*Earth's* interior. In 1906 he was able to show that the Earth has a fluid *\*core*, through the existence of the S-wave shadow zone, and was able to estimate its size.

**Old Red Sandstone Continent** The continental facies of the *\*Devonian* in the British Isles. It is characterized by red *\*sandstones* and *\*conglomerates* which were deposited under terrestrial conditions.

**Olduvai** A normal *\*polarity subchron* which occurs within the *\*Matuyama* reversed *\*chron*. It is dated at 1.76–1.98 Ma ago.

**Olenekian** A *\*stage* of the *\*Triassic* period, from 249.7–245 Ma ago, preceded by the *\*Induan* and followed by the *\*Anisian*. See SCYTHIAN.

**Olenelloidea** See REDLICHIIA.

**oligo-** From the Greek *oligos* meaning ‘small’ and *oligoi* meaning ‘few’, a prefix meaning few or small; in ecology it is often used to denote a lack, e.g. ‘*\*oligotrophic*’ meaning ‘nutrient-poor’ and ‘*\*oligomictic*’ meaning ‘subject to little mixing’.

**Oligocene** An *\*epoch* (33.9–23.03 Ma ago) of the *\*Palaeogene* period. It follows the *\*Eocene* and precedes the *\*Miocene* epochs. The Oligocene epoch comprises the *\*Rupelian* and *\*Chattian \*ages*.



<http://www.geologypage.com/2014/04/oligocene-epoch.html>

- Oligocene Epoch.

**Oligochaeta** (phylum *\*Annelida*) Class of annelid worms which possess very well-developed *\*metameric* segmentation. The segments have bristles, but parapodia (movable, paired lateral appendages) are not present. They are all hermaphrodites; asexual reproduction is predominant in aquatic forms. Eyes and tentacles are absent. A few marine forms occur but most are freshwater or terrestrial. There are fifteen families, and the class is first recorded from the Upper *\*Ordovician*.

**oligoclase** See *PLAGIOCLASE FELDSPAR*.

**oligohaline water** See *HALINITY*.

**oligomictic** 1. A *\*conglomerate* containing *\*clasts* of only a few different rock types. Compare *POLYMICTIC*. 2. Applied to lakes that are thermally almost stable, mixing only rarely. This condition is characteristic of tropical lakes with very high (20–30 °C) surface temperatures.

**oligotaxic times** A period of low biological diversity among marine organisms, associated with low sea levels (lowstand), sharp temperature gradients, and stronger ocean currents. Compare *POLYTAXIC TIMES*.

**oligotrophic** Applied to waters poor in nutrient and with low *\*primary productivity*. Compare *EUTROPHIC*.

**olisthostrome** See *OLISTOSTROME*.

**olistolith** See *OLISTOSTROME*.

**olistostrome (olisthostrome)** A sedimentary deposit which consists of a chaotic mass of rock and contains large *\*clasts* composed of material older than the enclosing sedimentary sequence. The clasts may be gigantic and are then called ‘olistoliths’. Such deposits are generally formed by *\*gravity sliding* of material, sometimes into oceanic *\*trenches*. Olistostromes have also been called ‘sedimentary mélange’ (see *MÉLANGE*).

**olivine** A major rock-forming *\*mineral* group belonging to the *\*nesosilicates*, forming a complete *\*solid solution* series between forsterite ( $Mg_2SiO_4$ ) and fayalite ( $Fe_2SiO_4$ ); sp. gr. 3.22–4.39 increasing with

increasing iron content; *\*hardness* 6–7; *\*orthorhombic*; usually olive-green, but white or yellowish in forsterite, brown or black in fayalite; colourless *\*streak*; *\*vitreous* *\*lustre*; crystals rare, short, *\*prismatic*, usually develops as granular aggregates; *\*cleavage* poor {010}; occurs in silica-poor, *\*igneous* rocks (e.g. *\*basalt*, *\*gabbro*, *\*troctolite*, and *\*peridotite*), extensively with *\*pyroxene* in *\*dunites*, and in *\*stony* meteorites and lunar basalts; alters readily to *\*serpentine* during weathering or *\*hydrothermal* alteration.

**olivine dolerite** See DOLERITE.

**omphacite** A rare *\*clinopyroxene* of isolated composition (Ca,Na)(Mg,Fe<sup>2+</sup>,Fe<sup>3+</sup>,Al)[Si<sub>2</sub>O<sub>6</sub>] and with similar properties to *\*jadeite* and *\*augite*; sp. gr. 3.16–3.43; *\*hardness* 5–6; green to dark green; *\*massive* or *\*granular*; occurs in *\*eclogites* at high temperatures and pressures in association with *\*pyrope* garnet.

**oncoid** See ONCOLITE.

**oncolite (oncoid, oncolith)** Spherical or subspherical particle, up to 5 cm across, produced by the accretion of sedimentary material on to a mobile grain through the action of *\*algae*.

**oncolith** See ONCOLITE.

**one-circle reflecting goniometer** See GONIOMETRY.

**Onesquethawian (Onondagan)** See ULSTERIAN.

**one-way travel time** The time taken for a seismic wave to travel one way through a medium. In a reflection event it is half the *\*two-way* travel time. Compare TRANSIT TIME.

**onion weathering** See SPHEROIDAL WEATHERING.

**onlap** The progressive spreading of successive beds of *\*sediment* over an increasingly wider area, usually as a result of rising sea level or the gradual subsidence of a land area. Onlap results in an overlapping (see OVERLAP) stratigraphical relationship between the sedimentary succession and the underlying *\*basement* rocks. The upper, younger beds ‘pinch out’ updip

against the older, inclined surface on which they rest. *Compare* OFFLAP. *See also* BASELAP; COASTAL ONLAP; DOWNLAP; TOPLAP.

**Onnian** A regional *\*stage* (453–449 Ma ago) of the Late *\*Ordovician* in the Upper *\*Caradoc*, underlain by the *\*Actonian* and overlain by the *\*Pusgillian*.

**Onondagan (Onesquethawian)** *See* ULSTERIAN.

**Ontarian** *See* TELYCHIAN.

**ontogenetic heterochrony** In colonial animals, *\*heterochrony* that affects individuals, rather than the colony as a whole. *Compare* ASTOGENETIC HETEROCHRONY and MOSAIC HETEROCHRONY.

**ontogeny** The development of an individual from fertilization of the egg to adulthood.

**Onverwacht** *See* SWAZIAN.

**onyx (sardonyx)** A mixture of chalcedonic (*see* CHALCEDONY) silica  $\text{SiO}_2$  and hydrous silica  $\text{SiO}_2 \cdot n\text{H}_2\text{O}$ ; occurs as flat, banded varieties with white, grey, brown, red, and black bands.

**oo-** Prefix in the *\*Folk* classification, for *\*limestones* containing *\*ooids*, e.g. *\*oosparite*.

**oobiosparite** *See* FOLK LIMESTONE CLASSIFICATION.

**ooid (oolith)** Subspherical, sand-sized, *\*carbonate* particle that has concentric rings of calcium carbonate surrounding a nucleus of another particle. *See also* OOLITE.

**oolite** Commonly occurring *\*limestone*, consisting largely of *\*ooids*.

**oolith** *See* OOID.

**oolitic** Composed, or largely composed of *\*ooids*.

**oolitic limestone** *See* OOLITE.

**oomicrite** A *\*limestone*, defined by the Folk classification as comprising *\*ooids* set in a *\*micrite* *\*matrix*.

**oomoldic porosity (oomouldic porosity)** A form of *\*secondary* porosity developed in oolitic *\*limestone* (see OOLITE), due to the preferential dissolution of the *\*ooids*, leaving a series of empty spaces. See POROSITY.

**oomouldic porosity** See OOMOLDIC POROSITY.

**<sup>18</sup>O:<sup>16</sup>O ratio** See OXYGEN-ISOTOPE RATIO.

**Oort cloud** A spherical zone surrounding, but gravitationally bound to the *\*solar* system, at distances between 20 000 and 100 000 AU from the Sun, which contains about 10<sup>12</sup> comets. Its existence was first proposed in 1950 by the Dutch astronomer, Jan Hendrik Oort (1900–92). Gravitational perturbations by passing stars or molecular clouds result in comets acquiring orbits within the orbit of *\*Jupiter*.

**oosparite** A *\*limestone*, defined by the *\*Folk* classification as comprising *\*ooids* together with sparry calcite *\*cement* (*\*sparite*).

**ooze** A *\*pelagic* mud consisting of the calcareous or siliceous remains of pelagic organisms (e.g. *\*coccoliths*, and *\*tests* of *\*foraminiferids* and diatoms (*\*Bacillariophyceae*)), and *\*hemipelagic* *\*clay minerals*. *\*Calcareous* oozes will accumulate in the deep oceans at depths shallower than the *\*carbonate compensation depth (CCD)*. See also GLOBIGERINA OOZE; DIATOM OOZE; PTEROPOD OOZE; RADIOLARIAN OOZE.

**opacus** From the Latin *opacus* meaning ‘shady’, a variety of extensive cloud that obscures the Sun or Moon. It is used to describe *\*stratus*, *\*stratocumulus*, *\*altostratus*, and *\*altocumulus*. See also CLOUD CLASSIFICATION.

**opal** Hydrrous silica SiO<sub>2</sub>.nH<sub>2</sub>O associated with the chalcedonic (see CHALCEDONY) varieties of silica. A layer of water molecules trapped near the mineral surface causes the *\*iridescence* (*\*opalescence*) which is a diagnostic property of opal; sp. gr. 1.99–2.25; *\*hardness* 5.5–6.5; amorphous; colourless, or milky-white to grey, red, brown, blue, green, to nearly black; resinous *\*lustre*; normally *\*massive*, but can be stalactitic, *\*botryoidal*, and also in veinlets, the various varieties depending on the amount of water contained in the mineral, which can vary from 6% to 10%; no *\*cleavage*; *\*conchoidal* fracture; normally deposited at low temperatures

from silica-bearing waters, and occurs as fissure fillings in rocks of any kind, and especially near geysers and hot springs. The variety known as precious opal has a milky-white or sometimes black body colour which exhibits a brilliant play of colours, usually blues, reds, and yellows. The colours can often disappear with the loss of water when the mineral is exposed to air.

**opalescence** In *\*mineralogy*, a pearly or milky mineral *\*lustre* resembling that of *\*opal*. It results from the reflection and refraction of light from the surface and possibly subsurface layers of the mineral. In *\*opal*, water molecules are trapped in the subsurface layers, giving the opalescent quality.

**opaque mineral (ore mineral)** In transmitted-light microscopy, a mineral which appears black in *\*thin* section in *\*plane-polarized* light. The term is often used synonymously with 'ore mineral' although neither term is strictly correct; for example, *\*pyrite* is opaque but rarely an ore, and *\*sphalerite* is often an ore but rarely opaque.

**open-cast mining (strip mining)** Mining in which large strips of land are excavated in order to extract materials without sub-surface tunnelling. It may result in short- and long-term environmental damage.

**open fold** As defined by M. J. Fleuty (1964), a *\*fold* whose inter-limb angle (see *FOLD ANGLE*) is between 90° and 170°. The classification is based on the degree of tightness of folds.

**open form** In *\*crystallography*, *\*crystal* faces are often described in groups ('forms'); if the form is open at each end (i.e. it does not totally enclose space), it is called an 'open form', e.g. the four prismatic faces of a *\*tetragonal* crystal. *Compare* *CLOSED FORM*.

**open hole** A *\*borehole* that has not yet been cased.

**open-pit mining** A method of extraction used where the *\*overburden* is limited and easily stripped, but where waste has to be transported to external dumps. It is generally used where deposits are limited laterally but are thicker than in *\*open-cast* mining.

**open system** In *\*geomorphology*, a *\*system* in which both energy and matter enter and leave across the system boundary.

**operculum** 1. In animals, a lid or cover, sometimes hinged, occurring, for example, in some cylindrical *\*rugose* corals, some *\*Bryozoans*, and in *\*gastropods*. 2. A flap of skin covering the gills in bony fish (*\*Osteichthyes*).

**Ophelia (Uranus VII)** One of the lesser satellites of *\*Uranus*, with a diameter of 16 km. It was discovered in 1986.

**ophicalcite** See SERPENTINE.

**Ophiocystioidea (ophiocystioids)** (phylum *\*Echinodermata*, subphylum Echinozoa) Class of extinct, free-living echinozoans, which possess a low, dome-shaped *\*test* lacking arms. The *\*peristome* is centrally placed on the *\*ventral* surface and is enclosed in a complex jaw system. The *\*ambulacra* are confined to the ventral surface and are composed of three rows of plates. The *\*interambulacra* are narrow and consist of one plate column only. The class is known from the *\*Ordovician* to *\*Devonian*.

**ophiocystioids** See OPHIOCYSTIOIDEA.

**ophiolite (ophiolite complex)** Sequence of rock types, consisting of deep-sea *\*sediments* lying above basaltic *\*pillow* lavas, *\*dykes*, *\*gabbro*, and *\*ultramafic peridotite*. Some are the remnants of main *\*oceanic* crust, others of crust formed in *\*back-arc* basins.

**Ophiomorpha** With *\*Skolithos*, an *\*ichnoguild* of *\*trace fossils* characterized by vertical or steeply inclined *\*burrows* with thick, knobby walls. The interior of the burrow system is smooth. *Ophiomorpha* ranges from the *\*Permian* to the present day and is associated with arthropods (*\*Arthropoda*). Individual tubes have a diameter of 3–6 cm, and may be branched, with large, bulbous terminations.

**ophitic texture** Large *\*crystals* of *\*augite* enclosing, either wholly or partially, laths of *\*plagioclase* feldspar, and found within *\*dolerites* and *\*basalts*. The texture is a particular type of *\*poikilitic* texture.

**Ophiuroidea (brittle-stars)** (subphylum *\*Asterozoa*, class Stellerioidea) Subclass of brittle-stars in which a central, rounded disc is sharply demarcated from the long, slender arms. The body cavity of the arms is filled almost completely with axial skeleton. The subclass is known from the *\*Ordovician* to Recent.

**Opisthobranchia** (phylum *\*Mollusca*, class *\*Gastropoda*) Subclass of marine organisms, which are often ‘naked’, that includes the sea slugs and pteropods, and ranges from the *\*Cretaceous* to the present day. They have only one gill and are thought to represent an evolutionary grade between the *\*prosobranch* (single-gilled) gastropods and the pulmonate gastropods (i.e. gastropods in which the *\*mantle cavity* functions as a lung).

**opisthodontic** Applied to the ligament of a bivalve when it is in a position posterior to the umbones (see *UMBO*). Compare *AMPHIDETIC*.

**opisthogyral** See *OPISTHOGYRATE*.

**opisthogyrate (opisthogyral)** Applied to the umbones (see *UMBO*) of bivalves (*\*Bivalvia*) where these are so curved as to point in a posterior direction.

**opisthoparian suture** See *CEPHALIC SUTURE*.

**opisthosoma** See *ARACHNIDA*; *CHELICERATA*.

**Opoitian** A *\*stage* (4.8–3.6 Ma ago) in the Early *\*Pliocene* of New Zealand, underlain by the *\*Kapitean*, overlain by the *\*Waipipian*, and roughly contemporaneous with the upper *\*Zanclean* (Tabianian) stage.

**Oppel, Albert** (1831–65) A German geologist and palaeontologist, Oppel made extensive studies of the *\*Jurassic* rocks of Europe. He devised a scheme to divide geologic formations into zones, each based on the period of existence of a single organic species, later to be termed ‘*\*index fossils*’. See *OPPEL ZONE*.

**Oppel zone** A *\*concurrent* range zone as originally conceived by *Albert \*Oppel*. The lower boundary of the zone is usually determined by the first appearance of one diagnostic *\*taxon* and the upper boundary by the last appearance of another (either might extend above or below into a region of overlap with the other taxon at the upper or lower end of their range). Other characteristic taxa may be contained within the zone or extend to either side, although long-ranging and slowly evolving lineages are not usually included in the diagnostic assemblage. Not all the significant taxa are required to be present at all levels and in all places. The zone is named after one of the diagnostic species. Oppel zones are thus more flexible and subjective than concurrent range zones in the rigorously applied sense of

that term, although the term ‘concurrent range zone’ is commonly used for what is, in fact, an Opper zone.

**optical continuity** In optical *\*mineralogy*, the situation that occurs where two mineral grains are oriented such that their optical properties are consistent and in the same orientation with respect to their fundamental crystallographic properties.

**optical emission spectrum** See SPECTRUM.

**optical goniometry** See GONIOMETRY.

**optical indicatrix** See INDICATRIX.

**Optical Satellite-3000 (OptSat-3000)** A generation of *\*minisatellites* built by Israeli Aerospace Industries that provide high-resolution imagery. The first in the series was launched on 2 August 2017, from Kourou, French Guiana, into a Sun-synchronous orbit at an altitude of 450 km.

**optic axis** In mineral optics, the direction perpendicular to a circular cross-section of an *\*indicatrix*. Sections cut parallel to this circular cross-section are *\*isotropic*.

**OptSat-3000** See OPTICAL SATELLITE-3000.

**opx** See ORTHOPYROXENE.

**Or** See ALKALI FELDSPAR.

**o-ray** See EXTRAORDINARY RAY.

**orbicular** Disc-shaped, circular, or globular.

**orbicular texture** Concentric shells of contrasting mineralogy and texture which form spherical masses of diameter 2–15 cm in *\*basic* and *\*felsic* *\*plutonic* *\*igneous* rocks.

**orbit** 1. The bony socket of the eye. 2. The path described by a body moving around another under gravitational attraction. See EQUATORIAL ORBIT; GEOSTATIONARY ORBIT; GEOSYNCHRONOUS ORBIT; LOW EARTH ORBIT; POLAR ORBIT; SUN-SYNCHRONOUS ORBIT.

**orbital forcing** The influence on climate and sea level of changes in the orbit and rotation of the Earth. See [MILANKOVITCH CYCLES](#).

**orbit period** The time taken by a body to complete a single orbit.

**Orbiting Carbon Observatory-2 (OCO-2)** A \*NASA mission that studies the distribution of sources and sinks of atmospheric CO<sub>2</sub>. The satellite was launched on 2 July 2014, from California, into a Sun-synchronous orbit at an altitude of 705 km. It replaces OCO-1, which failed at launch.

**OrbView-5** See [GEOEYE-1](#).

**order** See [CLASSIFICATION](#).

**Ordian** A \*stage (520–510 Ma ago) of the Lower to Middle \*Cambrian of Australia, underlain by the ‘Lower Cambrian’ and overlain by the \*Templetonian.

**ordinary ray** See [EXTRAORDINARY RAY](#).

**Ordnance Survey (OS)** The national mapping agency in the United Kingdom, which publishes general and specialized maps of the country at several scales. The OS began in 1791, when the British government, fearing invasion following the French Revolution, instructed its board of ordnance (equivalent to a defence ministry) to survey and map the south coast of England. That operation led to the surveying and mapping of the entire country. While remaining a government agency, since 1999 the OS has had the status of a ‘trading fund’, which gives it greater responsibility for its own finances and planning, and more freedom to develop new initiatives.

**Ordovician** The second (488.3–443.7 Ma ago) of six \*periods that constitute the \*Palaeozoic era, named after an ancient Celtic tribe, the Ordovices. The Ordovician follows the \*Cambrian and precedes the \*Silurian. It is noted for the presence of various rapidly evolving \*graptolite genera and of the earliest jawless fish.



<https://ucmp.berkeley.edu/ordovician/ordovician.php>

- The Ordovician Period.

**Ordovician V** A *\*stage* in the Late *\*Ordovician* epoch, 460.9–455.8 Ma ago, preceded by the *\*Darriwilian* and followed by the *\*Ordovician VI*.

**Ordovician VI** A *\*stage* in the Late *\*Ordovician* epoch, 455.8–445.6 Ma ago, preceded by the *\*Ordovician V* and followed by the *\*Hirnantian*.

**ore** A *\*mineral* or *\*rock* that can be worked economically.

**orebody** Accumulation of *\*minerals*, distinct from the host rock, and rich enough in a metal to be worth commercial exploitation.

**ore genesis** Process by which a mineral deposit forms. Metalliferous mineral deposits may be syngenetic (formed at the same time as the host rocks) or epigenetic (deposited later than the host rocks). Deposits may be classified according to their processes of formation into *\*igneous*, *\*sedimentary*, metamorphic (see *METAMORPHISM*), or *\*hydrothermal*.

**ore grade** The concentration of an element of interest in a potentially mineable *\*ore* deposit.

**ore microscope** See *POLARIZING MICROSCOPE*.

**ore microscopy** See *REFLECTED-LIGHT MICROSCOPY*.

**ore mineral** A metalliferous *\*mineral* that may be extracted profitably from an *\*orebody*. In mineralogy the term is applied to those minerals that may polish with a *\*metallic \*lustre* even when dispersed. See also *OPAQUE MINERAL*.

**Oretian** A New Zealand *\*stage* (225–218 Ma ago) of the Late *\*Triassic \*epoch*, preceded by the *\*Kaihikuan* and followed by the *\*Otamitan*.

**organic soil** Soil with a high content of organic matter and water. The term usually refers to *\*peat*. The *\*USDA* defines an organic soil as one with a minimum of 20–30% organic matter, depending on the *\*clay* content.

**organic weathering** Weathering caused by the action of living organisms.

**Oriental faunal realm** Area that encompasses India and Asia south of the Himalayan–Tibetan mountain barrier, and the Australasian archipelago, excluding New Guinea and the Celebes. There are marked similarities with the *\*Ethiopian* realm (e.g. both have elephants and rhinoceroses) but there are endemic groups (see *ENDEMISM*), e.g. pandas and gibbons.

**orientation survey** The first stage in a *\*geochemical soil survey*, concerned with the choice of sampling scheme, type of sample, and analytical method.

**original horizontality** See *LAW OF ORIGINAL HORIZONTALITY*.

**origination** The first appearance of a new species. Comparing the rate at which new species appear (origination rate) with the rate at which species go extinct (extinction rate) within an *\*ecosystem* reveals whether the system was stable (origination = extinction), diversifying (origination > extinction), or losing species (origination < extinction).

**Oriskanyan (Deerparkian)** See *ULSTERIAN*.

**Ornithischia** One of two orders of *\*Mesozoic \*dinosaurs*, distinguished primarily on the basis of a bird-like pelvis (see *PELVIC GIRDLE*; *PUBIS*). Ornithischian dinosaurs were exclusively vegetarian and produced both bipedal forms (*\*Ornithopoda*) and four-footed forms.

**ornithischian dinosaur** See *ORNITHISCHIA*.

**ornithomimid** One of the *\*coelurosaurs*, which were *\*theropod \*dinosaurs* from the Upper *\*Cretaceous*. They are referred to as the 'ostrich' dinosaurs because of their general build, and their name means 'bird imitators'.

**Ornithopoda** Suborder of *\*ornithischian* dinosaurs which had an essentially bipedal gait, including several families, e.g. iguanodonts (*\*Iguanodontidae*) and hadrosaurs (*\*Hadrosauridae*). The ornithopods are regarded as the most primitive of the ornithischian suborders.

**orogen** The body of rock involved in one or more *\*orogenies*. Many authors use the term as a synonym for *\*orogenic* belt.

**orogenesis** See *OROGENY*.

**orogenic belt (mobile belt)** A linear or arcuate zone, on a regional scale, which has undergone compressional tectonics. The histories of many orogenic belts have been interpreted using *\*plate-tectonics* models involving the *\*subduction* of oceanic *\*lithosphere* (e.g. the *\*Andean* orogenic belt), the collision of major continental masses (e.g. the *\*Himalayan* orogenic belt), or the accretion of *\*terranes* (e.g. the Cordillera of the western USA and Canada). Orogenic belts were formerly expected to show the phases of the *\*orogenic cycle*, but this has now been replaced by a

search for identifiable stages of the **\*Wilson** cycle to see if the belt can be interpreted using a plate-tectonics model.

**orogenic cycle** An obsolete concept, linked to geosynclinal theory, that several sequential phases were involved in the formation of an **\*orogenic** belt. Many authors produced variations, but most cycles began with a geosynclinal phase (mainly subsidence and sedimentation), followed by an orogenic phase (mainly compression), and ended with a post-orogenic phase (mainly uplift).


**orogeny (orogenesis)** Mountain building, especially when a belt of the Earth's **\*crust** is compressed by lateral forces to form a chain of mountains. There have been many orogenic episodes in the evolution of the crust, each extending over many millions of years. See **FOLD BELT**; **OROGEN**; **OROGENIC BELT**; **OROGENIC CYCLE**.

**orographic** Applied to the rain or cloud caused by the effects of mountains on air streams that cross them. Orographic cloud and rain are produced by the forced uplift of moist air and the consequent condensation when this is cooled to saturation point.

**Orosirian** A period of the **\*Proterozoic** eon that began 2050 Ma ago and ended 1800 Ma ago, both dates being fixed. The Orosirian followed the **\*Rhyacian** period and was followed by the **\*Statherian** period.

**orpiment** **\*Sulphide**,  $As_2S_3$ ; sp. gr. 3.5; **\*hardness** 2.0; yellow or yellowish-orange; foliated or powdery **\*aggregate**; occurs in low-temperature veins and hot-spring deposits, commonly associated with **\*realgar** and **\*stibnite**.

**Orthida (orthids)** (class **\*Articulata**) Extinct order of brachiopods (**\*Brachiopoda**), which have biconvex shells, a straight **\*hinge** line, and well-developed cardinal areas on both valves (see **CARDINAL TOOTH**). The **\*pedicle** opening notches into both **\*valves** and is not usually restricted by plates. The shells are usually **\*impunctate**. Orthida range from the Lower **\*Cambrian** to the Upper **\*Permian**, e.g. *Orthis* (**\*Ordovician**).



**Orthida**

**orthids** See ORTHIDA.

**orthite** See ALLANITE.

**orthoamphiboles** See AMPHIBOLES.

**orthochemical** 1. The term used in the *Folk* classification to describe the *micrite matrix* and sparry calcite (*sparite*) *cement* in *limestones*. 2. Applied to rocks consisting of micrite without allochemical particles (see ALLOCHEM). 3. Applied to a *carbonate* particle formed by direct chemical precipitation.

**orthoclase** See ALKALI FELDSPAR.

**orthoconglomerate** A *clast-supported conglomerate* containing less than 15% *matrix* between the pebbles. The clasts are bound together mainly by mineral *cement*.

**orthoconic** Applied to the conch of a *cephalopod* when it is a straight, tapering cone.

**orthoferrosilite** See ENSTATITE.

**orthogenesis** Evolutionary trends that remain fairly constant over long periods of time and so appear to lead directly from ancestor organisms to their descendants. This was once explained as the result of some internal directing force or 'need' within the organisms themselves. Such

metaphysical interpretations have been displaced by the concepts of **\*orthoselection** and species selection.

**orthogonal thickness (t)** The thickness of a folded layer, measured perpendicularly to parallel tangents to the inner and outer fold surfaces.


**orthomagmatic** Applied to the stage during which the main mass of **\*silicates** crystallizes from a **\*magma**. The stage can be divided into an early orthomagmatic stage, during which anhydrous silicate minerals crystallize, and a late orthomagmatic stage, during which anhydrous and hydroxyl-bearing silicate minerals crystallize.

**orthophotograph** An aerial photograph in which distortions due to the ground topography, such as elevation and tilt, have been removed, giving the appearance of every object being viewed directly from above, like a map. Distortions are removed by a **\*photogrammetric** system.

**orthopyroxene (opx)** A series of **\*pyroxenes** which crystallize in the **\*orthorhombic** system. They consist of the **\*enstatite** ( $\text{MgSiO}_3$ ) and ferrosilite ( $\text{FeSiO}_3$ ) **\*end-members**, with a number of other minerals between, including **\*bronzite**, **\*hypersthene**, and eulite.

**orthoquartzite** See QUARTZITE.

**orthorhombic** Applied to a **\*crystal** system where the **\*Bravais** lattices have three sets of edges at right angles, but all are of different lengths. They can be referred to three **\*crystallographic axes**, *a*, *b*, and *c* or *x*, *y*, and *z*, of unequal length and at right angles to each other.



**Orthorhombic**

**orthorhombic amphiboles** See AMPHIBOLES.

**orthoscopic** Applied to observations in optical microscopy which use parallel beams of light passing through the atomic structure of a **\*crystal**. This is the normal mode of observation while **\*conoscopic** (i.e. convergent light) observations are used for specialized techniques of **\*mineral** identification.

**orthoselection** Primary selective pressure of a directional kind, which results in a self-perpetuating evolutionary trend. Species selection has been advanced as an alternative explanation for such trends. See also **DOLLO'S LAW**.

**orthosilicates** See NESOSILICATES.

**ortstein** **\*Indurated** **\*soil horizon** in the B horizon of **\*podzols** (**\*spodosols**), in which the **\*cementing** materials are mainly iron oxide and organic matter.

**oryctognosy** See WERNER, ABRAHAM GOTTLÖB.

**Osagean** A **\*series** (348–340 Ma ago) in the **\*Mississippian** of N. America, underlain by the **\*Kinderhookian**, overlain by the **\*Meramecian**, and containing the Ivorian stage of the **\*Tournaisian** series and the Chadian stage of the **\*Visean** series.

**Osborn, Henry Fairfield** (1857–1935) An American palaeontologist who taught at Princeton and Columbia Universities, Osborn was an evolutionary theorist who developed the concept of *\*adaptive* radiation. He also arranged the mammalian palaeontology exhibits at the American Museum of Natural History.

**oscillation ripple (wave ripple mark)** Small ridge of sand formed by wave action on the floor of a sea or lake. Such *\*ripples*, which are usually less than 10 cm in height, commonly display rounded symmetrical troughs and rounded, or sometimes sharp, symmetrical peaks, although many wave-generated ripples are not symmetrical owing to an inequality in forward and backward motion associated with the wave action. The *\*ripple index* of oscillation ripples is usually between 6 and 10.

**oscillatory wave** Wave that causes a mass of water to move to and fro about a point but not to undergo any appreciable net displacement in the direction of wave advance. The wave-form advances, but the individual water particles move in closed or nearly closed orbits.

**oscillatory zoning** See CRYSTAL ZONING.

**osculum** See PORIFERA.

**osmosis** The movement of water or of another solvent from a region of low solute concentration to one of higher concentration through a partially permeable membrane. It is an important mechanism in the uptake of water by plants.

**osmotic potential** See WATER POTENTIAL.

**ossicles** 1. In *\*Echinodermata*, irregularly *\*fenestrated* calcareous plates, rods, and crosses arranged to form a lattice and bound together by connective tissue; together they comprise the skeleton. 2. Small bones.

**Osteichthyes (bony fish)** Superclass of fish in which the internal skeleton is largely ossified, and which also have teeth, plates, or scales of *\*dermal* bone. Characteristic features are: the terminal mouth; *\*homocercal* tail; *\*operculum* covering the gills; usually many flat, bony scales embedded in the skin, with epidermis over them; and the presence of a swim bladder (although sometimes this has been lost in the course of later evolution). With about 28 000 species it is the largest class of *\*vertebrate* animals. The

Osteichthyes have a fossil history going back to the Lower *\*Devonian*; they include the *\*Crossopterygii*, the *\*Dipnoi*, the *\*Chondrostei*, the *\*Holostei*, and the *\*Teleostei*.

**osteostracans** See *OSTEOSTRACI*.

**Osteostraci (Cephalaspida; osteostracans)** (superclass *\*Agnatha*) Order of extinct, fish-like, jawless *\*vertebrates*, ranging from the *\*Silurian* to the *\*Devonian*. They had somewhat flattened bodies, a broad head covered in a bony shield, dorsally located eyes with an opening between them for the pineal eye, and a single median nostril in front. At the sides and back of the *\*head shield* were areas covered in polygonal *\*plates*, possibly covering sense organs. The internal structure of the head resembled that of the extant lampreys. The body was enclosed in bony scales arranged in a series of vertical rows, and there were either one or two dorsal fins and a *\*heterocercal* tail. The internal skeleton was partly ossified. Osteostracans were usually small, around 30 cm in length; the dorsally placed eyes and flat belly suggest a bottom-dwelling way of life. See also *CEPHALASPIS*.

**ostia** See *PORIFERA*.

**Ostracoda (ostracods)** (subphylum *\*Crustacea*) A class of crustaceans that are typical *\*Arthropoda* but laterally compressed and enclosed within a bivalved carapace. This pair of calcareous valves is an integral part of the epidermis and is closed by a series of muscles that leave scars (see *MUSCLE SCAR*) on the valve interiors. The appendages are typically *\*biramous* but may be modified for digging, swimming, etc. The group includes herbivores, carnivores, and scavengers, and occurs in most aquatic habitats. Most are small (less than 1 mm long) and about 70 000 species have been described, occurring from the *\*Cambrian* to the present day. The biological *\*classification* of Recent forms is based on the soft-part anatomy and *\*fossil* forms are classified by the nature of the preserved carapaces. Such characters as the nature of the *\*hinge*, the pattern of the muscle scars, as well as overall shape and ornamentation are all used in species determination. Ostracods have considerable stratigraphic use, and are also used to demonstrate variations in *\*salinity* and fluctuations in the positions of shorelines.

**Ostracodermi (ostracoderms)** The name often used in older textbooks for the fossil, armoured, jawless, agnathan fish of the \*Ordovician to early \*Carboniferous. Probably the name indicates only a grade of development; some ostracoderms are close to the line of ancestry of the living \*Agnatha (lamprey and hagfish), others to the origin of the jawed vertebrates. The informal version of the name survives.

**ostracoderms** See OSTRACODERMI.

**ostracum** See SHELL STRUCTURE; SKELETAL MATERIAL.

**ostrich dinosaur** See ORNITHOMIMID.

**Ostwald's step rule** The observation, first made by the Latvian-born Prussian chemist (Friedrich) Wilhelm Ostwald (1853–1932), that during crystallization from a melt it tends to be the least stable polymorph that crystallizes first.

**Otaian** A \*stage (23.03–21 Ma ago) in the \*Miocene \*epoch of New Zealand, underlain by the \*Waitakian, overlain by the \*Hutchinsonian, and roughly contemporaneous with the \*Aquitainian and lower \*Burdigalian stages.

**Otamitan** A New Zealand \*stage (218–206 Ma ago) of the Late \*Triassic \*epoch, preceded by the \*Oretian and followed by the \*Warepan.

**Otapirian** A New Zealand \*stage (203–199.6 Ma ago) of the Late \*Triassic \*epoch, preceded by the \*Warepan and followed by the \*Aratauran.

**Otariidae** See CARNIVORA.

**Oteke** A \*series in the Upper \*Jurassic of New Zealand, underlain by the \*Kawhia and overlain by the Mokoiwian. It comprises the uppermost Ohauan \*stage and the Puroan stage, which are roughly contemporaneous with the \*Tithonian.

**ottrelite** See CHLORITOID.

**outburst flood** An occasional, large-magnitude flood that is caused by the failure of the side of a lake or a dam containing the lake. It is often associated with the failure of a dam containing a glacial lake.

**outcrop** That part of a rock formation which is exposed at the Earth's surface.

**outerarc** A ridge, or uplifted section, sometimes extending above sea level, found in some **\*arc-trench** gaps.

**outer planet** See JOVIAN PLANET.

**outgassing** 1. The removal of gas, usually by heating. 2. The release of gases by volcanic activity that resulted in the formation of the Earth's atmosphere and hydrosphere.

**outgroup** In **\*phylogenetics**, a species which is the least related to the species under analysis. The inclusion of a known outgroup allows the identification of **\*plesiomorphic** and **\*apomorphic** **\*character** states, which might otherwise remain unclear, a situation that can give rise to **\*topological** errors. A tree that includes an outgroup is said to be rooted.

**outlet glacier** A tongue of ice that extends radially from an **\*ice** dome. It may be identified within the dome as a rapidly moving ribbon of ice (an 'ice stream'), while beyond the dome it typically occupies a shallow, irregular depression. The 700 km long Lambert Glacier, Antarctica, is one of the world's largest outlet glaciers.

**outlier** Area where younger rocks are surrounded completely by older rocks. It may be produced by **\*erosion**, **\*faulting**, or **\*folding**, or any combination of these. Compare INLIER.

**out-of-phase component** See IMAGINARY COMPONENT.

**outwash** 1. The stratified **\*sands** and **\*gravels** deposited at or near to ice margins. 2. Meltwater escaping from the terminal zone of a **\*glacier**. The resulting streams are typically **\*braided**, and show marked seasonal variations in discharge.

**outwash plain (sandur, pl. sandar)** Extensive accumulation of rock debris built up by **\*outwash** in front of a **\*glacier**. Its constituents are very coarse close to the ice, and diminish in size further away. Its surface may be dissected by **\*braided** channels. Fossil outwash plains are found at the margins of many **\*Pleistocene** glaciers. The name 'sandur' is Icelandic.

**outwelling** The enrichment of coastal seas by nutrient-rich estuarine waters.

**overbank deposit** A *\*floodplain* sediment that lies beyond the limits of the river channel and was left by floodwaters that had overflowed the river banks.

**overbreak** Excess material removed by blasting, in *\*excavations*. As rock cannot always be cut exactly to straight lines, more must be removed than is required.

**overburden** **1.** Any loose material which overlies bedrock. **2.** In a sedimentary deposit, the upper strata which cover, compress, and consolidate those beneath. **3.** Any barren material, consolidated or loose, that overlies an *\*ore* deposit. The depth and type of overburden may control whether an ore deposit is worked by underground or *\*open-cast* methods. The proportion of overburden thickness to mineral deposit is called the overburden ratio.

**overburden ratio** See OVERBURDEN.

**over-consolidated clay** Clay that has been more compacted than would be expected from the existing *\*overburden*, e.g. it has been subjected to pressure from overburden that has subsequently been removed by *\*erosion*. Compare NORMALLY CONSOLIDATED CLAY.

**overflow channel (spillway)** Channel cut by meltwater escaping from a *\*proglacial* lake. Generally it is trough shaped in form, lacks tributaries, and is not integrated with the local drainage pattern. It is often difficult to distinguish from other varieties of *\*glacial* drainage channel.

**overflowing well** See ARTESIAN WELL.

**overflow levée** See LAVA LEVE´E.

**overfold (overturned fold)** A *\*fold* in which the *\*axial plane* is inclined so that the *\*fold limbs* *\*dip* in the same direction, although not necessarily by the same amount. One limb is thus ‘overturned’.

**overland flow** See SURFACE RUNOFF.

**overlap** An unconformable relationship in which a transgressive sequence (see TRANSGRESSION) of progressively younger members of an upper series of strata *\*onlap* and rest upon the underlying (oldest) stratum. Strictly, ‘onlap’ refers to the process and ‘overlap’ to the resulting structural

relationship, but the two terms are often used synonymously. The opposite of overlap is ‘\*offlap’. *See also* [OVERSTEP](#).

**overpressure** Overpressured zones occur in subsurface \*horizons in which the fluid pressure is greater than the normal hydrostatic pressure (*see* [HYDROSTATIC STRESS](#)) for the depth in question. Overpressures develop in rapidly deposited \*sediments which have been sealed by impermeable horizons, which prevent the \*pore fluids from escaping. Drilling into overpressured horizons can cause hazardous blowouts and well-caving.

**overspecialization** An old theory which held that straight-line evolution or \*orthogenetic trends might proceed to the point at which the lineage was at an adaptive disadvantage. Overspecialization was therefore considered as one of the causes of extinction. There is no reason to believe, however, that \*natural selection would permit evolution to proceed beyond maximum adaptation. More recently, the term has been applied to highly specialized organisms which have proved incapable of responding to environmental change and so have become extinct.

**overstep** **1.** An unconformable relationship in which a younger series of rocks rests upon progressively older rocks, suggesting the series below the \*unconformity is tilted. *See also* [OFFLAP](#); [OVERLAP](#). **2.** The progressive development of \*hanging-wall \*thrusts.

**overthrust** A \*thrust fault in which the horizontal displacement is large and the \*hanging wall is the relatively active element and is thrust over the \*footwall.

**overturned fold** *See* [OVERFOLD](#).

**overvoltage** *See* [INDUCED POLARIZATION](#).


**Owen, Richard** (1804–92) An anatomist and palaeontologist, Owen worked on fossil mammals and reptiles, including those brought back from South America by \*Darwin. He coined the name ‘\*dinosaur’, and reconstructed fossil reptiles such as \*Iguanodon. He believed that animals within a major group (e.g. vertebrates) were variations on a single theme, or ‘archetype’. His crowning achievement was the founding of the Natural History Museum in South Kensington, London, in 1881.

**oxbow** *See* [CUT-OFF](#).

**Oxfordian** A *stage* in the European Late *Jurassic* (161.2–155.7 Ma ago, Int. Commission on Stratigraphy, 2004), preceded by the *Callovian* and followed by the *Kimmeridgian*. Note, however, that the Oxford Clay (Britain) is predominantly of *Callovian* age. *See also* MALM.

**oxic horizon** Mineral subsoil *soil horizon* that is at least 30 cm thick and is identified by the almost complete absence of weatherable primary minerals, by the presence of *kaolinite* clay, insoluble minerals such as *quartz*, hydrated oxides of iron and aluminium, and small amounts of exchangeable bases, and by low *cation-exchange* capacity. It is the distinguishing subsoil horizon (B horizon) of an *oxisol*.

**oxidation** Specifically, a reaction in which oxygen combines with, or hydrogen is removed from, a substance. More generally, any reaction in which an atom loses *electrons*. For example, in the reaction between zinc and copper sulphate:



the zinc has lost two electrons and been oxidized. Conversely the copper has undergone *reduction*.

**oxidation potential (electrode potential, reduction potential;  $E^\theta$ )** The energy change, measured in volts, required to add or remove *electrons* to or from an element or compound. The reference reaction is the removal of electrons from hydrogen in a standard hydrogen half-cell (i.e.  $\text{H}_2(\text{gas})$  at 1 atm pressure delivered to a 1.0 M solution of  $\text{H}^+$  *ions* at 25 °C, into which a platinum electrode has been inserted):  $\text{H}_2 \rightarrow 2\text{H}^+ + 2\text{e}^-$ , This energy change is given the value of zero. The oxidation potentials of other species are determined relatively by measuring the potential difference between a half-cell containing an aqueous solution of the oxidized and reduced forms of the test substance, and the standard hydrogen half-cell. For example, for  $\text{Fe}^{2+} \rightarrow \text{Fe}^{3+} + \text{e}^-$ ,  $E^\theta = 0.77$ , for  $\text{Mn}^{2+} \rightarrow \text{Mn}^{3+} + \text{e}^-$ ,  $E^\theta = 1.51$ . With decreasing values of oxidation potential, the reduced form of a couple (e.g.  $\text{Fe}^{2+}$ ) will itself reduce the oxidized form of a couple with a higher oxidation potential (e.g.  $\text{Mn}^{3+}$ ). The oxidation potentials obtained under these controlled conditions are called standard electrode potentials, or sometimes standard reduction potentials. *Compare* REDOX POTENTIAL.

**oxidation–reduction** See REDOX REACTION.

**oxides** A group of minerals in which oxygen is combined with one or more metals to give simple and multiple oxides respectively. Simple oxides include **\*hematite** ( $\text{Fe}_2\text{O}_3$ ), **\*rutile** ( $\text{TiO}_2$ ), and zincite ( $\text{ZnO}$ ). Multiple oxides include the **\*spinels** ( $\text{MgAl}_2\text{O}_4$ ) and hydrated oxides, e.g. **\*goethite** ( $\text{FeO}\cdot\text{OH}$ ). Oxides are economically important and are the principal sources of tin ( $\text{SnO}_2$ ), iron ( $\text{Fe}_2\text{O}_3$ ,  $\text{Fe}_3\text{O}_4$ ), chromium ( $\text{FeCr}_2\text{O}_4$ ), titanium ( $\text{TiO}_2$ ), manganese ( $\text{MnO}_2$ ), and aluminium ( $\text{Al}_2\text{O}_3\cdot 2\text{H}_2\text{O}$ ). Oxides are relatively high-temperature minerals occurring in association with a variety of **\*igneous** rocks. They may also form as chemical precipitates in oxidized environments.

**oxisol** In the **\*USDA \*soil taxonomy**, an order of **\*mineral soils**, comprising soils with an **\*oxic** horizon within 2 m of the soil surface, or with **\*plinthite** close to the soil surface, and without a **\*spodic** or **\*argillic** horizon above the oxic horizon.

**oxycone** See INVOLUTE.

**oxygen ‘burning’** A stage of advanced ‘burning’ in stellar evolution, following **\*helium ‘burning’**, that takes place at a temperature around  $2 \times 10^9$  K.  $^{16}\text{O}$  nuclei react to form  $^{32}\text{S}$ ,  $^{31}\text{S}$ ,  $^{31}\text{P}$ , and  $^{28}\text{Si}$ . See CARBON ‘BURNING’; NUCLEOSYNTHESIS.

**oxygen-isotope analysis** Method for estimating past ocean temperatures. The ratio of the stable **\*oxygen isotopes**,  $^{18}\text{O}$  and  $^{16}\text{O}$ , is temperature dependent in water,  $^{18}\text{O}$  increasing as temperature falls. Oxygen incorporated in the calcium-carbonate shells of marine organisms will reflect the prevailing  $^{18}\text{O}:^{16}\text{O}$  ratio (see OXYGEN-ISOTOPE RATIO). Acidification (to release oxygen) of **\*fossils** of these organisms under carefully controlled conditions can therefore be used, with appropriate calibration, to indicate the record of past ocean temperatures.

**oxygen-isotope curve** A graphical plot of values for the relative proportions of two **\*isotopes** of oxygen. Oxygen can exist in several isotope forms, but only  $^{16}\text{O}$  and  $^{18}\text{O}$  are important in the analysis of oxygen isotopes (see OXYGEN-ISOTOPE RATIO). In nature, the present-day average

ratio of  $^{18}\text{O}$  to  $^{16}\text{O}$  is about 1:500, and measurements are made against this value as a standard. There is now evidence that this ratio changed in ocean waters in a cyclical fashion in succeeding glacial and *\*interglacial* periods (see [OXYGEN-ISOTOPE ANALYSIS](#)). At the height of the last (*\*Devensian*) glaciation it appears that deep-ocean waters may have been enriched in  $^{18}\text{O}$  by about 1.6‰. This would be equivalent to a lowering of sea level of about 165 m compared to the present level.

**oxygen-isotope ratio ( $^{18}\text{O}$ : $^{16}\text{O}$  ratio)** The abundance ratio between two of the three *\*isotopes* of oxygen. They have similar chemical properties because they have the same electronic structure, but because of the differences in mass between their nuclei they have different vibrational frequencies which cause them to behave slightly differently in physico-chemical reactions. These differences can provide information, e.g. regarding the source of water in a past environment or the temperature at which various interactions have taken place. For example, surface waters vary in their oxygen isotopes; light water ( $\text{H}_2^{16}\text{O}$ ) has a higher *\*vapour pressure* than  $\text{H}_2^{18}\text{O}$ , and therefore is concentrated by evaporation so that fresh water and polar ice are light but sea water is heavy.  $\text{CaCO}_3$  or  $\text{SiO}_2$  are richer in the heavier isotope when precipitated from sea water than when precipitated from fresh water. Moreover, because of meteorological cycles of evaporation/condensation, there is a steady depletion of  $^{18}\text{O}$  in sea water towards the poles. See also [ISOTOPE FRACTIONATION](#); [OXYGEN-ISOTOPE ANALYSIS](#).

**oxygen isotopes** There are three *\*isotopes* of oxygen,  $^{16}\text{O}$ ,  $^{17}\text{O}$ , and  $^{18}\text{O}$ . The most important in geology are  $^{16}\text{O}$  and  $^{18}\text{O}$ , as both these isotopes are found together in *\*carbonate* rocks and *\*minerals*. The  $^{18}\text{O}$ : $^{16}\text{O}$  ratio (see [OXYGEN-ISOTOPE RATIO](#)) varies with the temperature and chemical composition of the water in which shelly organisms grew, or of the subsurface waters from which diagenetic carbonate *\*cements* crystallized. The  $^{18}\text{O}$ : $^{16}\text{O}$  ratio is therefore a valuable tool for palaeothermometry (see [OXYGEN-ISOTOPE ANALYSIS](#)) and for diagenetic studies (particularly when used together with a study of  $^{13}\text{C}$ : $^{12}\text{C}$  isotope ratios), provided that there have not been subsequent changes to the isotopic composition. See [DIAGENESIS](#).

**oxygen-isotope stage** One of the glacial or *\*interglacial* *\*stages* revealed by *\*oxygen-isotope curves*. Curves from Atlantic and Pacific deep-sea cores were divided by C. Emiliani into 16 stages, their fluctuations being correlated with ice-sheet growth and decay. N. J. Shackleton and N. D. Opdyke extended these subdivisions in 1973, from their studies of a core from the western Pacific in which 23 stages were recognized. These are presumed to represent a continuous record from about 870 000 years BP. In 1976, J. Van Donk obtained curves from a core in the equatorial Atlantic which yielded 42 isotope stages, representing 21 glacial and 21 interglacial stages.

**oxyhornblende** See KAERSUTITE.

**oxyluminescence** See THERMOLUMINESCENCE.

**Oyashio current** Western *\*boundary* current in the subpolar gyre of the N. *\*Pacific*. It originates in the Bering Sea and flows south-west off the Kuril Islands to meet the *\*Kuroshio* current east of northern Japan. The current flows at less than 0.5 m/s, it is cold (4–5 °C at 200 m depth), and has a low *\*salinity* (33.7–34.0‰).

**ozone layer** The atmospheric layer at 15–30 km altitude, in which ozone (O<sub>3</sub>) is concentrated at 1–10 parts per million. Ozone also occurs in very low concentration at altitudes of 10–15 km and 30–50 km. Generally, atmospheric ozone is produced by the photochemical dissociation of oxygen (O<sub>2</sub>), resulting from absorption of ultraviolet solar radiation, to form atoms of oxygen (O). These atoms collide with molecular oxygen (O<sub>2</sub>) to form ozone (O<sub>3</sub>), which in turn absorbs solar radiation for further dissociation to O and O<sub>2</sub>. The ozone layer limits the amount of ultraviolet radiation reaching the ground surface. See also ATMOSPHERIC STRUCTURE.

**ozonesonde** A package of instruments carried by a balloon and used to acquire data from the *\*ozone* layer.

# P

**Pa** See PASCAL.

**Pachycephalosauridae** Family of 'bone-headed' \*dinosaurs of the Upper \*Cretaceous, which were \*ornithopods and apparently may have lived in herds in upland regions.

**pachydont** Applied to a type of \*hinge \*dentition, found in certain cemented bivalves (\*Bivalvia) in which the teeth are heavy, very large, and generally few in number.

**Pacific- and Indian-Ocean common water (PIOCW)** The deep waters of the \*Pacific and \*Indian Oceans are so similar in character that they are usually classed as one mass, with a mean temperature of 1.5 °C and a \*salinity of 34.70‰.

**Pacific-Antarctic Ridge** The \*ridge which lies between the \*Pacific and \*Antarctic Plates and joins the South-East Indian and \*East Pacific Rises.

**Pacific Ocean** The largest of the world's major oceans ( $179.7 \times 10^6$  km<sup>2</sup>). It is also the coldest (average 3.36 °C), deepest (4028 m average), and least saline (34.62‰) of the ocean basins.

**Pacific Plate** Although it is the largest present-day lithospheric \*plate, the Pacific Plate is shrinking because most of its margins, apart from the \*East Pacific Rise and the \*Pacific-Antarctic Ridge, are \*subduction zones (e.g. the Aleutian, Kuril, Japan, Izu-Bonin, and Marianas subduction zones). The Pacific Plate is the only major plate to consist of nothing but \*oceanic crust, parts of which date from the \*Triassic period.

**Pacific Province (American Province)** The name given to the trilobite (\*Trilobita) and graptolite (\*Graptolithina) faunas characteristic of the

northern and western margins of the *\*Iapetus Ocean* in the Early *\*Palaeozoic*. See also ATLANTIC PROVINCE.

**Pacific Ring of Fire** See RING OF FIRE.

**Pacific-type coast (longitudinal-type coast)** A coast that borders or lies within a mountain chain, so its subsidences and fractures follow the grain of the folding. This type of coast was first recognized by *Eduard \*Suess*. Compare ATLANTIC-TYPE COAST; see also DALMATIAN-TYPE COAST.

**Pacific-type margin** See ACTIVE MARGIN.

**packed biomicrite** See FOLK LIMESTONE CLASSIFICATION.

**packer test (Lugeon test)** Test for measuring the *\*permeability* of ground in sections of *\*boreholes*. An inflatable tube ('packer') is lowered down a borehole and expanded so that the sections above and below are isolated. Alternatively, two packers may be used to isolate a certain section. Water is pumped into the section under investigation and leakage can be measured. The rate at which water is absorbed per metre length of hole is measured in units of lugeon, named after the French geologist Maurice Lugeon (1870–1953). One lugeon is approximately equal to  $1.0 \times 10^{-5}$  cm/s permeability.

**packstone** As defined by the *\*Dunham classification*, a *\*limestone* characterized by a *\*grain-supported* texture, together with a *\*lime-mud \*matrix*.

**paedomorphosis** Evolutionary change that results in the retention of juvenile characters into adult life. It may be the result of *\*neoteny* or *\*progenesis*. It permits an 'escape' from *\*specialization*, and has been invoked to account for the origin of many *\*taxa*, from subspecies to phyla.

**pahoehoe** See LAVA.

**Paibian** A *\*stage* (501–496 Ma ago) of the Late *\*Cambrian*, preceded by the Middle Cambrian.

**paired metamorphic belts** Juxtaposed zones of metamorphic rocks; a high pressure–low temperature (*\*blueschist facies*) zone which lies parallel to a low pressure–high temperature (*\*greenschist facies*) zone. Paired metamorphic belts were first recognized in Japan, where the greenschists facies is nearer the *\*subduction zone*, and similar belts in different

**\*Phanerozoic \*orogenic** belts have been identified and are thought to indicate **\*destructive** margins. Recently it has been suggested that the Japanese belts are not contemporaneous and are separated by a major transcurrent fault.

**pakakota** A rock platform on a small island (**\*motu**) in an **\*atoll** that resists erosion. A pakakota is made from **\*conglomerate** or **\*phosphate** rock.

**Palaeartic faunal realm** Region coincident with Europe, Asia north of the Himalayan–Tibetan physical barrier, N. Africa, and much of Arabia. The region is similar at the family level, and rather less so at the generic level, to the **\*Nearctic** faunal realm: the **\*Bering \*land** bridge connected the two for much of the **\*Cenozoic** era.

**palaeo- (paleo-)** From the Greek *palaios* meaning ‘ancient’, a prefix meaning ‘very old’ or ‘ancient’.

**Palaeoarchaean** An **\*era** in the **\*Archaean \*eon**, 3600–3200 Ma ago, preceded by the **\*Eoarchaeon** and followed by the **\*Mesoarchaeon** eras. Fossil **\*stromatolites** and other indirect evidence of life are dated to this time, and it is possible that the first **\*supercontinent**, Vaalbara, formed around 3200 Ma ago, its name derived from the Kaapvaal **\*craton** in southern Africa and the Pilbara craton in Western Australia; palaeomagnetic data indicates that during the Palaeoarchaeon these were close together.

**palaeoautecology** The study of past populations, usually comprising only one or two species.

**palaeobiogeography** The scientific study of the geographic distribution of **\*fossils**. *See also* BIOGEOGRAPHY.

**palaeobiology** The attempt to interpret the biology of **\*fossil** organisms. It is sometimes combined with functional morphological studies in the more general study of **\*palaeoecology**.

**palaeobotany** The study of **\*fossil** plants.

**Palaeocene (Paleocene)** The lowest epoch of the **\*Palaeogene** period, about 65.5–55.8 Ma ago. The name is derived from the Greek *palaios* ‘ancient’, *eos* ‘dawn’, and *kainos* ‘new’, and means ‘the old part of the **\*Eocene**’ (the subsequent epoch).



<http://palaeos.com/cenozoic/paleocene/paleocene.html>

- The Palaeocene Epoch.

**Palaeocene-Eocene thermal maximum (PETM; Initial Eocene Thermal Maximum, IETM)** A *\*climatic optimum* that occurred approximately 55 Ma ago and lasted for less than 100 000 years, at the boundary between the Palaeocene and Eocene epochs. At the commencement of the PETM the global mean temperature rose rapidly by 5–8 °C and the oceans warmed throughout their depths. A second, briefer rise in temperature occurred about 2 million years later. The cause of the rise in temperature is not known.

**palaeoclimatic indicator** One of the sources from which evidence concerning past climates can be obtained. For example, glacial, *\*periglacial*, and pluvial deposits provide morphological information related to climate; cave deposits, dunes, and dunefields yield lithologic information; and plants (including pollen), molluscs (*\*Mollusca*), foraminifera (*\*Foraminiferida*), beetles, and ostracods (*\*Ostracoda*) are among the organisms that have been used to derive biotic information.

**palaeoclimatology** The study of past climates from the traces left behind in the geologic record. It is assumed that the *\*uniformitarian* principle has obtained, but this may not be the case, and the geologic data are always insufficient for palaeoclimatological purposes. Dating methods and *\*palaeogeographic* reconstructions are suspect, and often *\*fossil* faunas and floras cannot be related easily in terms of time. Most glacial *\*strata* can be dated only within wide limits and many climatic criteria leave no mark in the rocks.

**palaeocurrent analysis** The collection, presentation, and interpretation of directional data measured from *\*sedimentary structures* and textures which were formed under flowing water, the wind, or moving ice. Ranging from the study of a single structure to data collected across a whole *\*sedimentary basin*, palaeocurrents can yield a hierarchy of information, from the local direction of *\*flow* which yielded a single *\*ripple train*, through the direction of migration of a *\*bar* or *\*channel*, to the movement of *\*sediment* through a

river or deltaic network (see [DELTA](#)), and the regional pattern of sediment [\\*provenance](#) and dispersal through a basin.

**[Palaeodictyon](#)** A representative of the [\\*Nereites](#) [\\*trace fossil](#) assemblage, associated with deep-water [\\*turbidite](#) facies. *Palaeodictyon* was originally thought to be a complex grazing trace but is now thought to represent a permanent or semi-permanent [\\*burrow](#) system. The burrows form a complex, hexagonal pattern with a single, vertical tube reaching towards the bed surface. It is probable that the organism concerned ‘farmed’ bacteria or algae within the burrows; water currents passed down the vertical tubes to irrigate the burrow system.

**[palaeoecology](#)** The application of ecological concepts to [\\*fossil](#) and sedimentary evidence in order to study the interactions of the Earth’s surface, [\\*atmosphere](#), and [\\*biosphere](#) in pre-historic and geologic times.

**[palaeoflow](#)** The river flow that cut a stream bed that later filled; its direction can be determined from the [\\*cross-stratification](#) it produced.

**[palaeofluminology](#)** The study of ancient stream beds.

**[Palaeogene](#)** The earlier of the two [\\*periods](#) which comprise the [\\*Cenozoic](#) era, preceded by the [\\*Cretaceous](#), followed by the [\\*Neogene](#), and dated at 65.5–23.03 Ma ago (Int. Commission on Stratigraphy, 2004). The Palaeogene period is subdivided into the [\\*Palaeocene](#), [\\*Eocene](#), and [\\*Oligocene](#) [\\*epochs](#).



<http://www.geologypage.com/2014/04/paleogene-period.html>

- The Palaeogene Period.

**[palaeogeography](#)** The reconstruction of the physical geography of past geologic ages. A palaeogeographical map would normally show the [\\*palaeolatitude](#) of the area under discussion together with the location of inferred shorelines, drainage areas, [\\*continental shelves](#) and depositional environments. At the present time the base map would normally be a reconstruction based on palaeomagnetic data (see [PALAEOMAGNETISM](#)), although many maps in earlier publications used the present geographical positions of the continents as a foundation. With the advent of [\\*balanced sections](#) in structural geology, many palaeogeographical maps are now

being produced that take account of the physical shortening involved and attempt to restore areas to their 'depositional' condition.

**palaeoguild** A *\*guild* that existed in the past.

**Palaeohelikian** A sub-*\*stage* of the Lower *\*Helikian* of the Canadian Shield region.

**palaeohydraulics** The analysis of the geometry of ancient *\*fluvial* features, e.g. preserved channel forms, lateral accretion surfaces, and channel-fill sequences, to estimate the hydraulic parameters which were developed at the time of formation of the channel system.

**palaeoichnology** See *ICHOLOGY*.

**palaeolatitude** The position, relative to the equator, of a geographic or geologic feature at some time in the past. The evidence for the position may come from palaeoclimatic data (see *PALAEOCLIMATIC INDICATOR*; *PALAEOCLIMATOLOGY*) or, more usually a palaeomagnetic determination (see *PALAEOMAGNETISM*).

**palaeomagnetic pole** The pole position calculated from palaeomagnetic or *\*archaeomagnetic* directions, assuming an *\*axial* geocentric-dipole model of the Earth's *\*geomagnetic field*. It should be calculated on the mean direction of sufficient observations so that past *\*secular* variations of the geomagnetic field will have been averaged out. See *PALAEOMAGNETISM*.

**palaeomagnetism** Field of *\*geophysics* concerned with the measurement and interpretation of *\*remanent magnetism* or the record of the Earth's past magnetic field (see *GEOMAGNETIC FIELD*). Much valuable information has been obtained concerning *\*polar* wandering and *\*continental drift*.

**palaeoniscids** A group of primitive fish (*\*Actinopterygii*), most of which were of modest size (about the size of a herring), but possessed thick, rhomboidal, enamel scales. There is great variety in structural detail and they were most abundant in the *\*Carboniferous*. The group diminished rapidly after the early *\*Triassic* and finally became extinct in the *\*Cretaceous*.

**palaeontology** The study of *\*fossil \*flora* and *\*fauna*. Information thus gained may be used to establish ancient environments.



<http://palaeos.com/>

- Describes conditions and life throughout Earth history.

**palaeoplacer** A *\*placer* deposit of great antiquity. For example, the Witwatersrand gold placer deposits in South Africa are about 2.7 billion years old.

**Palaeoproterozoic** The earliest part of the *\*Proterozoic*, lasting from about 2500–1600 Ma.

**Palaeopterygii** Name formerly given to a supposed subclass that included some of the more primitive fish, but whose members are no longer held to be closely related. They were *\*bony* fish with many fossil relatives, including, for example, the still-extant families Acipenseridae (sturgeon), Polyodontidae (paddlefish), and Polypteridae (bichir).

**palaeoseismology** The study of ancient *\*earthquakes*, primarily by mapping zones of previous earthquake destruction by observation or from historical records.

**palaeosol (paleosol)** Soil formed during an earlier period of *\*pedogenesis*, and which may have been buried, buried and exhumed, or continuously present on the landscape until the current period of pedogenesis.

**palaeosome** That part of a composite rock body which appears to be older than an associated element. The term is commonly used with reference to that part of a *\*migmatite* which remained solid during *\*anatexis*.

**palaeospecies** Group of biological organisms, known only from *\*fossils*, which differs in some respect from all other groups.

**palaeosynecology** The study of past plant and animal communities, including all the species comprising them.

**palaeotaxodont** A type of *\*taxodont* dentition in bivalves (*\*Bivalvia*). In early *\*Palaeozoic* forms teeth are formed in two distinct sizes, an anterior row with large teeth and a posterior row with much smaller teeth.

***Palaeotaxus rediviva*** The earliest recorded species of the Taxales (yew) family. Found in the Lower *\*Jurassic* of Europe, it is distinguished by the form of the leaf and the structure of the *\*cuticle*. The Taxales, unlike the

conifers, have a single ovule surrounded by a berry-like structure. They bear terminal seeds. *See also* CONIFERALES.

**Palaeotethys** There is confusion over the relationship between *\*Tethys* (or Neotethys) and Palaeotethys. Some authors refer to the existence of a late *\*Palaeozoic* Palaeotethys which was a gulf at the margin of *\*Panthalassa* between the parts of *\*Pangaea* that were later to separate as *\*Laurasia* and *\*Gondwana*. This ocean, or large parts of it, were consumed in a Palaeocimmerian-Indosinian *\*subduction* zone. In the *\*Jurassic* a new ocean, or oceans, opened along the northern margin of Palaeotethys. This Jurassic-*\*Cretaceous* ocean has been called Neotethys or Tethys (*sensu stricto*).

**Palaeozoic** The first (542–251 Ma ago) of the three *\*eras* of the *\*Phanerozoic*. The *\*Cambrian*, *\*Ordovician*, and *\*Silurian* *\*periods* together form the Lower Palaeozoic sub-era; the *\*Devonian*, *\*Carboniferous*, and *\*Permian* the Upper Palaeozoic sub-era. During the Palaeozoic, two major *\*orogenies* occurred: the *\*Caledonian* during the Lower Palaeozoic, and the *\*Hercynian* in late Palaeozoic times. The faunas of the Palaeozoic are noted for the presence of many invertebrate organisms, including trilobites (*\*Trilobita*), graptolites (*\*Graptolithina*), brachiopods (*\*Brachiopoda*), cephalopods (*\*Cephalopoda*), and *\*corals*. By the end of the era, *\*amphibians* and *\*reptiles* were major components of various communities and giant tree-ferns, horsetails, and *\*cycads* gave rise to extensive forests.



<http://www.geologypage.com/2013/10/paleozoic-era.html>

- Palaeozoic Era.

**palagonite** Pale yellow to yellow-brown, hydrated, fragmental, basaltic *\*glass*, formed by the *\*hydration* of *\*metastable* *\*hyaloclastite* and *\*ion exchange*, due to the action of *\*groundwater* percolating in the subaerial environment or to the action of sea water in the case of submarine environments. The high surface area to volume ratio of the *\*lapilli*-size fragments of glass facilitates efficient chemical exchange with water. Much of the original calcium, potassium, and sodium, and some of the aluminium and silicon, are lost, iron is oxidized, and water is gained. Elements such as

iron, titanium, magnesium, and aluminium may be either lost or added. The type locality is at Palagonia, Sicily. Some planetary scientists think that the whole surface of *\*Mars* is covered with dust of palagonite origin.

**palate** The roof of the mouth; it separates the nasal and mouth cavities.

**paleo-** Alternative spelling for the prefix *\*palaeo-*.

**Paleocene** See PALAEOCENE.

**palimpsest** A landscape that bears the imprint of two or more sets of geomorphological *\*processes*. For example, much of the Sahel region of Africa shows land-forms resulting from former wet and dry episodes. The word is derived from the Greek *palimpsestos*, 'to rub smooth again', and is also used for a re-used parchment, paper, or ornamental brass whose original writing or engraving has been only partially erased and for a sedimentary structure in which one set of *\*trace fossils* has been partly overwritten by another. See RELAXATION.

**palingenesis** The partial or complete melting of pre-existing rocks to generate new *\*magma*.

**palinspastic map** A map on which the depiction of folded or faulted strata shows them restored as closely as possible to their original geographical positions prior to folding or faulting. *\*Balanced* sections are used to ensure the palinspastic reconstruction accounts for the entire volume of the materials represented and respects the lengths of lines and the thicknesses of individual layers. The term 'palinspastic' is derived from the Greek *palin* meaning 'again', and *spastikos* meaning 'pulling'.

**Pallas** The second largest (after *\*Ceres*) *\*solar system* asteroid (No. 2), diameter 538 km; approximate mass  $25 \times 10^{19}$  kg; rotation period 7.811 hours; orbital period 4.61 years. It was discovered in 1802 by Wilhelm Olbers. It is possible that Pallas will be reclassified as a minor planet.

**pallasite** A striking and unusual kind of *\*meteorite* in which *\*olivine* and *\*iron* are intergrown in roughly equal proportions; these meteorites are thus intermediate between *\*stony* and *\*iron meteorites*. Only 35 examples are known.

**pallial line** See MUSCLE SCAR.

**pallial sinus** See MUSCLE SCAR.

**pallium** See MANTLE.

**Palmer method** See GENERALIZED RECIPROCAL METHOD.

**palsa** Mound or ridge, largely made from \*peat, containing a perennial ice lens, and found in the damper sites of bogs in \*periglacial areas. Widths are in the range 10–30 m, lengths 15–150 m, and heights 1–7 m. Probably palsas are a result of heaving associated with the growth of segregated ice. See also PINGO.

**paludal** Applied to organisms, soils, etc., that are of or associated with a marsh.

**palus** From the Latin *palus* meaning ‘marsh’, a term introduced by Giovanni B. Riccioli in 1651 for small patches of lunar \*mare \*basalt. The Apollo 15 mission landed in Palus Putredinis, the ‘Marsh of Decay’.

**palustrine** Pertaining to a marsh.

**palynology** The study of living and \*fossil \*pollen grains, \*spores, and certain other \*microfossils (e.g. \*dinoflagellates and \*coccolithophorids). Palynology was developed from \*pollen analysis and deals principally with structure, classification, and distribution. It has many applications, e.g. in medicine, archaeology, petroleum exploration, and \*palaeoclimatology.

**palynomorph** A \*microfossil, 5–500 µm in size, made from an organic substance that is highly resistant to chemical attack.

**pampero** Regional storm of the \*line-squall type which affects the Pampas of Argentina and Uruguay. Sometimes the storm brings rain, thunder, and lightning, and its passage is marked by a fall in temperature. The storm moves ahead of a south-westerly wind, which follows a \*depression.

**Pan (Saturn XVIII)** One of the minor satellites of \*Saturn, with a radius of 10 km; visual albedo 0.5.

**pan (clay pan)** \*Soil horizon, usually in the subsoil, that is strongly compacted, \*indurated, \*cemented, or very high in \*clay content.

**Panama Isthmus** The narrow neck of land that connects N. America and Mexico with S. America. During the \*Mesozoic there was an open marine

connection between the **\*Atlantic** and **\*Pacific Oceans**. At the culmination of the Andean–**\*Laramide** orogeny (Late **\*Cretaceous**) a temporary connection between N. and S. America was formed, which explains the similarity between the floras of California and S. America at that time. Some early mammals also migrated southwards across the isthmus. During the early **\*Cenozoic** the land connection was broken but it became re-established in the **\*Pliocene**, thereby allowing another 14 mammal families to migrate from north to south. Some families, notably the elephants (see **PROBOSCIDEA**), appeared to be unable to migrate southwards, perhaps being constrained by the climatic zones that had to be traversed in such a migration.

**Pancrustacea** A **\*clade** that includes the **\*Crustacea** and **\*Hexapoda**.

**pandemic distribution** See **COSMOPOLITAN DISTRIBUTION**.

**Pandora (Saturn XVII)** One of the lesser satellites of **\*Saturn**, discovered in 1980 by **\*Voyager 1**, with a radius measuring  $55 \times 44 \times 31$  km; mass  $0.0013 \times 10^{20}$  kg; mean density  $420 \text{ kg/m}^3$ ; visual albedo 0.9.

**Pangaea** A single supercontinent which came into being in late **\*Permian** times and persisted for about 40 million years before it began to break up at the end of the **\*Triassic** period; or, in the view of some people, which existed throughout most of the Earth's history prior to the Triassic. It was surrounded by the universal ocean of **\*Panthalassa**.

**panhole (gnamma, kamenitza, solution pan, weathering pit)** A small depression or basin on the surface of **\*limestone** or **\*silicate** rock, and less commonly on **\*granite**, that is caused by weathering. Panholes are rarely more than 15 cm across, commonly have flat bottoms, and are sometimes wider at the base than at the top.

**panidiomorphic** Applied to a rock that has well-developed crystals; i.e. it is wholly or predominantly **\*euhedral** (idiomorphic).

**panning** The concentration of heavy minerals by hand in a shallow pan. Loose material is shaken backward and forward in water; heavy minerals settle and the **\*gangue** is washed out. It is a simple method for testing in the field.

**Pannotia** A \*supercontinent that existed during the \*Neoproterozoic era (1000–542 Ma ago). Early in the Neoproterozoic the supercontinent \*Rodinia began to break apart and sections of \*cratons reassembled to form Pannotia, at the same time opening a sea that would eventually become the Pacific Ocean. Pannotia occupied much of the southern hemisphere, with a narrow arm of the sea separating a section that included East \*Gondwana and another section that included \*Laurasia. The supercontinent began to break up about 250 Ma later.

**pannus** From the Latin *pannus* meaning ‘shred’, an accessory cloud term applied to ragged cloud either beneath or attached to another cloud, usually \*cumulonimbus, \*cumulus, \*nimbostratus, or \*altostratus. *See also* CLOUD CLASSIFICATION.

**panplain (panplane, planplain)** Area of very subdued relief that consists of coalesced \*floodplains. It is therefore due to lateral stream migration and is a component of a \*peneplain. Good examples are found in the Carpentaria region of Australia.

**panplane** *See* PANPLAIN.

**Panthalassa** The name given to the vast oceanic area that surrounded \*Pangaea when that supercontinent was in existence. \*Tethys was a minor arm of this ocean. Once Pangaea began to rift in the \*Triassic, the names of the modern oceans are normally applied to the developing ocean basins even though they were, at that time, still very small.

**Pantotheria** (class \*Mammalia, subclass Theria) Extinct infraclass of primitive Middle and Upper \*Jurassic mammals, known from N. America, Europe, and E. Africa. They were egg-laying insectivores of shrew-like appearance, and are believed, on the basis of the structure of the jaw and teeth, to include the ancestors of all later placental and \*marsupial mammalian groups. By the later Jurassic, they were widely distributed, numerous, and varied in form.

**papagayo** *See* NORTE.

**paper shale** A dark grey to black \*shale composed of thin, parallel laminae that tend to separate on \*weathering into tough, slightly flexible sheets reminiscent of sheets of paper.

**para-** From the Greek *para* meaning ‘beside’ or ‘beyond’, a prefix meaning ‘beside’ or ‘beyond’.

**Parablastoidea** (subphylum *Pelmatozoa*) A class of echinoderms (*Echinodermata*) restricted to the Lower and Middle *Ordovician*, which had *biserial brachioles*, well-developed *pentamerous symmetry*, and *ambulacra* composed of *biserial* plates.

**parabolic dune** See *DUNE*.

**paraclade** A group of *evolutionary lineages*; paraclades may be *paraphyletic* or *monophyletic*, but not *polyphyletic*.

**paraconglomerate** A *conglomerate* in which the pebbles are supported by a fine-grained *matrix*. The matrix constitutes more than 15% of the rock.

**Paracrinoidea** (subphylum *Pelmatozoa*) Class of echinoderms (*Echinodermata*) with *uniserial brachioles*, and a box-like *theca* composed of numerous *plates* tending to *bilateral symmetry*. They are restricted to the Middle and Upper *Ordovician*.

**paracycle** In the *genetic stratigraphic sequence model* used in *sequence stratigraphy*, a sequence of changes in sea level, lasting about 2 million years, and forming part of a hierarchy of sedimentation cycles.

**paradigm** Essentially, a large-scale and generalized *model* that provides a viewpoint from which the real world may be investigated. It differs from most other models, which are abstractions based on data derived from the real world.

**paragenesis** From the Greek *para* (‘beside’) and *gen* (‘be produced’), a particular assemblage of *minerals* all of which formed at the same time. See also *PARAGENETIC SEQUENCE*.

**paragenetic sequence** The chronological order of crystallization of minerals in a rock or ore deposit. See also *ORE GENESIS*; *PARAGENESIS*.

**paragonite** See *MUSCOVITE*.

**paralic** Applied to sediments deposited in a shallow freshwater basin on the landward side of a coast that is periodically inundated by the sea.

**parallax** The apparent change in position of an object in relation to another when the viewpoint is changed. *See also* STEREOPTIC VISION.

**parallel evolution** Similar evolutionary development that occurs in lineages of common ancestry. Thus the descendants are as alike as were their ancestors. The nature of the ancestry imposes or directly influences the development of the parallelism.

**parallel extinction** *See* STRAIGHT EXTINCTION.

**parallel fold** A *\*fold* in which the *\*orthogonal* thickness of layers remains constant throughout, so that adjacent, bounding, fold surfaces are parallel. Because of their geometry, parallel folds cannot maintain their profiles over long distances; their form changes along the *\*axial plane*.

**parallel twins** A twinned crystal (*see* CRYSTAL TWINNING) where the twin axis lies in the *\*composition plane* parallel to a *\*zone axis*, e.g. a *\*Carlsbad twin* in *\*orthoclase* feldspar.

**paramagnetism** Magnetization developed in atoms or *\*ions* that have permanent intrinsic magnetic moments, in the same direction as an applied field. It is usually somewhat larger than that resulting from the *\*diamagnetism* of the material, and may have *\*ferromagnetism* superimposed upon it.

**parameter (intercept ratio)** In *\*crystallography*, the ratio of the intercepts made by a plane (the parametral plane), parallel to a *\*crystal face*, which intersects the *\*crystallographic axes*, and which has been chosen to define a unit length of intersection along each axis. The *\*form* of which the face is a member is called the ‘unit form’, ‘fundamental form’, or ‘parametral form’. *See also* AXIAL RATIO.

**parametral form** *See* PARAMETER.

**parametral plane** *See* PARAMETER.

**paramorph** A crystal formed by the conversion of one *\*mineral* polymorph (*see* POLYMORPHISM) to another. The polymorphs of silica provide a good example. If the high-temperature form, *\*cristobalite*, converts to a lower-temperature form, e.g. *\*tridymite*, the tridymite would form a paramorph.

**parapatric** Applied to species whose habitats are separate but adjoining. Compare ALLOPATRIC; SYMPATRIC.

**paraphyletic** Of a \*taxon, including some but not all descendants of the common ancestor (i.e. not \*holophyletic).

**Parapithecidae** (order Primates, suborder Simiiformes) An extinct family of primates which lived during the \*Eocene and \*Oligocene in Egypt; Eocene fossils from Burma are sometimes included in the family in addition. They showed certain similarities in dentition to \*Condylarthra, but had short faces and jaws shaped like those of tarsiers.

**parasequence** In the \*genetic stratigraphic sequence model used in \*sequence stratigraphy, a relatively conformable sequence of genetically related beds bounded by surfaces formed by marine flooding. A succession of genetically related parasequences forms a parasequence set.

**parasequence set** See PARASEQUENCE.

**parasitic cone (adventive cone)** A conical mound of ejecta accumulated around an eruptive vent on the lower flanks of a large \*volcano. Parasitic cones sometimes grow into large volcanic centres themselves, and may lie on the line of a fissure which radiates a great distance from the main volcanic conduit.

**parasitic fold** A \*fold of small \*wavelength and \*amplitude which usually occurs in a systematic form superimposed on folds of larger wavelength. Parasitic folds usually show typical S and Z asymmetric profiles on their limbs and M profiles in the \*hinge regions.

**parasitic magnetization** \*Ferromagnetic behaviour resulting from the imperfect cancellation of antiparallel magnetic lattices in an \*antiferromagnetic substance. \*Hematite is a mineral exhibiting this magnetic behaviour.

**PARASOL** One of the five spacecraft in the \*A-Train, launched in December 2004 by the Centre National d'Études Spatiales (CNES) of France to measure \*aerosols on what was planned as a two-year mission. It flew within about 30 seconds of the \*CALIPSO and \*CloudSat spacecraft until 2 December 2009, when CNES began to manoeuvre it out of the A-Train; it left the A-Train altogether at the end of 2012. Its place was to have

been taken by Glory, but that mission failed at launch and was lost. The satellite was shut down on 18 December 2013.



[https://parasol.cnes.fr/en/PARASOL/GP\\_mission.htm](https://parasol.cnes.fr/en/PARASOL/GP_mission.htm)

- Mission.

**parastratotype** An additional stratigraphic section, designated and described at the time of the establishment of a stratigraphic unit, to augment the definition given by the principal *\*stratotype* (the *\*holostratotype*). The parastratotype is usually selected from within the *\*type area*. *Compare* HYPOSTRATOTYPE. *See also* LECTOSTRATOTYPE; NEOSTRATOTYPE.

**parataxon** Artificial classification suggested for certain common organisms of doubtful affinities, or as yet unknown origins, e.g. *\*fossil \*spores*, *\*dinosaur* footprints.

**Paratethys (Central European Sea)** A large, arcuate seaway that, at its maximum development, extended a distance of 4500 km from just north of the Alps to just east of the Aral Sea. By the end of the *\*Oligocene* it had been separated from the Boreal Sea by closure of the Ural, Polish, and Alsace Straits. Towards the end of the *\*Miocene* it became more lagoonal in character and by the *\*Pliocene* was represented only by a series of land-locked lakes: Lake Balaton in Hungary, the Black Sea (rejoined to the Mediterranean Sea by *\*Quaternary* faulting), the Caspian Sea, and the Aral Sea.

**paratype** Specimen, other than a *\*holotype*, used as the 'type' material by an author at the time of the original description, and designated as such by the author. *Compare* LECTOTYPE; NEOTYPE.

**parcel of air** Quantity of air with more or less uniform characteristics.

**Pareiasauridae** The largest (up to 3 m in length) of the stem reptiles (*\*cotylosaurs*), the pareiasaurs are found in Middle and Upper *\*Permian* rocks in Europe and Africa. They were herbivorous, and perhaps lived semi-aquatically in swamps. They were distinctive among cotylosaurs in that the limbs were rotated in towards the body, thus supporting the body in a more upright fashion.

**parent** See RADIOACTIVE DECAY; RADIOMETRIC DATING.

**parent material** The original material from which the **\*soil profile** has developed through **\*pedogenesis**, usually to be found at the base of the profile as weathered but otherwise unaltered mineral or organic material.

**pargasite** See HORNBLLENDE.

**parivincular** In **\*Bivalvia**, applied to a type of ligament that is elongate and cylindrical in shape. It occurs in a position posterior to the **\*shell beaks**.

**parna** An Australian term describing a silty-clay, carbonate-rich mantle that is deposited evenly over much of the plain of south-eastern Australia and elsewhere in the country. It is similar to **\*loess**, but derived from clay pellets and silt transported downwind from dry lakebeds and **\*alluvial** plains rather than resulting from glacial activity.

**parsimony** In **\*cladistic analysis**, the convention whereby the simplest explanation is preferred, because it requires the fewest conjectures, although the most parsimonious explanation is not always the correct one.

**partial annealing** In **\*fission-track dating**, a variation in the length of fission tracks that results from the sealing of tracks in a mineral that has remained for some time at just below its **\*closure temperature** (typically at 80–125 °C).

**partial melting** Incomplete melting of parent rock, characteristically producing a **\*melt** whose chemical composition differs from that of the parent material. It is thought that partial-melting processes play a major role in generating more-defined liquids from less-evolved ones, so that many **\*basalts** may be the result of partial melting in the (**\*ultrabasic**) upper **\*mantle**, and many **\*granites** may have derived partly or completely from the partial melting of **\*continental crust** (**\*anatexis**). Partial melting preferentially enriches melts with **\*incompatible elements**. In a **\*subduction zone**, rocks of **\*intermediate** composition may form (e.g. **\*andesites**). With increasing temperature and pressure, the subducted **\*oceanic crust** (of **\*basic** composition) first undergoes **\*metamorphism** and then begins to melt or release watery fluids; this material rises into the overlying mantle, which may also begin to melt, giving rise to intermediate **\*magma**.

**partial pressure** In a mixture of gases, the partial pressure of any one is the pressure it would exert alone in the same space as that occupied by the whole. The interaction of the partial pressures of each of the gases equals the total pressure of the mixture.

**partial range zone** A body of *\*strata* containing the documented lowest occurrence of one taxon and the documented highest occurrence of another taxon, but with no stratigraphic overlap of the taxa. *See also* [INTERVAL ZONE](#). *Compare* [CONCURRENT RANGE ZONE](#).

**particle density** The mass per unit volume of soil particles, usually expressed in grams per cubic centimetre. *Compare* [BULK DENSITY](#).

**particle shape (grain shape)** Particle shape is defined by the relative dimensions of the long, intermediate, and short axes of the particle. The ratio of intermediate to long diameter and the ratio of short to intermediate diameter is used to define four shape classes. These are: *\*oblate* (tabular or disc-shaped forms); *\*prolate* (rod-shaped); *\*bladed*; and *\*equant* (cubic or spherical forms).

**particle size (grain size)** The diameter or volume of the *\*grains* in a *\*sediment* or *\*sedimentary rock*. The particle size can be determined by sieving, by measuring the settling velocity, or (for pebbles, boulders, and cobbles) by direct measurement of individual *\*clasts*. The smaller particles are normally defined by their volume diameters, i.e. the diameter of a sphere with the same volume as the particle. There are numerous particle-size classifications. Two commonly used ones are the Udden–Wentworth scale (Wentworth scale) and the British standard classification. In the Udden–Wentworth scale the sediment *\*grades* are: more than 256 mm, boulder; 64–256 mm, cobble; 2–64 mm, pebble; 62.5–2000 µm, sand; 4–62.5 µm, silt; less than 4 µm, clay. In the British classification: more than 200 mm, boulder; 60–200 mm, cobble; 2–60 mm, gravel; 600–2000 µm, coarse sand; 200–600 µm, medium sand; 60–200 µm, fine sand; 2–60 µm, silt; less than 2 µm, clay. *See* [PHI SCALE](#).

**particle velocity** The velocity of a particle in a medium affected by the propagation of a seismic wave. It depends on the amplitude of the wave and for weak seismic events may be of the order of  $10^{-8}$ m/s. *Compare* [SEISMIC VELOCITY](#).

**parting lineation** A series of parallel, linear, low-relief ridges and hollows, spaced a few millimetres apart, seen on **\*bedding planes** in parallel-laminated **\*sandstones**. The **\*lineation** is the preserved trace of primary current lineations formed when **\*plane-bedded** sand is deposited from a high-velocity and usually shallow flow of water. The lineation is oriented parallel to the flow direction. See **BEDFORM**.

**partition coefficient (distribution coefficient)** **1.** If a substance is dissolved in two immiscible liquids standing in contact with each other, the substance will partition or distribute itself between them in a constant ratio, called the partition coefficient. The value of this constant is dependent on the temperature, and on the identities of the solute and the solvents. (The number of solute molecules in solvent A divided by the number of solute molecules in solvent B is a constant.) **2.** The ratio of the concentration (by weight) of an element (e.g. Ti) in a crystallizing **\*mineral** to its concentration in the **\*magma**. For example,  $k_{\text{Ti}} = [\text{Ti}]_{\text{min.}} / [\text{Ti}]_{\text{magma}}$ , where  $k_{\text{Ti}}$  is the partition coefficient for Ti, and  $[\text{Ti}]_{\text{min.}}$  and  $[\text{Ti}]_{\text{magma}}$  are the concentrations of Ti in the mineral and magma respectively. The value of  $k$  is dependent on temperature, pressure and the composition of the crystallizing mineral and magma.

**pascal** **1. (Pa)** The derived SI unit of pressure, equal to 1 N/m<sup>2</sup>. **2.** A high-level computer programming language. Both are named after the French mathematician Blaise Pascal (1623–62).

**pascichnia** One of five groups of **\*trace fossils** established in a behavioural (ethological) classification by A. Seilacher (1953). Grazing traces (pascichnia) result from the distinctive behavioural pattern of an animal during feeding. **\*Deposit feeders**, e.g. **\*Nereites**, leave a recognizable trace through the exploitation of **\*sediment** for nutrition.

**Pasiphae (Jupiter VIII)** One of the lesser satellites of **\*Jupiter**, with a diameter of 36 km; its orbit is **\*retrograde**.

**pass band** The range of frequencies that can be passed through a band-pass filter with negligible **\*attenuation**. See **BAND FILTER**.

**passive margin (trailing edge)** A continental margin which is not also a **\*plate margin**. Such margins are also known as ‘aseismic margins’ or ‘Atlantic-type margins’ and are contrasted with **\*active margins**. Passive

margins are characterized by rifted and rotated blocks of usually thick sedimentary sequences. These rocks are often highly prospective for oil and gas, with a variety of traps, including those related to the diapiric (see [DIAPIRISM](#)) rise of the rock salt formed during the initial separation of the continents.

**passive margin basin (Afrotype drainage)** A continental [\\*drainage](#) system in which water moves from a cratonic (see [CRATON](#)) interior with no mountain belt younger than [\\*Mesozoic](#) and into a [\\*passive margin](#). This type of drainage system is common in Africa, hence its alternative name; the Zaire River system is an example.

**passive margin-distal orogen basin (Amerotype drainage)** A continental [\\*drainage](#) system in which water moves to a [\\*passive margin](#) from a mountain belt of [\\*Mesozoic](#) or [\\*Cenozoic](#) age on the [\\*distal](#) side of the basin; the Amazon River system is an example.

**passive microwave** [\\*Electromagnetic radiation](#) with a wavelength between 1 mm and 1 m, which is emitted by all objects at temperatures higher than [\\*absolute zero](#).

**passive remote sensing** [\\*Remote sensing](#) which is based on the illumination of a scene by [\\*electromagnetic radiation](#) from a natural source. An example is photography. *Compare* [ACTIVE REMOTE SENSING](#).

**Pasteur effect** The transition from an anaerobic to an [\\*aerobic](#) life-style, which occurs among certain organisms when the oxygen content of the atmosphere is 1% of that of the present day. The critical point of transition is the 'Pasteur point'. The gradual oxygen enrichment of the Earth's atmosphere during the [\\*Precambrian](#) passed through the Pasteur point approximately 700 Ma ago, resulting in a general transition to an aerobic life-style.

**Pasteur point** *See* [PASTEUR EFFECT](#).

**Pastonian** A Middle [\\*Pleistocene](#) [\\*stage](#) represented by estuarine silts and fresh-water peat, revealed as marine clays in the [\\*borehole](#) at Ludham, Norfolk, England. *See also* [ANTIAN](#); [BAVENTIAN](#); [LUDHAMIAN](#); [THURNIAN](#).

**patch** *See* [GROUP](#).

**patch reef** See REEF.

**‘patella’ beach** Old term referring to a raised beach standing about 3 m above the present high-water mark and found locally on the south coast of England, in southern and western Ireland, and along the Channel coast of France. Traditionally it was given an *\*interglacial age*, but the term is now little used, owing to problems of correlations. It is named after the large number of limpets (*Patella vulgata* is the common limpet) it typically contains.

**patellate** See SOLITARY CORALS.

*Patella vulgata* See ARCHAEOGASTROPODA; ‘PATELLA’ BEACH.

**patera** An irregular or complex martian structure, with scalloped edges, usually a volcanic *\*caldera*. The classic example is Alba Patera (110° W, 40° N), the volcano of largest areal extent on Mars, but only 3 km high. On the jovian satellite Io, the term is used for a volcanic crater surrounded by irregular flows. The name is derived from a type of shallow, ceremonial dish used by the Romans.

**paternoster lake** Body of water in a formerly glaciated environment that is aligned with neighbouring lakes, so that it looks like a paternoster in a rosary. It is caused by irregular glacial scouring along a zone of weakness.

**patterned ground** An assemblage of small-scale, geometric features typically found at the surface of a *\*regolith* that has been disturbed by frost action. The group includes circles, polygons, and nets, which normally occur on level or gently sloping surfaces, and steps and stripes which are found on steeper gradients. Both sorted and non-sorted varieties are recognized. The sorted varieties are typically outlined by coarse, stony material, and so are termed ‘stone circles’, ‘stone polygons’, ‘stone nets’, ‘stone steps’, and ‘stone stripes’. The origin of patterned ground involves a complex interaction of several geomorphological *\*processes*, including ground cracking, frost sorting, *\*frost heaving*, and *\*mass movement*. The ‘ice-wedge polygon’ is an important member. It is usually 15–30 m in diameter and bounded by *\*ice wedges* up to 3 m wide and about 10 m deep which occupy contraction cracks that form under very low temperatures. The wedges define raised zones (when freezing is active) or depressions (due to thaw). The ‘stone garland’ is a variety of sorted step, which ends in

a stony riser (less than 1 m high) supporting a relatively bare tread (less than 8 m long) upslope. It is found on gradients of 5–15° (in Alaska) and may be due to a combination of **\*frost pull and frost push**, which heave stones to the surface, and mass movement. Patterned ground may also be found in areas underlain by **\*montmorillonitic** soils experiencing markedly seasonal rainfall, where the microrelief forms are called **\*gilgai**.

**Pauling's rules** Rules relating to chemical bonding based on certain regular features shown by ionically bonded, usually simple solids, and first summarized by L. Pauling (1960).

**Pauropoda** See MYRIAPODA.

**pavement** **1.** In geology, bare rock surface resembling a road, e.g. limestone pavement. **2.** The floor of a coal seam. **3.** In roadway construction, any material spread on the subgrade to distribute load and protect against erosion and traffic wear. The types of material used and thickness depend on the type and use of road. Generally the pavement has four layers. (a) The wearing course is the top layer of the carriageway. It must be durable, impermeable, skid-resistant, and resistant to polishing. On the most heavily trafficked roads the polished stone value (PSV) (see **AGGREGATE TESTS**) must exceed 60%. The wearing course lies on a surface provided by the basecourse. (b) The basecourse is a layer of gravel or crushed rock (**\*aggregate**) of specific dimensions to provide drainage, distribute load on to the road base, and protect against freezing. (c) The road base provides the main load-bearing foundation or ballast and consists of irregular-sized rock aggregate, either bonded or loose. (d) The subbase is a layer of coarse aggregate below the road base. It provides extra support or drainage. In cold climates large pore spaces are necessary to prevent water rising by capillary action, leading to the collapse of the structure after thawing. (e) The subgrade is the rock or subsoil on which the subbase is laid.

**Payntonian** A **\*stage** (491–488.3 Ma ago) of the Upper **\*Cambrian** of Australia, underlain by the Pre-Payntonian and overlain by the **\*Datsonian** (**\*Ordovician**).

**paystreak** Profitable part of a **\*mineral** deposit. In **\*alluvial** **\*gold** mining, the term often refers to pockets of gold concentrated at or near bedrock.

**P-band** The radar frequency band between 225 and 390 MHz. It is commonly used in *\*remote sensing* because of its good vegetation penetration.

**PBRs** See PERIODIC BEDROCK RIDGES.

**PcP-wave** See SEISMIC-WAVE MODES.

**PE** See POTENTIAL EVAPOTRANSPIRATION.

**peacock ore** See BORNITE.

**peak shear strength** See SHEAR STRENGTH.

**peak zone** See ACME ZONE.

**pearl-necklace lightning (chain lightning, beaded lightning)** Infrequent type of discharge, in which variations in luminosity resemble pearls on a string.

**pearlspar** See DOLOMITE.

**pearly** Of *\*mineral \*lustre*, milky, translucent, like a pearl.

**peat** An organic soil or deposit; in Britain, a soil with an organic *\*soil horizon* at least 40 cm thick. Peat formation occurs when decomposition is slow owing to *\*anaerobic* conditions associated with waterlogging. Decomposition of cellulose and hemicellulose is particularly slow for *Sphagnum* plants, which are characteristic of such sites, and hence among the principal peat-forming plants. Fen and bog peats differ considerably. In fen peats the presence of calcium in the groundwater neutralizes acidity, often leading to the disappearance of plant structure, giving a black, structureless peat. Bog peats, formed in much more acidic waters, vary according to the main plants involved. Species identification of constituents (including animals as well as plants) remains possible after long periods. Recent bog-moss (*Sphagnum*) peat is light in colour, with the structure of the mosses perfectly preserved.

**peat-borer** Implement designed to extract *\*peat* cores with the minimum of disturbance. The most familiar is the Hiller peat-borer (or corer), which consists of a short screw *\*auger* head to ease penetration of the peat, backed by a chamber which can be opened and closed at the required sample depth,

the sharp cutting edge of the chamber assisting detachment of the sample in more consolidated peats. The principal alternatives are the piston sampler, which is particularly good for loose peats, and the Russian borer, which allows easier removal of the complete peat core than is possible with the Hiller borer, but which is more difficult to use in compacted peats since it has no screw auger head.

**peat podzol** \*Podzol \*soil profile distinguished by having a surface \*mor (peaty) \*humus up to a maximum thickness of 30 cm, and usually with an \*iron pan at the top of the B \*soil horizons. The term occurs in most of the classification systems derived originally from the work of V. V. \*Dokuchayev, published in 1886. It has been superseded in the \*USDA \*soil taxonomy and podzols now fall within the order \*spodosols, but the term is retained as a reference group in the \*World Reference Base for Soil Classification.

**peat soil** A major group in the soil classification developed by the Soil Survey for England and Wales, peat soils are rich in organic matter, which forms a layer at least 40 cm thick.

**pebble** See PARTICLE SIZE.

**pectolite** See WOLLASTONITE.

**pectoral girdle** In vertebrates (see CRANIATA), the skeletal structure that provides support for the fore limbs or fins.


**ped** Unit of soil structure (e.g. an \*aggregate, crumb, granule, or prism) that is formed naturally. Compare CLOD.

**pedestal rock (mushroom rock)** An unstable, mushroom-shaped landform found typically in arid and semi-arid regions. The undercut base was formerly attributed to wind \*abrasion, but is now believed to result from enhanced \*chemical weathering at a site where moisture would be retained longest. A famous example is Pedestal Rock, Utah, USA.

**pedicel** See ARACHNIDA.

**pedicle** A fleshy stalk which attaches most brachiopods (\*Brachiopoda) to the sea floor. It emerges either posteriorly from between the two valves of the shell or from a triangular notch (the \*delthyrium) in one of them. In

some species it issues from a circular **\*pedicle foramen** (in the ventral valve). In other species it is either not developed or atrophied.



### **Pedicle**

**pedicle foramen** A rounded, slit-like, or closed opening in the **\*pedicle** valve of a brachiopod (**\*Brachiopoda**) for the passage of the pedicle to the exterior.

**pediment (concave slope, waning slope)** Surface of low relief, partly covered by a skin of rock debris, that is concave-upward and slopes at a low angle (normally less than 5°) from the base of a mountain zone or **\*scarp**. Classically it is developed and has been investigated in the arid and semi-arid regions of the western USA.

**pedion** A **\*crystal** form consisting of only one face.

**pediplain** Extensive plain, best developed in arid and semi-arid regions, showing gently concave or straight-slope profiles and terminated abruptly by uplands. A result of scarp recession rather than of surface lowering, it consists of coalesced **\*pediments**.

**pedofacies** A **\*facies**, the composition of which can be used to relate the maturity of a **\*palaeosol** formed by **\*avulsion** to its distance from the active

river channel.

**pedogenesis** The natural process of soil formation, including a variety of subsidiary processes such as humification, *\*weathering*, *\*leaching*, and *\*calcification*.

**pedogeomorphology** See SOIL GEOMORPHOLOGY.

**pedology** The scientific discipline devoted to the study of the composition, distribution, and formation of soils, as they occur naturally.

**pedon** Three-dimensional sampling unit of soil, with depth to the *\*parent material* and lateral dimensions great enough to allow the study of all *\*soil-horizon* shapes and *\*intergrades* below the surface.

**pedoturbation** Any disturbance (perturbation) of soil caused by a process other than *\*illuviation*.

**peel technique** Originally a technique for palaeobotanical work, but then refined and now used extensively in carbonate sedimentology and palaeontological work. Calcareous material is etched in a weak solution of hydrochloric acid and differences in relief are produced. After washing, the surface is flooded with acetone, and polyvinylacetate (PVA) sheeting is rolled on to the surface. The acetone softens the sheeting and moulds it to the etched rock surface. After drying the sheeting is peeled from the surface, bringing a thin layer of the surface with it. This 'peel' can then be examined in transmitted light. A series of peels can be taken to reveal and reconstruct buried structures and staining with various chemicals may reveal additional details.

**pegmatite** Very coarse-grained, *\*igneous* rock, usually of granitic composition, in which the individual *\*crystals* are at least 2.5 cm long. The crystals are often more than 1 m in length, and may be far larger. Crystallization occurs at a late stage, when the *\*magma* is enriched in *\*volatiles* and *\*trace elements*. Pegmatites may concentrate some rare elements (lithium, boron, fluorine, tantalum, niobium, *\*rare-earth* elements, and uranium) to economic proportions.

**pegmatitic** Applied to an *\*igneous* rock with a texture resembling *\*pegmatite*. As pegmatitic rocks cooled from *\*magma*, some of their crystals grew very large, the largest being more than 12 m long. This

texture develops in magmas that contain much water, allowing the elements that make up the crystals to migrate readily to the crystallization sites.

**pel-** The prefix used in the *\*Folk* limestone classification to denote a *\*limestone* containing *\*pellets* as the major constituent of the particles present.

**pelagic** In marine ecology, applied to the organisms that inhabit open water, i.e. *\*plankton*, *\*nekton*, and neuston (although neuston are fairly unimportant in such environments).

**pelagic ooze** Deep-ocean *\*sediments* that accumulate by the settling out of materials from the overlying ocean waters. The dominant constituents are microscopic *\*pelagic* organisms, e.g. the calcareous *\*globigerina* and pteropods, and the siliceous *\*diatoms* and *\*radiolaria*. Minor amounts of fine volcanic, terrigenous, and extraterrestrial debris also contribute to pelagic ooze.

**pelagic sediment (pelagite)** A sediment formed in the open sea by the slow settling of calcareous and siliceous biogenic particles, biogenic material comprising more than 75% of the volume. *Compare* [HEMIPELAGIC SEDIMENT](#).

**pelagite** *See* [PELAGIC SEDIMENT](#).

**Peléean eruption** A very violent type of volcanic activity characterized by the release of *\*nuées ardentes* (glowing clouds of gas-fluidized material). Highly viscous *\*lavas* are typically involved and the *\*eruptions* may be preceded by the growth of a lava dome. Nuées ardentes may be formed by the explosive disruption of a growing dome, or by mechanical collapse of a dome growing on a steep slope. *See* [VOLCANO](#). *Compare* [HAWAIIAN ERUPTION](#); [PLINIAN ERUPTION](#); [STROMBOLIAN ERUPTION](#); [VESUVIAN ERUPTION](#); [VULCANIAN ERUPTION](#).

**Pelecypoda** *See* [BIVALVIA](#).

**Pelé's hair** Thin filaments of basaltic glass formed from chilled *\*lava* spray thrown out from a *\*volcano* during *\*Hawaiian \*eruptions*, and named after Pelé, the Hawaiian goddess of volcanoes. The lava is so fluid that on eruption it forms droplets, shaped primarily by surface-tension forces,

drawing behind them the long filaments which break to form the Pelé's hair. The filaments may be metres long, and drift downwind for many kilometres. The solidified droplets themselves are known as 'Pelé's tears'. Fragments formed from solidified lava spray are given the general name 'achneliths'.

**Pelé's tears** See PELÉ'S HAIR.

**pelite** (*adj.* pelitic) An aluminium-rich, \*metamorphic rock formed by the \*metamorphism of clay-rich \*sedimentary rocks, e.g. \*shales and \*mudstones. The type of aluminium-bearing \*silicate minerals seen in the rock depends on the pressure and temperature of the metamorphism but usually includes one of the \*mica group minerals. \*Quartz is ubiquitous in pelites.

**pelitic** See PELITE.

**pellet limestone** See LEIGHTON—PENDEXTER CLASSIFICATION.

**pellets** \*Grains of faecal origin, commonly found in \*limestones and \*phosphorites. Structureless grains resembling pellets which cannot be shown to be of faecal origin are termed 'peloids'. See FAECAL PELLETT.

**pellicular water (film water)** Thin films of water that cling to soil and rock particles above the \*water-table.

**pelmatozoan** Applied to echinoderms that are attached to the substratum (e.g. primitive \*Crinoidea). The term is used informally, and in some classifications Pelmatozoa is ranked as a subphylum of the \*Echinodermata. Compare ELEUTHEROZOAN.

**pelmicrite** See FOLK LIMESTONE CLASSIFICATION.

**peloid** See PELLETS.

**pelosols** A major group in the soil classification developed by the Soil Survey for England and Wales, pelosols are soils with a high concentration of \*clay that crack when they are dry.

**pelsparite** A \*limestone defined in the \*Folk limestone classification as comprising pellets together with a \*sparite \*cement.

**pelvic girdle** 1. In vertebrates (see CRANIATA), the skeletal structure that provides support for the hind limbs or fins. 2. The part of the abdomen that is surrounded by the bony pelvis.

**pelvis** In vertebrates, part of the appendicular skeleton that is fused to the sacral vertebrae and provides support for the hind limbs or fins.

**Pelycosauria** Order of \*synapsid reptiles dating from the Upper \*Carboniferous and Lower \*Permian, e.g. *Varanosaurus*, *Edaphosaurus*, and *Dimetrodon*. Several sported large, sail-like, dorsal fins, and both carnivorous and herbivorous types appeared. They gave rise to and were replaced by the mammal-like reptiles (\*Therapsida).

**Penck, Albrecht** (1858–1945) A German mineralogist from the University of Berlin, Penck's main interest was in \*Quaternary glacial land-forms, which he classified according to shape. Some of his work was done in cooperation with his son **Walther \*Penck**.

**Penck, Walther** (1888–1923) A German geologist, Penck assisted his father **Albrecht \*Penck** in his studies of land-forms. Independently, he worked on the structure of mountain regions, especially the Alps, modifying \*Suess's ideas on continental uplift.

**pendent** Applied to the attitude of the \*stipes that hang downwards from the \*sicula in graptolites (\*Graptolithina). This is the more \*primitive condition, from which other graptoloid (\*Graptoloidea) stocks arose.

**Pendleian** The first regional \*stage (326.4–326 Ma ago) of the \*Mississippian epoch, preceded by the \*Visean and overlain by the \*Arnsbergian.

**pene-** From the Latin *paene* meaning 'almost', a prefix meaning 'almost' or 'nearly' (e.g. a \*peneplain is almost a plain).

**peneplain (peneplane)** Literally, almost a plain: an extensive area of low relief, dominated by convex-up ('bulging') hillslopes mantled by a continuous \*regolith, and by wide, shallow river valleys. Locally, \*monadnocks may occur. A peneplain is the end-product of a cycle of erosion, produced by the action of down-wearing over a long period of time, and it is the end-product of the \*Davisian cycle.

**peneplane** See PENEPLAIN.

**penesaline** A level of *\*salinity* intermediate between normal marine and hypersaline, ranging from 72‰ to 352‰. These salinity levels are high enough to be toxic to normal marine organisms, and can be tolerated by only a restricted range of fauna and flora. The characteristic *\*sediments* of penesaline zones are evaporitic *\*carbonates* (see EVAPORITE) interbedded with *\*anhydrite* or *\*gypsum*. Penesaline environments are often encountered in the back-*\*barrier* and *\*backreef* zones.

**penetration test** A standard penetration test of *\*sand* or *\*silt* determines the number of blows of a standard weight from a standard height that are needed to produce a standard penetration. A dynamic penetration test determines the *\*density* of layered *\*sediments* by recording the penetration per blow or series of blows. See also CONE PENETROMETER.

**penetration twin** See INTERPENETRANT TWIN.

**penitentes** (*sing. penitent*) Thin blades of snow or ice that stand upright, up to 2 m tall, surrounding a bowl-shaped depression several tens of centimetres in diameter, found at high altitudes on tropical mountains. First described by Charles *\*Darwin* in 1839 in Chile, they are named for the pointed caps worn by Roman Catholics doing penance.

**Pennales** See BACILLARIOPHYCEAE.

**pennate diatoms** See BACILLARIOPHYCEAE.

**Pennatulacea** See OCTOCORALLIA.

**penninite** See CHLORITE.

**Pennsylvanian** 1. The Late *\*Carboniferous* *\*epoch*, preceded by the *\*Mississippian*, followed by the Early *\*Permian*, and comprising the Early, Middle, and Late Pennsylvanian sub-epochs, dated at 318.1–299 Ma ago (Int. Commission on Stratigraphy, 2004). 2. The name of the corresponding N. American sub-system, comprising the *\*Morrowan*, *\*Atokan* (Derryan), *\*Desmoinesian*, *\*Missourian*, and *\*Virgilian* *\*series*, and roughly contemporaneous with most of the *\*Silesian* sub-system (i.e. above *\*Namurian* A).


**pentadactyl** Applied to a limb possessing five digits, or to one modified evolutionarily from an ancestral form which possessed five digits;

characteristic of all **\*tetrapods**.

**pentagonal dodecahedron** See PYRITOHEDRON.

**pentamerall symmetry** Five-sided symmetry, found in most members of the phylum **\*Echinodermata**. In some groups this symmetry has a **\*bilateral symmetry** superimposed on it.

**Pentamerida (pentamerids)** (class **\*Articulata**) Extinct order of brachiopods (**\*Brachiopoda**), with thick, strongly biconvex shells. A **\*spondylium** developed in the **\*pedicle valve** and brachial processes and supporting plates in the brachial (**\*dorsal**) valve, forming an internal calcareous box containing the adductor muscles (which close the valves) and diductor muscles (which open the valves). The shell is **\*impunctate**. Pentamerida first appeared in the Middle **\*Cambrian** and became extinct in the Upper **\*Devonian**.



**Pentamerida**

**pentamerids** See PENTAMERIDA.

**pentectic point** See BOWEN'S REACTION PRINCIPLE.

**Pentevrian** A **\*stage** of the Lower and Middle **\*Proterozoic** of Brittany, from about 2600 to 1000 Ma ago, overlain by the **\*Brioverian**.

**pentlandite** An iron and nickel sulphide, with the formula  $(\text{FeNi})_9\text{S}_8$ , and the main **\*ore mineral** for nickel; sp. gr. 4.5–5.0; **\*hardness** 3–4; **\*cubic**;

light bronze-yellow; greenish-black *\*streak*; *\*metallic \*lustre*; very rarely found in well-formed crystals, usually occurring as irregular grains and inclusions; no *\*cleavage*, parting {111}; occurs with the other *\*sulphide* ores genetically associated with *\*basic* and *\*ultrabasic* rocks, together with *\*pyrrhotite* and *\*chalcopyrite*.

**Penutian** A *\*stage* (53–51 Ma ago) in the Early *\*Eocene* of the west coast of N. America, underlain by the *\*Bulitian*, overlain by the *\*Ulatisian*, and roughly contemporaneous with the lower *\*Ypresian* stage.

**peperite** A mixture of solidified *\*magma* and sediment found along the margin of an *\*intrusion* or at the base of a larval flow.

**peralkaline** A chemical classification applied to *\*felsic \*igneous* rocks in which there are more molecules of (Na<sub>2</sub>O + K<sub>2</sub>O) than of Al<sub>2</sub>O<sub>3</sub>. Rocks with such a chemistry crystallize alkali-rich *\*ferromagnesian* minerals such as *\*aegirine*, aegirine-*\*augite*, *\*barkevikite*, and *\*arfvedsonite*. *See also METALUMINOUS*.

**peraluminous** *See METALUMINOUS*.

**peramorphosis** Evolutionary change that results in the descendant incorporating all the ontogenetic stages of its ancestor, including the adult stage, in its *\*ontogeny*, so that the adult descendant ‘goes beyond’ its ancestor. It may occur by *\*acceleration*, *\*hypermorphosis*, or *\*predisplacement*.

**perched aquifer** *See AQUIFER*.

**perched block** Glacially transported boulder resting on bedrock where it was deposited by melting ice. Some examples are spectacular and give rise to legends locally.

**percolation** In *\*pedology*, the downward movement of water through soil, especially through soil that is saturated or close to saturation.

**percussion boring** Drilling with a chisel-shaped *\*bit* which is repeatedly struck against rock to form a hole. It produces very fine debris, which can be washed away by drilling fluid. If only the drill bit can oscillate up and down to produce the percussive effect, the drilling equipment is called a ‘down-the-hole hammer’. If the whole series of drilling rods and bit are

oscillated from outside the drill hole, the equipment is called a 'drifter'. See [DOWN-HOLE HAMMER DRILLING](#).

**Perdita (Uranus XXV)** A lesser satellite of [\\*Uranus](#) with a radius of 10 km.

**perennial stream** A stream that would normally be expected to flow throughout the year, albeit with only small dry-weather flows in some cases.

**pergelic** The lowest of the soil-temperature classes for family groupings of soils in the soil taxonomy system, applied to soils in temperate regions. The assessment of soil temperature is based on the mean annual temperature, and on the difference between mean summer and mean winter temperature, measured at a depth of 50 cm or at the surface of the underlying rock, whichever is shallower. In order of ascending temperature, the higher-temperature classes in temperate-region soils are called cryic, frigid, mesic, thermic, and hyperthermic, and in tropical regions the scale from cold to hot is isofrigid, isomesic, isothermic, and isohyperthermic.

**pergelisol** See [PERMAFROST](#).

**peri-** From the Greek *peri* meaning 'around', a prefix meaning 'around' or 'enveloping'.

**periclase** Magnesium oxide (MgO), a member of the [\\*oxide](#) group of minerals; sp. gr. 3.5; [\\*hardness](#) 5.5; white, yellow, or green; crystalline or granular; occurs in contact-metamorphosed (see [THERMAL METAMORPHISM](#)) [\\*limestones](#) and may be altered to [\\*brucite](#) (Mg(OH)<sub>2</sub>) by hydration.

**periclinal fold** See [PERICLINE](#).

**pericline (periclinal fold)** A [\\*fold](#) shown on a map as concentrically arranged contour patterns whose amplitude decreases to zero in a regular fashion in two directions. An [\\*anticlinal](#) pericline is called a 'dome', a [\\*synclinal](#) pericline a 'basin'.

**pericline twinning** One of the many types of twins (see [CRYSTAL TWINNING](#)) shown by the [\\*plagioclase feldspars](#). The twin plane is along the *b* (or *y*) axis and produces polysynthetic [\\*lamellar](#) twinning which results in a series of fine striations on the [\\*cleavage](#).

**pericontinental sea (marginal sea)** A sea that surrounds a continent.

**periderm** The thin, sheet-like material that makes up the skeleton in graptolites (*\*Graptolithina*). Long thought to be *\*chitin*, it is now considered perhaps to be *\*collagen* (a scleroprotein). It is bi-layered, with an inner fusellar layer and outer cortical layer which have distinct structural differences.

**Peridiniales** See DINOPHYCEAE.

**peridotite** A coarse-grained, *\*ultrabasic*, *\*igneous* rock consisting of *\*essential* magnesium-rich *\*olivine* accompanied by lesser amounts of other *\*ferromagnesian minerals*, e.g. *\*orthopyroxene* (*\*enstatite*-*\*bronzite*), *\*clinopyroxene* (chromium-*\*diopside*), and *\*chromite*, with or without *\*pyrope* *\*garnet*. Peridotites are found in large, layered, *\*ultrabasic*, *\*intrusive* bodies, in *\*ophiolite* complexes, and as *\*xenoliths* brought up in *\*alkali* *\*basalts* and *\*kimberlites*. As well as forming by *\*crystal* fractionation processes in igneous bodies, their mineral chemistry and presence as xenoliths in kimberlites suggest that much of the Earth's *\*mantle*, and therefore of the mass of the Earth, is peridotite. *\*Meteorites* are also composed largely of peridotite, suggesting that peridotites are probably the commonest rock in the *\*solar system*.

**peridotite model** A model for the composition of the Earth's upper *\*mantle* based on compressional and *\*S-wave* velocities and gravity data. The probable composition is *\*peridotite*, *\*eclogite*, or an intermediate between the two. *\*Xenoliths* in *\*kimberlites*, and *\*mafic* volcanic rocks found at the surface, also suggest this composition.

**perigee** The point nearest the planet in the orbit of a *\*satellite*, originally defined for the lunar orbit. *Compare* APOGEE.

**periglacial** Applied strictly to an area adjacent to a contemporary or *\*Pleistocene* *\*glacier* or *\*ice sheet*, but more generally to any environment where the action of freezing and thawing is currently, or was during the Pleistocene, the dominant surface process.

**periglacial morphoclimatic zone** A *\*morphoclimatic* region in which the rate of *\*mechanical* weathering is very high, especially that due to *\*frost* action, and the rate of *\*chemical weathering* is low to moderate. *\*Fluvial*

processes are active seasonally, wind action is high locally, and there is much *\*mass-wasting*.

**perignathic girdle** Continuous or discontinuous ring of internal processes around the *\*peristome* of echinoids (*\*Echinoidea*), to which the muscles which support and control the lantern (the chewing mechanism, *see ARISTOTLE'S LANTERN*) are attached.

**perihelion** The point in the Earth's elliptical orbit at which the planet is closest to the Sun. During the present epoch this occurs about the beginning of January, when the average amount of incident solar radiation is at a maximum. *See also APHELION*.

**period 1.** Second-order geologic time unit which is the equivalent of the chronostratigraphic unit '*\*system*'. Periods are subdivided into *\*epochs*; together, several periods constitute an *\*era*. When used formally the initial letter of the term is often capitalized, e.g. the *\*Devonian* Period. **2. (T)** The time that elapses between repetitions of the same phase of a wave-form, i.e. the time taken to complete one cycle. For a simple harmonic function,  $T = 2\pi/\omega$  where  $\omega$  is the angular velocity; for a wavetrain of single *\*frequency*  $f$ ,  $T = 1/f = \lambda/V$  where  $\lambda$  is the *\*wavelength* and  $V$  the *\*phase velocity*.

**periodic bedrock ridges (PBRs)** A landform on Mars that consists of parallel ridges and troughs with an average wavelength of 18–51 m. They resemble sand dunes, but have formed by wind erosion of softer bedrock from a more coherent rock. PBRs are aligned transverse to the prevailing wind and the scientists who identified them believe they were formed by high winds that eroded the windward sides of the ridges and were deflected into the air, causing erosion on the lee side as the wind sank.

**periostracum** *See SHELL STRUCTURE; SKELETAL MATERIAL.*

**peripatric speciation** *See FOUNDER EFFECT.*

**peripheral foreland basin (pro-foreland basin)** A *\*foreland basin* that forms on the outer arc of an *\*orogen* on the *\*plate* that is subducted or underthrust during a plate collision.

**periproct** In *\*Echinoidea*, the area surrounding the anus and covered in leathery skin in which small, calcitic plates are loosely embedded.

**Perissodactyla** (cohort **\*Ferungulata**) Odd-toed ungulates. Order of mammals that comprises those ungulates in which the number of functional toes is reduced to three or one, and a fourth, if present, is reduced, the weight of the animal being borne by the central digit. The order includes the three suborders Ceratomorpha (tapirs, rhinoceroses, and their extinct relatives); Ancylopoda (or Chalicotheres), which are extinct forms with claw-like extremities in place of hoofs, e.g. *Moropus*; and Hippomorpha (horse-like forms). They appeared in the **\*Eocene**, derived from the **\*condylarths**, and reached their zenith in the mid-**\*Tertiary**, when they were the most numerous of ungulates. So many **\*fossil** remains of them have been found that their evolutionary history is known in greater detail than that of any other mammalian group. Since then the order has declined dramatically, having been displaced by the **\*artiodactyls**, and the group as a whole is moving towards extinction.

**peristome** (*adj.* peristomal) In **\*Echinoidea**, the area around the mouth, covered in a leathery skin studded with small plates.

**peritectic point** See **BOWEN'S REACTION PRINCIPLE**.

**peritidal** Applied to the zone extending from above the level of the highest **\*tide** to below that exposed at the lowest tide, and thus somewhat wider than the **\*intertidal zone**.

**perlite** A **\*glass** formed mainly by the **\*hydration** of **\*obsidian** that has the property of expanding when heated. It softens at 850–900 °C, and water held in its structure vaporizes causing the perlite to expand to seven to sixteen times its original volume. Expansion reduces its **\*bulk density** from about 1100 kg/m<sup>3</sup> to 30–150 kg/m<sup>3</sup>. Expanded perlite has many uses in construction and horticulture.

**perlitic** Displaying curved or sub-spherical cracks. The texture is found in glassy or devitrified **\*igneous** rocks and forms by contraction during rapid cooling of the **\*magma**.

**perlucidus** From the Latin *perlucidus* meaning 'allowing the passage of light', a variety of cloud comprising extensive layers or sheets with holes which allow a view beyond the cloud. The term is applied to **\*stratocumulus** and **\*altocumulus**. See also **CLOUD CLASSIFICATION**.

**permafrost (pergelisol)** Permanently frozen ground which occupies some 26% of the Earth's land surface under thermal conditions where temperatures below 0 °C have persisted for at least two consecutive winters and the intervening summer. Considerable thicknesses may develop, e.g. 600 m on the North Slope of Alaska, and 1400 m in Siberia, but these are partly relict from the last glaciation. Permafrost may contain an unfrozen unit, called 'talik', and may be overlain by an *\*active layer*.

**permafrost table** The upper limit of *\*permafrost*. Compare TJAELE.

**permanent strain (non-recoverable strain)** The deformation that is retained in a rock after a distortional strain is removed, forming a *\*shear zone*.



<http://www.see.leeds.ac.uk/structure/rheology/index.htm>

- Describes the concept of rheology.

**permanent wilting percentage (permanent wilting point, wilting coefficient, wilting point)** The percentage of water remaining in the soil after a specified test plant has wilted under defined conditions, so that it will not recover unless it is given water.

**permanent wilting point** See PERMANENT WILTING PERCENTAGE.

**permeability 1. (hydraulic conductivity)** In general, the ability of a rock, *\*sediment*, or soil to permit fluids to flow through it. More precisely, the hydraulic conductivity is the volume flow rate of water through a unit cross-sectional area of a porous medium under the influence of a *\*hydraulic gradient* of unity, at a specified temperature. It is measured in units of m/s or m/day and varies with temperature. Typical values range from  $10^{-6}$  m/day for *\*clay* to  $10^3$  m/day for coarse gravel. The magnitude of hydraulic conductivity depends on the properties of both the fluid and the medium. An alternative measure, used in the oil industry, is intrinsic permeability, measured in  $m^2$  (or in industrial units called *\*darcies*), which depends on the properties of the rock alone. **2.** Property of a membrane or other barrier, being the ease with which a substance will diffuse or pass across it. **3.** Capacity of a material to transmit fluids, expressed as hydraulic conductivity. In soils, the ease with which gases, liquids, or plant roots

penetrate into or pass through a layer of soil. **4. (magnetic permeability)** The ratio of the **\*magnetic flux** density in a medium to the magnetizing force. In free space (i.e. air) this is a constant,  $\mu_0$  equal to  $4\pi \times 10^{-7}$ .

**permeameter** A laboratory device for measuring the **\*hydraulic conductivity** of rock and soil samples or the coefficient of **\*permeability** of soil. Two main types are commonly used: those that require movement of water and those that do not, known respectively as falling head and constant head permeameters.

**Permian** Final **\*period** (299–251 Ma ago) of the **\*Palaeozoic \*era**, which is named after the central Russian province of Perm. The period is often noted for the widespread continental conditions that prevailed in the northern hemisphere and for the extensive nature of the southern hemisphere glaciation. Many groups of animals and plants, including the rugose corals (**\*Rugosa**), trilobites (**\*Trilobita**), and blastoid echinoderms (**\*Blastozoa**), vanished at the end of the Permian, in a mass **\*extinction** that was one of the great crises in the history of life. The period is divided into nine **\*ages**: the **\*Asselian**, **\*Sakmarian**, **\*Artinskian**, and **\*Kungurian** ages comprise the Cisuralian epoch; the Roadian, Wordian, and Capitanian ages comprise the Guadalupian epoch, and the Wuchiapingian and Changhsingian ages comprise the Lopingian epoch.



<https://ucmp.berkeley.edu/permian/permian.php>

- The Permian Period.

**permineralization** See FOSSILIZATION.

**permitted intrusion** The 'passive' intrusion of **\*magma** which ideally produces no tensional forces and involves no forceful emplacement. The intrusion occurs by **\*stopping**, with the melting and assimilation of wall rock.

**permittivity** See DIELECTRIC CONSTANT; DIELECTRIC PERMITTIVITY.

**perovskite** A **\*mineral \*oxide**  $\text{CaTiO}_3$ ; sp. gr. 3.98–4.26; **\*hardness** 5.5; **\*monoclinic** or **\*orthorhombic**; tiny, **\*cubic \*crystals**; yellow-brown to black; common **\*accessory mineral** in undersaturated (see SILICA SATURATION) **\*alkaline \*igneous** rocks found with **\*nepheline** or **\*melilite**.

Occasionally occurs in some ultramafic (see [ULTRABASIC](#)), [\\*plutonic](#), igneous rocks.

**perovskite model** A theoretical model of the Earth's composition suggesting that the mineral perovskite ( $\text{CaTiO}_3$ ) makes up most of the lower part of the [\\*mantle](#) between 600 and 2800 km depth, at 2000–2500 °C. Perovskite has been made artificially by compressing [\\*olivine](#), [\\*pyroxene](#), and [\\*garnet](#), and has been shown to withstand temperatures of 2000 °C in the laboratory. The properties of [\\*mantle](#) perovskites strongly influence the transport properties of rocks in the lower mantle. Perovskites are expected to occur throughout the universe as an important constituent of rocky planets larger than [\\*Mars](#).

**persistence** Continuation of an [\\*anomaly](#) beyond its expected limits in time or space.

**perthite** A series of layers that occurs as intergrowths in [\\*alkali feldspars](#) which form at high temperatures and cool slowly, unmixing as they cool. Most alkali feldspars in such slowly cooled rocks (e.g. [\\*plutonic](#) and [\\*igneous](#)) consist of an alkali feldspar host with an exsolved, sodium-rich, [\\*plagioclase feldspar](#) phase that segregated from the alkali feldspar during cooling. Perthitic intergrowths have textures ranging from macroperthite (visible to the naked eye) through microperthite (visible under a microscope) to cryptoperthite (which can be detected only by [\\*X-ray](#) diffraction and other techniques). The nature of the perthite depends upon the ordering of the silicon and aluminium atoms achieved by the mineral lattice as it cools.

**Peru–Chile Trench** The oceanic [\\*trench](#) which marks the boundary between the subducting [\\*Nazca Plate](#) and the [\\*South American Plate](#).

**Peru current (Humboldt current)** Oceanic water flowing northwards along the west coast of Chile and Peru, driven by the westward flow of the S. [\\*Equatorial](#) current (itself driven by south-east trade winds). It is essentially a 'continuity current': water flows into the low-sea-level region left by the S. Equatorial current. This eastern [\\*boundary](#) current is slow-moving, broad, and shallow, and is noted for a prominent area of [\\*upwelling](#) bringing cold bottom waters to the surface. As the near-surface concentrations of nutrient elements are high there is an abundance of marine life associated with this current.

**PeruSat-1** A Peruvian Earth-observation *\*minisatellite* mission that provides high-resolution imagery. It was launched on 16 September 2016, from Kourou, French Guiana, into a Sun-synchronous orbit at an altitude of 695 km.

**petals** See IRREGULAR ECHINOIDS.

**petit-spot volcano** A *\*volcano* that forms where oceanic *\*lithosphere* flexes as an ocean plate approaches a *\*subduction zone*. Flexure produces fractures, which cause melting in the *\*mantle*, allowing *\*magma* to rise to produce a line of volcanoes parallel to and some distance from the subduction zone. Petit-spot volcanism was first discovered offshore from Japan but has since been identified in association with subduction zones in other parts of the world.

**petrification** See FOSSILIZATION.

**petrocalcic horizon** *\*Indurated \*calcic \*soil horizon* that is *\*cemented* by a high concentration of calcium carbonate, often comprising 40% by weight of the mineral material, and that is impenetrable to plant roots or to spade digging.

**petrofabric** See FABRIC.

**petrogenesis** The origin and evolutionary history of rocks, especially *\*igneous* rocks. Compare PETROGRAPHY. See PETROLOGY.

**petrogenetic grid** A diagram that links the stability ranges of metamorphic minerals or mineral assemblages to the conditions of *\*metamorphism*. Experimentally determined mineral or mineral-assemblage stability ranges are commonly plotted as reaction boundaries in pressure–temperature space to produce a petrogenetic grid for a particular range of *\*rock* compositions. The region of overlap of the stability fields of those minerals forming the equilibrium mineral assemblage define the pressure–temperature conditions of the equilibrium. This corresponds to the pressure and temperature of the metamorphism.

**petrographic microscope** See POLARIZING MICROSCOPE.

**petrography** Systematic description and interpretation of rock *\*textures* and *\*mineralogy* in *\*thin* section and as hand specimens. Compare

PETROGENESIS. See PETROLOGY.

**petrogypsic horizon** Surface or subsurface **\*soil horizon** cemented by **\*gypsum** so strongly that dry fragments will not slake in water. The **\*cementation** restricts penetration by plant roots. This is a **\*diagnostic horizon**.

**petroleum (crude oil)** In geology, a term used generally to describe naturally occurring liquid **\*hydrocarbons** formed by the anaerobic decay of organic matter. Oil is rarely found at its original site of formation but migrates to a suitable **\*structural** or **\*lithological trap**. Petroleum is frequently associated with salt water and with gaseous hydrocarbons. See also NATURAL GAS; OIL SHALE.

**petroleum geology** Study of the mode of origin and conditions of accumulation of **\*hydrocarbons** in the Earth's crust; their exploration, involving the application of the techniques of **\*geophysics**, **\*geochemistry**, **\*palaeontology**, **\*stratigraphy**, and **\*tectonics**; and their evaluation.

**petrological microscope** See POLARIZING MICROSCOPE.

**petrology** The study of rocks in general, including their occurrence, field relations, structure, origins and history (**\*petrogenesis**), and their **\*mineralogy** and **\*textures** (**\*petrography**). It may usefully be qualified as 'igneous petrology', or 'sedimentary petrology'.

**petrophysics** The study of the character of rocks using data from down-hole geophysical **\*well** logs. Petrophysical analysis allows an estimate of **\*porosity**, shaliness, rock density, gas content, water saturation, and **\*lithology**, using the suites of **\*logs** normally run in a well.

**p-form** A smooth, plastically sculptured feature (e.g. a groove, crescentic depression, or pothole), generally less than 20 m in size, cut into an exposed rock surface by the action of a **\*glacier**. It may be attributed to abrasion when it is linear and shows striations, but the origin of other types is uncertain.

**PGF** See PRESSURE-GRADIENT FORCE.

**pH** The negative logarithm of the hydrogen **\*ion** concentration in a solution,  $\text{pH} = \log_{10} 1/\text{H}^+$ . If the hydrogen ion concentration of a solution

increases, as happens with increasing acidity, the pH will decrease, and vice versa. The pH is measured on a scale of 0–14; a neutral medium (such as pure water) has a pH of 7.0, numbers above 7.0 indicate relative alkalinity, numbers below 7.0 indicate relative acidity. Most pH values in natural systems lie in the range 4.0–9.0. Human blood has a pH of 7.4, ocean water 8.1–8.3, water in saline environments may have a pH around 9.0 or higher, while for water in acidic soils it may be as low as 4.0 or less.

**phaceloid** See COMPOUND CORALS.

**phacoidal** Lens-shaped.

**phacolith** A curved, lensoidal *\*intrusion* which is *\*concordant* and follows the arches and troughs of folded *\*strata*.

**Phacopida** An order of *\*Trilobita* that lived from the Lower *\*Ordovician* to Upper *\*Devonian*. In most, the *\*cephalic suture* is *\*proparian*. There are three suborders.

**Phaeophyceae (brown algae)** Class of *\*algae* which includes no single-celled species; almost all are marine, growing mostly in the intertidal regions. They are the dominant seaweeds in the colder waters of the northern hemisphere. They are typically olive brown or greenish in colour (at least when wet) due to the presence of the pigment fucoxanthin in the chloroplasts. Some fossil plants resembling Fucales (an order of brown seaweeds) have been reported from the *\*Palaeozoic*, but their fossil record is equivocal owing to their lack of calcification.

**phaeozems** A reference soil group in the *\*World Reference Base for Soil Resources*. Phaeozems comprise all soils with a *\*mollic* A horizon, except for *\*chernozems* and *\*kastanozems*.

**phaneritic** Applied to an *\*igneous* rock in which the mineral grains are large enough to be visible to the naked eye.

**phanerocrystalline** Having a crystal structure in which the crystals are visible to the naked eye.

**Phanerozoic** Period of geologic time comprising the *\*Palaeozoic*, *\*Mesozoic*, and *\*Cenozoic* *\*eras*, it began approximately 542 Ma ago at the start of the *\*Cambrian* *\*period* and continues to the present day. It is marked by the accumulation of *\*sediments* containing the remains of

animals with mineralized skeletons. Although the name is derived from the Greek *phaneros* meaning ‘visible’ and *zoion* meaning ‘animal’, the term is no longer used in the sense of ‘visible life’, but its commencement merely defines the base of the Cambrian.



<http://www.geologypage.com/2013/12/phanerozoic.html>

- Phanerozoic.

**phase 1.** An individually distinct and homogeneous part of a system. For example, liquid water and water vapour are each single phases; a mixture of the two constitutes a two-phase system. Similarly *\*minerals* crystallizing from a *\*melt* form separate phases within it. A ‘phase boundary’ is the line marking the contact between two constituent or liquid phases. **2.** A short unit of time, or an episode of development or change, usually within the context of a longer period. The term has been used informally (see *INFORMAL*) in this sense in many branches of the Earth sciences, e.g. ‘a phase of *\*igneous* activity’, or ‘a cold phase’ during a generally warmer period.

**phase angle** The angle, usually termed  $\alpha$ , formed by lines joining the centres of the *\*Earth*, the *\*Sun*, and another *\*solar system* object. Aristarchus of Samos (3rd century BC) measured the Earth–Sun–Moon angle for the first lunar quarter and thus established that the Sun was much more distant than the Moon.

**phase boundary** The boundary between two *\*phases* of the same chemical substance (e.g. between liquid and solid).

**phase diagram** Graph to show fields of stability for *\*phases* in a heterogeneous system; variables such as temperature, pressure, and concentration are plotted against each other.

**phase layering** See *MINERAL LAYERING*.

**phase rule** Rule stating that for any system in equilibrium,  $F = (C - P) + 2$ , where  $F$  is the number of *\*degrees* of freedom,  $C$  the number of components, and  $P$  is the number of phases.

**phase transitions in the mantle** Measurement of the velocity of *\*seismic waves* and determination of the Earth's *\*moment* of inertia indicate that at depths of 80–100 km there appears to be a *\*low-velocity* zone and from 400–900 km velocities increase more rapidly than had been expected. These variations in the *\*mantle* may be due to changes in composition or to *\*phase* change, e.g. a transition to a *\*cubic* form of *\*olivine*.

**phase velocity (V)** The speed with which a particular *\*phase* (e.g. peak or trough) of a wave travels. *Compare* GROUP VELOCITY.

**phena** See PHENON.

**phenoclast** A large and conspicuous *\*clast* set in a generally finer-grained *\*sediment*, e.g. a large *\*boulder* in a fine-grained *\*conglomerate*, or a *\*granule* in *\*siltstone*.

**phenocryst** Large and often well-formed *\*crystals* set in a finer *\*groundmass* or *\*matrix*. Rocks containing phenocrysts are said to be porphyritic. *Compare* PORPHYROBLAST.

**phenogram** A tree-like diagram used in analysis to show similarity or dissimilarity among specimens or groups of specimens.

**phenon** (*pl.* **phena** or **phenons**) A group of organisms that are similar in appearance. They may or may not belong to the same taxon.

**phenotype** Those observable properties of an organism that are produced by the *\*genotype* in conjunction with the environment. Organisms with the same overall genotype may have different phenotypes because of the effects of the environment and of *\*gene* interaction. Conversely, organisms may have the same phenotype but different genotypes.

**Philippine Plate** One of the present-day minor lithospheric *\*plates*, the Philippine Plate is surrounded by *\*subduction zones* (the Ryukyu, Philippine, Marianas, and Izu-Bonin), and is splitting along the Marianas Islands, in the *\*back-arc* of the *\*Marianas Trench-arc* system, at about 60 mm/year.

**Philippine Scientific Earth Observation Microsatellite-1** See DIWATA-1.

**phi scale** The expression of *\*grain* sizes on a logarithmic scale. The phi value ( $\phi$ ) is related to grain diameter ( $d$ ) by the expression  $\phi = -\log_2 d$ .

Increasing positive phi values are for grain diameters progressively finer than 1000 μm, and increasing negative values for grain diameters progressively coarser than 1000 μm (e.g.  $2\phi = 250\ \mu\text{m}$ ;  $1\phi = 500\ \mu\text{m}$ ;  $0\phi = 1000\ \mu\text{m}$ ;  $-1\phi = 2\ \text{mm}$ ;  $-2\phi = 4\ \text{mm}$ ).

**PHL-Microsat-1** See [DIWATA-1](#).

**phlogopite** A member of the *\*mica* group of *\*minerals*, phlogopite is the Mg-rich variety of *\*biotite*  $\text{K}_2(\text{Mg,Fe}^{2+})_6[\text{Si}_3\text{AlO}_{10}]_2(\text{OH,F})_4$  with Mg/Fe more than 2/1; sp. gr. 2.76–2.90; *\*hardness* 2.0–2.5; subtransparent; pale brown; small, tabular *\*crystals*; occurs in metamorphosed, impure, magnesian *\*limestones* by reaction between *\*dolomite* and K-*\*feldspar* or *\*muscovite*, in *\*kimberlites*, and in some *\*leucite-bearing* rocks.

**Phobos** 1. Two Soviet missions to *\*Mars*, launched in 1988; both were lost.  
2. One of the two satellites of Mars.



<http://nssdc.gsfc.nasa.gov/planetary/phobos.html>

- An RSA mission to study the Sun, Mars, and Phobos, a moon of Mars.

**Phocidae** See [CARNIVORA](#).

**Phoebe (Saturn IX)** One of the lesser satellites of *\*Saturn*, with a radius measuring  $115 \times 110 \times 105\ \text{km}$ ; visual albedo 0.06. Its orbit is *\*retrograde*. It was discovered in 1898 by W. Pickering.

**Phoenix Plate** See [NAZCA PLATE](#).

**Pholas** See [BORING](#); [DOMICHNIA](#).

**phonolite** A fine-grained, *\*porphyritic*, *\*extrusive*, *\*igneous* rock consisting of *\*essential \*alkali feldspar* (*\*sanidine* or *\*anorthoclase*), *\*nepheline*, sodic *\*pyroxene*, and sodic *\*amphibole* (with or without iron-rich *\*olivine*), with *\*accessory \*sphene*, *\*apatite*, and *\*zircon*. Where present, *\*phenocrysts* can consist of alkali feldspar, sodic pyroxene, or sodic amphibole. Phonolites are the extrusive equivalent of *\*nepheline-syenites* and are found on off-axis ocean islands and in continental regions subjected to anorogenic upwarping and rifting. The name, from the Greek

*phone* meaning 'sound', refers to the fact that the rock rings like a bell when struck with a hammer.

**phosphates** Rock or deposit made up largely of inorganic phosphate, commonly calcium phosphate (e.g. the *\*minerals* *\*apatite*, *\*autunite*, *\*monazite*, *\*pyromorphite*, *\*torbernite*, *\*turquoise*, *\*vivianite*, and *\*wavellite*).

**phosphorescence** The property of some *\*minerals* of emitting light during exposure to X-rays, ultraviolet light, or cathode radiation and continuing to do so after the exposure has ceased (if light emission ends when the radiation source is switched off, the property is called 'fluorescence'). The colour of the emitted light varies with the wavelength of the radiation to which the mineral is exposed, and is thought to be due to the presence of traces of organic material or *\*cations* within the atomic structure of the mineral.

**phosphorite** A *\*sedimentary rock* rich in phosphate, usually in the form of carbonate hydroxyl fluorapatite ( $\text{Ca}_{10}(\text{PO}_4\text{CO}_3)_6\text{F}_{2-3}$ ). Phosphorites occur as *\*nodules* and crusts formed in oceanic areas where sedimentation rates are very low, by the early, near-surface diagenetic alteration (see *DIAGENESIS*) of *\*ooids*, *\*pellets*, and *\*bioclasts*, or as accumulations of *\*bones* and fish scales in *\*fluvial* to shallow marine environments. See also *GUANO*.

**photo-** From the Greek *phos photos* meaning 'light', a prefix meaning 'light'.

**photochemical smog** Hazy condition of the atmosphere due to the reaction of hydrocarbons with molecules of nitrogen oxide in sunlight which produces complex organic molecules of peroxyacetyl nitrates (PAN). In humid conditions these molecules produce *\*smog*. Such phenomena are common in large urban areas (e.g. the Los Angeles Basin and Athens) where there are stable atmospheric conditions and a high level of hydrocarbon input from incomplete combustion in car engines. Natural photochemical reactions occur in the high atmosphere with the absorption of radiation by oxygen to produce ozone. See also *OZONE LAYER*.

**photodisintegration** Decomposition of a compound in the presence of light, particularly sunlight.

**photodissociation** The splitting of a molecule into atoms or other molecules as a result of its absorption of radiation.

**photogeology** The determination of the overall geology of an area through the interpretation of photographic data (see [AERIAL PHOTOGRAPHY](#)) by noting variations in colour, tone, geometry, relative relief, and surface texture of land-forms, rock [\\*outcrop](#) boundaries, vegetation patterns, etc.

**photogrammetry** In [\\*remote sensing](#), the use of aerial photographs or satellite images to measure distances between ground objects accurately.

**photographic infrared** See [NEAR-INFRARED](#).

**photohydrometer** An instrument that calculates the size of sediment grains by measuring changes in the intensity of a light beam passed through a column of settling particles.

**photometer** In a [\\*reflected-light](#) microscope, an attachment which is used to measure the [\\*reflectance](#) of [\\*ore minerals](#). It consists of a photomultiplier tube that has high sensitivity throughout the visible spectrum. It is used in conjunction with a stabilized light source and high quality [\\*monochromators](#); its readings conform to [\\*Commission on Ore Mineralogy](#) (COM) standards.

**photometry** A general term for the physical measurement of the ultraviolet, visible, and infrared portions of the electromagnetic spectrum.


**photon log** The record of a scintillometer that has passed through a [\\*borehole](#). It is identical to a [\\*gamma-ray log](#), except that the sonde is central within the hole and therefore sensitive to the size of the borehole. (The gamma-ray sonde is held against the borehole wall and is insensitive to the size of the borehole.)

**photopolarimeter-radiometer (PPR)** A [\\*remote-sensing](#) instrument that measures the intensity and polarization of sunlight in the visible part of the spectrum, from which the temperature and cloud formation in planetary and satellite atmospheres can be determined, as well as some surface details.

**photosphere** The thin shell of light around the [\\*Sun](#) from which light escapes and within which the solar [\\*spectra](#) originate.

**photosymbiosis** A symbiotic (see SYMBIOSIS) relationship between two organisms (e.g. \*Foraminiferida and algae) one of which (in this example the \*alga) is capable of \*photosynthesis.

**photosynthesis** Term given to the series of metabolic reactions that occur in certain autotrophic organisms, whereby organic compounds are synthesized by the reduction of carbon dioxide using energy absorbed by chlorophyll from sunlight. In green plants, where water acts both as a hydrogen donor and as a source of released oxygen, photosynthesis may be summarized by the empirical equation:



(oxygen being released as a gas). Photosynthetic bacteria are unable to utilize water and therefore do not produce oxygen. Instead they may use hydrogen sulphide (purple and green sulphur bacteria) or organic compounds (purple non-sulphur bacteria) as a source of hydrogen.

**phragma** See DINOPHYCEAE.

**Phragmites cliffwoodensis** Early representative of the reed-grasses, recorded from the mid-\*Cretaceous of New Jersey, USA. It has rather uncertain affinities but if accepted would be the first of the family Poaceae (grasses).

**phragmocone** The septate shell of a cephalopod (\*Cephalopoda). The term is also applied to the septate portion of a belemnite (\*Belemnitida) skeleton which, although reduced in size, is \*homologous to the external shell of other cephalopods.

**phreatic activity** Volcanic \*eruptions generated by the interaction between hot \*magma and surface lake water, sea water, or \*groundwater. The water immediately surrounding the magma is heated and volatilized. Its expansion builds up pressure on the envelope of water surrounding it. When the pressure exceeds the confining pressure of the overlying water column the water vapour expands explosively to produce a steam-dominated, phreatic (i.e. subsurface water) eruption. Where significant amounts of magmatic

material are ejected in addition to steam the activity is said to be 'phreatomagmatic'.

**phreatic zone (zone of saturation)** The soil or rock zone below the level of the \*water-table, where all voids are saturated.

**phreatomagmatic activity** See PHREATIC ACTIVITY.

**phreatoplinian deposit** A widely dispersed layer of fine-grained, very poorly sorted, volcanic material resulting from a \*Plinian eruption, in which \*acid and intermediate \*magma reacted with water as it passed through the vent.

**Phycosiphon** An \*ichnoguild of structures made by deposit feeders that moved freely some distance below the surface.

**phyletic** Pertaining to a line of descent.

**phyletic evolution** Evolutionary change within a lineage, as a result of gradual adjustment to environmental stimuli.

**phyletic gradualism** Theory holding that \*macroevolution is merely the operation of \*microevolution over relatively long periods of time. Thus gradual changes will eventually accumulate to the point at which descendants of an ancestral population diverge into separate species, genera, or higher-level \*taxa.

**phyllic alteration** A type of \*alteration that is found in \*country rocks of copper and molybdenum \*porphyry deposits.

**phyllite** Fine-grained (less than 0.1 mm), low-grade \*metamorphic rock, of \*pelitic composition, with a well-developed \*schistosity, that often has a silky sheen due to the parallel orientation of \*phyllosilicate minerals (e.g. \*chlorite, \*muscovite, and \*sericite). Compare SLATE; SCHIST.

**phyllonite** A \*slate-like, \*dynamic metamorphic rock formed in \*fault zones and containing a penetrative \*cleavage orientated parallel to the \*fault plane. The cleavage is formed by the \*recrystallization and alignment of sheet \*silicates during fault movement.

**phyllosilicate (sheet silicate, layered silicate)** A large group of \*silicate minerals which are characterized by possessing layers of  $[\text{SiO}_4]^{4-}$  tetrahedra linked together to form a flat sheet with the composition  $[\text{Si}_4\text{O}_{10}]_n$ . This

group includes *\*micas*, *\*chlorite*, *\*clays*, *\*talc*, and *\*serpentine*, which are soft minerals and of variable but generally low density. They form at low temperatures and some, particularly the clays, replace *\*primary minerals* as a result of *\*hydrothermal* alteration or *\*weathering*. They are an essential constituent of *\*argillaceous* sedimentary rocks and some low-grade *\*metamorphic rocks*.

**phylogenetics** The taxonomical classification of organisms on the basis of their degree of evolutionary relatedness.

**phylogenetic tree** A variety of *\*dendrogram* in which organisms are shown arranged on branches that link them according to their relatedness and evolutionary descent.

**phylogenetic zone** See *LINEAGE-ZONE*.

**phylogeny** The evolutionary relationships within and between taxonomic levels, particularly the patterns of lines of descent, often branching, from one organism to another, i.e. the relationships of groups of organisms as reflected by their evolutionary history. See *TAXONOMY*.

**phylozone** See *LINEAGE-ZONE*.

**phylum** In animal *\*taxonomy*, one of the major groupings, coming below subkingdom and kingdom, and comprising superclasses, classes, and all lower *\*taxa*.

**-phyre** A suffix applied to an *\*igneous* rock which is *\*porphyritic*.

**physico-theology** See *NATURAL THEOLOGY*.

**phytogeography (floristics)** The study of the geography of plants, particularly their distribution at different taxonomic levels, i.e. family, genus, and species. Patterns of distribution are interpreted in terms of climatic and anthropogenic influence, but above all in terms of earlier continental configurations and migration routes.

**phytophagous** Feeding on plants.

**Piacenzian** The middle *\*stage* in the *\*Pliocene \*epoch*, preceded by the *\*Zanclean* (Tabianian), followed by the *\*Gelasian* (Late Pliocene), and dated at 3.6–2.588 Ma ago (Int. Commission on Stratigraphy, 2004). The corresponding European stage was originally described as a subdivision of

the (then recognized) Astian stage and is roughly contemporaneous with the upper *\*Delmontian*, *\*Repettian*, and *\*Venturian* (N. America), *\*Waipipian*, *\*Mangapanian*, and lowermost *\*Nukumaruan* (New Zealand), and upper *\*Kalimnan* and *\*Yatalan* (Australia). The type area is around Castell'Arquato, Italy.

**pick** To select one feature from among others, e.g. *\*first arrivals* on a seismic refraction record or reflection events in a seismic section. The feature so selected becomes a 'pick'. For example, on a seismic refraction record the selected first arrivals are the picks for each trace, providing *\*travel-time* data for the compilation of a *\*travel-time* graph.

**pickup** See GEOPHONE.

**pico-** From the Spanish *pico* meaning 'beak', or 'peak' (i.e. a point), a prefix (symbol p) used with SI units to denote the unit  $\times 10^{-12}$ .

**picoplankton** Marine planktonic organisms, 0.2–2.0  $\mu\text{m}$  in size, and consisting mainly of *\*bacteria* and *\*cyanobacteria*.

**picosatellite** A space satellite with a *\*wet mass* of 0.1–1.0 kg.

**Picosatellite for Remote-sensing and Innovative Space Missions (PRISM)** A *\*minisatellite* mission from the Intelligent Space Systems Laboratory of the University of Tokyo to demonstrate high- and medium-resolution imaging. The satellite was launched on 23 January 2009, from the Tanegashima Space Center, Japan, into a Sun-synchronous orbit at an altitude of 800 km.

**picrite** A strongly *\*porphyritic*, *\*olivine*-rich *\*basalt*. Picrites on the islands of Hawaii are formed during high-discharge-rate eruptions and may reflect mobilization of an olivine-rich crystal-cumulate layer near the base of the *\*magma chamber*.

**pi diagram** A stereographic projection, used in the analysis of the orientation of *\*folds*, in which poles (points representing lines) perpendicular to folded bedding ( $\pi$  poles) are plotted, and the diagram rotated on an overlay to achieve a best-fit *\*great* circle (a  $\pi$  circle). A pole to the plane of the  $\pi$  circle plane records the *\*plunge* and *\*trend* of the *\*fold axis*.

**piedmont** The tract of country at the foot of a mountain range, e.g. the Po Valley, Italy, at the foot of the Alps. The word is derived from the Italian *piemonte*, meaning 'mountain foot'.

**piedmont glacier** A lobe of *\*ice* formed when a *\*valley* glacier extends beyond its restraining valley walls and spreads out over the adjacent lowland, or *\*piedmont* zone. Much of the glacier surface is therefore at a low altitude and may show rapid *\*ablation*. An example is the Malaspina Glacier, Alaska.

**piercing fold** A *\*fold* which develops by the forcible upward movement of mud or salt *\*diapirs*.

**piezoelectricity** An electric charge induced by a flow of *\*electrons* when pressure is applied at the ends of a polar axis of a crystal which lacks a centre of symmetry (see **CRYSTAL SYMMETRY**) and which has different crystal forms at opposite ends. This property was first detected in 1881 by Pierre and Jacques Curie.

**piezometer** An *\*observation* well designed to measure the elevation of the *\*water-table* or *\*hydraulic* head of *\*groundwater* at a particular level. The well is normally quite narrow and allows groundwater to enter only at a particular depth, rather than throughout its entire length.


**piezometric surface** See **POTENTIOMETRIC SURFACE**.

**piezoremanent magnetization** The magnetization acquired when material is subjected to prolonged pressure. See also **SHOCK-REMANENT MAGNETIZATION**.

**pigeonite** A member of the *\*clinopyroxenes* with a very low calcium content and the composition  $(Ca,Mg,Fe(Mg,Fe)Si_2O_6)$ ; sp. gr. 3.30–3.46; *\*hardness* 6.0; properties similar to *\*augite*, but occurs in rapidly chilled *\*igneous* rocks. In slowly cooled rocks, pigeonite changes to an *\*orthopyroxene* as cooling proceeds.

**piggyback basin (thrust-sheet-top basin)** A type of *\*foreland basin* formed, in addition to a *\*foredeep*, on top of a thrust sheet by the progressive development of *\*thrusts* by collapse of the *\*footwall*; the newly formed thrust becomes the active thrust surface and older thrusts, with their thrust slices, are carried forward on it passively.

**piggyback thrust sequence** A *\*thrust* sequence formed by the progressive development of thrusts by collapse of the *\*footwall*; the newly formed thrust becomes the active thrust surface and older thrusts, with their thrust slices, are carried forward on it passively.



**Piggyback thrust sequence**

***Pikaia*** An early chordate (*\*Chordata*) from the *\*Burgess Shale* that was possibly related to the modern amphioxus (lancelet).

**pile** Timber, steel, or concrete sheet or column sunk into loose ground or cast in a *\*borehole* to carry vertical or horizontal loads and provide support under earth or water pressure.

**pileus** From the Latin *pileus* meaning ‘cap’, an accessory cloud occurring as a small cap on or above a cumuliform cloud. The cloud is associated with *\*cumulus* or *\*cumulonimbus*. See also CLOUD CLASSIFICATION.

**pillar and stall (bord and pillar, room and pillar; in Scotland: stoop and room)** Method of mining in which large chambers are excavated, leaving pillars of *\*ore*, rock, or *\*coal* to support the roof.

**pillar structure** Pipe-like, near vertical tubes, with diameters of a few centimetres, found in association with *\*dish structures*, and formed by the upward escape of water from liquefied *\*sediment*.

**pillow lava** Piles of elongate basaltic *\*lava* pods, having the general appearance of a stacked accumulation of discrete stone pillows, often many hundreds of metres in thickness. Each ‘pillow’ is surrounded by a chilled, fine-grained, lava skin and sags into the pillows below it. The pillows are rarely more than one metre in diameter and in cross-section each one has a

convex upper surface, radial and concentric fractures, and commonly a central cavity or tube which once fed lava to the front of the advancing finger. The morphology indicates that the pillows continued to behave as fluid bodies after the chilled carapace had formed. This provides good evidence of submarine eruption: lava entering water acquires a glassy outer skin as heat is conducted rapidly from the surface. Because water absorbs heat more readily than air, with little increase in its own temperature, the rapid surface cooling allows the molten plastic state of the pillow interior to be maintained longer than it would be in air. Pillows have been observed forming under water from lava entering the sea off Hawaii. *See also* [HYDROVOLCANIC PROCESSES](#).

**pilotaxitic** A felted mass of [\\*acicular](#) or lath-shaped crystals found in fine-grained [\\*igneous](#) rocks; the crystals may be aligned to produce a flow structure.

**pinacoid** Applied to a [\\*crystal face](#) which cuts the vertical  $c$  (or  $z$ ) [\\*crystallographic axis](#) and is parallel to the horizontal  $a$  (or  $x$ ) and  $b$  (or  $y$ ) axes. A horizontal plane of symmetry (*see* [CRYSTAL SYMMETRY](#)) repeats the face at the opposite end of the crystal. The resulting form is a pair of parallel faces with the index (001).

**pinch-and-swell** *See* [BOUDINAGE](#).

**pinger** A high-frequency, high-resolution, shallow penetration device used in marine seismic-reflection profiling and offshore [\\*engineering geophysical](#) investigations.

**pingo** Ice-cored, dome-shaped hill, oval in plan, standing 2–50 m high, and 30–600 m in diameter, developed in an area of [\\*permafrost](#). The larger examples have breached crests in which ice may be exposed. They are probably due to local freezing of water that has migrated from adjacent uplands, or to the late freezing of the ground beneath a lake. *See also* [PALSA](#).

**pinite** A fine-grained mixture of [\\*muscovite](#) and [\\*chlorite](#) with some [\\*serpentine](#) or iron oxide, which forms as an alteration product of [\\*cordierite](#). It is colourless to bluish-green and similar to [\\*mica](#) in structure and composition.

**pinnacle reef** *See* [REEF](#).

**pinnate fractures** Extension fractures that occur along a **\*fault** in **\*en échelon** arrays that point in the direction of fault movement.

**pinnular plates** See BRACHIA.

**pinnules** See BRACHIA.

***Pinus longaeva* (bristlecone pine)** Pine species from California, famous for its longevity and one of those used to develop an exceptionally long, arid-site, tree-ring chronology. The oldest living specimens date back more than 4600 years, but **\*cross-dating** these with remnants of dead bristlecone pines has extended the arid-site chronology to more than 8200 BP. A 5500-year chronology has been developed for bristlecone pines at the upper tree limit. These pines are also used to calibrate the **\*radiocarbon-dating** method to allow for fluctuations in atmospheric  $^{14}\text{C}:^{12}\text{C}$  ratios as revealed by measuring  $^{14}\text{C}:^{12}\text{C}$  ratios of individual tree rings in the long, absolutely dated, tree-ring series. *P. longaeva* is Great Basin bristlecone pine and *P. aristata* is mountain bristlecone pine. Both are used for dating. See also DENDROCHRONOLOGY.

**PIOCW** See PACIFIC- AND INDIAN-OCEAN COMMON WATER.

**Pioneer** A series of **\*NASA** spacecraft that conducted explorations of the solar system. Pioneer 10, launched on 2 March 1972, was the first spacecraft to reach the edge of the solar system, in 1987; its last signal was received on 23 January 2003, following the decay of its radioisotope power source. Pioneer 11 was launched on 5 April 1973, for Saturn; the last communication with it was in November 1995. Pioneer Venus, launched in 1978 to **\*Venus**, comprised an orbiter and probes sent to the surface.



<http://www.nasa.gov/centers/ames/missions/archive/pioneer.html>

- A NASA programme of unmanned space probes launched for planetary exploration.

**pipe** **1.** Nearly vertical, cylindrical body or opening in rock. **2.** In mining, an ore shoot at the intersection of two barren veins. **3.** At Kimberley, South Africa, pipes of diamond-bearing **\*breccia**. **4.** In sedimentology, tube often filled with mud, particularly in **\*limestones**. **5.** In volcanology, vertical

channel-ways below a **\*volcano** through which **\*magma** flows towards the surface.

**pipette analysis** A standard method for measuring the size of small particles. Sediment is stirred into suspension in a measured volume of water in a **\*sedimentation tube**, aliquots of uniform size are withdrawn by pipette at specified intervals and depths, oven-dried, and weighed. The grain diameter ( $D$ ) is then calculated by  $D = \sqrt{C/x/t}$ , where  $C$  is a constant depending on the particle density and density and viscosity of the fluid,  $x$  is the depth (in cm) from which the particles are withdrawn,  $t$  is the time elapsed (in seconds), and  $x/t$  is the settling velocity.

**piprake (needle ice)** Columnar ice found beneath individual stones or patches of earth in the **\*periglacial** environment. It is a result of the relatively high thermal conductivity of such materials, leading to freezing under them when the temperature falls. Heaving of less than 0.1m may occur by this process.

**Piripauan** See **CAMPANIAN**.

**pisoid** See **PISOLITH**.

**pisolith** (*adj.* **pisolitic**, **pisoid**) A spherical to subspherical, inorganic **\*carbonate** particle, larger than 2  $\mu\text{m}$  in diameter, and in some cases as much as 10 cm in diameter, characterized by an internal concentric lamination. Some pisoliths are said to form in the same manner as **\*ooids**, but others ('vadose pisoliths'), form in subaerial environments in **\*calcrete** profiles, Pisoliths should not be confused with oncoids (**\*oncolites**) which although superficially similar, are of organic origin.

**pisolitic** See **PISOLITH**.

**pistacite** See **EPIDOTE**.

**piston corer** See **HYDRAULIC CORER**.

**piston sampler** See **PEAT-BORER**.

**Pistosaurus grandaevus** A likely ancestor for all **\*plesiosaurs** although it is known only by its skull. It bears some similarities to both **\*nothosaurs** and plesiosaurs, and is found in rocks of Middle **\*Triassic** age.

**pitch (rake)** The angle made by a *\*lineation* with the *\*strike* of the surface on which it occurs.

**pitchblende** See URANINITE.

**pitchstone** A glassy *\*igneous* rock, rather like *\*obsidian* but with a waxy, resinous *\*lustre* owing to the absorption of water.

**pitting** The digging of a pit; in Scotland, mining an outcrop by making shallow pits. In exploration, sampling of *\*alluvial \*sediments* by shallow (down to 15–20 m) trial pits.

**pixel** 1. A picture element. Commonly the smallest component of a *\*multispectral* image as determined by a single optic fibre. 2. In *\*remote sensing*, a single sample of data in an image. A pixel has a spatial attribute corresponding to its location within the image and the area of ground represented, and a spectral attribute corresponding to the *\*intensity* of a particular wavelength. See also PIXEL COLOUR.

**pixel colour** In *\*remote sensing*, the colour of each *\*pixel*. Pixel colour is dependent on three parameters: the intensity, which is the brightness of the colour; the saturation, which is the perpendicular distance to the *\*achromatic* axis (the closer to the achromatic axis the more pastel the colour, the further away the more vivid the colour); and the *\*hue*. See also INTENSITY-HUE-SATURATION PROCESSING.

**pixie** A *\*transient luminous* event that is a pinpoint of light, lasting less than 16 milliseconds, on the upper surface of a convective cloud that has produced *\*gnomes*.

**PKP-wave** See SEISMIC-WAVE MODES.

**placental mammals** See EUTHERIA.

**placer deposit** Deposit of materials (e.g. gold, diamonds, tin, or platinum) that has been concentrated by mechanical action. Placer minerals generally have high density and resistance, and therefore may concentrate during various types of *\*weathering*.

**place value** The economic importance attached to the location of a mineral deposit. Minerals or metals with a high intrinsic value, e.g. diamonds or gold, have a low place value as transport costs add little to the eventual

market price, so they may be worked anywhere on Earth. In contrast, sands and gravels have a high place value and must be worked near the place of use.

**placic horizon** Subsurface **\*soil horizon**, formed most readily in humid tropical or cold conditions, that is **\*cemented** by iron and organic matter, by iron and manganese, or by iron alone.

**Placodermi** Class of archaic, jawed (**\*Gnathostomata**), and heavily armoured fish, which appeared in the **\*Devonian** and which were virtually extinct by the end of that period. They were rather diverse in body form. They all possessed a **\*head shield** formed by bony plates. Pectoral and pelvic fins appear to have been present. Most of them (e.g. *Coccosteus*) were bottom-dwelling fish, with a depressed body terminating in a heterocercal tail.

**Placodontidae** **\*Triassic** **\*euryapsid** reptiles, which were specialized shellfish feeders. Some were heavily armoured and strongly resembled turtles; while the more lightly armoured varieties, e.g. *Placodus*, were analogous in general form to the **\*nothosaurs**, except for modifications relating to their molluscan diet.

**placoid scale (dermal denticle)** Type of scale that comprises the basic unit of the hard skin cover of sharks. It consists of a hard base embedded in the skin and a spiny process (cusp); these are covered by a hard, enamel-like substance (vitrodentine), and the scale projects outwards and backward. The bulk of the denticle consists of **\*dentine** surrounding a central pulp cavity. Unlike the scales of **\*bony fish**, placoid scales stop growing after they reach a certain size and new denticles are added instead.

**plaggen** A man-made **\*soil horizon** more than 500 mm deep, resulting from long-continued manuring, often enriched by phosphate.

**plagioclase feldspar** One of the most important rock-forming silicate **\*minerals** with the general formula  $(\text{Na,Ca})(\text{Al})_{1-2}(\text{Si})_{2-3}\text{O}_8$ . There is a **\*solid solution** series (see **PLAGIOCLASE SERIES**) between the two **\*end-members** albite (Ab) ( $\text{NaAlSi}_3\text{O}_8$ ) and anorthite (An) ( $\text{CaAl}_2\text{Si}_2\text{O}_8$ ), and the percentage of calcium end-member present is used to subdivide the series into a number of individual minerals: albite (0–10 mol % An); oligoclase (10–30 mol % An); andesine (30–50 mol % An); labradorite (50–70 mol %

An); bytownite (70–90 mol % An); and anorthite (90–100 mol % An). Albite; sp. gr. 2.61; **\*hardness** 6.0–6.5; is whitish, **\*vitreous**; **\*tabular** or irregular; with two **\*cleavages** {010} and {001} meeting at almost right angles on the (100) face; and occurs in **\*acid \*igneous** rocks and **\*spilites**. Oligoclase; sp. gr. 2.64; **\*triclinic**; cleavage perfect basal {001}, good {010}; is similar to albite except in the percentage An it contains. Andesine; sp. gr. 2.66; triclinic; cleavage perfect basal {001}, good {010}; has properties similar to those of albite but tends to occur in more **\*intermediate** igneous rocks. Labradorite; sp. gr. 2.67; crystals thin and **\*tabular**, flattened parallel to (010); cleavage perfect basal {001}, good {010}; is greyish-white but may show iridescence due to lattice imperfections on cleavage faces and occurs in **\*basic** igneous rocks. Bytownite; sp. gr. 2.72; triclinic; crystals often tabular prismatic but normally form irregular grains; cleavage perfect basal {001}, good {010}; also greyish-white, is a constituent of basic and **\*ultrabasic** igneous rocks. Anorthite; sp. gr. 2.75; triclinic; crystals tabular and prismatic but normally form irregular grains; cleavage perfect basal {001}, good {010}; also greyish-white, occurs in basic and ultrabasic igneous rocks and in metamorphosed **\*limestones**. These plagioclase feldspars cannot be distinguished from one another in hand specimens, but under the microscope their **\*extinction** angles vary; this is a useful property in their identification, together with the nature of the multiple twinning (see **CRYSTAL TWINNING**) which is a very characteristic feature and serves to distinguish them from **\*alkali feldspars**. **\*Exsolution** of potassium-feldspar in a plagioclase feldspar host is called ‘antiperthite’.

**plagioclase series** **\*Isomorphous** series of **\*feldspars** ranging in composition between albite ( $\text{NaAlSi}_3\text{O}_8$ ) and anorthite ( $\text{CaAl}_2\text{Si}_2\text{O}_8$ ). Between the two **\*end-members** there is a continuous **\*solid solution** series: albite, oligoclase, andesine, labradorite, bytownite, anorthite. Usually, the higher the temperature of formation the more likely the feldspar is to be calcium-rich, whereas at lower temperatures sodium-rich plagioclase tends to form. The preservation of electrical neutrality in the plagioclases is achieved by concomitant substitution of  $\text{Ca}^{2+}$  and  $\text{Al}^{3+}$  by  $\text{Na}^+$  and  $\text{Si}^{4+}$ . See **PLAGIOCLASE FELDSPAR**.

**plagiogranite** See **GRANODIORITE**.

**plana** See **PLANUM**.

**planar cross-stratification** See CROSS-STRATIFICATION.

**planar slide** See TRANSLATIONAL SLIDE.

**planar slump** See TRANSLATIONAL SLIDE.

**planation surface** 1. Synonym for \*erosion surface. 2. The final stage produced by the erosion of folded \*sedimentary rocks, when inverted relief is planed across by erosion.

**plane bed (flat bed)** A near-horizontal surface of \*sand or gravel. Two types of plane bed are found. Upper-stage plane beds are produced by the intense transport of \*sediment by high-velocity, shallow flows (upper-flow-regime conditions), and characterized by primary current lineation on the \*sediment surface. Lower-stage plane beds are produced only in coarse sands and gravels by flow conditions broadly similar to those which generate current \*ripples in finer sand. The lower-stage plane bed exhibits a series of shallow scours on the sediment surface. The accumulation of plane-bedded sediment gives rise to an internal \*sedimentary structure of horizontal \*lamination.

**plane of projection (equatorial plane)** The horizontal surface on which points on a spherical projection can be represented in two dimensions. A \*stereographic projection can be compared to a slice through the Earth's equator as seen from the N. or S. Pole, and for this reason is often called the 'equatorial plane', and its circumference is called the 'primitive circle'. Since the plane passes through the centre of the sphere its circumference is a \*great circle, the angle between the plane of projection and the N–S axis of the sphere being 90°.

**plane of symmetry** See CRYSTAL SYMMETRY.

**plane-polarized light (PPL)** As light travels, it normally vibrates in all directions at right angles to the line of transmission. If a strongly absorbing crystal, e.g. \*polaroid or \*tourmaline, is placed in the light path, the rays are strongly absorbed in all directions except one, and the rays of light that emerge are confined to this one plane of vibration; i.e. the light is 'plane polarized'. Polarization may also result from double refraction (see NICOL PRISM) or reflection.

**plane strain** The *\*strain* state which occurs when a reference sphere is extended along its X axis and shortened along its Y axis, leaving its Z axis unchanged. This strain changes the shape from a sphere to a triaxial ellipsoid.

**planetary boundary layer** The layer of the lower atmosphere extending to a height of about 500 m above the ground, in which frictional effects of the underlying surface generate *\*turbulence*. *See also* TURBULENT BOUNDARY LAYER.

**planetary geodesy** The measurement of the shape and gravitational field of the Moon and other bodies in the *\*solar system*, other than Earth.

**planetary geology (planetary geoscience, astrogeology, exogeology)** The scientific study of planetary surfaces, employing the techniques of geology. It was first used formally to establish the stratigraphic sequence on the visible face of the *\*Moon* by *\*Shoemaker* and Hackman in 1962, but was used informally by earlier workers, such as G. K. *\*Gilbert* and R. B. Baldwin, to interpret lunar features as lava flows, craters, etc.

**planetary geomorphology** The study of surface structures and the processes that formed them on the Moon and other bodies in the *\*solar system*, other than Earth.

**planetary geoscience** *See* PLANETARY GEOLOGY.

**Planet-B (NOZOMI)** A Japanese *\*ISAS* mission that was launched in 1998. The spacecraft suffered a series of failures and the mission was finally abandoned in December 2003. The aim had been to study the martian atmosphere and the effect of the *\*solar wind* from orbit.

**planetismal** Small, solid body, a few kilometres in diameter, accreted by gravitational attraction, after condensation of *\*mineral \*phases* from the primitive solar nebula. The *\*solar-system* planets are thought to have formed by the further accretion of planetismals.

**planetismal hypothesis** A cataclysmic theory, propounded early in the 20th century, to explain the origin of the *\*solar system*. This was attributed to tidal interactions produced by a close encounter between the *\*Sun* and a star. The immense gaseous tides produced in the Sun pulled matter away, in

the direction of the star, to form a long, gaseous filament. This then broke into several parts, each contracting to form a planet.

**plane wave** A wavefront which has no effective curvature because of its distance from the source. At a very large range from the source, the degree of curvature over a short local distance is negligible and the wavefront can be regarded as planar.

**planèze** Sloping triangular facet on the flank of a **\*volcano**, underlain and protected by **\*lava**. Its form is a result of the radial dissection by streams of a complex volcanic cone.

**planispiral** Applied to the condition in which a univalve gastropod (**\*Gastropoda**) shell is coiled (see **COILING**) in a single horizontal plane and the diameter increases away from the axis of coiling. Many cephalopod (**\*Cephalopoda**) shells are also planispiral.

**planitia** (*pl.* **planitiae**) A plain that is low relative to the surrounding terrain. A typical example is Hellas Planitia on **\*Mars** (42° S, 293° W), a probable degraded impact basin, about 2000 × 1500 km.

**plankton** (*adj.* **planktonic**) Minute aquatic organisms that drift with water movements, generally having no locomotive organs. The phytoplankton (plants) comprise mainly **\*diatoms**, which carry out **\*photosynthesis** and form the basis of the aquatic food-chains. The zooplankton (animals) which feed on the diatoms may sometimes show weak locomotory powers. They include **\*protozoans**, small crustaceans (**\*Crustacea**), and in early summer the larval stages of many larger organisms. Plankton are sometimes divided into net-plankton (more than 25 µm diameter) and nanoplankton, which are too small to be caught in a plankton net (see **NANO-**; **NANOFOSSIL**).

**planktonic geochronology** The use of **\*planktonic** organisms, e.g. globigerinid **\*foraminiferids** or microscopic **\*algae**, to provide a relative dating of **\*sediments** deposited in marine waters. **\*Radioactive-decay** methods applied to planktonic organisms may also yield an absolute date or information on palaeoclimates.

**Planolites** An **\*ichnoguild** of structures made by deposit feeders that moved freely through shallow burrows.

**planosols** A reference soil group in the [\\*World Reference Base](#) for Soil Resources. Planosols have an E [\\*soil horizon](#) that results from prolonged submersion beneath stagnant water within 100 cm of the surface. The E horizon often has a markedly different texture from the overlying horizons.

**Plantae (Metaphyta)** A kingdom that includes all the plants. The earliest true plants were probably unicellular green algae (see [ALGA](#)), which first appeared in the [\\*Precambrian](#). The [\\*Bryophyta](#) (mosses and liverworts) are known from the [\\*Devonian](#), and the first recorded vascular plants ([\\*Tracheophyta](#)) date from the [\\*Silurian](#).

**planum** (*pl. plana*) A [\\*plateau](#) or high-elevation plain on the surface of an extraterrestrial body.

**plasma** A low-density, high-temperature, completely ionized gas, consisting of free atomic nuclei and free electrons. Overall it is electrically neutral. It is sometimes referred to as the ‘fourth state of matter’.

**plasma engine** A propulsion device used in spacecraft in which electrons are stripped from atoms of a neutral gas, creating a cloud of positive [\\*ions](#). The ions and electrons are then accelerated by an electric field to generate thrust. *Compare* [ION ENGINE](#).

**plasma instrument** An instrument carried by spacecraft that measures [\\*ions](#) and [\\*electrons](#).

**plasma wave sub-system (PWS)** An instrument carried on the [\\*Galileo](#) spacecraft that measures the properties of varying electric fields at frequencies of 5–5.6 MHz and magnetic fields at 5–160 kHz in the atmosphere of [\\*Jupiter](#). [\\*Plasma](#) particles are bound to the magnetic field, and motions within the plasma perturb the electric and magnetic fields around them producing changes known as ‘plasma waves’. Scientists use the PWS to investigate the way plasma waves and radio emissions control the scattering and/or loss of trapped radiation in the jovian atmosphere, and to derive fundamental plasma parameters, such as [\\*electron](#) density.

**plastic deformation** A deformation process which proceeds elastically at low [\\*stress](#) values and becomes viscous when a critical stress value is reached. Plastic deformation occurs in rocks under conditions of high pressure and temperature, producing a permanent alteration of their shape but without failure by rupture.

**plasticity index** See ATTERBERG LIMITS.

**plastic limit** See ATTERBERG LIMITS.

**plastron** 1. The lower, bony shell of a turtle. 2. See IRREGULAR ECHINOIDS.

**plate** 1. A segment of the \*lithosphere, which has little volcanic or seismic activity but is bounded by almost continuous belts (known as \*plate margins) of \*earthquakes and, in most cases, by volcanic activity and young subsea or subaerial mountain chains. Most Earth scientists consider there are currently seven large, major plates (the \*African, \*Antarctic, \*Eurasian, \*Indo-Australian or Indian, \*North American, \*Pacific, and \*South American Plates). There are also several smaller plates (e.g. the \*Arabian, \*Caribbean, \*Cocos, \*Nazca, and \*Philippine Plates) and an increasingly long list of \*microplates (e.g. the \*Gorda, \*Hellenic, and \*Juan de Fuca Plates). The positions of the boundaries of some present-day plates are disputed, particularly within and adjacent to \*collision zones, e.g. the Alpine–Himalayan belt, so it is not surprising that very little agreement has been reached about the histories of plates in the geologic past. 2. A general term applied to plane pieces of \*skeletal material usually formed from calcium carbonate. Plates occur in groups of several types, e.g. the \*delthyrium in some brachiopods (\*Brachiopoda) is closed by a pair of \*deltidial plates. 3. The outer covering of a crinoid (\*Crinoidea) body, which consists of a series of rows of plates. 4. The bony covering, often fused to the ribs, on the upper and lower surfaces of the body of a turtle. The upper surface is the ‘carapace’, the lower is the ‘plastron’.

**plateau basalt** An extensive, thick, smooth flow or succession of flows of high-temperature, fluid \*basalt erupted from fissures, flooding topographic lows, and accumulated to form a plateau. The Deccan Traps in India covers 260 000 km<sup>2</sup> and the Columbia River Plateau basalt, in Washington State, USA, covers 130 000 km<sup>2</sup> and is more than 1800 m thick. Individual flows may have volumes of the order of 100 km<sup>3</sup>. The original area of the Thulean Plateau which formed in the north-eastern Atlantic Ocean region 30 Ma ago, was  $1.8 \times 10^6$  km<sup>2</sup>. The lava plains are built up by many thousands of individual flows from numerous, coalescing, \*shield-type volcanoes with extremely low angles of slope.

**plate bearing test** Static test to measure deformability in terms of the theory of elasticity (see [ELASTIC REBOUND THEORY](#)). Values may be assigned to the ground for [\\*Young's modulus](#) and [\\*Poisson's ratio](#).

**plate boundary** See [PLATE MARGIN](#).

**plate kinematics** The study of the geometry of [\\*plate margins](#) and the relative velocities of [\\*plate motions](#).

**plate margin (plate boundary)** The boundary of one of the [\\*plates](#) that form the upper layer (the [\\*lithosphere](#)) and together cover the surface of the Earth. Plate margins are characterized by a combination of tectonic and topographic features: oceanic [\\*ridges](#), [\\*Benioff zones](#), young fold mountains, and [\\*transform faults](#). Plate margins are of three main types: (a) [\\*constructive](#) margins where newly created lithosphere is being added to plates which are moving apart at oceanic ridges; (b) [\\*convergent margins](#) which can be either [\\*destructive margins](#), where one plate is carried down into the [\\*mantle](#), beneath the bordering plate, at a [\\*subduction zone](#), or a [\\*collision zone](#), where two [\\*island arcs](#) or continents, or an arc and a continent, are colliding; or (c) [\\*conservative margins](#), where two plates are moving in opposite directions to each other along a [\\*transform fault](#). All three margins are seismically active, with volcanic activity at constructive and destructive margins. Some plate margins exhibit features of more than one of the three main types and are known as [\\*combined plate margins](#). See [PLATE TECTONICS](#); [SEA-FLOOR SPREADING](#).

**plate motions** The movement of tectonic [\\*plates](#) is expressed in terms of rotations relative to a [\\*Euler pole](#). Such motions during the last 200 million years have been determined mainly from [\\*magnetic](#) anomaly patterns in the ocean basins. Determination of older motions is based on [\\*palaeomagnetic](#) studies and evidence for continental collisions and separations.

**plate stratigraphy** The study of sedimentary strata in order to reconstruct the geographic positions and water depths to which they have been subjected as a consequence of the movement of the [\\*plates](#) on which they lie.

**plate tectonics** The unifying concept that has drawn [\\*continental drift](#), [\\*sea-floor spreading](#), seismic activity, crustal structures, and volcanic activity (see [VOLCANICITY](#)) into a coherent model of how the outer part of

the **\*Earth** evolves. The theory proposes a model of the Earth's upper layers in which the colder, brittle, surface rocks form a shell (the **\*lithosphere**) overlying a much less rigid **\*asthenosphere**. The shell comprises several discrete, rigid units (tectonic **\*plates**) each of which has a separate motion relative to the other plates. The **\*plate margins** are most readily defined by present-day **\*seismicity**, which is a consequence of the differential motions of the individual plates. The model is a combination of continental drift and sea-floor spreading. New lithospheric plates are constantly forming and separating, and so being enlarged, at **\*constructive margins** (ridges), while the global circumference is conserved by the **\*subduction** and recycling of material into the **\*mantle** at **\*destructive margins** (trenches). This recycling results in andesitic volcanism and the creation of new **\*continental crust**, which has a lower density than the **\*oceanic crust** and is more difficult to subduct. Many features of the Earth's history are explicable within this model which has served as a unifying hypothesis for most of the Earth sciences. Previous mountain systems are now recognized as the sites of earlier subduction, often ending with continental crustal collision: the movement of plates has been used with varying success in interpreting **\*orogenic belts** as far back as the early **\*Proterozoic**. **\*Plate motions** are driven by mantle **\*convection** and are likely to have occurred throughout Earth history, although the resultant surface features are likely to have changed with time. See RIDGE-PUSH; SLAB-PULL.

**platform** See SHELF.

**platform conodonts** See CONODONTOPHORIDA.

**platy** Applied to **\*minerals** which develop a **\*crystal** form consisting of thin, leaf-like layers. The **\*mica** group of minerals provides a good example; the mineral splits along **\*cleavage** planes which are parallel to rows of alkali atoms in the crystal structure.

**platykurtic** See KURTOSIS.

**play 1.** The combination of factors that makes possible the accumulation of oil and gas in a particular area. **2.** A **\*shale** formation that contains significant accumulations of **\*natural gas**.

**playa (salina)** The lowest part of an **\*intermontane** basin or **\*bolson**, which is frequently flooded by runoff from adjacent highlands or by local rainfall.

\*Sediments consist largely of \*colloids, \*clays, and \*evaporites, e.g. \*halite, \*gypsum, and sodium sulphate. The surface is generally flat, with \*mudflats and locally small \*dunes. The name (*playa* is the Spanish word for 'beach' and *salina* for 'salt-mine') was first applied to the arid basin-and-range province between the Colorado Plateau and Sierra Nevada, in the western USA, but is now used to describe such areas throughout the world.

**Playfair, John** (1748–1819) Scottish mathematician, who popularized and promoted the plutonist and uniformitarian theories of James \*Hutton, rewriting his work to make it more easily readable (1802).

***Plectronoceras cambria*** One of the first \*nautiloids, which was small and horn shaped. As its name suggests, the species is discovered in rocks of \*Cambrian age, in Europe and Asia.

**Pleiades-HR** A constellation of two satellites of the French Space Agency that provide high-resolution, panchromatic, and multispectral optical imagery over the globe, with every part of the surface covered daily. Pleiades-1A was launched on 17 December 2011 and Pleiades-1B on 2 December 2012, both from Kourou, French Guiana, into Sun-synchronous orbits at an altitude of 694 km, phased 180° apart.

**Pleistocene** The first of two \*epochs of the \*Quaternary, held conventionally to have lasted from approximately 1.806 Ma ago until the beginning of the \*Holocene 11 430 years ago. The epoch is marked by several glacial and \*interglacial episodes in the northern hemisphere.



<http://www.geologypage.com/2014/05/pleistocene-epoch.html>

- Pleistocene Epoch.

**Pleistocene refugium** Favourable area where species have survived periods of glaciation during the \*Pleistocene era. Such species are termed \*relicts.

**Pleistogene** The current informal \*period of the \*Cenozoic \*era, comprising the \*Pleistocene and \*Holocene \*epochs. The Pleistogene began 1.806 Ma ago. The International Commission on Stratigraphy regards this as an archaic period name and prefers to extend the \*Neogene to the present day.

**pleochroic halo** The dark, strongly pleochroic (see PLEOCHROISM) zone often seen around certain radioactive inclusions (e.g. \*zircon, \*apatite, and \*sphene) in some \*minerals (e.g. \*biotite, \*tourmaline, \*amphibole, \*chlorite, \*muscovite, \*cordierite, and \*fluorite). The halo is created by the interaction of alpha particles with the atoms of the crystal \*lattice. Some haloes contain concentric rings with radii of 10–15 µm, the radii indicating the various kinetic energies of the different sets of alpha particles.

**pleochroism** In optical microscopy, the differential absorption of light in different crystallographic orientations by a coloured mineral when it is rotated on the stage in \*plane-polarized light. The mineral may show variations in shade of the same colour or even a different colour. See also DICHROISM.

**pleonaste** See SPINEL.

**plesiomorph** Primitive (of a character state); the opposite of \*apomorph.

**plesiomorphic** Applied to features that are shared by different groups of biological organisms and are inherited from a common ancestor. The term means ‘old-featured’ and the features to which it is applied were formerly called ‘primitive’.

**plesion** In taxonomy, a group of superfamilies within a suborder.

**Plesiorycteropus** A genus of apparently burrowing, ant-eating mammals, known only from incomplete material found in sub-Recent deposits in Madagascar. Individuals weighed about 10 kg or less. *Plesiorycteropus* has been placed in a new order, \*Bibymalagasia.

**Plesiosauroidea (plesiosaurs)** Suborder of aquatic \*reptiles which enter the fossil record in the late \*Triassic, and which are common in many \*Jurassic and \*Cretaceous \*sediments. In appearance they were likened by \*Buckland to ‘a snake strung through the body of a turtle’, and some grew up to 15 m in length. There were also short-necked types, as well as the swan-necked. The former are defined as the superfamily Plesiosauroidea (e.g. *Plesiosaurus* and *Muraenosaurus*), the latter as the super family Pliosauroida (e.g. *Pliosaurus* and *Trinacromerum*), the two superfamilies making up the suborder.

**plesiosaurs** See PLESIOSAUROIDEA.

**pleura** See PLEURON.

**pleural** See PLEURON.

**pleuron** (*pl.* **pleura**, *adj.* **pleural**) The lateral portion of a single thoracic segment in a trilobite (*\*Trilobita*). Each pleuron is indented by an oblique pleural furrow and the outer ends are downturned. The inner edges are attached to a central axial ring, so that each thoracic segment consists of a central axial ring and a pair of laterally placed pleura.

**plicate** Folded or wrinkled.

**Pliensbachian** A *\*stage* in the European Early *\*Jurassic*, preceded by the *\*Sinemurian*, followed by the *\*Toarcian*, and roughly contemporaneous with the lower *\*Ururoan* (New Zealand) and dated at 189.6–183 Ma ago (Int. Commission on Stratigraphy, 2004). See LIAS.

**Plinian eruption** An explosive volcanic *\*eruption* of *\*pyroclastic* ejecta forming an eruption column which may be up to 55 km high, dispersing ejecta over an area of 500–5000 km<sup>2</sup>. The eruption column has the distinctive, spreading, branched shape of the stone pine (*Pinus pinea*) native to the Mediterranean region. A Plinian eruption may produce thick, airfall, *\*pumice* deposits, or the eruption column may collapse to generate a *\*pyroclastic flow*. This type of eruption is named after Pliny the elder, who died at Pompeii in AD 79 during the eruption of Vesuvius that destroyed the city.

**plinthic horizon** In the *\*World Reference Base* for Soil Resources classification, an iron-rich *\*soil horizon* more than 15 cm thick and containing more than 25% plinthite.

**plinthite** Portion of *\*mineral soil* containing a large proportion of iron and aluminium oxides, *\*clay*, and *\*quartz*, which has developed through a combination of *\*leaching* and *\*gleying* in well-weathered tropical soils. On drying, it changes irreversibly to an ironstone *\*hardpan*.

**plinthosols** A reference soil group in the *\*World Reference Base* for Soil Resources. Plinthosols have a *\*plinthic horizon* that hardens on exposure to air within 50 cm of the surface.

**Pliocene** The more recent (5.332–1.806 Ma ago) of the two **\*Neogene** **\*epochs**, comprising the **\*Zanclean**, **\*Piacenzian**, and Gelasian ages.



<http://www.geologypage.com/2014/05/pliocene-epoch.html>


- Pliocene Epoch.

**plough mark** A depression, in some cases many metres deep and tens of metres wide, made on the sea floor by the base of an iceberg that is driven by wind and tide. Many plough marks are found near the edges of **\*continental shelves** in high latitudes, marking the passage of ancient icebergs.

**plug** See **VOLCANIC PLUG**.

**plunge** The angle between a line and a horizontal datum plane; the term is commonly used in respect of the inclinations of **\*fold axes**.

**plunging breaker** Wave that breaks by plunging forward in the direction of motion, so that its crest falls into the preceding trough and encloses a pocket of air. The wave-form is then lost. Typically it occurs when a fairly low wave approaches a steep shingle beach. It is characterized by a smooth forward and under face, and a convex top.



**Plunging breaker**

**plus–minus method (Hagedoorn method)** A method of interpreting **\*seismic refraction** profiles over irregular layers whose slope angles do not exceed  $10^\circ$ . It uses both forward and reverse shooting to obtain matched **\*time–distance graphs**. The plus component of the method allows the perpendicular depths to the refractor to be calculated; the minus component is used to determine the **\*seismic velocity** in the refracting medium.

**Pluto** Formerly the ninth and outermost planet of the *\*solar system*, but now classified as a minor planet, Pluto's orbit is an average 39.44 AU from the *\*Sun* and highly eccentric, sometimes carrying it inside the orbit of *\*Neptune*. Its distance from Earth ranges from  $4293.7 \times 10^6$  km to  $7533.3 \times 10^6$  km. Pluto is much smaller than the *\*Moon*, with a radius of 1137 km; volume  $0.616 \times 10^{10}$  km<sup>3</sup>; mass  $0.0125 \times 10^{24}$  kg; mean density 2050 kg/m<sup>3</sup>; surface gravity 0.66 (Earth = 1); visual albedo 0.3; black-body temperature 42.7 K. The atmosphere is very thin, with a surface pressure of about 0.003 bar, and composed of methane and nitrogen. The average surface temperature is about 50 K. It has one satellite, *\*Charon*, so large that some astronomers consider Pluto and Charon a double-body system. It was visited by NASA's New Horizons space probe in July 2015. Pluto was discovered in 1930 by Clyde Tombaugh.

**pluton** General term applied to a body of *\*intrusive \*igneous* rock, irrespective of its shape, size, or composition.

**plutonic** A loosely defined term with a number of current usages. Many petrologists use it to describe *\*igneous* rock bodies which have crystallized at great depth, there being no real agreement about what is meant by 'great depth'. Other petrologists use the term to describe any igneous rock that forms in largish *\*intrusions* (i.e. excluding *\*dykes* and *\*sills*). 'Plutonic' has also been used to describe the origin of *\*magmas* and gas derived from near the base of the *\*crust* or in the upper *\*mantle*.

**plutonism** A theory of the formation of the Earth, prominently advocated by *\*Hutton*, and based on cycles of the growth, decay and regeneration of land masses. The forces involved in these cycles were heat and gravity, the heat deriving from subterranean fires. This generated volcanic rocks and granites, which rose molten from the Earth's interior, to form new land.

**pluvial period** Prolonged phase of markedly wetter climate that occurs in a normally dry or semi-arid area.

**pneumatolysis** (*adj.* **pneumatolytic**) Changes in rock mineralogy and chemistry that are initiated by the action of a hot, chemically active, gaseous solution derived from a *\*magma* during its final stages of crystallization. The gaseous solution, rich in *\*volatile* elements such as fluorine, chlorine, boron, and hydrogen, is released from the crystallizing interior of the *\*granite* body and rises through cracks and fissures in the

crystalline and cooler upper carapace to initiate greisenization and \*[tourmalinization](#) of the roof-zone granite.

**poached soil** See [PUDDLED SOIL](#).

**Podolskian** A Russian \*stage (308–307.2 Ma ago) in the \*[Pennsylvanian](#) \*epoch and a sub-stage of the \*[Moscovian](#), preceded by the \*[Kashirskian](#) and followed by the \*[Myachkovskian](#).

***Podopteryx mirabilis*** Named ‘marvellous foot wing’, this is the species that is the probable ancestor of the \*[pterosaurs](#). Unlike them it was a glider, relying on the large membrane stretched between its hind legs and tail. It is known from the Lower \*[Triassic](#) of Soviet Kirgizstan.

**podsol** See [PODZOL](#).

**podsolization** See [PODZOLIZATION](#).

**podzol (podsol)** \*Soil profile formed at an advanced stage of \*[leaching](#) by the process of \*[podzolization](#), and identified by its acid \*[mor](#) \*[humus](#), \*[eluviated](#) and bleached E \*[soil horizon](#), and an iron-coloured B horizon, enriched with a variety of translocated materials. See also [SPODOSOLS](#).

**podzolic soils** A major soil group in the classification system developed by the Soil Survey for England and Wales, podzolic soils generally form in acidic environments and have accumulations of organic matter and of iron and aluminium oxides. There may be an iron pan in the B \*[soil horizon](#).

**podzolization (podsolization)** An advanced stage of \*[leaching](#), podzolization is the process of removal of iron and aluminium compounds, \*[humus](#), and \*[clay minerals](#) from the surface \*[soil horizons](#) by an organic leachate solution, and the deposition of some of these translocated materials in lower B horizons.

**podzols** A reference soil group in the \*[World Reference Base](#) for Soil Resources. Podzols have a \*[spodic](#) B horizon, and usually a bleached surface \*[soil horizon](#) and an iron pan at depth.

**Pogonophora (beard worms)** A phylum comprising deep-sea worms, first encountered in early \*[Cambrian](#) rocks but discovered only in the 20th century. Beard worms bear a superficial resemblance to \*[Polychaeta](#) and it has recently been proposed that they are actually very highly specialized

polychaetes. They live at great depths inside chitinous tubes they secrete for themselves in soft substrates, often in the vicinity of **\*hydrothermal vents**. The body is **\*coelomate**, partly segmented, has 'bristles' (chaetae), and is divided into three parts, the anterior crowned with tentacles. Their most remarkable feature is the complete absence of a gut. This has led to difficulties with classification because it is impossible to distinguish the ventral and dorsal surfaces. The animals are believed to obtain nourishment through a chemosymbiotic association (see **CHEMOSYMBIOSIS**) with bacteria. There are two groups, one found near vents and **\*cold seeps**, the other occurring widely in all oceans.

**poikil-** From the Greek *poikilos*, meaning varied or variegated, a prefix with the same meaning.

**poikilitic** Applied to the **\*texture** produced when several orientated or unorientated crystals are enclosed within a larger crystal in an **\*igneous rock**. The larger crystals have more widely separated nuclei than the enclosed crystals, and may grow faster, thus enclosing surrounding grains. Where **\*augite** encloses a number of **\*plagioclase feldspar** crystals the poikilitic texture is given the specific name of 'ophitic texture'.

**poikiloaerobic** See **DYSAEROBIC**.

**poikiloblast** (*adj.* **poikiloblastic**) A large crystal which encloses several smaller, orientated or unorientated crystals in a **\*metamorphic rock**.

**poikilotherm (exotherm)** A vertebrate animal that has no behavioural or metabolic means of regulating its body temperature, which therefore fluctuates with the temperature of its surroundings. Fish and **\*amphibians** are poikilothermic. Poikilotherms are often termed 'cold-blooded', although when exposed to sunshine their bodies may be warmer than their surroundings. *Compare* **HOMIOOTHERM**.

**poikilotopic** Applied to the **\*fabric** of a **\*sedimentary rock** in which coarse crystals of **\*cement** enclose a number of smaller, detrital grains.

**point bar** See **BAR**.

**point counting** A method for determining the proportion of an area covered by a particular type of object. The area is covered by a grid of points and the underlying object identified for each point. The estimate for the

proportion of the total area covered by each type of object is then given by:  $p_i \approx \frac{h_i}{N}$ , where  $p_i$  is the true proportion of objects type I,  $h_i$  is the number of points in contact with an object type I, and  $N$  is the total number of points counted.

**point group** See CRYSTAL CLASS.

**point load index ( $I_s$ )** The force needed to fracture a sample of rock between conical points:  $I_s = P/D^2$ , where  $P$  is force and  $D$  is the distance between the points, both at failure.  $I_s$  is related to **\*uniaxial compressive strength** (approximately equal to  $I_s \times 24$ ).

**point load tester** Equipment for measuring **\*point load index**, consisting of a jack which closes  $60^\circ$  cones across a sample (usually a piece of drill **\*core**) and a gauge to measure pressure just before failing.

**point source** A source of energy which is considered to originate from a point and to have no effective size in relation to its observed effects, e.g. an explosive shot, a **\*current electrode** in **\*electrical** resistivity sounding, etc.

**Poisson distribution** In statistics, a discrete **\*probability distribution** which is applied to the number of times an event occurs.

**Poisson's ratio ( $Y$ )** The ratio of latitudinal to longitudinal **\*strain**, which describes the extent to which a material is distorted in a direction perpendicular to an applied **\*stress**:

$$Y = e_{(\text{normal to } \sigma_1)} / e_{(\text{parallel to } \sigma_1)},$$

where  $Y$  is Poisson's ratio,  $e$  the strain, and  $\sigma_1$  is the maximum **\*principal stress**. It is named after Simeon Poisson (1781–1840).

**polar air** **\*Air** mass originating in latitudes  $50\text{--}70^\circ$ , which may have a maritime or continental source. Maritime polar air has high relative humidity, is warmed in its passage equatorward or over warmer seas, and becomes unstable. Continental polar air is stable in its source region and is associated with very cold surface conditions in winter, e.g. Siberian air moving across Europe.

**polar-air depression** Non-frontal \*depression in the northern hemisphere, which typically results from the southward movement of unstable polar or arctic \*maritime air along the eastern side of a large-scale meridional high-pressure ridge. *See also* COLD LOW.

**polar climate** Climatic type associated with regions inside the Arctic and Antarctic Circles. A gradation of climatic characteristics exists towards the poles, from \*tundra conditions to those of perpetual frost. *See also* KÖPPEN CLIMATE CLASSIFICATION.

**polar-desert soil** \*Mineral soil without identifiable \*soil horizons, and with almost no surface \*humus. It is associated with arid, polar-desert environments where precipitation is less than 130 mm annually, plant cover less than 25%, and thawed, active soil is 20–70 mm deep.

**polar front** The main boundary line between polar and tropical \*air masses along which \*depressions develop in mid latitudes, especially over the oceans. The front is in general displaced equatorwards in winter and polewards in summer, though large displacements in either direction take place over shorter periods in individual sectors of the hemisphere.

**polar-front jet stream** The \*thermal wind observed in different positions over middle and higher latitudes at heights of 10–13 km, and associated with the boundary zone between polar and tropical air above the \*polar front. Maximum velocity averages 60m/s, but can be twice this in extreme cases. The jet is typically discontinuous but at times can extend almost around the globe. It is more persistent in winter in response to the stronger temperature gradient between north and south than in summer. The jet is related to the development of surface frontal depressions of middle latitudes.

**polar glacier** *See* GLACIER.

**polarimetry** The measurement of the degree of polarization of light reflected from a planetary surface. The degree of polarization depends on the composition of the surface material. Polarimetry measurements have been used to obtain information on particle size distribution in the atmosphere of Titan, for example.

**polarity** The direction of evolutionary change. The polarity of different states of a *\*character* means whether these are *\*primitive* or *\*derived*.

**polarity chron** The basic time interval in the *\*magnetostratigraphic timescale*, during which the Earth's *\*geomagnetic field* is constantly, or predominantly, of one polarity, e.g. the *\*Gauss* normal polarity chron and the *\*Matuyama* reversed polarity chron. The duration of polarity chrons is variable, but generally is longer than 0.1 Ma. Polarity chrons may be interrupted by *\*polarity subchron(s)*, and grouped to form a *\*polarity superchron*. The term 'polarity chron' has been proposed by the *\*ISSC* to replace polarity epoch; however, at present both terms are in use. The corresponding *\*chronostratigraphic unit* is *\*polarity chronozone*.

**polarity chronozone** The polarity *\*chronostratigraphic unit* denoting all the rocks, with or without magnetic minerals (see *FERROMAGNETIC*), formed during a specific *\*polarity chron*.

**polarity epoch** See *POLARITY CHRON*.

**polarity event** See *POLARITY SUBCHRON*.

**polarity excursion** A geomagnetic *\*polarity subchron*.

**polarity interval** 1. An *\*informal* term for any of the polarity *\*chronostratigraphic units*: *\*polarity chronozone*, *\*polarity subchronozone*, or *\*polarity super chronozone*. 2. A general, informal term, either spatial or temporal, describing the intervening span between *\*polarity reversals*.

**polarity reversal, geomagnetic** A change by 180° in the direction of the Earth's magnetic field. See *GEOMAGNETIC FIELD*.

**polarity subchron** A very short time interval (generally less than 0.1 Ma) of alternate polarity occurring within a *\*polarity chron*, e.g. the *\*Olduvai* normal subchron within the *\*Matuyama* reversed polarity chron. The term 'polarity subchron' has been proposed by the *\*ISSC* to replace '*\*polarity event*'; however, at present both terms are in use. The corresponding polarity *\*chronostratigraphic unit* is *\*polarity subchronozone*. See also *MAGNETOSTRATIGRAPHIC TIMESCALE*; *POLARITY SUPERCHRON*.

**polarity subchronozone** A polarity *\*chronostratigraphic unit* denoting all the rocks, with or without magnetic minerals (see *FERROMAGNETIC*),

formed during a *\*polarity subchron*.

**polarity subzone** See POLARITY ZONE.

**polarity superchron** The longest polarity time interval in the *\*magnetostratigraphic timescale*, comprising a number of *\*polarity chrons*, and indicating a period of time (varying between 30 and 100 Ma) during which the polarity of the Earth's *\*geomagnetic field* has a distinct bias. The bias may be towards a *\*normal field*, a reversed field, or an evenly alternating field (called a mixed field), e.g. the present *\*Cretaceous–\*Cenozoic* mixed-polarity superchron. The corresponding polarity *\*chronostratigraphic unit* is the *\*polarity superchronozone*.

**polarity superchronozone** The largest polarity *\*chronostratigraphic unit*, denoting all the rocks, with or without magnetic minerals (see FERROMAGNETIC), formed during a specific *\*polarity superchron*.

**polarity superzone** See POLARITY ZONE.

**polarity timescale** See MAGNETOSTRATIGRAPHIC TIMESCALE.

**polarity transition period** The time taken for a change of polarity of the *\*geomagnetic field*. It is thought to take 3000–5000 years for directional changes and about 12 000 years for associated intensity changes.

**polarity zone** The basic magnetic *\*lithostratigraphic unit*, defined by the measured magnetic polarity of a body of rock. A polarity zone is bounded above and below by a *\*polarity reversal* horizon or a *\*polarity transition zone*. Polarity zones may comprise a number of polarity subzones and be grouped into polarity superzones. When used formally the term is often capitalized, e.g. *\*Gauss Polarity Zone*. See MAGNETOZONE.

**polarization, electrode** The accumulation of *\*ions* around an electrode, causing the accumulation of a *\*charge*.

**polarization colours** See INTERFERENCE COLOURS.

**polarized radiation** *\*Electromagnetic radiation* which is orientated in a single plane.

**polarizer** A piece of *\*Polaroid* in a transmitted- or *\*reflected-light microscope* which is inserted into the light path between the light source

and the **\*mineral** section. Observations are made in **\*plane-polarized light** and the light emerging is confined to either an E–W or N–S plane of vibration.

**polarizing microscope** A microscope fitted with an **\*analyser** and **\*polarizer** to allow specimens to be examined in polarized light. The petrological (petrographic) microscope has a light source below the **\*stage** so that light can be transmitted through the specimen. It is used to examine **\*thin sections** of **\*rocks** and **\*minerals**. The ore (reflected light) microscope uses an incident light source, allowing light to be reflected from the polished surface of an **\*ore** (**\*opaque**) mineral. The system of lenses, diaphragms, rotating stage, polarizer, and analyser is broadly similar in both types of microscope, and dual microscopes are available for use with either transmitted or reflected light.

**Polaroid** The trade name of a plastic sheet impregnated with an organic iodide compound which strongly absorbs light in one **\*vibration direction** and allows it to pass freely in the other vibration direction. The compound, iodocinchonidine sulphate, was first discovered by W. D. Herapath in 1852 and was named ‘herapathite’ in his honour. In 1928 E. H. Land combined the compound with plastic to produce the Polaroid sheet. It is now used in the **\*polarizer** and **\*analyser** of all **\*polarizing microscopes** to produce **\*plane-polarized** and cross-polarized (see **CROSSED POLARS**) light respectively.

**polar orbit** An **\*orbit** which is inclined at 45° or more to the equatorial plane. *Compare* **EQUATORIAL ORBIT**.

**Polar Orbiting Passive Atmospheric Calibration Sphere (POPACS)** A US **\*nanosatellite** mission to measure changes in density of the auroral zone of the upper atmosphere caused by solar stimuli such as flares and coronal mass ejections. The mission involved deploying three aluminium spheres of 10 cm diameter into an elliptical **\*polar orbit** and tracking them as they make repeated **\*perigee** passes in the upper atmosphere. The satellite was launched on 29 September 2013, from California, into orbit at an altitude of about 340 km perigee, about 1510 km **\*apogee**.

**polar stratospheric cloud (PSC)** A cloud, consisting of ice crystals, which occurs in the **\*stratosphere** late in winter over Antarctica and, less commonly, over the arctic. The water vapour that freezes may be derived

from the dissociation of methane and the cloud forms in the very still, very cold air of a vortex over the pole (the less common occurrence of such clouds in the northern hemisphere is owing to the generally higher temperature of arctic stratospheric air and the briefer duration of the vortex). The reactions that deplete polar stratospheric ozone in the late winter and early spring take place on the surface of the ice crystals in polar stratospheric clouds.

**Polar units** See MARTIAN TERRAIN UNITS.

**polar wander path** The successive positions of the \*palaeomagnetic pole. The individual pole position is calculated assuming the observed field is that of an axial geocentric dipole. As the Earth's \*axis of rotation is fixed relative to the \*ecliptic, the changing positions of the pole, as a function of time, are due to the motion of the tectonic \*plate from which rock samples were obtained. The pole path is therefore, more strictly, an apparent polar wander path. Sudden changes in the pole path ('hairpins') are usually caused by continental plate collisions and 'superintervals', between hairpins, correspond to the motion of a plate with little or no collisional interactions with other plates.

**polder** A low-lying, flat area reclaimed from the sea and protected by embankments or dykes; especially along the Netherlands North Sea coast.

**pole of a face (face pole)** The normal (line perpendicular) to a \*crystal face, and the point at which the normal intersects the \*plane of projection. In a \*stereographic projection of a \*crystal, the crystal is imagined to lie at the centre of a sphere and a pole of each face intersects the surface of the sphere at a point. These points are then projected to either the N. or S. pole of the sphere, on to the horizontal (equatorial) plane to produce an accurate, two-dimensional representation of the faces and \*interfacial angles of the crystal.

**pole of rotation (Euler pole)** A point on the Earth's surface which defines a line through the centre of the Earth about which the relative motion of two \*plates may be described.

**'Polflucht' ('flight from the poles')** A concept that was invoked by \*Wegener to explain his ideas about \*continental drift. He suggested that a differential gravitational force (the Eötvös force), caused by the flattening

of the Earth at the poles, would cause continental masses to drift towards the equator. This force is now known to be far too weak to cause continental movements.

**polished section** A specimen of an *\*ore* (*\*opaque*) mineral after it has been prepared for examination under a reflected-light microscope by light reflected from its polished surface. A sample of the mineral is mounted in a cold-setting epoxy resin, sawn to produce a flat surface, and then inverted, ground, and polished in a number of stages using diamond-impregnated fluids and a rotating lap fitted to a polishing machine. Conventional 0.03 mm *\*thin sections* are prepared in the same way to give 'polished thin sections', used in the identification of opaque minerals. Polished sections are also required in the *\*electron-probe microanalyser* to enable analyses of minerals to be carried out.

**polished stone value** See AGGREGATE TESTS.

**polishing relief** In the preparation of a *\*polished section* the relief caused when harder *\*minerals* stand out slightly above the surface of softer minerals. Although not desirable, the presence of such relief may help to determine the relative hardness of adjacent minerals. See KALB LIGHT LINE.

**polje** Large, flat-floored depression bounded by steep valley walls and found in a *\*karst* environment. It is classically described for the Dinaric region of Yugoslavia, where it may result from faulting or from solutional processes controlled by a local base level.

**pollen** Collectively, the mass of microspores (pollen grains) produced by the anthers of a flowering plant (*\*angiosperm*) or the male cones of a *\*gymnosperm*. Different pollen types are described according to their shapes, apertures, etc. Furrows on the surface of the pollen grain are called 'sulci' (sing. sulcus) and monosulcate pollen has a single sulcus. Tricolpate pollens have three, furrow-like, germinal apertures arranged 120° apart and there are many variants of this type.

**pollen analysis** The study of fossil *\*pollen* and *\*spore* assemblages in *\*sediments*, especially when reconstructing the vegetational history of an area. The outer coat (*\*exine*) of a pollen grain or spore is very characteristic for a given family, genus, or sometimes even species. It is also very resistant to decay, particularly under *\*anaerobic* conditions. Thus virtually

all spores and pollen falling on a rapidly accumulating sediment, anaerobic water, or *\*peat* are preserved. Since both pollen and spores are generally widely and easily dispersed, they give a better picture of the surrounding regional vegetation at the time of deposition than do macroscopic plant remains, e.g. fruits and seeds, which tend to reflect only the vegetation of the immediate locality. With careful interpretation, pollen analysis enables examination of climatic change and human influence on vegetation, as well as sediment dating and direct study of vegetation character. The technique has also been applied, more controversially, to the pollen and spore contents of modern and *\*fossil \*soil profiles*. See also [PALYNOLOGY](#).

**pollen-assembly zone** See [POLLEN ZONE](#).

**pollen diagram** Standardized pictorial summary of the *\*pollen* record for a particular location. The vertical axis represents depth, and the proportions or absolute amounts of the various pollen types occurring at different levels are shown by bar histograms or by points on a continuous curve. Conventionally, similar patterns are grouped together on the diagram with arboreal types (i.e. trees) shown first followed in turn by shrubs, herbs, and *\*spores*.

**pollen zone (pollen-assembly zone)** Characteristic *\*pollen-and-\*spore* assemblage classically considered indicative of a particular type of climate which was assumed to be typical of a fairly extensive geographic region. Changes from one group of pollens to another characteristic assemblage are used to define pollen-zone boundaries. The standard British (Godwin, 1940) and very similar European *\*post-glacial* (i.e. late *\*Devensian* and *\*Flandrian*) chronology recognizes eight major pollen zones, Zones I–III being the characteristic late-glacial sequence: *\*Older Dryas (I)*, *\*Allerød (II)*, and *\*Younger Dryas (III)*. More recently, based on N. American work, the importance of regional variation in the typical zone floras has been acknowledged, and a more flexible approach to pollen-zone definition applied. Pollen-assembly zones are defined in terms of their pollen and spore profiles alone for a particular site, and initially without reference to or matching with the standard zone models with their strong climatic links. This has enabled local changes, often anthropogenic rather than climatic, to be elucidated more clearly. The scheme has largely been abandoned, however, owing to the lack of synchronicity even within the British Isles.

Pollen assemblage diagrams are now constructed for each individual site; regional comparisons and correlations can then be made on the basis of chronology.

**poloidal field** A magnetic field with radial and tangential components. The **\*geomagnetic field** detected at the Earth's surface is of this form and contrasts with the unmeasurable **\*toroidal field**.

**poly-** From the Greek *polus* meaning 'many', a prefix meaning 'many'.

**polyanions** Stable and complex **\*anions**, of various sizes, formed by aluminate and silicate anions linked by shared oxygen **\*ions**.

**Polychaeta (bristleworms)** (phylum **\*Annelida**) Class of annelid worms which possess distinct **\*metameric** segmentation. All have bristly parapodia (movable, paired appendages) on each body segment. Eyes may be present. Males and females occur. Most are marine, although some occur in fresh water and on land. The group first appeared in the **\*Cambrian**, and impressions of polychaete worms are among the fauna of the **\*Burgess Shale**. However, polychaetes are represented in the fossil record largely by **\*burrows** (e.g. **\*Skolithos**), tubes, and **\*scolecodont** assemblages.

**polycyclic landscape (polyphase landscape)** A land-form or landscape that has been acted on by the erosional processes associated with two or more partially completed cycles of erosion (see **DAVISIAN CYCLE**). A diagnostic feature is an abrupt break of slope (see **KNICK POINT**) in the profiles of both rivers and hillslopes.

**polygenetic** See **CONGLOMERATE**.

**polygonization** See **HOT WORKING**.

**polyhaline water** See **HALINITY**.

**polyhalite** Mineral,  $K_2Ca_2Mg(SO_4)_4 \cdot 2H_2O$ ; sp. gr. 2.8; **\*hardness** 2.5–3.0; **\*triclinic**; normally flesh-pink to brick-red, and translucent; silky to resinous **\*lustre**; usually occurs as fibrous or lamellar masses; **\*cleavage** {100}, parting {010}; occurs in bedded **\*evaporite** deposits and is one of the last minerals to be precipitated from saline waters, due to its high solubility; tastes bitter.

**polymetallic sulphide** A mineral deposit with three or more metals in commercial quantities; common metals include Cu, Pb, Zn, Fe, Mo, Au, and Ag. It may occur in magmatic, volcanogenic, or hydrothermal environments.

**polymetamorphism** Repeated episodes of heating and deformation (**\*metamorphism**) acting upon a rock system.

**polymictic** 1. Applied to lakes (e.g. in high altitudes in the tropics) whose waters are circulating virtually continuously. If periods of stagnation occur they are very short. 2. Applied to a **\*conglomerate** which contains **\*clasts** of many different rock types. *Compare* OLIGOMICTIC.

**polymineralic** See ROCK.

**polymorph** One of several possible **\*crystal** forms of an element, compound, or mineral, all of which have the same chemical composition. See POLYMORPHISM.

**polymorphic** Occurring in several different forms.

**polymorphic minerals** **\*Minerals** which have the same composition, but different atomic lattices, so their physical properties may differ.

**polymorphic transformation** Change in the atomic structure of a **\*mineral** (but not in its chemical composition) to produce a different form. In reconstructive transformation there are large energy barriers to overcome in order to break down and re-form bonds, and transition tends to be slow, e.g. **\*graphite** to **\*diamond**. In displacive transformation angles are changed between bonds, but no bonds are broken, so little energy is required, e.g. alpha to beta **\*quartz**.

**polymorphism** Ability of elements or compounds to exist in more than one crystal form, with each having the same chemical composition but different physical properties due to differences in the arrangement of atoms. Examples are **\*graphite** and **\*diamond** (both C); alpha and beta **\*quartz** (both SiO<sub>2</sub>); and **\*calcite** (**\*hexagonal**) and **\*aragonite** (**\*orthorhombic**), both forms of CaCO<sub>3</sub>. The terms 'dimorphism' and 'trimorphism' describe (respectively) two and three different forms.

**polypedon (soil individual)** Two or more contiguous *\*pedons*, which are all within the defined limits of a single *\*soil series*.

**polyphase landscape** See POLYCYCLIC LANDSCAPE.

**polyphyletism** (*adj.* **polyphyletic**) The occurrence in *\*taxa* of members that have descended via different ancestral lineages. True polyphyletism has traditionally been distinguished from errors of *\*classification*, especially at the higher taxonomic levels, where organisms, as a result of *\*convergent* or *\*parallel* evolution, have been placed wrongly in the same natural group; but modern phyletic taxonomists would hold that any taxon found to be polyphyletic is unnatural, and so an 'error', and must be disbanded.

**Polyplacophora (chitons)** (phylum *\*Mollusca*) Class of molluscs in which seven or eight *\*dorsal \*plates*, generally composed of calcium carbonate, are enclosed by a marginal girdle. The plates articulate with one another and overlap to a varying extent. The anterior and posterior plates differ from the others, the anterior plate often being ribbed. Polyplacophora are entirely marine, and first appeared in the Upper *\*Cambrian*.

**polysynthetic twin** See ALBITE TWIN.

**polytaxic times** A period of high biological diversity among marine organisms, associated with high sea level (highstand), equable climates, little convection in ocean waters, and an abundance of available niches.

**polytetrahedron** In *\*crystallography*, a *\*tetrahedron* in which negative and positive forms are developed to different degrees in the same crystal, together with the possibility of raised secondary faces on each principal tetrahedron face. In *\*mineralogy*, the term may refer to the ability of  $\text{SiO}_4$  tetrahedra in silicate structures to polymerize, giving a variety of structures as a result of oxygen-sharing. These may result in chains (*\*pyroxenes*), double chains (*\*amphiboles*), sheets (*\*micas*), and three-dimensional framework silicates (*\*quartz* and *\*feldspars*).


**Pomeranian** One of a series of *\*Weichselian* recessional *\*moraines* in northern Germany, which postdates the Frankfurt series of moraines and predates the Velgart series.

**ponente** Regional term for a westerly wind that blows in the Mediterranean.

**Pongola** See MESOARCHAEAN.

**ponor** A natural opening in some \*karst landscapes through which a lake or stream flows partially or completely underground.

**pool-and-riffle** An alternation between a deep zone (the pool) and a shallow zone (the riffle) along the sand and/or gravel bed of a stream. The pool-to-pool spacing is about 5–7 times the width of the channel. The sequence is found in both straight and \*meandering channels; in the latter case the pool occurs in the meander bend on the concave side, and the riffle between bends.



**Pool-and-riffle**

**poorly washed biosparite** See FOLK LIMESTONE CLASSIFICATION.

**POPACS** See POLAR ORBITING PASSIVE ATMOSPHERIC CALIBRATION SPHERE.

**Poperinge** See MOERSHOOFD.

**pop-shooting (secondary blasting)** A method of secondary blasting used to break large boulders into pieces of manageable size. A hole is drilled to just beyond the middle of the boulder so that the charge is central. The method is fairly quiet and economical.

**pop-up** The relatively uplifted section of a \*hanging wall which is formed by \*back thrusting. The structure is bounded by the back thrust and by the major \*thrust from which the back thrust originated.

**Porangan** A \*stage (47–44 Ma ago) in the \*Eocene \*epoch of New Zealand, underlain by the \*Heretaungan, overlain by the \*Bortonian, and roughly contemporaneous with the mid \*Lutetian stage.

**porcelain jasper** See JASPER.

**porcellanite** A highly siliceous *\*sedimentary rock* with a dull *\*lustre*, porous texture, and *\*conchoidal fracture*, similar in general appearance to porcelain. These rocks are less hard and vitreous than *\*cherts*. Porcellanites include certain *\*biogenic*, siliceous, deep-sea *\*sediments* and some fine-grained, *\*recrystallized*, acid *\*tuffs*.

**pore** A void surrounded completely by soil or rock materials and created by the packing of mineral and organic particles. Pores can be filled by any proportion of air or water.

**pore fluid pressure** Pressure of a fluid which fills the *\*pores* in a saturated soil or rock. It can indicate the degree of consolidation of an earthwork, zero pressure indicating complete consolidation. Compare PORE-WATER PRESSURE.

**pore space** The total continuous and interconnecting void space in the bulk volume of soil or rock.

**pore-water pressure** The pressure exerted on its surroundings by water held in *\*pore spaces* in rock or soil. The pressure is positive when a soil is fully saturated, and is then proportional to the height of the water measured in an open tube (a *\*piezometer*) above the point of interest. A buoyancy effect is achieved and the *\*shear* strength of the soil is reduced. The pressure is zero when the soil voids are filled with air, and is negative when the voids are partly filled with water (in which case surface-tension forces operate to achieve a suction effect and the shear strength of the soil is increased). Compare PORE FLUID PRESSURE.

**Porifera (Spongiaria, sponges)** A phylum of multicellular animals that are not included in the *\*Animalia*. Sponges are sessile, benthic, filter feeders with a bag-like body, a central cavity, and an outer surface pierced with tiny openings (ostia), through which water enters, and an upper, larger opening (osculum), through which it leaves. Many sponges have a skeleton that is either horny, or composed of spongin, or consists of calcareous or siliceous *\*spicules*. They may be fossilized as a complete skeleton or as isolated spicules. Their *\*fossil* record dates from the *\*Cambrian* (or possibly from the *\*Precambrian*) and they are a common source of *\*biogenic silica*.

**porosity** Absolute porosity is the total of all void spaces present within a rock, but not all these spaces will be interconnected and thus able to contain and transmit fluids. The **\*effective** porosity is thus defined as the proportion of the rock which consists of interconnected **\*pores**. Porosity is expressed as a percentage of the bulk volume of the rock. *See also* CHOQUETTE AND PRAY CLASSIFICATION; FENESTRAL POROSITY; FRACTURE POROSITY; MOLDIC POROSITY; OOMOLDIC POROSITY.

**porosity and permeability determination** **\*Porosity** is measured from rock plugs in a porosimeter. The method involves the extraction of air from the **\*pores** in the rock by a vacuum, and measurement of the bulk volume of the rock. The porosity is calculated by the expression: % porosity = (volume of gas extracted/bulk volume of the rock sample) × 100. Permeability is measured in a **\*permeameter**, by determining the pressure drop ( $P_1 - P_2$ ) from a fluid of known **\*viscosity** ( $\mu$ ) and **\*flow** rate ( $Q$ ), across a rock sample of known cross-section ( $A$ ) and length ( $L$ ). Permeability ( $K$ ) is then determined by **\*Darcy's law**:  $K = Q\mu L / (P_1 - P_2)A$ .

**porphyrin** Heterocyclic derivative of a porphin, which is composed of four linked rings, each containing nitrogen (a tetrapyrrole ring structure). As such it is capable of combining with a variety of metals and so forms part of the structure of many important biological molecules, including haemoproteins, chlorophyll, cytochromes, and vitamin B<sub>12</sub>.

**porphyritic** *See* PHENOCRYST.

**porphyroblast (metacryst)** A large, well-formed (**\*euhedral**) crystal which grew *in situ* during metamorphic **\*recrystallization** and which is surrounded by a finer-grained **\*groundmass** of other metamorphic crystals.

**porphyroclast** A large, intact, **\*mineral** fragment or **\*clast** which is surrounded by a finer-grained, crushed **\*groundmass** produced during **\*dynamic metamorphism**. The porphyroclast represents a relict of the original rock which has escaped crushing during the deformation process.

**porphyry** A medium- to coarse-grained, **\*intrusive**, **\*felsic**, **\*igneous** rock which is conspicuously **\*porphyritic**, containing more than 25% **\*phenocrysts** by volume. The phenocryst **\*mineral** is usually **\*alkali**

feldspar. The term can be used as a suffix to a specific name, e.g. *\*quartz porphyry*.

**porphyry copper** See PORPHYRY DEPOSIT (1).

**porphyry deposit** 1. Porphyry copper occurs as large copper deposits centred around *\*stocks* of *\*intermediate* to *\*acid*, *\*porphyritic*, *\*igneous* rocks. Most occur in *\*Mesozoic* and *\*Tertiary* *\*orogenic belts*. They show concentric zones of *\*minerals*; for example at Bingham, Utah, there is an inner zone of Cu/Mo and an outer zone of Pb/Zn/Ag. The deposits are also characterized by extensive *\*alteration* halos. Most deposits are 3–8 km across and several kilometres deep. They consist of disseminated *\*chalcopyrite* and other *\*sulphides* mined on a large scale from *\*open* pits. The *\*ore* is low grade (less than 1% Cu) but of great economic importance. It was probably formed by a sudden release of *\*volatiles* near the surface, with shattering of the enclosing rocks. 2. A deposit of molybdenum-bearing ore, usually *\*molybdenite*, associated with rocks of porphyritic texture, and containing minor amounts of copper. It is formed in a similar manner to porphyry copper deposits, but the orebodies are frequently shaped like inverted cups over the progenitor intrusive, e.g. those in the mineral belt of the USA. 3. A gold-enriched porphyry copper deposit frequently associated with *\*island-arc* environments. At the present time no true porphyry gold deposits are known, only those with *\*co-product* gold and copper.

**porphyry gold** See PORPHYRY DEPOSIT.

**porphyry molybdenum** See PORPHYRY DEPOSIT.

**Portia (Uranus XII)** One of the lesser satellites of *\*Uranus*, with a diameter of 55 km. It was discovered in 1986.

**Portlandian** The youngest *\*stage* (146–142 Ma ago) of the *\*Jurassic* in Britain, overlain by *\*sediments* of the *\*Purbeckian* stage and resting in turn on those of the *\*Kimmeridgian*. It is characterized in southern England by *\*mollusc-rich* *\*limestones*. See also MALM; TITHONIAN; VOLGIAN.

**positive inversion** See INVERSION.

**post-** From the Latin *post* meaning ‘after’, a prefix meaning ‘after’, ‘behind’, or ‘later’.

**post-depositional remanent magnetization** The magnetization acquired by a sedimentary rock after deposition and before undergoing \*metamorphism, mostly associated with \*chemical remanent magnetization as \*ferromagnetic minerals grow during diagenetic alteration (see **DIAGENESIS**), but it also results from the physical rotation of magnetic grains associated with the movement of fluids and gases through the rock.

**post-deuteric alteration** Changes to the \*fabric or composition of an \*igneous rock after the completion of \*deuteric changes at elevated temperatures.

**postdisplacement** In animal \*evolution, an alteration in the \*ontogeny of a descendant such that some developmental process commences later than it did in its ancestor, and may not have been completed by the time the animal reaches maturity.

**posterior (biological)** In animals, applied to the part of the body that is to the rear when the animal is moving forward; in bipedal animals the \*dorsal surface is the posterior surface. In plants, that part of the flower or axillary bud that faces towards the flower stalk or main stem.

**post-glacial** See **FLANDRIAN**; **HOLOCENE**.

**post-perovskite** A form of the magnesium \*silicate mineral  $\text{MgSiO}_3$  that is identical in composition to its more common \*perovskite (a group of minerals based on  $\text{CaTiO}_3$ ) form, but much denser. In perovskite, the atoms form a three-dimensional bonding structure, and in post-perovskite they form stacks of two-dimensional sheets. The existence of post-perovskite was discovered in 2004 when a group at the Tokyo Institute of Technology led by Kei Hirose reproduced the high temperature and pressure found in the lowermost \*mantle, in the \*D-layer close to the boundary between the mantle and outer \*core. Post-perovskite was the predominant mineral present and its density accounts for the previously mysterious changes in speed of \*seismic waves passing through this region.

**postseismic slip** See **AFTERSLIP**.

**post-tectonic** Applied to a process or event, e.g. the emplacement of \*plutons, which occurs after deformation. Compare **PRE-TECTONIC**; **SYNTECTONIC**.

**postzygapophyses** See VERTEBRA.

**potassium–argon dating (K–Ar method)** Geologic dating technique based on the *\*radioactive* decay of potassium ( $^{40}\text{K}$ ) to argon ( $^{40}\text{Ar}$ ). This potassium *\*isotope* has a half-life (see DECAY CONSTANT) of 1.3 billion ( $10^9$ ) years, making this a valuable dating method. The minimum age limit for this dating method is about 250 000 years.

**potassium–calcium dating** A *\*radiometric* dating method based on the decay of  $^{40}\text{K}$  to stable  $^{40}\text{Ca}$ . This is not a generally useful technique because  $^{40}\text{Ca}$  is the most abundant naturally occurring stable *\*isotope* of calcium (96.94%). The formation of radiogenic  $^{40}\text{Ca}$  atoms in a rock or mineral therefore increases its abundance only slightly. The ratio  $^{40}\text{Ca}:^{44}\text{Ca}$  (96.94%:2.08%) can be used to determine the amount of radiogenic  $^{40}\text{Ca}$  present, although the dominance of naturally occurring  $^{40}\text{Ca}$  makes it rather insensitive. Furthermore the determination of the isotopic composition of calcium by *\*mass spectrometry* is made difficult by the low efficiency of ionization of calcium atoms in a thermionic source, and by fractionation of isotopes during that process. Because of these disadvantages the  $^{40}\text{K}:^{40}\text{Ca}$  method of dating is really viable only for minerals that are strongly enriched in potassium and depleted in calcium, such as *\*micas* in *\*pegmatite* and *\*sylvite* in *\*evaporite* rocks.

**potassium feldspar** See ALKALI FELDSPAR.

**potential electrode** An electrode used as a ground contact in a voltage-measuring circuit. In electrical-resistivity and *\*induced-polarization* surveying, two potential electrodes are used, with a variety of possible *\*electrode configurations*; in *\*spontaneous* potential measurements *\*non-polarizable* (e.g. porous pot) electrodes are commonly used.

**potential energy** See ELEVATION POTENTIAL ENERGY; HYDRAULIC HEAD.

**potential evapotranspiration (PE)** The amount of water that would evaporate from the surface and be transpired by plants were the supply of water unlimited. It is calculated from the mean monthly temperature, with corrections for day length, and was devised by C. W. *\*Thornthwaite* as part of his system of *\*climate* classification (see THORNTHWAITE CLIMATE CLASSIFICATION). From *PE* minus precipitation an approximate index can

be calculated of the extent to which the water available for plants falls short of the amount they are capable of transpiring. Compare ACTUAL EVAPOTRANSPIRATION.

**potential instability (convective instability)** Atmospheric condition in which otherwise stable air would become unstable if forced to rise, e.g. over high ground, thereby reaching its saturation point. Large \*cumulus with much precipitation often results from the forced uplifting of such air. See also INSTABILITY; STABILITY.

**potential reserve** See RESERVE.

**potential temperature ( $\Phi$ )** 1. The temperature that \*mantle material would have were it raised to a specified depth where it was subjected to lower pressure. Decompression lowers the temperature and the potential temperature determines the depth at which melting will occur. 2. The temperature a volume of water would have if it were raised to a specified level, usually the ocean surface. 3. The temperature a \*parcel of air would have if it were brought to the surface, warming at the \*dry adiabatic lapse rate.

**potentiometric surface (piezometric surface)** A hypothetical surface defined by the level to which water in a confined \*aquifer rises in observation \*boreholes. In practice, the potentiometric surface is mapped by interpolation between borehole measurements. As with the \*water-table in an unconfined aquifer, the slope of the potentiometric surface defines the \*hydraulic gradient and the horizontal direction of \*groundwater flow. See HYDRAULIC HEAD.

**pot-hole** An approximately hemispherical depression made in the bedrock of a river channel by stones or boulders that have been spun rapidly by eddies. Each stone is washed away only to be soon replaced by another, so the drilling action is powerful and ceaseless.

**Potsdam gravity** The value of \*gravitational acceleration as measured at Potsdam, Germany, and previously used as a worldwide standard. It is now replaced by the \*International Gravity Formula.

**Potter's flood-peak formula** See FLOOD-PEAK FORMULAE.

**Poundian** A *\*stage* of the *\*Ediacaran* period, dated at about 570–542 Ma ago (Int. Commission on Stratigraphy, 2004), preceded by the *\*Wonokan* and followed by the Early *\*Cambrian \*epoch*.


**powder photograph** See X-RAY POWDER PHOTOGRAPH.

**powder technology** The study of the properties, behaviour, and uses of finely particulate materials.

**powellite** See TUNGSTATES.

**power-law creep** *\*Creep* resulting from the movement of *\*crystal* dislocations into systematic patterns, usually polygonal, within a stress field.

**pozzolan** Originally, a *\*tuff* found near Pozzuoli, in southern Italy, that was used in mortars in Roman times. Today the most widely used pozzolan is pulverized fly ash. A pozzolan is any substance containing silicic acid ( $\text{H}_4\text{SiO}_4$ ) that has the properties of a *\*cement* when mixed with calcium hydroxide ( $\text{Ca}(\text{OH})_2$ ). The cement is a calcium silicate hydroxide ( $\text{CaH}_2\text{SiO}_4 \cdot 2\text{H}_2\text{O}$ ):



**p-parameter** A measure of the prolateness of an ellipsoid: if  $p$  is less than 0.9 the ellipsoid is oblate, if  $p$  is greater than 1.1 it is prolate.

**ppb** Parts per billion ( $10^9$ ).

**PPL** See PLANE-POLARIZED LIGHT.

**ppm** Parts per million.

**PPR** See PHOTOPOLARIMETER-RADIOMETER.

**p-process** See PROTON-ADDING PROCESS.

**praecipitatio** From the Latin *praecipitatio* meaning ‘I fall’, a supplementary cloud feature in which precipitation from the cloud base falls to the ground. The feature is usually seen with *\*cumulus*, *\*cumulonimbus*, *\*stratus*, *\*stratocumulus*, *\*altostratus*, and *\*nimbostratus*. See also CLOUD CLASSIFICATION.


**praedichnia** \*Trace fossils comprising structures that have resulted from predation.

**Praghian (Siegenian)** A \*stage of the Early \*Devonian \*epoch, dated at 411.2–407 Ma ago, preceded by the Lochkovian and followed by the \*Emsian (Int. Commission on Stratigraphy, 2004).

**Prandtl number** A dimensionless number, formulated by the German mathematician and aerodynamicist Ludwig Prandtl (1875–1953), that describes the ratio of \*kinematic viscosity to thermal diffusivity. The Prandtl number (Pr) is equal to  $\nu/\alpha$ , where  $\nu$  is kinematic viscosity and  $\alpha$  is thermal diffusivity, both in units of  $\text{m}^2/\text{s}$ .

**Pratt, John Henry** (?1811–1871) Pratt was a mathematician and physicist, and Archdeacon of Calcutta. He attempted to calculate the mass of the Himalayas, after discrepancies were found in the Trigonometrical Survey of India. He proposed that high mountains had a lower density than other parts of the \*crust, and went on to develop his own version of the theory of \*isostasy. See PRATT MODEL.

**Pratt model** A model for the \*lithosphere that accounts for \*isostatic anomalies by assuming there is a level of compensation that lies at a constant depth everywhere. Below the level of compensation all rocks have the same density, but above it density decreases as topographic elevation increases. For a column of material anywhere on Earth, the mass lying above the level of compensation will be the same, and  $\rho_c h$  a constant, where  $\rho_c$  is the density of the crust and  $h$  the topographic elevation. If the rocks are on the sea bed, then  $\rho_c h + \rho_w d$  is a constant, where  $\rho_w$  is the density and  $d$  the depth, of sea water. See also AIRY MODEL.



**Pratt model**

**pre-** From the Latin *prae* meaning ‘before’, a prefix meaning ‘in front of’, ‘earlier than’, ‘more important than’, or ‘better than’.

**pre-adaptation** Adaptation evolved in one adaptive zone which, quite by chance, proves especially advantageous in an adjacent zone and so allows the organism to radiate into it. No selection for a future environment is implied.

**Preboreal** The first *\*Flandrian* (*\*Holocene*, or *\*post-glacial*) *\*stage*, a time of rapid forest spread, from about 10 300–9600 BP. ‘Preboreal’ refers to climatic conditions and in vegetational terms is equivalent to Pollen Zone IV (see **POLLEN ZONE**) of the standard British and European post-glacial pollen chronology.

**Precambrian** A name now used only informally to describe the *\*Hadean*, *\*Archaean*, and *\*Proterozoic*, which together comprise the longest period of geologic time that began with the consolidation of the Earth’s crust and ended approximately 4000 million years later with the beginning of the *\*Cambrian* period around 542 Ma ago. The rocks of this period of geologic time are usually altered and few *\*fossils* with hard parts or skeletons have been found within them. Precambrian rocks outcrop extensively in shield areas such as northern Canada and the Baltic Sea.

**precession** The action of a couple whose axis is perpendicular to the rotational axis (a torque) on a rotating body causing its *\*axis of rotation* to

trace out a path about an average position, instead of having a constant alignment, i.e. the axis of rotation itself revolves conically about a central point. The Earth's axis of rotation precesses as a result of several forces, e.g. changes in mass distribution on its surface, changes in the gravitational field due to changes in the relative positions of the Moon, Sun, and planets, etc.

**precession of the equinoxes** See [MILANKOVITCH CYCLES](#).

**precipitable water** The quantity of rainfall that would result from condensation and precipitation of the total moisture in a column of air in the atmosphere. Most atmospheric moisture is contained in the lower atmosphere, below about 5500 m. On average, an atmospheric column of 1 m<sup>2</sup> cross-section contains vapour equivalent to 5–25 mm depth of rainfall. The average *\*residence* time of moisture in the atmosphere is about nine days.

**precipitation** **1.** In meteorology, all the forms in which water (H<sub>2</sub>O) falls to the ground as rain, sleet, snow, hail, drizzle, or other more specialized forms, and also the amounts measured. Sometimes precipitation seen falling from clouds evaporates before reaching the ground. The term also includes dew, frost, clear ice, fog, and other forms in which atmospheric moisture is precipitated as water but without falling from clouds. **2.** The process of depositing dust or other substances (pollution) from the air. **3.** The deposition of solid particles out of a supersaturated solution.

**precipitation-efficiency index** Devised in 1931 by C. W. *\*Thornthwaite*, an index based on the ratio of mean monthly rainfall and temperature values to evaporation rates. Summation of monthly values gives an annual precipitation-efficiency index (P-E), which is used to define major climatic regions. See also [THORNTHWAITE CLIMATE CLASSIFICATION](#).

**precipitation fog** See [FRONTAL FOG](#).

**precision** See [ERRORS](#).

**predator** An organism that obtains its food by consuming and usually killing another organism (the prey).

**pre displacement** An alteration in the *\*ontogeny* of a descendant such that some developmental process begins earlier than in its ancestor, and so has

progressed further by the time maturity is reached.

**pre-ferns** A group of plants transitional between the **\*Psilophytales** and the true ferns, having some fern and some psilophyte characters. They had leaves and reproduced by **\*spores**, but had a variety of growth forms. The orders usually included in the group are the Protopteridales and Coenopteridales.

**preferred orientation** The alignment of inequidimensional **\*mineral** grains in a rock to define a two-dimensional, planar **\*fabric** or a three-dimensional, linear fabric. Alignment may be caused by deformation accompanying **\*metamorphism**, flow of crystal-laden **\*magma**, settling of **\*crystals** in a magma, or settling of flaky minerals in water.

**pre-Hadean** The earliest **\*era** of the **\*Priscoan**, covering the formation of the Earth and ending around 4550 Ma ago (Harland et al., 1989). The name is no longer used.

**prehnite** A hydrated **\*silicate** and member of the **\*phyllosilicates** (sheet silicates) with composition  $\text{Ca}_2\text{Al}[\text{Si}_3\text{AlO}_{10}]\text{OH}_2$ ; sp. gr. 2.9; **\*hardness** 6.0; colourless, white, or green; **\*tabular** or in granular **\*aggregates**; has a layered structure similar to the **\*micas**; occurs in association with the **\*zeolites** infilling cavities in **\*basic \*igneous** rocks and in contact metamorphosed (see THERMAL METAMORPHISM) **\*limestones**. See also PREHNITE-PUMPELLYITE FACIES.

**prehnite-pumpellyite facies** A set of metamorphic **\*mineral** assemblages produced by **\*metamorphism** of a wide range of starting rock types under the same metamorphic conditions and typically characterized by the development of the mineral assemblage prehnite ( $\text{Ca}_2\text{Al}[\text{Si}_3\text{AlO}_{10}](\text{OH})_2$ ), pumpellyite ( $\text{Ca}_2\text{Al}_2(\text{Mg}, \text{Fe}^{2+}, \text{Fe}^{3+}, \text{Al})[\text{SiO}_4][\text{Si}_2\text{O}_7](\text{OH})_2(\text{H}_2\text{O}.\text{OH})$ ) and **\*quartz**, with relict **\*plagioclase** and **\*pyroxene**, in rocks of **\*basic \*igneous** composition. Other rocks of contrasting composition (e.g. **\*shales** or **\*limestones**) would each develop their own specific mineral assemblage, even though they are all being metamorphosed under the same conditions. The variation of mineral assemblage with starting rock composition reflects a particular range of pressure, temperature, and  $P(\text{H}_2\text{O})$  conditions. Experimental studies of pressure–temperature stability fields indicates that the **\*facies** represents a range of moderate pressure (2.5–5.0 kb), low

temperature (150–300 °C) conditions usually developed during *\*burial* metamorphism in thick sedimentary sequences on *\*continental margins* and in intracontinental basins.

**Pre-Imbrian** *See* LUNAR TIMESCALE.

**Pre-Nectarian system** *See* LUNAR TIMESCALE.

**pre-splitting** A method of blasting in which a planar crack is propagated by blasting to determine the final shape of a rock face before holes are drilled for the final blast pattern; firing with a minimum time scatter can then be used. The crack helps to screen the surroundings from ground vibrations during the firing of the main round.

**pressure-depth profile** In the Earth, the relationship between pressure and depth is controlled by the density of the rocks and the *\*gravitational* acceleration at each point along the profile. Gravitational acceleration is almost constant through most of the *\*mantle*, though it rises slightly close to the *\*core*–mantle boundary, and then decreases linearly to zero at the centre of the Earth. The pressure rises to about 100 GPa at the core–mantle boundary and reaches 375 GPa at the centre of the Earth.

**pressure dissolution** *See* PRESSURE SOLUTION.

**pressure fringe** A growth of fibrous *\*quartz*, *\*calcite*, *\*chlorite*, or *\*muscovite*, normal (or sometimes parallel) to the face of a *\*porphyroblast* (usually *\*pyrite* or *\*magnetite*) in a *\*regional metamorphic* rock. During *\*metamorphism* and deformation *\*minerals* dissolved from the region of high pressure, where the *\*matrix* is squeezed against the *\*crystal*, are redeposited in the *\*pressure shadow* zone at the side of the crystal. The minerals nucleate on the face of the porphyroblast and regrow in a fixed crystallographic orientation with respect to the edge of the large porphyroblast. This results in the straight fibres which often intersect at segment junctions. Where the porphyroblast is rotated during the metamorphic event the pressure-fringe minerals grow as curved fibres, thus tracing the stages of detachment between porphyroblast and matrix.

**pressure-gradient force (PGF) 1.** The force acting on air due to pressure differences. Horizontal variations in pressure create a tendency for movement from higher to lower pressure, but as air moves it experiences the *\*Coriolis* effect (CorF), deflecting it. When the PGF and CorF are in

balance above the layer affected by friction with the surface, the air flows around the centre of low pressure, parallel to the **\*isobars**, forming the **\*geostrophic wind**. Near the surface, friction slows the moving air, reducing the magnitude of the CorF, and air flows at an angle across the isobars towards the centre of low pressure. **2.** The force acting on a water mass due to pressure differences over distance. Horizontal variations in pressure create a tendency for movement from higher to lower pressure areas. *See also* **GEOSTROPHIC CURRENT**.

**pressure head** The potential energy possessed by a unit weight of water at any point when compared with a pressure of one **\*atmosphere** at the same elevation. For **\*groundwater**, it is measured by the depth of submergence between the measurement point and the water level in a **\*piezometer**. *See also* **HYDRAULIC HEAD; ELEVATION POTENTIAL ENERGY; POTENTIOMETRIC SURFACE**.

**pressure melting** The melting of **\*ice** in response to **\*stress**, due to the fact that the freezing temperature of water falls as pressure increases, at about 1°C for every 140 bars ( $140 \times 10^5 \text{ N/m}^2$ ). The term ‘pressure melting point’ refers to the temperature at which ice just begins to melt under a given pressure.

**pressure ridge** A curved, elongated ridge on the surface of a basaltic **\*lava** flow, formed at right angles to the flow direction. It may have been pushed up by the dragging effect of mobile lava below a cooling surface, although collapse may contribute to its detached form.

**pressure shadow** The area in a **\*regional metamorphic** rock which is protected from deformation by the presence of a relatively rigid **\*porphyroblast** or **\*porphyroclast**. Randomly orientated **\*quartz** and/or **\*chlorite** concentrates in triangular regions next to the faces of the porphyroblast or porphyroclast which are themselves orientated perpendicular to the **\*schistosity** of the surrounding, deformed, metamorphic **\*fabric**. During **\*metamorphism** and deformation quartz and chlorite are dissolved from the region of high pressure, where the matrix is squeezed against the hard, unyielding porphyroblast or porphyroclast, and are redeposited in the no-stress shadow region on either side of the porphyroblast or porphyroclast, where the deformed fabric wraps around the crystal, so forming the pressure shadow.

**pressure solution (pressure dissolution)** The process that occurs preferentially at the contact surface of *\*grains* or *\*crystals* as a result of an excess of external pressure relative to the hydraulic pressure of pore fluids. Material in these zones is dissolved and removed, resulting in an increase in *\*compaction* and a decrease in *\*porosity*.

**pressure-tube anemometer** See ANEMOMETER.

**pressure wave** See P-WAVE.

**pressure welding** The suturing together of *\*grains* in a *\*sedimentary rock* as the result of pressure dissolution (see PRESSURE SOLUTION) taking place along the grain contact. The sutured margin appears as an irregular plane. Pressure welding of grains is particularly common when a *\*mineral \*cement* has developed at a late stage, after deep burial of the *\*sediment*, as this results in the main *\*overburden* being supported by the grain-to-grain contacts of the sediment.

**pre-tectonic** Applied to a process or event which occurs before deformation. Compare POST-TECTONIC; SYNTECTONIC.

**pre-Tolstojan** The earliest period of the *\*Mercurial geologic timescale*, lasting from 4568–3900 Ma ago. It marks the formation of the planet *\*Mercury* and the *\*late heavy bombardment*.

**prevailing wind** In a particular locality, the wind direction that is most frequent over time. For most areas the prevailing wind varies, sometimes quite markedly, according to season. It may also change, or have changed, when the climate changes (as in ice ages).

**prey** See PREDATOR.

**prezygapophyses** See VERTEBRA.

**Priabonian (Runangan)** The final *\*stage* in the *\*Eocene \*epoch*, preceded by the *\*Bartonian*, followed by the *\*Rupelian* (Stampian), and dated at 37.2–33.9 Ma ago (Int. Commission on Stratigraphy, 2004). It is roughly contemporaneous with the *\*Refugian* (California) and the Runangan (New Zealand) and part of the *\*Aldingan* (Australia).

**Priapulida** Phylum comprising the priapus worms, known from *\*Cambrian* times (e.g. *Ottoia* from the *\*Burgess Shale*, Canada), and ranging up to the

present.

**Pridoli** A sub-*epoch* (418.7–416 Ma ago) of the Late *Silurian*, underlain by the *Ludlow* and followed by the Early *Devonian* epoch.

**primary creep (transient creep)** The initial stage of *creep*, characterized by viscoelastic *strain*, in a material subjected to long-term, low-level *stress*.

**primary crushing** See CRUSHING.

**primary geochemical differentiation** A theory explaining the formation of the Earth's *core*, *mantle*, and *crust*, in which the formation of the nickel–iron core may have been accompanied by partitioning of the elements. It postulates that some of the elements were reduced, alloyed with iron, and concentrated in the core, while the remainder formed the mantle and primitive crust.

**primary geochemical dispersion** The movement of elements below the Earth's surface by metamorphic, magmatic, or hydrothermal processes (see METAMORPHISM; MAGMATIC DIFFERENTIATION; HYDROTHERMAL ACTIVITY), resulting in the formation of *igneous* and *metamorphic rocks*.

**primary magma** See PRIMARY MELT.

**primary melt (primary magma)** A liquid that forms in the *mantle* as a result of *partial melting*.

**primary migration** First stage in the upward migration of *hydrocarbons* within and then out of the *source rock*.

**primary mineral** A *mineral* which has crystallized from a *magma*. *Essential* primary minerals are those primary minerals whose presence is essential for the classification and naming of the rock, while *accessory* primary minerals are those primary minerals whose presence does not affect the classification or naming of the rock.

**primary porosity** See POROSITY; CHOQUETTE AND PRAY CLASSIFICATION.

**primary productivity** The amount of organic matter synthesized by organisms from inorganic substances. In the oceans, *photosynthesis* by phytoplanktonic *algae* in the upper 100 m (the euphotic zone) accounts for most primary production. Waters of tropical areas are less productive than

those of temperate regions, because in tropical areas the water column undergoes no seasonal vertical mixing. Areas of **\*upwelling** of nutrient-rich deep waters have high productivity.

**primary sedimentary structure** A structure that forms during or very soon after the deposition of the sediment of which it is composed. *Compare* SECONDARY SEDIMENTARY STRUCTURE.

**primary wave** *See* P-WAVE.

**primitive (evol.)** Preserving the character states of an ancestral stage. The term may be used of a character (as a synonym of **\*plesiomorphic**) or, occasionally, of a whole organism.

**primitive circle** In a **\*stereographic projection**, the circumference of the **\*plane of projection**.

**primordial** The period of time at, or just before, the final formation of the **\*Earth**.

**principal component analysis** A **\*multivariate** analysis which maximizes the spread of data by plotting **\*covariance** values on sets of axes in multidimensional space allowing correlations which may have been hidden in the data to be identified. The first principal component corresponds to the first axis in multidimensional space and describes the majority of the spread of the data, subsequent higher order principal component axes are orthogonal to the first axis. Higher order axes display progressively less variation, where the data is less correlated and more representative of statistical noise.

**principal point** The centre point of an aerial photograph.

**principal shock** The main **\*earthquake** event with the largest amplitude.

**principal strain axes** In a strained material, three mutually perpendicular axes (designated X, Y, and Z) which are parallel to the directions of greatest, intermediate, and least elongation, and which describe the state of **\*strain** at any particular point.

**principal strain ratio (ellipticity)** The ratio ( $R$ ) of the minimum **\*strain** axis to the maximum strain axis. In its simplest, two-dimensional form the long and short axes of initially circular markers are measured and plotted

against one another; the slope of the best-fit line through the origin and a number of points is the strain ratio for that plane of measurement. More complex methods are employed to determine *R* minimum, *R* maximum, and *R* tectonic for initially non-spherical strain markers, e.g. **\*conglomerate** **\*clasts**.

**principle stress axes** Three mutually perpendicular axes (designated,  $\sigma_1$ ,  $\sigma_2$ , and  $\sigma_3$ ) which are parallel to the directions of maximum, intermediate, and least principal **\*stress**. Their separate lengths and directions describe the state of stress at a particular point. The stress ellipse contains  $\sigma_1$  and  $\sigma_3$ ; the stress ellipsoid contains all three axes.

**principle of contained fragments** See PRINCIPLE OF INCLUDED FRAGMENTS.

**principle of included fragments (principle of contained fragments)** The principle that any rock containing fragments of another rock body must be younger than the rock unit from which those fragments were derived.

**principle of superposition** See LAW OF SUPERPOSITION.

**principle of uniformitarianism** See UNIFORMITARIANISM.

**Priscoan** See HADEAN.

**PRISM** See PICOSATELLITE FOR REMOTE-SENSING AND INNOVATIVE SPACE MISSIONS.

**prism** A **\*crystal** form composed of a number of repeat **\*crystal** faces all of which are parallel to one of the principal **\*crystallographic axes** (usually the vertical *c* (or *z*) axis). See also ACCRETIONARY PRISM.

**PRISMA** See HYPERSPECTRAL PRECURSOR AND APPLICATION MISSION.

**prismatic** Applied to elongate **\*minerals** which have well-developed **\*crystal faces** parallel to the **\*prism**.

**protactinium dating** See URANIUM-PROTACTINIUM DATING; PROTACTINIUM/THORIUM ISOTOPES.

**protactinium/thorium isotopes** The *\*isotopes*  $^{231}\text{Pa}$  and  $^{230}\text{Th}$  are products of the decay of the uranium isotopes  $^{235}\text{U}$  and  $^{234}\text{U}$ , respectively, dissolved in seawater and well mixed owing to uranium's long residence time, yielding a Pa:Th ratio that is constant and distributed uniformly.  $^{231}\text{Pa}$  and  $^{230}\text{Th}$  are adsorbed onto settling mineral particles, Th more efficiently than Pa, so Pa has a longer seawater residence time (~200 years) than Th (~30 years), which allows Pa to move laterally more than Th prior to deposition. The  $^{231}\text{Pa}$  deficit in the Pa:Th ratio in the mineral content of ocean sediment can be used to calculate the extent to which the water was moving laterally (and removing Pa) during the period of deposition and is, therefore, an indicator of past ocean circulation.

**probability density function** In statistics, the mathematical function which allocates probabilities of particular observations occurring. The probability density function may be used to construct a frequency distribution of certain events occurring either discretely, in the form of a *\*histogram*, or continuously.

**probability distribution** In statistics, the relative frequency distribution of different events occurring, as defined by the *\*probability density function*. Probability distributions may be discrete as in the cases of the *\*binomial* and *\*Poisson distributions*, or continuous as in the case of the *\*normal distribution*.

**PROBA-V** See PROJECT FOR ON-BOARD AUTONOMY – VEGETATION.

**'Problematica'** A named group into which geological structures are placed if their affinity is uncertain but their origin is considered to be organic. They are removed from the 'Problematica' when more certain evidence of their relationships is discovered.

**Proboscidea (proboscideans)** (infraclass *\*Eutheria*, cohort *\*Ferungulata*) Order comprising the elephants and their extinct relatives, e.g. mastodons (see MAMMUTIDAE), gomphotheres (see GOMPHOTHERIIDAE), and mammoths (see MAMMUTHUS). The order was formerly highly successful and occupied the Americas, Eurasia, and Africa. Proboscideans tend toward large size. Since the late *\*Miocene* most have possessed a long trunk: this is developed from the nose and upper lip. Teeth are reduced in number, young adults having three molars in each side of each jaw; these are used one at a

time, old teeth being shed and replaced by those behind. The upper incisors are enlarged to form tusks. The jaw muscles are large, and the skull short and high. The *\*vertebrae* and up to 20 ribs carry the weight of the abdomen, which is balanced on the fore limbs by the weight of the head, the hind limbs providing propulsion. The brain is well developed. Parental care of the young is prolonged, and social organization is complex.

**proboscidean** See PROBOSCIDEA.

**Procaryotae** A taxonomic kingdom, proposed in 1968, to include *\*Bacteria* (prokaryotes), divided into three divisions: those with a Gram-negative cell wall; those with a Gram-positive cell wall; and those with no cell wall. The name is no longer used.

**Procellarum system** See LUNAR TIME-SCALE.

**processes, geomorphological** The set of mechanisms that operate at and near the Earth's land surface, breaking down and transferring rock material and consequently fashioning land-forms at small and medium scales. Processes operating on the surface are termed 'exogenetic'; those originating below the surface 'endogenetic'. Two categories may be recognized: (a) zonal processes, which are broadly controlled by climate (e.g. the work of *\*glaciers*); and (b) azonal processes, which operate on a world-wide basis (e.g. *\*fluvial processes*). The rate at which such processes operate may be measured by direct field recording or by making use of historical records. See EROSION RATE.

**process-response system** In *\*geomorphology*, a natural *\*system* that is formed by the combination of at least one *\*morphological* and one *\*cascading system*. It therefore shows how form and geomorphological process are related. An example is the coastal process-response system, in which the cascading system of wave energy advances from deep water to the edge of the *\*swash* zone, and is linked to various morphological features of the shallowing zone.


**prochoanitic** See SEPTUM.

**Procolophonia** Relatively advanced cotylosaurs (*\*Cotylosauria*, 'stem reptiles'), distinguished from the lower types by their shorter jaws and

better jaw movement. These anapsids (*\*Anapsida*) lived in the *\*Triassic* and *\*Permian*.

**Procyonidae** See *CARNIVORA*.

**pro-delta** The furthest offshore portion of a *\*delta*, lying at the toe of the *\*delta front*, and characterized by a relatively slow rate of fine-grained deposition.



**Pro-delta**

**prod mark** A type of *\*tool* mark formed by the impact of an object with a soft, muddy surface. Prod marks are characterized by an asymmetrical impact mark, with the steeper margin of the depression being on the down-current side. See also *BOUNCE MARK*; *SKIP MARK*; *TOOL MARK*.

**production log** A *\*well* log which is run inside a casing, primarily to measure the behaviour of fluids within the pipe.

**production well** A well from which water, gas, or oil is actually to be recovered, as opposed to other wells, e.g. those designed to determine hydraulic characteristics, to recharge an *\*aquifer*, or to act as injection wells to push oil towards the production well.

***Productus giganteus* (*Gigantoproductus giganteus*)** A huge brachiopod (*\*Brachiopoda*) which attained a width of approximately 370 mm. It lived in the warm seas of the Lower *\*Carboniferous* (*\*Mississippian*), adopting a clam-like habit.

**Proetida** An order of *\*Trilobita* that lived from the *\*Ordovician* to *\*Permian*. The glabella (see *CEPHALON*) was large and clearly defined, often

with *\*genal spines*, the thorax had 8–10 segments, and the *\*pygidium* had furrows and was not spiny. The order contained two superfamilies.

**profiling** **1.** A method in which an *\*array* is moved progressively along a traverse line to obtain a continuous cover of measurements of the subsurface and so produce a profile. see [CONSTANT-SEPARATION TRAVERSING](#). **2.** In reflection seismology, the acquisition of data along a line, which then form a seismic section (see [SEISMIC RECORD](#)). See [COMMON DEPTH POINT](#).

**pro-foreland basin** See [PERIPHERAL FORELAND BASIN](#).

*Proganochelys quenstedii* (*Triassochelys quenstedii*) The oldest known turtle, first described in 1887 from the *\*Triassic* Stubensandstein of Württemberg, Germany. It possessed teeth, and stout ribs on the neck vertebrae.

**progenesis** The onset of sexual maturity at a younger and smaller stage in development than is usual.

**proglacial** Applied to the area between a *\*glacier* and adjacent high ground. A proglacial lake is a body of water impounded in such an area and is often inferred for areas of *\*Pleistocene* glaciation from the evidence of strandlines, lake sediments, and *\*overflow* channels.

**progradation** The outward building of a sedimentary deposit, such as the seaward advance of a *\*delta* or shoreline, or the outbuilding of an *\*alluvial fan*.

**prograde metamorphism (progressive metamorphism)** The *\*recrystallization* of a rock in response to an increase in the intensity of *\*metamorphism* as this is reflected by an increase of pressure, temperature, and/or  $P(\text{H}_2\text{O})$ . Compare [RETROGRADE METAMORPHISM](#).

**progressive deformation** The accumulation over time of incremental *\*strain* elements within a body as a response to *\*stress*. The accumulated distortions and rotations within the body add up to a final (*\*finite*) strain state.

**progressive evolution** Steady, long-term improvement of evolutionary grade, which has allowed plants and animals to become ever more

independent of the aquatic environment in which they first evolved. For example, the sequence *\*bryophyte* to *\*pteridophyte* to *\*gymnosperm* to *\*angiosperm* represents a progressive evolutionary trend.

**progressive metamorphism** See [PROGRADE METAMORPHISM](#).

**progressive wave** A wave that is typified by the progressive forward motion of the wave-form. The speed of propagation of the wave-form depends largely on the depth of the water. See also [CELERITY](#).

**Progymnospermopsida (progymnosperms)** The ancestors of the *\*gymnosperms*, which arose in the *\*Devonian* and dwindled to extinction in the latter part of the *\*Carboniferous*. They had trunks with wood resembling that of gymnosperms, but their fertile branches or leaves bore sporangia (see [SPORE](#)), and their foliage was often fern like. Probably seeds evolved in various different progymnosperms. See [ARCHAEOPTERIS](#).

**progymnosperms** See [PROGYMNOSPERMOPSIDA](#).

**Project for On-Board Autonomy – Vegetation (PROBA-V)** A mission by the *\*European Space Agency* and the French Space Agency to accommodate on a small satellite an improved version of the large optical vegetation-observing instrument on *\*SPOT-4* and *-5*. The satellite was launched on 7 May 2013, from Kourou, French Guiana, into a Sun-synchronous orbit at an altitude of 704 km.

**prokaryote** An organism, usually unicellular, in which the cells lack a true nucleus, the DNA being present as a loop in the cytoplasm. Other prokaryotic features include the lack of chloroplasts and mitochondria and the possession of small ribosomes. Compare [EUKARYOTE](#).

**prokaryotic** See [PROKARYOTE](#).

**prolate** Applied to *\*clasts* which are rod-shaped, and defined as having a ratio of short to intermediate diameters of more than 2:3, and a ratio of intermediate to long diameters of less than 2:3.

**prolate uniaxial strain** The *\*strain* state which occurs when a reference sphere is extended along its *X* axis and shortened in all other directions perpendicular to *X*. The resultant strain shape is a prolate ellipsoid.

**proloculus** The initial chamber of a foraminiferid (*\*Foraminiferida*) *\*test*.

**Prometheus (Saturn XVI)** One of the lesser satellites of *\*Saturn*, discovered in 1980 by *\*Voyager 1*, with a radius measuring  $74 \times 50 \times 34$  km; mass  $0.0014 \times 10^{20}$  kg; mean density  $270 \text{ kg/m}^3$ ; visual albedo 0.6.

**promontorium** On *\*Mars*, a cape, e.g. Prom. Deville, Prom. Kelvin.

**proparian** Applied to a trilobite (*\*Trilobita*) *\*cephalic suture* that runs around the front of the glabella (see *CEPHALON*), terminating in front of the *\*genal* angle.

**proper motion** The apparent movement of a star relative to the other 'fixed' stars, at right angles to the observer's line of sight. For example, Barnard's star moves 10 seconds of arc per year; this is the largest proper motion recorded. There is no absolute frame of reference, and the positions of stars in the familiar constellations change significantly within a few thousand years.

**propylitization** The process whereby original *\*plagioclase* in an *\*igneous* rock is altered to an *\*epidote*–*\*sericite*–secondary *\*albite* assemblage and original *\*ferromagnesian* minerals are altered to a *\*chlorite*–*\*calcite*–epidote–secondary iron-ore assemblage.

**proseptum** (*pl. prosepta*) See *SEPTUM*.

**prosobranch gastropods** Single-gilled gastropods (*\*Gastropoda*), thought to represent the first stage in the evolutionary development of the group from the ancestral forms, the *\*Archaeogastropoda*.

**prosogyral** See *PROSOGYRATE*.

**prosogyrate (prosogyral)** Applied to the umbones (see *UMBO*) of the *\*Bivalvia*, where the beaks are curved so as to point in an anterior direction.

**prosoma** See *ARACHNIDA*; *CHELICERATA*.

**Prospero (Uranus XVIII)** A lesser satellite of *\*Uranus* with a radius of 25 km and a visual *\*albedo* of 0.07.

**protalus rampart** Ridge of rock debris less than 10 m high and found near the base of a steep inland face. It consists of frost-shattered debris that has been carried some distance from the face down the steep surface of a basal snow bank.

**protaspis** The earliest stage recognized in larval trilobite (*\*Trilobita*) development. The larva is small, often spiny, and grows through successive moult stages. Initially it is a small disc but size and segmentation increase with each successive moult.

**Proterosuchus** Well-known representative of the *\*Thecodontia*, the ancestral archosaurian (*\*Archosauria*) stock. According to some authorities, *Proterosuchus* was the first crocodile. It lived in N. America during the *\*Triassic*.

**Proterozoic** The most recent (about 2500–542 Ma ago) of the three subdivisions of the former *\*Precambrian*. It is now classed as an eon following the *\*Archaean* and preceding the *\*Phanerozoic* eons.



<http://www.geologypage.com/2013/10/proterozoic-eon.html>

- Proterozoic Eon.

**Proteus (Neptune VIII)** A satellite of *\*Neptune*, measuring  $436 \times 416 \times 402$  km; visual albedo 0.06.

**protist** A single-celled, eukaryotic organism (see *EUKARYOTE*) that may resemble an animal or a plant. Animal-like protists include naked and shelled amoebas, foraminiferans, zooflagellates, and ciliates; plant-like protists include dinoflagellates, diatoms, and algae. In a five-kingdom system of classification protists were grouped as a kingdom, Protista; later some multicellular organisms with protist affinities but previously classed as fungi or plants were transferred into the Protista and the name of the kingdom was changed to *\*Protoctista*. In the modern three-*\*domain* classification, the Protista comprise a kingdom within the domain Eukarya.



<https://www.livescience.com/54242-protists.html>

- What Are Protists?

**Protista** See *PROTIST*.

**proto-** From the Greek *protos* meaning ‘first’, a prefix meaning ‘original’ or ‘primitive’.

**proto-Atlantic** See IAPETUS OCEAN.

**Protoaulopora ramosa** Small, colonial coral belonging to the subclass \*Tabulata, found in the Upper \*Cambrian rocks in Kazakhstan. With *Bijasibirica* (a doubtful stony coral) it marks the appearance of an important group of \*reef-forming organisms.

**Protoceratops andrewsi** The first of the horned (\*ceratopsian) \*dinosaurs, known from the Middle \*Cretaceous of Mongolia. It possessed the bony frill and beak characteristic of later forms, but was only 2m in length and 1.5 t in weight.

**protoconch** The initial (larval) shell of molluscs (\*Mollusca), often retained at the tip of the \*spire of the adult shell.

**Protoctista** In the five-kingdom classification of biological organisms, a kingdom within the superkingdom Eukarya. Protoctists are aquatic \*eukaryotes, but they are not plants, animals, nor fungi. The kingdom includes naked and shelled amoebae, the \*Foraminiferida, zooflagellates, ciliates, \*Dinophyceae, \*Bacillariophyceae, all algae (see ALGA), \*protozoa, slime moulds, and slime nets. The name is not used in the three-\*domain classification, the protoctists being placed in other kingdoms in the domain Eukarya.

**protolith** The parent rock from which a \*metamorphic rock is derived. See METAMORPHIC ROCK CLASSIFICATION.

**proton-adding process (p-process)** Nuclear process in red giant stars at very high temperatures, producing proton-rich, heavier elements. See also NUCLEOSYNTHESIS.

**proton magnetometer** A form of \*nuclear-precession magnetometer that is based on the \*precession of protons in water or alcohol.

**protoparian suture** See CEPHALIC SUTURE.

**protoplanet** An individual condensation, representing a very small proportion of the total cloud mass but of similar composition, that occurs during the condensation of an interstellar cloud, from which stars and planets are assumed to have originated. See PROTOSTAR.

**Protopteridales** See PRE-FERNS.

**protore** Rock containing sub-economic material from which economic \*mineral deposits may form by geologic concentration processes such as \*supergene enrichment. A protore may become profitable with technological advance or change in market value.

**protostar** Primitive star formed from the break-up of \*interstellar clouds. After a fragment becomes detached, it continues to shrink under the influence of its own gravitation, drawing in more gas and dust and increasing in temperature and pressure. Eventually, the outward pressure associated with the rising temperature balances the inward pressure due to gravitation and collapse ceases, perhaps 10 000 years after separation from the cloud. At this stage, the fragment is called a protostar. When internal temperature exceeds  $10^7$  K \*hydrogen 'burning' begins, marking the transition to a star.

**protozoa** (*sing.* protozoon; *adj.* protozoan) Formerly an animal phylum (Protozoa) of \*eukaryotic, single-celled micro-organisms. Protozoa are now included in the kingdom \*Protoctista in the five-kingdom classification and in the kingdom Protista (\*domain Eukarya) in the three-domain classification.

**protrusive** Applied to the downward extension into \*sediment of a vertical or oblique \*burrow, frequently reflected in the development of \*spreiten distal to the point of entry. In U-shaped burrows the spreiten occur inside the bend of the 'tube'. Protrusive spreiten are found in \*Diplocraterion and \*Rhizocorallium.

**proved reserve** See RESERVE.

**provenance** The source or origin of \*detrital \*sediments.

**province** 1. Region or area of large extent with similar features throughout and capable of being considered as a unit. 2. In geography, an area of land or sea in one climatic belt.

**provinciality** The association of species within well-defined biogeographic areas or provinces. Each province contains a distinct assemblage of species, some of which are endemic (i.e. confined to that area only). See ENDEMISM.

**proximal** Applied to a \*sediment or sedimentary environment close to the source or origin of the deposit. Compare DISTAL.

**proximity log** The record produced by one version of the *\*microlog* sonde.

**proxy** See IONIC SUBSTITUTION.

**Psamathe (Neptune X)** A satellite of *\*Neptune* with a radius of 20 km, visual *\*albedo* of 0.16, and a *\*retrograde* orbit with a period of 9115.9 days.

**psammite** A metamorphosed *\*sandstone*, *\*arkose*, or *\*quartzite*, extremely rich in the *\*mineral* *\*quartz*.

**PSC** See POLAR STRATOSPHERIC CLOUD.

**psephite** A general term used to describe coarse-grained, *\*detrital*, *\*sedimentary rock*, e.g. *\*gravels*, *\*conglomerates*, or *\*breccias*. G. W. Tyrell (1921) suggested the term be restricted to metamorphosed conglomerates and breccias, a suggestion followed by many authors describing *\*metamorphic rocks*.

**pseudobreccia** An irregularly *\*recrystallized* or partially *\*dolomitized* *\*limestone*, in which the selective growth of coarse crystals gives the rock an apparently fragmented *\*texture*.

**pseudoextinction** Within an *\*evolutionary lineage*, the disappearance of one taxon caused by the appearance of the next *\*chronospecies* in the series. The extinction is purely taxonomic.

**pseudofossil** A naturally occurring object that may resemble a *\*fossil*. If there is uncertainty the object may be referred to the *\*Problematica*; it is called a pseudofossil when the resemblance results purely from chance.

**pseudo-gravitational field** A gravitational field that is transformed to simulate a magnetic field for the purpose of computational analysis.

**pseudo-magnetic field** A magnetic field that is transformed to simulate *\*gravitational* acceleration for the purpose of computational analysis.

**pseudomorph** A *\*secondary mineral* or a random aggregate of secondary minerals (monomineralic or polymineralic) which have replaced an earlier mineral but have retained its shape. For example, a random aggregate of *\*chlorite* crystals which have replaced a *\*euhedral* *\*augite* crystal while preserving its shape is said to form a monomineralic pseudomorph of augite. Other examples include *\*quartz* after *\*fluorite*, *\*serpentine* after

\***olivine**, and \***limonite** after \***pyrite**, where the original \***cubic** form is preserved. *See also* METASOMATISM.

**pseudonodule** A ball-like body of \***sandstone**, with an internal lamination that is convoluted or upcurled at the edges, set in a bed of \***mudstone**. The sand ball is the result of the sinking of sand from the base of an overlying bed into the less dense, soft mud below.

**pseudopunctate** *See* PUNCTATE.

**pseudosection (quasi-section)** The plotting of data against position along a traverse line, to produce a display of \***resistivity** or \***induced-polarization** data in which values are given to the intersection point of 45° lines drawn from mid-points of the current and potential electrode pairs. Depths in the resulting 'section' below a transverse bear no simple relationship to the true geology; a pseudosection shows the variation of the measured parameter with position and with effective depth of penetration, rather than with true depth. It is used widely in displaying \***induced-polarization** data and \***apparent resistivities** obtained from \***constant-separation** traverses with different electrode separations, and apparent conductivities from electromagnetic traverses with different coil separations.

**pseudospar** *See* NEOMORPHISM.

**Pseudosycidium** The first of the Charales, a distinct evolutionary line of green \***algae** which arose during the Upper \***Silurian** and whose method of reproduction was sexual. Well-developed male (antheridia) and female (oogonia) organs can be identified, the secretion of calcium carbonate around the female organs assisting preservation. *See also* CHAROPHYCEAE; GYROGONITE.

**pseudotachylite** A rare, glassy rock produced by frictional melting during extreme \***dynamic metamorphism** in a \***fault** or \***thrust** zone, or by meteoric impact.

**psilomelane** A hydrated manganese oxide with the possible formula  $Ba_3(Mn^{2+}Mn_7^{4+})O_{16}(OH)_4$ ; sp. gr. 3.7–4.7; \***hardness** 5–6; grey-black; \***sub-metallic** or \***earthy lustre**; \***massive**, \***botryoidal**; occurs as a precipitate in \***secondary** manganese deposits and may be worked commercially for manganese.

**Psilonichnus** An *\*ichnoguild* comprising a *\*biofacies* developed on the *\*backshore* of former beaches.

**Psilophytales (psilophytes)** Primitive *\*pteridophytes*, which were the earliest vascular plants, from the *\*Silurian* and the *\*Devonian*. They had slender, tapering, leafless or scale-bearing stems up to 50 cm high, often with cone-shaped sporangia (see **SPORE**) at the top. Some authorities placed the fossil groups with the present-day psilophytes (*Psilotum*, *Tmesipteris*) in the class Psilopsida, comprising the orders Psilotales (living forms) and Psilophytales (fossil forms). In more recent classifications the members of the Psilophytales have been placed into three separate groups, usually ranked as subdivisions: Rhyniophytina (e.g. *Cooksonia* and *\*Rhynia*); Trimerophytina (e.g. *Psilophyton* and *Trimerophyton*); and Zosterophyllophytina (e.g. *\*Zosterophyllum*). The living forms are contained in a fourth subdivision, the Psilophytina.

**psilophytes** See **PSILOPHYTALES**.

**Psilopsida** See **PSILOPHYTALES**.

**PSV** See **AGGREGATE TESTS**.

**psychrometer** *\*Hygrometer* comprising a *\*wet-bulb* and a *\*dry-bulb* thermometer. See also **WHIRLING PSYCHROMETER**.

**psychrophile** An *\*extremophile* (domain *\*Archaea*) that thrives in environments where the temperature is low, usually below 15 °C.

**Pteraspida** See **HETEROSTRACI**.

**Pteraspis** See **HETEROSTRACI**.

**Pteridophyta (pteridophytes)** Phylum of the plant kingdom, comprising the true ferns. These first enter the *\*fossil* record in the *\*Silurian*. They are flowerless plants exhibiting an alternation of two distinct and dissimilar generations. The first is a non-sexual, *\*spore*-bearing, sporophyte generation. It usually appears as a relatively large plant, with stems containing vascular tissue that conducts water and dissolved solutes through the plant, and usually bears the leaves and roots. Spores are produced in sporangia that are either attached to the leaves (as in ferns) or are on specialized scales (sporophylls) grouped into cones (as in horsetails and

club mosses), or in the axils of leaves on unspecialized stems (as in some club mosses). The second is a sexual, gametophyte generation, in which the plants generally are relatively small, and without differentiation of stem, leaves, or roots. These plants bear male (antheridia) and female (archegonia) sex organs, together or on separate plants. When the eggs in the archegonia are fertilized by sperms from the antheridia, an embryo results; this can grow into a new sporophyte generation.

**pteridophytes** See PTERIDOPHYTINA.

**Pteridospermales (seed ferns)** Extinct \*gymnosperm order, which were the earliest seed plants and flourished in the \*Carboniferous, before disappearing in the \*Cretaceous. Their foliage was fern-like in appearance, but the fertile leaves bore seeds and \*pollen-producing organs. The first plant to be identified (in 1903) as a seed plant rather than a true fern was *Lyginopteridales oldhamia*.

**Pterobranchia** (phylum \*Hemichordata) Class of minute, fixed, colonial, deep-sea organisms which secrete an external cuticular skeleton in which they are housed (e.g. *Rhabdopleura*). Pterobranchs may be the nearest living relatives of the graptolites (\*Graptolithina). The mesosoma is small but carries one or more pairs of arms bearing tentacles (the lophophores). The metasoma consists of a long stalk (peduncle) by which the individual is attached. The branchial apparatus is rudimentary.

**pteropod ooze** Deep-sea ooze in which at least 30% of the sediment consists of the shells of \*planktonic small \*gastropods (known as pteropods or 'wing-footed' snails). The shells are aragonitic and, as \*aragonite solubility increases rapidly with depth, pteropod ooze is restricted to water depths less than 2500 m.

**Pteropsida** A former name of a subdivision of plant that included the ferns, \*gymnosperms, and \*angiosperms.

**Pterosauria (pterosaurs)** \*Mesozoic order of flying \*reptiles, which were particularly numerous in the \*Jurassic, but survived until late \*Cretaceous times. Their fossil skeletons suggest that they could not stand upright on land, and so it is assumed that their mode of life involved swooping over the sea to catch fish. Their remains are always associated with marine deposits. See QUETZALCOATLUS NORTHROPI.

**Ptilograptidae** See DENDROIDEA.

**Ptychopariida** An order of *\*Trilobita* that lived from the Lower *\*Cambrian* to Upper *\*Devonian*. It was a large, *\*paraphyletic* group, divided into two suborders.

**ptygmatic fold** An irregular, lobate *\*fold*, usually found where single *\*competent* layers are enclosed in a *\*matrix* of low competence. Typically, ptygmatic folds do not maintain their *\*orthogonal* thickness (i.e. they are *\*similar* folds). Characteristically their *\*axial planes* are curved.

**Puarooan** A New Zealand *\*stage* (150–145.5 Ma ago) of the Late *\*Jurassic* *\*epoch*, preceded by the *\*Ohauan* and followed by the *\*Mokoian*.

**pubis** In *\*tetrapods*, the anterior, ventral part of the *\*pelvic girdle*. In *\*ornithischian* *\*dinosaurs* the pubis lies alongside the *\*ischium*; in advanced ornithischians there is also a forward prong, the prepubis. In *\*saurischian* dinosaurs the pubis points downwards and forward from the hip socket.

**Puck (Uranus XV)** One of the lesser satellites of *\*Uranus*, with a diameter of 77 km. It was discovered in 1985.

**puddingstone** See CONGLOMERATE.

**puddled soil (poached soil)** Soil in which the structure has been destroyed by the physical impact of rain drops, by tillage when wet, or by trampling by animals.

**Puercan** A N. American *\*stage* of the *\*Palaeocene* (65.5–62.5 Ma ago), preceded by the *\*Maastrichtian* and followed by the *\*Torrejonian*.

**pull-apart basin** See STRIKE-SLIP FAULT.

**pulsations, geomagnetic (micropulsations)** Small, almost sinusoidal fluctuations of the *\*geomagnetic field*, usually with durations of seconds to minutes.

**pulse length** In radar terminology, the total length of an electromagnetic wave emission which is equal to the product of the wavelength, frequency, and time duration of emission.

**pumice** Extremely vesicular, frothy, natural *\*glass*, having a high (60–75%) *\*silica* content and low density. In some cases it will float on water. Usually, but not always, it is of *\*pyroclastic* origin. *See also* RETICULITE; VESICLE.

**pump** A device for moving liquids by adding to the pressure existing within them. For example, a centrifugal pump first increases the velocity of the fluid by the use of impellers; this velocity increase is then converted to an increase in pressure by the use of appropriately orientated guide vanes or the use of a volute casing. Other pump types include multistage turbine pumps, jet pumps, positive-displacement pumps, and suction lifts.

**pumpellyite** *See* PREHNITE-PUMPELLYITE FACIES.

**pumping test (aquifer test)** Water may be pumped from one or more wells to determine the particular hydraulic characteristics of an *\*aquifer* or of individual wells. The effect of pumping at known rates is assessed by the use of *\*observation* wells sunk at appropriate places to monitor the height of water in the aquifer or the wells of special interest.

**punctae** *See* PUNCTATE (2).

**punctate** 1. Applied to any structure that is marked by pores or by very small, point-like depressions. 2. Applied to a type of brachiopod (*\*Brachiopoda*) shell structure in which fine pores (punctae) extend from the inner to the outer surface. Three main shell types are recognized: impunctate, where the shell consists of an outer, lamellar layer of shell and an inner, fibrous layer; punctate (or endopunctate), in which the punctae extend through the shell and end beneath the organic periostracum; and pseudopunctate, in which solid rods of *\*calcite* (taleolae) are contained in the fibrous layer.

**punctuated equilibrium** The theory, first proposed in 1972 by Niles Eldredge and Stephen Jay Gould, that *\*evolution* is characterized by geologically long periods of stability during which little speciation occurs, punctuated by short periods of rapid change, species undergoing most of their morphological changes shortly after breaking from their parent species.

**pupaeiform** Literally, resembling the pupa of an insect, and applied to a gastropod (*\*Gastropoda*) shell where the shell is an elevated ovoid in which the later-formed *\*whorls* have decreasing radii of curvature.



### **Pupaeiform**

**Purbeckian** A *stage* in the British Upper *Jurassic* and Lower *Cretaceous*, underlain by the *Portlandian* and overlain by rocks of the *Wealden* Beds. The *type* section is found in southern England and consists of sediments deposited under *intertidal* to brackish freshwater conditions. *See also* TITHONIAN; RYZANIAN.

**pure shear (homogeneous non-rotational strain)** A flattening *strain* in which, during the deformation of a body, the *principal strain axes* (X, Y, and Z) remain parallel to their respective *principal stress axes* ( $\sigma_1$ ,  $\sigma_2$ , and  $\sigma_3$ ). *See also* HOMOGENEOUS STRAIN. *Compare* SIMPLE SHEAR.

**purga** A strong *blizzard*, similar to the *buran*, in the *tundra* of north-eastern Siberia.

**pushbroom system** In *remote sensing*, an imaging device consisting of a linear array of sensors (*charge-coupled devices*) which is swept across the area of observation. Pushbroom systems allow greater resolution of data to be assimilated than do *line scanner* systems.

**push moraine** *See* MORAINE.

**push-pull wave** *See* P-WAVE.

**Pusgillian** A *stage* (449–447.5 Ma ago) of the Late *Ordovician* *epoch* in the Lower *Ashgill*, underlain by the *Onnian* and overlain by the *Cautleyan*.

**puy** **1.** A volcanic hill in the Auvergne region of France. **2.** Any steep-sided tower of volcanic rock, e.g. the Devil's Tower, Wyoming, USA, and

Shiprock, New Mexico, USA. It consists of the resistant central plug or neck of a former *\*volcano*. See also VOLCANIC PLUG.

**P-wave (compressional wave, dilatational wave, irrotational wave, pressure wave, primary wave, push-pull wave)** An elastic *\*body wave* or sound wave. It is the wave most studied in *\*reflection* and *\*refraction* seismology. Particles oscillate about a fixed point, but in the direction of propagation of wave energy. In an *\*isotropic* and homogeneous medium the P-wave velocity ( $V_p$ ) is given by:  $V_p = [\psi/\rho]^{1/2}$ , where  $\psi$  is the *\*axial modulus* and  $\rho$  is the density of the material. P-waves are the fastest of the *\*seismic waves*.

**PWS** See PLASMA WAVE SUB-SYSTEM.

**pycnocline** Zone in the oceans where water density increases rapidly with depth in response to changes in temperature and salinity. The pycnocline tends to coincide with the *\*thermocline* and *\*halocline*, and separates the surface-water zone of ocean waters from the deep bottom water.

**pycnometer (pyknometer)** An apparatus used to determine the specific gravity or density of soils, or of *\*rock* or *\*mineral* fragments (see DENSITY DETERMINATION). It consists of a small bottle fitted with a ground glass stopper, with a capillary opening to allow air to escape. The bottle is filled with a reference liquid of known specific gravity, e.g. water or mercury, and weighed, then emptied and filled with the specimen liquid and weighed again. The specific gravity or density is calculated from the difference in weights. It is also possible to determine the specific gravity or density of a powder or gas using a pycnometer.

**pygidium** (*adj.* pygidial) The posterior part of the *\*exoskeleton* of a trilobite (*\*Trilobita*); it is generally formed by the fusion of several body segments but in some *\*Cambrian* forms consisting of a single segment. Many Cambrian trilobites have small pygidia and are said to be 'micropygous'. Most trilobites are either isopygous, where the *\*cephalon* and pygidium are the same size, or heteropygous, where the pygidium is the smaller. In some cases the pygidium is larger than the head (macropygous).

**pyknometer** See PYCNOMETER.

**pyralspite** See GARNET GROUP.

**pyramid** A *\*crystal* form composed of a number of non-parallel *\*crystal* faces which meet at a point. The crystallographic notation is frequently (111) or {111} if it is a form, whereby the crystal face intersects all the axes at their unit lengths.

**pyramidal peak** See HORN.

**pyrite (fool's gold)** A *\*sulphide* mineral, FeS<sub>2</sub>; sp. gr. 4.9–5.2; *\*hardness* 6.0–6.5; *\*cubic*; pale brass-yellow, does not tarnish; greenish-black *\*streak*; *\*metallic \*lustre*; crystals *\*cubic*, *\*pyritohedra* (pentagonal dodecahedra), *\*octahedra*, or combinations of the two; *\*cleavage* poor basal {001}; occurs with other *\*sulphide* ores genetically associated with *\*basic* and *\*ultrabasic* rocks, and together with *\*pyrrhotite* and *\*chalcopyrite*; very widely distributed in a great variety of environments, and found in *\*igneous* rocks as an *\*accessory mineral*, in *\*sedimentary rocks* (especially black *\*shales*), as nodules in *\*metamorphic rocks*, and common in hydrothermal veins, in replacement deposits, and contact *\*metamorphic rocks*; alters to iron sulphate and *\*limonite*. It was formerly used widely for the production of sulphuric acid. Compare MARCASITE.

**pyritohedron (pentagonal dodecahedron)** A *\*crystal* form within the *\*cubic* system consisting of a twelve-faced, *\*closed* form (a dodecahedron) with each *\*crystal* face having five sides, but not in the shape of a regular pentagon. It is a common form of the mineral *\*pyrite*.

**pyro-** From the Greek *pur* meaning 'fire', a prefix meaning 'fire'.

**pyrochlore** Mineral oxide (Na,Ca,U)<sub>2</sub>(Nb,Ta,Ti)<sub>2</sub>O<sub>6</sub> (OH,F); sp. gr. 4.3–4.5; *\*hardness* 5.0–5.5; *\*cubic*; normally brown to black, green when heated; light brown *\*streak*; vitreous to greasy *\*lustre*; crystals usually octahedra or irregularly shaped grains; *\*cleavage* octahedral when present; found in granitic rocks and *\*pegmatites*, with or near *\*alkaline rocks*, and in association with *\*zircon*, *\*apatite*, and *\*carbonatites*. The calcium and sodium in the crystal *\*lattice* can be substituted with uranium, thorium, or the *\*rare-earth elements*.

**pyroclastic** Literally, 'fire-broken'. Applied to volcanic rocks consisting of fragmented particles, generally produced by explosive action.

**pyroclastic flow (ash-flow)** General term for a hot, high-concentration flow of \*pumice or lithic clasts, entrained and transported in a fluidized \*ash matrix. Pyroclastic flows include a wide range of phenomena, from \*ignimbrites (large volume, \*pumiceous) to block-and-ash flows (small volume, lithic). The flow originates by the gravitational collapse of a dense, turbulent, eruption column at the source vent and moves downslope as a coherent flow. Fluidization of the ash matrix, which contributes to the high mobility of such avalanches, is achieved by (a) the diffusion and release of gas during breakage and the attribution of ash and pumice particles entrained within the flow and (b) air ingested and compressed at the front of the advancing flow margin. Where the ash matrix is the dominant component, the term 'ash-flow' is applied by American authors, although British authors prefer to use the term 'pyroclastic flow'.

**pyroelectricity** The positive and negative electrical charges developed at opposite ends of the polar axis of a \*crystal in response to a change in temperature. It is observed in \*tourmaline.

**pyrogenetic minerals** \*Minerals which are crystallized from a completely or almost completely anhydrous \*magma. These minerals, e.g. \*olivine, \*pyroxene, and \*plagioclase, have anhydrous compositions and are some of the first minerals to crystallize from the magma. If the original magma contained a small amount of water in solution, crystallization of pyrogenetic minerals enriches the residual magma in water.

**pyrolite model** Model for the composition of upper-\*mantle material proposed by Ringwood: one part \*basalt to three parts \*dunite (mainly \*olivine and \*pyroxene), \*partial melting of which gives basaltic \*magma.

**pyrolusite** Common manganese mineral,  $MnO_2$ ; sp. gr. 4.5–5.0; \*hardness 5–6 for crystals, decreasing to 2 when \*massive; \*tetragonal; black to bluish-grey; black \*streak; \*metallic \*lustre; crystals rare, \*acicular and rod like, but it is usually massive, \*cryptocrystalline, and \*dendritic on joints and bedding planes; \*cleavage \*prismatic when present; a \*secondary mineral found in the oxidized zone of manganese deposits, in \*quartz veins, and as nodules on the sea bed. Synthetic  $MnO_2$  is used for dry batteries, as a decolourizer for glass, and in the manufacture of chemicals.

**pyrolysis** The heating of organic molecules without oxygen, to produce \*hydrocarbons (\*char) which have a high calorific value. Fuel produced by

this process can be concentrated and stored. Organic waste may be used.

**pyrometasomatic deposit (skarn)** Deposit formed by *\*metasomatism*, usually in *\*limestones*, at or near *\*igneous* contacts, due to the passage of mineralizing solutions through reactive rocks.

**pyromorphite** Mineral,  $Pb_5(PO_4)_3Cl$ ; sp. gr. 6.5–7.1; *\*hardness* 3.5–4.0; *\*hexagonal*; various shades of green, yellow, and brown; white *\*streak*; vitreous *\*lustre*; crystals *\*prismatic* and often hollow, or barrel-shaped, forming aggregates or crusts, also occurs granular, fibrous or globular; *\*cleavage* prismatic in traces; a *\*secondary* mineral occurring in the oxidized zones of *\*veins* containing lead minerals, often associated with mimetite and *\*anglesite*, and *\*galena*.

**pyrope** Member of the *\*garnet* group of minerals,  $Mg_3Al_2(SiO_4)_3$ ; sp. gr. 3.51; *\*hardness* 6.0–7.5; *\*cubic*; dark red, pink-red, or black; vitreous *\*lustre*; crystals (dodecahedra, also occurs *\*massive*; occurs in high-grade *\*metamorphic rocks* (e.g. *\*eclogites*) and deep-seated *\*igneous* rocks (e.g. garnet *\*peridotites*), which are often *\*mantle* derived, although the pyrope molecule occurs to a variable extent in most *\*almandines*; it is also distributed in *\*alluvial* deposits. Transparent crystals are used as *\*gemstones*. The name is derived from the Greek *puropos*, meaning ‘fiery eyed’.

**pyrophyllite** Uncommon *\*silicate* mineral  $Al_2[Si_4O_{10}](OH)_2$  belonging to the *\*phyllosilicates* (sheet silicates) and with properties similar to *\*muscovite*; sp. gr. 2.65–2.90; *\*hardness* 1–2; occurs as a secondary product from the hydrothermal alteration of *\*feldspar*, and as foliated masses (see FOLIATION) in metamorphic *\*schists*. It has been mined as a substitute for *\*talc*.

**pyroxene** An important group of *\*inosilicates* (chain silicates) comprising the *\*orthorhombic* pyroxenes (*\*orthopyroxenes*) and the *\*monoclinic* pyroxenes (*\*clinopyroxenes*) with the general formula  $XYZ_2O_6$ , where X = Mg, Fe, Ca, or Na; Y = Mg, Fe,  $Fe^{3+}$ , or Al; and Z = Si (and some Al substitution). The main orthopyroxenes are *\*enstatite* and orthoferrosilite; the main clinopyroxenes include *\*diopside*, *\*hedenbergite*, *\*augite*, *\*pigeonite*, and *\*jadeite*, and also the alkali pyroxenes *\*aegirine* and *\*aegirine* augite (note that *\*wollastonite*, although similar to the pyroxenes,

has a different atomic structure); sp. gr. 3.0–3.5; \*hardness 5–6; colours variable, but usually dark greens, browns, or black; \*vitreous \*lustre; \*crystals usually short or columnar \*prisms; well-marked \*cleavage; widely distributed in both \*igneous and \*metamorphic rocks.

**pyroxene gneiss** See GRANULITE.

**pyroxene hornfels facies** A set of metamorphic \*mineral assemblages (produced by the \*metamorphism of a wide range of starting rock types under the same metamorphic conditions), typically characterized by the development of the mineral assemblage \*clinopyroxene–\*labradorite–\*quartz in rocks of \*basic \*igneous composition. Other rocks of contrasting composition, e.g. \*shales or \*limestones, would each develop their own specific mineral assemblage, even though they are all being metamorphosed under the same conditions. The variation of mineral assemblage with starting rock composition reflects a particular range of pressure, temperature, and  $P(\text{H}_2\text{O})$  conditions. Experimental studies of mineral pressure–temperature stability fields indicates that the \*facies represents a range of low pressures (0–2 kb) and moderate temperatures (550–750 °C) characteristic of contact metamorphic conditions.

**pyroxenite** An \*ultrabasic, \*igneous rock consisting of \*essential \*clinopyroxene, \*orthopyroxene, and \*olivine. The \*ferromagnesian minerals are magnesium-rich, the pyroxenes dominating olivine, which is usually less than 40% by volume. Different types of pyroxenites are defined using the ratio of clinopyroxene to orthopyroxene to olivine. These include orthopyroxenite, websterite, clinopyroxenite, and olivine-bearing versions of these. Pyroxenites are found as \*cumulate horizons within layered \*basic intrusions and as components of the sub-oceanic \*mantle in obducted fragments of \*ocean crust known as \*ophiolites.

**pyroxenoid** See WOLLASTONITE.

**pyrrhotite** Mineral, FeS; sp. gr. 4.6–4.7; \*hardness 3.5–4.5; \*hexagonal; bronze-yellow but darkens rapidly to reddish-brown on exposure; greyish-black \*streak; \*metallic \*lustre; \*crystals rare, and can be \*platy or \*tabular, but the mineral is usually \*massive or granular; no \*cleavage; magnetic; occurs in \*igneous rocks, e.g. \*gabbro or \*norite, as disseminated grains, and also in \*metamorphic contact zones in association

with *\*chalcopyrite* and *\*pyrite*. Pyrrhotite was used formerly for the production of sulphuric acid.

**p-zone** A biostratigraphic *\*zone* distinguished by *\*pelagic \*fossils*, e.g. ammonites (*\*Ammonoidea*) and graptolites (*\*Graptolithina*). The term was proposed in 1965 by T. G. Miller. *Compare B-ZONE*.



**Q 1.** See **KÖNIGSBERGER RATIO**. **2.** The ratio of the frequency at the midpoint of a **\*band-pass filter** to the filter width. **3.** The ratio of the peak energy in a waveform to the energy lost by dissipation. As a measure of seismic absorption it has been used effectively in delimiting **\*Benioff zones** using natural **\*earthquake** events (high Q implies low absorption and is associated with **\*crust** and deep **\*mantle**; low Q implies high absorption and is associated with the sub-crustal **\*low-velocity layer**).

**QAPF classification** A modal classification scheme for **\*igneous** rocks with a **\*colour index** less than 90, based on the relative proportions of: **\*quartz** to quartz plus total **\*feldspars** (Q); **\*alkali** feldspar to total feldspars (A); **\*plagioclase** to total feldspars (P); and **\*feldspathoids** to feldspathoids plus total feldspar (F).

**QAP triangle** A three-component, modal classification scheme for **\*igneous** rocks of granitic affinity based on the relative proportions of: **\*quartz** (Q); **\*alkali** feldspar (including albite if the anorthite content is less than 5%) (A); and **\*plagioclase** feldspar (containing more than 5% anorthite (P)).


**QBITO** A 2-unit **\*CubeSat** mission by the Universidad Politécnica de Madrid, Spain, to study the properties of the lower **\*thermosphere**. It was launched on 18 April 2017, from Cape Canaveral, to the International Space Station and deployed from there into orbit at an altitude of about 400 km.

**Q days** See **QUIET DAYS**.

**Q-factor** See **KÖNIGSBERGER RATIO**.

**QFL diagram** (**QFL triangle**) A triangular diagram that shows the composition of **\*sandstones** and **\*sands** as the proportions of **\*Quartz**,

\*Feldspar, and \*Lithic fragments, as determined by \*point counting using the \*Gazzi-Dickinson method.



QFL diagram

**QFL triangle** See QFL DIAGRAM.

**Q<sub>n</sub>** See KÖNIGSBERGER RATIO.

**QSat-EOS** See KYUSHU SATELLITE FOR EARTH OBSERVATION SYSTEM DEMONSTRATION.

**quadrature** See IMAGINARY COMPONENT.

**quaquaversal** Applied to dome structures that dip from a centre in every direction.

**quarrying** 1. An *\*exogenetic process* involved in coastal *\*erosion*, in which waves dislodge material by striking rocks with great force and compressing the air in cracks and the spaces between loose particles. Unconsolidated or loosely consolidated (see *CONSOLIDATION*) *\*sedimentary rocks* and rocks with many *\*joints* are especially susceptible. 2. See *GLACIAL PLUCKING*.

**quartz (rock crystal)** Widely distributed rock-forming *\*silicate* mineral  $\text{SiO}_2$  with the related semi-precious varieties rose quartz (pink), amethyst (purple), cairngorm (dark brown and similar to smoky quartz, although the original cairngorms from the Scottish mountain range were of *\*topaz*), citrine (light brown); sp. gr. 2.65; *\*hardness* 7; *\*trigonal*; commonly colourless or white, but can occur in a variety of colours; *\*vitreous lustre*; *\*crystals* usually six-sided *\*prisms* terminated by six-faced pyramids, the prisms often striated, also occurs extensively in *\*massive* form; no *\*cleavage*; *\*conchoidal* fracture; found in many *\*igneous* and *\*metamorphic rocks*, extensively in *\*clastic rocks*, and a common *\*gangue* mineral in mineral *\*veins*. See also *AGATE*; *CHALCEDONY*; *COESITE*; *CRISTOBALITE*; *FLINT*; *JASPER*; *ONYX*; *OPAL*; *STISHOVITE*; *TRIDYMITE*.

**quartz arenite** See *DOTT CLASSIFICATION*.

**quartz dolerite** See *DOLERITE*.

**quartzite** A *\*metamorphic rock* composed mainly of *\*quartz* and usually formed by the *\*metamorphism* of quartz *\*sandstones*. If deformation has accompanied metamorphism the individual quartz crystals are elongate and take on a preferred orientation, defining a planar or linear *\*fabric* within the rock. Unmetamorphosed sedimentary quartzites (*\*sandstones* with a quartz *\*cement*) are known as 'orthoquartzites' to distinguish them from metamorphic quartzites (metaquartzites).

**quartz overgrowth** The development of *\*quartz cement* around *\*detrital grains*, the quartz cement growing in *\*optical* continuity with the grains which they have enclosed.

**quartz porphyry (elvan)** A *\*porphyritic microgranite*, *\*microgranodiorite*, or microtonalite.

**quartz sandstone** See *ARENACEOUS*.

**quartz wacke** A *sandstone* containing more than 15% mud *matrix*, with over 95% of the *grains* being *quartz*. See DOTT CLASSIFICATION.

**quartz wedge** See ACCESSORY PLATE.

**quasi-equilibrium** See EQUILIBRIUM.

**quasi-section** See PSEUDOSECTION.

**Quaternary (Pleistogene, Anthropogene)** Traditionally, either a sub-era of the *Cenozoic era* of geologic time that began 1.806 Ma ago and continues to the present day; it is sometimes known as the Pleistogene period and includes the *Pleistocene* and *Holocene epochs*, during which there have been numerous ice-sheet advances in the northern hemisphere. However, the boundary between the Quaternary and the preceding *Tertiary* was never clearly defined and the International Commission on Stratigraphy proposed abandoning both terms, Tertiary and Quaternary, as formal names. Quaternary researchers objected strongly, so that name (but not Tertiary) has been retained, commencing either 2.588 Ma ago at the start of the Gelasian stage (2.588–1.806 Ma ago) or about 3.6 Ma ago at the start of the preceding *Piacenzian* stage (3.6–2588 Ma ago).

**quaternary system** A *mineral* system with four components, for example, the four-component system for *diopside* (Di)–*anorthite* (An)–*albite* (Ab)–*pyroxene* (Fs), diagrammatically represented as a tetrahedron. See COTECTIC SURFACE; PHASE; PHASE DIAGRAM.

**quenching** The sudden chilling of *magma*, which may provide a ‘snapshot’ of the *mineral phases* that were in equilibrium under a particular set of temperature and pressure conditions. Quenching is much used in experimental *petrology* to investigate the course of mineral reactions.

***Quetzalcoatlus northropi*** A pterosaur (*Pterosauria*) first discovered in 1975 in Texas. It was huge, with a wing-span of 10m or more, and lived as do vultures, using warm air thermals to soar high over the *Cretaceous* plains.

**quick clay** *Clay* which becomes movable or semi-liquid and loses all *shear strength* when disturbed, especially when saturated. This may happen naturally due to seismic activity.

**quickflow** The part of a storm rainfall which moves quickly to a stream channel via *\*surface runoff* or *\*interflow*, and forms a flood wave in the channel.

**quiet days (Q days)** The geomagnetic intensity and directions for the five least magnetically disturbed days in each month are averaged to determine the quiet-day variations of the *\*geomagnetic* field. See **DIURNAL VARIATION** (geomagnetic).

**quiet zone** See **MAGNETIC QUIET ZONE**.

**QuikSCAT** A NASA satellite that was launched on 19 June 1999 into an elliptical *\*Sun-synchronous orbit* at a maximum height of 800 km. The name is an abbreviation of Quick Scatterometer and the satellite carries the SeaWinds instrument, which is a specialized microwave radar that measures low-level wind speed and direction. The purpose of the mission is to gather all-weather, high-resolution measurements of near-surface winds, observe atmosphere–ocean interactions, study daily and annual changes in the Arctic and Antarctic sea ice, and study changes in rain forest vegetation. The resulting data will help improve weather forecasts in coastal areas and improve storm warning and monitoring. The mission was still operating in 2018.



<https://winds.jpl.nasa.gov/missions/quikscat/>

- Missions: QuikSCAT.

**Qungzusian** A Chinese *\*stage* (532–523 Ma ago) of the Early *\*Cambrian*, preceded by the *\*Meishuchuan* and followed by the *\*Changlangpuan*.

**Q-wave** See **LOVE WAVE**.



**Ra** See RAYLEIGH NUMBER.

**radar** Acronym for *radio detection and ranging*; the use of electromagnetic energy for the detection of objects which are capable of reflecting it. For example, cloud-detection radars are extensively used in meteorological forecasting and rainfall measurement, while sideways-looking radars are used for topographic mapping. Radar has been used to probe through **\*ice sheets** in order to determine ice thickness and to detect internal reflection events; in polar regions the radar equipment has usually been airborne, but the instruments can be mounted on sledges for surface use. Radar can also be used in arid environments to probe through sand in search of water. Ground radar is being developed for use in engineering site investigations, but has very limited depth penetration where the moisture content is high because of the high **\*dielectric loss** associated with water.

**radar altimetry** The measurement of the surface relief of a planetary body by recording the travel time of reflected **\*radar** waves. The most spectacular success of the technique has been the establishment of the detailed surface topography of **\*Venus** by the US probes Pioneer Venus 1 and 2 launched in 1978, and the Soviet Venera 15 and 16 launched in 1983. Radar scanning of the Earth's surface from aircraft or satellites provides information on topography, vegetation cover, and **\*lithology**, and is particularly valuable where cloud cover prevents normal surveying. Over the oceans, Seasat altimetry data provide information on wave height and motion, and on ocean **\*bathymetry**, and assist in definition of the **\*geoid**.

**radar cross-section** In **\*radar** terminology, the hypothetical area of a perfect **\*diffuse** reflecting surface which would be required to return the same amount of energy to a radar antenna as is observed from a point

target. The radar cross-section serves as a measure of the *\*intensity* of the energy that is backscattered from a point target.

**radar imaging** The use of *\*radar* wavelengths to determine the location, size, and reflectivity characteristics of objects. It is particularly useful in regions with dense cloud cover. *See also* REMOTE SENSING.

**Radar Imaging Satellite (RISAT)** A reconnaissance satellite of the Indian Space Research Organization that uses *\*synthetic-aperture radar* with an all-weather, 24-hour monitoring capability. RISAT-1 was launched in 2012 and broke up in 2017. RISAT-2 was launched on 20 April 2009, from Satish Dhawan Space Centre, India, into a near-circular orbit at an altitude of 548 km.

**Radar in a CubeSat** *See* RAINCUBE.

**RADARSAT** A Canadian satellite mission to monitor the environment, provide data to help manage natural resources, and perform coastal surveillance. RADARSAT-1 was launched on 4 November 1995 and retired on 9 May 2013. RADARSAT-2 was launched on 14 December 2007, from Baikonur, Kazakhstan, into a Sun-synchronous *\*polar orbit* at an altitude of 798 km. The RADARSAT Constellation Mission (RCM), complementary of RADARSAT-2 and concentrating on disaster management and ecosystem monitoring, is scheduled for launch in May 2019.



<http://www.asc-csa.gc.ca/eng/satellites/radarsat/default.asp>

- RADARSAT Constellation Mission home page.

**radar scattering coefficient** The average *\*radar cross-section* value per unit area of a large target. This is the fundamental measurement of the radar properties of a surface, allowing discernment of the texture of the ground.

**radar scatterometer** A device which measures the electromagnetic energy backscattered from a target as a function of the *\*depression angle*.

**radial drainage** *\*Drainage* pattern consisting of streams that extend radially from a central zone. It is typical of the patterns developed on freshly constructed land-forms (e.g. *\*volcanoes*) and on areas of domed uplift.

**radial dykes** Vertical to subvertical **\*dykes** which radiate from a central volcanic **\*plug**. Radial dykes associated with the emplacement of **\*plutons** at depth may theoretically lie along the minimum **\*principal** stress axis  $\sigma_3$  within the stress field that is generated.

**radial fault** One of a system of **\*faults** which radiate from a central point.

**radial fibrous** A **\*mineral texture** seen where crystals are aligned perpendicularly to the curved surface on which they are growing.

**radial relief displacement** The apparent leaning away from the **\*centre** point of vertical objects in an aerial photograph, due to the conical field of view of the camera lens.

**radial symmetry** The condition in which the body of an organism is repeated in a circular manner. In **\*corals** the repetition occurs around the mouth, and in some echinoderms (**\*Echinodermata**), where the five rays of the animal are symmetrically placed, the symmetry is also radial.

**radiance** The **\*radiant** flux density of electromagnetic radiation measured by a **\*remote** sensing detector as it travels through a given solid angle.

**radiant flux density** The power of **\*electromagnetic** radiation falling on or emanating from a body, measured as watts per square metre.

**Radiata** The former subkingdom of animals (kingdom Animalia) that includes the coelenterates (**\*Coelenterata**) and Placozoa (a phylum of flat, amoeboid animals 2–3 mm in size). Members of the Radiata possess **\*radial symmetry**. The term is no longer accepted.

**radiating** A crystal form in minerals, where the crystals grow outwards from a central point, frequently diverging to give a concentric pattern of growth.

**radiation** Any form of emitted wave phenomena, usually the **\*electromagnetic spectrum**, sound, or heat.

**Radiation Belt Storm Probes** See VAN ALLEN PROBES.

**radiation budget (energy budget)** The difference between the amount of incoming solar radiation and the amount of outgoing terrestrial radiation. The balance is in deficit (i.e. more energy leaves the Earth's surface than

reaches it from the Sun) at night. Overall, the highest positive net balance is found in low latitudes.

**radiation densimeter** An instrument employed to measure the intensity of radiation.

**radiation fog** Condensation effect over land surfaces on clear nights with light breezes, due to surface radiation cooling. Favoured initial conditions are very humid air, with wet and cold surfaces, e.g. marshes. The fog forms when moist air crosses a cold land or sea surface, reducing the air temperature to below the *\*dew-point*. The fog, most common in winter, is generally cleared by the Sun's warmth in the morning, but thick fog over wet surfaces in winter may persist much longer, particularly if an upper cloud layer screens the Sun.

**radiation inversion** *\*Temperature* inversion in the lower atmosphere due to radiation cooling of the ground at night. *See also* [RADIATION NIGHT](#).

**radiation night** A night with clear skies, when terrestrial long-wave radiation cannot be partly returned to the surface by cloud. There is rapid cooling of the air close to the ground, particularly when there is little wind, giving low minimum surface temperatures.

**radiation tracks** *See* [FISSION-TRACK DATING](#).

**radiatus** From the Latin *radiatus* meaning 'with rays', a variety of cloud with parallel bands which, due to perspective, appear to meet at the horizon. The effect is seen in *\*cumulus*, *\*stratocumulus*, *\*altocumulus*, *\*altostratus*, and *\*cirrus*. *See also* [CLOUD CLASSIFICATION](#).

**radiaxial** Radially-axial, applied to *\*crystals*, seen particularly in *\*calcite*, which form a radiating crust of the *\*mineral* within cavities.

**radioactive decay** Process by which a radioactive 'parent' element loses elementary particles from its nucleus and in doing so becomes a stable 'daughter' element. The rate of decay is constant for a given element and is a very precise and accurate device for the measurement of geologic time. *See also* [DECAY CONSTANT](#); [DECAY CURVE](#).

**radioactive logging** *See* [DENSITY SONDES](#); [GAMMA-GAMMA SONDE](#); [GAMMA-RAY SONDE](#); [NEUTRON-GAMMA SONDE](#); [NEUTRON-NEUTRON](#)

SONDE.

**radioactive survey** A survey to measure the natural radioactivity of a region, usually by means of a scintillometer, **\*spectrometer**, or **\*Geiger counter**. It may be a ground survey, or made from a low-flying aircraft, usually a helicopter.

**radioactive waste** Any discarded substance that is radioactive. Wastes are classified as high-, intermediate-, or low-level according to their level of radioactivity. Low-level waste includes clothing and materials which have been used when handling radioactive sources, e.g. in hospitals. It can be safely buried in trenches 9 m deep beneath a covering of 2 m of clay; no alpha or beta radiation could penetrate the clay cover. Intermediate- and high-level wastes are mainly from the **\*fission** process in nuclear power stations or from military waste. High-level waste is hot and intensely radioactive. It is stored, usually in ponds of water, for up to 50 years, during which time it cools and its short-lived isotopes decay until it can be classed as intermediate-level. It can then be incorporated in a borosilicate glass or synthetic rock (a **\*synroc**), sealed in a container which corrodes at a known rate, and stored in a secure surface or underground facility. After 500 to 1000 years the radioactivity will have decayed sufficiently for the waste to emit no more radiation than many naturally occurring rocks.

**Radio Aurora eXplorer (RAX)** A US 3-unit **\*CubeSat** mission to study **\*plasma** instabilities that lead to irregularities in the alignment of the **\*geomagnetic** field. RAX-1 was launched on 30 November 2010, from Kodiak Island, Alaska, into a circular orbit at an altitude of 650 km. A follow-on mission, RAX-2, was launched on 20 October 2011, into a Sun-synchronous near-circular **\*polar orbit** at an altitude of 824 km.

**radiocarbon dating (<sup>14</sup>C dating)** A dating method for organic material that is applicable to about the last 70 000 years. It relies on the assumed constancy over time of atmospheric <sup>14</sup>C:<sup>12</sup>C ratios (now known not to be valid), and the known rate of decay of radioactive carbon, of which half is lost in a period (the 'half-life') of every 5730 ± 30 years. (The earlier 'Libby standard', 5568 years, is still widely used.) In principle, since plants and animals exchange carbon dioxide with the atmosphere constantly, the <sup>14</sup>C content of their bodies when alive is a function of the radiocarbon content of the atmosphere. When an organism dies, this exchange ceases

and the radiocarbon fixed in the organism decays at the known half-life rate. Comparison of residual  $^{14}\text{C}$  activity in fossil organic material with modern standards enables the age of the samples to be calculated. Since the method was first devised it has been realized that the atmospheric  $^{14}\text{C}$  content varies, as the cosmic-ray bombardment of the outer atmosphere that generates the  $^{14}\text{C}$  varies. Correction for these fluctuations is possible for about the last 8000 years by reference to the  $^{14}\text{C}$  contents of long tree-ring series, e.g. those for bristlecone pine (*\*Pinus longaeva*).

**radiogenic** Applied to *\*isotopes* produced by the process of radioactive decay.

**radiogenic heating** The thermal energy released as a result of spontaneous nuclear disintegrations. In the *\*Earth*, the major *\*isotopes* concerned today are of the elements uranium, thorium, and potassium, but various short-lived isotopes may have been important during the early formation of the Earth.

**radioimmunoassay** A technique for the very precise analysis of proteins, based on the ability of unlabelled proteins to inhibit competitively the binding of labelled protein by specific antibodies (i.e. an immunological reaction). The protein concentration of the unknown sample is determined by comparing the degree of inhibition with that produced by a series of standards containing known amounts of the protein. The technique has been adapted to assay non-proteins.

**Radiolaria (Radiozoa)** A large group of marine sarcodinids (*\*protists*), characterized by having a shell with a perforated, membraneous capsule containing the endoplasm, and a siliceous or strontium sulphate skeleton consisting of a lattice shape of variable morphology made up of *\*spicules*, bars, and spines. Radiolarians live mainly in surface waters and the earliest forms are *\*Cambrian* in age. They are used in the biostratigraphic correlation of oceanic sediments, particularly where calcareous *\*microfossils* have been dissolved.

**radiolarian earth** An unconsolidated or semi-consolidated bed of *\*radiolarian ooze*. It is a relatively rare deposit, as most ancient radiolarian oozes have been compacted into hard *\*radiolarites*.

**radiolarian ooze** Deep-sea ooze in which at least 30% of the sediment consists of the siliceous radiolarian (*\*Radiolaria*) *\*tests*. Radiolarian-rich oozes occur in the equatorial regions of the *\*Pacific* and *\*Indian Oceans* where the depth exceeds the *\*carbonate compensation depth* (around 4500 m in the central Pacific). *See also* DIATOM OOZE.

**radiolarite** A compacted, siliceous, *\*sedimentary rock* composed mainly of the siliceous *\*tests* of the marine zooplankton called *\*Radiolaria*. The sediment from which radiolarites are formed is called a *\*radiolarian ooze*.

**radiometer** An instrument to measure radiation, but usually a device to monitor infrared radiation. It can be precisely tuned for specific frequencies, but usually has poor spatial resolution.

**Radiometer Assessment using Vertically Aligned Nanotubes (RAVAN)** A *\*NASA* 3-unit *\*CubeSat* mission to demonstrate a radiometer that will measure Earth's radiative diurnal cycle and absolute energy imbalance to climate accuracies (globally at  $0.3 \text{ Wm}^{-2}$ ). It was launched on 11 November 2016, from California, into a Sun-synchronous near-circular orbit at an altitude of 617 km.

**radiometric dating (radioactive dating)** The most precise method of dating rocks, in which the relative percentages of 'parent' and 'daughter' *\*isotopes* of a given radioactive element are estimated. Early methods relied on uranium and thorium minerals (*see* URANIUM–LEAD DATING), but *\*potassium–argon*, *\*protactinium–thorium*, *\*rhenium–osmium*, *\*rubidium–strontium*, *\*samarium–neodymium*, and *\*carbon-14–carbon-12* are now of considerable importance. Uranium-238 decays to lead-206 with a half-life of 4.5 billion ( $10^9$ ) years, rubidium-87 decays to strontium-87 with a half-life of 50.0 billion years, and potassium-40 decays to argon-40 with a half-life of 1.5 billion years. For carbon-14 the half-life is a mere  $5730 \pm 30$  years (*see* RADIOCARBON DATING), and beyond about 70 000 years the amount of carbon-14 remaining in organic matter is beyond accurate measurement. *Compare* INITIAL STRONTIUM RATIO.

**radiometry** The measurement of incident radiation by a *\*radiometer*. A typical usage was the infrared scanning radiometer on the Apollo 17 lunar mission which used a *\*thermistor* mounted in a small telescope on the orbiting lunar command vehicle to record lunar surface temperatures.

**radio occultation** The technique of using radio waves transmitted from a spacecraft to probe the atmospheres of planets. The signals received on the Earth as the spacecraft swings behind a planet provide information about both the vertical structures of the planetary **\*ionospheres** and the atmospheric density. Monochromatic radio waves provide more accurate measurements than those obtained using stellar occultations.

**radiosonde** Instrument, comprising an **\*aneroid** barometer and sensors for temperature and humidity, that is carried aloft to the upper atmosphere by a balloon that ascends at about 5 m/s. Data is transmitted to the surface by radio. *See also* **RAWINSONDE**.

**Radiozoa** *See* **RADIOLARIA**.

**radius** In **\*tetrapods**, the bone of the fore limb on the side of the first digit (in humans, the thumb). *Compare* **ULNA**.

**radius ratio** Radius of a **\*cation** divided by that of an **\*anion**. In **\*silicate** minerals, the radius of a cation divided by that of the oxygen anion. Most minerals have bonds that are mainly ionic, therefore radius ratios are most useful in studying mineral structures. **\*Ions** tend to pack closely together in a low-energy, stable state and the most stable arrangements are classified in terms of the cation-to-anion radius ratio. Oxygen is the major anion in most minerals, so the radius ratio of various cations to oxygen is particularly important. This controls the **\*coordination numbers** and therefore the structures of silicates, e.g. chains, frameworks, etc.

**raft foundation** Type of **\*foundation** used for heavy loading, or construction on soft ground, composed of a continuous slab of reinforced **\*concrete** below the entire surface. In Venice, many of the mediaeval raft foundations underneath the large churches are made of wood.

**rainbow** Phenomenon in which an arc of colours of the spectrum results from the double reflection and **\*refraction** of the Sun's rays by water droplets in the atmosphere. A primary bow is seen at a limiting angle of 42° to the observer's shadow. A secondary bow, at 51°, may occur; as can multiple bows due to some light being reflected within the raindrops before refraction. The colour separation in the primary bow produces a spectrum (violet, indigo, blue, green, yellow, orange, and red) with red on the outer edge. The colours are reversed in the secondary (outer) bow. The colour and

intensity vary with the size of raindrops, large drops producing bright colours dominated by red.

**RainCube (Radar in a CubeSat)** A *\*nanosatellite* mission of *\*NASA* and the Jet Propulsion Laboratory to use profiling radar to monitor the short-time evolution of weather systems. It was launched on 21 May 2018, from Virginia, into a near-circular orbit at an altitude of about 400 km.

**raindrop** A water droplet, formed by condensation of water vapour in a cloud, that is heavy enough to fall from the cloud and large enough to reach the surface of land or sea before evaporating in the unsaturated air beneath the cloud. Droplets reaching the surface range in diameter from about 100  $\mu\text{m}$  in fog to 0.2 mm in drizzle and 5.0 mm in a heavy shower.

**rain-gauge** A device for measuring the rainfall in a specified period, usually one day. A standard gauge comprises a cylinder 20 cm in diameter, with an upper surface in the shape of a funnel, set vertically on open ground with its upper rim 1m above ground level. Rain flows through the funnel into a measuring tube 6.32 cm in diameter. The depth of water in the tube is 10 times the rainfall amount. In an alternative design, water passes from the first funnel to a second, smaller funnel, and from there to one of two 'tipping buckets'. When one bucket is full, it descends, makes an electrical contact, and empties itself, while the second bucket is positioned to collect water. Because the buckets make an electrical contact each time they collect a predetermined weight of water the rain gauge provides a record of rainfall and does not require regular emptying.

**rain-making** Attempt to induce rainfall by 'seeding' supercooled water clouds. *See also* [CLOUD SEEDING](#).

**rain print** Small, crater-like pits on the surface of a sedimentary rock, made by rain that fell while the surface was still soft.

**rain-shadow** The reduction of rainfall to the lee side of a mountain barrier, which results in relatively dry surface conditions, e.g. in the mountains of the south-western USA, where the wetter western slopes of the Coast Range and Sierra Nevada contrast with the desert areas of Nevada and eastern California on the lee side of the mountains. Moist air rising on the windward side cools at the *\*saturated adiabatic lapse rate*, its moisture condensing to produce *\*precipitation*. Air subsiding on the lee side is dry

and warms at the **\*dry adiabatic lapse rate**, which further reduces its **\*relative humidity**.

**rain-splash** See RAIN-WASH.

**rain-wash** A general term for the transfer of material across the surface and down a hillslope as a result of rainfall. Normally it consists of two components: rain-splash, which is the detachment and subsequent downslope transfer of small soil particles by raindrop impact; and soil-wash, which is the downslope movement of material by surface water flow. Rain-wash and **\*creep** are the two main hillslope processes.

**raised beach** Former beach, now found above the level of the present shoreline as a result of earth movement or of a general fall in sea level. Such beaches are frequently described and correlated in terms of height above present sea level.

**raised bog** A peat-forming community of plants, typically dome-shaped, that develops on former lake **\*sediments**, **\*estuaries**, uniform **\*clay** substrates, and sometimes on the surface of **\*valley bogs**.

**rake** See PITCH.

**Ramapithecus** Late **\*Miocene** and early **\*Pliocene** ape, known from fragmentary **\*fossils** from E. Africa, south-eastern Europe, and northern India and Pakistan, dating from 14–10 Ma ago, and apparently identical or very similar to the E. African *Kenyapithecus*. *Ramapithecus* is regarded by many as transitional between the true **\*Miocene** apes (the Dryopithecinae) and the later **\*Hominidae**. If this is so, then the human and ape lines diverged prior to the late Miocene, 15–25 Ma ago. More recent evidence, however, suggests that *Ramapithecus* and the related or identical **\*Sivapithecus** are nearer to the evolutionary line that led to the orang-utan.

**rammer** 1. Part of the equipment for tunnelling, which pushes into the face. 2. A tree trunk of suitable dimensions fitted with iron handles and used to ram an **\*auger** into bedrock.

**ramose** Branched.

**ramp** 1. That part of a staircase **\*thrust** trajectory which forms the steeply dipping sections between the **\*flats**. Where thrusting occurs in horizontally bedded strata the ramps cut up-section, obliquely to the bedding. A thrust

belt may contain several types of ramp, classified as frontal, oblique, and lateral according to their respective perpendicular, oblique, and parallel *\*strike* orientations in relation to the main direction of transport. 2. See SHELF.

**rampart craters** Martian impact craters with diameters of 5–15 km, with *\*ejecta* sheets extending about one crater radius and terminating in a low ridge or escarpment. The rampart is a primary feature, and not due to secondary modification.

**Ramsden eyepiece** See EYEPIECE.

**Randian** A South African *\*stage* of the Upper *\*Archaean*, from about 3000–2500 Ma ago, preceded by the *\*Swazian* and followed by the *\*Vaalian*, that includes the Ventersdorp, Witwatersrand, and Dominion Reef sequences.

**random sampling** See SAMPLING METHODS.

**range** In *\*radar* terminology, the distance of radar propagation. In order to avoid interference between consecutive electromagnetic pulses, the interval between pulses must be adequate to allow the return of the previous pulse. Pulse frequency is inversely proportional to the range. Range is measured as either *\*slant* range or *\*ground* range.

**Ranger** A series of *\*NASA* lunar missions, which ran from 1964 to 1965, sending instruments to the surface.



<http://nssdc.gsfc.nasa.gov/planetary/lunar/ranger.html>

- A NASA mission to obtain close-up images of the surface of the Moon.

**range zone** Unit of *\*strata* defined by the presence and time range of a particular *\*fossil* *\*taxon*. A range zone comprises the entire vertical and horizontal extent of the given organism. Range zones may be local (*\*teilzone*, or local range zone), or the term may be used to refer to the total stratigraphic range of a particular taxon (*\*taxon range zone*). When used formally (see FORMAL) the term is often capitalized and the qualifying fossil name given in italics, with the generic name capitalized and the specific

name in lower-case letters, e.g. the stratigraphic range of the Late **\*Jurassic** **\*ammonite** *Cardioceras cordatum* delimits the *Cardioceras cordatum* Range zone.

**rank** 1. Category of **\*stratigraphic unit** or **\*geologic-time** unit, classed according to magnitude or duration. 2. The grade or purity of a substance (referring particularly to **\*coal**).

**rapakivi texture** Rounded crystals of pink, potassic **\*feldspar**, mantled by white rims of sodic **\*plagioclase** feldspar (occasionally rhythmically zoned with **\*alkali** feldspar), and found as large crystals, up to 4 cm in diameter, in a finer-grained, **\*igneous** **\*groundmass** of granitic composition. The term was applied first to **\*granites** from eastern Finland.

**RapidEye Earth Observation Constellation** A US commercial mission of five **\*minisatellites** to provide Earth observation relating to agriculture, environment, forestry, and mapping, as well as intelligence and defence applications. The constellation was launched on 29 August 2008, from Baikonur, Kazakhstan, into a Sun-synchronous orbit, with the satellites evenly spaced in a single orbital plane, at an altitude of 630 km.

**rapid flow (shooting flow)** See **CRITICAL FLOW**; **FROUDE NUMBER**.

**rapid-neutron process (r-process)** Neutron-capture chain which takes place on a very short timescale. It is believed to result from the gravitational collapse of supernovae, leading to a thermonuclear explosion which is capable of forming very large **\*nuclides** in a matter of seconds. See also **NUCLEOSYNTHESIS**.

**Rapitan-type granular iron formation** A **\*formation** of **\*hematite** and **\*jasper** associated with glacially derived **\*turbidites** and **\*diamictites** that was deposited in a **\*rift** basin, first recognized in northwestern Canada. It is the rarest of the granular iron formation types. See also **ALGOMA-TYPE GRANULAR IRON FORMATION**; **SUPERIOR-TYPE GRANULAR IRON FORMATION**.

**rare-earth element (REE; lanthanide)** One of those elements with atomic numbers between 57 and 71, that have closely similar chemical properties. The ionic radius decreases with increasing atomic number, a phenomenon referred to as the lanthanide contraction. Rare-earth elements occur in

minerals only in trace amounts, sometimes replacing  $\text{Ca}^{2+}$  in *\*apatite* and *\*hornblende*. They tend to become concentrated in the residual fluid of *\*magmas*, and in some *\*pegmatites* the REE cerium replaces the calcium in *\*epidote* to form the mineral *\*allanite*. Lunar rocks, apart from *\*anorthosite*, show considerable enrichment in most of the rare-earth elements relative to the REE *\*cosmic abundance*. See *EUROPIUM ANOMALY*.

**RASAT** A remote-sensing *\*microsatellite* imaging mission by the Space Technologies Research Institute of the Scientific and Technological Research Council of Turkey. It was launched on 17 August 2011, from the Yasný Cosmodrome, Russia, into a Sun-synchronous orbit at an altitude of 700 km.

**raster** In *\*remote* sensing, a grid of *\*pixels* used to store and display digitally recorded, remotely sensed images, produced by a series of lines scanned using a *\*pushbroom*, *\*line scanner*, or *\*radar* system at a regular rate of repetition. A separate raster image is used to represent each spectral band.

**ratio** In *\*remote* sensing, the *\*digital* number value of one band of a multispectral image divided by the digital number value of another band. The ratio allows analysis of relative differences between *\*channels*.

**Raukumara** A *\*series* in the Upper *\*Cretaceous* of New Zealand, underlain by the *\*Clarence* and overlain by the *\*Mata*. It comprises the Arowhanan, Mangaotanean, and Teratan *\*stages* which are roughly contemporaneous with the upper *\*Cenomanian*, *\*Turonian*, *\*Coniacian*, and lower *\*Santonian*.

**RAVAN** See *RADIOMETER ASSESSMENT USING VERTICALLY ALIGNED NANOTUBES*.

**ravinement surface** In *\*sequence* stratigraphy, the first surface to have been formed by flooding due to rising sea level, at or close to the shoreline.

**ravine wind** Wind that passes through an upland barrier along a narrow valley or ravine. The wind is generated by a pressure gradient between the two ends of the valley and the force of the wind is often enhanced as a result of the channelling effect caused by constriction in the valley.

**rawinsonde** \*Radiosonde that is tracked by radio or \*radar to observe wind characteristics as well as to record temperature, pressure and humidity.

**Rawtheyan** A \*stage (446.5–445.6 Ma ago) of the Late \*Ordovician in the \*Ashgill, underlain by the \*Cautleyan and overlain by the \*Hirnantian.

**RAX** See RADIO AURORA EXPLORER.

**Rayleigh, Lord** See STRUTT, JOHN WILLIAM.

**Rayleigh criterion** In \*remote sensing, a method of estimating surface behaviour as either \*specular or \*diffuse with regard to a particular wavelength of \*electromagnetic radiation. A surface is rough if the square root of the mean of the squares of the height of surface irregularities is greater than one-eighth of the wavelength divided by the cosine of the angle of incidence.

**Rayleigh number (Ra)** A dimensionless value used to estimate when convection commences in a fluid. The Ra depends on the density and depth of the fluid, the coefficient of thermal expansion, the \*gravitational field, the temperature gradient (in excess of the \*adiabatic gradient), the thermal diffusivity, and the kinematic \*viscosity. Convection usually starts when Ra is 1000 or more, while heat transfer is entirely by conduction when Ra is less than 10.

**Rayleigh scattering** The scattering of electromagnetic radiation by spherical particles with radii that are less than 10% that of the wavelength of the incident radiation. Such scattering by air molecules produces the blue effect of the sky. Particles such as dust and smoke, that are significantly smaller than 0.4 $\mu\text{m}$  (the wavelength of the blue/violet or lower limit of the visible spectrum) can also scatter visible radiation. Reddish colours at sunset and sunrise result from Rayleigh scattering; these longer wavelengths pass directly through the atmosphere to the observer, while particles in the air scatter out radiation of shorter wavelengths. See also MIE SCATTERING.

**Rayleigh wave** A type of \*surface wave which travels along a free \*interface. Particle motion is elliptical in a plane perpendicular to the interface and retrograde (at the top of the elliptical orbit movement is in the opposite direction to that in which energy is travelling). Rayleigh waves

travel at about 90% of the speed of *\*S-waves* in the same medium. See [GROUND ROLL](#).

**rays** Bright streaks radiating from young lunar craters. They have no surface relief and darken with age, probably due to mixing with the *\*regolith* and to exposure to radiation. They are most probably composed of fine rock powder and glass produced during the impact. The classic example is the spectacular rays emanating from the lunar crater Tycho (85 km diameter), conspicuous through binoculars at full Moon. Hundreds of rays radiate from the crater and some extend across the visible face of the Moon.

**RBSP** See [VAN ALLEN PROBES](#).

**R<sub>E</sub>** Earth radius; 1 R<sub>E</sub> = 6371 km.

**reaction time 1.** In *\*geomorphology*, the time taken for a *\*system* to react to a sustained change in external conditions. Representative reaction times are difficult to define, because of variations both in the resistance of systems to change and in the magnitude of the external change. For example, a sand-bed river channel reacts more readily to change than does a rock-floored channel. See also [RELAXATION TIME](#). **2.** See [CORONA](#).

**reactivation surface** A discontinuity cutting across a *\*foreset*, generated by *\*erosion* or changing flow strength before the resumption of the forward migration of the foreset.

**real-aperture radar** A *\*radar* system where the *\*azimuth resolution* is determined by the physical length of the antenna, *\*wavelength*, and *\*range*. Compare [SYNTHETIC-APERTURE RADAR](#).

**real component** See [IN-PHASE COMPONENT](#).

**realgar** Minor sulphide ore for arsenic, As<sub>2</sub>S<sub>2</sub>, associated with *\*orpiment* (As<sub>2</sub>S<sub>3</sub>) to which realgar changes on exposure; sp. gr. 3.6; *\*hardness* 1.5–2.0 (can be cut with a knife); *\*monoclinic*; red, varying to orange and yellow, transparent to translucent; orange-red *\*streak*; resinous *\*lustre*; crystals rare, short, striated, and *\*prismatic*, also occurs granular, compact, or *\*massive*; *\*cleavage* good *\*pinacoidal*; also found at hot springs and in *\*limestones* and *\*dolomites*; alters to yellow powder.

**realistic reaction** A chemical reaction which, from criteria based on texture and changing *\*mineral* assemblage, can be convincingly demonstrated to have occurred in a *\*rock* during *\*metamorphism*.

**recapitulation of phylogeny** Theory, due to E. Haeckel (1834–1919), asserting that *\*ontogeny* (the development of the individual) recapitulates or reflects the *\*phylogeny* (the evolutionary history of the group). The theory as such has been rejected as not of general applicability, von Baer's '*\*biogenetic law*' being sufficient explanation for the observations on which it was based. However, either *\*hypermorphosis* or retardation of sexual maturity can result in individual cases of recapitulation.

**Recent** See HOLOCENE.

**recessional moraine** See MORAINE.

**recharge** 1. The downward movement of water from the soil to the *\*water-table*. 2. The volume of water added to the total amount of *\*groundwater* in storage in a given period of time.

**recharge area** 1. The geographical area of an *\*aquifer* in which there is a downward movement of water towards the *\*water-table*. 2. The area that acts as a *\*catchment* for any particular aquifer.

**reclined** See STIPE.

**reclined fold**. As defined by M. J. Fleuty (1964), a dipping *\*neutral fold* in which the *\*axial plane* *\*dips* between 10° and 80° and the *\*pitch* of the *\*hinge line* on the axial plane is more than 80°.

**reconstructive transformation** See POLYMORPHIC TRANSFORMATION.

**recovery factor** 1. Measure of extraction efficiency. 2. In mining, the percentage of metal derived from an ore, or *\*coal* from a coal seam, etc. 3. In *\*petroleum geology*, the percentage of the *in situ* oil that is recoverable. Between 20% and 40% is common for primary recovery techniques, but enhanced recovery by use of water injection, detergents to reduce viscosity, etc., may increase the recovery factor to 75%. Recovery factors for *\*natural gas* can be as high as 90%.

**recrystallization** 1. The growth of new *\*mineral* grains from pre-existing mineral grains by the solid-state *\*diffusion* of *\*ions* in response to a change

in temperature, pressure, or composition of the rock system. **2.** The changing of crystal *\*fabric* or crystal size without an accompanying change in mineral chemistry. **3.** See FOSSILIZATION.

**rectangular drainage** See DRAINAGE PATTERN.

**rectilinear slope** That part (or 'segment') of a hillslope profile that is straight. It is usually the steepest part of the profile. Its gradient varies little in an area of uniform rock types, when it stands at the 'characteristic angle'. More generally, it has been seen (by W. *\*Penck*) as that hillslope profile that develops above a river that is down-cutting at a constant rate.

**rectimarginate** Applied to a brachiopod (*\*Brachiopoda*) shell where a planar *\*commissure* is present.

**recumbent fold** A *\*fold* in which the *\*hinge* line and *\*axial plane* are horizontal or subhorizontal. M. J. Fleuty (1964) suggests that the term recumbent fold be restricted to a fold whose axial plane does not *\*dip* more than 10°.

**red algae** See RHODOPHYCEAE.

**red-bed copper** Conformable copper deposits in *\*sandstones* laid down under terrestrial conditions, usually red in colour. The sands are usually porous and copper minerals, normally *\*chalcocite*, develop in the pores. Such deposits are found in many parts of the world, e.g. in the *\*Permian* of the Urals, and in the *\*Triassic* of central England, Nova Scotia, and the south-western USA.

**red beds** *\*Sedimentary rocks*, generally *\*sandstones*, which are red due to their *\*grains* being coated with *\*hematite*.

**red clay (brown clay)** Brown or red, very fine-grained, deep-sea deposit composed of finely divided *\*clay* material that is derived from the land, transported by winds and ocean currents, and deposited far from land in the deepest parts of the ocean basin, especially in mid-latitudes. Red-clay deposits cover about a quarter of the *\*Atlantic* and *\*Indian* ocean floors and almost half the *\*Pacific ocean* floor.

**red copper ore** See CUPRITE.

**red edge** In *\*remote* sensing, the sharp increase in spectral reflectance of wavelengths in the red and very-*\*near* infrared (700–750 nm) part of the spectrum associated with healthy, green-leaved vegetation. *See also* VEGETATION INDEX.

**red iron ore** *See* HEMATITE.

**Redlichiida** An order of *\*Trilobita* that lived in the Lower to Middle *\*Cambrian*. They had large eyes, a large, semicircular *\*cephalon* with strong *\*genal spines*, many small, often spiny, thoracic segments, and a tiny *\*pygidium*. There were four suborders. Some, especially members of the suborder Olenelloidea, are important stratigraphic markers.

**redox potential ( $E_H$ )** A scale of values, measured as electric potential in volts, indicating the ability of a substance or solution to cause *\*reduction* or *\*oxidation* reactions under non-standard conditions. The term is sometimes used interchangeably with the term *\*oxidation potential*, but in either case the symbol  $E^\theta$  would refer to standard conditions, while  $E_H$  signifies non-standard conditions, usually processes in natural systems such as sea water or soils. The higher the value of  $E_H$ , the more oxidizing the conditions. The redox potential is important in *\*weathering* in terms of oxidation and reduction; if the environment will accept *\*electrons* it can precipitate  $Fe(OH)_3$ , if not the  $Fe^{2+}$  *\*ions* will remain in solution. Values in natural environments are closely linked, and vary, with changes in *\*pH*.

**redox reaction (oxidation–reduction)** Reaction involving the transfer of *\*electrons* from a donor molecule, the reducing agent, to an acceptor molecule, the oxidizing agent.

**red podzolic soil** *\*Soil profile* formed at an advanced stage of *\*weathering* and *\*leaching* by the process of *\*podzolization*; it is similar in appearance and properties to a *\*podzol* but associated with the greater degree of *\*chemical weathering* and higher iron-oxide concentrations of a humid, tropical environment. *See also* ULTISOLS.

**Red Sea** An elongate basin, 2000 km long, with shorelines 360 km apart at the widest point and only 28 km apart where it joins the Gulf of Aden. The Red Sea has an inner *\*median valley*, associated with a positive *\*gravity*

**anomaly**, containing **\*basalts** and hot brines, and the sea is thought to be at the young stage of the **\*Wilson** cycle of an ocean.

**red sprite** A **\*transient luminous event** consisting of a weak flash directly above a thunderstorm that coincides with a strong cloud-to-ground lightning stroke. It extends above the cloud as a single or multiple vertical lights to a height of up to about 90 km.

**reduction** Chemical reaction in which atoms or molecules either lose oxygen, or gain hydrogen or **\*electrons**. *Compare* OXIDATION.

**reduction potential** *See* OXIDATION POTENTIAL.

**reduction to pole** The simplification of the interpretation of **\*magnetic** anomalies by modifying the anomaly pattern to that which it would be in a vertical field, i.e. if the locality were at the north (or south) magnetic pole; induced magnetic effects would then be symmetrical. The anomaly is directly analogous to that of a **\*gravity anomaly** (in which the gravitational force is also vertical).

**REE** *See* RARE-EARTH ELEMENT.

**reef 1.** A rigid, wave-resistant build-up constructed by carbonate organisms. Types of reef include patch reefs (small and circular in shape); pinnacle reefs (conical in form); barrier reefs (separated from the coast by a lagoon); fringing reefs (attached to a coast); and atolls (isolated reefs enclosing lagoons). Factors influencing reef growth include: (a) water temperature (optimum 25 °C); (b) water depth (must be less than 10 m); (c) **\*salinity** (normal marine salinity is necessary); (d) wave action (intense wave action favours coral growth); and (e) turbidity (coral growth requires clear water and an absence of terrigenous suspended sediment). The diversity of species found in a reef will be a function of salinity and water temperature, with stressful conditions resulting in a reduction of species present. **2.** In mining, certain palaeoplacer gold deposits in Australia and South Africa.

**reef flat** A pavement of naturally cemented, large skeletal debris and **\*reef** debris to the protected rear (leeward) of a reef crest. Water depth in this zone is shallow, only a few metres at most. Sand shoals and small islands or **\*cays** may be present on the reef flat.

**reef front** An irregularly sloping ramp extending from the \*surf zone to a depth of approximately 100 m on the windward, open-sea side of a \*reef. Abundant skeletal growth grades downward into sediment of the \*fore-reef zone. The robust, branching form of the coral *Acropora palmata*, common in present-day reefs, typifies this reef area.

**reef trap** \*Stratigraphic oil or gas trap produced by porous \*reef \*limestones (\*reservoir rock) covered by impermeable strata. \*Porosity of limestones depends on post-depositional diagenetic changes. A reef trap may also host lead and zinc mineralization in material deposited from migrating \*brines. Compare ANTICLINAL TRAP; FAULT TRAP; STRATIGRAPHIC TRAP; STRUCTURAL TRAP; UNCONFORMITY TRAP.

**re-entrant** An embayment or recess in the flank of a main valley, often of sufficient length to form a tributary.

**reference ellipsoid** An imaginary ellipsoid (a solid body formed when an ellipse is rotated about an axis) that is used when expressing the height of the \*geoid as the perpendicular distance between the geoid and the reference ellipsoid. At one time different countries used their own reference ellipsoids, but nowadays a standard reference ellipsoid is used, with its origin at the Earth's centre of mass.

**reference section** See HYPOSTRATOTYPE.

**refertilization** The replenishment of the upper \*mantle with elements transported upward from the convecting mantle through weak zones and faults.

**reflectance (reflectivity)** 1. The ratio of \*electromagnetic radiation reflected by a surface to that which is incident upon it. 2. In \*ore microscopy, the amount of incident polarized light which is reflected off the surface of an \*opaque mineral. The value of the percentage reflectance ( $R\%$ ) at a specified wavelength is a useful quantitative aid to mineral identification and is determined by:  $R\% = (\text{intensity of reflected light} / \text{intensity of incident light}) \times 100$ .

**reflectance spectrometry** A number of methods for determining the amount of light reflected from a plane surface. In diffuse \*reflectance, this is detected and recorded as a function of wavelength by such instruments as reflectometers, spectroreflectometers, or colorimeters.

**reflected infrared** In *\*remote* sensing, infrared which is solar-generated *\*electromagnetic* radiation that has been reflected from an object. Characteristically, reflected infrared radiation has a wavelength between 0.7  $\mu\text{m}$  and 3  $\mu\text{m}$  and is therefore *\*near-infrared*. Compare THERMAL INFRARED.

**reflected-light microscopy (ore microscopy)** The scientific study of *\*ore* (*\*opaque*) minerals by means of a *\*polarizing* reflected-light microscope. Systematic observations of *\*reflectance* and *\*hardness* are made in order to identify individual *\*minerals*, and the interpretation of textural relationships and *\*paragenetic* studies may reveal the sequence of mineral formation. It is an important technique in the study of metallic mineral deposits.

**reflection** The rebounding of an object or wave (light, heat, sound, seismic, etc.) from a surface; the object or wave so reflected. In geophysics, a signal reflected from a *\*reflector* according to *\*Snell's law*. A seismic reflection occurs as a result of a contrast in *\*acoustic impedances*; an electromagnetic reflection occurs because of a contrast in electrical and *\*dielectrical* properties.

**reflection coefficient** 1. (**R**) The ratio of the amplitude of a reflected ray ( $A_1$ ) to that of the incoming ray ( $A_0$ ), such that  $R = A_1/A_0$ . In the case of a normally incident ray,  $R$  can be expressed in terms of the *\*acoustic* impedances of the two media above and below the *\*reflector*,  $Z_1$  and  $Z_2$ , so that  $R = (Z_2 - Z_1)/(Z_2 + Z_1)$ . The range of values for  $R$  lie between  $-1$  and  $+1$ . If  $R$  is negative a phase reversal ( $\pi$ ) in the wave occurs at the reflector. For water/air  $R$  has a typical value of  $-1$ ; for rocks  $R$  has an average value of  $0.2$  or less. See also TRANSMISSION COEFFICIENT. The reflection coefficient can also be expressed in terms of energy ( $R'$ ), when  $R' = R^2$ . 2. (**k**) A ratio of true resistivities, such that  $k = (\rho_2 - \rho_1)/(\rho_2 + \rho_1)$ , where  $\rho_1$  and  $\rho_2$  are the true resistivities above and below an *\*interface*.

**reflection pleochroism** In *\*ore microscopy*, a change in colour exhibited by a mineral on rotation of the *\*stage* in *\*plane-polarized* light. Some minerals, e.g. *\*covellite*, show a strong variation in colour and this is a useful property in their identification.

**reflectivity** See REFLECTANCE.

**reflector** 1. The surface from which an object or wave (light, heat, sound, seismic, etc.) is reflected. 2. An **\*interface** which gives rise to a contrast in geophysical properties between the media above and below the boundary. See REFLECTION. 3. A component of an **\*ore microscope**, used in **\*reflected-light microscopy**. There are two types. The glass plate reflector is oriented at 45° to direct the horizontal light source vertically on to the polished surface of the mineral specimen and then to allow the reflected light to pass through it vertically up to the observer. The half-field prism (or mirror system) reflects light downwards through one-half of the aperture of the **\*objective** or lens; the light is then reflected back from the mineral specimen through the other half of the objective, passing behind the prism to the observer.

**reflexed** See STIPE.

**reflux theory** A proposed mechanism for the continued movement of dense, hypersaline water through a **\*sediment**, causing mineralization or alteration. Reflux is thought to occur by evaporation of salt water producing dense, saline water which sinks through the sediment and displaces lighter waters of normal **\*salinity**. In such a way high concentrations of minerals can be flushed through sediment, so causing effects such as large-scale **\*dolomitization** of buried **\*limestones**.

**refolded fold** A **\*fold** which, subsequent to the original folding, has undergone one or more further episodes of folding to produce a complex structure in which the folds, or successive folding episodes, are commonly designated chronologically as  $F_1, F_2, F_3, \dots, F_n$ . The **\*outcrop** manifestations of refolding are **\*interference patterns**.

**refraction** The bending of a ray which travels obliquely from one medium to another, at the **\*interface** separating the two; it is caused by the contrast in velocities with which the ray travels in the two media, and described by **\*Snell's law**. See also REFRACTION SURVEY.

**refraction survey** A field investigation in which seismic **\*head** waves are used to study subsurface geologic structures. Seismic waves travel down from a source to an **\*interface**, where they are critically refracted along the boundary and reradiated back to the surface, and detected by a **\*geophone \*array**. The **\*travel times** of the **\*first breaks** are plotted on a **\*travel-time**

graph, from which depths to the refractor, its dip, and the velocities of the layers encountered can be calculated. See [SNELL'S LAW](#). See also [CROSS-OVER DISTANCE](#); [INTERCEPT TIME](#).

**refractive index (*n*)** When light travels from air into a substance its velocity is reduced. The light path is also refracted into the substance, and the relationship between the angle of incidence (*i*) and the angle of refraction (*r*) is a constant (*\*Snell's law*). This constant (*n*) is the refractive index of the mineral and is determined by:  $\sin i / \sin r = n$ . The refractive index is also the ratio of the velocity of light in air (*V*) to the velocity of light in the mineral (*v*):  $n = V/v$ .

**refractometer** An instrument used to determine *\*refractive index*. Several types are available. (a) The Herbert Smith refractometer measures the *\*critical angles* by total *\*internal reflection*, using a glass hemisphere through which light is directed upwards and reflected from the mineral surface. The reflected beam produces an image in the observing telescope, where a graduated scale measures the critical angle. The refractive index is then calculated by *\*Snell's law*. (b) The Abbé refractometer is used primarily for determining the refractive index of liquids. It consists of a pair of glass prisms with a film of liquid between them. The line of the critical angle is measured through a fixed telescope and the refractive index is read off a graduated scale calibrated against a glass plate of known refractive index. (c) The Leitz–Jelley refractometer is used to determine the refractive index of small amounts of liquid. It consists of a glass prism cemented to a glass slide which can hold the liquid. A beam of light is directed at right angles to the glass and is refracted by the liquid on to a graduated scale from which the refractive index can be read.

**refractory mineral** Mineral resistant to decomposition by heat, pressure, or chemical attack. Most commonly applied to heat resistance.

**refugia** Small isolated areas where extensive changes, most typically due to changing climate, have not occurred. Plants and animals formerly characteristic of the region in general now find a refuge from the new unfavourable conditions in these areas. An example might be a mountain summit projecting above a glaciated lowland region. See also [RELICT](#).

**Refugian** A *\*stage* (35–33.5 Ma ago) in the *\*Palaeogene* of California, underlain by the *\*Narizian*, overlain by the *\*Zemorrian*, and roughly contemporaneous with the *\*Priabonian* stage.

**reg 1.** Stony desert. **2.** Gravel veneer, normally consisting of small, rounded pebbles, that mantles a Saharan plain and has a gradient as low as 1:5000. The pebble layer may be underlain by a stony soil, or it may be a *\*lag deposit*. Compare *SERIR*.

**regelation** A process by which water that has been released by *\*pressure melting* beneath a temperate *\*glacier* is refrozen (*see also* *BASAL SLIDING*; *ICE*). The process takes place in a relatively thin zone, the ‘regelation layer’, which may be only a few centimetres thick. Regelation is associated with the incorporation of bedrock materials in the debris-rich ‘sole’ of the glacier, which is restricted to the thickness of the regelation layer.

**regio** (*pl. regionis, regiones*) A term applied to any feature on a planetary surface that is not clearly defined or understood, usually because of insufficient resolution. Examples include dark regions on *\*Ganymede* and *\*Iapetus*. On *\*Venus*, the term was originally used for radar-bright features such as Beta Regio, which is probably a volcanic construct. Its usage is now extended to cover elevated terrain smaller than continents.

**regional field** The values in gravity and magnetic surveys that can be attributed to sources within the lower *\*crust* or *\*basement*. These are usually of much longer wavelengths than those associated with near-surface bodies. *See* *RESIDUAL GRAVITY MAP*.

**Regionally Important Geological/Geomorphological Sites (RIGS)** A British network of sites selected and conserved by informally constituted groups of volunteers working closely with statutory and voluntary conservation bodies. The scheme began in 1990.

**regional metamorphism** The *\*recrystallization* of pre-existing rocks in response to simultaneous changes of temperature, lithostatic pressure, and in many cases *\*shear stress*, occurring in *\*orogenic belts* where lithospheric *\*plates* are converging. The broad areas covered by orogenic belts cause the associated *\*metamorphism* to be developed on a regional scale, hence the name attached to this type of metamorphism. Regional metamorphism can be pre-, syn-, or post-tectonic, depending whether the metamorphic event

(or events) is (or are) before, synchronous with, or after the orogenic deformation event (or events). Typical rock *\*fabrics* produced during regional metamorphism are, in order of increasing grain size (reflecting increasing *\*metamorphic grade*), slaty, phyllitic, schistose, and gneissose fabrics. Increase of metamorphic grade in regional *\*terrains* typically produces a *\*prehnite–pumpellyite–\*greenschist–\*amphibolite–\*granulite* facies series. However, each regional metamorphic terrain is characterized by a unique mineral zonal sequence reflecting a particular pressure–temperature gradient during metamorphism.

**regional stratigraphic scale** See STRATIGRAPHIC SCALE.


**regolith 1.** General term for the layer of *\*unconsolidated* (non-cemented), weathered material, including rock fragments, mineral grains, and all other superficial deposits, that rests on unaltered, solid bedrock. It reaches its maximum development in the humid tropics, where depths of several hundreds of metres of weathered rock are found. Its lower limit is the *\*weathering front*. Soil is regolith that often contains organic material and is able to support rooted plants. Compare SAPROLITE. **2.** The continuous layer of incoherent fragmental material, produced by *\*meteorite* impact, that typically forms the surface blanket on planets, *\*satellites*, and *\*asteroids* where the atmosphere is thin or lacking. The classic example is the lunar regolith, typically several metres thick, with components ranging from metre-sized blocks to micron-sized dust and glass particles.

**regosols** A reference soil group in the *\*World Reference Base for Soil Resources* classification. Regosols are all soils not included in another group.

**regression (marine)** The withdrawal of water from parts of the land surface due to a fall in sea level relative to the land. Shallow-water *\*sediments* overlie sediments characteristic of deeper water. See OFFLAP. Compare TRANSGRESSION.

**regressive systems tract (RST; falling stage systems tract)** In the *\*genetic stratigraphic* sequence model used in *\*sequence stratigraphy*, a sigmoid *\*clinoform* produced by *\*onlap* under conditions of rising sea level due to the rapid subsidence of the basin. It is a type of *\*lowstand systems tract* produced under special conditions. If the rate of subsidence was at no

time high enough to outpace the rate of sedimentation, thus allowing transgression, the tract is known as a midstand (or forced regressive) systems tract.



#### Regressive systems tract

**regular echinoids** Informal term for sea-urchins, of the class *\*Echinoidea*, in which the anus is enclosed within the apical system. The term includes the Perischoechinoidea, Diadematacea, and Echinacea.

**Regulares (Monocyathea)** (phylum *\*Archaeocyatha*) Class of animals that were usually solitary, only rarely colonial, found in Lower and Middle *\*Cambrian* deposits. The conical cup varies from cylindrical to saucer-shaped, and usually consists of two porous walls (a single porous wall in the order Monocyathida). The *\*intervallum* may contain *\*tabulae* alone or with *\*septa*. There is a fan-like divergence of longitudinal pore-rows in the septa. *\*Dissepiments* may be present. Some single-walled monocyathids had a flap (pelta) over the central cavity. *Compare* IRREGULARES.

**Reid, Harry Fielding** (1849–1944) An American geophysicist, Reid proposed the '*\*elastic rebound*' theory of earthquake motion after studying the 1906 San Francisco earthquake. He was a vigorous opponent of

**\*continental drift** theory, describing **\*Wegener's** work as 'pseudo-scientific'.

**REIMI** A **\*microsatellite** mission, known as INDEX (Innovative Technology Demonstration Experiment) and renamed after launch, by the **\*Japan Aerospace Exploration Agency** (JAXA) and Institute of Space and Astronomical Science (ISAS) to observe the structure of the **\*aurora**. *Reimi* means dawn. The microsatellite was launched on 23 August 2005, from Baikonur, Kazakhstan.

**rejuvenation** The marked increase in the rate of **\*erosion** that takes place when a land mass is relatively elevated. Streams respond by incision, with the development of **\*terraces** and **\*knick** points, and finally a **\*polycyclic** landscape emerges.

**relative age** The position within a time sequence (in the Earth sciences usually the **\*stratigraphic timescale**) held by an event, **\*fossil**, **\*mineral**, or **\*rock**, compared with others of its kind, e.g. 'an early **\*Cambrian \*trilobite**' or a 'Late **\*Jurassic** marine **\*transgression**'. No age in years is implied. *Compare* ABSOLUTE AGE. *See also* DATING METHODS.

**relative humidity** The water-vapour content of air at a given temperature, expressed as a percentage of the water-vapour content that would be required for saturation at that temperature. Generally the relative humidity decreases during the day, with increase in temperature, and increases at night as the temperature falls.

**relative permittivity** *See* DIELECTRIC CONSTANT.

**relative plate motion** The motion of one lithospheric **\*plate** relative to another. This can be described by the **\*pole of rotation** and the angular velocity about this pole.

**relative pollen frequency (RPF)** Expression of **\*pollen** data from sediments for each species, genus, or family, as a percentage of the total pollen count, or the total tree pollen. It is the traditional and most widely used method for preparing **\*pollen diagrams**. *Compare* ABSOLUTE POLLEN FREQUENCY. *See also* POLLEN ANALYSIS.

**relative timescale** *See* DATING METHODS; GEOCHRONOLOGY.

**relative vorticity** See VORTICITY.

**relaxation** A term used to describe the 'fading', or loss of topographic relief, of craters on icy **\*satellites**. Such craters are generally shallower than those on rocky satellites, due to viscous flow of the icy crust. Some disappear completely, leaving a discoloured patch or **\*palimpsest** on the surface.

**relaxation frequency ( $f_r$ )** The frequency at which the **\*dielectric loss factor** ( $\epsilon''$ ) reaches a maximum, for a dielectric material that has no static (d.c.) conductivity and that is subjected to an alternating electromagnetic field.

**relaxation time** **1.** The time taken by a disturbed system to reach equilibrium, or the time taken for the magnitude of some parameter to decrease to about 37% of its initial value. For example, the temperature-dependent relaxation time ( $\tau$ ) of a **\*dielectric** is related to the **\*relaxation frequency** ( $f_r$ ) such that  $\tau = 1/(2\pi f_r)$ . Physically, it is the time taken for an ionic defect to move within a **\*crystal \*lattice** under the influence of an applied alternating electromagnetic field. **2.** In **\*geomorphology**, the time taken for a **\*system** to become adjusted to a sustained change in the nature and/or intensity of external **\*processes**. Such an adjustment normally involves a change in the shape of the land-form or landscape constituting the system. Relaxation times vary. The width of a river channel may adjust in response to an increase in discharge in, perhaps, 10 years, while a glaciated mountain range may require  $10^5$ – $10^6$  years to lose the imprint of ice.

**releasing bend** A bend in an offset **\*strike–slip** fault (i.e. the fault is not straight) that allows the sides of the fault to separate, sometimes to form a pull-apart basin or **\*graben**. The releasing bend accommodates local extension.

**relict** Applied to organisms that have survived while other related ones have become extinct. Often the term refers to **\*species** that have survived periods of unfavourable conditions (e.g. **\*glacial** periods or land submergence) by existing in regions called **\*refugia**, while becoming extinct elsewhere (e.g. some arctic-alpine plants). It may also refer to a surviving species of a group, the other species of which have become extinct (e.g. **\*coelacanth** fish). See also RELICT SEDIMENT.

**relict land-form** A land-form that developed under climatic conditions very different from those obtaining today.

**relict sediment** \*Sediments of the \*continental shelf deposited by processes no longer active in the area where the sediments now occur. Relict sediments are remnants from an earlier environment and are now in disequilibrium. Approximately 50% of the present continental shelves are covered by relict sediments deposited during the period of lower sea levels in the \*Pleistocene.

**relict structure** A textural or structural feature inherited from an original \*igneous or \*sedimentary rock and preserved as a \*relict in a low-grade \*metamorphic rock which has suffered little or no deformation.

**relief** In \*thin-section microscopy, variations due to the difference in \*refractive index between a \*mineral and its mounting medium. If the differences are small the mineral appears flat and featureless, with faint outlines. If the differences are large the mineral appears to stand out, with strongly marked outlines and conspicuous \*cleavages or fractures. The nature of the relief is determined by the \*Becke line test.

**Relizian** A \*stage (16.5–15.5 Ma ago) in the \*Miocene of California, underlain by the \*Saucasian, overlain by the \*Luisian, and roughly contemporaneous with the Lower \*Langhian stage.

**remanent magnetization** The magnetization remaining after the removal of an externally applied field, and exhibited by \*ferromagnetic materials. See NATURAL MAGNETIZATION; ISOTHERMAL MAGNETIZATION; GYROMAGNETIZATION; and ANHYSTERETIC MAGNETIZATION.

**remanié** See DERIVED.

**remanié beds** See CONDENSED BED.

**remnant arc** A member of a complex type of \*back-arc basin in which a number of remnant arcs are separated from the \*magmatic arc, and from each other, by \*inter-arc basins.

**remote sensing** The gathering of information without actual physical contact with what is being observed. This involves the use of \*radars, sonars, spectroscopy, and the use of airborne and \*satellite photography.

See BISTATIC RADAR; IMAGING; LASER RANGING; POLARIMETRY; RADAR ALTIMETRY; RADIOMETRY; RADIO OCCULTATION.

**remoulding** The change that occurs in *\*clay* which has been disturbed and lost *\*shearing strength* but gained *\*compressibility*.

**removal time** See RESIDENCE TIME.

**rendzina** A *\*brown earth* soil of humid or semi-arid grassland that has developed over calcareous *\*parent material*. Rendzinas may fall within the orders *\*inceptisols* or *\*mollisols*.

**renewable resource** Resource produced as part of the functioning of natural systems at rates comparable with its rate of consumption, e.g. food production by *\*photosynthesis*. Limits to renewable resources are determined by flow rate and such resources can provide a sustained yield. Compare NON-RENEWABLE RESOURCE.

**reniform** Kidney-shaped.

**repeated twinning** See LAMELLAR.

**Repettian** A *\*stage* (2.9–2.2 Ma ago) in the *\*Pliocene* of California, underlain by the *\*Delmontian*, overlain by the *\*Venturian*, and roughly contemporaneous with part of the *\*Piacenzian* stage.

**repichnia** A behavioural category of *\*trace fossils* that result from locomotion. Animals may leave distinct *\*tracks* through walking or crawling across soft *\*sediment* surfaces; repichnia are the fossilized traces of those tracks.

**replacement** Widely used geologic term denoting a process that involves some kind of transformation. In a petrological sense it refers to the partial or complete alteration of an original mineral to an aggregate of *\*secondary minerals*, by the diffusion of *\*ions* between the solid *\*phases* and an introduced, fluid (usually water-rich) phase. Such diffusion takes place easily when the temperature of the rock system is below the stability limit of an individual mineral and a fluid is present to act as a catalyst to initiate the diffusion reactions. The secondary minerals may be all of one type, or a combination of mineral types. For example, high-temperature, magnesium-rich *\*olivine* can be replaced by an aggregate of secondary *\*serpentine* and

\*chlorite, while \*plagioclase can be replaced by a fine aggregate of white mica (\*sericite). See also FOSSILIZATION.

**reptation** A mode of particle transport in which grains are lifted or ejected only weakly and do not rebound or eject other particles when they return to the bed.

**Reptilia (reptiles)** Large and varied class of \*poikilothermic \*vertebrates, which arose in the \*Carboniferous from \*labyrinthodont amphibians. They were the dominant animals of the \*Mesozoic world and gave rise to the \*birds and \*mammals. Reptiles have a body covering of ectodermal scales, sometimes supported by bony scutes. There is no gilled larval phase; development is by \*amniote egg, but ovovivipary is common. Reptiles are air-breathing from hatching onwards.

**reptiles** See REPTILIA.

**Reptilomorpha** A superorder of reptile-like amphibians that lived during the \*Carboniferous. It is thought to be one of the supporting stems of the \*crown group \*Tetrapoda.

**resequent** Applied to a land-form whose orientation is similar to that of the inferred original feature, but which has passed through a complex subsequent history. For example, a resequent \*fault-line scarp faces the same way as the original fault scarp.

**resequent fault-line scarp** See RESEQUENT.

**reserve** Resources of coal, ore, or minerals which can be mined legally and profitably under existing conditions. The indicated reserve is the estimate of ore computed from \*boreholes, \*outcrops, and developmental data, and projected for a reasonable distance on geologic evidence. An inferred reserve is an estimate based on relationships, character of deposit, and past experience, without actual measurements or samples; it should include the limits between which the deposit may lie. A potential reserve is ore not yet discovered but whose presence is suspected; the term is sometimes used for ore not commercially viable at present time. A proved reserve is a resource reliably established by tunnels, boreholes, or mining.

**reservoir** 1. A surface body of water whose flow is artificially controlled by means of dams, embankments, or sluice gates in such a way that the

water remains static until it is allowed to flow for a specific purpose, e.g. flood control or public water supply. **2.** An underground rock formation with sufficient void space to act as a store for water, natural gas, or oil.

**reservoir pool** Large and usually *\*abiotic* store of a nutrient in a *\*biogeochemical cycle*. Exchanges between the reservoir pool and the *\*active pool* are typically slow by comparison with exchange within the active pool. Human activity, such as the mining of mineral resources, may profoundly alter this exchange rate, generally releasing an excess into the active pool which can be accommodated only by establishing a new equilibrium. This may in turn produce unfavourable conditions, manifested as chemical pollution, e.g. excess phosphorus in eutrophication, excess sulphur in acid rainfall, and lake acidification.

**reservoir rock** Any porous rock in which oil, gas, or water may accumulate; usually *\*sandstone*, *\*limestone*, or *\*dolomite*, but sometimes fractured *\*igneous* or *\*metamorphic rock*.

**reshabar** Regional south-easterly *\*katabatic* wind affecting mountain slopes in southern Kurdistan (the plateau and mountains in south-eastern Turkey, northern Iraq, northern Syria, and western Iran). The strong, swirling wind is hot and dry in summer but brings cold conditions in winter.

**residence time** **1. (removal time)** The time that a given substance remains in a particular compartment of a *\*biogeochemical cycle*. **2.** The time during which water remains within an *\*aquifer*, lake, river, or other water body before continuing around the *\*hydrological cycle*. The time involved may vary from days for shallow gravel aquifers to millions of years for deep aquifers with very low values for *\*hydraulic* conductivity. Residence times of water in rivers are a few days, while in large lakes residence time ranges up to several decades. Residence times of continental *\*ice* sheets is hundreds of thousands of years, of small *\*glaciers* a few decades. **3.** The average time a particular element of sea water spends in solution between the time it first enters and the time it is removed from the ocean. **4.** The average time that a water molecule or particulate pollutant spends in the atmosphere. For pollutants (e.g. dust from a volcanic *\*eruption*), the residence time may range from a few weeks in the lower *\*troposphere* to several years in the upper *\*stratosphere*, before it is scavenged out by

**\*precipitation.** For water molecules the overall average is believed to be 9–10 days.

**residual deposit** 1. Weathered material remaining *in situ* after soluble constituents have been removed. 2. **\*Ore** deposit in **\*clay** formed by near-surface oxidation, e.g. **\*bauxites** (aluminium ore), residual nickel, extensive iron **\*laterites**, and **\*soil**.

**residual gravity map** Usually a map of the Earth's **\*gravitational acceleration** remaining after allowing for all distorting effects. It is known as a **\*Bouguer anomaly** map when a **\*regional field** (usually a gradient), attributable to gravitational sources within the lower **\*crust** or **\*basement**, has been removed. See SMITH'S RULE.

**residual shear strength** See SHEAR STRENGTH.

**resinite** See COAL MACERAL.

**resinous** Of a mineral **\*lustre**, translucent yellowish to brown.

**resistate mineral** A mineral which is not readily weathered by chemical attack, e.g. **\*quartz**, **\*zircon**, and **\*muscovite**. The relative ability of minerals to resist **\*chemical weathering** is expressed in the Goldich stability series.

**resistivity logging** See LATEROLOG SONDE.

**resistivity methods** Geophysical methods in which very-low-frequency or direct electrical current is injected into the ground and its potential distribution is measured in order to obtain information about the Earth's resistivity. Loosely, the term may also include **\*electromagnetic methods**, since **\*apparent conductivities** ( $\sigma_a$ ) can be used to derive **\*apparent resistivities** ( $\rho_a$ ) by:  $\sigma_a = 1/\rho_a$ . See CONSTANT-SEPARATION TRAVERSING; ELECTRICAL SOUNDING; ELECTRODE CONFIGURATION; INDUCED POLARIZATION.

**resonance** 1. Condition of very large wave amplitude, occurring when the frequency of an external wave-generating force matches and amplifies a natural frequency for waves moving to and fro in an enclosed space such as an **\*estuary**. 2. The relationship in which the orbital period of one body is related to that of a second by a simple integer fraction (e.g.  $\frac{1}{2}$ ,  $\frac{3}{5}$ ). Such

orbits are common in the **\*solar system**. Well-known examples include the Kirkwood Gaps in the Asteroid Belt and the Cassini Division in Saturn's ring system (a particle moving in the Cassini Division has a period  $\frac{1}{2}$  that of Mimas, and  $\frac{1}{3}$  that of Enceladus).

**resorption** The partial fusion of a **\*euhedral \*phenocryst** in a **\*magma** in response to a change in magma temperature, pressure, and/or composition. If the magma is erupted rapidly the partially fused phenocryst can be preserved as a large, **\*anhedral \*crystal** with a lobate outline set in a fine-grained **\*groundmass**.

**ResourceSat-2 and -2A** A data continuity mission of the India Space Research Organization carrying three electrooptical cameras that supply images relevant to a variety of applications, including agricultural crops, crop yields, water resources, forest mapping, and disaster management. ResourceSat-2 was launched on 20 April 2011, from Satish Dhawan Space Centre, into a Sun-synchronous orbit at an altitude of 817 km. A follow-on mission, ResourceSat-2A, was launched on 7 December 2016, from the same location, into a similar orbit.

**restraining bend** A bend in an offset **\*strike-slip fault** (i.e. the fault is not straight) where the two sides of the fault converge to produce a **\*horst**. The restraining bend accommodates local contraction.

**resurgence** 1. See **SPRING**. 2. See **CALDERA**.

**resurgent caldera** See **CALDERA**.

**Resurs-DK1** The first Russian civil Earth-observation satellite producing high-resolution (1 m) images. It was launched on 15 June 2006, from Baikonur, Kazakhstan, into an elliptical semi-polar orbit at an altitude of 360–610 km. As its imaging showed signs of deteriorating, its place was taken by Resurs-P, launched on 25 June 2013, from Baikonur, into a Sun-synchronous near-circular orbit at an altitude of 475 km.

**reticula** See **RETICULUM**.

**reticulated** Applied to a meshwork of intercalating crystals which may give rise to lattice-like groups of crystals in a mineral.

**reticulite** A gold-brown, foam-like type of glass produced by **\*Peléean eruptions** and found near Hawaiian volcanoes, often in considerable

quantities.

**reticulum** (*pl.* **reticula**) A netlike (i.e. reticular) feature on the surface of **\*Venus**.

**retro-arc basin** A type of **\*back-arc** basin which is floored by **\*continental crust**. The main **\*sediments** are fluvial, deltaic, or marine, derived from the uplifted area behind the arc.

**retro-arc foreland basin** A **\*foreland basin** that forms on the overriding **\*plate** at a **\*convergent margin** where one plate is being subducted.

**retrochoanitic** See SEPTUM.

**retrograde** Used in a planetary context to denote a body moving in the opposite sense to that of most **\*solar-system** bodies, i.e. clockwise rather than anticlockwise. The classic example is the retrograde rotation of **\*Venus**. **\*Triton** is in retrograde orbit around **\*Neptune**. At least four of the outer **\*satellites** of **\*Jupiter**, and one of **\*Saturn**, are also in retrograde orbits.

**retrograde metamorphism** (**diaphthoresis**, **retrogressive metamorphism**) The **\*recrystallization** of pre-existing rocks in response to a lowering of **\*metamorphic grade** in the presence of a fluid **\*phase**. After reaching a metamorphic climax, lowering of metamorphic grade does not usually cause retrograde reactions to occur because all the water in the rock system has been expelled at the metamorphic climax, thus preserving high-grade **\*mineral** assemblages. If some water remains in the system, however, or is introduced as the grade decreases, the water can act as a catalyst to initiate retrograde reactions. The reactions produce hydrated mineral types (see **HYDRATION**), in contrast to the dehydration reactions of **\*prograde metamorphism**.

**retrogression** A progressive failure of **\*quick clay** during a quick-clay landslide (called a flow slide). A flow slide often begins with a small slump of clay from the bottom of an embankment that has been eroded or undercut. The removal of this material places a **\*shear stress** on the material above it on the slope. This material also slumps and a series of failures advances up the slope, sometimes at a rate of several metres per second.

**retrogressive metamorphism** See **RETROGRADE METAMORPHISM**.

**retrosiphonate** A condition in some cephalopods (*\*Cephalopoda*) in which the septal (see *SEPTUM*) necks point back, towards the *\*protoconch*.

**retrusive** Applied to the direction of *\*spreiten* that are extended upwards through the *\*sediment* and are therefore proximal to the point of entry.

**return flow** See *INTERFLOW*.

**return period** The frequency, based on statistical analysis of past records, with which a particular environmental hazard may be expected.

**Réunion** Two normal *\*polarity subchrons* which occur within the *\*Matuyama* reversed *\*chron*.

**reverberation (ringing)** An oscillatory effect seen on seismic wave-forms and produced by short-path *\*multiples*.

**reversal** **1.** A change of direction, usually by 180°. It commonly refers to a change of *\*polarity* of the *\*geomagnetic field*. **2.** A form of *\*homoplasy*; resemblance between two taxa because one of them has gained a new character, then lost it again, and the other taxa has never gained it. Reversal by character loss is common; there is much doubt about whether it ever occurs by regaining a lost character.

**reversal timescale** See *MAGNETOSTRATIGRAPHIC TIMESCALE*.

**reversed field** See *GEOMAGNETIC FIELD*; *POLARITY REVERSAL*. Compare *NORMAL FIELD*.

**reversed kink band** See *CONTRACTIONAL KINK BAND*.

**reverse fault** A low-angle, *\*dip-slip fault* in which the relative displacement of the *\*hanging wall* is upwards. A *\*thrust* is a type of reverse fault.

**reverse zoning** See *CRYSTAL ZONING*.

**reversing dune** A *\*seif dune* that has asymmetrical ridges.

**revolving storm** See *TROPICAL CYCLONE*.

**reworked** See *DERIVED*.

**Reykjanes Ridge** The part of the **\*Mid-Atlantic Ridge** to the south-west of Iceland, whose axis, marked by a **\*median valley**, continues into the active **\*graben** across Iceland.

**Reynolds number** A dimensionless number expressing the balance of viscous and interstitial forces on a small element of moving fluid. The transition from **\*laminar** to **\*turbulent flow** depends on the Reynolds number ( $R$ ) which is equal to  $\rho v d / \eta$ , where  $\rho$  is the fluid density,  $v$  the fluid velocity,  $d$  the diameter of the **\*pore space** through which flow occurs, and  $\eta$  the **\*viscosity**. For laminar flow, the Reynolds number is less than 500, while turbulent flow occurs when  $R$  is greater than 1000. **\*Darcy's Law** for **\*groundwater flow** is valid for values of  $R$  less than about 1–10.

**rhabdosome** In **\*Graptolithina**, a complete colony.

**Rhaetian** A Late **\*Triassic stage**, preceded by the **\*Norian**, followed by the **\*Hettangian** (**\*Lias**) and dated at 203.6–199.6 Ma ago (Int. Commission on Stratigraphy, 2004). It is roughly contemporaneous with the Erchiao (China) and **\*Warepan** (New Zealand). Some authors have placed the Rhaetian in the **\*Jurassic**. Others have questioned its status as a stage, suggesting that it should be a biostratigraphic **\*zone** within the underlying Norian.

**Rhea (Saturn V)** One of the major satellites of **\*Saturn**, with a radius of 764 km; mass  $23.1 \times 10^{20}$  kg; mean density  $1240 \text{ kg/m}^3$ ; visual albedo 0.7. It was discovered in 1672 by G. D. Cassini.

**rheid flow** Flow that occurs in a solid material that is subjected to stress. See **RHEIDITY**.

**rheidity** The length of time that it takes for a solid material to flow when subjected to pressure. The rheidity of ice is measured in weeks and that of **\*igneous** rocks close to the surface in tens of thousands of years. See **RHEID FLOW**.

**rhenium–osmium dating** A method of radiometric dating based on the decay of rhenium-187 ( $^{187}\text{Re}$ ) to osmium-187 ( $^{187}\text{Os}$ ), with a half-life of  $4.16 \times 10^{10}$  years. Since both rhenium and osmium are strongly siderophile and chalcophile, they are useful in dating sulphide ores.

[http://www.geochronometria.pl/pdf/geo\\_27/Geo27\\_05.pdf](http://www.geochronometria.pl/pdf/geo_27/Geo27_05.pdf)


- A review of the physical foundations of the rhenium–osmium method.

**rheology** (*adj.* **rheological**) **1.** Study of deformation and flow in materials, including their elasticity, **\*viscosity**, and plasticity. **2.** In geology, the study of flow in water, **\*ice**, **\*magma**, and during rock deformation.

**Rhine graben** The **\*rift valley** which contains the river Rhine and which lies between the Ardennes, the Vosges, and the Black Forest. The uplift occurred in the late **\*Mesozoic**, with rifting in the mid-**\*Eocene** and production of **\*alkaline \*magmas** in the **\*Oligocene**. In places, 3 km of **\*sediment** have been deposited. The Rhine graben formed synchronously with the Alpine collision and has been called an ‘impactogen’, i.e. a collisional rift that forms at the end of the **\*Wilson Cycle**.

**Rhipidistia** Group of **\*crossopterygian** (‘tassel-finned’) fish ranging from the **\*Devonian** to the **\*Permian**. They possessed two dorsal fins, lobate or stalked pectoral and pelvic fins, and internal nostrils. Distantly related to the living **\*coelacanth**, they are considered by some to be ancestral to the **\*Tetrapoda** (terrestrial **\*vertebrates**).

**Rhizocorallium** U-shaped feeding structures excavated during the search for food, and abundant in the **\*Jurassic**, *Rhizocorallium* is typically elongate and is found parallel or slightly oblique to the bedding surface. It may attain a length of 1 m and the parallel tubes usually have a diameter of 2–3 cm. Delicate **\*spreiten** may reveal the direction of excavation. See **FODINICHNIA**.



**Rhizocorallium**

**rhodochrosite** Mineral,  $\text{MnCO}_3$ ; sp. gr. 3.4–3.7; \*hardness 3.5–4.5; \*trigonal; translucent rose-pink, and sometimes light grey to brown, developing a brown or black crust on exposure; white \*streak; vitreous \*lustre; crystals rare, but form as \*rhombohedra, and rod-like and curved, but it is usually \*massive or granular; \*cleavage perfect rhombohedral; occurs in \*hydrothermal veins containing silver, lead, and copper, and in \*metamorphic and \*metasomatically altered rocks of sedimentary origin, often a \*secondary mineral after manganese oxide; soluble, with effervescence, in hot, dilute hydrochloric acid. It is used in the manufacture of ferromanganese, being added to blast-furnace charges, and in the chemical industry.

**rhodonite** Member of the \*inosilicates with composition  $(\text{Mn,Ca})\text{Si}_2\text{O}_6$  and related to the \*pyroxenes but with a different atomic structure, and termed a pyroxenoid (*see also* WOLLASTONITE); sp. gr. 3.57–3.76; \*hardness 5.5–6.0; pink to brownish-red; \*tabular or \*massive; occurs in manganese-rich mineral deposits, e.g. in Franklin, New Jersey.

**Rhodophyceae (red algae)** A class of marine \*algae, most of them red in colour, whose basic shape is filamentous or membranaceous. They tend to occur at greater depths than the green algae (\*Chlorophyta) and they are among the oldest groups of \*eukaryotic algae, known from the \*Cambrian upwards. *Epiphyton* (Cambrian to \*Devonian) formed mounds and *Solenopora* (Lower Cambrian to \*Cretaceous) formed nodular masses made up of close-packed tubes. The coralline red algae, e.g. *Lithothamnion*, are important rock builders, constructing rigid structures and contributing \*lime mud to \*sediments.

**rhombododecahedron (rhombic dodecahedron)** A 12-faced \*crystal form (110) which has \*cubic \*crystal symmetry. Each face is rhombus-shaped and intersects two \*crystallographic axes equally and is parallel to the third.

**rhombic dodecahedron** *See* RHOMBODODECAHEDRON.

**rhombochasm** A chasm, rhomboid in shape, that forms deep in the crust through transverse faulting of two blocks.

**rhombohedral** See TRIGONAL.

**rhombohedron** A six-faced, \*closed \*crystal form which belongs to the \*trigonal system. There are three upper faces and three lower faces arranged about the vertical *c* (or *z*) axis. The three horizontal axes emerge in the middle of each edge. The mineral \*calcite frequently occurs in this crystal form.

**rhourd** A large, star-shaped or pyramidal sand \*dune (a ‘sand mountain’) that may be 100–200 m high and that has been described for the Algerian Sahara. It may form where two zones of sand-laden wind cross one another. See DRAA.


**Rhuddanian** A \*stage (443.7–439 Ma ago) of the Early \*Silurian, underlain by the \*Hirnantian (\*Ordovician) and overlain by the \*Aeronian.

**Rhyacian** A \*period of the \*Proterozoic \*eon that began 2300 Ma ago and ended 2050 Ma ago. Both its commencement and end are defined by fixed dates. The Rhyacian followed the \*Siderian period and was followed by the \*Orosirian period.

**Rhynchocephalia** (‘beak-heads’, rhynchocephalians) (class \*Reptilia) Order of primitive, lizard-like reptiles dating from the \*Triassic and often cited as a \*living fossil. The order contains only one species, *Sphenodon punctatus*, the tuatara of New Zealand. This survives only in the Bay of Plenty, and it has protected status. The skull is of the primitive \*diapsid type, with a fixed quadrate bone. The teeth are fused to the edge of the jaw (acrodont), with a tendency to develop a beak-like structure anteriorly.

**rhyncholite** One of the beak-like structures which are considered to be the upper jaw structures of \*fossil cephalopods (\*Cephalopoda). They are approximately rhomboidal in shape, with a slightly concave lower surface. The anterior portion is termed the hood, the posterior portion the shaft. The first examples are found in the \*Carboniferous.

**Rhynchonellida** (rhynchonellids) (class \*Articulata) Order of brachiopods (\*Brachiopoda), with rostrate shells, a functional \*pedicle, and a \*delthyrium partly restricted by a pair of \*deltidial plates. The shell is usually \*impunctate. They appeared first in the Middle \*Ordovician. The order contains about 250 genera, most of which are extinct.



### Rhynchonellida

**rhynchonellids** See RHYNCHONELLIDA.

**Rhynia** An early vascular plant, first known from the Lower \*Devonian Rhynie Chert, Aberdeenshire, Scotland, and described by R. Kidston and W. H. Lang in a series of papers between 1917 and 1921. *Rhynia* was a simple, leafless plant with a creeping, horizontal stem (rhizome) from which the upright, aerial shoots arose. The tips of fertile shoots bore oval-shaped sporangia (see SPORE) and the prostrate, horizontal axis was supported by rhizoids rather than true roots. Originally two species were included in the genus: *R. gwynne-vaughanii* (up to 20 cm tall) and *R. major* (20–50 cm). Recent work by David S. Edwards (1986) has shown *R. major* to vary in its branching pattern from *R. gwynne-vaughanii* and to lack the tracheids necessary for it to qualify as a vascular plant. Consequently *R. major* has been transferred to a new genus of uncertain affinity and is now termed *Aglaophyton major*. See also COOKSONIA; PSILOPHYTALES.

**Rhyniophytina** See PSILOPHYTALES.

**rhyodacite (toscanite)** A fine-grained, \*extrusive, \*igneous rock characterized by an \*adamellite \*mineral assemblage and composition. Most rhyodacites are \*porphyritic, with \*quartz and \*plagioclase as common \*phenocryst types. The term ‘toscanite’ was used originally by H. S. Washington in 1897 to describe rocks of rhyodacite composition from

Tuscany, Italy; this older term is now little used. Rhyodacites are erupted above subducted *\*plates* and belong to the calc-alkaline *\*magma* series.

**rhyolite** A fine-grained, *\*extrusive*, *\*igneous* rock, often with a sugary texture, consisting of *\*essential quartz*, *\*alkali feldspar*, and one or more *\*ferromagnesian minerals*. Alkali rhyolites are the most common type, being characterized by the ferromagnesian mineral *\*biotite* with or without *\*pyroxene*, and are found in calc-alkaline *\*terrains*. *\*Peralkaline* rhyolites are characterized by alkali pyroxenes (*\*aegirine*, aegirine–*\*augite*) and alkali *\*amphiboles* (*\*riebeckite*, *\*arfvedsonite*), and are found as *\*end-members* of alkaline *\*magma* series on oceanic islands and rifted *\*continental crust*.

**rhythmic sedimentation** See *CYCLOTHEM*.

**rhythmite** A sequence of fine-textured, regularly repeated bands laid down by a sequence of cyclical or rhythmic sedimentation (see *CYCLOTHEM*). Rhythmites are most commonly associated with freshwater environments, but can also be deposited by tidal movements.

**ria** Drowned river valley in an area of high relief. Classic examples are found in some of the peninsulas of western Europe, notably western Ireland, where they have resulted from the post-glacial rise in sea level.

**ribbon bomb** See *VOLCANIC BOMB*.

**ribbon jasper** See *JASPER*.

**ribbon lakes** See *TUNNEL VALLEY*.

**ribbons** Straight to sinuous, thin bodies of *\*sand*, with a narrow width in relation to their length. Sand ribbons develop on *\*sediment-poor*, tide-swept shelves (see *SHELF*), oriented parallel to the tidal stream. The term is also used more generally to describe the large-scale geometry of a preserved *\*sandbody* with a width to length ratio in excess of 1:100, and a thickness to width ratio greater than 1:10.

**Richmondian** A regional *\*stage* (449–445.6 Ma ago) of the Late *\*Ordovician* in the Upper *\*Cincinnatian series* of N. America, preceded by the *\*Caradoc* and followed by the *\*Hirnantian*.

**Richter, Charles Francis** (1900–1985) An American physicist and geologist, Richter is best known for his logarithmic scale of *\*earthquake magnitudes*. First proposed in 1927, the scale was later refined with the assistance of *\*Gutenberg*, with whom Richter also cooperated in a study of the world's greatest earthquakes, and other seismological work. See *MOMENT MAGNITUDE SCALE; RICHTER SCALE*.

**Richter denudation slope** A hillslope that develops at the foot of a cliff that is retreating fairly rapidly, chiefly by *\*rock fall*. The slope has a uniform gradient, is cut across bedrock, and stands at the angle at which the *\*talus* accumulates. With each unit of cliff retreat the related rock fall builds up on older talus, and so the foot of the cliff steadily rises. The Richter slope is revealed when the talus is removed, or it may remain hidden beneath a thick skin of mobile debris. The Austrian geographer and glaciologist Eduard (Edward) Richter (1847–1905) described such slopes in the Alps in 1900, and they are named after him.


**Richter scale** The measurement of the intensity of an *\*earthquake* using the amplitude of *\*seismic waves*. As the amplitude depends on the depth of the earthquake *\*focus*, the distance of the recording station from the focus, the travel path, and local geology at both the source and receiver, such magnitude estimates need to be constrained by several determinations. At any given recording station, the magnitude ( $M$ ) of a shallow earthquake is given by the equation:  $M = \log(A/T) + 1.66 \log \Delta + 3.3$ , where  $A$  is the maximum amplitude,  $T$  is the period, and  $\Delta$  is the *\*epicentral* angular distance between the earthquake and receiver. For deeper earthquakes, the magnitude is given using 20-second-period *\*Rayleigh* waves by  $M = \log(A/T) + af \Delta h + b$ , where  $h$  is the depth of the focus, and  $a$  and  $b$  are empirically determined constants for each seismic station.

**Ricker pulse** A seismic wavelet caused by the passage of a seismic pulse through an ideal viscoelastic medium, where the attenuation is proportional to the square of the frequency.

**ridge 1. (wedge)** An extension of high pressure from an *\*anticyclone* into a zone where generally lower pressure prevails. **2.** The poleward meanders of the flow of the upper westerly winds over mid latitudes. See also *LONG WAVE*. **3.** See *MID-OCEAN RIDGE*. **4.** See *RIDGE-AND-RAVINE TOPOGRAPHY; RIDGE AND RUNNEL*.

**ridge-and-ravine topography** A landscape consisting of a monotonous network of branching valleys and intervening low ridges, and which is similar to that of a maturely dissected *\*peneplain* (see *DAVISIAN CYCLE*). The term has been introduced, however, to avoid any genetic implications. It is well displayed in the central Appalachians, USA.

**ridge and runnel** A series of asymmetrical ridges running parallel to the coast and separated by shallow troughs (runnels) 100–200 m wide. This topography is developed on the foreshore of *\*mesotidal* or *\*macrotidal* beaches. The development of these forms is favoured by moderate wave-energy conditions acting on a flat *\*beach* with an abundant *\*sediment* supply.



**Ridge and runnel**

**ridge crest** The highest part of a *\*ridge*, typically 2–3 km above the level of the *\*abyssal plains*. With slow-spreading ridges (e.g. the *\*Mid-Atlantic Ridge*) the crest is split by a *\*median valley*, whereas fast-spreading ridges (e.g. the *\*East Pacific Rise*) have no median valley and the crest has more subdued topography.

**ridge-push** The hypothetical force, caused by the horizontal spreading of the near-surface *\*asthenosphere* at *\*constructive margins*, which is thought to be one of the two main driving forces for the movement of lithospheric *\*plates* (the other is *\*slab-pull*).

**riebeckite** A member of the alkali *\*amphiboles*  $\text{Na}_2(\text{Fe}^{2+}_3\text{Fe}^{3+}_2)[\text{Si}_4\text{O}_{11}]_2(\text{OH},\text{F})_2$  and *\*end-member* of an *\*isomorphous* series with *\*glaucophane*  $\text{Na}_2(\text{Mg}_3\text{Al}_2)[\text{Si}_4\text{O}_{11}]_2(\text{OH},\text{F})_2$ ; sp. gr. 3.43; *\*hardness* 5;

forms either tiny *\*prismatic* crystals or large, *\*poikilitic*, *\*subhedral*, prismatic crystals; dark bluish-green or black; occurs in *\*alkaline \*igneous* rocks, especially *\*granites*, in association with *\*aegirine*. Fibrous riebeckite (called crocidolite or blue *\*asbestos*) is formed by *\*metamorphism* of massive *\*ironstone* deposits. When infiltrated with silica it constitutes the semi-precious cat's eye or tiger's eye.

**Riedel shear structure** A network of *\*shear bands* that forms in the early stages of the formation of a *\*strike-slip fault*. The principal set of bands, called R shears, are at about 15° to the main fault, and a second set, R' forms at about 30°.

**riegel** A rock bar that extends across the floor of a *\*glacial trough*. It may be caused by a local reduction in the erosive ability of a *\*valley glacier* or by a local increase in bedrock strength, perhaps due to a reduction in *\*joint* density. It may alternate with a rock basin to give an irregular long profile.

**riffle** See POOL-AND-RIFFLE.

**rift 1.** A breach or split between two bodies that were once joined. See RIFT VALLEY. **2.** In quarrying, a split in *\*granite*, whose plane is oblique or perpendicular to the sheeting.

**rift valley** An elongate trough, of regional extent, bounded by two or more *\*faults*. Many rifts on land are associated with alkaline *\*volcanicity* and, because their margins are uplifted, many are starved of *\*clastic \*sediments* and so contain lakes; the E. African rift system is an outstanding example. Some rifts are thought to be at the embryonic stage of ocean development of the *\*Wilson cycle*, whilst others may become 'failed rifts' (or 'failed arms') and fill with sediment to become *\*aulacogens*. The rift valley developed along the axis of slow-spreading oceanic *\*ridges* is known as the *\*median valley* (or axial rift or axial trough) and is associated with the production of basaltic *\*magmas*. Tibetan rifts form at the end of the Wilson cycle as a result of the northward indentation of India into Asia and the spreading of the thickening Tibetan crust. *\*Graben* (the German word for 'ditch') can be used synonymously for 'rift valley' and also for an infilled, fault-bounded trough of any size, with or without topographic expression.

**right lateral fault** See DEXTRAL FAULT.

**rigidity modulus** See SHEAR MODULUS.

**RIGS** See REGIONALLY IMPORTANT GEOLOGICAL/GEOMORPHOLOGICAL SITES.

**rill** A shallow, narrow watercourse on sloping ground that is incised into the soil or the surface of easily eroded rock (e.g. **\*limestone**). Although small, a rill can transport significant amounts of soil and they are an early sign of **\*erosion**.

**rille (rima)** A small valley on the **\*Moon**. Three types are recognized. (a) Straight rilles are typically 1–5 km wide and hundreds of kilometres long, unrelated to surficial topography, analogous to terrestrial fault **\*grabens**. (b) Curved or arcuate rilles are variants of straight rilles, with similar dimensions, and form concentrically to major ringed basins (e.g. Mare Humorum). (c) Sinuous and meandering rilles, formed by thermal erosion by flowing **\*lava**. Hadley Rille, 1.2 km wide, 270 m deep, and 135 km long, visited by Apollo 15, is the type example. See also **RIMA**.

**rill-wash** Eroded material that is concentrated into more or less intermittent trickles and rills on inclined slopes, due to runoff of water.

**rima** On **\*Mars**, a rille or cleft, e.g. Rima Bradley, Rima Sirsalis.

**rime** The white deposit of ice that results from **\*crystal** growth on objects that are at a temperature below the freezing point. Supercooled water droplets in fog freeze on contact with such surfaces.

**rimmed shelves** See CARBONATE PLATFORM.

**rim syncline** An approximately circular depression that surrounds, or nearly surrounds, a **\*dome**.

**ring canal** See CIRCUM-ORAL CANAL.


**ring-dyke** Steeply dipping **\*dyke** of arcuate outcrop formed by the uprise of **\*magma** along a steep conical or cylindrical fracture which bounds central collapsed blocks. See also CAULDRON-SUBSIDENCE.

**ringed basin** See MULTI-RING BASIN.

**ring fracture** A steep-sided, outwardly dipping, fault pattern or fracture, circular or sub-circular in plan view, commonly associated with **\*ring-dykes**. Ring fractures surround collapsed volcanic depressions, and are thought to form due to circular **\*stress trajectories** created by a parent **\*pluton** at depth. See CALDERA; CAULDRON-SUBSIDENCE.

**ringing** See REVERBERATION.

**Ring of Fire (Pacific Ring of Fire)** A series of oceanic **\*trenches**, island arcs, and active **\*volcanoes** that surrounds the **\*Pacific Ocean** in an almost continuous chain, about 40 000 km long. It is a consequence of the movement of tectonic plates. On the eastern (American) side of the ocean, the **\*Nazca** and **\*Cocos** plates are being subducted beneath the **\*South American Plate**, the Cocos Plate is also being subducted beneath the **\*Caribbean Plate**, and part of the **\*Juan de Fuca** and Pacific plates are being subducted beneath the North American Plate. On the western (Asian) side of the ocean, the Pacific Plate is being subducted beneath the **\*Philippine** and **\*Indo-Australian plates**, and in the south the Pacific Plate is being subducted beneath the **\*Antarctic Plate**. The Ring of Fire contains 452 volcanoes and experiences about 90% of all the world's **\*earthquakes**.



**Ring of fire:** The triangles mark the locations of major volcanoes associated with subduction zones.


**ringshine** Reflected light from the rings of *\*Saturn* that illuminates features at the top of the saturnian atmosphere on the night side of the planet.

**ring silicate** See *CYCLOSILICATE*.

**ringwall** An archaic term that refers to the ring-like walls enclosing lunar craters or *\*mare* basins, as observed telescopically.

**Ringwood's rule** A rule stating that: 'Wherever *\*diadochy* in a *\*crystal* is possible between two elements with appreciably different *\*electronegativities*, the element with the lower electronegativity will be preferentially incorporated because it forms a stronger and more *\*ionic bond*.'

**rip current** Strong, narrow current usually of short duration, flowing seaward from the shore. The presence of a rip current can be detected as a visible band of agitated water flowing seawards, usually as a gap in the line of the incoming waves. Rip currents mark the swift return movement of water piled up on the shore by incoming waves and onshore winds.



**Rip current**

**Riphean** A *stage* of the Middle *Proterozoic*, lasting from about 1400 to 800 Ma ago, that was preceded by the *Animikian* and followed by the *Sinian*. Russian usage extends the era to about 680 Ma ago.

**rippability** A measure of the ease with which earth materials can be broken by mechanical ripping equipment to facilitate their removal by other equipment. Rippability is related to the *seismic velocity* of the material. Rock with a seismic velocity of less than 2 km/s can usually be ripped.

**ripple** Small-scale ridge of sand produced by flowing water, wind motion, or wave action. The wavelength or spacing of ripple crests is usually less than 50 cm and the heights are less than 20 cm. Ripple form can be

described by the wavelength to height ratio, referred to as the **\*ripple index**. The migration of ripples leads to the formation of **\*cross-lamination** in sands. *See also* **DUNE BEDFORM**.

**ripple-drift cross-lamination** A form of **\*cross-stratification** (sometimes called climbing-ripple cross-lamination) characterized by set boundaries which **\*dip** in the opposite direction to the **\*foresets**, giving the impression of one set climbing upwards over the underlying set. In some cases the ripple stoss slope (i.e. the gently dipping backslope) is preserved. Ripple-drift cross-lamination forms when there is a rapid rate of net deposition, with the angle at which the sets climb over one another being a function of increasing rate of sedimentation.

**ripple index** A measure of the symmetry of a ripple form, expressed by the ratio of ripple wavelength to ripple height. Flowing-water ripples (current ripples) have an index of 8–20 and are asymmetric, with a steeper face downstream (lee) and a gentle upstream-facing (stoss) side. The ripple index of wind ripples is 30–70; the structures are much flatter, reaching heights of 1 cm only. Wave-formed ripples (oscillation ripples) have a ripple index of 4–16 and have a more symmetrical profile form.

**ripple mark** A repeating, wave-like feature with peaks and symmetrical slopes found on the surface of a **\*sedimentary rock**. It results from the action of waves or wind.

**ripple-symmetry index** A measurement of the ratio of the horizontal extent of the **\*stoss** side of a **\*ripple** to the horizontal extent of the lee side. Wave ripples have ripple symmetry index values of less than 2.5, whereas current ripples have values of more than 3.0.

**ripple train** The collective term applied to a series of **\*ripples** which lie one behind the other, as found on a rippled surface.

**rip-rap** Loose foundation layer of large, irregular, unscreened rock fragments used under water or in soft material for protection and to prevent erosion of dams, sea walls, bluffs, or other structures exposed to wave action. Used extensively in irrigation works and river improvements.

**RISAT** *See* **RADAR IMAGING SATELLITE**.

**Rising-2** A Japanese Earth-observation *\*microsatellite* with a 5-m resolution that monitors *\*cumulonimbus* and *\*sprites*. It was launched on 24 May 2014, from the Yoshinobu Launch Complex, Japan, into a Sun-synchronous near-circular orbit at an altitude of 628 km.

**Riss** The third of four glacial episodes named after Alpine rivers and established in 1909 by A. *\*Penck* and E. Bruckner. It is perhaps the equivalent of the *\*Saalian* of northern Europe and the *\*Wolstonian* of the East Anglian succession.

**Riss/Würm Interglacial** An Alpine *\*interglacial \*stage* that may be the equivalent of the *\*Eemian* stage of northern Europe or the *\*Ipswichian* of the East Anglian succession. It is the last interglacial and immediately precedes the last glaciation, the *\*Devensian* or in European usage *\*Weichselian*.


**river capture** Process whereby a stream is able to tap and so capture the discharge of a neighbour. The capturing stream normally extends by headward *\*erosion* along an outcrop of soft rock until it meets and diverts a second, less-favoured, transverse system. A right-angled bend, the 'elbow of capture' is typical of the junction between capturing and captured streams. *See also* AVULSION.

**river grade** *See* GRADE.

**river profile** The slope of the long profile of a river, expressed as a graph of distance-from-source against height. It is generally concave-up, and this downstream reduction in gradient may be a consequence of the decreasing energy needed to transfer greater discharge and finer load. In detail, however, it is typically compound, with the profiles of individual segments reflecting the local rock types. It may be broken by *\*knick* points.

**river-sediment analysis** *See* STREAM-SEDIMENT ANALYSIS.

**river terrace (stream terrace)** Fragment of a former valley floor that now stands well above the level of the present *\*floodplain*. It is caused by stream incision, which may be due to uplift of the land, to a fall in sea level, or to a change in climate.



**River terrace**

**river water, major constituents** The average composition of river water is very different from that of sea water. The **\*pH** varies greatly from areas of high organic activity, e.g. the equatorial rain forests, to zones of little activity around the poles; the average is usually between 6.0 and 8.0. Large rivers, e.g. the Amazon, remove about  $10^5$  kg of material per  $\text{km}^2$  of **\*catchment** area every year. Of this,  $2 \times 10^4$  kg is dissolved and the rest is carried as solid particles of colloidal debris. The composition of typical river water is:

<i>Ion</i>	<i>Parts per thousand</i>	<i>Percentage of dissolved material</i>
bicarbonate $\text{HCO}_3^-$	58.5	48.6
calcium $\text{Ca}^{2+}$	15.0	12.5
silicate $\text{SiO}_2^-$	13.1	11.0
sulphate $\text{SO}_4^{2-}$	11.2	9.3
chloride $\text{Cl}^-$	7.8	6.5
sodium $\text{Na}^+$	6.3	5.3
magnesium $\text{Mg}^{2+}$	4.1	3.4
potassium $\text{K}^+$	2.3	2.0
nitrate $\text{NO}_3^{2+}$	1.0	0.8
iron $\text{Fe}^{2+}$ or $\text{Fe}^{3+}$	0.7	0.6
TOTAL	120.0	100.0

See also [SEA WATER, MAJOR CONSTITUENTS](#).

**RMQ** See [ROCK-MASS QUALITY](#).

**road base** See PAVEMENT.

**Roadian (Ufimian)** A \*stage (270.6–268 Ma ago) of the Middle \*Permian \*epoch, preceded by the \*Kungurian and followed by the \*Wordian.

**roadstone** Unconsolidated \*aggregate of strong rock which will withstand crushing and abrasion. It is used in the construction of roads.

**roadway construction** See PAVEMENT.

**roaring forties** Popular maritime term for the prevailing westerly winds which are commonly strong over the oceans in temperate latitudes of the southern hemisphere, particularly between about 40° S and 50° S. See also FURIOUS FIFTIES; SHRIEKING SIXTIES.

**Roche limit** The distance within which the tidal forces exerted by a planet are sufficient to disrupt a \*satellite or smaller body. For bodies in circular orbits with zero tensile strength and the same mean density as the primary, the Roche limit is  $2.46 \times$  (primary-body radius). In the case of the Earth–Moon system, the critical distance is 2.89 Earth radii (18 400 km).

**roche moutonnée (glaciated rock knob, stoss-and-lee topography)** Mound-like land-form of glacial \*erosion, consisting of a smoothed, streamlined, up-glacier surface and a broken, shattered, lee flank. It probably results from a combination of abrasion, frost-shattering, and the plucking out of blocks by the \*glacier, although crushing has been suggested as a contributory mechanism.

**rock** A consolidated or unconsolidated aggregate of \*minerals or organic matter. The minerals may be all of one type, in which case the rock is ‘monomineralic’, or of many types, in which case it is ‘polymineralic’. The aggregate of minerals can form by: (a) accretion or precipitation of grains during Earth surface processes, to give \*sedimentary rocks; (b) crystallization of \*magma to give \*igneous rocks; and (c) solid-state \*recrystallization in response to changes in external conditions (e.g. pressure and temperature) to give \*metamorphic rocks. The grain relationships (\*textures) of these three rock types contrast. Sedimentary rocks are characterized by one of the following: (i) rounded or angular grains held together by an intergrain precipitate or a fine intergrain mud; (ii) fine aggregates of \*clay minerals displaying a preferred orientation of their long axes; (iii) a crystalline aggregate of minerals (e.g. \*calcite) displaying

straight edges and triple junctions between the grains; (iv) an aggregate of *\*fossil* fragments held together by an interfragment precipitate of *\*calcite* or a fine interfragment mud; or (v) an aggregate of organic material (e.g. *\*lignite* or *\*coal*). All igneous rocks are characterized by an aggregate of minerals displaying an interlocking texture. Metamorphic rocks are characterized by one of the following: (i) a crystalline aggregate of minerals which display a preferred orientation of their long axes; (ii) a crystalline aggregate of equidimensional and randomly oriented non-equidimensional minerals; or (iii) an extremely fine-grained aggregate of sutured, *\*anhedral*, or sometimes elongate minerals.

**rock bench** See VALLEY-SIDE BENCH.

**rock bolt** A form of support for broken or jointed rock in mines or excavations. The far ends of metal bolts are secured in solid rock, with a face plate and nut at the external end. Rock bolts are usually used in conjunction with steel roof bars, plates, and mesh, combined with *\*grouting*.

**rock burst** Sudden, often explosive, breaking of rock walls when stressed beyond their limits. It is a hazard encountered in some deep mines (below 1000 m), and may be accompanied by shocks, rock falls, and air concussion. It usually occurs in rocks having high elasticity and strength, so that when a face is exposed the unbalanced *\*stress* is great enough to cause *\*failure*. It can be controlled by systematic *\*stoping*, reduction of open spaces, use of strong supports, and retreating mining methods.

**rock crystal** See QUARTZ.

**rock drumlin** See DRUMLIN.

**rock fall** The detachment of rock masses of variable size from a steep slope or cliff and their descent, chiefly through the air, by free fall, bounding, or rolling. Movement is very rapid. It is normally caused by *\*weathering*, or by the undercutting of a steep slope by *\*fluvial* or marine processes.

**rock fill** Waste rock used to back-fill worked-out sections of mines and to support the roof.

**rock flour** Finely ground rock debris produced chiefly by abrasion beneath a *\*glacier*. It may be removed by meltwater streams, which consequently

develop a typically milky appearance.

**rock glacier** A tongue-like mass of large, angular blocks, finer debris, and ice, found especially in middle latitude Alpine regions (there are about 1000 active examples in the Swiss Alps). It may have a core of ice, in which case it may have originated as a debris-covered glacier, or it may be ice-cemented, when it may be a *\*permafrost* phenomenon.

**rock head** 1. Surface between overlying unconsolidated material and solid bedrock below. 2. Bedrock on which sediments are deposited. 3. Base of gold-rich gravel where gold concentrates.

**Rocklandian** A regional *\*stage* (459–458 Ma ago) of the Late *\*Ordovician* in the Middle *\*Champlainian* *\*series* of N. America, preceded by the *\*Costonian* and followed by the *\*Soudleyan*.

**rock mass** A large and indistinct body of solid earth materials, containing features on the scale of *\*jointing*, *\*folding*, *\*schistosity*, etc. The term would not be used to describe a rock only the size of a hand specimen.

**rock-mass quality (RMQ)** A classification of *\*rock* for engineering purposes, based on the number of major *\*joints* or other discontinuities (planes of weakness) in it, their orientation, and their spacing.

**rock-mass strength** An estimate of the ability of a mass of rock to withstand *\*shear stress*. It is calculated on the basis of a *\*Schmidt hammer test* to determine the *\*intact* rock strength, the extent of *\*weathering*, the spacing, width, continuity, infill, and orientation of *\*joints*, and the outflow of *\*groundwater*. Each of these factors is given a weighting and rocks are classified as very strong, strong, moderately strong, weak, or very weak.

**rock mechanics** 1. The study of the physical behaviour of rocks, including crushing, bending, and *\*shear-strength* testing, and also their elasticity, *\*internal angle* of friction, *\*density*, *\*permeability*, and *\*porosity*. See ELASTIC DEFORMATION; ELASTIC LIMIT. 2. In geology, the study of the mechanics of rock structures, their physical properties, and forces acting on strata. 3. In engineering, the study of rocks as raw materials, and their behaviour in tunnels, quarries, and mines; and the stability of buildings on rock foundations.

**rock pavement** See RUWARE.

**rock-quality designation (RQD)** Rough measure of the degree of *\*jointing* or fracture in a *\*rock mass*, measured as a percentage of the drill core in lengths of 10 cm or more. High-quality rock has an RQD of more than 75%, low quality of less than 50%.

**rock salt** See HALITE.

**rock slide** A landslide made up of fragments and blocks of rock (slide-rock), and which is typical of steep slopes. The failure unit may consist of the *\*weathered zone*, when the form of the unweathered bedrock surface determines the shape of the shear plane. Elsewhere, failure may be defined by major *\*joints* or *\*bedding planes*.

**rock:soil ratio (RSR)** A measure of the relative proportions of rock and soil. It is a field criterion for separating fresh and weathered rock, which is often difficult to establish, especially in *\*sedimentary rocks*.

**rock-stratigraphic unit** See LITHOSTRATIGRAPHIC UNIT.

**rock-structure rating (RSR)** The nature and distribution of structural features within a *\*rock mass* which may have a dominant effect on the response of the rock mass to mining. The structure can influence the choice of mining methods and the design of the mine layout.

**rock unit** See LITHOSTRATIGRAPHIC UNIT.

**Rodda's flood-peak formula** See FLOOD-PEAK FORMULAE.

**rodding structure** A very coarse *\*lineation* of *\*minerals*, rock streaks, lines (striations), and *\*mullions*, which develops in strongly deformed rocks. The rods are cylindrical structures which *\*strike* parallel to the *\*hinge line*.


**Rodebaek** A glacial *\*stage* in Poland that may be equivalent to the *\*Amersfoort* of the Netherlands. It postdates sediments of *\*Eemian* age.

**rodingite** A *\*gabbro* or *\*dolerite* which has suffered calcium-*\*metasomatism* to produce a rock consisting of *\*grossular* *\*garnet* and *\*prehnite*, with or without *\*wollastonite*, *\*diopside*, and hydrogrossular.

**Rodinia** A *\*supercontinent* that existed during the *\*Mesoproterozoic* era (1600–1000 Ma ago). It began to form about 2200 Ma ago, as smaller continents were drawn together and started to unite. When fully developed,

Rodinia extended from about 60°N to the South Pole. The supercontinent began to break apart early in the **\*Neoproterozoic** era, leading to the later formation of another supercontinent, **\*Pannotia**.

**rogen moraine** A field of morainic (see **MORAINE**) ridges that lie at right angles to the direction of a former ice advance. Individual ridges may vary between 10 and 30 m in height, may be more than 1 km in length, and may be 100–300 m apart. They are often linked by cross-ribs. This landscape probably formed beneath a **\*glacier**, but the details of its origin are uncertain. It is named for its fine development around Lake Rogen, Sweden.



### Rogen moraine

**roll along** A seismic-**\*reflection** field method used to acquire data for a **\*common-depth-point** stack.

**rollover** The downwarping of the **\*hanging wall** block along a **\*listric fault**, which occurs when a space opens next to the fault. The resulting structure is usually called a 'rollover anticline'.

**rollover anticline** See **ROLLOVER**.

**roll-type uranium ore** C-shaped deposit, most commonly in **\*sandstones**, formed by advancing mineralizing fluids. Deposition takes place where **\*groundwaters** encounter reducing conditions, e.g. where rocks contain

organic material. They are usually no more than a few metres thick, and cut across bedding.

**Romer, Alfred Sherwood** (1894–1973) An American palaeontologist and comparative anatomist, who specialized in the evolution of vertebrates. He obtained his Ph.D. in 1921 from Columbia University, then taught anatomy at the Bellevue Hospital Medical College, New York, and from 1923 to 1934 he was an associate professor at the University of Chicago. In 1934 he was appointed professor of biology at Harvard University, in 1945 director of the biological laboratories, and in 1946 director of the Museum of Comparative Zoology, posts he held until his death.

**roof pendant** A downward projection of *\*country rock* into a *\*batholith*. The removal of roof rocks by *\*erosion* may leave remnants of roof pendants which resemble *\*xenoliths* in the intruding rock.

**roof thrust** In a *\*duplex* system, the structurally highest bounding *\*thrust* surface, which joins the *\*floor thrust* at the leading and trailing edges of the duplex.

**room and pillar** See PILLAR AND STALL.

**rooted tree** A *\*phylogenetic tree* in which the common ancestor is identified, usually by the incorporation of a known outgroup, thus resolving the direction of evolution.

**rootlet** Strictly, a small plant root, but the term is also commonly used to describe the traces of plant roots preserved in a *\*fossil \*soil* or *\*sediment*.

**root-mean-square velocity ( $V_{\text{rms}}$ )** The velocity of waves through subsurface media down to the  $n$ th interface, such that  $V_{\text{rms}} = (\sum V_{\text{int}}^2 t_i / \sum t_i)^{1/2}$ , where  $V_{\text{int}}$  and  $t_i$  are the interval velocity and single-travel time through the  $i$ th interval.  $V_{\text{rms}}$  is usually derived from normal *\*moveout* measurements from a  $T^2-X^2$  graph, and is commonly several per cent larger than the corresponding average velocity.

**root zone** A region from which *\*nappes* appear to be derived, and which has undergone extensive compression. Root zones are characterized by highly flattened and steepened structures.

**rope bomb** See VOLCANIC BOMB.

**Rosalind (Uranus XIII)** One of the lesser satellites of \*Uranus, with a diameter of 29 km. It was discovered in 1986.

**rose diagram** A circular histogram plot which displays directional data and the frequency of each class. Rose diagrams are commonly used in sedimentary geology to display palaeocurrent data (see PALAEOCURRENT ANALYSIS), or the orientation of particles. In structural geology rose diagrams are used to plot the orientation of \*joints and \*dykes. Wind directions and frequencies can also be plotted on rose diagrams.

**Rose Garden** A \*hydrothermal vent site at the \*Galápagos Rift, discovered in 1979, where blood-red tubeworms protruding from narrow, white tubes resembled a garden of long-stemmed roses. By 1985 the tubeworms had been replaced by mussels and clams.

**rose quartz** See QUARTZ.

**Rosetta** An \*ESA mission, to comet 67P/Churyumov–Gerasimenko, launched in March 2004. Rosetta swung by \*Mars early in 2007 and began to observe \*Jupiter. After ten years, it went into orbit around the comet and its lander, named Philae made the first-ever landing on the surface of a comet in November 2014. It transmitted much data for a few days before its battery ran out of power; but having moved closer to the sun in June 2015, it briefly began transmitting again. Rosetta remained with the comet until August 2015, with the mission terminating in September 2016 when Rosetta landed on the comet.



<http://sci.esa.int/rosetta/>

- ESA mission website.

**Rossby waves** Named after the Swedish-American meteorologist Carl-Gustav Rossby (1898–1957), Rossby waves are equatorward \*troughs and poleward \*ridges forming long waves in the circumpolar flow of the upper air, particularly in the mid and upper \*troposphere, with a typical wavelength of around 2000 km. Three or four waves usually occur in the circumpolar westerly wind flow over mid latitudes. They may remain stationary (as \*standing waves) when wind speed and wavelength have a given relationship. The waves may be initiated by lower winds over

mountain barriers, e.g. the Rocky Mountains, or by heating over warm oceans in winter or over land in summer. They are then amplified by vorticity (due to the Earth's rotation) in anticyclonic curvature (in ridges) and in cyclonic curvature (in troughs). Characteristic positions for the main troughs in the upper westerlies over the northern hemisphere are about 70° W and 150° E. The Rossby waves influence the formation of surface \*depressions which tend to develop on a \*frontal wave ahead of an upper trough. Rossby waves also occur in the oceans.

**Rossi–Forel scale** The first scale for reporting \*earthquake intensity, introduced in 1883 by the geophysicists Michele Stefano Conte de Rossi (1834–98) of Italy and François-Alphonse Forel (1841–1912) of Switzerland, and based on scales each of them had devised independently in 1874 and 1871, respectively. It remained in widespread use until the introduction of the \*Mercalli scale in 1902 and it is still used in some countries. The Rossi–Forel scale has 10 levels: I microseismic shock; II extremely feeble shock; III very feeble shock; IV feeble shock; V shock of moderate intensity; VI fairly strong shock; VII strong shock; VIII very strong shock; IX extremely strong shock; X shock of extreme intensity.

**rostral suture** See CEPHALIC SUTURE.

**rostrum** 1. (guard) A massive deposit of fibrous \*calcite which makes up part of a belemnite (\*Belemnitida) skeleton. 2. In a brachiopod (\*Brachiopoda), a beak-like process formed by the drawing out of the \*umbo.


**rotary drilling** The process of cutting a \*borehole in which ground is cut or crushed by a rotating drill system, using a drill \*bit turned by means of a \*kelly. Engineering and shallow studies usually involve either a mast or an A-frame. Deep drilling is usually from a derrick. See also CABLE DRILLING. Compare PERCUSSION DRILLING.

**rotating-cups anemometer** See ANEMOMETER.

**rotational remanent magnetism (RRM)** The magnetization acquired by rotation of a specimen within an alternating magnetic field. The process is unclear, but related to the movement of \*magnetic domains with a magnetic field. See also GYROREMANENT MAGNETIZATION.

**rotational shear** See SIMPLE SHEAR.

**rotational slip (rotational slump)** Variety of landslide characterized by movement along a concave-up failure surface, so the moving material partly rotates about an axis at right angles to the slope. Typically, the upper part of the block fractures into a series of \*scarps arranged like steps tilted backward, and the lower part (toe) forms a fan-shaped bulge covered with radial cracks and ridges. Surface water may be retained in the depressed zone. The slump may be caused by basal undercutting, or by the oversteepening of artificial embankments. It is normally found in a uniform, relatively weak material, e.g. \*clay.



Rotational slip


**rotational slump** See ROTATIONAL SLIP.

**rotation of the Earth** See EARTH ROTATION; TIDAL FRICTION.

**Rotliegende** The Lower \*Permian \*red beds underlying the \*Zechstein. Although old, the word is still very much in use, notably in the oil industry, since much of northern Europe's natural gas is obtained from this rock unit.

**Rotliegenden** An alternative name for the Cisuralian or Early \*Permian epoch, 299–270.6 Ma ago (Int. Commission on Stratigraphy, 2004).

**rotor cloud** Cloud formed in moist air by condensation in the upper part of a rotating eddy that has been generated beneath the wave-form in stable air on the lee side of a mountain barrier. The closed eddy system can result in local reversal of wind direction in the general air stream.



**Rotor cloud**

**roughness** See BED ROUGHNESS.

**rounded biosparite** See FOLK LIMESTONE CLASSIFICATION.

**roundness index** The average radius of curvature of the corners of a particle, divided by the radius of the maximum inscribed circle for a two-dimensional image of the particle, i.e.  $(\Sigma r/N)/R$ , where  $\Sigma r$  is the average radius of curvature at the corners,  $N$  is the number of corners, and  $R$  is the radius of the largest inscribed circle. In practice, it is used empirically and other techniques are also used. For example, a pebble may be compared with a set of standard silhouettes. The resulting index figure allows inferences to be made about the nature of the depositing process.

**Rowe cell** Instrument for testing the \*consolidation of large \*soil or loose \*sediment samples. The sample is compressed under hydraulic pressure and drained from below. Pore pressure and change in thickness of the sample can be measured.

**RPF** See RELATIVE POLLEN FREQUENCY.

**r-process** See RAPID-NEUTRON PROCESS.

**RQD** See ROCK-QUALITY DESIGNATION.

**RRM** See ROTATIONAL REMANENT MAGNETISM.

**RRR junction** A \*triple junction formed by the meeting of three \*ridges.

**R-selection** The *\*natural selection* of those organisms that breed in such a way as to maximize their intrinsic rate of increase (*r*) so that when favourable conditions occur (e.g. in a newly formed habitat) the species concerned can rapidly colonize the area. Such an opportunist strategy, based on producing large numbers of seeds, spores, eggs, or offspring most of which perish, is advantageous in rapidly changing environments, as in the early stages of a succession. *Compare* K-SELECTION.

**RSR** See ROCK:SOIL RATIO; ROCK-STRUCTURE RATING.

**RSS** Regional stratigraphic scale. See STRATIGRAPHIC SCALE.

**RST** See REGRESSIVE SYSTEMS TRACT.

**rubble levée** See LAVA LEVÉE.

**rubefaction** The reddening of some *\*soils* in the humid tropics as a result of the dehydration of iron compounds in the process of *\*ferralization*.

**rubidium–strontium dating** A *\*radiometric* dating method based on the *\*radioactive* decay of  $^{87}\text{Rb}$  to  $^{87}\text{Sr}$ . Rubidium has two *\*isotopes* ( $^{85}\text{Rb}$  72.15%,  $^{87}\text{Rb}$  27.85%), but only  $^{87}\text{Rb}$  is radioactive.  $^{87}\text{Rb}$  disintegrates in a single step to  $^{87}\text{Sr}$  by the emission of a low-energy beta particle (see BETA DECAY). Unfortunately this low-energy disintegration makes it very difficult to assess the half-life (see DECAY CONSTANT) and two values ( $5.0 \times 10^{10}$  years or  $4.88 \times 10^{10}$  years) have been in common use. When a mineral crystallizes, it will usually incorporate both rubidium and strontium *\*ions* and the ratio of Rb to Sr will vary depending on the mineral involved. This initial strontium in the mineral is known as ‘common strontium’ (see INITIAL STRONTIUM RATIO) and is normally in the proportion of  $^{88}\text{Sr}$  82.56%,  $^{87}\text{Sr}$  7.02%,  $^{86}\text{Sr}$  9.86%, and  $^{84}\text{Sr}$  0.56%. Using these proportions it is possible to identify the amount of radiogenic  $^{87}\text{Sr}$  present. Originally the above proportions were assumed, but today it is more usual to plot  $^{87}\text{Sr}:$  $^{86}\text{Sr}$  against  $^{87}\text{Rb}:$  $^{86}\text{Sr}$  to produce a straight-line *\*isochron* from which the age of the mineral can be determined. When using the  $^{87}\text{Rb}:$  $^{87}\text{Sr}$  method it is customary to use *\*whole-rock* samples in the analysis, because although  $^{87}\text{Sr}$  may leak from one mineral to adjacent minerals over time it usually remains in the system. *\*Micas* and potassium *\*feldspars* are the

most suitable minerals for  $^{87}\text{Rb}$ : $^{87}\text{Sr}$  age determinations and the results can commonly be compared with  $^{40}\text{K}$ : $^{40}\text{Ar}$  age determinations from the same sample (see [POTASSIUM–ARGON DATING](#)). The method has particularly been applied to ancient [\\*metamorphic rocks](#).

**ruby** See [CORUNDUM](#); [SPINEL](#).

**rudaceous rock** General term applied to [\\*sedimentary rocks](#) having a grain size of 2 mm or more.

**rudist bivalves** An extinct group of coral- and horn-shaped bivalves ([\\*Bivalvia](#), subclass [\\*Heterodonta](#)). They have a variable [\\*morphology](#) and many species are not easily seen to be bivalves. They are adapted to a sedentary mode of life. Some forms have a larger lower left valve and an upper right valve that forms a flat lid; other forms have a larger lower right valve. The lower valve is often very thick and may have a complex wall structure. There is a tendency to develop a cylindrical and coral-like form. Rudist bivalves are found in [\\*Cretaceous](#) rocks, sometimes forming extensive mounds.

**rudite** General name given to [\\*rudaceous rocks](#).

**rudstone** See [EMBRY AND CLOVAN CLASSIFICATION](#).

**Rugosa (tetracorals)** (subclass [\\*Zoantharia](#)) A [\\*Palaeozoic](#) order of solitary and colonial corals which appears in the [\\*Ordovician](#) and becomes extinct in the [\\*Permian](#). The [\\*corallum](#) contains radial plates (septa) and horizontal plates (tabulae), and sometimes oblique plates (dissepiments). The septa developed in insertion cycles of four and many species retain a degree of [\\*bilateral symmetry](#). Solitary rugosans seem to have preferred soft substrates since they had no means of cementation and colonial forms seem to have relied for stability on the weight of the skeleton.

**rugose** Applied to a shell that has a rough or wrinkled texture. The term is commonly used to describe the appearance of the [\\*epitheca](#) of some corals (e.g. [\\*Rugosa](#)), but is also applied to any shell texture where the surface is raised in a wrinkled manner to produce ridges. The word is derived from the Latin *rugosa*, meaning wrinkle.

**rule of Vs** A rule which describes the relationship between the [\\*attitude](#) of a bed and its [\\*outcrop](#) pattern for a given topography. In summary, if the

outcrop pattern of a bed forms a V-shape pointing down-gradient, that is the direction in which the bed *\*dips*. The rule is most usefully applied when studying maps on which outcrop patterns show moderately dipping beds traversing valleys.

**Rumford, Count** See THOMPSON, BENJAMIN.

**Runangan** See PRIABONIAN.

**Runcorn, Stanley Keith** (1922–95) A geophysicist and head of the School of Physics at the University of Newcastle, Runcorn was best known for his studies of *\*polar wander* curves, but he also did work on convection currents in the *\*mantle*. In 1959 he was able to show that the curves for N. America and Europe could be reconciled if it were assumed that the Atlantic Ocean had once been closed.

**runnel** See RIDGE AND RUNNEL.

**Rupelian (Stampian)** The first *\*stage* in the *\*Oligocene \*epoch*, preceded by the *\*Priabonian* (Late *\*Eocene*), followed by the *\*Chattian*, and dated at 33.9–28.4 Ma ago (Int. Commission on Stratigraphy, 2004). It is roughly contemporaneous with the lower *\*Zemorrian* (N. America), most of the *\*Whaingaroan* (New Zealand), and the upper *\*Aldingan* and lower *\*Janjukian* (Australia).

**rupes** A term for a straight escarpment. The classic example is Rupes Recta, the ‘Straight Wall’ in Mare Nubium on the *\*Moon*.

**Rusophycus** An *\*ichnogenus* of *\*cubichnia* that are believed to have been formed beneath a thin layer of sand.

**Russian borer** See PEAT-BORER.

**Rutherford, Ernest** (1871–1937) A New Zealand-born physicist, Rutherford worked at the Universities of Cambridge and Manchester, and at McGill University in Canada. He made important studies of radiation and the structure of the atom, developing the concept of a *\*half-life* for the decay of radioisotopes. This concept led directly to the techniques of *\*radiometric dating*.


**rutile** An *\*accessory mineral*, TiO<sub>2</sub>; iron, niobium, and tantalum may also be present; sp. gr. 4.2–5.6; *\*hardness* 6.0–6.5; *\*tetragonal*; normally

reddish-brown, but also yellowish-red, or black; pale brown *\*streak*; adamantine *\*lustre*; crystals normally square *\*prisms* terminated by pyramids; *\*cleavage* prismatic {110} and {100}; occurs in a variety of *\*igneous* rocks, *\*schists*, *\*gneisses*, and *\*metamorphic* *\*limestones* and *\*quartzites*, and becomes concentrated in *\*alluvial* deposits and beach sands. It is an important *\*ore mineral* for titanium and its compounds.

**ruware (rock pavement)** An area of bare rock with a slightly domed profile that locally *\*outcrops* at the surface of a tropical plain. It is formed when the *\*weathered profile* is stripped from a sound rock surface, and may be seen as the first stage in the emergence of a *\*dome* or *\*tor*.

**Ryukyu Trench** The oceanic *\*trench* which forms the boundary between the oceanic *\*Philippine Plate* and the *\*continental crust* of the *\*Eurasian Plate*. The Philippine Plate is subducting obliquely under the Eurasian Plate.

**Ryzanian** A regional *\*stage* (142–136.5 Ma ago) in the British Early *\*Cretaceous*, preceded by the *\*Volgian* and overlapping with the *\*Valanginian* (140.2–136.4 ma ago). It is roughly of *\*Berriasian* age and contains the uppermost Purbeck Beds.



**S** See SIEMENS.

**Saalian** A northern European glacial *\*stage* dating from about 0.25 to 0.1 Ma ago that may be equivalent to the *\*Wolstonian* of the East Anglian succession.

**sabkha** Wide area of coastal flats bordering a *\*lagoon*, where *\*evaporites*, dominated by carbonate–sulphate deposits, are formed. It is named after such an area on the Trucial coast of Arabia.

**saccate** Applied to *\*spores* and *\*pollens* where a separation of the *\*exine* layers produces an air sac (saccus).

**Saccominopsis** Known from the *\*Ordovician*, this genus is thought to be the first representative of the very important *\*protist* group the fusulinids. See also FORAMINIFERIDA.

**saccus** See SACCATE.

**saddle** See SUTURE.

**safe bearing capacity** The *\*bearing capacity* that is used when planning the foundations of a building. The safe bearing capacity ( $q_s$ ) is calculated by dividing the *\*ultimate bearing capacity* ( $q_f$ ) by a chosen safety factor ( $F$ ):  $q_s = q_f/F$ .

**Saffir–Simpson Hurricane Wind Scale** A standard scale, introduced in 1955 by meteorologists of the US Weather Bureau, for reporting tropical cyclones. It adds a further five categories to the *\*Beaufort scale*, together with the damage likely at each level. The former inclusion of the surface

atmospheric pressure at the centre of the low pressure system and the size of the *\*storm surge* it causes has been dropped. See [WIND STRENGTH](#).



<http://www.nhc.noaa.gov/aboutsshws.php>

- Describes the Saffir–Simpson Hurricane Wind Scale.

**sag and swell topography** See [KNOB AND KETTLE](#).

**sagduction** The gravitational descent of dense crustal rock into the *\*mantle* in the form of *\*diapirs*. The mechanism is proposed as an explanation for the formation of dome-and-basin (see [INTERFERENCE PATTERN](#)) *\*granite–\*greenstone* belts and as an alternative to *\*plate tectonics*, which some believe could not have occurred under the hot conditions early in Earth history, as a means of returning crustal material to the mantle.

**SAGE-III** See [STRATOSPHERIC AEROSOL AND GAS EXPERIMENT-III](#)).

**sag pond** A pool of water, typically a few tens of metres across and a few hundred metres long, that forms in an elongated depression caused by compressional or extensional *\*stresses* along a *\*strike-slip fault*.

**St David's** An *\*epoch* (513–501 Ma ago) of the Middle *\*Cambrian*, underlain by the Early Cambrian and overlain by the Late Cambrian.

**Sakian** A Russian-Kazakhstanian *\*stage* (494.5–493 Ma ago) of the Late *\*Cambrian \*epoch*, preceded by the *\*Ayusokkanian* and followed by the *\*Aksayan*.

**Sakigake** A Japanese *\*ISAS* mission to comet *\*Halley*, launched in 1985, carrying instruments to measure plasma wave spectra, solar wind ions, and interplanetary magnetic fields. All its instruments worked correctly. Sakigake flew by Halley on 11 March 1986, and swung by Earth three times, on 8 January 1992, 14 June 1993, and 28 October 1994. Communication with the satellite ended on 7 January 1999.



<http://nssdc.gsfc.nasa.gov/nmc/spacecraftDisplay.do?id=1985-001A>

- A Japanese mission to send a test spacecraft close to Halley's Comet, when it last approached the Earth in 1986.

**Sakmarian** A *stage* in the Early *Permian epoch*, preceded by the *Asselian*, followed by the *Artinskian*, and dated at 294.6–284.4 Ma ago (Int. Commission on Stratigraphy, 2004). Originally it was regarded as the basal Permian age, incorporating the Asselian age, from which it is now separated. It is roughly contemporaneous with parts of the *Rotliegendes* (western Europe), the upper *Wolfcampian* (N. America), and the upper Somoholoan (New Zealand).

**salcrete** A crust of salt–cemented mineral particles that forms on sandy beaches or other permeable surfaces which are periodically washed by salt water that dries rapidly. Continued exposure to the wind can produce *deflation hollows* resembling *sastrugi*, with their sides supported by the salcrete.

**salic** Applied to the silicon- and aluminium-bearing *CIPW* normative components of an *igneous* rock. Salic components include the normative *minerals* *quartz* (Q), *corundum* (C), *orthoclase* (Or), *albite* (Ab), *anorthite* (An), *nepheline* (Ne), *leucite* (Lc), and *kaliophilite* (Kp).

**salic horizon** *Soil horizon*, usually below the surface, containing not less than 2% salt and with a figure of 60 or more for the value calculated as the thickness of the horizon in centimetres multiplied by the percentage of salt. It is a *diagnostic horizon*.

**salina** **1.** A salt flat or similar place where salt deposits form by evaporation or are found. **2. (solar pond)** A highly saline pond or other body of water where salt crystals are produced by evaporation. The pond may be natural or artificial. **3.** See *PLAYA*.

**salination** See *SALINIZATION*.

**saline giant** A thick and extensive salt deposit, produced by the evaporation of a large hypersaline sea. One example of a saline giant is the *Miocene evaporites* of the Mediterranean, which formed by the repeated evaporation of the Mediterranean Sea. Another is the *Permian*-aged Zechstein salts of north-western Europe (see *ZECHSTEIN SEA*), which formed as a result of the repeated evaporation of a partially barred marine basin which covered more than 250 000 km<sup>2</sup>.

**saline-sodic soil** Soil that contains more than 15% exchangeable sodium, a saturation extract with a conductivity of more than 0.4 siemens per metre at 25 °C, and in the saturated soil it usually has a \*pH of 8.5 or less. Either high concentration of salts or high pH, or both, interfere with the growth of most plants.

**saline soil** Soil that contains enough soluble salt to reduce its fertility. The lower limit is usually defined as 0.4 siemens per metre.

**salinity** Measure of the total quantity of dissolved solids in sea water in parts per thousand ( ‰ ) by weight when all the carbonate has been converted to oxide, the bromide and iodide to chloride, and all the organic matter is completely oxidized. Ocean-water salinity varies in the range 33–38‰, with an average of 35‰.

**salinization** (US salination) The process of accumulating soluble salts in soil, usually by an upward capillary movement from a saline \*groundwater source, followed by evaporation from the surface.

**SALR** See SATURATED ADIABATIC LAPSE RATE.

**salt** The product, with water, of the reaction between an \*acid and a \*base.

**saltation** Major process of particle transport in either air or water, which involves an initial steep lift followed by travel and then a gentle descent to the bed. An essential requirement for the process is \*turbulent flow that can lift particles into the zone of relatively high downstream (downwind) velocity.

**salt dome** See DOME.

**salt-dome trap** Salt \*diapir which has pushed up existing \*sediments into a dome structure, and which may result in the trapping of gas, oil, or water in the \*pores of the permeable rocks adjacent to and above the salt dome if a suitable \*cap rock is present. The rocks ahead of the salt diapir are often severely faulted and may give rise to \*fault traps. Oil may also accumulate in the porous top of the salt diapir.

**salt fingering** A suggested mixing process between layers of saline and less saline water in the ocean. Where warm, saline water overlies cooler, less saline water, e.g. where the saline Mediterranean water flows out into the less saline Atlantic, mixing by this process is thought to take place. The

vertical water movements between the water masses of different **\*salinity** occur in small columns or fingers a few millimetres across. These fingers penetrate only a small distance, producing a mixed layer. The mixing process may then be repeated at the two interfaces that are present, and a number of layers may develop.

**salt flat** Extensive flat surface, found in hot deserts, consisting of salts that have accumulated in a shallow saline lake or **\*playa**. Evaporation then produces a crust of varying hardness.

**salt lake** A lake with a concentration of mineral salts typically of the order of 100‰ or greater, dominated by dissolved chlorides, e.g. the Dead Sea, which contains 64‰ NaCl and 164‰ MgCl<sub>2</sub>.

**salt marsh** Vegetation often found on mud banks formed at river mouths, showing regular zonation reflecting the length of time different areas are inundated by tides. Sea water has a high salt content which produces problems of osmotic pressure for the vegetation, so only plants that are adapted to this environment (halophytes) can survive there.


**salton** A sand grain that moves close to the ground, carried by the wind, and then collides with another grain and bounces upward, to where stronger winds at the higher level accelerate it.

**salt pan** A basin in a semi-arid region where chemical precipitates (**\*evaporites**) are deposited due to the concentration by evaporation of natural solutions of salts. The least soluble salts (calcium and magnesium carbonates) precipitate first, on the outside of the pan, followed by sodium and potassium sulphates. Finally, in the centre, sodium and potassium chlorides and magnesium sulphate are deposited. This pattern, slightly distorted through tilting, is seen in Death Valley, California, USA.

**saltpetre** See NITRE.

**salt weathering** A type of **\*weathering** that is most active in arid environments, where there is little **\*precipitation** and a high rate of evaporation. Precipitation fills voids and crevices with water, which dissolves **\*salts** from the adjacent rock. As the water evaporates, the salts crystallize. Subsequent heating or **\*hydration** causes the crystals to expand, widening the spaces they occupy and detaching loose rock fragments that are washed away with the dissolved salts the next time it rains.

**salt wedge** An intrusion of sea water into a tidal **\*estuary** in the form of a wedge along the bed of the estuary. The lighter fresh water from riverine sources overrides the denser salt water from marine sources unless mixing of the water masses is caused by estuarine topography. Salt wedges are found in estuaries where a river discharges through a relatively narrow channel.



**Salt wedge**

**samarium–neodymium dating** The change from  $^{147}\text{Sm}$  to  $^{143}\text{Nd}$  is the result of **\*alpha decay** ( $^{147}\text{Sm}$  has a **\*half-life** of  $2.5 \times 10^{11}$  years) and provides one of the newest methods of dating used in **\*geochronology**. The decay enriches the material in  $^{143}\text{Nd}$  relative to the stable isotope  $^{144}\text{Nd}$ . The ratio  $^{143}\text{Nd} : ^{144}\text{Nd}$  that is measured is highly resistant to secondary processes of alteration and **\*metamorphism**. It can be used on terrestrial and extraterrestrial materials and gives valuable secondary information on **\*petrogenesis** in the **\*crust** and **\*mantle**.

**sampling frequency (station frequency)** The frequency at which a data set is sampled is determined by the number of sampling points per unit distance or unit time, and the sampling frequency is equal to the number of samples (or stations) divided by the record (or traverse) length. For example, if a wave-form is sampled 1000 times in one second the sampling frequency is 1k Hz (and the **\*Nyquist** frequency 500 Hz); if a traverse is 500 m long with 50 stations, the sampling frequency is one per 10 m.

**sampling interval (station interval)** The distance between points at which measurements are taken or the time which elapses between measurements; it is equal to the traverse (or record) length divided by the number of stations (or samples). For example, a 250 m ground traverse with 25

stations along it has a sampling interval of 10 m; a wave-form might be sampled every two milliseconds, i.e. with a sampling interval of 2 ms (and a **\*sampling frequency** of 500 Hz).

**sampling methods** Techniques for collecting representative sub-volumes from a large volume of geologic material. The particular sampling method employed depends on the nature of the material being sampled and the kind of information required. Common methods are: (a) random sampling, a non-systematic or haphazard distribution of sampling locations; (b) systematic sampling, a regularly spaced distribution of sampling locations; (c) stratified sampling, a sequentially spaced, vertical distribution of sampling locations based on sampling each **\*stratigraphic unit** in a succession; (d) grab sampling, the collection of material using an externally controlled, mechanical grab (a method often used when sampling the sea floor or the surface of another planet); and (e) chip sampling, the collection of **\*borehole** drilling chips that are carried to the surface by lubricating mud and are representative of the material traversed by the borehole.

**sand 1.** In the commonly used Udden–Wentworth scale, particles between 62.5 and 2000  $\mu\text{m}$ . Other classifications exist (see **PARTICLE SIZE**). In pedology, sand is defined as mineral particles of diameter 2.0–0.02 mm in the international system, and as 2.0–0.5 mm diameter particles in the **\*USDA** (American) system. **2.** A class of soil texture. **3.** See **SHARP SAND**.

**sandbody** A finite unit of **\*sand** (or **\*sandstone**) that usually accumulated in response to one type of depositional process (e.g. as a **\*channel**, beach-bar, or **\*barrier** system). The distribution of the sand and the three-dimensional geometry (**\*architecture**) is controlled largely by the nature of the depositional regime under which it accumulated (i.e. channel sands may be sinuous, and beach or shore-bar sands may be linear and shore-parallel, etc.).

**Sander's symmetry principle** An interpretation of the **\*Curie symmetry principle** introduced by the German geophysicist B. Sander in 1930. It states that: Whatever the nature of the contributing factors, the symmetry that is common to them cannot be higher than the symmetry of the observed geologic texture, and symmetry elements absent in the texture must be absent in at least one of the contributing factors.

**sand line** The line on an *\*electrical* log marking the usual *\*apparent resistivity* of a clean *\*sand*.

**sand ribbon** Longitudinal strip of *\*sand* up to 15 km long, 200 m wide and less than 1m thick, standing on, and surrounded by, an immobile gravel floor. Sand ribbons are developed on the sea floor of the *\*continental shelf* where there are a paucity of sand, water depths of 20–100 m, and fast-flowing currents. See RIBBONS.

**sand sheet** See SHEET SAND.

**sandstone (arenite)** *\*Sedimentary rock* type, formed of a lithified *\*sand*, comprising grains between 63  $\mu\text{m}$  and 1000  $\mu\text{m}$  in size, bound together with a mud *\*matrix* and a mineral *\*cement* formed during burial *\*diagenesis*. The main constituents are *\*quartz*, *\*feldspar*, *\*mica*, and general rock particles, although the proportions of these may vary widely.

**sandstone dyke** See NEPTUNIAN DYKE.

**sandstorm** Phenomenon in which *\*sand* and dust particles are uplifted, often to great altitude, by turbulent winds. Visibility is greatly reduced.

**sandur** (*pl. sandar*) See OUTWASH PLAIN.

**sand volcano** A conical body of *\*sand*, resembling the form of a small volcano, rarely more than a few metres wide and less than 50cm high. Internally the sand volcano consists of a massive central plug, surrounded by laminated sand paralleling the external form. Sand volcanoes are formed by the extrusion of liquefied sand through a local vent at the surface. The extrusion usually results from a highly liquefied sand below a confining surface layer.

**sand wave** Large-scale, transverse ridge of *\*sand*, characteristic of *\*continental-shelf* areas such as the southern North Sea. The external morphology is identical to that of the smaller-scale ripple-and-*\*dune bedform* (megaripple). The wavelength or spacing of sand-wave crests is 30–500 m and the height is 3–15 m. The down-current migration of sand waves leads to the formation of large-scale cross-bedding.

**Sangamonian** The third (0.55–0.001 Ma ago) of four *\*interglacial \*stages* recognized in mid-continental N. America. It followed the *\*Illinoian* glacial

episode and is the approximate equivalent of the **\*Riss/Würm Interglacial** of the Alps. Early warm climates and later cooling climates are represented in well-exposed **\*pollen** spectra.

**sanidine** See ALKALI FELDSPAR.

**Santerian** See QUATERNARY.

**Santonian** A **\*stage** in the European Late **\*Cretaceous** (85.8–83.5 Ma ago, Int. Commission on Stratigraphy, 2004), for which the **\*stratotype** is at Saintes, France. It was preceded by the **\*Coniacian** and followed by the **\*Campanian**. See also SENONIAN.

**Sao (Neptune XI)** A satellite of Neptune with a radius of 20 km, visual **\*albedo** of 0.16, and an orbital period of 2914.1 days.

**sapphire** See CORUNDUM.

**saprolite** Chemically rotted rock *in situ*. The term is often applied to the lower portion of a **\*weathering profile**. The saprolite on **\*granite** is locally called ‘grus’ or ‘growan’, although the latter term may include material broken down by mechanical weathering. Compare REGOLITH.

**sapropel** Organic ooze or sludge accumulated in **\*anaerobic** conditions in shallow lakes, swamps, or on the sea bed. It contains more **\*hydrocarbons** than **\*peat**; when dry it is dull, dark, and tough; it may be a source of oil and gas.

**sapropelic coal** See COAL.

**sapropelite** A sapropelic **\*coal**, consisting of organic material, particularly **\*algae**, which accumulated in stagnant lake bottoms or the floors of **\*anoxic** shallow seas.

**SAR** See SODIUM-ADSORPTION RATIO.

**SARAL** See SATELLITE WITH ARGOS AND ALTIKA.

**SAR-Lupe** A German constellation of five reconnaissance satellites that uses **\*synthetic-aperture radar** (SAR) that provides surface observation independently of weather and illumination conditions to monitor crisis and emergency situations and to assist the military. The constellation was

launched between 2006 and 2008 at six-monthly intervals from the Plesetsk Cosmodrome, Russia, into three orbital planes in near-[\\*polar orbits](#) at an average altitude of about 500 km.

**Sarcopterygii** Class of fleshy-finned fish comprising two main groups (superorders): the coelacanth ([\\*Crossopterygii](#)), and the lungfish ([\\*Dipnoi](#)). The Crossopterygii appeared in the [\\*Devonian](#), were widely distributed in the [\\*Mesozoic](#), and are now represented by a single genus, *Latimeria*. One order of the Crossopterygii, the [\\*Rhipidistia](#), is credited with being the root stock for more advanced vertebrates. The Dipnoi also appeared in the Devonian and are now represented by three species. They inhabit fresh water and have developed organs by which they can breathe atmospheric air.

**sardonyx** See [ONYX](#).

**Sargasso Sea** Calm centre of the anticyclonic [\\*gyre](#) in the N. [\\*Atlantic](#). This large eddy of surface water has boundaries demarcated by major current systems such as the [\\*Gulf Stream](#), Canary Current, and [\\*North Atlantic Drift](#). The Sargasso Sea is a large, warm (18 °C), saline (36.5–37.0‰) lens of water, which is characterized by an abundance of floating brown seaweed (*Sargassum*).

**sarl** A [\\*pelagic](#) or [\\*hemipelagic sediment](#) (an [\\*arl](#)), typically found interbedded with purer oozes in beds up to 1.5 m thick, with a composition intermediate between a non-biogenic sediment and a calcareous or siliceous ooze. It is 30% clay and 70% microfossils, at least 15% of its volume being calcareous microfossils. Compare [MARL](#); [SMARL](#).

**saros unit** The term for the period of 18 years (inclusive of leap years) and 10.3 days in which the [\\*Earth](#), [\\*Moon](#), and [\\*Sun](#) return to the same relative positions, so that solar and lunar [\\*eclipses](#) repeat themselves. Thus the solar eclipse of 30 June 1973 was repeated on 11 July 1991. Each saros period contains about 43 solar and 28 lunar eclipses.

**sarsen stone** Boulder or block of [\\*sediment](#) consisting of a siliceous [\\*cement](#) binding one of a range of materials, and found widely in south-eastern England, especially in the chalk country. Typical cemented components include angular [\\*quartz](#) grains and rounded to angular [\\*flints](#)

(the 'Hertfordshire puddingstone'). Sarsens are of Palaeogene and Neogene age, and probably formed as a **\*duricrust** under tropical conditions.

**Sartan** See VALDAYAN/ZYRYANKA.

**sastrugi (zastrugi)** Irregular, small grooves and ridges that occur on the surface of wind-blown snow as a result of deposition and erosion. They resemble sand dunes, but unlike dunes they are aligned with the wind direction. The word is Russian and plural; singular is sastruga (zastruga).

**Satélite de Coleta de Dados (SCD)** A Brazilian satellite programme to gather environmental, mainly meteorological and hydrological, data that is distributed by about 700 terrestrial platforms. SCD-1 was launched on 9 February 1993, from Kennedy Space Center, into a near-circular orbit at an altitude of 750 km.

**satellite** A minor body orbiting a planet in the **\*solar system**. About 180 are known, divided into three general classes. (a) Regular satellites form miniature solar systems and include all the classical major satellites, e.g. the **\*Galilean satellites**. (b) Collisional shards are tiny, craggy chunks, probably remnants of larger satellites, e.g. **\*Amalthea**, which is embedded in **\*Jupiter's** planetary ring system. (c) Irregular satellites have elongate, highly inclined orbits, mostly far from the planet, suggestive of capture, e.g. the outer satellites of Jupiter. Three bodies, the Earth's **\*Moon**, **\*Triton** (orbiting **\*Neptune**), and **\*Charon** (orbiting **\*Pluto**) do not fit into any of the above classes, and each has to be regarded as a unique case. See Appendix I for list of satellite missions.

**satellite photography** The production of photographic images, using the visible, infrared, thermal infrared, and other wavebands, by means of satellite-based cameras or sensors. Such images are extensively used in the environmental sciences for the study of the oceans, the atmosphere, and land masses.

**satellite sounding** Remote sensing of atmospheric properties by an orbiting Earth satellite. For example, indirect sensing is carried out by spectral analysis of outgoing long-wave radiation with infrared photography of clouds and determination of cloud-top height from the temperature variations.

**Satellite with ARgos and ALtiKa (SARAL)** A *\*minisatellite* altimetry mission of the Indian Space Research Organization and the French space agency that aims to produce precise, repetitive, global measurements of sea surface height, significant wave heights, and wind speeds to assist oceanography, climate, and operational meteorology. It was launched on 25 February 2013 into a Sun-synchronous near-circular orbit at an altitude of about 800 km.

**satin spar** See GYPSUM.

**saturated** See SILICA SATURATION.

**saturated adiabatic lapse rate (SALR)** The *\*adiabatic* cooling rate of a rising *\*parcel* of air which is saturated (see SATURATED AIR), and in which condensation is taking place as it rises, so that the energy release of the latent heat of vaporization moderates the adiabatic cooling. The reduction of the rate of cooling below the *\*dry* adiabatic lapse rate of 9.8 °C/km varies with temperature. This results from the greater energy release by condensation from air at higher temperatures. Thus at a given atmospheric pressure, air at 20 °C may have an SALR as low as 4 °C/km, whereas at -40 °C the SALR may be close to 9 °C/km. The *\*stability* or *\*instability* of the atmosphere at any given time for vertical motion is determined by whether the *\*environmental lapse rate* of temperature within it is less than or greater than the adiabatic lapse rate (i.e. less than or greater than the rate of decrease of temperature of rising parcels of air). An average SALR value is 6 °C/km.

**saturated air** Air that contains the maximum amount of water vapour that is possible at the given temperature and pressure, i.e. air in which the *\*relative humidity* is 100%.

**saturated flow** The movement of water through a soil that is temporarily saturated. Most of the loosely held water moves downward, and some moves more slowly laterally.

**saturation** In *\*remote sensing*: 1. the maximum digital number value which can be assigned to a *\*pixel*; 2. a point between the *\*achromatic line* and a pure hue of a *\*pixel colour* corresponding to the relative mixture of hues going to make up a colour.

**saturation deficit** At a given temperature, the difference between the actual **\*vapour pressure** of moist air and the **\*saturation vapour pressure**.

**saturation magnetization and moment ( $M_{\text{sat}}$ )** The maximum remanent magnetization (see **REMANENT MAGNETISM**) that a material can acquire after being placed in a direct magnetic field. If uncorrected for either volume or weight, the observed magnetization is the saturation magnetic moment.

**saturation moisture content (SMC)** The maximum amount of water that can be contained in a rock, when all **\*pore spaces** are filled with water; it is expressed as the percentage of the dry weight of the rock.

**saturation vapour pressure** The **\*vapour pressure** at which the layer of air immediately above the surface of liquid water is saturated with moisture at a given temperature.

**Saturn** The sixth planet in the **\*solar system**, distant 9.52 AU from the **\*Sun**. Its radius is 60 000 km, density 704 kg/m<sup>3</sup>, mass 95 × Earth mass, volume 833 × Earth volume, and it has an equatorial inclination to the **\*ecliptic** of 29°. An outer zone of hydrogen and helium is underlain by a zone of metallic hydrogen, around an ice–silicate core. It has 62 **\*satellites** and is famous for its ring system. The orbiter **\*Cassini** studied Saturn's atmosphere, its mission ending on 15 September 2017, when it descended into the planet's atmosphere.

**saturnian satellites** **\*Saturn** has 62 known satellites, of which 18 are described here. See **ATLAS (SATURN XV)**; **CALYPSO (SATURN XIV)**; **DIONE (SATURN IV)**; **ENCELADUS (SATURN II)**; **EPIMETHEUS (SATURN XI)**; **HELENE (SATURN XIII)**; **HYPERION (SATURN VII)**; **IAPETUS (SATURN VIII)**; **JANUS (SATURN X)**; **MIMAS (SATURN I)**; **PAN (SATURN XVIII)**; **PANDORA (SATURN XVII)**; **PHOEBE (SATURN IX)**; **PROMETHEUS (SATURN XVI)**; **RHEA (SATURN V)**; **TELESTO (SATURN XII)**; **TETHYS (SATURN III)**; **TITAN (SATURN VI)**.

**Saucesian** A **\*stage** (22–16.5 Ma ago) in the **\*Miocene** of California, underlain by the **\*Zemorrian**, overlain by the **\*Relizian**, and roughly contemporaneous with the upper **\*Aquitania** and **\*Burdigalian** stages.

**saurian** Of, or resembling a lizard. Loosely refers to lizard-like animals, and applied to **\*fossils**, life habits, etc. of the extinct **\*reptiles**.

**Saurischia** The 'lizard-hipped' \***dinosaurs**, one of the two dinosaur orders. They included bipedal carnivores (\***Theropoda**) and herbivorous \***tetrapods** (\***Sauropoda**). The theropods produced the largest known terrestrial carnivores, and the sauropods yielded the largest known land animals.

**saurischian dinosaur** See SAURISCHIA.

**Sauropoda** \***Jurassic** and \***Cretaceous** quadrupedal \***dinosaurs** of herbivorous habit. They included *Diplodocus*, from the Upper Jurassic, one skeleton of which, measuring 26.6 m, at one time made *Diplodocus* the longest terrestrial animal known, but larger sauropods have since been discovered. *Brachiosaurus*, from the Late Jurassic of the USA and Tanzania, is estimated to have weighed 80 t, and *Supersaurus* is estimated to have been 15 m tall and possibly 30 m long. \***Apatosaurus** (*Brontosaurus*) was also a sauropod, and although shorter than *Diplodocus* its skeleton was more massively built.

**Sauropterygia** (subclass \***Archosauria**) An extinct reptilian order, that comprises the two suborders \***Plesiosauria** (plesiosaurs) and \***Nothosauria** (nothosaurs).

**Saussure, Horace Bénédict de** (1740–99) A Swiss naturalist, Saussure made an extensive study of the structure of the Alps, described in the four volumes of *Voyages dans les Alpes* (1779–96). His theory was \***neptunian**, but with \***uniformitarian** overtones.

**saussuritization** The complete or partial alteration of calcium-rich \***plagioclase** to a fine-grained aggregate of secondary, sodic-rich plagioclase, \***epidote**, \***muscovite**, \***calcite**, \***scapolite**, and \***zeolites**. The process commonly takes place during the low-grade \***regional metamorphism** of \***gabbros** and \***basalts**, both of which contain plagioclase as an \***essential** component.

**Saxonian** The European stratigraphic term that is equivalent to the Upper \***Rotliegendes** and Weissliegende. See also AUTUNIAN.

**S-band** The radar frequency band between 1550 and 5200 MHz.

**scalenohedron** A \***crystal** form consisting of a number (usually 6 or 12) of triangular faces, all with unequal sides. Its \***crystal symmetry** is either \***tetragonal** or, more frequently, \***hexagonal**. Each face cuts the vertical *c*

(or z) axis and the form is often developed in the mineral *\*calcite*, where it is called 'dog-tooth spar'.

**scandent** In a graptoloid graptolite (*\*Graptoloidea*), applied to the condition where the *\*stipes* are united back to back with the nema in between.

**Scandinavian ice sheet** An *\*ice cap* which developed over Scandinavia during the *\*Quaternary*. It has been suggested, largely on the basis of the degree of downwarp and the partial recovery of the land surface in Scandinavia and the surrounding areas, that the ice was 2600 m thick.

**scanning electron microscope (SEM)** A microscope that operates by scanning a finely focused beam of *\*electrons* across the specimen. The reflected electron intensity is measured and displayed on a cathode-ray screen to produce an image. The SEM enables magnifications of up to 100 000 times to be made and provides a much better depth of field than a conventional light microscope (which suffers from focus limitations), making the three-dimensional structure of small objects (e.g. *\*Foraminiferida*) spectacularly visible. In geology it is used extensively for *\*micropalaeontology*, diagenetic studies (see *DIAGENESIS*), and *\*grain* textural examination. When coupled with an *\*electron* probe, semi-quantitative determinations of grain chemistry can be made.

**scapolite** A member of the *\*feldspathoids*, of composition  $(\text{Na,Ca,K})_4[\text{Al}_3(\text{AlSi})_3\text{Si}_6\text{O}_{24}] (\text{Cl,SO}_4,\text{CO}_3,\text{OH})$ , forming a *\*solid solution* series between the two *\*end-members* marialite (Na and Cl) and meionite (Ca and  $\text{CO}_3$ ); sp. gr. 2.5 (mar) to 2.7 (me); *\*hardness* 5–6; white or pale bluish or greenish; small *\*prisms* or *\*massive*; occurs in some *\*pegmatites* replacing *\*quartz* or *\*plagioclase*, but mainly in *\*metamorphic* or metasomatic rocks (see *METASOMATISM*). Scapolite is found in association with *\*sphene*, *\*grossular*, *\*diopside*, and *\*epidote*.

**scar** Steep, cliff-like slope of bare rock, developed in the near-horizontally bedded *\*Carboniferous \*limestone* of the Yorkshire Dales, England. The steepest and highest scars are normally associated with the *\*outcrop* of the purest and most massively bedded limestone. Often a *\*scree* is formed at the base.

**scarp (abbreviation of escarpment)** Steep slope or cliff found at the margin of a flat or gently sloping area. Many varieties are recognized, and distinguished in terms of origin. A 'fault scarp' results when a *\*fault* displaces the ground surface so that one side stands high. A 'fault-line scarp' is produced by *\*erosion* on one side of an ancient fault: *\*obsequent* and *\*resequent* varieties are recognized. A 'composite fault-line scarp' results from a combination of erosion and faulting. Erosional scarps result from vertical incision, or from the headward enlargement of *\*pediments*.

**scarp, lobate** A fault *\*scarp* characterized by lobes. They are common on the surface of *\*Mercury*, where they are relatively steep, with crest heights from 0.5 to 3 km, lengths from 20 to 500 km, with a broadly lobate outline on a scale of a few to tens of kilometres. They are probably due to *\*reverse* or *\*thrust* faulting resulting from compressive stress. They predate the close of the massive bombardment, and so are probably more than four billion years old. If so, they record an early period of planetary contraction.

**scarp-and-vale topography** A landscape consisting of a roughly parallel sequence of *\*cuestas* (*\*scarps* and *\*dip* slopes) and intervening valleys ('vales'). It is typically found on uniclinal (homoclinal) structures whose beds show differing *\*lithological* composition and consequently varied resistance to *\*denudation*. It dominates most of lowland Britain, which is characterized by *\*Mesozoic* *\*sediments* dipping gently towards the east and south-east.

**scarp-foot knick** An abrupt change in gradient that often occurs in semi-arid environments between a *\*pediment* and the adjacent *\*scarp*. It is the boundary between the zones of *\*scarp retreat* and pedimentation. It may be *\*joint* controlled, when it is often so abrupt that a boot may barely be placed in it.

**scarp retreat** The recession of the relatively steep hillslope that terminates a *\*butte*, *\*mesa*, *\*cuesta*, or any elevated, plateau-like surface. Several geomorphological *\*processes* may be involved, including undercutting by an adjacent stream, *\*spring sapping*, *\*mass movement*, *\*rain-wash*, and *\*weathering*. Under semi-arid conditions it may give rise to a hillslope that retreats parallel to itself.

**scarp slope** The relatively steep face of a *\*scarp*. Its steepness is maintained by the *\*erosion* of a relatively weak *\*stratum* that typically

underlies the resistant **\*cap rock** that maintains the form of the scarp. Erosion may be achieved by **\*spring sapping**, **\*sheetwash**, and **\*mass-wasting**.

**SCATSat-1** See SCATTEROMETER SATELLITE-1.

**scatter diagram** In statistics, the diagram obtained when two sets of observations are plotted against each other. Scatter diagrams are usually employed to visualize any correlations that may occur between two sets of observations.

**scattering** Diffusion of incident radiation by atmospheric particles, e.g. by haze and water droplets as well as by molecules. Such diffused radiation is often refracted many times in passing through the atmosphere. See also **MIE SCATTERING**; **RAYLEIGH SCATTERING**.

**Scatterometer Satellite-1 (SCATSat-1)** A gap-filling mission between **\*OceanSat-2** and -3 by the Indian Space Research Organization to monitor ocean surface winds. It was launched on 26 September 2016, from Satish Dhawan Space Centre, India, into a Sun-synchronous orbit at an altitude of 720 km.

**scavenger** An animal that feeds mainly on dead animals or animal products (e.g. faeces).

**scavenging** The capture and removal by rain or snow of particulate matter in the atmosphere.

**SCD** See SATÉLITE DE COLETA DE DADOS.

**scheelite** Mineral,  $\text{CaWO}_4$ ; sp. gr. 5.9–6.1; **\*hardness** 4.5–5.0; **\*tetragonal**; white, or sometimes shades of yellow, green, brown, or red; white **\*streak**; **\*vitreous** **\*lustre**; **\*crystals** usually **\*bipyramidal**, also occurs **\*massive** or granular; **\*cleavage** good {111}; **\*fluorescent**; occurs in **\*pegmatite** **\*veins** and high-temperature mineral veins, often in association with **\*wolframite**, **\*cassiterite**, **\*molybdenite**, **\*fluorite**, and **\*topaz**, also in contact **\*metamorphic zones** together with **\*axinite**, **\*garnet**, **\*wollastonite**, and metamorphic calcium minerals. It is one of the chief ores of tungsten, and named after the Swedish chemist K. W. Scheele (1742–86).

**Scheinwoodian** (**Tonawandian**) A substage of the N. American **\*Niagaran** **\*stage** (Early **\*Silurian** **\*epoch**), preceded by the **\*Llandovery** sub-epoch and followed by the **\*Homerian**.

**Scheuchzer, Johann Jacob** (1672–1733) A Swiss mathematician and physician, Scheuchzer is best known for his work on fossil fish and plants which he attributed to a universal deluge. He is notorious for finding, in 1726, the fossil remains of *Homo Diluvii Testis* (the man who witnessed the deluge), which was taken as important evidence for the Mosaic Flood, but was later shown by **\*Cuvier** to be a giant salamander.

**schiller** A metallic, bronze-coloured **\*lustre**, seen for example in moonstone (see **ALKALI FELDSPAR**).

**Schindewolf, Otto H.** (1896–1971) A German palaeontologist at the University of Tübingen, Schindewolf made extensive studies of ammonites (**\*Ammonoidea**) and their evolutionary history. These studies led him to propose a concept of discontinuous evolution, a theory which is not now generally accepted.

**schist** A **\*regional metamorphic** rock of pelitic (see **PELITE**) composition which displays a **\*schistosity**. Schists are coarser-grained than **\*phyllites**, having a grain size greater than 1 mm. The minerals defining the schistosity may be **\*muscovite** **\*mica**, **\*biotite** mica, and/or elongate **\*quartz**, depending on the composition and the pressure and temperature of formation. When a **\*basic** **\*igneous** rock is metamorphosed it forms a **\*hornblende-schist** (**\*amphibolite**) or **\*greenschist** if it contains a planar **\*fabric**, or a **\*greenstone** if no fabric is present. Thus in this latter context 'schist' refers to the fabric component and not to the overall rock type.

**schistosity** The planar alignment of platy **\*micas** and elongate **\*amphiboles** in a **\*regional metamorphic** rock. The alignment of these minerals into a planar **\*fabric** is caused by (a) the physical rotation of the mineral grains under the influence of a **\*shear stress**, or (b) the syntectonic, metamorphic growth of new minerals with their long axes perpendicular to the principal compressive stress direction.

**schizochroal** See **TRILOBITE EYE**.

**schizodont** Applied to a type of **\*hinge** **\*dentition**, found in certain members of the pelecypod order Trigonioidea, in which the teeth are large

and possess parallel ridges at right angles to the axis of the teeth. The left *\*valve* bears a single tooth.

**Schizomycophyta** A name formerly given to the *\*bacteria*. See EUBACTERIA.

**schlieren** An *\*igneous fabric* consisting of streaked-out, linear, planar, or discoidal aggregates of *\*mafic* minerals more densely concentrated than in the host rock, with which they have diffused boundaries. Some schlieren may represent partly remobilized, mafic-rich *\*xenoliths*, streaked out by the flow of the enclosing *\*magma*.

**Schlumberger array** An *\*electrode* configuration in which the spacing of the two *\*potential electrodes* is less than one-fifth of the distance between the centre of the *\*array* and one *\*current electrode*. See also GEOMETRIC FACTOR.

**Schmidt hammer** An instrument for measuring the compressive strength of a surface by hammering it with a spring-loaded metal piston. It is used to test rock hardness and abrasiveness before drilling.

**Schmidt hammer test** In *\*geomorphology*, a technique for comparing the surface toughness or hardness of rocks. For example, it has been used to compare the mechanical strength of the surface layers of weathered (see WEATHERING) *\*limestones* and *\*sandstones* with that of the unaltered rock beneath. See SCHMIDT HAMMER.

**Schmidt–Lambert net** See EQUAL-AREA NET.

**schorl** A black, opaque variety of *\*tourmaline* which may occur as radiating, needle-like crystals, as seen, for example, in the *\*granites* of Cornwall, Britain.

**Schroeder Van Der Kolk method** See HALF-SHADOW TEST.

**schuppen structure** See IMBRICATE STRUCTURE.

**Schwassmann–Wachmann 3** A *\*comet* with an orbital period of 5.35 years; *\*perihelion* date 2 June 2006; perihelion distance 0.933 AU.

**Science Satellite/Atmospheric Chemistry Experiment (SciSat-1/ACE)** An atmospheric science mission by the Canadian Space Agency that allows

scientists to define and conduct space experiments in Earth sciences, space astronomy, and solar-terrestrial relations. It was launched on 13 August 2003, from California, into circular, high-inclination orbit at an altitude of 650 km.

**scintillation counter (scintillometer)** An instrument which measures gamma radiation and is extensively used in airborne and ground radiometric surveys. It utilizes the flash of light emitted when the atoms of a suitable 'phosphor' (e.g. a large sodium iodide crystal 'doped' with thallium) are energized by \*gamma rays. The scintillations are detected by the light-sensitive cathode of a photomultiplier tube and are converted by the succession of electrodes in the tube into a stream of electrons which are collected and recorded on a meter. The scintillation counter has now been developed into the gamma-ray spectrometer (see [GAMMA-RAY SPECTROMETRY](#)) for portable and airborne use. It analyses the complex gamma-ray spectrum of uranium, thorium, and potassium, and indicates the relative gamma-ray contribution of each element to ground gamma-ray emission on a continuous readout.

**Scintillation Observations and Response of the Ionosphere to Electrodynamics (SORTIE)** A \*NASA 6-unit \*CubeSat mission to study wave-like \*plasma perturbations in the \*ionosphere. The \*nanosatellite was launched on 21 May 2018, from Virginia, into a near-circular orbit at an altitude of about 400 km.

**Scintillation Prediction Observations Research Task (SPORT)** A collaborative 6-unit \*CubeSat mission between \*NASA, the Brazilian National Institute for Space Research, and the Brazilian Air Force to study the preconditions leading to equatorial \*plasma bubbles. It is scheduled for launch in 2019.

**scintillometer** See [SCINTILLATION COUNTER](#).

**scintillometer survey** A geophysical prospecting method using a scintillation detector in which radioactivity causes crystals to emit flashes of light which can be recorded by a photomultiplier tube. It is much more sensitive than methods using a \*Geiger counter and can distinguish different types of radiation.

**scirocco (sirocco)** Regional term for one of the genera of warm winds from south of the Mediterranean. It moves ahead of an eastward-travelling depression and brings hot, dry, dusty conditions to Algeria and the Levant. To the north, where its humidity increases very rapidly as it crosses the sea, it brings moist air to the coast of Europe.

**SciSat-1/ACE** See SCIENCE SATELLITE/ATMOSPHERIC CHEMISTRY EXPERIMENT.

**Scleractinia (Madreporaria; stony corals)** (subclass *\*Zoantharia*) Order of solitary or, more commonly, colonial corals, which always possess an external calcareous skeleton consisting essentially of radial partitions (septa, see SEPTUM). Septa develop following the pattern of the radial infoldings in the body wall (*\*mesenteries*) in cycles of 6, 12, 24, 48, etc. The order first appeared in the Middle *\*Triassic*.

**sclerotinite** See COAL MACERAL.

**SCLM** See SUB-CONTINENTAL LITHOSPHERIC MANTLE.

**scolecodont** The pharyngeal jaws or maxillae of annelid worms (*\*Annelida*), commonly found in paired assemblages. They are usually black and chitinous and are known from most geologic *\*systems*.

**scolecoïd** See SOLITARY CORALS.

**Scolicia** An *\*ichnoguild* of structures made by motile, chemosymbiotic organisms (see CHEMOSYMBIOSIS) found in sands and muddy sands dating from the *\*Jurassic* or possibly earlier. Recent ones were inhabited by *Echinocardium cordatum* (sea potato, *\*Miocene* to the present), a species of heart urchins, remains of which have been found within them.

**scopuli** See SCOPULUS.

**scopulus** (*pl. scopuli*) An irregular or lobate scarp on the surface of an extraterrestrial body.

**scoria** Loose, rubbly, basaltic ejecta that accumulate around *\*Strombolian eruptive* volcanic vents, eventually building up as a *\*scoria cone*, whose height may range from a few tens of metres to up to 300 m, and whose slope is determined by the *\*angle* of repose of the loose material. The scoria

**\*clasts** range widely in size, and have a light, frothy texture, being full of **\*vesicles**. They are mainly drab grey in colour, although when fresh they may be iridescent, but often the scoria oxidizes by reaction with steam escaping from the vent, when it becomes a deep reddish brown.

**scoriaceous** Applied to a vesicular **\*lava** or **\*pyroclastic** rock, to describe a frothy, bubbly texture. See SCORIA; VESICLE.

**scoria cone (cinder cone, ash cone)** Volcanic cone built of **\*pyroclastic** material (cinders and **\*scoriaceous** ejecta) usually of **\*basaltic** or **\*andesitic** composition, dominated by fragments 2–64 mm diameter. These air-fall deposits build a straight-sided cone with slopes of about 30° (the **\*angle** of repose). Complex varieties may occur, including breached or gaping, and cone-within-cone types.

**scorpions** See CHELICERATA.

**Scotia Plate** A tectonic **\*plate** that borders the **\*South American Plate** to its north, the **\*Antarctic Plate** to its south and west, and the South Sandwich **\*Microplate** to its east. **\*Transform fault** boundaries border the Scotia Plate to the north and south and there is a spreading boundary between it and the South Sandwich Microplate. The western boundary with the Antarctic Plate is more complex.

**scour and fill** A **\*sedimentary structure** characterized by a concave-upwards **\*erosion surface** cut into the underlying bed by a high-velocity **\*flow** of water, and filled by a **\*sediment**, which is usually coarse, during the waning stage of the flow that cut the scour.

**Scourian** A sub-**\*stage** of the **\*Lewisian**, from about 2600 to 2300 Ma ago, characterized by Scourian **\*dykes** and **\*gneiss** formation, and named after Scourie, north-west Scotland. See also INVERIAN.

**Scourian orogeny** A mountain-building episode that occurred about 2600 Ma ago, during the **\*Archaean**, prior to the **\*Laxfordian orogeny** but possibly representing its first stages. It is marked by NW–SE trending folds immediately to the south of the Laxford area, affecting part of the Lewisian **\*gneisses** in what is now the extreme north-west of Scotland.

**scour lag** A coarse-grained **\*sediment** deposited immediately above a scour surface (see SCOUR AND FILL).

**scree** Accumulation of coarse rock debris that rests against the base of an inland cliff. It is added to by the *\*weathering* and release of fragments from the cliff face. Scree is widely found in upland areas that are affected by past or present *\*periglacial* conditions, and in hot, rocky deserts.

**screw dislocation** See DISLOCATION.

**scroll bar** One of a series of long ridges of sand on a point *\*bar*, lying approximately parallel to the contours.

**ScS-wave** See SEISMIC-WAVE MODES.

**scud** Fragments of low cloud, usually nimbostratus, moving quickly beneath rain clouds. See also FRACTUS.

**scute** In fish, an enlarged, bony, dermal plate or scale.

**Scyphozoa (jellyfish)** (phylum *\*Cnidaria*) Class of marine, mainly *\*pelagic* medusoids, usually with four-part *\*radial symmetry*, in which the polyp stage is reduced or absent. Their fossil record is in general scanty, owing to the absence of hard parts, but jellyfish formed an important component of the *\*Precambrian* *\*Ediacaran* fauna. See also MEDUSINA MAWSONI.

**Scythian (Skythian)** An Early *\*Triassic* *\*epoch* in the Alps (251–245 Ma ago), preceded by the Late *\*Permian* and followed by the Middle Triassic.

**sea 1.** Large body of usually saline water which is smaller in size than an ocean. **2.** Chaotic waves generated by the action of the wind on the surface layers of the ocean. See also OCEAN WAVE; SWELL.

**sea-anemones** See ANTHOZOA.

**sea breeze** See LAND AND SEA BREEZES.

**sea-floor spreading** The theory that the ocean floor is created at the spreading (accretionary) *\*plate margins* within the ocean basins. *\*Igneous* rocks rise along conduits from the *\*mantle*, giving rise to volcanic activity in a narrow band along the *\*mid-ocean* ridges. As these cool, the basaltic *\*lavas* and *\*dykes* form the upper part of the *\*oceanic crust*, and the underlying *\*magma chamber* solidifies to form layer 3 of the oceanic crust. The newly formed oceanic crust spreads perpendicularly away from the

ridge, probably in response to mantle convective motions (see [PLATE TECTONICS](#)). As the [\\*basalts](#) originally cooled, they became magnetized by the ambient [\\*geomagnetic field](#). As this field reverses [\\*polarity](#), oceanic crust formed at different times is characterized by oceanic [\\*magnetic anomalies](#) that are parallel to the ridge at which they originally formed (Vine and Matthews, 1963). These anomalies allow the dating of the oceanic crust and the determination of its past relative motion. The creation of new ocean floor was implicit in the previous concept of [\\*continental drift](#), but is mainly characterized by the narrowness of the zone within which the new ocean floor is formed. It is now a fundamental concept within the plate-tectonic theory.

**sea fret** Popular local term for sea fog, common in spring and summer in Cornwall and on the south, east, and north-east coasts of England.

**sea ice** In polar regions the surface of the sea freezes, due to the low air and water temperatures: the product is known as 'sea ice'. It exists year-round in the central Arctic and in some Antarctic bays, extending in winter across the entire Arctic and far out to sea around Antarctica. As ice crystals form from sea water, so salt is excluded and eventually returned to the sea. Sea ice therefore contains no salt, except where pockets of sea water become trapped in the ice.

**seamount** Isolated, submarine mountain rising more than 1000 m above the ocean floor. The sharp, crested summits of seamounts are usually 1000–2000 m below the ocean surface. Seamounts are of volcanic origin. They are increasingly coming under study with modern methods of submarine acoustic imagery, e.g. GLORIA (Geological *L*Ong Range Inclined Asdic) [\\*side-scan](#) sonar. See also [GUYOT](#).

**sea pens** See [ANTHOZOA](#).

**seatearth** A [\\*clay](#)-rich [\\*fossil](#) soil, found immediately beneath a [\\*coal](#) seam, representing the soil in which the coal-forming vegetation grew.

**sea water, major constituents** 99.9% of dissolved material in sea water can be accounted for by eleven constituents: Na, Mg, Ca, K, Sr, Cl, SO<sub>4</sub>, HCO<sub>3</sub>, Br, BO<sub>3</sub>, and F, whose relative proportions are almost constant in all oceans regardless of [\\*salinity](#). The [\\*pH](#) of sea water is maintained at 8.0–

8.4 by a buffering system. Because the water is alkaline, most of the  $\text{Ca}^{2+}$  and  $\text{HCO}_3^-$  form insoluble  $\text{CaCO}_3$ . Aluminium and iron oxides coagulate to form **\*colloids** because of the high concentration of **\*electrolytes**, and sink to the sea bed, silica is taken up by organisms, and thus Cl, Na,  $\text{SO}_4$ , Mg, and K, for which no removal mechanism exists, account for almost all the dissolved material, at concentrations much higher than those found in rocks.

<i>Ion</i>	<i>Parts per thousand by weight</i>	<i>Percentage of dissolved material</i>
chloride, $\text{Cl}^-$	18.980	55.05
sodium, $\text{Na}^+$	10.556	30.61
sulphate, $\text{SO}_4^{2-}$	2.649	7.68
magnesium, $\text{Mg}^{2+}$	1.272	3.69
calcium, $\text{Ca}^{2+}$	0.400	1.16
potassium, $\text{K}^+$	0.380	1.10
bicarbonate, $\text{HCO}_3^-$	0.140	0.41
bromide, $\text{Br}^-$	0.065	0.19
borate, $\text{H}_3\text{BO}_3^-$	0.026	0.07
strontium, $\text{Sr}^{2+}$	0.008	0.03
fluoride, $\text{F}^-$	0.001	0.00
TOTAL	34.477	99.99

**SeaWiFS** An instrument (Sea-viewing Wide Field-of-view Sensor) carried on NASA's SeaStar spacecraft that was launched on 1 August 1997 into a **\*Sun-synchronous orbit** at a height of about 700 km. It orbited the Earth in 99 minutes, observing a swathe 2800 km wide. SeaWiFS monitored the colour of the ocean surface, providing data from which the concentration and movement of phytoplankton (see **PLANKTON**) could be calculated. The mission ended on 11 December 2010.



<https://oceancolor.gsfc.nasa.gov/SeaWiFS/>

- SeaWiFS Project.

**Secchi disc** A device used in a simple method for measuring the transparency of water. The disc is 20 cm across, and divided into alternate black and white quadrants. It is lowered into the water on a line until the

difference between the black and white areas just ceases to be visible, and this depth is recorded. The Secchi disc provides a convenient method for comparing the transparencies of water at different times.

**second arrival** The next coherent seismic event to follow the *\*first break*.

**secondary blasting** See POP-SHOOTING.

**secondary creep** The second stage of *\*creep*, characterized by viscous strains, in a material being deformed as a consequence of long-term, low *\*stress*.

**secondary crushing** See CRUSHING.

**secondary depressions** *\*Depressions* initiated as part of a 'family' or sequence of frontal systems that develop to the rear of the first depression of the series below the *\*polar-front jet stream*, which drags them along an approximately eastward path. See also FRONTAL WAVE.

**secondary enrichment** See SUPERGENE ENRICHMENT.

**secondary front** A *\*front* that forms along a frontal wave behind the primary front; there may be more than one secondary front.

**secondary geochemical differentiation** A theory concerning the processes which may have taken place in and on the Earth's *\*crust* and in the atmosphere, including the organic reactions leading to the development of the *\*biosphere*.

**secondary geochemical dispersion** The movement of elements at or just below the Earth's surface, which results from *\*weathering*, *\*erosion*, and deposition.

**secondary matrix** The *\*matrix* of a *\*sedimentary rock* that has been formed by the diagenetic (see DIAGENESIS) alteration of the grains.

**secondary migration** Movement of *\*hydrocarbons*, usually laterally, into and within the *\*reservoir rock* where they accumulate.

**secondary mineral** A *\*mineral* formed by the subsolidus alteration of a pre-existing *\*primary mineral* in an *\*igneous* rock. Minerals which have crystallized from a *\*magma* are stable only at high temperature and can readily alter to low-temperature, secondary minerals when a fluid, e.g.

water, is introduced into the rock system. The fluid acts as a catalyst to initiate the alteration reaction. Most secondary minerals are hydrated **\*silicates**. A typical example is the alteration of primary **\*olivine** to secondary **\*chlorite** and **\*serpentine**.

**secondary porosity** The forms of **\*porosity** which develop in a rock during and after **\*consolidation**, by the selective dissolution or alteration of mineral **\*grains** and **\*cements**, or by the fracturing of the rock by tectonic processes (see **TECTONISM**). See **CHOQUETTE AND PRAY CLASSIFICATION**.

**secondary quartz** **\*Quartz** formed as a **\*cement** during **\*diagenesis**.

**secondary recovery methods** In the petroleum industry, techniques by which recovery has been increased to about 50% since the 1940s, including the injection of natural gas above the oil in a **\*reservoir** to force the oil downwards, or pumping water in below the oil to force it upwards.

**secondary sedimentary structure** A structure formed by the precipitation of minerals in the pores of a sedimentary rock during or following its consolidation, or by chemical replacement of some of its constituents. Compare **PRIMARY SEDIMENTARY STRUCTURE**.

**secondary wave** See **S-WAVE**.

**second derivative** Acceleration due to gravity ( $g$ ) is the first derivative of the gravity potential field. Where  $g$  varies with horizontal distance (due to anomalies), then the gradient in the direction of the variation is the second derivative. Such computations exaggerate noise, as well as highlighting maxima and minima in the gravity field.

**secular variation** Any long-term variation over a period of the order of 10 to 20 years or longer. In **\*geomagnetism**, a variation that operates on a timescale greater than one year, but excluding the variation associated with the sunspot cycle.

**SEDEX** See **SEDIMENTARY EXHALATIVE PROCESSES**.

**Sedgwick, Adam** (1785–1873) Woodwardian Professor of Geology at Cambridge, where he modernized the teaching of geology. He is best known for unravelling the stratigraphy of North Wales (the **\*Cambrian** system) which resulted in a quarrel with R. I. **\*Murchison** over the

boundary between the Cambrian and **\*Silurian** systems. He made important contributions to structural geology, distinguishing between stratification, jointing, and **\*slaty cleavage**.

**sedigraph** An instrument that calculates the size of sediment particles by measuring the attenuation of a finely collimated X-ray beam through a settling suspension as a function of height above the base of the **\*sedimentation tube** and time.

**sediment** Material derived from pre-existing rock, from **\*biogenic** sources, or precipitated by chemical processes, and deposited at, or near, the Earth's surface.

**sedimentary basin** A subsiding area of the Earth's **\*crust** which permits the net accumulation of **\*sediment**.

**sedimentary cycle** A cycle which comprises the **\*weathering** of an existing rock, followed by the **\*erosion** of **\*minerals**, their transport and deposition, then burial. First-cycle **\*sediments** are characterized by the presence of less resistant minerals and rock fragments. If this material is reworked through a second cycle, the less resistant minerals will be eliminated, or altered to more stable products. The more sedimentary cycles that a sediment has passed through, the more mature it will become and it will be dominated by well-rounded, resistant minerals (see **RESISTANT MINERAL**).

**sedimentary environments** Environments in which **\*sediments** are deposited on land include **\*alluvial fans**, river beds, lake beds, sand **\*dunes**, marshes, **\*deltas**, **\*barrier beaches**, **\*lagoons**, and **\*tidal flats**. Sediments also accumulate on **\*reefs**, continental shelves, **\*continental rises**, **\*continental slopes**, and **\*abyssal plains**. The type of sedimentary deposit varies according to the type of rock. Knowledge of the sedimentary processes active today, the environments in which they occur, and the types of rock they produce facilitate the interpretation of ancient deposits.

**sedimentary exhalative processes (SEDEX)** Processes associated with the upwelling of mineralizing fluids into submarine sedimentary environments, whereby **\*mineral** deposits, usually of base-metal sulphides, are formed. See **'BLACK SMOKERS'**.

**sedimentary mélange** See **OLISTOSTROME**.

**sedimentary rock** Rock formed by the deposition and compression of mineral and rock particles, but often including material of organic origin and exposed by various agencies of **\*denudation**. Sedimentary rocks may be classified as terrigenous (i.e. derived from the breakdown of pre-existing rocks exposed on the land), organic (i.e. produced either directly or indirectly by organic processes such as shell production or **\*peat** formation), chemical (i.e. produced by precipitation from water, e.g. some **\*carbonates** and all **\*evaporites**), or volcanogenic (**\*pyroclastic**, e.g. **\*tuffs** and **\*bentonites**). They may also be described according to their chemical properties and behaviour and their environmental deposition, and each scheme complements the others.

**sedimentary rock classification** The categorization of **\*sedimentary rocks** according to the way they formed, as either **\*clastic** or chemical. Clastic rocks consist of fragments of weathered (see **WEATHERING**) rock cemented together (see **CEMENTATION**). Chemical sedimentary rocks form when **\*minerals** dissolve in water and are later deposited as solids. The tables show a selection of both categories.

### Clastic Sedimentary Rocks

Rock	Type of sediment	Texture
Conglomerate	Gravel with rounded fragments	Coarse (>2 mm)
Breccia	Gravel with angular fragments	Coarse (>2 mm)
Sandstone	Sand	Medium (0.06–2 mm)
Siltstone	Mud	Fine (4–62.75 µm)
Shale	Mud	Very fine (<4 µm)

### Chemical Sedimentary Rocks

Rock	Composition	Texture
Crystalline limestone	Calcite (CaCO <sub>3</sub> )	Coarse to fine crystalline
Fossiliferous limestone	Calcite (CaCO <sub>3</sub> )	Visible shell fragments
Chalk	Calcite (CaCO <sub>3</sub> )	Microscopic shells and clay
Chert	Quartz (SiO <sub>2</sub> )	Very fine crystalline
Gypsum	Gypsum (CaSO <sub>4</sub> .2H <sub>2</sub> O)	Fine to coarse crystalline
Rock salt	Halite (NaCl)	Fine to coarse crystalline
Bituminous coal	Organic matter	Fine

**sedimentary structure** The external shape, the internal structure, or the forms preserved on bedding surfaces, generated in *\*sedimentary rocks* by sedimentary processes or contemporaneous *\*biogenic* activity. Internal sedimentary structures include: those formed by physical depositional processes (*\*cross-stratification*, flat bedding (see *PLANE BED*), *\*lamination*, and *\*heterolithic* structures); those due to post-depositional deformation (convolute bedding, *\*slump structures*, dish and pillar structures, flame structures, *\*ball* and pillow structures, etc.); those caused by organic disturbance (*\*bioturbation*, *\*trace fossils*); or by post-depositional chemical disturbance (*\*enterolithic structures*, collapse and solution structures, *\*concretions*, etc.). Structures preserved on the tops of beds include: those formed by depositional processes (*\*ripple* marks, primary current lineations); erosional structures (*\*flutes* and scour marks, see *SCOUR AND FILL*); structures caused by the transportation of an object over the bed (*\*tool marks*); and other features such as *\*desiccation* and *\*syneresis cracks*, *\*sand volcanoes*, adhesion ripples and warts, rain prints, and biogenic traces and *\*trails*. Structures preserved on the bases of beds (sole marks) include *\*load casts*, the casts of flutes, trails and tool marks, and the fill of erosional scours. The external form of sedimentary units (sheet-like, *\*channel-fill*, *\*reef* or mound (see *MUD MOUND*), lenticular, etc.) is a function of the depositional environment and sometimes of post-depositional *\*compaction*.

**sedimentation, rate of** See *SUBSIDENCE*.

**sedimentation coefficient** Measure of the rate of sedimentation of a molecule or particle; it is equal to the velocity per unit centrifugal field (acceleration), and is measured in *\*Svedberg units*.

**sedimentation tube** Apparatus designed to determine rapidly the grain-size distribution of *\*sand*-sized *\*sediment*. The sedimentation-tube method is based on the principle of *\*Stokes's law* that particles of different sizes, shapes, and densities will settle through a column of fluid at different settling velocities. Sediment is introduced into the top of the settling tube, and the time is recorded for the various sediment fractions to settle to the bottom of the tube.

**sedimentology** The scientific study, interpretation, and classification of sediments, sedimentary processes, and sedimentary rocks.

**seed plants** The group that comprises the **\*gymnosperms** and **\*angiosperms**, plants distinguished by their production of seeds rather than spores. They arose in the **\*Devonian** from **\*pteridophyte** forebears, probably of a heterosporous character (i.e. bearing microspores and megaspores (see **SPORE**) on the same plant). These early seed plants were gymnosperms. Angiosperms are not certainly represented in the fossil record until about 130 Ma ago (Early **\*Cretaceous**).

**seep** See **SPRING**.

**seepage** The slow but often steady flow of water between one water body and another. As a term, it is often used to describe leakage to underlying **\*aquifers** through stream beds or the emergence of **\*groundwater** into a stream channel, but it may also relate to flow between different aquifer units.

**seepage velocity** The velocity of **\*groundwater** calculated from **\*Darcy's law**. It is not the actual velocity of the water in the pores, but the apparent velocity through the bulk of the porous medium. Actual velocity is higher than seepage velocity by a factor which combines the effects of porosity and the tortuosity of the actual flow path among and around the mineral grains.

**seiche** A stationary or **\*standing** wave in an enclosed body of water, e.g. a bay or lake. Seiches are usually the product of intense storm activity.

**seif dune** Linear **\*dune** consisting of curved, sword-like components, and found in hot deserts. Typically it is developed by the elongation of a **\*barchan** arm, and built up by winds blowing from two principal directions.



Seif dune

**seism-** From the Greek *seismos* meaning ‘earthquake’, a prefix meaning ‘pertaining to earthquakes’.

**seismic anisotropy** The situation in which the speed at which a **\*seismic wave** travels varies according to the direction of travel. While most **\*minerals** are anisotropic (see **ANISOTROPY**) the mineral grains forming a rock are randomly aligned, so the anisotropies usually cancel out, leaving the rock isotropic at the wavelengths of seismic waves (metres to hundreds of metres). **\*Foliation** tends to align mineral grains, however, so large masses of foliated rocks are often anisotropic. Rocks such as **\*shale** are also seismically anisotropic because they are composed of elongate grains that align during the deposition that eventually forms the rock.

**seismic blind zone** A layer which cannot be detected by seismic refraction methods because it is too thin or its velocity is lower than that in overlying strata. See **HIDDEN LAYER**.

**seismic gap** **1.** An area within a known active **\*earthquake** zone within which no significant earthquakes have been recorded. It is not always clear whether this gap represents a zone where gradual motion takes place continually so there is no strain accumulation, or where motion is locked and strain is accumulating. **2. (shotpoint gap)** In split-spread seismic **\*reflection** shooting, the distance between the shot point and the nearest **\*groups** of **\*geophones**, which is larger than that between subsequent geophone groups at larger **\*offsets**. **3. (inter-record gap)** A blank space on

a magnetic tape which signals the end of one block of seismic data (one seismic record) and heralds the start of another. It is used to facilitate the transfer of magnetically recorded data into a computer system.

**seismicity** The likelihood that an *\*earthquake* will be felt in a particular area; i.e. earthquakes are frequent in zones of high seismicity (e.g. Japan and California).

**seismic margin** See ACTIVE MARGIN.

**seismic moment** A measure of the size of an *\*earthquake*, based on the concept of rupture and slippage along a fault plane as a rotational motion about a point on the fault. It is given by:  $M_0 = \mu Ad$ , where  $\mu$  is a constant varying according to the material,  $A$  is the area of the ruptured fault, and  $d$  is the average slip.

**seismic record (seismogram)** The output of a seismic recording system (*\*seismograph*), e.g. a paper or film record, showing the seismic *\*wobble traces*, usually for a single *\*shot \*spread*. In *\*refraction surveys*, many shots into one spread may be summed by a signal-enhancement seismograph to produce a single record. When records are processed they can be placed side by side along a profile to form a 'seismic section'.

**seismic reflection** See REFLECTION.

**seismic-reflection profiling** See PROFILING.

**seismic refraction** See REFRACTION.

**seismic section** See SEISMIC RECORD.

**seismic stratigraphy (seismostratigraphy)** The study and interpretation of information obtained by seismic-*\*reflection \*profiling* in order to construct subsurface *\*stratigraphic cross-sections*. Analysis of seismic reflections on a seismic section (see SEISMIC RECORD) can identify buried stratal surfaces that, when traced laterally and continuously, represent surfaces of synchronous deposition or their correlative *\*unconformity* surfaces. The character of a reflection may vary as the seismic profile moves across a *\*facies* boundary, but the continued presence of the reflection is of *\*chronostratigraphic* significance. More detailed information regarding the age and *\*lithology* of the subsurface strata may be gathered by means of

geophysical **\*well logging**. *See also* CHRONOSTRATIGRAPHIC CORRELATION CHART; DEPOSITIONAL SEQUENCE.

**seismic survey** The exploration of a subsurface geologic structure by means of **\*seismic waves** which are generated artificially. **\*Reflection** profiling is the most common surveying method, and **\*refraction surveys** are particularly important in land surveys for **\*static corrections**. Small seismic refraction surveys are commonly carried out for **\*engineering geophysics** site investigations.

**seismic tomography** A range of methods in which the subsurface is divided into a box-grid whose elements are illuminated by seismic rays and the physical character of each element computed. The results are displayed as colour-coded contour maps of subsurface planes. In **\*borehole \*tomography**, two holes are used; a source is moved up one while detectors record along the length of the other. By using a succession of **\*shot** positions, most of the box elements in the plane linking the two boreholes are illuminated satisfactorily. Seismic tomography is used to study geologic structures in detail. It has been extended to investigate the **\*mantle** and, increasingly, the effectiveness of extraction techniques in hydrocarbon reservoirs.

**seismic velocity** The speed with which an elastic wave propagates through a medium. For non-dispersive **\*body waves**, the seismic velocity is equal to both the **\*phase** and **\*group velocities**; for dispersive **\*surface waves**, the seismic velocity is usually taken to be the phase velocity. Seismic velocity is assumed usually to increase with increasing depth and when measured in a vertical direction it may be 10–15% lower than when measured parallel to strata. *See* ANISOTROPY.

**seismic wave** A packet of elastic strain energy which travels away from a seismic source, e.g. a **\*shot** or **\*earthquake**. *See also* BODY WAVE; SURFACE WAVE.

**seismic-wave modes** The conventional notation ascribed to **\*seismic waves** on the basis of their travel times from their **\*earthquake** sources. Letters are used to designate the type of wave along the different portions of its travel path: P and S refer to **\*P-waves** and **\*S-waves** that do not travel through the Earth's **\*core**; K and I refer to P-waves only, which travel through the core

and inner core respectively. The letter J refers to S-waves which are generated by the conversion of incident P-waves at the boundary between the inner and outer cores, and which travel only in that mode within the solid inner core. As they propagate towards the Earth's surface, J-waves are reconverted back into P-waves at the inner core boundary. A repetition of the letters (e.g. PP, SS) indicates that the waves have been reflected at the Earth's surface; reflection at the outer edge of the core is indicated by 'c' (e.g. PcP, PcS). For example, the wave PKIKP travels from the surface, through the \*mantle to the outer core, into the inner core, and then back to the surface; the wave PPP has travelled only in the \*crust and upper mantle, and has been reflected twice off the surface.

**seismic zone** A region of high \*seismicity.

**seismogram** See SEISMIC RECORD.

**seismograph** A device which records seismic information. Usually the term is used to describe the entire system, including amplifiers and means for filtering data and transferring them to magnetic tape or computer disk, but occasionally it describes only \*geophones. An enhancement seismograph can sum successive \*hammer impacts or \*shots fired into one geophone \*spread in order to enhance the signal-to-\*noise ratio, and is commonly used in \*engineering geophysics site investigations.

**seismology** The study of elastic (seismic) waves and how they are produced. Global seismology is the study of \*seismic waves from \*earthquakes (and to a lesser extent nuclear explosions), to investigate the structure of and processes within the Earth. In exploration seismology, artificially generated seismic waves are used in the search for resources (e.g. hydrocarbons, etc.) and the study of the Earth's surface and near-surface. Planetary seismology is the use of seismic waves to investigate the structure of and processes within planets and natural \*satellites in the \*solar system.

**seismometer** 1. A device used to detect \*seismic waves originating from \*earthquakes. 2. In exploration \*seismology, a \*geophone.

**seismostratigraphy** See SEISMIC STRATIGRAPHY.

**seismotectonics** The study of recent \*earthquakes, especially with their spatial and temporal distribution and magnitudes, and their relationship

with *\*plate tectonics*. See also NEOTECTONICS.

**Selandian** A *\*stage* of the *\*Palaeocene* epoch, 61.7–58.7 Ma ago, preceded by the *\*Danian* and followed by the *\*Thanetian*.

**SELENE (KAGUYA)** A mission that was launched from the Tanegashima Space Center on 14 September 2007 by the *\*Japan Aerospace Exploration Agency*, that obtained data on the origin and evolution of the Moon and developed technologies needed for future lunar exploration. Its name is derived from SELEnological Engineering Explorer. SELENE orbited at about 100 km and released two small satellites into polar orbits, Okina and Ouna, which entered elliptical orbits at  $100 \times 2400$  km and  $100 \times 800$  km, respectively. The main orbiter was manoeuvred into a controlled crash into the Moon on 11 June 2009.



[http://www.jaxa.jp/projects/sat/selene/index\\_e.html](http://www.jaxa.jp/projects/sat/selene/index_e.html)

- A JAXA mission to study the Moon's origins and evolution.

**selenite** See GYPSUM.

**selenizone** See ARCHAEOGASTROPODA.

**selenology** The astronomical study of the *\*Moon*. The term is derived from the Greek word *selene*, meaning 'moon'.

**self diffusion** See DIFFUSION (2A).

**self-exciting dynamo** A dynamo that produces a magnetic field around itself whereby motions of an electrical conductor, carrying magnetic lines of force, generates further current, eventually resulting in a stable external magnetic field. It is generally considered that the *\*geomagnetic field* is produced by two self-exciting dynamos, the interaction of which results in *\*reversals* of geomagnetic polarity.

**self-potential method** See SPONTANEOUS-POTENTIAL METHOD.

**self-potential sonde (spontaneous potential sonde, SP sonde)** A *\*well-logging* instrumental package that measures the electrochemical activity of the rocks within a *\*borehole*. Formations yielding a high and true SP log response include *\*clay-rich* *\*sediments* and *\*sulphide* minerals. The SP

method cannot be used for logging wells in offshore areas, where saltwater-based drilling muds are used.

**self-reversal** The ability of certain *\*ferromagnetic* materials to acquire a thermal remanence in the opposite direction to the ambient magnetic field. Usually this requires the interaction of two or more magnetic lattices, or of minerals with different *\*Néel* or *\*Curie temperatures*.

**SEM** See SCANNING ELECTRON MICROSCOPE.

**Senecan** A *\*stage* (388–370 Ma ago) in the *\*Devonian* of N. America, underlain by the *\*Erian*, overlain by the *\*Chautauquan*, and comprising the Fingerlakian and Cohoktonian *\*stages*. It is roughly contemporaneous with the *\*Frasnian* of Europe.

**Senonian** A European *\*epoch* within the Late *\*Cretaceous* which is dated at 89.3–65.5 Ma ago (Int. Commission on Stratigraphy, 2004), preceded by the *\*Gallic*, followed by the *\*Danian*, and comprising the *\*Coniacian*, *\*Santonian*, *\*Campanian*, and *\*Maastrichtian* stages. Some authors do not include the Maastrichtian age within the Senonian.

**Sensitive High Resolution Ion MicroProbe (SHRIMP)** A device for dating granite *\*batholiths in situ* by analysing zircon grains taken from several samples.

**sensitive tint** See ACCESSORY PLATE.

**sensitivity** 1. The consistency of *\*clay* as this is affected by *\*remoulding*. The effect depends on the type of clay and the amount of pore water. In a sensitive clay, *\*shear strength* is decreased dramatically on remoulding when moisture content remains constant. Sensitivity is measured as the ratio of the *\*unconfined compressive strength* to the strength in the remoulded state at the same water content. 2. In chemical analysis, the smallest change in concentration which can be discriminated by the analytical method.

**Sentinel** See COPERNICUS.

**SEOSat/Ingenio** A Spanish optical high-resolution imaging satellite mission to supply multispectral land images to Spanish agencies and to European users of *\*Copernicus*. The satellite is scheduled to launch in 2019 into a Sun-synchronous circular orbit at an altitude of 670 km.



<http://www.aerospace.sener/products/seosat-ingenio-spanish-earth-observation-satellite>

- SEOSAT/INGENIO: Spanish Earth Observation Satellite.

**septa** See RUGOSA; TABULATA; SEPTUM.

**septarian nodule** A *\*concretion*, roughly spheroidal in shape, usually of *\*clay* *\*ironstone*, and characterized by an internal structure of angular blocks separated by radiating mineral-filled blocks. The mineral filling the cracks is usually *\*calcite*. The structure results from the formation of a hard exterior to the nodule due to the development of an aluminous *\*gel* on the exterior, followed by dehydration of the colloidal mass (see COLLOID) in the interior, leading to cracking and subsequent mineral infilling of the radiating pattern of cracks.

**septomaxilla** A bone at the front of the upper jaw in reptiles; monotremes (*\*Monotremata*) are the only mammals in which this bone maintains its separate identity. See OBDURODON.

**septum** (*pl. septa*) A cross wall or partition. **1.** A morphological term used particularly with reference to the *\*Cephalopoda*, whose shells are divided internally into a series of chambers (camerae) by septa which are generally concave towards the anterior. There is an opening (foramen) in each septum and this is usually bordered by a collar (septal neck). Septal necks are forward-pointing (prochoanitic) in ammonoids (*\*Ammonoidea*) and backward-pointing (retrochoanitic) in nautiloids (*\*Nautiloidea*). In many ammonoids the septum becomes fluted and there has been much discussion concerning the function of the complex fluting. **2.** The radially arranged *\*plates* that occur in the *\*corallum* of corals. The first-formed septa are called 'prosepta' and are usually larger than the metasepta, which occur between them.

**sequence stratigraphy** The analysis, within the framework of a *\*chronostratigraphic scale*, of depositional units that are related to one another genetically (i.e. they were formed by similar processes). There are two schools: the Exxon Production Research (EPR) *\*depositional sequence model* and the *\*genetic stratigraphic sequence model* (proposed by W. E.

Galloway in 1989), differing mainly in the type of boundaries by which they define the strata.

**serac** Ice pinnacle found on the surface of a **\*glacier** and resulting from tensional failure in the more rigid upper crust. This is due to the stretching that occurs when a glacier moves over a convex-up slope, spreads out over a plain, or passes round a bend in its valley.

**SERB** See SOLAR IRRADIANCE AND EARTH RADIATION BUDGET.

**serein** The fall of rain from an apparently clear sky. The phenomenon may be explained by the evaporation of cloud particles following the formation of rain droplets, or by the movement of cloud away from the overhead position as the rain approaches the ground.

**sericite** A white variety of **\*muscovite** or paragonite and member of the **\*phyllosilicates** (sheet silicates) with the formula  $K_2Al_4[Si_3AlO_{10}]_2(OH)_2$  formed from the alteration of **\*feldspar** by either hydrothermal alteration or later-stage **\*weathering**. It appears as small scales or flakes or compact aggregates which give the feldspar a 'cloudy' appearance in **\*thin** section.

**series** The major subdivision of a **\*system**, and the chronostratigraphic equivalent of an **\*epoch**. It denotes the layers of strata or the body of rock formed during one epoch. A series may itself be divided into **\*stages**. When used formally the initial letter of the term is often capitalized, e.g. Lower **\*Cretaceous** Series.

**serir** Veneer of mixed sand and gravel mantling a Saharan plain and transported originally by **\*sheetwash** and **\*braided-stream** activity. Subsequently it was weathered and modified under more arid conditions.

**serpens** (*pl. serpentes*) A sinuous, undulating feature on the surface of an extraterrestrial body.

**serpentes** See SERPENS.

**serpenticone** See INVOLUTE.

**serpentine** A group of **\*minerals** belonging to the 1:1 group of **\*phyllosilicates** (sheet silicates) with the composition  $Mg_6[Si_4O_{10}](OH)_8$  and including the minerals chrysotile (the asbestiform variety), lizardite, and antigorite; sp. gr. 2.55–2.60; **\*hardness** 2.0–3.5; **\*monoclinic**; chrysotile

is fibrous whereas lizardite and antigorite occur as flat **\*tabular** crystals or **\*massive**; various shades of green, also brown, grey-white, or yellow; **\*greasy** to **\*waxy** **\*lustre**, occasionally **\*silky**; formed from altered **\*olivine** and **\*orthopyroxene**. It results from the alteration of ultramafic (see **ULTRABASIC**) rocks either by **\*hydrothermal** action at a late stage or by alteration during **\*metamorphism**, chrysotile forming first and then altering to antigorite; it is a constituent of ophicalcites, a serpentine-**\*calcite** rock derived from the dedolomitization (see **DEDOLOMITE**) of a siliceous **\*dolomite**. It is used extensively as a facing stone and for ornament; chrysotile has been used as a source of commercial **\*asbestos** in Thetford, Canada. Lizardite occurs on the Lizard Peninsula, Cornwall, UK.

**serpentine barrens** Impoverished vegetation, often dominated by scrub or heath, associated with **\*serpentine** rocks. On **\*weathering** these rocks release an excess of magnesium into the soil, and this often inhibits the development of the natural climax in the areas concerned.

**serpentinite** Altered rock formed from an **\*ultrabasic** precursor by low temperature and water interaction. Such rocks are compact, variously coloured, and may have considerable ornamental value. They consist mainly of hydroxyl-bearing magnesium silicates formed from original **\*olivine** and **\*pyroxenes**.

**serpentinization** The process whereby high-temperature **\*primary** **\*ferromagnesian minerals** in an **\*igneous** rock undergo alteration to a member of the **\*serpentine** group of minerals. The process is initiated by the introduction of low-temperature water into the rock system, the water acting as a catalyst for the reaction in which the high-temperature primary ferromagnesian minerals are converted to low-temperature, **\*secondary**, serpentine-group minerals. Serpentinization is extremely common in **\*ultrabasic** rocks, especially those found in **\*ophiolites** where the entire rock may be converted to serpentine-group minerals, forming a **\*serpentinite** rock. Where the original mineral assemblage can be inferred from relict minerals or **\*pseudomorph** textures, the original mineral name can be added to the rock name. Thus a serpentinized **\*dunite** would be an **\*olivine-serpentinite**.

**Serpukhovian** A **\*stage** of the Late **\*Mississippian** (326.4–318.1 Ma ago), comprising the **\*Pendleian**, **\*Arnsbergian**, **\*Chokierian**, and **\*Alportian**

stages. It is preceded by the **\*Visean** and followed by the **\*Bashkirian** (Early **\*Pennsylvanian**). It is roughly contemporaneous with the upper **\*Chesterian** (N. America) and **\*Namurian A** (western Europe).

**Serravallian** A **\*stage** in the Middle **\*Miocene \*epoch** preceded by the **\*Langhian**, followed by the **\*Tortonian**, and dated at 13.65–11.608 Ma ago. It is roughly contemporaneous with the Helvetian (Europe), upper **\*Luisian** and lower **\*Mohnian** (California), **\*Lillburnian** (New Zealand), and **\*Bairnsdalian** (Australia). The **\*stratotype** is in the Scrivia Valley, Italy.

**sesquioxides** General term for the hydrated oxides and hydroxides of iron and aluminium.

**sessile** 1. Lacking a stalk. 2. Attached to a substrate; non-motile.

**seston** Particulate matter suspended in sea water.

**seta** Stiff, hair-like or bristle-like structure.

**Setebos (Uranus XIX)** A lesser satellite of Uranus with a radius of 24 km and a visual **\*albedo** of 0.07.

**settlement** 1. Gradual subsidence of a structure, caused by the compression of soil below **\*foundation** level. Normally a uniform amount of settlement can be accepted but damage may occur when different parts settle disproportionately. See **DIFFERENTIAL SETTLEMENT**. 2. In mining, the lowering of overlying strata due to extraction of material.

**settling lag** The period of time it takes for a fine-grained particle to settle through a water body after the cessation of transportation in suspension. The settling time will be dependent on the **\*particle size** and **\*particle shape**, and governed largely by Stokes's law of settling.

**sexual dimorphism** Phenomenon of morphological differences (besides primary sexual characters) that distinguish the males from the females of a species. For example, male deer often have larger antlers than females, and the males of many birds have differing plumage (often more brightly coloured). Sexual dimorphism is known to have been common in ammonites (**\*Ammonoidea**), and many **\*fossils** originally thought to have represented separate species are now recognized as dimorphs within one species (e.g. the **\*Jurassic** *Kosmoceras jason* and *K. gulielmi* are dimorphs within *K. jason*). See also **DIMORPHISM**.

**Seymouria** One of the more problematic of fossil *\*amphibians*, possessing a combination of amphibian and reptilian characters so that it has been referred to both groups on different occasions. One of the features separating it and its relatives from other amphibians is the presence of a large, forward-extended optic notch (the primitive site of the ear drum). A medium-sized amphibian, it existed during the Lower *\*Permian* in N. America.

**sferics** Natural electromagnetic signals, generated in the atmosphere by discharges of lightning, which propagate around the Earth between its surface and the *\*ionosphere*. The word is derived from ‘atmospherics’.

**S-fold** An asymmetrical *\*parasitic fold* whose approximately S-shaped profile, when observed down the *\*plunge* of the *\*fold axis*, indicates its position on the right limb of the major *\*anticline*, but not on the *\*syncline*. See MINOR FOLD.

**SG** The older abbreviation for *\*specific gravity* (sp. gr.).

**SGCS** Standard Global Chronostratigraphic (Geochronologic) Scale. See STANDARD STRATIGRAPHIC(AL) SCALE.

**shade temperature** The temperature of the air, conventionally measured in a standard shelter or screen that protects the thermometer from rain and from direct sunshine, but that allows the free passage of air.

**shadow test** See HALF-SHADOW TEST.

**shadow zone** 1. A region of the subsurface from which seismic *\*reflections* cannot be detected because their ray-paths do not emerge to the surface. 2. A zone over the Earth’s surface in which *\*P-waves* and/or *\*S-waves* generated by an *\*earthquake* are detected only weakly, or are absent, because of *\*refraction* within the various layers deep within the Earth, and especially in association with the Earth’s *\*core*. Seismic signals refracted from the *\*mantle* are usually poorly discernible at a range of up to about 10°. At a range of 103–142° from the earthquake P- and S-waves are observed only very weakly; between 110° and 142° some very weak P-waves can be measured and these are thought to have been refracted from the inner core. Between 142° and 180° from an earthquake no S-waves are recorded because the waves cannot travel through the liquid outer core.

**shaft well** A well sunk by a mining technique rather than by drilling. Normally it is larger in cross-section than a drilled well.

**shale** Fine-grained, fissile, **\*sedimentary rock** composed of **\*clay**-sized and **\*silt**-sized particles of unspecified mineral composition. The noun may be qualified by an adjective (e.g. **\*black shale**, **\*paper shale**, and **\*oil shale**).

**shale line** The line on an electrical log marking the usual **\*apparent resistivity** for a **\*shale**.

**shallowing-upward carbonate cycle** A stratigraphic sequence found on platforms, shelves (see **SHELF**), and some former lake beds, where carbonates have been deposited in progressively shallower water. The sequence develops where the rate of carbonate deposition exceeds the rate at which the receiving basin sinks, so the sediment surface repeatedly rises towards the water surface, while at the same time the deposit progrades (see **PROGRADATION**).

**shamal** Regional north-westerly wind which brings hot, dry conditions in summer, especially in June and July, to Iraq, Iran, and the Arabian Peninsula. It blows with great force during the day.

**shape fabric** A **\*fabric** in which the orientation of mineral grains is determined by their shape, so the fabric may be modelled by the shapes of various **\*strain ellipsoids**. Three **\*tectonite** fabrics are commonly used to demonstrate shape fabrics. L-tectonites, produced by uniaxial stretching with planar fabrics, correspond to prolate ellipsoids (see **PROLATE UNIAXIAL STRAIN**). S-tectonites, produced by flattening with planar fabrics, correspond to oblate ellipsoids (see **OBLATE UNIAXIAL STRAIN**). L-S-tectonites, produced by stretching and flattening with intermediate fabrics, correspond to triaxial ellipsoids.

**sharp sand** **\*Sand** composed of angular, not rounded, grains, with little foreign material; it is used in mortar.

**shatter cones** Striated, conical, nested fracture pattern that has apparently occurred in response to shock waves of the magnitude (2–25 GPa) generated by a **\*meteorite** impact. Individual cone shapes may vary from less than a centimetre to several metres in size. The apices of the cones tend to point towards the centre of impact. Shatter cones were first discovered by


R. S. *\*Dietz* in the 1940s and have been observed at many postulated impact *\*crater* sites, e.g. the Sudbury Structure, Ontario, Canada; the Ries and Steinheim Basins, southern Germany; and Gosses Bluff, Australia.

**shear box** Laboratory equipment used to test the *\*shear strength* of soil or rock. The upper and lower halves of a sample are subjected to horizontal shear by lateral pressure, and the effects measured.

**shear direction** See *SHEAR PLANE*.

**shear modulus ( $\mu$ , Lamé's constant  $\mu$ , rigidity modulus)** The ratio of the shearing stress ( $\tau$ ) to the resultant shear strain component ( $\tan \theta$  in the case of simple shear). It can also be given by  $\mu = \frac{1}{2}E/(1 + \sigma)$  where  $E$  is *\*Young's modulus* and  $\sigma$  is *\*Poisson's ratio*.

**shear plane** A plane that is parallel to the walls of a *\*shear zone* and contains the shear direction (the displacement vector).



Shear plane and zone

**shear strain** See *ANGULAR SHEAR STRAIN*.

**shear strength** The internal resistance of a material to *\*shear stress*; this varies depending on temperature, *\*confining pressure*, shape, size, loading rate, and amount of pore fluid present. It is measured as the maximum shear stress on an original cross-sectional area that can be sustained. In soils, it is the maximum resistance of a soil to shearing forces under specified conditions. The peak shear strength is the highest stress sustainable just prior to complete *\*failure* of a sample under load; after this, stress cannot be maintained and major strains usually occur by displacement along failure surfaces. The residual shear strength is the ultimate strength along a surface

or parting in soil or rock after shearing has occurred. For material not previously sheared there is a rapid decline in strength with increasing shear until the residual shear strength is reached.

**shear stress ( $\tau$ )** The *\*stress* which acts parallel to a plane on which a force has been applied. Shear stresses tend to promote sliding along a plane. Conventionally they are designated by the symbol  $\tau$ ;  $+\tau$  indicates a left-handed shear sense,  $-\tau$  a right-handed shear sense.

**shear wave** See *S-WAVE*.

**shear zone** A region, narrow compared to its length, within which rocks have undergone intense deformation. There are two end-members. Brittle shear zones (*\*faults*), marked by a surface of rupture, are a common feature of higher crustal levels. In ductile shear zones deformation is continuous and characterized by high ductile *\*strains* due to the rocks having formed under high temperatures and pressures at deeper crustal levels. Both brittle and ductile shear zones may occur as subparallel or *\*conjugate* sets.

**sheath fold** A *\*fold* with extreme curvature at the *\*hinge*, often close to 180°.

**sheep-walk** See *TERRACETTE*.

**sheet flood** See *FLASH FLOOD*.

**sheet lightning** The appearance of lightning flashes as a general flash when cloud cover causes the diffusion of their light. It may be due to flashes between two clouds seen through an intervening cloud, or flashed between separate charges within a single cloud.

**sheet sand (sand sheet)** A flat or gently undulating area of sand surrounding a *\*dune*.

**sheet silicate** See *PHYLLOSILICATE*.

**sheetwash** A geomorphological *\*process* by which a thin, mobile sheet of water flows over the surface of a hillslope and may transport the surface *\*regolith*. It is important in semi-arid regions, and may also be significant in temperate zones if the vegetation cover has been removed. See *RAIN-WASH*.

**Sheinwoodian** A *\*stage* (428.2–426.2 Ma ago) of the Early *\*Silurian*, underlain by the *\*Telychian* and overlain by the *\*Homerian*.

**shelf** A gently sloping or near-horizontal, shallow, marine platform. Horizontal shelf areas, particularly in carbonate-dominated areas, are referred to as ‘platforms’ if they have precipitate margins; more uniformly sloping shallow marine carbonate areas are termed ‘ramps’.

**shell beak** In *\*Bivalvia*, the oldest part of the shell, located near the *\*hinge*.

**shell bed** A *\*lag deposit* of shells and skeletal remains that are too heavy to be transported by sea currents, but from which the currents have winnowed away the sedimentary particles.

**shell structure** Shell growth is similar in *\*Mollusca* and *\*Brachiopoda*. The shell is divided into an outer periostracum, which is organic, and an inner ostracum. Both parts are secreted by the *\*mantle* and shells grow by marginal increments which are commonly visible on the exterior as growth lines. Shell thickening takes place and in some groups accessory or additional shell material may be laid down. External ornament, e.g. ribs, often reflects the shape of the mantle edge or structures contained within it and quite complex patterns of ornament are sometimes produced.

**shelly limestone** *\*Carbonate* rock that contains a large proportion of shell or shell fragments.

**shelter porosity** See CHOQUETTE AND PRAY CLASSIFICATION.

**Shenzhou (SZ)** A series of Chinese satellites hosting research into space life science, space material, Earth observation, and monitoring of Earth and space environment. There have been seven satellites. SZ-1 operated from 20 to 21 November 1999; SZ-2 from 9 January until 24 August 2001; SZ-3 from 25 March until 12 November 2002; SZ-4 from 29 December 2002 until 5 January 2003; SZ-5, the first Chinese manned mission, from 15 to 16 October 2003; SZ-6 (manned) from 12 to 16 October 2005; and SZ-7 from 25 to 28 September 2008.

**shergottite/nakhlite/chassignite meteorites (SNC)** Association of about a dozen *\*meteorites*, much younger (1.3 billion years) than the majority found (4.6 billion years). They have *\*igneous* *\*textures* and contain iron-rich *\*silicates* and iron oxides, therefore must come from an oxygen-rich

environment, and also small amounts of water-bearing minerals. Probably they all come from the same planetary body; the most favoured possibility is the planet [\\*Mars](#).

**Shermanian** A regional [\\*stage](#) (457–454 Ma ago) of the Late [\\*Ordovician](#), preceded by the [\\*Soudleyan](#) and followed by the [\\*Actonian](#).

**shield 1.** A tunnel borer consisting of a conventional shield with thrust rams and erector system. The cutter head and support are inside the shield. This type of machine is usually employed in soils or variable materials, e.g. a sand–rock–gravel sequence. **2.** See [CRATON](#).

**shield volcano** See [HAWAIIAN ERUPTION](#).

**shingle** Beach pebbles, normally well rounded as a result of abrasion, whose diameters are typically 0.75–7.5 cm. They are made of resistant materials such as [\\*flint](#), which is the dominant constituent of the shingle beaches of south-eastern England. They may also show lateral sorting, e.g. the shingle of the Chesil Beach, Dorset, England, steadily increases in size over 29 km from west to east.

**Shipka** A [\\*solar system](#) asteroid (No. 2530), with an orbital period of 5.25 years. It was to have been visited in October 2008 by the [\\*Rosetta](#) spacecraft, but the plans were revised and Shipka was dropped from the itinerary.

**shoal** A mound or other structure raised above the sea bed in shallow water that is composed of, or covered by, unconsolidated material and may be exposed at low water.

**shoaling** The behaviour of waves as they approach a shelving shore. The waves cease to be symmetrical and sinusoidal and become asymmetrical and solitary. Wavelength and wave velocity decrease, wave height and wave steepness increase, and wave period remains constant.

**shoal retreat massif** Large [\\*sand](#) accumulation preserved on the [\\*continental shelf](#) during and after a marine [\\*transgression](#). The massifs represent former [\\*estuary](#)-mouth sand bars (inlet-associated shoals) or former zones of [\\*longshore-drift](#) convergence (cape-associated shoals). In the Middle Atlantic Bight of the east coast of the USA the massifs are up to 70 km long by 20 km wide and can be found offshore from the present

Hudson, Delaware, and Chesapeake estuaries, and longshore-drift convergence zones such as Cape Hatteras.

**shock metamorphism** Changes included in *\*rocks* and *\*minerals* when they are subjected to pressures from shock waves travelling at 10–13 km/s resulting from *\*meteorite*, cometary, or asteroidal impact. Pressures ranging to more than 50 Mpa are experienced during such events. Materials subjected to less than 10 Mpa are severely disrupted and fractured; between 10 and 25 Mpa, framework *\*silicates* exhibit planar deformation features; maskelynite (vitrified *\*plagioclase feldspar*) forms between 25 and 40 Mpa; *\*quartz* and *\*feldspar* melt from 40 to 60 Mpa; while rocks subjected to pressures of more than 60 to 70 Mpa become molten. Peak pressures are sufficient to vaporize material.

**shock-remanent magnetization** The magnetization acquired as a result of rapidly applied stress while being held in a magnetic field. It is usually attributed to meteoritic impacts. *See also* [PIEZOREMANENT MAGNETIZATION](#).

**Shoemaker, Eugene Merle** (1928–97) An American geologist who founded the scientific study of impact cratering on Earth, the Moon, and on other solar system planets and their satellites, and who also pioneered the study of near-Earth comets and asteroids, often in collaboration with his wife Carolyn. This led him to recognize the importance of the cratering process in the history of the solar system. Shoemaker worked for the US Geological Survey ([USGS](#)) from 1948 to 1993 then continued to serve as scientist emeritus. In the 1960s he established the Astrogeology Branch of the USGS and its centre at Flagstaff, Arizona. Studies of the Moon and other bodies lacking atmospheres led him to propose that their surfaces should be covered with a layer of ejecta he termed ‘regolith’, and that the age of a surface could be inferred from its cratering. In March 1993, Gene and Carolyn Shoemaker and David Levy discovered a fragmented comet close to Jupiter. The comet, named Shoemaker-Levy 9, collided with Jupiter spectacularly in July 1994. Gene Shoemaker died in a car crash in the Australian outback. He and Carolyn, who survived, had been on their way to study a crater.

**shoestring sand** An irregular, sinuous *\*sandbody* having a form resembling a shoe-lace, often representing the preserved sandy deposit of a meandering river channel (*see* [MEANDER](#)).

**shonkinite** A dark coloured, coarse-grained, *\*igneous* rock consisting of *\*essential \*diopside* (making up about 50% of the rock), *\*alkali feldspar*, and *\*biotite*, with or without *\*olivine* and/or *\*nepheline*. Where olivine or nepheline is abundant the rock name is prefixed with these *\*mineral* names (e.g. olivine-shonkinite). Shonkinites are essentially a type of alkali *\*syenite*.

**shooting flow** See CRITICAL FLOW; FROUDE NUMBER.

**shoreface** The subtidal coastal zone between the low-water mark and a depth of about 10–20 m, within which wave action governs the sedimentary processes. Below the lower limit of the shoreface waves do not affect the sea bed. See WAVE BASE.

**shore platform (marine platform, marine terrace, marine flat, marine bench, wave-cut bench, wave-cut platform)** Intertidal bench cut into a land mass by the action of waves and associated processes. It is terminated landward by a sea cliff, and slopes gently seaward at about 1°.

**shortening** The reduction in length of a line as a result of *\*strain*, e.g. by thickening, folding, thrusting, or the loss of material by solution. Simple shortening may be calculated using the same equation as for *\*extension* ( $e$ ):  $e = (L_F - L_O)/L_O$ , where  $L_F$  is the final length and  $L_O$  the original length; if  $e$  has a negative value when calculated from this equation, the line has been shortened.

**short wavelength infrared** See NEAR-INFRARED.

**shot** A source of seismic shock waves that are produced for experimental purposes, e.g. by a *\*hammer*, an explosion, an *\*airgun*, or a *\*water gun*.

**shot bounce** The noise on a *\*seismic record* that is generated by the physical and mechanical motion of a recording vehicle.

**shotcrete** A type of *\*concrete*, sometimes reinforced with metal or glass fibre, which is sprayed on to the face of an excavation to form a protective lining and support, usually a few tens of millimetres thick. It is most useful in protecting soft or weak material and can be adapted to suit varying conditions. It is often used in conjunction with other forms of support, e.g. *\*rock bolts* and *\*arching*.

**shot depth** The depth below ground level of a seismic source. If the source is non-explosive, it is the depth at which the source impulse is generated; if it is explosive, it is the depth to the top of the explosive, or to its mid-point if the length of the explosive material is small in relation to the depth of the hole, or to both top and bottom of the explosive charge (i.e. two shot depths) if it is very large.

**shotpoint gap** See SEISMIC GAP.

**shower** \*Precipitation of short duration that begins and ends abruptly, associated with convective clouds which usually do not form a continuous cover of the sky. Intensity varies, e.g. showers may be slight, moderate, or heavy. Hail or snow may form part of the precipitation.

**shrieking sixties** Popular maritime term for the prevailing westerly winds which are commonly strong over the oceans in temperate latitudes of the southern hemisphere, south of 60 °S. See also FURIOUS FIFTIES; ROARING FORTIES.

**SHRIMP** See SENSITIVE HIGH RESOLUTION ION MICROPROBE.

**shrimps** See MALACOSTRACA.


**shrinkage** In \*concrete, a measurement of the reduction in dimensions due to the loss of water. Fine-grained \*cement has a low coefficient of permeability and behaves in a similar manner to fine-grained soil.

**shrinkage cracks** See DESICCATION CRACKS.

**shrinkage joint** See JOINT.

**shrub-coppice dune (nabkha, nebkha)** A \*sand \*dune that is irregular or approximately elliptical in shape, and streamlined in the downwind direction, that results from the accumulation of sand around a clump of vegetation and downwind of it. It is the commonest type of sand dune.

**shutter ridge** A ridge that forms when a \*strike-slip fault occurs across a \*gully, at an angle to the axis of the gully. The movement shifts the raised side of the gully on one side of the fault to a position opposite the lowered side of the gully on the other side of the fault.



**Shutter ridge**

**SH-wave** See [LOVE WAVE](#).

**SI (SI units)** *Système International d'Unités*, an internationally agreed group of units of measurement, used especially in scientific work. The system comprises seven basic and two supplementary units. The basic units are the: metre (m); kilogram (kg); second (s); ampere (A); kelvin (K); mole (mol); and candela (cd). The supplementary units are the: radian (rad); and steradian (sr). A further 18 units are derived from these: becquerel (Bq); coulomb (C); farad (F); gray (Gy); henry (H); hertz (Hz); joule (J); lumen (lm); lux (lx); newton (N); ohm ( $\Omega$ ); pascal (Pa); siemens (S); sievert (Sv); tesla (T); volt (V); watt (W); and weber (Wb). See [SI UNITS, CONVERSIONS, AND MULTIPLES \(APPENDIX D\)](#).

**Siberian high** The region of high average pressure in winter over Siberia, centred to the south of Lake Baikal.

**sichelwannen** Crescent-shaped [\\*p-forms](#) cut into a flat or sloping rock surface by the action of a [\\*glacier](#). They are typically 1–10 m long and 5–6 m wide. The horns of the crescents point down-glacier. Their origin is unclear; the action of saturated [\\*till](#) and of subglacial meltwater have both been invoked. The name is German, its literal meaning being 'sickle tubs'.

**Sicilian** A European [\\*stage](#) (0.781–0.26 Ma ago) of the Middle [\\*Pleistocene](#), preceded by the Early Pleistocene and followed by the [\\*Tyrrenian](#).

**sicula** The skeleton of the initial zooid of a graptolite ([\\*Graptolithina](#)) colony.

**side-looking airborne radar (SLAR)** An airborne \*radar system which scans sideways from the flight track and detects the \*backscattered radar \*reflections from the ground surface in order to produce a radar image of the ground much as \*side-scan sonar produces an image of the sea floor. Since different surface materials have different radar reflectance characteristics the radar image can be interpreted in terms of surface features.

**sidereal day** The time it takes for the Earth to complete one rotation on its axis such that a particular point on the surface returns to its former position in relation to the position of the fixed stars. The sidereal day, of 23 hours 56 minutes and 3.93 seconds, is 3 minutes 55.9 seconds shorter than the mean solar day, because the movement of the Earth in its solar orbit is imposed on its rotational motion.

**sidereal month** A period of 27 days, 7 hours, and 43 minutes, which is the average period of revolution of the \*Moon around the \*Earth, as determined by using a fixed star as a reference point.

**Siderian** A period of the \*Proterozoic eon that began 2500 Ma ago and ended 2300 Ma ago, its beginning and end being defined by fixed dates. The Siderian was the first period of the \*Palaeoproterozoic, following the end of the \*Archaean eon, and was followed by the \*Rhyacian period.

**siderite (chalybite, spathose iron)** Mineral,  $\text{FeCO}_3$ ; sp. gr. 3.8–4.0; \*hardness 3.5–4.5; \*trigonal; grey to grey-brown or yellowish-brown, translucent when pure; white \*streak; vitreous \*lustre; uneven \*fracture; crystals \*rhombohedral with curved faces, but also occurs \*massive, granular, fibrous, compact, \*botryoidal, and \*earthy in habit; \*cleavage perfect rhombohedral {1011}; widespread in \*sedimentary rocks, especially \*clays and \*shales where it is \*concretionary and makes clay into \*ironstone, also as a \*gangue mineral in \*hydrothermal veins together with other metallic ores (e.g. \*pyrite, \*chalcopyrite, and \*galena) and as a replacement mineral in \*limestone; dissolves slowly in cold, dilute hydrochloric acid, which effervesces when warmed.

**siderolite** See STONY-IRON METEORITE.

**siderophile** Applied to elements with a weak affinity for oxygen and sulphur and soluble in molten iron. For example, Ni, Co, Pt, and Ir, which

are found in iron *\*meteorites* and are probably concentrated in the Earth's *\*core*. Compare *ATMOPHILE*; *BIOPHILE*; *CHALCOPHILE*; *LITHOPHILE*.

**side-scan sonar** A sideways-looking acoustic-survey system which uses the *\*reflection* of high-frequency (30–110 kHz) sound waves from a surface to map the texture of that surface. It is most commonly used to map sea-floor features (sand *\*ripples*, rock *\*outcrops*, pipes, wrecks, etc.) but it can also be suspended vertically from a vessel to investigate vertical surfaces, e.g. the submerged parts of icebergs and major ice fronts.

**sidewall corer** A device for coring samples from the side of a drill hole. See also *CORE SLICER*.

**Sidufjall** A normal *\*polarity subchron* which occurs within the *\*Gilbert* reversed *\*chron*.

**Siegennian** See *PRAGHIAN*.

**siemens (S)** The derived SI unit of electrical conductance in a circuit with a resistance of one ohm, and thus equal to a reciprocal ohm (mho, or  $\Omega^{-1}$ ). It is used in measuring conductivities, where the unit is S/m. The siemens is named after Sir William Siemens (1823–83).

**sieve** A circular-framed container with a meshed base. The mesh size is accurately machined and the sieve will permit the passage only of particles finer than the mesh size.

**sieve deposit** A well-sorted, *\*matrix*-free *\*conglomerate*, which forms where the *\*sediment* transported and deposited comprises only *\*pebble* and gravel grades. Sieve deposits are found mainly on *\*alluvial fans* whose source areas consist of well-jointed, resistant *\*lithologies* such as *\*quartzites*.

**sieving** A method of grain-size analysis (see *PARTICLE SIZE*) in which the sample is passed through a stack of sieves, arranged with the coarsest mesh at the top and finest at the base. The weight of *\*sediment* trapped on each mesh is recorded, and percentage weight plotted either on a histogram or cumulative frequency curve (see *CUMULATIVE PERCENTAGE CURVE*).

**sigma-t density ( $\sigma$ -t density)** The density of a sea-water sample measured at atmospheric pressure, i.e. at the sea surface. It is defined as density minus

1000, where the density is measured in kg/m<sup>3</sup>. Thus sea water at 0°C and 35‰ **\*salinity** has a density of 1028 kg/m<sup>3</sup>, or a  $\sigma$ -t density, at a pressure of 0.1 MPa, of 28.

**signature** The characteristics of geophysical anomalies within a region or along a profile. It is often based on **\*Fourier** or power-spectrum analyses of either gravity (gravity signature) or magnetic (magnetic signature) **\*residual** anomalies.

**significant wave height** The average height of the highest one-third of waves in a given group of waves. This height is usually used as a standard when describing the wave characteristics of a given area.

**silcrete** See DURICRUST.

**silent quake** See SLOW SLIP.

**Silesian** A European **\*epoch** of the **\*Carboniferous** (326.4–299 Ma ago), underlain by the **\*Dinantian**, followed by the **\*Asselian**, and comprising the **\*Namurian**, **\*Westphalian**, and **\*Stephanian** **\*series**. It is roughly contemporaneous with the uppermost **\*Mississippian** (**\*Serpukhovian**) plus the **\*Pennsylvanian**.

**silex** See FLINT.

**silica** Silicon dioxide (SiO<sub>2</sub>) which occurs naturally in three main forms: (a) crystalline silica includes the **\*minerals** **\*quartz**, **\*tridymite**, and **\*cristobalite**; (b) **\*cryptocrystalline** or very finely crystalline silica includes some **\*chalcedony**, **\*chert**, **\*jasper**, and **\*flint**; and (c) amorphous hydrated silica includes **\*opal**, **\*diatomite**, and some chalcedony. **\*Coesite** and **\*stishovite** are two high-density **\*polymorphs** of quartz which rarely occur in nature but have been synthesized experimentally.

**silica-oversaturated rock** See SILICA SATURATION.

**silica saturation** The concentration of silica (SiO<sub>2</sub>) in an **\*igneous** rock, relative to the concentration of other chemical constituents in the rock which combine with the silica to form **\*silicate** minerals. On this basis, three classes of igneous rock are recognized: (a) silica-oversaturated rocks (e.g. **\*granite**), in which there is more than enough silica to satisfy the requirements of all the major silicate minerals, the free silica appearing as

\*quartz in the rock; (b) silica-saturated rocks (e.g. \*diorite), in which there is just enough silica present to satisfy the requirements of all the major silicate minerals, there being neither an excess nor deficiency of silica, resulting in a lack of both quartz and \*feldspathoid minerals in the rock; and (c) silica-undersaturated rocks (e.g. \*nepheline-syenite), in which there is not enough silica present to satisfy the requirements of all the major silicate minerals, the silica deficiency being accommodated by the crystallization of feldspathoids (\*nepheline, \*leucite) in place of \*feldspar, the feldspathoids containing less silica in their structure than feldspars.

**silicates** The most important and abundant group of rock-forming minerals, which can be classified according to the structural arrangement of the fundamental  $\text{SiO}_4$  \*tetrahedra which are the main building blocks of the group. (a) \*Nesosilicates have independent  $\text{SiO}_4$  \*tetrahedra linked by \*cations, e.g. \*olivine group. (b) Sorosilicates have two  $\text{SiO}_4$  \*tetrahedra sharing one oxygen, e.g. \*epidote group. (c) \*Cyclosilicates have rings of three, four, or six linked  $\text{SiO}_4$  \*tetrahedra, e.g. \*axinite and \*tourmaline. (d) \*Inosilicates (chain silicates) have  $\text{SiO}_4$  \*tetrahedra linked either into single chains by sharing two oxygens, e.g. \*pyroxene group, or into double chains (band silicates) by alternately sharing two or three oxygens, e.g. \*amphibole group, (e) \*Phyllosilicates (sheet silicates) share three oxygens to form a flat sheet, e.g. \*mica group. (f) Tectosilicates have  $\text{SiO}_4$  \*tetrahedra linked into a three-dimensional framework by sharing all the oxygens, e.g. \*feldspar and \*quartz groups.

**silica-undersaturated rock** See SILICA SATURATION.

**siliceous ooze** Fine-grained \*pelagic deposit of the deep-ocean floor with more than 30% siliceous material of organic origin. \*Radiolaria and \*diatom remains are the major constituents of the siliceous oozes, which tend to occur at depths in excess of 4500 m.

**siliceous sinter** A \*silica-rich precipitate found around the mouth of a \*geyser or hot spring whose waters carry large amounts of dissolved minerals which precipitate when the water cools suddenly on exposure to the atmosphere.

**siliciclastic** Applied to a \*sediment which comprises particles composed of \*silicate minerals and rock fragments, i.e. \*mudstones, \*sandstones, and

\*conglomerates.

**silicification** The introduction of \*cryptocrystalline \*silica into a non-siliceous rock via \*groundwater or fluids of \*igneous origin. The introduced silica either fills \*pore spaces in the rock or replaces pre-existing minerals.

**silicon 'burning'** In stellar evolution, the process whereby silicon, with magnesium and sulphur, 'burns' at a temperature of around  $3 \times 10^9$  K, producing elements of the 'iron peak' (e.g. Cr, Mn, Fe, Co, and Ni). See CARBON 'BURNING'; HELIUM 'BURNING'; HYDROGEN 'BURNING'; OXYGEN 'BURNING'; NUCLEOSYNTHESIS.

**silky** Applied to the mineral \*lustre of some fibrous minerals, e.g. \*satin spar (see GYPSUM).

**sill** A tabular \*igneous \*intrusion having \*concordant surfaces of contact.

**sillar** A local Peruvian name for \*ignimbrites that have been pervasively altered by \*vapour-phase crystallization. The alteration of original \*pumice and \*glass shards involves deposition of \*tridymite, \*cristobalite, and \*alkali feldspar that occurs as \*drusy infills of the \*matrix and pumice cavities in the upper parts of ignimbrite cooling units. The gas is derived by diffusion from \*juvenile glassy fragments within the flow and from heated \*groundwater percolating through the flow. The resulting material makes excellent, lightweight, easily worked building stone, usually white. The city of Arequipa, southern Peru, is renowned for its buildings of sillar.

**sillimanite (fibrolite)** A \*nesosilicate and one of an important group of three \*mineral \*polymorphs with the same composition  $\text{Al}_2\text{SiO}_5$ , the other two being \*andalusite and \*kyanite; sp. gr. 3.23; \*hardness 6.5–7.5; \*orthorhombic; occurs as elongate \*prismatic \*crystals with a diamond-shaped cross-section or as \*fibrous and felted masses; colourless; occurs under conditions of high-grade \*metamorphism either in the innermost zones of thermal aureoles (see CONTACT AUREOLE) or in high-grade regional metamorphic rocks (see REGIONAL METAMORPHISM) formed under high temperatures and pressures. It is named after the American mineralogist B. Silliman.

**silt** **1.** In the commonly used Udden–Wentworth scale, particles between 4µm and 62.57 µm in size. Other classifications exist. In *\*pedology*, silt refers to mineral soil particles that range in diameter from 0.02 to 0.002 µm in the international system, or from 0.05 to 0.002 µm in the *\*USDA* system. See *PARTICLE SIZE*. **2.** A class of soil texture.

**siltstone** A lithified *\*silt*, comprising *\*grains* between 4 µm and 62.5 µm in size.

**Silurian** Third (443.7–416 Ma ago) of six *\*periods* of the *\*Palaeozoic \*era*. The end of the period is marked by the climax of the *\*Caledonian orogeny* and the filling of several Palaeozoic basins of deposition.



<https://ucmp.berkeley.edu/silurian/silurian.php>

- The Silurian Period.

**silva** See *CLOUD CLASSIFICATION*.

**silvagenitus** See *CLOUD CLASSIFICATION*.

**silver, native** Malleable metallic element, Ag; sp. gr. 10.5; *\*hardness* 2.5; white; *\*cubic* crystals, thin sheets, or scales; occurs in association with other silver minerals in the oxidized zone of silver-rich mineral deposits and in *\*hydrothermal* vein deposits.

**silver glance** See *ARGENTITE*.

**silver iodide** Substance used in the form of fine particles to act as nuclei in *\*cloud seeding*.

**silver spike** In stratigraphy, a time-line selected as a reference point in a regional succession of *\*strata*. Compare *GOLDEN SPIKE*.

**silver thaw** **1.** Phenomenon resulting from the formation of ice on cold surfaces (e.g. due to rain) during a period of rapid thaw after a severe frost. See also *GLAZE*. **2.** A name given to freezing rain in the Pacific Northwest of the USA; it is called ‘thaw’ because warmer weather often follows it.

**Simiiformes (Anthropoidea)** (cohort Unguiculata, order Primates) Infra-order comprising the monkeys, apes, and humans. Monkeys and apes have

a common ancestor and diverged in the **\*Oligocene**. The dryopithecines of the succeeding **\*Miocene** were undoubted apes. The traditional palaeontological view is that these Miocene apes gave rise in turn to three new lines, one leading to the gibbons, another to the great apes, the third to humans. However, on the basis of anatomical characteristics and genetic criteria, it has long been maintained that ancestors of both the gibbons and the orang-utan diverged from the ancestral line of the advanced Primates at an early date and that only subsequently did that line split to give one group comprising humans and another comprising the gorilla, chimpanzee, and bonobo; recent palaeontological evidence now tends to support this second view.

**similar fold** A **\*fold** in which the **\*orthogonal thickness** changes systematically such that the thickness of the fold remains constant when measured parallel to the **\*axial surface**. Single layers are thicker at the **\*hinge** and thinner in the limb regions. Unlike a **\*parallel fold**, a similar fold can maintain its form throughout a sequence.

**simoom** Regional wind that blows in spring and summer over the south-eastern Sahara and Arabia. The wind blows for only a short period at a time and its whirlwind effect carries sand and brings very hot, dry conditions.

**simple shear (homogeneous rotational strain, rotational shear)** A rotational **\*strain** in which the maximum and minimum strain axes are re-oriented in relation to their original positions. *See also* **HOMOGENEOUS STRAIN**. *Compare* **PURE SHEAR**.

**Simpson, George Gaylord** (1902–84) An American palaeontologist, specializing in mammalian evolution, who also studied mammal migrations, Simpson obtained his Ph.D. from Yale University in 1926, for a thesis on **\*Mesozoic** mammals, and then joined the staff of the American Museum of Natural History, in New York City, becoming curator in 1942. In 1945 he moved, as a professor, to Columbia University. From 1959 to 1970 he was Alexander Agassiz Professor of Vertebrate Palaeontology at the Museum of Comparative Zoology at Harvard University, and in 1967 was appointed professor of geosciences at the University of Arizona. His studies of fossil mammals, especially those of Madagascar, led him initially to oppose the theory of **\*continental drift**. Simpson proposed that the dispersion of species occurred along **\*sweepstakes dispersal routes**.

**Sinemurian** A *\*stage* (196.5–189.6 Ma ago) in the European Early *\*Jurassic*, preceded by the *\*Hettangian*, followed by the *\*Pliensbachian*, and contemporaneous with the upper Aratauran (New Zealand). *See also* LIAS.

**single couple (Type I earthquake source)** A *\*seismic wave* pattern, consisting of four lobes (for *\*P-waves*) and two lobes (for *\*S-waves*) of alternate compression and dilation, which is generated by movement along a single *\*fault plane*. *Compare* DOUBLE COUPLE.

**single-stage lead** *\*Common lead* (Pb) removed at a particular time from a single, constantly evolving, uranium–thorium–lead environment. It can be used to provide the basis for the common-lead method of dating (*see* LEAD–LEAD DATING). After the removal of this lead in solution and its deposition as a lead ore (such as *\*galena*), it is then assumed not to have been altered in isotopic composition by the further addition of *\*radiogenic* products.

**singularity** In meteorology, a short seasonal episode lasting a few days and commonly occurring about specific dates of the year. One example is the periods of fine, dry, warm weather commonly occurring in September and early October in Britain and central Europe, and known as ‘old wives’ summer’ (or ‘Indian summer’), which are due to slow-moving *\*anticyclones*.

**Sinian** The Chinese final *\*stage* of the *\*Neoproterozoic*, dated at about 800–542 Ma ago (Int. Commission on Stratigraphy, 2004), preceded by the *\*Riphean* and followed by the *\*Meishuchuan*. The name is derived from a region of central China.

**sinistral coiling** *See* COILING.

**sinistral fault (left lateral fault)** The sense of displacement in a *\*strike-slip fault* zone where one block is displaced to the left of the block from which the observation is made.

**sink** A natural reservoir that can receive energy or materials without undergoing change.

**sink-hole** *See* DOLINE.

**sinking** *See* DOWNWELLING.

**Sinope (Jupiter IX)** One of the lesser satellites of *\*Jupiter*, with a diameter of 28 km; its orbit is *\*retrograde*.


***Sinornis*** An early *\*Cretaceous* bird from Liaoning Province, China. It is one of the earliest to show the advanced avian features of an extensive *\*synsacrum* and complete fusion of the distal *\*tarsal bones* with *\*metatarsals*.

**sinter** A concentration of material, especially *\*silica* and calcium carbonate, formerly dissolved in a hydrothermal solution, that was precipitated as the solution cooled.

**sinus** A Latin word meaning ‘curve’ or ‘bay’. **1.** The word ‘sinus’ was used by Giovanni B. Riccioli in 1651 to designate bay-like features on the lunar maria (see *MARE*). The best known example is Sinus Iridum, the ‘Bay of Rainbows’, on the north-western margin of the ‘Sea of Rains’ (Mare Imbrium). **2.** In certain bivalves (*\*Bivalvia*), a recess or embayment in the pallial line; most bivalves with a pallial sinus are burrowers.

**siphon** In bivalve molluscs (*\*Bivalvia*) and gastropods (*\*Gastropoda*), a tube that connects the mollusc to the world outside, funnelling water towards and away from the gills. In bivalves siphons may occur in pairs.

**siphonal canal** In some gastropod (*\*Gastropoda*) shells, an indentation or channel that accommodates the *\*siphon*.



### Siphonal canal

**siphonate** Applied to the *\*aperture* of a gastropod (*\*Gastropoda*) shell when a canal or notch for the *\*siphon* is present.

**siphonostomatous** See APERTURE.

**siphuncle** A long tube present in those cephalopods (*\*Cephalopoda*) that possess external shells. It runs internally, through all the chambers, and contains the siphuncular cord of body tissue, extending from the visceral mass through a perforation in each *\*septum* of the shell. The siphuncle releases gas into the unoccupied chambers of the shell, making the animal buoyant.

**Sistema Satelital para la Observación de la Terra (SSOT)** An optical high-resolution *\*microsatellite* imaging mission by the Chilean Space Agency (Agencia Chilena del Espacio) and the Chilean Air Force. The mission was launched on 17 December 2011, from Kourou, French Guiana, into a Sun-synchronous near-circular orbit at a nominal altitude of 620 km.

**sister groups** The twin products of *\*cladogenesis* which, under the rules of *\*phylogenetic* systematics, must be classified at the same rank

taxonomically (i.e. the two daughter lineages that result from the splitting of a single parent species).

**sister taxa** In *\*phylogenetics*, two taxa connected through a single *\*internal node*.

**site investigation** The geologic examination of a potential development site in order to design the foundations of surface buildings, roads, etc. It includes geophysical surveys, trial pits, and *\*boreholes*.

***Sivapithecus*** Genus of early *\*hominoid* Primates, which probably includes the so-called *\*Ramapithecus*. They are known from E. Africa, south-eastern Europe, Turkey, Arabia, Pakistan, northern India, and southern China, from the Middle and Late *\*Miocene* 15–8 Ma ago. The genus may include ancestors of great apes and humans, but certainly early relatives of the orang-utan have been identified among species of the genus. Although there are similarities between the teeth of *Sivapithecus* and those of humans, these are probably misleading, and *Sivapithecus* is now regarded as an ape that was ancestral either to the orang-utan alone, or to all living great apes and humans.

**skarn** A contact metamorphic rock (see *CONTACT METAMORPHISM*) composed of calcium, magnesium, and iron silicates (with or without iron, copper, and manganese sulphides and oxides) which has been derived from *\*limestone* or *\*dolomite* by the metasomatic (see *METASOMATISM*) introduction of large amounts of silicon, aluminium, iron, and magnesium from a nearby *\*igneous \*intrusion*, usually a *\*granite*. Many skarns serve as host rocks for economic deposits of *\*magnetite* and copper sulphides.

**skeletal limestone** See *LEIGHTON–PENDEXTER CLASSIFICATION*.

**skeletal material** In most vertebrates the skeleton is made from *\*bone* (calcium phosphate); among *\*invertebrates* it is more varied. *\*Calcite* or *\*aragonite* in various forms is common in such groups as the brachiopods (*\*Brachiopoda*) and molluscs (*\*Mollusca*). Normally the invertebrate skeleton is made up from several layers, and often each layer has a distinctive structure. The main, calcified portion of the shell is called the ‘ostracum’; an outer layer, the ‘periostracum’ (made from layers of protein), disappears after death. Calcitic skeletons also occur in corals and *\*Bryozoa*. Echinoderms (*\*Echinodermata*) have skeletons made up of a number of

elements. Each element is permeated with living tissue but the hard material is calcite, which forms in optical continuity to form a single *\*crystal*. Chitin (a hydrocarbon related to cellulose) is the principal component of insect cuticle. It has been assumed that chitin impregnated with calcium carbonate forms the exoskeletal material in trilobites (*\*Trilobita*), but although organic material occurs its nature is still unknown. Similarly, recent studies of graptolite (*\*Graptolithina*) skeletal material have shown that the material is not chitin but a scleroprotein (a fibrous, insoluble protein). In some simple animals, e.g. *\*Radiolaria* and some sponges (*\*Porifera*), the skeleton is composed of opaline silica.

**skeletal micritic limestone** See LEIGHTON–PENDEXTER CLASSIFICATION.

**skewness** An expression of the degree of asymmetry shown on a frequency distribution. A non-skewed distribution is a perfect, normal (*\*Gaussian*) distribution. *\*Grain*-size distributions are skewed positively where larger particles are more numerous than fine ones, or negatively where finer particles exceed coarse within the distribution.

**skin depth ( $z_s$ )** The effective depth of penetration ( $m$ , in metres) of an *\*electromagnetic wave* of frequency  $f$  Hz, at which the amplitude of the wave has been attenuated by  $1/e$  (37%) of its surface value. It can be calculated by:  $z_s = 503.8/\sqrt{(\sigma f)} = 503.8\sqrt{(\rho/f)}$ , where  $\sigma$  is the conductivity (in S/m) and  $\rho$  the resistivity (in  $\Omega/m$ ) of the medium.

**skip mark** A type of *\*tool* mark formed by the intersection at a low angle of an object with a muddy *\*sediment* surface, so that the object skips across the surface, producing a linear series of depressions. See also BOUNCE MARK; PROD MARK; SEDIMENTARY STRUCTURE; TOOL MARK.

**Skolithos** With *\*Ophiomorpha*, an *\*ichnoguild* of cylindrical or U-shaped, vertical or steeply inclined *\*burrows*, permanently occupied by suspension feeders. The burrows lack *\*spreiten*, and were possibly formed in shallow, sublittoral environments. Burrows of this type are recorded from the *\*Cambrian* to Recent.

**SKS-wave** See SEISMIC-WAVE MODES.

**skull** See CRANIUM.

**Skylab** The first US manned Earth-orbiting space station; Skylab-1 (unmanned) was launched on 14 May 1973, Skylab-2 (manned) on 25 May 1973, Skylab-3 (manned) on 28 July 1973, and Skylab-4 (manned) on 16 November 1973, all from the Kennedy Space Center, into a near-circular orbit at an altitude of 435 km.

**SkySat** A commercial *\*microsatellite programme* to provide customers with reliable and frequent high-resolution images. SkySat-1 was launched on 21 November 2013, from the Yasnny Cosmodrome, Russia, into a Sun-synchronous near-circular orbit at an altitude of 600 km. SkySat-2 was launched on 8 July 2014, from Baikonur Cosmodrome, Kazakhstan, into a Sun-synchronous circular orbit at an altitude of about 635 km. SkySat-3 was launched on 22 June 2016, from Satish Dhawan Space Centre, India, into a Sun-synchronous orbit at an altitude of 515 km.

**Skythian** See SCYTHIAN.

**sky-view factor** The extent of sky observed from a point as a proportion of the total possible sky hemisphere.

**slab-pull** The force, caused by the sinking of the cold, dense *\*lithosphere* into the *\*asthenosphere* at a *\*destructive margin*, which is hypothesized to be one of the two major driving forces for the movement of *\*plates* (the other is *\*ridge-push*).

**slake-durability test** A test to estimate the resistance of rocks, particularly *\*argillaceous* rocks, to a combination of wetting and abrasion. Test results are expressed as a slake-durability index for each particular rock.

**slaking** Breaking up of earth materials when exposed to water or air. The susceptibility of rock to slaking is measured with a *\*slake-durability* test.

**slant range** In *\*radar* terminology, the distance between the radar platform and an object on the ground.

**slant-range resolution** The minimum distance by which two objects on the ground must be separated in order to be resolved by a *\*radar*. The minimum distance must be equal to or greater than half the *\*pulse* length of a radar. See also AZIMUTH RESOLUTION.

**SLAR** See SIDE-LOOKING AIRBORNE RADAR.

**slate** Low-grade, *\*regionally metamorphosed* rock, which is highly fissile and fine grained. The *\*fissility* (*\*slaty cleavage*) results from the parallel alignment of numerous fine *\*phyllosilicate* minerals (e.g. *\*muscovite* and *\*chlorite*) induced by compressive tectonic deformation. The smooth, hard, impermeable surface produced when slate is split makes it commercially valuable for roofing, cladding of buildings, and for making such items as billiard-table tops, laboratory benches, and blackboards. *Compare* PHYLLITE; SCHIST.

**slaty cleavage** A continuous *\*cleavage* formed by the preferred alignment of homogeneously distributed *\*phyllosilicates* throughout a rock. Slaty cleavage is commonly produced during lower grade *\*metamorphism*. See METAMORPHIC GRADE.

**sleet** **1.** In UK usage, *\*precipitation* in the form of a mixture of rain and melting snow. **2.** In N. American usage, ice pellets of less than 5 mm in diameter that form when rain freezes as it falls.

**sleeve exploder** A seismic source, used in marine investigations, in which a mixture of gases (e.g. propane and oxygen) is detonated inside a rubber sleeve which expands rapidly to accommodate the products of the explosion, thus generating a shock wave in the surrounding water. The exhaust gases are vented to the surface through a pipe to reduce problems caused by a *\*bubble* pulse.

**slick** Quiescent area on the surface of a water body. The relative smoothness of these areas compared to adjacent waters is usually caused by a thin surface film of oil which changes the *\*surface tension*.

**slickenfibres** Clusters of *\*mineral* fibres that grow along *\*fault planes*, marking the direction of relative displacement of the faulted blocks.

**slickenside** **1.** A *\*lineation* on a *\*fault* or *\*bedding plane* caused by the frictional movement of one rock body against another. The plane may be coated by a *\*mineral*, often *\*quartz* or *\*calcite*, which itself shows striations in the direction of movement. **2.** The polished surface left by the passage of a mud slide. **3.** In soils, the natural crack surfaces produced by swelling and shrinkage in clayey soils that are high in swelling *\*clays*.

**slide** See GRAVITY SLIDING.

**slide-rock** See ROCK SLIDE.

**slingram method** A dual-coil, \*electromagnetic \*profiling system in which both the transmitter and receiver can be moved while maintaining a constant distance between them.

**sling psychrometer** See WHIRLING PSYCHROMETER.

**slip** 1. The relative displacement to either side of a \*fault plane of points which were originally coincident. The total displacement (i.e. the sum of the \*dip-slip and \*strike-slip components) is called the 'net slip'. 2. **(translation gliding)** The gliding of intracrystalline zones in relation to one another over distances which are integers of the unit pattern.

**slipface** The lee face of a sand dune where the surface is at the \*angle of repose for sand (30–34°).

**slip-off slope** See MEANDER.

**sloc** (*pl.* **slocan**) From the Gaelic word for hollow. Long, deep, parallel-sided depressions at the coast, common on the Hebridean islands of Mull and Skye, caused by the erosion of \*dykes cutting through the ancient \*gneisses.

**slope angle** The gradient of a stable hillslope. In part this may reflect the \*particle-size characteristics of the mantling \*regolith. For example, maximum slope angles of about 26° are common in areas where the regolith consists of a mixture of rock rubble and coarse soil particles.

**slope processes** The set of geomorphological \*processes that act on and below the surface of a hillslope, affecting the \*regolith and bedrock. Some of the most important are \*rain-wash, \*sheetwash, \*weathering, \*mass movement, piping, and linear \*erosion by \*rill and gully activity. Each process tends to be associated with a distinctive \*slope profile.

**slope profile** The two-dimensional form of a hillslope when measured down the steepest gradient. Traditionally it has been divided into a number of units, each of which reflected a distinctive geomorphological \*process. For example, in 1957 L. C. King identified four elements in his ideal profile: a crest (or 'waxing slope' or 'convex slope') dominated by \*creep; then a \*scarp (or 'free face') affected by \*rill activity and \*mass movement; followed by a debris (or 'constant') slope where \*talus accumulated; and

succeeded finally by a **\*pediment** (or ‘waning slope’) modified by **\*sheetwash**. Subsequently, a nine-unit model has gained some acceptance.

**slope stability** The likelihood that a slope will fail. This depends on the relative strengths of the forces tending to make slope material move and the forces resisting such movement. The safety factor for a slope is expressed as the ratio between the **\*shear strength** and **\*shear stress**.  $F = s/t$ , where  $F$  is the safety factor,  $s$  is total shear strength along a specified **\*shear plane**, and  $t$  is the total shear stress along that plane. For shallow slopes,  $F = c + [(gz \cos^2 b - u) \tan F]/gz \sin b \cos b$ , where  $c$  is **\*cohesion**,  $g$  is the unit weight of the **\*regolith**,  $z$  is the vertical depth of the shear plane,  $b$  is the angle of the shear plane,  $u$  is the **\*pore-water pressure** at the shear plane, and  $F$  is the angle of internal friction. A slope is stable if  $F > 1.3$  and unstable if  $F < 1$ .

**slope stabilization** The stability of slopes is important in the design of such excavations as open pits, quarries, and foundations, and in natural slopes forming cliffs, valley sides, and reservoirs, where movement may have serious consequences. Investigations into slope stability include measurements of shape, geologic structures, and soil strengths. Stabilization can be achieved by regrading, grassing, **\*dewatering**, meshing, **\*grouting**, **\*shotcreting**, **\*rock** bolting, or combinations of these.

**slow earthquake** See **SLOW SLIP**.

**slow-neutron process (s-process)** In a second-generation star, successive neutron capture on a slow timescale, termed the ‘s-process’, is considered to account for the abundance distribution of the heaviest nuclides. The neutrons are furnished by the basic sequence of energy-producing reactions, i.e. **\*hydrogen** ‘burning’, **\*helium** ‘burning’, etc., thus continuing the element-building process beyond Fe. See also **NUCLEOSYNTHESIS**.

**slow slip (silent quake, slow earthquake)** An episodic movement along a segment of a major fault in which the opposing sides are able to slide past each other slowly. A slow slip generates no seismic waves and can be detected only by extremely sensitive measurements of movements in or on the crust.

**sluff** A small avalanche (see **MASS-WASTING**) of loose material.

**slump** A type of *mass-wasting* in which loosely consolidated material moves down a slope. The material may be rock, debris, or soil, and the movement is usually rotational (see [ROTATIONAL SLUMP](#)) and occurs slowly or at a moderate speed.

**slump structure** A *sedimentary structure* consisting of overturned  *folds*, formed by the mass sliding of the semi-consolidated *sediment* downslope under the influence of gravity.

**Smålfjord** A *stage* of the *Cryogenian* period, dated at 650–640 Ma ago (Int. Commission on Stratigraphy, 2004), preceded by the *Sturtian* and followed by the *Mortensnes*.

**small circle** A circle on the surface of a sphere whose centre is not coincident with the centre of the sphere, so that a plane whose circumference is a small circle does not bisect the sphere. In a *stereographic projection* a small circle projects on to the *plane of projection* as a circle whose centre lies inside the *primitive circle*. If the small-circle plane is horizontal it will project as a circle smaller than the primitive circle but concentric with it. If it is vertical it will project as an arc, concave towards the centre of the primitive circle.

**small satellite** See [MINISATELLITE](#).

**SMAP** See [SOIL MOISTURE ACTIVE/PASSIVE MISSION](#).

**smarl** A *pelagic* or *hemipelagic* sediment (an *arl*), typically found interbedded with purer oozes in beds up to 1.5 m thick, with a composition intermediate between a non-biogenic sediment and a calcareous or siliceous ooze. It is 30% clay and 70% microfossils, with siliceous and calcareous microfossils present in approximately equal amounts. Compare [MARL](#); [SARL](#).

**SMC** See [SATURATION MOISTURE CONTENT](#).

**smectite** A family of *clay minerals* that includes *montmorillonite* and *bentonite*.

**Smith, William** (1769–1839) A land surveyor engaged on canal construction whose work led him to see that strata could be identified and

correlated by their **\*fossil** content. In 1815 he published the first geologic map and sections of England, showing the sequence of strata.

**Smithian (Nammalian)** A regional **\*stage** (249.7–247.4 Ma ago) of the Early **\*Triassic \*epoch**, preceded by the **\*Dienerian** and followed by the **\*Spathian**.

**smithsonite (calamine)** Mineral,  $ZnCO_3$ ; sp. gr. 4.4; **\*hardness** 4.5; **\*trigonal**; colour variable, shades of grey, brown, or greyish-white, but green, brown, and yellow types also occur; grey **\*streak**; vitreous **\*lustre**; crystals rare, but when they develop **\*rhombohedral** with curved faces, more usually occurs as **\*botryoidal** and stalactitic masses; **\*cleavage** perfect rhombohedral; occurs in the oxidized zone of zinc ore deposits, commonly associated with **\*sphalerite**, **\*galena**, and **\*calcite**, also as a **\*replacement** in **\*limestone**, and in **\*hydrothermal** veins; soluble in dilute hydrochloric acid, with effervescence. The green variety is used to make ornaments. It is named after the British mineralogist James Smithson (founder of the Smithsonian Institution in Washington, DC).

**Smith's rule** The formula by which the maximum possible depth of a body of unknown shape can be determined using gravity data. The body can have either a positive or negative density contrast with the surrounding rocks. The maximum depth,  $d_{\max}$ , is given by:  $d_{\max} = Ag_{\max}/(\delta g/\delta x)$ , where  $\delta g/\delta x$  is the maximum horizontal gradient of the **\*gravity anomaly**,  $g_{\max}$  is the value of the anomaly peak, and  $A$  varies between 0.86 for a three-dimensional body to 0.65 for an essentially two-dimensional body.

**smog** Naturally occurring fog mixed with visible (smoke) and/or invisible pollutants. *See also* **PHOTOCHEMICAL SMOG**.

**smoker** *See* **HYDROTHERMAL VENT**.

**SMOS** *See* **SOIL MOISTURE AND OCEAN SALINITY MISSION**.

**SMOW (Standard Mean Ocean Water)** A sea-water sample which comprises the international standard for **\*D:H** and **\*<sup>18</sup>O:<sup>16</sup>O ratios**. Differences in isotopic composition are expressed as parts per mille (‰) deviations from the isotopic composition of this standard.

**smudging** The burning of materials (e.g. oil) to produce a smoke layer that reduces the effect of radiation cooling of the air above the ground surface. It is used as a protective measure (e.g. in fruit-growing areas) especially in **\*frost hollows**.

**SNC** See **SHERGOTTITE/NAKHLITE/CHASSIGNITE METEORITES**.

**Snell's law** The ratio of the sine of the **\*angle of incidence** ( $i$ ) to the sine of the **\*angle of refraction** ( $r$ ) is a constant for any two **\*isotropic** media bounded by a common **\*interface**. The refractive index  $n$  is given by:  $n = \sin i / \sin r$ , and  $n_1 \sin i = n_2 \sin r$ , where  $n_1$  and  $n_2$  are the refractive indices of the two media. The law also indicates that a **\*P-wave** incident upon a boundary will be reflected and refracted partly as a P-wave and partly as an **\*S-wave**. The law was formulated by the Dutch astronomer and mathematician Willebrord Snell (1591–1626).

**snout** The steep, terminal zone of a **\*glacier**. It is usually heavily loaded with debris.

**snowball Earth** A period during which ice covers the entire surface of the Earth, except for the highest mountains, and mean temperatures are about  $-50^{\circ}\text{C}$ . This condition is believed to have occurred four times between 750 and 580 Ma ago, on each occasion lasting for approximately 10 million years.

**snowblitz theory** A theory which proposes that following a bad winter with heavy snowfall, snow persists in lowland areas throughout the summer. This increases the **\*albedo** and thus reduces the amount of solar warming of the ground. More snow is added during the next winter, and more snow may thus accumulate year by year. An **\*ice cap** may develop, and glaciation may occur after only a few hundred years. Such a sequence of events is more likely in high than in low latitudes.

**snowflake** The result of the growth of ice crystals in a varied array of shapes. Very low temperatures usually result in small flakes; formation at temperatures near freezing point produces numerous crystals in large flakes.

**snow-gauge** A modified **\*rain-gauge** used for collecting and measuring snowfall. The recording is made after the snow has melted and reported as the rainfall equivalent to correct for the varying amounts of air entrained between **\*snow grains** in different types of snow.

**snow grain** Small ice particle precipitated as a usually flattish grain.

**snow line** **1.** The lower limit of permanent snow cover. The height of the line varies with latitude; locally it also varies with aspect, because of the relationship to prevailing winds and quantity of snow deposited, insolation, and to summer temperatures, etc. **2.** The minimum distance from a star at which volatiles will condense.

**soapstone** See TALC.

**soda lake** See NATRON LAKE.

**sodalite** An important group of *\*silicate* minerals belonging to the *\*feldspathoids* and including sodalite  $\text{Na}_8[\text{Al}_6\text{Si}_6\text{O}_{24}]\text{Cl}_2$ , *\*nosean*, *\*häuyne*, *\*cancrinite*, *\*lazurite*, and *\*scapolite*; sp. gr. 2.27–2.88; *\*hardness* 5.5–6.0; greyish to bluish, yellow; occurs as *\*rhombohedra* or *\*massive*; *\*vitreous* *\*lustre*; occurs in *\*nepheline-syenites* in association with *\*nepheline* and *\*fluorite*, and in metasomatized (see METASOMATISM) calcareous rocks near *\*alkaline* *\*igneous* intrusions.

**soda nitre (Chile saltpetre, nitratine)** Very soluble *\*mineral*,  $\text{NaNO}_3$ ; sp. gr. 2.2–2.3; *\*hardness* 1–2; *\*trigonal*; normally colourless or white, with various darker colours due to impurities; white *\*streak*; *\*vitreous* *\*lustre*; *\*crystals* rhombohedral, also occurs *\*massive*; *\*cleavage* perfect, rhombohedral; occurs in arid regions as surface deposits together with *\*gypsum*, *\*halite*, and other soluble nitrates and *\*sulphates*; deliquescent. It is worked as a source of nitrate.

**sodication** In soils, an increase in the percentage of exchangeable sodium. Sodium adsorbs on to *\*cation-exchange* sites in the soil, causing soil *\*aggregates* to disperse, which closes soil *\*pores* and renders the soil impermeable to water. See also SODIUM-ADSORPTION RATIO.

**sodic soil** **1.** Soil with a sodium content sufficiently high to interfere with the growth of most crop plants. **2.** Soil with more than 15% exchangeable sodium.

**sodium-adsorption ratio (SAR)** Describes the tendency for sodium *\*cations* to be adsorbed at *\*cation-exchange* sites in soil at the expense of other cations, calculated as the ratio of sodium to calcium and magnesium in the soil; more precisely, it is the amount of sodium divided by the square

root of half the sum of the amounts of calcium and magnesium, where *\*ion* concentrations are given in milliequivalents per litre. A low sodium content gives a low SAR value. In practice, allowance must be made for other reactions within the soil that do not involve sodium but that affect concentrations of calcium and magnesium. The SAR value is most likely to be changed by irrigation water.

**sodium feldspar** See [ALKALI FELDSPAR](#).

**sodium-sulphate soundness test** A method of testing the *\*weathering* resistance, particularly to frost action, of building materials. A sample is soaked in saturated sodium-sulphate solution, drained, and dried. This is repeated and the sample examined for cracks. The method simulates the stresses due to frost action.

**SOFAR channel (sound channel)** Acronym for the *SOund Fixing And Ranging* channel, a zone in the oceanic water column at a depth of about 1500 m where the velocity of sound is at a minimum value. Sound passing through the zone is refracted upwards or downwards back into the zone, with little loss of energy, causing sound energy to be trapped in a zone of well-defined depth. The SOFAR channel may be used for the transmission of sound over long distances, exceeding 28 000 km, and can be used to track free-drifting, subsurface, *\*neutrally buoyant floats*.

**soil** **1.** The natural, unconsolidated, mineral and organic material occurring above bedrock on the surface of the Earth; it is a medium for the growth of plants. **2.** In engineering geology, any loose, soft, and deformable material, e.g. unconsolidated *\*sands* and *\*clays*.

**soil air** The soil atmosphere, comprising the same gases as in the atmosphere above ground, but in different proportions: it occupies the *\*pore space* of an unsaturated *\*soil*.

**soil anchor** See [ANCHOR](#).

**soil association** **1.** Group of *\*soils* forming a pattern of soil types characteristic of a geographical region. **2.** Mapping unit used to denote the distribution of soil types where the scale of the map does not require or permit the identification of individual soils. See also [SOIL COMPLEX](#).

**soil-atmosphere survey** A method for finding geochemical anomalies, which uses geochemical features of the gases trapped in *\*soil \*pore space*. The method can be used only to find *\*ore minerals* which, on *\*weathering* or as a result of their radioactive decay, liberate gases which can readily be analysed (e.g. radon, released by the decay of uranium).

**soil borrow** The transference of material from elsewhere for refilling excavations, etc.

**soil complex** Mapping unit used to denote the distribution of *\*soils*: it is more precise than a *\*soil association*, and is used where soils of different types are mixed geographically in such a way that the scale of the map makes it undesirable, or impractical, to show each one separately.

**soil conservation** The protection of the *\*soil* by careful management, to prevent physical loss by *\*erosion* and to avoid chemical deterioration (i.e. to maintain soil *\*fertility*).

**soil formation** The action of the combined primary (*\*weathering* and *\*humification*) and secondary processes to alter and to rearrange mineral and organic material to form *\*soil*, involving the differentiation of *\*soil profiles* and the formation of loose soil from consolidated rock material. *See also PEDOGENESIS.*

**soil geomorphology (pedogeomorphology)** The study of the relationship between types of *\*soil* and land-forms. This involves analysing the way geomorphological and pedological processes interact.

**soil grading curve** A graph of *\*grain size* plotted on a horizontal logarithmic axis against percentage on an arithmetical vertical axis. A point on the curve gives the percentage by weight of material smaller in size than that at the given point on the graph. *See GRADING CURVE.*

**soil horizon** A relatively uniform *\*soil* layer which lies at any depth in the *\*soil profile*, which is parallel, or nearly so, with the soil surface, and which is differentiated from adjacent horizons above and below by contrasts in mineral or organic properties. Soil horizons are grouped primarily into O, A, B, and C horizons. O horizons (formerly known as Ao horizons) comprise organic material at the surface. A horizons are surface horizons of mixed organo-mineral composition. Where mineral matter has been lost, the A horizon is sometimes called the E (for eluviated) horizon. Where they are

present, B horizons are usually located in the middle of the sequence, and are horizons into which material (mineral and organic) is deposited, thus altering the character of the horizon. C horizons are soil **\*parent materials**, weathered but not otherwise altered by pedogenic processes. The underlying unweathered material is sometimes called the D or R horizon. In addition to these, surface litter may form an L horizon, above a layer of fermented material (F horizon) and, below that, humified material (H horizon), and a mineral crust, often cemented, is sometimes called the K horizon. See **PEDOGENESIS**.

**soil individual** See **POLYPEDON**.

**soil line** In **\*remote sensing**, the line which runs at 45° on a plot of **\*digital number** values of red wavelengths of light against digital number values of very-**\*near infrared** wavelengths of light. Soils plot very close to this line, vegetation plots away from this line having a higher tendency to reflect very-near infrared. See also **VEGETATION INDEX**.

**soil lithology** The relationship between key characteristics of a soil (e.g. organic carbon content, pH, **\*cation-exchange capacity**, percentage **\*clay**, total phosphorus, etc.) and the **\*parent material** from which it is derived, giving rise to fourteen soil types: **\*alluvial**, basaltic (see **BASALT**), **\*boulder clay**, **\*calcareous**, **\*chalk**, **\*clay**, **\*diorite**, glacial **\*till**, **\*granite**, **\*loess**, sandy, **\*schist**, **\*shale**, and volcanic.



<http://www.environmentdata.org/archive/vocabpref:19348>

- Lithological soil types.

**soil management** A variety of practices and operations with respect to **\*soil**, that aid the production of plants; normally they are planned to allow for sustained yield in the future.

**soil mechanics** Study of the mechanical properties of loose or unconsolidated particles, especially their composition, shear resistance, and the effects of water. It is applied to **\*soils** to determine their suitability for building sites, mining, etc., and to engineering problems dealing with the stability of **\*foundations** due to mechanical and chemical **\*weathering** of rocks. See **SHEAR STRENGTH**.

**Soil Moisture Active/Passive mission (SMAP)** A \*NASA mission to monitor global soil-moisture mapping. The satellite was launched on 31 January 2015, from California, into a Sun-synchronous dawn–dusk orbit at an altitude of 685 km.

**Soil Moisture and Ocean Salinity mission (SMOS)** A satellite mission by the \*European Space Agency to improve understanding of the hydrological cycle, supplying data for modelling of weather and climate. It was launched on 2 November 2009, from the Plesetsk Cosmodrome, Russia, into a Sun-synchronous \*polar orbit at a mean altitude of 755 km.

**soil-moisture content** The ratio of the volume of contained water in a \*soil compared with the entire soil volume. When a soil is fully saturated, water will drain easily into the underlying unsaturated rock. When such drainage stops, the soil still retains \*capillary moisture and is said to contain its \*field-capacity moisture content. Further drying of the soil (e.g. by evaporation) creates a soil-moisture deficit, which is the amount of water which must be added to the soil to restore it to field capacity, measured as a depth of \*precipitation.

**soil-moisture deficit** See SOIL-MOISTURE CONTENT.

**soil-moisture index** See MOISTURE INDEX.

**soil-moisture regime** The changing state of soil moisture through the year, which reflects the changing balance of monthly \*precipitation and \*potential evapotranspiration above the ground surface. When the latter exceeds the former the period is one of soil-moisture deficit in the annual regime.

**soil profile** A vertical section through all the constituent \*horizons of a soil, from the surface to the relatively unaltered \*parent material.

**soil separates** Size divisions of mineral particles (\*sand, \*silt, and \*clay) that comprise the fine earth, each particle being less than 2 mm in diameter.

**soil series** Basic unit of soil mapping and classification: all \*soils in a series have similar profile characteristics and have developed from the same \*parent material.

**soil structure** Grouping of individual \*soil particles into secondary units of \*aggregates and \*peds.

**soil survey** **1.** Systematic examination and mapping of *\*soil* in the field. **2.** Chemical analysis of soil for the detection of geochemical anomalies. See [GEOCHEMICAL SOIL SURVEY](#); [ORIENTATION SURVEY](#).

**soil taxonomy** The classification of types of soil, in a manner similar to that used for biological classification. In one of the most widely used systems, devised by workers at the US Soil Survey, within the *\*USDA*, soils are divided into 11 orders: *\*alfisols*, *\*andosols*, *\*aridisols*, *\*entisols*, *\*histosols*, *\*inceptisols*, *\*mollisols*, *\*oxisols*, *\*spodosols*, *\*ultisols*, and *\*vertisols*. These orders are further subdivided into suborders, great groups, families, and soil series, defined by *\*diagnostic* horizons. The *\*World Reference Base* for Soil Resources is the principal international classification.

**soil variant** *\*Soil* sufficiently different in properties from adjacent soils to warrant the use of a new *\*soil-series* name, but occupying a geographical area too small to warrant the issuing of such a new name.

**soil-wash** See [RAIN-WASH](#).

**soil-water zone (unsaturated zone, vadose zone)** The zone between the ground surface and the *\*water-table*. Water is able to pass through this zone to reach the water-table, but while in the zone it is not given up readily to wells because it is held by *\*soil* or rock particles and capillary forces. See [GROUNDWATER](#).

**Sojourner** See [MARS PATHFINDER](#).

**sol** **1.** Colloidal solution (see [COLLOID](#)) or dispersion of solid particles in a liquid, as in a completely fluid mud. Compare [QUICK CLAY](#); [GEL](#). **2.** One martian day (= 24.7 hours).

**sola** See [SOLUM](#).

**solar abundance of elements** Studies of the solar spectrum have determined the relative abundances of about 70 elements in the *\*Sun*'s atmosphere. Hydrogen and helium predominate, and in general abundance decreases with increasing atomic number. A few exceptions, however, e.g. silicon and iron, have a high abundance, probably correlated with nuclear binding energies and nuclear stability. Spectra originate from the outer layers of the Sun, so their value in determining the total solar abundance of

elements should not be over-emphasized, but total abundances of elements are much less important than their relative proportions.

**solar constant** The mean intensity of the solar beam in free space (i.e. before penetrating the Earth's atmosphere) at the average distance of the *\*Earth* from the *\*Sun*. The intensity is not strictly constant for all the wavelengths of radiation involved. The amount of variability is still a subject of debate, but is certainly very small apart from the long-term development in the history of the Sun.

**solar cosmic rays** See COSMIC RADIATION.

**solar flare** A short-lived, cataclysmic outburst of solar material, driven by magnetic forces, from a relatively small area of the solar surface and generating particles with energies in the range 1–100 MeV which produce track records, e.g. in exposed lunar minerals.

**solarimeter** Instrument for measuring *\*solar radiation*.

**Solar Irradiance and Earth Radiation Budget (SERB)** A French 3-unit *\*CubeSat* mission to study the Earth–Sun relationship. It is scheduled for launch in 2020 into a Sun-synchronous dawn–dusk orbit at an altitude of  $680\pm 30$  km.

**solar magnetic variation** See DIURNAL VARIATION (2).

**solar nebula** See SOLAR SYSTEM.

**solar pond** See SALINA.

**solar radiation** Electromagnetic radiated energy from the *\*Sun*. This is the dominant energy input to the *\*Earth* and is intercepted by the *\*atmosphere* and absorbed at the surface. See also RADIATION BUDGET.

**Solar Radiation and Climate Experiment (SORCE)** A NASA-sponsored *\*minisatellite* mission, launched on 25 January 2003, from Cape Canaveral, into a circular orbit that crosses the equator at  $40^\circ$ , at a height of 645 km. SORCE carries four instruments, the total irradiance monitor (TIM), spectral irradiance monitor (SIM), solar stellar comparison experiment (SOLSTICE), and the extreme ultraviolet photometer experiment (XPS). These provide precise measurements of solar radiation in the UV and

visible and near-infrared (VNIR) wavebands, providing data to help understand the influence of variations in solar radiation on climate.



<https://earth.esa.int/web/eoportal/satellite-missions/s/sorce>

- **SORCE** (Solar Radiation and Climate Experiment).

**solar system** The system that consists of the central **\*Sun** (G spectral type star), around which orbit eight planets, three dwarf planets (Pluto, Ceres, and Eris), about 110 **\*satellites**, about 3000 discovered **\*asteroids**, and probably  $10^{12}$  **\*comets**. Most bodies lie close to the plane of the **\*ecliptic**. The age of the solar system, 4.56 billion years obtained from **\*meteorites**, marks the formation of the system from a rotating cloud of dust and gas (the solar nebula).

**solar wind** General term for the stream of high-energy particles (mainly protons, electrons, and alpha particles) emitted by the **\*Sun**. The particles have velocities of hundreds of kilometres per second and 'wind' strength is greatest during periods of maximum solar activity. In the neighbourhood of the **\*Earth** the solar wind has velocities in the range of 300–500 km/sec and an average density of  $10^7$  ions/m<sup>3</sup>. *See also* **COSMIC RADIATION**.

**sole mark** A mark by which a **\*sole structure** may be recognized and identified.

**Solenopora** *See* **RHODOPHYCEAE**.

**sole structure** The term applied to a group of **\*sedimentary structures** found on the base of beds. Sole structures are mostly formed by the scouring action of a current, or by the passage of an implement (**\*tool**) over a muddy substrate, followed by the infilling of the scour by **\*sands** (**\*scour** and fill). Sole structures include **\*flute marks**, produced by turbulent flows of water; **\*skip**, prod, bounce, drag, and **\*groove marks**, caused by the transport of an object over the mud; and **\*load casts**, formed by the sinking of dense sand into underlying layers of less dense mud. Sole structures provide an important means of determining the 'way-up' of beds, and in many cases give an important indication of palaeocurrent direction. *See* **PALAEOCURRENT ANALYSIS**.

**sole thrust (basal thrust)** In a *\*thrust* terrain, the lowest regional thrust surface. *See also* FLOOR THRUST.

**solfataric activity** The quiet escape of hot, sulphur-rich gases from recently emplaced volcanic bodies. The name is derived from the fields of sulphurous gas vents at the Solfatara crater, north of Naples, Italy. When cooled by the atmosphere the escaping gases deposit many minerals, including chlorides, sulphur and *\*hematite*.

**solid** Applied to a map (e.g. in Britain one published by the Geological Survey) which depicts the *\*outcrop* patterns of rocks unobscured by their cover of recent superficial deposits (e.g. glacial, *\*alluvial*, or marine *\*sediments*).

**solid-melt equilibrium** The degree of *\*partial melting* which, in any particular rock, is a function of temperature, pressure, and the availability of water, and is usually controlled by the abundance and thermal stability of hydroxyl-bearing minerals, e.g. *\*muscovite*, *\*biotite*, and *\*hornblende*.

**solid solution** Solid crystalline *\*phases* representing a mixture of two or more *\*end-members* and which may vary in composition within finite limits without the appearance of another phase. A mineral may exhibit solid solution involving atoms of an *\*isomorphous* mineral; for example, magnesium and iron *\*ions* are similar in size, and complete solid solution occurs between *\*forsterite* ( $Mg_2SiO_4$ ) and *\*fayalite* ( $Fe_2SiO_4$ ) end-members of the *\*olivine* series, in the most common type of solid solution, known as 'substitutional solid solution'. *See also* IONIC SUBSTITUTION.

**solid-state imaging camera (SSI)** In *\*remote sensing*, an instrument that operates in the visible spectrum and uses a *\*charge-coupled* device to enhance its images.


**solidus** The position of points marking the boundary between complete solid and liquid/solid at equilibrium, in a *\*temperature–composition diagram*. In *\*binary systems* the solidus is a straight or curved line, in *\*ternary systems* a flat plane or curved surface.

**solifluction (solifluxion)** Downhill movement of *\*regolith* that has been saturated with water. It was originally described in *\*periglacial* regions (*see* GELIFLUCTION), but the term was subsequently widened to include all

environments. The thick regolith of the humid tropics is particularly prone to solifluction after intense rainfall.

**solifluxion** See SOLIFLUCTION.

**solitary corals** Those corals where a single *\*corallite* makes up the *\*corallum*. Shape and size range from an extremely low cone with an apical angle of 120° or more (patellate), through discoid (button-shaped), to horn-shaped and slender with an apical angle of 20° or less (ceratoid). A steep, conical shape with an apical angle of 40° is called ‘trochoid’; a subcircular, parallel-sided corallite ‘cylindrical’; cylindrical corallites that bend crookedly in a worm-like manner are ‘scoleoid’; a corallite shaped like the toe of a slipper is ‘calceolid’. Other shapes are also described.



**Discoid**



**Patellate**

**solodic soil** Leached, formerly *\*saline soil*, associated with semi-arid tropical environments, in which the A *\*soil horizon* has become slightly acid, and the B horizon is enriched with sodium-saturated *\*clay*. The term was used in soil classification systems derived from early Russian systems based on the work of V. V. *\*Dokuchayev*, but is no longer used in soil classification.

**solonchaks** A reference soil group in the *\*World Reference Base for Soil Resources* classification. Solonchaks have a *\*soil horizon* more than 15 cm

deep that is enriched with soluble **\*salts**, located at the surface or a little way below it. These soils often form from recent **\*alluvial** deposits.

**solonetz** Mineral soil at a transitional stage of **\*leaching** or a **\*saline soil** during solodization, in semi-arid, tropical environments; it has a sandy, acid, A **\*soil horizon** and a B horizon partially enriched with sodium-clay. The term was used in earlier systems of soil classification. Solonetz soils fall within the order **\*aridisols** in the US **\*soil taxonomy** and solonetz is a soil group in the **\*World Reference Base for Soil Resources**.

**solstice** The time of most northerly or southerly declination of the sun from the equator. In the northern hemisphere, the summer solstice is around 22 June and the winter solstice around 22 December. These dates are reversed for the southern hemisphere.

**solubility product** Constant which describes the product of ionic concentrations of any slightly soluble salt. The solubility of a substance is the maximum amount of that substance which can be dissolved in a **\*solution** that is in equilibrium with a solid source of the solute. The main technique for studying solubility involves the use of the solubility product, which is the total number of **\*ions** of each type in a compound that can coexist in a solution. At a given temperature and at equilibrium the value for the solubility product will always be the same. For example, in a solution of silver chloride, AgCl dissociates to  $\text{Ag}^+$  and  $\text{Cl}^-$ . The solubility product  $K_{\text{sp}}$  is the product of their concentrations:  $K_{\text{sp}} = [\text{Ag}^+][\text{Cl}^-]$  mole<sup>2</sup>/litre<sup>2</sup>. The solubility of AgCl in moles per litre is equal to the activity of  $\text{Ag}^+$  or  $\text{Cl}^-$  as one mole of AgCl dissolves in water to give one mole of each in solution. The value of  $K_{\text{sp}}$  for AgCl at 25°C is  $10^{-9.8}$ .

**solum** (*pl. sola*) The upper part of a **\*soil profile**, above the **\*parent material**, in which processes of soil formation occur, and within which most plant roots and soil animals are found.

**solution 1.** A physically homogeneous mixture of two or more substances in which solid, liquid, or gaseous **\*phases** may combine in one of those phases. A constituent of a solution can be separated out by changing its phase, e.g. boiling, condensing, or freezing. Where a solution is formed by dissolving a quantity of one substance in a larger quantity of another, the smaller quantity is called the 'solute', the larger quantity, the 'solvent'.

Compare COLLOID. 2. A **\*weathering** process by which weakly bonded ionic components of minerals are detached through the attraction of water molecules (which carry a positive electrical charge at one end and a negative charge at the other, although they are neutral overall), and then carried away from the weathering environment. **\*Halites** and the sulphates and carbonates of magnesium and calcium are especially vulnerable. Solution is usually the first stage of **\*chemical weathering**.

**solution channel** An elongate void within a rock, which has been enlarged by the **\*solution** action of moving **\*groundwater** on the rock itself. Solution channels are most commonly associated with **\*carbonate** rocks, and groundwater in them can flow as fast as in a river on the ground surface. See KARSTIC AQUIFER.

**solution cleavage** A spaced, usually disjunctive **\*cleavage** which is common in **\*quartzites** and limestones. Solution cleavages commonly contain zones of relatively insoluble minerals indicating that **\*pressure solution** has operated. Some authorities (e.g. C. McA. Powell, 1979) suggest that the term 'solution cleavage' should not be used in descriptions of cleavage as it implies the mode of origin.

**solution pan** See PANHOLE.

**solution pipe** A cylindrical, near-vertical pipe that is developed at a **\*joint** intersection in a **\*karst** environment. It is caused by a local increase in the rate of **\*carbonation** resulting from enhanced drainage.

**Solván** A Russian-Kazakhstanian **\*stage** of the Middle **\*Cambrian**, underlain by the **\*Toyonia** and overlain by the **\*Mayan** and dated at 513–502 Ma ago (Int. Commission on Stratigraphy, 2004).

**solvus** In a geochemical system, a line or surface which separates a homogeneous **\*solid solution** from a field of several **\*phases** which may form by **\*exsolution** or **\*incongruent melting**.

**solvus temperature** See HYPERSOLVUS GRANITE.

**Somali Plate** See AFRICAN PLATE.

**sombric horizon** Subsurface **\*soil horizon** of well-drained, mineral, tropical and subtropical soils into which **\*humus** has leached downward.

Base saturation is less than 50%. It is a *\*diagnostic horizon*.

**Somoholoan** See ASSELIAN; SAKMARIAN.

**sonar** See SIDE-SCAN SONAR; SONIC LOG; ECHO-SOUNDING.

**sonde** See RADIOSONDE; RAWINSONDE; SELF-POTENTIAL SONDE; SONIC SONDE; WELL LOGGING.

**sonde self-potential** See SELF-POTENTIAL SONDE.

**sonic log** See CONTINUOUS VELOCITY LOGGING; SONIC SONDE.

**sonic sonde** An instrumental package containing two seismic-energy sources and four *\*geophones* which allow the measurement of the *\*seismic velocities* of the rocks in a *\*borehole* as the sonde is pulled through it. The record is a sonic log (velocity log). See also WELL SHOOTING.

**sonobuoy** A disposable, free-floating buoy used in large-scale marine *\*seismic-refraction* surveys. One or more hydrophones are suspended from the buoy and detect the *\*head waves* which are then transmitted back to the firing ship, timed, and recorded. After a certain time the buoys are designed to sink automatically.

**sonograph** A graphic presentation of reflected sound waves from a sonar scanner.

**Sorby, Henry Clifton** (1826–1908) An English amateur scientist, Sorby studied estuarial and inland waters of England, but he is best known for developing the study of rocks in *\*thin sections*, using the techniques invented by *\*Nicol*. He was the first to show that individual mineral crystals and grains could be identified using this process. He also used the method to study *\*meteorite* sections.

**SORCE** See SOLAR RADIATION AND CLIMATE EXPERIMENT.

***Sordes pilosus*** Discovered in 1971, one of the first *\*pterosaurs* known to have been covered in thick fur: the name means ‘hairy filth’. The indication is that this reptile and its close relatives were *\*homoiotherms*. It was found in Upper *\*Jurassic* sediments of Chimkent, Kazakhstan, and was a small, toothed pterosaur with a long tail.

**sorosilicate** See SILICATES.

**sorption** A blanket term covering \*absorption, \*adsorption, and \*ion exchange.

**sorted biosparite** See FOLK LIMESTONE CLASSIFICATION.

**SORTIE** See SCINTILLATION OBSERVATIONS AND RESPONSE OF THE IONOSPHERE TO ELECTRODYNAMICS.

**sorting** An expression of the range of grain sizes (see PARTICLE SIZE) present in a \*sediment. A well-sorted sediment is characterized by a narrow range of grain sizes, whereas a poorly sorted sediment contains a wide range of grain sizes.

**Soudleyan (Kirkfield)** A \*stage (457–458 Ma ago) of the \*Ordovician in the Middle \*Caradoc, underlain by the \*Harnagian and overlain by the \*Longvillian.

**sound channel** See SOFAR CHANNEL.

**sounder (radio)** An instrument which emits a continuous series of short pulses of electromagnetic energy towards a planetary or \*satellite surface. The return signal provides a map of the subsurface electrical conductivity, which can be used to infer subsurface structure, as well as information on topography. This technique was used on the Apollo 17 lunar mission to study subsurface structure.

**sound speed** The rate at which sound energy moves through a medium. In sea water this is between 1400 and 1550 m/s. In sea water, the speed of sound is a function of temperature, \*salinity, and pressure due to depth. At a salinity of 34.85‰ and a temperature of 0 °C, the speed of sound is 1445 m/s. It increases by approximately 4 m/s for each degree Celsius rise in temperature, by 1.5 m/s for each 1‰ increase in salinity, and by 18 m/s for each 1000 m increase in depth. It is calculated by  $a = \sqrt{\gamma RT}$ , where  $a$  is the speed of sound,  $\gamma$  is the ratio of specific heats (= 1.4 for air at standard temperature and pressure),  $R$  is the gas constant (286 m<sup>2</sup>/s<sup>2</sup>/K for air), and  $T$  is the temperature in kelvins. In air at 20 °C the speed of sound is approximately 343 ms<sup>-1</sup>.

**source region (for air masses)** Extensive areas of essentially uniform surface conditions over land or water, typically of large-scale air \*subsidence and lateral divergence, where \*air masses acquire their initial properties.

**source rock** 1. \*Sediment (usually \*shale or \*limestone) in which \*hydrocarbons originate; it contains more than 5% organic matter and has the potential to generate \*petroleum. 2. Any parent rock from which later sediments are derived.

**South American Plate** One of the major present-day lithospheric \*plates, extending from the \*Mid-Atlantic Ridge in the east to the subducting \*Nazca Plate in the west, with most of the boundaries with other plates (i.e. \*Antarctic, Scotia, \*Caribbean, and \*North American) being \*transform faults.

**South-east Pacific Plate** A lithospheric \*plate which is now coupled with the \*Antarctic Plate, but whose oceanic \*lithosphere is interpreted to have been subducted under southern Chile and the Antarctic Peninsula.

**southerly burster** Regional wind in southern and south-eastern Australia, characterized by a rapid shift in direction from north-west to south in the rear of a \*cold front. Such winds are especially prevalent between October and March. The change to a southerly wind can bring a great increase in wind speed accompanied by a rapid and marked fall in temperature. Such conditions are akin to \*line squalls, and related to the S. American \*pamperos.

**Southern Ocean** See ANTARCTIC OCEAN.

**southern oscillation** A fluctuation of the intertropical atmospheric circulation, in particular in the \*Indian and \*Pacific Oceans, in which air moves between the SE Pacific subtropical high and the Indonesian equatorial low, driven by the temperature difference between the two areas. The general effect is that when pressure is high over the Pacific Ocean it tends to be low in the Indian Ocean, and vice versa. The pressures are measured at Tahiti and Darwin, Australia. The phenomenon is strongly linked to \*El Niño.

**sövite** A type of \*carbonatite that consists largely of \*calcite accompanied by minor \*magnetite, and \*apatite with or without \*phlogopite.

**SP** See SELF-POTENTIAL SONDE.

**spaced cleavage** See FRACTURE CLEAVAGE.

**space lattice** See LATTICE.

**space–time substitution** A technique for analysing land-forms that is based on a theory of how particular land-forms develop over time as a result of geomorphological processes. Thus, the appearance (i.e. shape) of a landscape changes over time and this fact can be used to determine its past and future appearance. The technique was pioneered by W. M. *\*Davis* (see *DAVISIAN CYCLE*), but was introduced by Charles Darwin, in his study of the development of *\*atolls*, fringing *\*reefs*, and *\*barrier reefs*.

**spallation** 1. (nuclear) A nuclear reaction involving the ejection of many particles from an atomic nucleus, following a collision with a high-energy particle. Both the mass number and the atomic number of the target nucleus are changed by the event. 2. (planetary geol.) The removal of the surface layers of a rock by the interaction of a compressional shock wave with the surface, caused by micrometeorite impact. See ‘*ZAP PITS*’.

**spandrels of San Marco** An analogy used, in a classic paper by Stephen Jay Gould and Richard Lewontin, to indicate how non-adaptive characters may arise in evolution. Spandrels are the spaces left between the tops of neighbouring arches in churches (in this case, St Mark’s cathedral, in Venice); these spaces, not related to the functional architecture, are free to be decorated in a non-functional fashion.

**spar** Any crystal that breaks easily into fragments with clearly visible faces. By extension, a mining term describing any white or light–coloured, non-metallic mineral with *\*vitreous \*lustre* and well developed *\*cleavage*, resembling *\*gypsum*, for which the Old English name was *spærstān*.

**sparite** Sparry *\*calcite \*cement*. Sparite is the coarse crystalline calcite cement which fills *\*pore spaces* in many *\*limestones* after deposition, formed by the *\*precipitation* of calcite from *\*carbonate*-rich solutions passing through the pore spaces in the *\*sediment*.

**-sparite** See FOLK LIMESTONE CLASSIFICATION.

**sparkler** A seismic source created by the rapid formation of a gas-plasma bubble from the ionization of sea water after the discharge of a high-voltage spark from a comb of electrodes contained within a frame. It provides good resolution (1–2 m) with limited depth penetration (less than 100 m).

**sparse biomicrite** See FOLK LIMESTONE CLASSIFICATION.

**spastolith** A *\*grain* which, being composed of soft material, became squashed and deformed by mechanical *\*compaction* during burial.

**Spathian** The final *\*stage* (247.4–245 Ma ago) of the Early *\*Triassic \*epoch*, preceded by the *\*Smithian* and followed by the *\*Anisian*.

**spathose iron** See *SIDERITE*.

**spatial frequency** In *\*remote sensing*, the frequency of change per unit distance across an image. High spatial frequencies include those changes which occur in very close proximity, such as fine lines, low spatial frequencies include those changes which occur over greater distances, such as broad bands. The ability of the human eye to discern spatial frequency is limited and so selective removal of certain spatial-frequency ranges within an image may result in a more interpretable image with less noise. *See also* SPATIAL-FREQUENCY FILTER.

**spatial-frequency filter** In *\*remote sensing*, a filter used to enhance the appearance of the spatial distribution of data in an image to make it more interpretable to the human eye. Spatial-frequency filters examine the spatial variations in *\*digital number* of an image and are used to modify the image by selectively suppressing or separating certain *\*spatial-frequency* ranges. Spatial-frequency filters include *\*directional filter*, *\*high-pass filter*, medium-pass filter, and low-pass filter. *See also* MEDIAN FILTER.

**spatter** Fluid basaltic *\*pyroclasts* which accumulate by fallout from a *\*Strombolian* volcanic *\*eruption* column to form a rampart around the vent. The individual clots are so fluid when they land that they often mould together, like flattened pancakes.

**spatter cone (dribble cone)** Small (usually 5–20 m high), volcanic cone built from *\*tephra* blown out as clots of relatively fluid basaltic *\*lava*.

**spatter-fed flow** See HAWAIIAN ERUPTION.

**specialization** Degree of adaptation of an organism to its environment. A high degree of specialization suggests both a narrow habitat or niche and significant interspecific competition.

**species** (*sing.* and *pl.*) **1.** A class of particular chemical individuals all of which are similar, e.g. *\*ions*, atoms, or molecules. **2.** See **CLASSIFICATION**.

**species longevity** The persistence of species for long periods of time, characterizing, for example, members of the *\*Gastropoda* and *\*Bivalvia*.

**species selection** A postulated evolutionary process in which selection acts on an entire species population, rather than individuals. This might occur, for example, as a consequence of the geographical range of a population, which affects the population as a whole and, possibly, its longevity or development.

**species zone** See **TAXON RANGE ZONE**.

**specific gravity (sp. gr.)** The ratio of the weight of a substance to the weight of an equal volume of water, expressed as a number. For example, the weight of a given volume of *\*quartz*, with a specific gravity of 2.65, is 2.65 times that of the same volume of water. The average sp. gr. of metallic minerals is about 5. See also **DENSITY**.

**specific-gravity determinations** In soils, for engineering computations, grains are weighed in a calibrated glass container (*\*pycnometer*), carefully excluding air, to relate the mass of the sample to its volume. The *\*specific gravity* of solid rocks can be determined using a Walker balance, and can be carried out on dry or water-saturated samples. See also **DENSITY DETERMINATION**.

**specific humidity** The ratio of the mass of water vapour to a unit mass of air including the water vapour. See also **HUMIDITY**; **MIXING RATIO**.

**specific retention** The ratio of the undrained water to the total water in a rock, the undrained water being water contained in rock voids or *\*pore spaces*, from which it cannot be recovered by drainage or pumping. It is retained against the action of gravity by molecular attraction and *\*capillarity*.

**specific yield** **1.** The ratio of the water drained from a rock under the influence of gravity, or removed by pumping, to the total volume of the rock voids or **\*pore space** in the drained rock. The difference is caused by the retention of water in the rock, due to molecular attraction and **\*capillarity**. See **SPECIFIC RETENTION**. **2.** The volume of water released by a falling **\*water-table** from a given volume of fully saturated rock.

**spectra** See **SPECTRUM**.

**spectral hue** A **\*hue** which is present in the spectrum of colours produced by splitting white light with a prism. Spectral hues include red, green, and blue. Compare **NON-SPECTRAL HUE**.

**spectral radiance** The **\*radiance** of a specified wavelength of **\*electromagnetic radiation**.

**spectrochemical analysis** An analytical technique in which a sample is heated to a high temperature, usually in a carbon arc, to produce emission lines whose intensities are proportional to the abundance of elements present. Line intensities may be recorded on photographic plates or measured directly by a photosensitive device, e.g. a photomultiplier. See also **SPECTRUM**.

**spectrograph** Analytical instrument used mainly for elemental analysis.

**spectrometer** An instrument which, with associated equipment, furnishes the ratio, or some function of a ratio, of the radiant power of two electromagnetic beams as a function of their spectral wavelength. It requires a source of radiation (in emission spectrometers the sample serves as its own source), a means of distinguishing between different radiation frequencies, and includes narrow band filters, a prism, a diffraction grating, a system of slits to isolate a narrow band of radiation, a sample-containing system, and a photodetector, amplifier, and output device (meter, recorder, VDU, etc.) See **GAMMA-RAY SPECTROMETRY**; **MASS SPECTROMETRY**.

**spectrophotometer** An instrument for determining the intensity of the light absorbed by a compound (usually in **\*solution**). The light absorbed at a given wavelength is proportional to the concentration of the compound in the solution.

**spectroradiometer** A *spectrometer* which measures very narrow wavelengths of the *electromagnetic radiation* radiated or reflected by a surface. It is used in *remote sensing* to establish the spectral characteristics of a surface material.

**spectroscope** An instrument used in spectroscopy, whose main features are a slit and collimator, prism, telescope, and counter. A parallel beam of radiation is passed through the prism, so dispersing different wavelengths through different angles of deviation, which can be measured.

**spectroscopic binary** See STAR PAIR.

**spectrum** (*pl.* spectra; **optical emission spectrum**) A series of lines (line spectra), produced as *electrons* return to their original energy levels and emit excess energy as infrared, visible, or ultraviolet light of characteristic wavelengths, after atoms have been heated strongly and valence electrons in the outer shell have moved to higher energy levels. Each element has a characteristic line spectrum. The intensity of each line is related to the concentration of the element being excited.

**specular** See SPLENDENT.

**specularite** See HEMATITE.

**specular reflection** *Reflection* of light or a radar beam, as from a mirror or a plane faceted surface such as an angular boulder. See BACKSCATTER; compare DIFFUSE REFLECTION.

**spelean** Pertaining to caves.

**speleothem** See DRIPSTONE.

**Spermatophyta (seed plants)** Depending on the classification, a phylum or in most modern systems a superphylum of the plant kingdom comprising the *angiosperms* and *gymnosperms*, the gymnosperm superphylum comprising five groups of plants each of which is classed as a phylum. It may also be ranked as a subphylum (Spermatophytina) of the *Tracheophyta* (vascular plants).

**spessartine (spessartite)** Member of the *garnet* group of *minerals*, that has the formula  $Mn_3 Al_2 (SiO_4)_3$ ; sp. gr. 4.18; *hardness* 6.5–7.5; *cubic*; dark red to orange-yellow or brown; greasy to vitreous *lustre*; crystals

commonly *\*dodecahedra*; widely distributed in *\*metamorphic* and *\*igneous* rocks, and in beach and river *\*sands*. It is named after the Spessart Mountains, Bavaria, Germany.

**spessartite** 1. A type of *\*lamprophyre*, characterized by *\*essential \*hornblende* and *\*plagioclase feldspar*. *\*Aphyric* varieties are called '*\*malachites*'. 2. See SPESSARTINE.

**sp. gr.** See SPECIFIC GRAVITY.

**sphaericone** See INVOLUTE.

**sphalerite (black jack, zinc blende)** Mineral, ZnS; sp. gr. 3.9–4.1; *\*hardness* 3.5–4.0; *\*cubic*; colour variable, but commonly yellow, brown, or black, and crystals can be transparent to translucent; brownish red to bright yellow or white *\*streak*; resinous to near-metallic *\*lustre*; crystals *\*tetrahedral* or *\*dodecahedral*, with curved faces, but also granular, *\*fibrous*, or *\*botryoidal*; *\*conchoidal fracture*; *\*cleavage* perfect {011}; the most common *\*ore mineral* for zinc metal, Zn–Pb is common in strata-bound veins and massive sulphide deposits, frequently associated with *\*galena* in *\*hydrothermal* veins, and in *\*limestones* where it occurs by *\*replacement*, commonly with *\*pyrite*, *\*pyrrhotite*, and *\*magnetite*; dissolves in concentrated nitric acid with the separation of sulphur.

**sphene (titanite)** A *\*nesosilicate*  $\text{CaTiSiO}_4$  (O,OH,F); sp. gr. 3.45–3.55; *\*hardness* 5; *\*crystals* wedge-shaped, occasionally *\*massive*; brown, grey, reddish brown, yellow, or black; *\*monoclinic*; *\*adamantine* to *\*resinous \*lustre*; occurs as a primary *\*accessory mineral* in *\*calc-alkaline \*igneous* rocks and alkaline igneous rocks in which it may occur as a major constituent along with *\*apatite*, *\*nepheline*, and *\*aegirine*. It is common in *\*contact* metamorphosed *\*limestones*, particularly *\*skarns*.

**sphenoid** A wedge-shaped, four-faced, *\*closed form*, normally occurring in the *\*tetragonal* or *\*orthorhombic \*crystal systems*, but with a special form (the 'monoclinic sphenoid', or 'dihedron') occurring in the *\*monoclinic* system. The terminology for sphenoidal forms is complex and inconsistent, although in general the *\*crystallographic axes* emerge at the centre of the edges where pairs of triangular faces meet.

**Sphenopsida (horsetails)** A subphylum of the *\*Pteridophyta* that first appeared in the *\*Devonian* and reached the peak of its abundance and diversity during the *\*Carboniferous*, forming a major component of the coal-swamp vegetation. Sphenopsids are characterized by jointed stems with whorls of leaves and branches borne at the joints (or nodes). The internodal part of the stem is vertically ridged and *\*spores* are produced in rings of sporangia arranged in cones, usually at the tips of the fertile shoots. The only living genus, *Equisetum*, is a comparatively small plant (different species ranging between 4 or 5 cm and 12 m), but one of the best-known fossil genera, *Calamites*, included tree-like forms that grew up to 30 m in height. Another common fossil sphenopsid, *Sphenophyllum*, was a slender plant with a ribbed stem only 1–7 mm in diameter but up to several metres in length, that probably scrambled over other vegetation. *See also* [ARCHAEOCALAMITES RADIATUS](#); [CALAMITES CISTIIFORMES](#); [EQUISITITES HEMINGWAYI](#).

**sphericity** An expression of how closely the shape of a *\*grain* resembles the shape of a sphere. Sphericity can be determined by examining the relation between the long (*L*), intermediate (*I*), and short (*S*) axes of the particle, the maximum projection sphericity,  $\Psi$ , being given by the expression  $\Psi = \sqrt[3]{(S^2/LI)}$ . For a perfect sphere,  $\Psi = 1$ . Values less than one relate to increasingly less spherical shapes. *See also* [PARTICLE SHAPE](#); [ROUNDNESS INDEX](#).

**spheroid** The surface produced when an ellipse is rotated about one of its principal axes.

**spheroidal oscillation** *See* [FREE OSCILLATION](#).

**spheroidal weathering (onion weathering)** The development of concentric shells of normally chemically weathered material in the outer zone of a joint-bounded mass of rock. The spheroidal or onion-skin appearance results from enhanced *\*weathering* at *\*joint* intersections together with the expansion resulting from chemical change.

**spherule** A small spherical particle. *\*Glass* spherules resulting from impact-induced melting or volcanic *\*fire-fountaining* are common in the lunar *\*regolith*. A typical diameter is 100  $\mu\text{m}$ . Nickel–iron spherules,

usually less than 30  $\mu\text{m}$  in diameter, derived from the impacting *\*meteorite*, are common in impact glasses.

**spherulite** (*adj.* **spherulitic**) A spherical to ellipsoidal aggregate of radiating, fibrous crystals, usually *\*quartz* and *\*alkali feldspar*, found in glassy or *\*felsitic*, *\*aphanitic* *\*groundmasses* of *\*igneous* rocks. Spherulites range in diameter from less than 1 mm to about 1 m and are formed by the devitrification of quenched glassy igneous rocks which are usually silicic in composition.

**spicular chert** A very fine-grained siliceous *\*sedimentary rock*, the *\*silica* originating from the accumulation of sponge (*\*Porifera*) *\*spicules*. Sponge spicules can be an important source of *\*biogenic* silica in shelf-sea environments. See **CHERT**.

**spicule** A small needle or spine.

**spiculite** A *\*sedimentary rock* or *\*sediment* composed largely of sponge (*\*Porifera*) *\*spicules*.

**spider diagram** A useful plot, used in *\*igneous* *\*petrology* to show variations between two rocks or rock types for a wide range of elements. Usually, one rock is a standard type, e.g. a *\*mid-oceanic-ridge basalt* or a *\*carbonaceous chondrite* meteorite. Data for the test rock are 'normalized' to the standard by dividing the abundance for each element by the abundance in the standard and plotting the quotient, usually in order of atomic number. Elements which have the same abundance in the test rock as in the standard yield a quotient of 1; if the two compositions were identical the plot would be a straight line through 1 on the y axis. Elements more abundant in the test rock yield numbers greater than 1; those less abundant yield numbers less than 1. By connecting the points plotted by a single line, a spidery diagram of peaks and troughs is obtained which illustrates immediately systematic differences in composition, e.g. relative enrichment in *\*incompatible elements*.


**spiders** See **CHELICERATA**.

**spike** A solution (liquid or gaseous) containing a known concentration of a particular element whose isotopic concentration has been changed by the enrichment of one of its naturally occurring *\*isotopes*. The spike is used in *\*isotope dilution* analysis, being mixed in known proportions with the

sample solution, prior to isotopic determination by means of a mass spectrometer (see [MASS SPECTROMETRY](#)).

**spilite** A low-grade [\\*metamorphic rock](#) composed of [\\*albite](#), [\\*chlorite](#), [\\*actinolite](#), [\\*sphene](#), and [\\*calcite](#), with or without [\\*epidote](#), [\\*prehnite](#), and laumontite, and formed by sea-floor [\\*metasomatism](#) of [\\*mid-oceanic-ridge basalts](#). Sea water circulating through the [\\*oceanic crust](#) is heated by the cooling [\\*basalt \\*dykes](#) and [\\*lavas](#) and reacts with them, introducing sodium and water into the rock system and converting the basalt [\\*mineral assemblage](#) into a typical spilite assemblage.

**spilling breaker (surf wave)** Oversteepened wave in which the unstable top of the wave spills down the front of the wave-form as it advances into shallower water. Consequently it gradually diminishes in height until it moves up the beach as [\\*swash](#).



**Spilling breaker**

**spillway** General term for a [\\*glacial drainage channel](#) cut by water during glaciation, and normally including three varieties: (a) channels cut by water escaping from a glacially impounded lake (see [OVERFLOW CHANNEL](#)); (b) channels cut by meltwater released from a decaying [\\*glacier](#) (see [MELTWATER CHANNEL](#)); and (c) channels cut by a stream deflected by an advancing glacier. Impressive examples were developed in central Europe (the Urstromtäler of northern Germany) when the Scandinavian [\\*ice sheet](#) diverted streams flowing north from the southern highlands.

**spinal column** See [VERTEBRA](#).

**spindle bomb** See [VOLCANIC BOMB](#).


**spinel** Important group of non-silicate mineral *\*oxides*, including the subgroups spinel series, *\*magnetite* series, and *\*chromite* series. Members of the spinel series have the general formula  $XAl_2O_4$ , where X = Mg in spinel, X =  $Fe^{2+}$  in hercynite, X = Zn in gahnite, and X = Mn in galaxite. Complete *\*solid solution* exists between all four *\*end-members* and also magnesiochromite ( $MgCr_2O_4$ ); spinels with appreciable  $Fe^{2+}$  (Mg:Fe = 1:3) are called pleonaste and those with  $Fe^{2+}$  and  $Cr^{3+}$  are called mitchellite; sp. gr. 3.5–4.1; *\*hardness* 7.5–8.0; spinel is dark green and hercynite is dark bluish-green; all occur as small octohedra. Unlike chromite, which occurs in ultramafic (see **ULTRABASIC**) *\*igneous* intrusions, spinel may occur in metamorphic *\*schists* and *\*gneisses* with *\*sillimanite*, *\*garnet*, and *\*cordierite*. In *\*contact* metamorphosed impure *\*limestones* spinel occurs with *\*chondrodite*, *\*olivine*, and *\*orthopyroxene*, and in *\*emery* deposits it occurs with *\*corundum*. Spinel may develop in residual aluminous *\*xenoliths* enclosed in *\*basic* magmatic rocks. Spinel is found in *\*alluvial* deposits and gem quality spinel also occurs. Hercynite occurs in metamorphosed *\*laterites*; gahnite in *\*granite \*pegmatites*; and galaxite in manganese vein deposits.

**spinifex texture** An array of criss-crossing sheafs of subparallel, blade- or plate-like, skeletal, magnesium-rich *\*olivine* or aluminous *\*pyroxene*, between which is found a finer-grained aggregate of devitrified *\*glass*, skeletal pyroxene, and skeletal *\*chromite*. The *\*texture* is usually found as the product of extreme *\*undercooling* of magnesium-rich *\*komatiite \*lava*.

**spiny sharks** See **ACANTHODII**.

**spire** In the shell of a gastropod (*\*Gastropoda*), all the *\*whorls* other than the body whorl.

**Spiriferida (spiriferids)** (class *\*Articulata*) Order of *\*Brachiopoda*, whose members have spiral brachidia, *\*punctate* or *\*impunctate* biconvex shells, and a large body cavity. They first appeared in the Middle *\*Ordovician*, and are last known from the Lower *\*Jurassic*.



## Spiriferida

**spiriferids** See SPIRIFERIDA.

**spissatus** From the Latin *spissatus* meaning 'thickened', a species of \*cirrus cloud which has sufficient thickness to appear grey even when the cloud is between the Sun and the observer. See also CLOUD CLASSIFICATION.

**spit** Elongated accumulation of sand or gravel projecting from the shore into a water body. \*Longshore drift of material is usually responsible for the development of a spit.

**splanchnocranium** See CRANIUM.

**splay fault** One of a series of branching \*synthetic faults near the termination of a major fault which spread the displacement over a large area.

**splendent (specular)** Applied to the \*lustre of a \*mineral if it reflects light intensely to give a bright, shining surface. The reflectivity may be related to the high \*refractive index of the mineral (e.g. in \*gemstones).

**SP method** See SPONTANEOUS-POTENTIAL METHOD.

**spodic horizon** Subsurface \*soil horizon in which organic matter together with aluminium and often iron compounds have accumulated amorphously. It is a \*diagnostic horizon in the \*USDA \*soil taxonomy.

**spodosols** Order of soils in which subsurface \*soil horizons contain amorphous materials comprising organic matter and compounds of

aluminium and often iron that have accumulated illuvially. Such soils form in acid material, mainly coarse in texture, in humid, cool to temperate climates.

**spodumene** Unusual, lithium-bearing *\*pyroxene*,  $\text{LiAlSi}_2\text{O}_6$ ; sp. gr. 3.0–3.2; *\*hardness* 6.5–7.0; *\*monoclinic*; usually greyish-white, often with a greenish or yellowish-green tinge, occasionally violet, transparent to translucent; vitreous *\*lustre*; crystals *\*prismatic*, often striated, etched, and corroded, also occurs *\*massive* and *\*columnar*; *\*cleavage* perfect prismatic {110}, parting {100}; typically occurs in lithium-rich *\*granites* and *\*pegmatites*, associated with *\*lepidolite*, *\*tourmaline*, and *\*beryl*. The green variety (hiddenite) and the lilac variety (kunzite) are used as *\*gemstones*, otherwise it is an *\*ore mineral* of lithium.

**spondylium** Curved platform for muscle attachment in the *\*shell* beak region of some brachiopods (*\*Brachiopoda*).

**sponges** See DEMOSPONGEA; PORIFERA.

**Spongiaria** See PORIFERA.

**spontaneous potential** See SELF-POTENTIAL SONDE.

**spontaneous-potential method (self-potential (SP) method)** A method which measures the naturally occurring potential differences between two *\*non-polarizable electrodes*. It is often used in exploration for massive *\*sulphide* and *\*graphite* *\*orebodies*.

**sporangium** See SPORE.

**spore** A propagative plant body consisting of a gametophyte enclosed in a non-cellular coat. Spores are enclosed within a capsule (sporangium) and are produced in groups of four (tetrads) when the parent cell divides meiotically. In more primitive plants the spores are identical (isospore) and the condition is called 'homospory'. In more advanced, vascular plants spores of two sizes are produced and the condition is called 'heterospory'. Small, male microspores are contained within a microsporangium; larger, female megaspores within megasporangia. Where spores occur in tetrads the contact surfaces produce a 'trilete' mark on each of the four spores, marking the point for the germination of the prothallus. Less commonly, two contact surfaces are produced, resulting in a 'monolete' marking.

Spores which were probably produced singly are 'alete', with no obvious marking.

**SporeSat** A cooperative 3-unit *\*CubeSat* mission between *\*NASA* and Ames Research Center, California, that supplies data to help studies into gravity detection by plant cells. The *\*nanosatellite* was launched on 18 April 2014, from Cape Canaveral, into a near-circular orbit at an altitude of about 400 km.

**sporinite** See COAL MACERAL.

**SPORT** See SCINTILLATION PREDICTION OBSERVATIONS RESEARCH TASK.

**SPOT** See SYSTÈME PROBATOIRE D'OBSERVATION DE LA TERRE.

**spotting** Dark-coloured, rounded areas, up to 2 mm in diameter and sometimes larger, found on the surface of *\*slaty cleavage* in low- to medium-grade contact *\*metamorphic rocks* (see CONTACT METAMORPHISM) of pelitic composition. Many spots are *\*graphite*-rich, having formed by metamorphic aggregation from organic material originally disseminated through the starting sedimentary *\*shale*. Rocks may also have a spotty appearance when *\*andalusite* is beginning to appear in contact-metamorphosed *\*pelites*.

**spread** A pattern of *\*geophone* *\*groups* used simultaneously to record data from a single *\*shot*. Examples of spreads include in-line, offset, interlocking, L-spread, reversed, split-spread, and T-spread. See ARRAY.

**spreading rate** The rate, usually in tens of millimetres a year, at which two adjacent lithospheric *\*plates* are separating. The spreading rate varies along a *\*constructive margin* and is at a maximum of 90° from the *\*pole of rotation*. Some authors use 'spreading rate' when 'half spreading rate' (i.e. the rate of movement of a plate from the relevant *\*ridge*) would be more accurate.

**spreiten** (*sing.* **spreite**) From the German *spreiten*, meaning 'to spread out' or 'to extend', sedimentary *\*laminae* that result from the behaviour of an animal during feeding, excavation, or locomotion. They may be U-shaped, sinuous, blade-like, or spiralled, and they are always repeated over a small area. They reflect the intensive working of *\*sediment* for food. Well-defined spreiten are associated with *\*Diplocraterion*, *\*Rhizocorallium*, and

*Daedalus*. In the latter they are arranged spirally around a single trunk. See FUGICHNIA.

***Spriggina*** See EDIACARAN FOSSILS.

**spring** A flow of water above ground level that occurs where the **\*water-table** intercepts the ground surface. Where the flow from a spring is not distinct (i.e. it does not give rise to obvious trickles) but tends to be somewhat dispersed, the flow is more correctly termed a 'seep'. The reappearance of surface water that had been diverted underground in a **\*karst** region is a type of spring known as a 'resurgence'. A major variety is the 'Vauclisian spring', named after the Fontaine de Vaucluse, southern France, and descriptive of the upward emergence of an underground river from a flooded **\*solution channel**.

**spring balance** A weighing scale, consisting of a pan suspended below a vertical spiral spring, used to measure the weight and often the density of rocks and minerals. When a specimen is placed on the pan the spring stretches along a calibrated scale and a pointer indicates the weight. The Jolly balance (see DENSITY DETERMINATION) is a spring balance.

**Springerian** See CHESTERIAN.

**spring sapping** A set of geomorphological **\*processes** that erode a hillslope around the site where a **\*spring** emerges. The processes may include the collapse of saturated material, surface stream **\*erosion**, and **\*chemical weathering**. It occurs towards the bases of **\*chalk** escarpments in southern England, where its effect may have been enhanced by frost activity under former **\*periglacial** conditions.

**spring tide** **\*Tide** of greater range than the mean range; the water level rises and falls to the greatest extent from the mean tide level. Spring tides occur about every two weeks, when the Moon is full or new. Tides are at their maximum when the Moon and the Sun are in the same plane as the Earth. Compare NEAP TIDE.

**sprite halo** A **\*transient luminous event** consisting of a luminous disc seen above a thunderstorm, lasting about 0.001 seconds, that propagates downward from a height of about 85 km to about 70 km, often preceding a **\*red sprite**.

**s-process** See SLOW-NEUTRON PROCESS.

**SPS** A satellite positioning system that uses interaction-geostationary and low, short-period satellite passes to determine the location of stations on the Earth's surface. See GLOBAL POSITIONING SYSTEM.

**spur** A ridge that descends towards a valley floor from the higher ground above. It may be due to an outcrop of resistant rock, or it may develop on the concave side of a winding stream as a result of incision.

**squall** Short-lived storm with strong winds, which increase by up to 50% to at least 16 knots (30 km/hr) for two minutes before dying away again. It may include thunder and heavy precipitation. See also LINE SQUALL; SQUALL LINE.

**squall line** A series of very vigorous \*cumulonimbus clouds that merge to form a continuous line, up to 1000 km long, that advances at right angles to the line, bringing severe \*squalls associated with thunderstorms and sometimes triggering \*tornadoes. Squall lines usually develop along a \*cold front from which they become detached, advancing ahead of the front.

**square array** An \*electrode configuration in which the \*current and \*potential electrodes are positioned at the four corners of a square of side  $a$ . Its \*geometric factor is  $K_g = 3.41\pi a$  metres.

**squeeze-up** The extrusion of a small volume of viscous \*lava from a crack or opening on the solidified surface of a lava flow, in response to the pressure of fluid lava within the flow interior. Squeeze-ups are generally bulbous or linear in form, range from a few centimetres to several metres in height, and may have vertical grooves down their length.

**squeezing ground** Weak ground, such as \*clay, which has deformed under surrounding loads and has been squeezed into an \*excavation as a result of overstraining.

**Sq variation** See DIURNAL VARIATION.

**SSI** See SOLID-STATE IMAGING CAMERA.

**SSOT** See SISTEMA SATELITAL PARA LA OBSERVACIÓN DE LA TERRA.

**SSS** See STANDARD STRATIGRAPHIC SCALE.

**stability** **1.** Atmospheric condition in which air that is forced to rise tends to return to its pre-existing level in the absence of the uplifting force. If the **\*adiabatic \*lapse** rate of uplifted air is greater than the **\*environmental lapse rate**, then the vertically displaced air will become colder than the surrounding air and as its density increases it will tend to sink back. *See also* **INSTABILITY**. **2.** In engineering, the resistance of a structure to collapse or sliding, dependent upon the **\*shearing strength** of the material. **3.** In geochemistry, the state of equilibrium towards which a system will move from any other state under the same conditions. **4.** In thermodynamics, the condition when a slight disturbance of temperature, pressure, or composition does not result in the appearance of a new **\*phase**.

**stability field** Range of temperature and pressure within which a particular mineral or mineral assemblage is stable.

**stable isotope** Any naturally occurring, non-radiogenic **\*isotope** of an element. Many elements have several stable isotopes.

**stable-isotope studies** Study of non-radiogenic isotopic ratios of selected elements, e.g.  $^{18}\text{O}:^{16}\text{O}$ ,  $^{32}\text{S}:^{34}\text{S}$ , which are fractionated in different proportions during different geologic processes (*see* **ISOTOPE FRACTIONATION**). Thus natural waters may be 'fingerprinted' by reference to their D:H and  $^{18}\text{O}:^{16}\text{O}$  ratios as being of **\*meteoric**, magmatic, or metamorphic origin; sulphur in **\*sulphide \*ores** may be characterized as **\*sedimentary** or **\*igneous** by reference to its  $^{32}\text{S}:^{34}\text{S}$  ratio. *See also* **OXYGEN-ISOTOPE ANALYSIS; OXYGEN-ISOTOPE RATIO**.

**stack** **1.** Pillar or block of rock, with near-vertical sides, standing adjacent to a present or former sea cliff. Typically it has been isolated from the main cliff by wave erosion concentrated along steeply inclined **\*joints** or **\*faults**. **2.** The product of **\*stacking**. A 'brute' or 'final' stack is the end product of the standard processing of seismic-**\*reflection** data; the data can be processed further, e.g. by migration programs to produce a 'migration' stack.

**stacking** The summing of traces from a variety of **\*seismic records** to increase the signal-to-**\*noise** ratio and enhance coherent signals into a composite record (a **\*stack**). *See also* **COMMON-DEPTH-POINT STACK; VERTICAL STACKING**.

**stacking fault** An abnormality in the arrangement of the rows of atoms affecting the structure of a *\*crystal* (e.g. *\*crystal twinning*), caused by changes in the physical and chemical conditions of its immediate surroundings while it was growing.

**stacking velocity** In seismic investigations, the velocity determined from normal *\*moveout* measurements using *\*common-depth-point* gathers prior to *\*stacking*.

**stade (stadial in continental-European usage)** A term that is difficult to define with precision, but which refers to a single period of increased cold or advancing ice, which forms a subdivision of a cold *\*stage* within the overall division of a glacial period into periods of cold interspersed with warm, or warmer, periods.

**stadial** See *STADE*.

**staff gauge** A graduated pole or board placed in or beside a water course, from which it is possible to measure directly the height of the water surface relative to a known datum elevation.

**stage** 1. The elevation of the water surface of a river with reference to a fixed datum level. Hence 'rising' and 'falling' stages. 2. The major subdivision of a *\*series*. A stage is the fourth order unit in chronostratigraphy, the equivalent of *\*age* in terms of geologic time units. It refers to the body of rock accumulated during one age unit. When used formally the initial letter of the term is often capitalized, e.g. *\*Frasnian Stage*. 3. In palaeoclimatology, a climatic, and partly geologic-climatic, term usually defined by a series of *\*sediments* or a sequence of *\*fossil assemblages* and named at a type locality. For example, the *\*Hoxnian* (a temperate stage) is named for organic *\*interglacial* deposits at Hoxne, Suffolk, England. 4. The degree of development of a land-form or landscape over time, and which traditionally has been described by the terms 'youthful', 'mature', and 'old age' (see *DAVISIAN CYCLE*). The recognition of such stages implies an orderly evolution and this is now seen as unlikely for many parts of the Earth's land surface. 5. The part of a microscope on which the specimen to be examined is placed. Normally it is flat and may be fixed, as in biological or metallurgical microscopes, or rotating with a 360° calibrated scale as in geologic microscopes. Transmitted-light microscopes have a hole in the centre of the stage through

which light passes up to the observer from below. **\*Reflected-light** microscopes have an incident light system, whereby light is directed on to the stage from above and is reflected from the specimen to the observer.

**stage hydrograph** See HYDROGRAPH.

**stagnosols** Soils related to **\*luvisols** that occur mainly in humid temperate regions of western and central Europe. They have reducing conditions within 50 cm of the surface and strong mottling in 50% of the reducing layer that develops under temporary saturation. The topsoil may be bleached and **\*concretions** may occur in the subsoil. They are a soil group in the **\*World Reference Base for Soil Resources**.

**stains and staining techniques** Various chemical staining techniques are used to identify **\*minerals**. The procedure followed is to etch the specimen, and then to expose it to a range of organic and/or inorganic compounds which form distinctive coloured complexes with certain minerals. **\*Feldspars** can be identified by etching with hydrofluoric acid and then treating with sodium cobaltinitrate, barium chloride, and potassium acid rhodizonate. **\*Plagioclases** stain red and K-feldspars are stained yellow after this treatment. For **\*carbonates**, a range of stains, including alizarin red S, Feigl's solution, potassium ferricyanide alizarine cyanic green, and titan yellow are used. A number of staining combinations allow the differentiation of **\*calcite**, high-Mg calcite, **\*dolomite**, **\*anhydrite**, and **\*gypsum**.

**stalactite** Elongated body of **\*dripstone** descending from the roof of a cave in a **\*karst** environment. It is produced by **\*calcite** **\*precipitation** as excess carbon dioxide diffuses from water droplets entering a cave environment.

**stalagmite** Pinnacle of **\*dripstone** rising from the floor of a cave in a **\*karst** environment. It is produced by the **\*precipitation** of **\*calcite** as excess carbon dioxide diffuses when water droplets strike the floor.

**Stampian** See RUPELIAN.

**standard deviation** A measure of the normal variation within a set of data. In any given measurement, two-thirds of the samples fall within one standard deviation on either side of the mean, 95% between two standard deviations, and so on; the proportion falls off sharply because of the bell-

curve effect. The standard deviation is calculated as the root-mean-square deviation.

**Standard Global Chronostratigraphic Scale (SGCS)** See STANDARD STRATIGRAPHIC SCALE.

**standard mean open water** See SMOW.

**Standard Stratigraphic Scale (Standard Stratigraphical Scale, SSS; Standard Global Chronostratigraphic Scale, SGCS)** Globally standardized \*stratigraphic scale whose \*chronostratigraphic units will ultimately all be delimited by \*boundary stratotypes.

**standing wave** Type of wave in which the surface oscillates vertically between fixed points called 'nodes', without any forward progression. The crest at one moment becomes the trough at the next and so on. The points of maximum vertical rise and fall are called 'antinodes'. At the nodes particles show no vertical motion but exhibit the maximum horizontal motion. Standing waves may be caused by the meeting of two similar wave groups that are travelling in opposing directions. See SEICHE.


**stand of the tide** Period at high or low water during a tidal cycle when there is little or no change in the height of the \*tide. The water level is almost stationary and the tidal currents fall away to zero velocity before reversing.

**stannite** Comparatively rare mineral,  $\text{Cu}_2\text{FeSnS}_4$ ; sp. gr. 4.3–4.5; \*hardness 3–4; \*tetragonal; steel-grey with an olive-green tinge on fresh surfaces, which become yellowish on exposure; black \*streak; \*metallic \*lustre; crystals rare, \*cubes or \*tetrahedra, but normally the grains are irregular or \*massive; occurs in \*hydrothermal tin-\*ore deposits and in association with \*cassiterite, \*chalcopyrite, and \*wolframite, also in stanniferous \*sphalerite–\*galena ores, and associated with sphalerite, \*pyrrhotite, and galena.

**stapes** In \*Mammalia, the inner auditory ossicle of the ear, stirrup-shaped because it is pierced by an artery. It is derived from the hyomandibular bone in fish, which connects the cranium and the upper jaw.

**star dune** A complex \*aeolian \*dune form characterized by a series of slip faces radiating about a central point, producing a rough star shape. Such

dunes are the product of highly variable wind directions and thus have a highly variable palaeocurrent pattern preserved in the dune cross-bedding (see [CROSS-LAMINATION](#)).



**Star dune**

**Stardust** A [\\*NASA](#) mission to return coma samples from comet [\\*Wild 2](#). Launched in 1999, it reached the comet in January 2004.



<https://www.jpl.nasa.gov/missions/stardust/>

- Stardust.

**Starlette and Stella** Two virtually identical passive satellites of the French Centre National d'Études Spatial (CNES) to assist geodetic (see [GEODESY](#)) and geophysical studies. Each satellite is covered with sixty laser retroreflectors. The satellites were launched on 26 September 1993, from Kourou, French Guiana, Starlette into a near-circular orbit at an altitude of 812 km, and Stella into a [\\*polar orbit](#) at a mean altitude of 800 km.

**star pair (binary star, double star)** Binary stars are among the commonest stellar systems in our galaxy, accounting for about 50% of all stars. They consist of two stars in orbit around their common centre of gravity. Those which can be resolved by telescope are referred to as visual binaries. Spectroscopic binaries comprise star pairs too close to be resolved visually,

but which exhibit **\*Doppler shifts** in spectral lines because of varying velocities of the two stars in the line of sight. Other stars reveal the presence of an invisible companion by changes in their **\*proper motion**. Double stars appear to be visually close to each other, but may be at vastly different distances from the observer.

**star phylogeny** In a **\*phylogenetic tree**, the occurrence of a multifurcation with many short branches connected at the **\*internal node**. Such topologies are often inferred to represent a recent population expansion event from a common ancestor (the founder lineage). This is often seen in populations which have undergone a **\*founder effect**.

**star twinkling** See **ATMOSPHERIC SHIMMER**.

**stasigenesis** Situation in which an evolutionary lineage persists through time without splitting or otherwise changing. So-called '**\*living fossils**' are examples of stasigenesis.

**stasis** Period of little or no evolutionary change; the 'equilibrium' that alternates with 'punctuations' in the theory of **\*punctuated equilibrium**.

**Statherian** The final period of the **\*Palaeoproterozoic \*era** that began 1800 Ma ago and ended 1600 Ma ago, both its beginning and end being defined by fixed dates. The Statherian followed the **\*Orosirian** period and was followed by the **\*Calymmian** period, marking the start of the **\*Mesoproterozoic** era.

**static correction (statics)** A correction applied to geophysical data, especially seismic data, to compensate for the effect of irregular topography, differences in the elevation of **\*shots** and **\*geophones** relative to a datum, low-velocity surface layers (weathering correction), and the horizontal geometry of shots and receivers (geophones or **\*hydrophones**), or any correction which applies to the geometry of the source and receiver(s). A static correction provides some form of direct-current shift (e.g. in seismic-**\*reflection** surveys), usually a time element added to or subtracted from the travel times, in contrast to a dynamic correction which involves an operation on the data. See **ELEVATION CORRECTION**; **MOVEOUT**.

**statics** See **STATIC CORRECTION**.

**stationary front** Condition in which the frontal boundary between cold and warm air is stationary, or moves slowly and erratically, with air on either side of the front moving approximately parallel to the front.

**station frequency** See SAMPLING FREQUENCY.

**station interval** See SAMPLING INTERVAL.

**stauroilite** A member of the *\*nesosilicates* and an important metamorphic *\*index mineral* with the approximate composition  $(\text{Fe}^{2+}, \text{Mg})_2(\text{Al}, \text{Fe}^{3+})_9\text{O}_6[\text{Si}_4\text{O}_{16}](\text{O}, \text{OH})_2$ ; sp. gr. 3.74–3.85; *\*hardness* 7.5; *\*monoclinic*; *\*crystals* *\*prismatic*; shades of brown; occurs in regionally metamorphosed *\*schists* and *\*gneisses*, such as iron-rich *\*pelites*, with a high  $\text{Fe}^{3+}/\text{Fe}^{2+}$  ratio at moderate grades of *\*metamorphism* and in association with *\*garnet* (*\*almandine*) and *\*kyanite*; it may develop from *\*chloritoid* as the metamorphic grade increases.

**steady flow** The condition in which flow velocities do not vary with time. This is applicable to both *\*groundwater* and channel flows. In respect of flow to a pumped well, it is sometimes called ‘equilibrium’ flow. Non-steady (unsteady or transient) flow changes its velocity and/or direction with time.

**steam fog** See ARCTIC SEA SMOKE.

**steatite** See TALC.

**S-tectonite** See SHAPE FABRIC.

**Stefan–Boltzmann law** The law stating that energy radiated from a *\*black body* is proportional to the fourth power of its *\*absolute temperature*.

**Stegosauridae** Suborder of quadrupedal, *\*ornithischian* *\*dinosaurs*, mainly *\*Jurassic* in age, characterized by a double row of plates and spines along the back and tail.

**steinkern** See FOSSILIZATION.

**Steinmann trinity** *\*Spilites*, *\*serpentine* rocks, and radiolarian *\*cherts*, which, as G. Steinmann observed in 1905, often occur together in mountains such as the Alps, comprising rocks formed as deep-sea sediments.

**Stella** See STARLETTE AND STELLA.

**stem group** In *\*cladistic analysis*, those taxa descended from the point where an ancestral taxon split into two *\*sister groups* to the point at which a further split gave rise to an extant *\*crown group*.

**stem reptiles** See COTYLOSAURIA; CAPTORHINOMORPHA.

**Stenian** The final period of the *\*Mesoproterozoic* era, which began 1200 Ma ago and ended 1000 Ma ago, both its beginning and end being defined by fixed dates. The Stenian followed the *\*Ectasian* period and was followed by the *\*Tonian* period, marking the start of the *\*Neoproterozoic* era.

**Steno, Nicolaus** (**Nicolaus Niels Stensen Stenonis**) (1638–87) A Danish physician who moved to Florence in 1665. He opposed the prevailing idea that *\*fossils* grew within the Earth, proposing instead that they were organic relics of an earlier period. Steno also distinguished between fossils and inorganic remains such as crystals. He had some conception of stratigraphy, describing strata in Tuscany as being formed sequentially.

**stenothermal** Unable to tolerate a wide range of temperature.

**stenotopic** Able to tolerate only a narrow range of several factors.

**Stensen, Niels** See STENO, NICOLAUS.

**step faulting** The faulting process in which separate *\*fault blocks* are downthrown systematically in one direction, forming a stepped sequence.

**Stephanian** The uppermost *\*stage* in the *\*Silesian* (Upper *\*Carboniferous*) of Europe, underlain by the *\*Westphalian*, followed by the Early *\*Permian*, dated at 306.5–299 Ma ago and roughly contemporaneous with the uppermost *\*Moscovian*, *\*Kasimovian*, and *\*Gzhelian* stages. Originally, the lower part of the Stephanian was known as Westphalian E.

**Stephano** (**Uranus XX**) A lesser satellite of *\*Uranus* with a radius of 16 km and a visual *\*albedo* of 0.07.

**stepout** See MOVEOUT.

**steptoe** A type of *\*kipuka* named after Steptoe Butte (itself named after Lt. Col. Edward Steptoe, 1815–65), in Washington State, USA. It is an isolated hill of older rock that protrudes above an extensive *\*lava* field.

**stereogram** The two-dimensional plot of a *\*stereographic projection*, in which points on the surface of a sphere are represented on a *\*plane of projection*, as points on the Earth's surface might be projected on to a plane representing a slice through the equator by joining these points to either the N. or S. poles. If the plane of projection (equatorial plane) is constructed with *\*great circles* and *\*small circles* drawn at 2° intervals and with an overall diameter of about 20 cm, it is called a 'Wulff stereographic net'. If it is graduated at 10° intervals and includes small circles concentric to the *\*primitive circle* and radii representing vertical great circles, it is called a 'Federov net'. Plotting is done on a sheet of tracing paper laid over the printed net and pivoted about a pin at its centre.

**stereographic net** See STEREOGRAM.

**stereographic projection** A two-dimensional graphic representation of a three-dimensional solid object, in which the angular relationships of lines and planes of the object are drawn in terms of their relationship to the *\*great circle* formed by the intersection of the equatorial plane with the surface of an imaginary sphere in which the object is contained. Stereographic projections are used widely in structural geology and *\*crystallography*.

**stereom (stereome)** In *\*Echinodermata*, a mesh, mainly of magnesian *\*calcite*, from which the skeleton is constructed. In *\*Scleractinia*, a secondary structure, composed of bundles of *\*aragonite* crystals arranged transversely, that thickens and strengthens the *\*epitheca*.

**stereonet** A two-dimensional, circular representation of a sphere in which the lines of longitude and latitude form a system of coordinates (a 'net'), on which projections of *\*great* and *\*small circles* occupy the equatorial plane of a reference sphere. Two types of stereonet are used in the analysis of structural data: an equal-angle net (Wulff net); and an *\*equal-area* net (Schmidt–Lambert net), which is the net preferred for the contouring of data and the evaluation of clusters of data suggesting preferred orientations.

**stereophotography** The taking of two pictures (stereo pairs), e.g. of a landscape or *\*fossil*, from slightly different angles so that when the two images are viewed through a *\*stereoscope* a three-dimensional image is produced, giving greater detail than an ordinary photograph.

**stereoptic vision** The perception of depth and three dimensions accompanying binocular vision resulting from differences in **\*parallax** producing different images on the retina of each eye.

**stereoscope** An optical device which allows a pair of overlapping, two-dimensional photographs to be examined with three-dimensional (**\*stereoptic**) vision, thus permitting more detailed interpretation.

**Steropodon** The earliest Australian mammal, a monotreme known only from opalized jaws and teeth found in **\*Cretaceous** deposits at Lightning Ridge, central Australia (the name is from the Greek *sterope*, meaning 'lightning'). The pattern of the molar teeth suggests that monotremes may be derived from Theria and not from a separate protomammalian stock.

**Stettin** An end **\*moraine** and a variety of **\*sediments** occurring in Poland and European Russia. Its stratigraphic position is uncertain but it is probably equivalent to the early **\*Weichselian** or **\*Rodebaek**.

**stibnite (antimonite, antimony glance)** One of the main **\*ore minerals** for antimony, with the formula  $Sb_2S_3$ ; sp. gr. 4.6; **\*hardness** 2.0–2.5; **\*orthorhombic**; normally lead-grey with a bluish tarnish; **\*metallic** **\*lustre**; crystals **\*prismatic**, columnar, **\*acicular**, and vertically striated; perfect lengthwise **\*cleavage** {010}, imperfect {100}, {110}; occurs mainly in hydrothermal deposits, at the low-temperature end, often forming sheet-like bodies, associated with **\*fluorite**, **\*quartz**, and **\*barite**, and with sulphides of lead, zinc, and other metals.

**stick-slip** A discontinuous, jerky pattern of movement along a **\*fault plane**, which is thought to be consistent with **\*earthquake** phenomena.

**Stigmaria** The **\*form-genus** for the underground axes of Lepidodendrales (**\*Carboniferous**).

**Stille, Wilhelm Hans** (1876–1976) A German geologist from the Universities of Göttingen and Berlin, Stille's main work was in the field of **\*orogeny**. He believed that mountain building occurred in phases and that the continents were formed from the accretion of mountain belts around ancient **\*cratons**.

**stillstand** A period of geologic time characterized by unchanging sea levels, i.e. a state neither of **\*regression** nor **\*transgression**.

**stilpnomelane** A **\*phyllosilicate** (sheet silicate) similar to **\*biotite** with the formula  $(K,Na,Ca)_{0-1.4} (Fe^{2+},Fe^{3+},Mg,Al,Mn)_{5.9-8.2} [Si_8O_{20}](OH)_4(OH,F)_{3.6-8.5}$ ; sp. gr. 2.59–2.96; **\*hardness** 3.0–4.0; properties similar to those of biotite; found in metamorphosed iron- and manganese-rich sedimentary deposits and in some glaucophane schists.

**stipe** A branch of a rhabdosome (graptolite colony, *see* **GRAPTOLITHINA**). A stipe or stipes originate(s) from an initial conical cup (**\*sacula**). The number of stipes can range from one to as many as 64. Their attitudes also vary. In the primitive condition the stipes hang downwards from the sicula and are said to be ‘pendent’. If they grow out horizontally from the sicula they are ‘horizontal’; if they grow upwards along the nema they are ‘scandent’; if they are straight and grow downwards they are ‘declined’; if they are curved and slope downwards they are ‘deflexed’; if they are straight and grow upwards and outwards they are ‘reclined’; and if they are curved and grow upwards and outwards they are ‘reflexed’.

**stishovite** A high-density form of crystalline silica (*see also* **QUARTZ**) with the same formula  $SiO_2$ , but formed at pressures greater than 10 GPa; discovered at some **\*meteorite** impact sites, but rarely in terrestrial rocks; sp.gr. 4.3; minerals possessing a similar atomic lattice may exist in the Earth’s upper **\*mantle**.

**stock** **\*Igneous** **\*intrusion**, approximately circular in plan, that has steep contacts with the **\*country rocks** and a surface area of 20 km<sup>2</sup> or less.

**stockwork** Mineral deposit formed of a network of small, irregular **\*veins** so closely spaced that it may be mined as a unit.

**Stokes’s law** A law describing the rate at which suspended particles settle, formulated in 1845 by the physicist Sir George Gabriel Stokes (1819–1903). The settling velocity ( $V$ ) in cm/s is calculated by  $V = CD^2$ , where  $C$  is a constant related to the density and viscosity of the fluid and the density of the suspension and  $D$  is the diameter of the particles (assumed to be spheres) in cm.

**stolon** In colonial invertebrates, the stalk-like structure by which individuals are attached to the substrate.

**stolotheca** One of the three types of graptolite (*\*Graptolithina*) *\*thecae*, which encloses the main *\*stolon* and the earliest parts of the daughter stolotheca (the authotheca, and *\*bitheca*).

**stomodeum** See ANTHOZOA.

**-stone** Suffix for many different kinds of lithified *\*sediments*, e.g. *\*siltstone*, *\*limestone*, *\*sandstone*, *\*grainstone*, *\*packstone*, and *\*ironstone*.

**stone canal** In *\*Echinodermata*, a canal with walls strengthened by calcareous matter, which connects the *\*madreporite* with the water-vascular system.

**stone circle** See PATTERNED GROUND.

**stone garland** See PATTERNED GROUND.

**stone net** See PATTERNED GROUND.

**stone polygon** See PATTERNED GROUND.

**stone steps** See PATTERNED GROUND.

**stone stripes** See PATTERNED GROUND.

**stony-iron meteorite (siderolite)** Relatively rare *\*meteorite* type with approximately equal quantities of nickel–iron and basic *\*silicates*, usually *\*pyroxene* and *\*olivine*.

**stony meteorite (siderite)** *\*Meteorite* type consisting mainly of rock-forming *\*silicates* (*\*olivine*, *\*pyroxene*, and *\*plagioclase*) with some nickel–iron; more than 90% of meteorites seen to fall are of this type. Stony meteorites are referred to as *\*chondrites* or *\*achondrites*, depending on the presence or absence of *\*chondrules*.

**stoop and room** See PILLAR AND STALL.

**stopping** 1. The method of emplacement of an *\*igneous \*intrusion* in which percolating *\*magma* detaches blocks of *\*country rock* which sink, allowing the magma to move upwards. 2. In underground mining, the breaking and removal of rock in an *\*orebody*.

**storage coefficient (storativity)** The volume of water given up per unit horizontal area of an *\*aquifer* and per unit drop of the *\*water-table* or

**\*potentiometric surface.** It is a dimensionless ratio and always less than unity. In unconfined aquifers it is equal to the specific yield (see **SPECIFIC YIELD (2)**), but in confined aquifers the storage coefficient depends on elastic compression of the aquifer, and is usually less than  $10^{-3}$ .

**storativity** See **STORAGE COEFFICIENT**.

**storm** Common term for gales, squalls, rainstorms, or thunderstorms. It is used specifically for conditions associated with the active areas of low-pressure systems. ‘Storm-force winds’ are, by definition, strong gales or winds, with speeds exceeding 30 m/s. See **APPENDIX C: THE BEAUFORT SCALE OF WIND STRENGTH**.

**storm beach** Accumulation of coarse beach sediments built above the high-water mark by storm action. Gravel, shell debris, and other coarse materials are thrown into ridge or bank structures by waves during heavy storms.

**storm bed** A bed of **\*sediment** deposited by a storm event. Storm beds are usually the product of shallow marine wave activity, and are often referred to as ‘event deposits’, that is, they are the product of a short-lived, high-energy, sedimentary environment.

**storm deposit** See **TEMPESTITE**.

**storm surge** Rise or piling-up of water during a storm, as a result of wind stresses acting on the surface of the sea and of atmospheric-pressure differences. If a storm surge occurs at the time of highest **\*spring tides**, flooding of coastal areas may result, as happened in Holland and East Anglia, England, in 1953.

**storm wave-base** The sea depth, typically 15–40 m, above which the sea bed is affected by waves that occur during storms. During storms, sand is eroded and deposited as **\*hummocky** cross-bedding. Compare **FAIRWEATHER WAVE-BASE**.

**stoss** Describes the side of a sloping surface that faces into the flow of ice, water, or wind.

**stoss and lee** Terms referring to the up-**\*glacier** and down-glacier slopes respectively of a rocky obstacle that has been glaciated. The stoss slope is

smoothly abraded, the lee slope roughly plucked. A landscape dominated by such features is said to have 'stoss-and-lee topography'.

**stoss-and-lee topography** See ROCHE MOUTONNÉE; STOSS AND LEE.


**Strahler climate classification** A system for describing climates, devised in 1969 by A. N. Strahler, in which world climates are related to the main *\*air masses* that produce them, as: (a) equatorial/tropical air masses, producing low-latitude climates; (b) tropical and polar air masses, producing mid-latitude climates; and (c) polar and arctic air masses, producing high-latitude climates. Subsets of these are based on variations in temperature and *\*precipitation* to give 14 regional types, plus upland (highland) climates which are regarded as a separate category. See also KÖPPEN CLIMATE CLASSIFICATION; THORNTHWAITE CLIMATE CLASSIFICATION.

**straight extinction (parallel extinction)** In optical *\*mineralogy*, the phenomenon which occurs when the *\*vibration direction* of the light ray is parallel to the *\*crystal face* or *\*cleavage* traces within the mineral. As the mineral *\*thin section* is rotated between *\*crossed polars*, the vibration direction is brought parallel to the plane of the *\*polarizer*. The light passes through the polarizer and is eliminated by the *\*analyser*. *\*Extinction* occurs four times in a 360° rotation of the mineral.

**strain** The dimensional change in the shape or volume of a body as a result of an applied *\*stress* or stresses. Strain is the ratio of the altered length, area, or volume to its original value, and may be *\*homogeneous* or *\*inhomogeneous*, and involve distortion, *\*dilation*, and rotation. See HOOKE'S LAW; POISSON'S RATIO; PURE SHEAR; SIMPLE SHEAR; SHEAR MODULUS.

**strain ellipse** A two-dimensional figure used to describe the magnitude and orientation of the maximum and minimum *\*principal strain axes*  $x$  and  $z$  when a reference circle of unit radius is deformed by *\*homogeneous strain*.

**strain ellipsoid** A three-dimensional version of the *\*strain ellipse* which describes the magnitude and orientation of the maximum, intermediate, and minimum *\*principal strain axes* when a reference sphere of unit radius is deformed by *\*homogeneous strain*.



**Strain ellipsoid**

**strain gauge** A device for measuring **strain**.

**strain marker** A natural object whose original geometry is known and from which the magnitude and orientation of the **principal strain axes** may be determined after deformation. Many objects have been used, e.g. **fossils** (**crinoids**, **belemnites**, **corals**, **ammonites**, **brachiopods**, and **trilobites**), rock **textures** (i.e. **ooids**), **conglomerate** pebbles and **clasts**, single crystals (e.g. **feldspar megacrysts**), spots in **contact aureoles**, volcanic **lapilli**, **spherulites**, and **xenoliths**.

**strain parallelepiped** The three-dimensional shape produced when a small cube of unit dimensions is deformed.

**strain rate** The rate of change of the size and shape of a body under an applied **stress**. The duration of the stress is very important in determining strain behaviour. Most geologic strain rates are relatively low, with stresses applied over millions of years, leading to a great reduction in yield strength. Evidence for geologically high strain rates is provided by **pseudotachylite** glasses whose formation results from ‘instantaneous’ frictional melting during deformation.

**strain-slip cleavage** A term synonymous with **crenulation cleavage**, but which some authorities (e.g. C. McA. Powell, 1979) have suggested should not be used in cleavage descriptions, owing to the implications it carries of the mode of origin.

**strain–time diagram** A diagram which illustrates a change in *\*strain* over time for a given *\*stress*. Strain–time diagrams are used to delimit fields of *\*elastic* behaviour, viscous behaviour and *\*failure*, and fields of *\*primary*, *\*secondary*, and *\*tertiary creep*.

**strandflat** A *\*shore platform*, up to 60 km wide, found along the coasts of Greenland, Iceland, Norway, and Spitzbergen. It may be the result of combined glacial and marine processes.

**strandline** The shoreline of a marine or *\*lacustrine* environment. The term is applied most commonly to ancient shorelines. The development of a strandline requires that the relative positions of land and water remain stable long enough for features to form. Subsequent displacement may be caused by a change in the level of the water or of the land.

**strandplain** A linear accumulation of *\*sand* or *\*gravel* that lies parallel to a coast, forming the commonest type of *\*beach*.

**strata** See STRATUM.

**strath terrace** A *\*river terrace* that is cut into bedrock and is covered by only a thin layer of sediment.

**stratified sampling** See SAMPLING METHODS.

**stratiform deposit** A mineral deposit that is *\*concordant* with *\*bedding*; usually in sheets but it may be ribbon-like.

**stratiformis** From the Latin *stratus* meaning ‘flattened’ or ‘spread out’ and *forma* meaning ‘appearance’, a species of cloud consisting of an extensive level sheet or layer, found in *\*altocumulus*, *\*stratocumulus*, and sometimes *\*cirrocumulus*. See also CLOUD CLASSIFICATION.

**stratigraphic** See STRATIGRAPHY.

**stratigraphic column** 1. A succession of rocks laid down during a specified interval of *\*geologic time*. The phrase ‘the stratigraphic column’ often refers to the whole sequence of *\*strata* deposited throughout geologic time. 2. A simplified columnar diagram relating a succession of named *\*lithostratigraphic units* from a particular area to the subdivisions of geologic time.

**stratigraphic correlation** Geologic study concerned with establishing geochronological relationships between different areas, based on geologic investigations of many local successions.

**stratigraphic cross-section** A section, usually with the vertical scale considerably exaggerated with relation to the horizontal scale, that is designed to show the thicknesses and stratigraphic relationships of successions of named **\*lithostratigraphic units**. For simplicity, the upper part of one of the units is restored to a horizontal position and topography is ignored. Items such as **\*facies** changes, **\*interdigitating** of units, **\*unconformities**, and breaks in succession are shown. The units and their boundaries are related to the subdivisions of **\*geologic time**.

**stratigraphic nomenclature** The naming of **\*stratigraphic** and **\*geologic-time** units according to established practices and principles. **\*Formal** naming of a stratigraphic unit occurs when the unit is first proposed and described from a type section (see **STRATOTYPE**), which acts thereafter as the standard reference for that unit. Ideally, the name given is binomial, and in the case of **\*chronostratigraphic** and **\*lithostratigraphic units** consists of a preceding geographic name taken from the **\*type locality** (plus lithological description where appropriate), followed by the name of the unit, e.g. Ludlow series and Elk Point Group. The names of **\*biostratigraphic units** consist of the name of the characteristic **\*fossil** plus the relevant unit term, e.g. *Monograptus uniformis* Range zone. The name chosen for a stratigraphic unit should be unique to that unit. When used as a proper name, as above, the initial letters are often capitalized. Except in very special circumstances the first formal name given has priority and is adhered to. In practice many well-known units, e.g. Coal Measures, Millstone Grit, were named long before the present conventions were established, and to avoid confusion these names are preserved in their original form. Geologic-time units generally take their preceding name from that of the corresponding chronostratigraphic unit, plus the name of the unit (**\*period**, **\*epoch**, **\*age**, etc.), e.g. the **\*Jurassic** period, from the Jurassic system (named after the **\*type area** in the Jura Mountains). The names of **\*eons** and **\*eras** (e.g. **\*Phanerozoic** eon, **\*Mesozoic** era) were proposed independently, so that the names for the corresponding **\*eonthems** and **\*erathems** are derived from the time units. See also **INFORMAL**.

**stratigraphic reef** A name proposed in 1970 by R. J. Dunham to describe a \*reef that comprises only a thick mass of pure or nearly pure \*carbonate rock. *Compare* ECOLOGIC REEF.

**stratigraphic scale** A general term to denote a timescale that incorporates both the traditional elements of the \*geologic time-scale as it has evolved over the last century and a half, and, as they are agreed, the reference points of an ideal and globally standardized \*chronostratigraphic scale that is defined by \*boundary stratotypes. The ideal, globally standardized stratigraphic scale has been termed both the \*Standard Stratigraphic Scale (SSS), and the \*Standard Global Chronostratigraphic Scale (SGCS). W. B. Harland (1978) has suggested that, for clarity, the older geologic timescale, which evolved through the designation of type sections (*see* STRATOTYPE) and is gradually being superseded, should be referred to as the Traditional Stratigraphic Scale (TSS). A local stratigraphic scale, from which a standard reference point might be selected later, is termed a Regional Stratigraphic Scale (RSS).

**stratigraphic trap (lithologic trap)** Oil or gas trap resulting from lithologic variations, e.g. interbedded lenses of \*sands and \*silts in a deltaic environment. *See* DELTA; NATURAL GAS; PETROLEUM. *Compare* ANTICLINAL TRAP; FAULT TRAP; REEF TRAP; STRUCTURAL TRAP; UNCONFORMITY TRAP.

**stratigraphic unit** A body of rock forming a discrete and definable unit. Such units are determined on the basis of their lithology (\*lithostratigraphic units), or their \*fossil content (\*biostratigraphic units), or their time span (\*chronostratigraphic units). It is unlikely that any rock succession will form a unit that accords with all three categories of classification. All stratigraphic units are defined by a \*type section. \*Geologic-time units are abstract concepts, not actual rock sequences, so do not class as stratigraphic units.

**stratigraphy** 1. The branch of the geologic sciences concerned with the study of stratified rocks in terms of time and space. It deals with the correlation of rocks from different localities. Correlation methods may involve the use of \*fossils (\*biostratigraphy), rock units (\*lithostratigraphy), or \*geologic-time units or intervals (\*chronostratigraphy). 2. The relative spatial and temporal arrangement of rock strata.

**stratocumulus** From the Latin *stratus* meaning ‘flattened’ or ‘spread out’ and *cumulus* meaning ‘heap’, a genus of low clouds composed of patches, sheets, or layers of grey to whitish appearance, always with dark patches, rolls, or rounded masses, which are not fibrous. *See also* CLOUD CLASSIFICATION.

**stratomere** A general term for any *\*chronostratigraphic unit* within the Stratomeric Standard hierarchy. *See* CHRONOSTRATIGRAPHY.

**stratopause** The level that marks the maximum height of the *\*stratosphere*, at around 50 km. After high temperatures in the upper stratosphere (about 0 °C at the stratopause) temperature decreases with increasing altitude in the *\*mesosphere* above. *See also* ATMOSPHERIC STRUCTURE.

**stratophenetic classification** *See* STRATOPHENETICS.

**stratophenetics (stratophenetic classification)** In *\*cladistics*, a method for determining the evolutionary relationships among organisms that exist only as fossils. It is based on quantitative assessments of morphological (i.e. phenetic) similarities and geologic age (derived from stratigraphy).

**stratosphere** The atmospheric layer above the *\*troposphere*, which extends on average from about 10 to 50 km above the Earth’s surface. The stratosphere is a major stable layer whose base is marked by the *\*tropopause*, and where temperatures overall average approximately –60 °C. Temperature in the lower stratosphere is isothermal but increases markedly in the upper part, to reach a maximum of about 0 °C at the *\*stratopause*. High stratospheric temperatures result from absorption of ultraviolet radiation (0.20–0.32 μm wavelengths) by ozone concentrated at 15–30 km. Due to the very low air density, even the small amount of ozone concentrated in the upper stratosphere is extremely effective in absorbing radiation, thus giving high temperatures at 50 km. The isothermal condition at the base of the stratospheric inversion layer creates stability, which generally limits vertical extensions of cloud and leads to the lateral spreading of high *\*cumulonimbus* cloud with characteristic anvil heads. *See also* ATMOSPHERIC STRUCTURE.

**Stratospheric Aerosol and Gas Experiment-III (SAGE-III)** A component of the *\*NASA* Earth Observing System, comprising a grating

spectrometer instrument that measures near-UV, visible, and near-IR energy through the Earth's limb during solar and lunar occultations and, during the daytime part of its orbit, limb scattering. The instrument was launched to the International Space Station on 19 February 2017.

**stratotype (type section)** An actual rock succession, chosen at a particular locality (the *\*type locality*) to act as the standard comparison for all other *\*chronostratigraphic* or *\*lithostratigraphic units* of its ilk. Generally, the type section of a *\*stratigraphic unit* should be the rock succession originally so designated and described (the *\*holostratotype*), but circumstances may require amendment (see [LECTOSTRATOTYPE](#); [NEOSTRATOTYPE](#)), or amplification (see [PARASTRATOTYPE](#); [HYPOSTRATOTYPE](#)). However, although many similar sequences may exist, only a single stratotype at any one time can act as the standard. See also [BOUNDARY-STRATOTYPE](#); [COMPOSITE-STRATOTYPE](#); [COMPONENT-STRATOTYPE](#); [TYPE AREA](#).

**stratovolcano (composite volcano)** *\*Volcano* built up of layers of *\*lava* alternating with beds of *\*ash* and other *\*pyroclastics* and with material eroded from higher slopes of the cone. Many of the world's highest volcanoes are of this type, and include Mt Fuji (Japan) and Mt. Egmont (North Island, New Zealand).

**stratum** (*pl. strata*) Lithological term applied to rocks that form layers or beds. Unlike 'bed', 'stratum' has no connotation of thickness or extent and although the terms are sometimes used interchangeably they are not synonymous.

**stratus** From the Latin *stratus* meaning 'flattened' or 'spread out', a genus of low clouds of flat, uniform base and of grey appearance, through which the Sun may be outlined clearly when the cloud is not too dense. When stratus forms at surface level it is called *\*fog*. See also [CLOUD CLASSIFICATION](#).

**streak** The colour of a mineral when in the form of powder, which is usually produced by scratching the solid mineral on an unglazed porcelain plate (streak plate). The colour may be different from the mineral's colour in mass.

**streak lightning** An electric discharge with a branching appearance of the main channel. It may flash between cloud and air or between cloud and ground.

**streak plate** See STREAK.


**streamer** A long (up to several kilometres) tube containing a number (sometimes many hundreds) of \*hydrophones and filled with oil, designed to be towed by a ship, and used in marine \*seismic surveying. It may be balanced to give it neutral buoyancy or have depth controllers to maintain it at a constant depth. Compass devices are attached at intervals to monitor the \*feather angle of the cable.

**stream flood** See FLASH FLOOD.

**stream grade** See GRADE.

**streamline** 1. In a flowing fluid, a hypothetical line which indicates the local direction of flow. See also HYDRAULIC GRADIENT; POTENTIOMETRIC SURFACE. 2. A shape which allows a body to offer minimum resistance to a fluid through which it moves; to impart such a shape to a body.

**stream order** Measure of the position of a stream (defined as the reach between successive tributaries) within the hierarchy of the \*drainage network. A commonly used approach allocates order '1' to unbranched tributaries, '2' to the stream after the junction of the first tributary, and so on. It is the basis for quantitative analysis of the network.



### Stream order

**stream power** The rate at which a stream can do work, especially the transport of its load, and measured over a specific length. It is largely a function of channel slope and discharge and is expressed by  $\Omega = \gamma Qs$ , where  $\Omega$  is the power,  $\gamma$  is the specific weight of water,  $Q$  is the discharge, and  $s$  is the slope. Streams tend to adjust their flow and channel geometry in order to minimize their power (see [LEAST-WORK PRINCIPLE](#)).

**stream-sediment analysis (river-sediment analysis, drainage-sediment survey)** A technique used in geochemical exploration and analysis, in which semi-mobile and immobile elements are measured from river or stream sediments, although under some circumstances highly mobile elements, e.g. molybdenum, may be used. Anomalies in sediments are not always accompanied by anomalies in the water, there may be seasonal variations in the composition of stream water, and it is easier to collect, carry, and store sediment samples than water samples. Anomalies may occur in the active sediment and in the banks and [\\*floodplains](#), so sampling of any of these can give satisfactory results. Immobile elements are determined by total-metal analysis; mobile and semi-mobile elements by total- or cold-extractable metal analysis.

**stream terrace** See [RIVER TERRACE](#).

**stress** A measure of the intensity of a force ( $F$ ) acting upon a body as a function of its area ( $A$ ), such that  $\text{stress} = F/A$ , in units of  $\text{N/m}^2$ . Stress can be resolved into two important components. Compressive (tensile) stress ( $\sigma$ ) acts normal to the surface (see **NORMAL STRESS**) and changes the volume of the body; **\*shear stress** ( $\tau$ ) acts parallel to the surface and changes the shape of the body. See also **STRAIN**.

**stress axial cross** Three mutually perpendicular **\*stress** axes whose lengths are proportional to the magnitudes of the principal stresses they represent. The maximum ( $\sigma_1$ ) and minimum ( $\sigma_3$ ) principal stress axes define the stress ellipse; the addition of the intermediate ( $\sigma_2$ ) principal stress axis defines the stress ellipsoid.

**stress difference** ( $\sigma_d$ ; **differential stress**) The simple difference between the greatest and least principal stresses, and used in the construction of **\*Mohr stress diagrams**. The diameter of a Mohr stress circle represents the stress difference for a given combination of  $\sigma_3$  and  $\sigma_1$  (see **STRESS AXIAL CROSS**). In **\*stress–strain diagrams** the stress axis is given as  $\sigma_d$  and is usually plotted against percentage **\*strain**. See **PRINCIPAL STRESS AXES**.

**stress field** The spatial change in the orientation of **\*stress** throughout a body of material, as opposed to the simple stress at a point. A stress field is represented as a grid of **\*stress trajectories** which may or may not be superimposed.

**stress meter** Instrument for measuring pressure changes in rocks that result from mining operations. The stress meter has a steel shaft with a groove containing glycerine. When pressure is exerted the glycerine is squeezed on to a diaphragm which pushes a strain gauge to measure the movement.

**stress–strain diagram** A diagram which illustrates the change in **\*strain** under an applied **\*stress**, usually as a function of the progressive change in temperature, pressure, and **\*strain rate**. The percentage strain is plotted on the x axis, and differential stress (the difference between the maximum ( $\sigma_1$ ) and minimum  $\sigma_3$ ) **\*principal stress** axes) on the y axis.

**stress trajectory** A line showing the continuous change in the orientation of a principal **\*stress** throughout a body. Although trajectories may curve,


their intersections with other principal stresses remain perpendicular.

**stretch** In *\*strain* and *\*stress* analysis, a measure of the extent by which an object of known dimensions has lengthened, expressed as the ratio of the new length to the old.

**strewnfield** An area which is associated with a specific group of *\*tektites* and microtektites that can be distinguished according to their age and chemical composition, and which probably represents a particular impact event. Four major strewnfields are known: the Australasian (formed 0.7 Ma ago), which is the largest and covers an area of about  $5 \times 10^7$  km<sup>2</sup> around Australia and South-east Asia; the Ivory Coast (formed 1.3 Ma ago) covering an area at least  $4 \times 10^6$  km<sup>2</sup> around and off the coast of W. Africa; the Central European (formed 14 Ma ago) without proven associated microtektites as yet; and the N. American strewnfield (formed 34 Ma ago) forming a belt stretching across the Pacific from Southeast Asia to the western Atlantic, and of unknown latitudinal extent.

**striation** Narrow groove or scratch cut in exposed rock by the abrasive action of hard rock fragments embedded in the base of a sliding *\*glacier*. Striation provides a useful clue to the direction of ice movement in formerly glaciated areas.

**strike** **1.** (*noun*) The compass direction of a horizontal line on an inclined plane. (*verb*) To lie in the direction of such a line. **2.** (*noun*) The discovery of an economically valuable source of a mineral. (*verb*) To make such a discovery.



**Strike**

**strike fault** A \*fault which strikes parallel with the \*strike of the layering (i.e. \*bedding or \*cleavage) in adjacent rocks.

**strike ridge** An elongated hill developed along the \*strike of a bed that is more resistant than its adjacent strata. *See also* CUESTA.

**strike-slip fault (wrench fault, tear fault, transcurrent fault)** A \*fault in which the major displacement is horizontal and parallel to the \*strike of a vertical or subvertical \*fault plane. Movements along such a fault may be \*dextral or \*sinistral. Localized zones of deformation due to pressures and tensions across the fault occur at bends in the fault strike and give rise to the formation of pull-apart basins and \*grabens, which are rhombic in shape. \*Flower structures are also recognized features of such regimes. There are certain geometrical similarities between strike-slip faults and \*transform faults, but also significant differences (e.g. displacement along the fault plane is equal and unlimited in a transform fault but not in a strike-slip fault). *See* TRANSPRESSION; TRANSTENSION.

**strike stream** *See* SUBSEQUENT STREAM.

**strike valley** *See* SUBSEQUENT STREAM.

**string 1. (flyer)** Up to ten \*geophones which are connected together permanently but have only one lead on to the seismic cable. **2. (drill string)** The rods (flights) and tools from the drill collar to the \*bit which, when connected together, enable a \*borehole to be drilled.

**strip mining** *See* OPEN-CAST MINING.

**stromatactis** A series of elongated cavities, with curved or irregular tops and flat bases, filled with \*calcite \*cements. Stromatactis cavities occur most commonly in carbonate and \*mud mounds. They were originally believed to be of organic origin, but currently they are thought to result either from the dewatering of \*lime muds or from the development of cavities beneath local cemented crusts on the sea floor.

***Stromatocystites walcotti*** The earliest known representative of the echinoderm (\*Echinodermata) group the Edrioasteroidea. It had a pentagonal shape with five distinct 'arms' apparent on the surface of a flexible, many-plated \*test. It existed during the Lower \*Cambrian.

**stromatolite** A laminated, mounded structure, built up over long periods of time by successive layers or mats of *\*cyanobacteria* that trapped sedimentary material. Stromatolites are found in shallow marine waters in warmer regions. Some are still in the process of being formed, e.g. those in Shark Bay, Western Australia; *\*fossil* stromatolites dating from the Early *\*Precambrian* are also known, although it is not certain that these were formed by cyanobacteria.

**Stromatoporoidea** Extinct group that has been attributed to the *\*Hydrozoa*, sponges (*\*Porifera*), foraminifera (*\*Foraminiferida*), *\*Bryozoa*, or *\*algae*, or regarded as a phylum with no modern representatives. Stromatoporoids are calcareous masses built up of horizontal layers (latilaminae) and vertical pillars. The calcareous skeleton is called the *\*coenosteum*. The upper surfaces show a pattern of polygonal markings and may have swellings (mammelons) and stellate grooves (astrorhizae). They are found in *\*limestones* of *\*Cambrian* to *\*Cretaceous* age, often forming *\*reefs* in *\*Ordovician* to *\*Devonian* times.


**Strombolian eruption** A type of volcanic activity which produces frequent, moderate *\*eruptions*. The *\*lava* is basaltic, but sufficiently viscous for entrapped gases to build up a pressure which is released in continuous small explosions. Lava, flung into the air, falls back to build up a steep-sided cone of interbedded *\*lava* and *\*tephra*. Lava flows are commonly erupted through breaches in the flanks of the cone. See *VOLCANO*. Compare *HAWAIIAN ERUPTION*; *PELÉÉAN ERUPTION*; *PLINIAN ERUPTION*; *SURTSEYAN ERUPTION*; *VESUVIAN ERUPTION*; *VULCANIAN ERUPTION*.

**strontianite** Mineral, SrCO<sub>3</sub>; sp. gr. 3.7; *\*hardness* 3.5–4.0; *\*orthorhombic*; white to pale green, grey, or pale yellow; white *\*streak*; *\*vitreous* *\*lustre*; crystals *\*prismatic* or *\*acicular*, but also fibrous and *\*massive*; *\*cleavage* prismatic, good {110}; occurs in low-temperature *\*hydrothermal* veins, often in *\*limestone*, and in association with *\*celestite*, *\*barite*, and *\*calcite*; soluble, with effervescence, in dilute hydrochloric acid. It is named after the type locality of Strontian, Highland Region, Scotland.

**strophic** See *HINGE*.

**Strophomenida (strophomenids)** (class *\*Articulata*) The largest order of brachiopods (*\*Brachiopoda*), now extinct, in which one *\*valve* is usually

convex and the other flat or concave. There is a straight **\*hinge** line. The **\*pedicle foramen** is filled by one **\*plate** in each valve in adult shells, the **\*pedicle** having degenerated and been lost. Attachment is by cementation of the pedicle valve, with or without tubular spines from the valves. The **\*shell structure** is **\*pseudopunctate**. Strophomenida appeared in the Lower **\*Ordovician** and became extinct in the Lower **\*Jurassic**. The order includes the **\*Carboniferous** **\*Productus giganteus** (*Gigantoproductus giganteus*), the largest of all brachiopods.



### Strophomenida

**strophomenids** See STROPHOMENIDA.

**structural contour map** A map on which structural features, e.g. **\*folds**, are represented in three dimensions. The map is read in the same way as a topographic contour map. The contours are based on a single **\*horizon** (e.g. the top of a bed), the position of that horizon being given with reference to a datum plane.

**structural geology** The study of the structure of rocks at all scales, and of the processes producing those structures.

**structural trap** Trap formed by deformation of porous and non-porous **\*strata** as a result of **\*folding**, **\*faulting**, etc., in which oil, gas, or water may accumulate. See NATURAL GAS; PETROLEUM; POROSITY. Compare ANTICLINAL TRAP; FAULT TRAP; REEF TRAP; STRATIGRAPHIC TRAP; UNCONFORMITY TRAP.

**structure grumeleuse** A *\*texture* in *\*limestones*, characterized by the presence of *\*micrite* clots completely surrounded by coarser, granular, *\*calcite* or *\*microspar*. This structure is thought to be produced by the selective *\*recrystallization* of the limestone, with larger crystals growing at the expense of smaller ones.

**Strutt, John William (Lord Rayleigh)** (1842–1919) A mathematician and physicist of Cambridge University, Rayleigh worked on optics, noble gases, wave mechanics, etc. In the field of Earth science, he studied radioactivity in rocks, and gave his name to a type of surface earthquake wave. See RAYLEIGH WAVES; RAYLEIGH NUMBER; RAYLEIGH SCATTERING.

**Sturtian** A stage in the *\*Cryogenian \*period*, from about 800 to 650 Ma ago (Int. Commission on Stratigraphy, 2004), preceded by the *\*Riphean* and followed by the *\*Varangian*.

**sturzstrom** A long-runout form of *\*mass-wasting*, in which the material moves a horizontal distance that is twenty–thirty times greater than the vertical distance. The word is German, *Sturz* meaning a fall and *Strom* a river.

**Stylasterina (branched hydrocorals)** (class *\*Hydrozoa*) Order of reef-building *\*Cnidaria* similar to *\*Milleporina*, but without free medusae. Stylasterina are known from the Upper *\*Cretaceous* to Recent.

**stylolite** An irregular, suture-like contact, produced by pressure dissolution (see PRESSURE SOLUTION) of rock under deep burial conditions. Stylolites are most commonly found in *\*limestones*, and may be picked out by the concentration of insoluble *\*clay* residues along the stylolite surface. Up to 40% of the original thickness of a limestone sequence can be dissolved through stylolitization.

**stylolitization** See STYLOLITE.

**sub-** From the Latin *sub* meaning ‘under’ or ‘close to’, a prefix meaning ‘beneath’ or ‘lying below’.

**Sub-arctic current** See ALEUTIAN CURRENT.

**subarkose** A *\*sandstone* characterized by the presence of less than 15% mud *\*matrix*, with between 5% and 25% of the *\*grains* being *\*feldspar*, and

there being more feldspar than rock fragments present. See [DOTT CLASSIFICATION](#).

**Sub-Atlantic** A colder, wetter climatic phase which followed more continental [\\*Sub-Boreal](#) times. The change from Sub-Boreal to Sub-Atlantic conditions in Britain is roughly coincident with the transition from Bronze to Iron Age cultures. The Sub-Atlantic marks a period of renewed [\\*peat](#) growth on bog surfaces that in late Sub-Boreal times were sufficiently dry and humified to support heath vegetation, e.g. *Calluna vulgaris* (ling, heather). This renewed peat growth gives a major recurrence surface, the Grenz horizon, which in Britain defines the Zone VIIb/Zone VIII (Sub-Boreal/Sub-Atlantic) boundary of the standard pollen stratigraphy. See [POLLEN ANALYSIS](#); [POLLEN ZONE](#).

**subbase** See [PAVEMENT](#).

**Sub-Boreal** From Scandinavian evidence, a cooler, drier, more continental climate phase that followed the [\\*Atlantic](#) climatic optimum (though with summers still warmer than those at present). In Britain, clear evidence for climatic change (e.g. differing [\\*peat](#) deposits) is lacking, and the [\\*Atlantic](#)/Sub-Boreal boundary is usually taken as the point of marked decline in elm pollen in the pollen stratigraphy (see [POLLEN ANALYSIS](#)). The reason for the elm decline (which, irrespective of elm species, is characteristic of pollen chronologies throughout Europe) has been the subject of much research. Hypotheses range from colder climate to epidemic disease to anthropogenic causes. The latter is linked to the selective use of elm as fodder for stalled livestock, now thought to have been a characteristic practice of Neolithic, forest-dwelling peoples. In many pollen chronologies this initial elm decline is followed quickly by a general decline in tree pollen caused by temporary forest clearance ('landnam') to provide land for slash-and-burn agriculture. The Sub-Boreal forms [\\*Pollen Zone VIIb](#), and lasted from about 5000 to 2800 BP.


**sub-continental lithospheric mantle (SCLM)** The uppermost part of the [\\*mantle](#), beneath continents, in which convection does not occur.

**subcritical reflection** In [\\*refraction surveying](#), [\\*seismic waves](#) which are incident on the refractor [\\*interface](#) at less than the [\\*critical angle](#) of incidence and are weakly reflected to the surface.

**subduction** The process of consumption of a lithospheric *\*plate* at convergent *\*plate margins*. See SUBDUCTION ZONE.

**subduction earthquake cycle** A sequence of events that occurs when two colliding *\*plates* become locked along some section of the subduction thrust (see SUBDUCTION ZONE). There is an interseismic period lasting for hundreds of years during which the upper plate is uplifted and shortened. Eventually, the accumulated *\*stress* exceeds the strength of the fault and the locked region fails, triggering a coseismic period, lasting a matter of minutes, during which the upper plate simultaneously subsides and extends horizontally, releasing the elastic *\*strain* and causing a major earthquake.

**subduction zone** The zone, at an angle to the surface of the Earth, down which a lithospheric *\*plate* descends. Most present-day subduction zones extend from *\*trenches* on the ocean floor, from where a zone of *\*earthquake \*hypocentres* (called a *\*Benioff zone*) extends, at an angle ranging from near-horizontal to near-vertical, to a depth of up to 700 km. Andesitic *\*volcanoes* form approximately 100 km above the subducting slab, and the presence of andesitic volcanoes in the geologic record is regarded as evidence of an ancient subduction zone and thus of a *\*destructive* plate margin.



**Subduction zone**

**subglacial** See ENGLACIAL.

**subgrade** See PAVEMENT.

**subgroup** A term that may be used for *\*formal* identification of a distinct and defined assemblage of *\*formations* within an already named *\*group* whose name is in use.

**subhedral (hypidiomorphic)** An *\*igneous* textural term applied to crystals which are only partly bounded by *\*crystal faces*. The irregular parts of the crystal surface may be caused by partial corrosion of the crystal or by partial intergrowth with other crystals.

**sublimate** A solid substance that has condensed directly from a gas.

**sublimation** Direct evaporation from ice. In meteorology, the term is also often applied to the reverse process, in which water vapour changes directly to the solid phase, a process more correctly called *\*deposition*. See also ABLATION.

**sublitharenite** A *sandstone* characterized by the presence of less than 15% mud *matrix*, with between 5% and 25% of the *grains* being rock fragments, and there being more rock fragments than *feldspar* present. See DOTT CLASSIFICATION.

**sublittoral zone** 1. In freshwater *ecosystems*, an alternative name for the limnetic zone. 2. The sea-shore zone lying immediately below the *littoral (intertidal) zone* and extending to a depth of about 200 m or to the edge of the *continental shelf*. Red and brown *algae* are characteristic of this area. Typical animals include sea anemones and corals on rocky shores, and shrimps, crabs, and flounders on sandy shores. The zone may alternatively be called the subtidal zone. It is approximately equivalent to the *circalittoral* zone.

**submarine canyon** Deep, steep-sided valley cut into the *continental shelf* or slope, whose axis slopes seaward at up to 80 m/km. The development of submarine canyons is thought to have originated through erosion by *turbidity currents*. However, few turbidity currents have been recorded from the submarine canyons that have been studied.

**submersible** Underwater vehicle, used for oceanographic investigation or offshore engineering. These small submarines may be manned or unmanned.

**sub-metallic** Applied to the *lustre* of a mineral which is intermediate between *metallic* and *non-metallic*. Varieties of *chromite* or *sphalerite* may give a rather dull, metallic appearance on occasions and might therefore be described as sub-metallic. There is no sharp distinction between 'metallic' and 'sub-metallic'.

**sub-Plinian eruption** See VESUVIAN ERUPTION.

**subpolar glacier** See GLACIER.

**subsequent stream** A stream that follows a line of geologic weakness, such as the *outcrop* of a soft bed, a sequence of major *joints*, a *fault trace*, or the axis of an *anticline*. Such a stream tends to extend headwards actively, and may acquire further tributaries through the process of *river capture*. It is called a 'strike stream' when its trace follows that geologic *strike*, and the associated valley is called a 'strike valley'.

**subsidence** **1.** A progressive depression of the Earth's *\*crust*, which allows *\*sediment* to accumulate and be preserved. Subsidence is caused by *\*mantle convection* and by sediment loading. The subsidence rate will control the proportion of deposited sediment which will be preserved in the subsiding area. Subsidence rates in *\*sedimentary basins* typically vary from 0.3 to 2.5 mm per year. **2.** Sinking or settling of the ground surface due to natural or anthropogenic causes. Surface material with no free side is displaced vertically downwards with little or no horizontal movement. **3.** Local sinking, due to underground mine workings. **4.** Downward movement of air, characteristically gentle (1–10 cm/s) and often in large *\*anticyclones*, which is related to the *\*divergence* in lower layers near the ground surface. Subsidence results from radiation cooling or from convergence of air horizontally in the upper *\*troposphere*. It typically brings settled weather, with evaporation of cloud drops by *\*adiabatic* warming in the subsiding *\*air mass*, causing cloudless skies above the friction layer near the surface, so that in winter fog and low cloud may prevail when moisture is sufficient.

**subsoiling** The breaking up of subsoils, usually because they are compacted, without inverting them. Subsoiling is usually performed with a chisel-like device that is pulled through the soil.

**subsolvus granite** An *\*igneous* rock of granitic composition characterized by the presence of two types of *\*alkali* feldspar: a potassium-rich type displaying perthitic (see **PERTHITE**) texture; and a sodium-rich type displaying antiperthitic texture. Subsolvus granites have a high water content, which depresses their liquidii sufficiently to intersect the subsolidus *\*solvus* surface and cause two extremes of feldspar composition to crystallize from the *\*melt*, instead of one intermediate-composition feldspar as in *\*hypersolvus* granites.

**subsolvus syenite** See **SYENITE**.

**substage** A subdivision of a *\*stage*. See **CHRONOZONE**.

**subsurface flow (interflow, throughflow)** The flow of water at a shallow depth beneath the ground surface, that occurs when rain falls faster than it can infiltrate downwards. The subsurface flow re-emerges at the surface at or near the base of ground slopes.

**subtidal** Applied to that portion of a **\*tidal-flat** environment which lies below the level of mean low water for **\*spring tides**. Normally it is covered by water at all states of the **\*tide**. The word is often used as a general descriptive term for a subaqueous but shallow-marine depositional environment.

**subtractive primary colours** The colours cyan, magenta, and yellow, which can be subtracted from white light to produce all other colours. *See also* **ADDITIVE PRIMARY COLOURS**.

**subtropical high** Surface high-pressure cells, especially prominent and persistent over oceans at around 30° latitude in both hemispheres. The **\*anticyclones** develop below the **\*subtropical jet stream** from subsiding air. The development tends to shift equatorward in winter and poleward in summer. The high pressure is most strongly developed in summer, when the highs frequently block depressions travelling eastward. *See also* **AZORES HIGH**; **BERMUDA HIGH**.

**subtropical jet stream** **\*Jet** stream of subtropical latitudes. The jet is related to a marked temperature gradient in the upper **\*troposphere**. The jet moves equatorward in winter and is associated with subsiding air and settled surface weather. In summer the jet moves poleward. At the seasonal extremes it tends at times to merge with the **\*polar-front jet**, but it persists throughout the year.

**subzone** Division of the fundamental unit (**\*zone**) used in **\*biostratigraphy**. The demarcation of a subzone is based on the **\*fossil** subspecies or assemblage contained within the rock sequence studied.

**sucrosic limestone** *See* **DUNHAM CLASSIFICATION**.

**Suess, Eduard** (1831–1914) Professor of geology in Vienna, Suess published his important work on structural geology, *Das Antlitz der Erde* ('Face of the Earth'), between 1833 and 1909. He studied mountain building, especially the Alps which he believed to have been formed in a **\*geosyncline**, which he named **\*Tethys**. He opposed the concept of **\*isostasy**, arguing that subsidence of the ocean floors had caused what he termed 'eustatic' changes in sea level. *See* **OROGENY**; **EUSTATIC**.

**Suess wiggles** Small oscillations, mostly lasting 2–3 years, between the  $^{14}\text{C}$  determinations and **\*dendrochronological** age determinations of the same piece of wood. These were thought originally to be instrumental, but are now considered to reflect genuine changes in  $^{14}\text{C}$  productivity of uncertain cause.

**suevite** A **\*breccia** of rock fragments in a **\*matrix** of **\*glass**, found within **\*meteorite** impact craters near the site of impact. The shock waves associated with the meteorite impact produce extremely high pressures and temperatures in the rocks for a few microseconds. Near the point of impact, these can brecciate the rock and also melt it. When the melt chills after the passage of the shock wave it produces the glass component known as '**\*impactite glass**'.

**suffusion** Spreading out of material on the substratum.

**Suisei** A Japanese **\*JAXA** mission to comet **\*Halley**, launched in 1985. It reached Halley on 8 March 1986 at a distance of 151 000 km and was then redirected to comet **\*Giacobini–Zinner** and from there to comet **\*Temple–Tuttle**. The spacecraft ran out of fuel in 1991 and contact with it was lost.



<http://www.isas.jaxa.jp/en/missions/spacecraft/past/suisei.html>

- Suisei.

**sulci** See **SULCUS**.

**sulcus** (*pl.* **sulci**) Latin for groove or furrow. **1.** A complex region of parallel ridges and furrows on a **\*satellite** surface, particularly well developed on Ganymede (e.g. Uruk Sulcus, an area of bright, grooved terrain bordering Galileo Regio). **2.** A major, rounded depression on the longitudinal mid-line of a brachiopod (**\*Brachiopoda**) shell. It usually occurs in the **\*ventral \*valve** and is usually accompanied by a major, rounded, shell elevation (fold) in the other valve. **3.** See **DINOPHYCEAE**. **4.** See **POLLEN**.

**sulphates** Group of non-silicate **\*minerals** in which the  $\text{SO}_4^{2-}$  radical is in combination with a number of metal **\*cations**. Examples include **\*barite** ( $\text{BaSO}_4$ ), **\*celestite** ( $\text{SrSO}_4$ ), **\*anglesite** ( $\text{PbSO}_4$ ), and **\*anhydrite** ( $\text{CaSO}_4$ ).

**\*Gypsum** ( $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ ) is the most common of a number of hydrated sulphates which also occur. Sulphates are normally colourless or white, soft (**\*hardness** about 3), **\*massive** or **\*earthy**, but **\*tabular** when crystalline. They are low-temperature minerals and occur as **\*gangue** minerals in hydrothermal veins and as chemical precipitates and **\*evaporites**.

**sulphides** A group of **\*minerals** in which the element sulphur (S) is in combination with one or more metallic elements. Simple sulphides include the common **\*ore minerals** **\*galena** (PbS), **\*sphalerite** (ZnS), and **\*pyrite** ( $\text{FeS}_2$ ). Two metallic **\*cations** may also be present, as in **\*chalcopyrite** ( $\text{CuFeS}_2$ ). More complex combinations may also occur to give 'double sulphides' or 'sulpho-salts' in which metallic and metalloid or non-metallic elements are present in combination with sulphur, e.g. **\*tetrahedrite** ( $\text{Cu}_{12}\text{Sb}_4\text{S}_{13}$ ) and **\*enargite** ( $\text{Cu}_3\text{AsS}_4$  or  $3\text{Cu}_2\text{S} \cdot \text{As}_2\text{S}_5$ ).

**sulpho-salts** See SULPHIDES.

**sulphur, native** Non-metallic element, S; sp. gr. 2.0; **\*hardness** 2.0; yellow; **\*massive**, or **\*tabular** when crystalline; produced by **\*fumarole** volcanic activity and by hot springs, and recovered commercially from bedded sedimentary deposits associated with **\*gypsum** and **\*salt domes**. Most sulphur is now obtained as a by-product of oil-refining, since it is a common contaminant of natural oil.

**sumatra** Regional squall, usually occurring at night during the south-west monsoon, in the Malacca Strait, accompanied by high winds which veer (see **VEERING**) from southerly to south-westerly and north-westerly. Extensive **\*cumulonimbus** cloud brings heavy rain, with thunder and lightning.

**Sun** The central star (G spectral type) in the **\*solar system**, 696 000 km in radius,  $333\,000 \times$  **\*Earth** mass,  $1\,300\,000 \times$  Earth volume, and with a mean density of  $1410 \text{ kg/m}^3$ . The equator is inclined at  $7.25^\circ$  to the plane of the **\*ecliptic**. It is principally composed of hydrogen and helium. The visible surface is the 'photosphere' (temperature 6000 K). The Si-normalized solar abundances for the terrestrially non-gaseous elements match those of the C1 **\*carbonaceous** chondrites.

**suncracks** See DESICCATION CRACKS.

**Sundaland** Name commonly given to the unit composed of Malaya, Sumatra, Java, and Borneo, with the intervening small islands; these are linked by the shallow-water (less than 200 m) Sunda shelf, which was exposed during periods of low sea level in the *\*Pleistocene*.

**Sundance Sea** A shallow marine embayment that extended over what are now Wyoming and S. Dakota during the late *\*Callovian* and *\*Oxfordian*. The southern edge of this sea (in modern Colorado) was bordered by *\*tidal flats*, the marine connection being northwards through the present-day mid-west and Canada. The *\*sediments* resulting from deposition in the Sundance Sea are characterized by a rich ammonite (*\*Ammonoidea*) fauna (*Quenstedtoceras* and *Cardioceras*), especially the Redwater Shale Member of the Sundance *\*Formation*.

**sunshine recorder** See CAMPBELL–STOKES SUNSHINE RECORDER.

**Sun-synchronous orbit** A satellite orbit that remains constant in relation to the Sun, passing close to both poles and crossing the meridians at an angle. The orbit, at a height of about 860 km (one-seventh of an Earth radius), takes about 102 minutes and carries the satellite over a different swathe of territory at each pass, so every point on the surface is overflowed every 12 hours, at the same solar times each day. Compare GEOSYNCHRONOUS ORBIT; see also POLAR ORBIT.

**sun-tan age** See EXPOSURE AGE.

**Suomi NPP** A National Polar-orbiting Partnership (NPP) mission between *\*NASA* and the Integrated Program Office of NOAA (National Oceanic and Atmospheric Administration) that is named for Verner Suomi (1915–95), recognized by many as the ‘father of satellite meteorology’. The mission supplies multispectral imagery on clouds, land, and oceans and on temperature and atmospheric humidity profiles and ozone distribution. The mission was launched on 28 October 2011, from California, into a Sun-synchronous near-circular orbit at an altitude of 824 km.

**super-** From the Latin *super* meaning ‘on top of’, a prefix meaning ‘directly over’, ‘over’, or ‘above’.

**super-adiabatic lapse rate** A fall of temperature with increasing altitude, which is greater than the usual *\*dry adiabatic lapse rate*; it occurs in

conditions of intense heating over land or sea.

**supercell** A very large convection cell that forms within a *\*cumulonimbus* by the merging of several smaller convective cells. Inside the supercell air may rise at 45 m/s and the cell may extend to a height of more than 16 km, breaking through the *\*tropopause*. Supercells last much longer than ordinary convective storm cells, because the warm updraughts rise at an angle to the vertical and the cold downdraughts associated with precipitation subside to the side of them, so the sinking cold air does not chill the rising warm air, thereby suppressing the vertical circulation. Supercells produce extremely violent storms, and they can trigger *\*tornadoes*. See also *MESOCYCLONE*.

**supercontinent** A continental mass which includes several of the *\*cratons* of the present-day continents. Examples include *\*Pangaea*, *\*Gondwana*, and *\*Laurasia*.

**supercooled cloud** Cloud containing pure water droplets at temperatures considerably below the nominal freezing temperature of 0 °C. With very pure water (i.e. free from pollutants), in the absence of freezing nuclei, supercooling of liquid drops can occur down to around -40 °C; *\*altocumulus* cloud, for example, is usually composed of water droplets at temperatures well below 0 °C. See also *CLOUD SEEDING*.

**supercooling (undercooling)** The cooling of a liquid to a temperature lower than its normal freezing temperature.

**supercritical fluid** See *CRITICAL TEMPERATURE*.

**supercritical reflection** In *\*refraction surveying*, *\*seismic waves* which are incident on the refractor *\*interface* at greater than the *\*critical angle* of incidence and are strongly reflected to the surface. Such waves travel with the *\*seismic velocity* of the first layer (therefore more slowly than the refracted *\*head waves*).

**supergene enrichment (secondary enrichment)** Re-precipitation of sulphides and oxides by descending acidic *\*groundwater* which has leached the surface zone of an *\*ore* deposit (see *GOSSAN*); this upgrades the deposits *in situ*, as in *\*porphyry copper* ores.

**supergroup** A term that may be used for *\*formal* identification of an assemblage of related and adjacent *\*groups*, or related and adjacent *\*formations* and groups.

**superimposed drainage (epigenetic drainage)** A *\*drainage pattern* that has been established on an earlier surface (perhaps conformable with the immediately underlying strata, and standing well above the present landscape). Subsequently the pattern was lowered by river incision so it now lies across geologic structures to which it bears no relation.

**superinterval** The time between *\*hairpins* on a palaeomagnetic apparent *\*polar wander path*, usually hundreds of millions of years.

**Superior-type granular iron formation** A *\*granular iron* deposit that is thick and extensive, sometimes covering  $10^5$  km<sup>2</sup>, originally recognized around Lake Superior, that formed in a large basin or across a stable shelf. *See also* ALGOMA-TYPE GRANULAR IRON FORMATION; RAPITAN-TYPE GRANULAR IRON FORMATION.

**superposition** *See* LAW OF SUPERPOSITION.

**supersaturation** The condition of air in which the humidity is above the level required for saturation at a given temperature (i.e. the *\*relative humidity* is greater than 100%). Supersaturation results when the temperature of air containing no *\*cloud condensation nuclei* falls below its *\*dew* point. *See also* SATURATED AIR.

**Supersaurus** *See* SAUROPODA.

**SuperView Earth Observation Constellation** *See* GAOJING.

**supplementary forms** *\*Crystals* whose faces have developed in different positions relative to their atomic structure but which nevertheless have the same general *\*crystal symmetry*. They may be distinguished as 'positive' or 'negative' forms, or as 'diploids'.

**suppressed layer** In *\*electrical \*resistivity* depth sounding, a thin layer whose true resistivity is intermediate between those of the layers above and below it, so it may be masked and its effects suppressed. It is analogous to the *\*hidden* layer in *\*refraction* seismology.

**supra-** From the Latin *supra* meaning ‘above’, ‘beyond’, or ‘earlier in time’, a prefix meaning ‘above’ or ‘in a superior position to’.

**supraglacial** See ENGLACIAL.

**supralittoral zone** The seashore zone immediately above the littoral fringe and beyond the reach of tidal submergence, though affected by sea spray.

**supratidal** Applied to that portion of a **\*tidal flat** which lies above the level of mean high water for **\*spring** tides. It is inundated only occasionally by exceptional **\*tides** or by tides augmented by a **\*storm surge**.

**surf** Breaking waves in the area between the shoreline and the outermost limit of breaking waves.

**surface inversion** A **\*temperature inversion** in the lower atmospheric layers, extending upwards from the Earth’s surface. The condition results, for example, from radiation cooling of the ground and the air above, or from **\*advection** of warm air over cold surfaces.

**surface runoff (overland flow, Hortonian flow)** The flow across the land surface of water that accumulates on the surface when the rainfall rate exceeds the **\*infiltration capacity** of the **\*soil**. The rate of infiltration, and therefore the possibility of surface runoff, is determined by such factors as soil type, vegetation, and the presence of shallow, relatively impermeable, **\*soil horizons**. Saturated overland flow can occur when a temporary rise of the **\*water-table** inhibits infiltration and causes flow over the surface.

**surface tension ( $\gamma$ )** Fluid surfaces may take on the behaviour of a stretched elastic membrane as a result of the tendency of a liquid surface to contract. The surface tension of a liquid is given as the tension across a unit length of the fluid surface. Surface tension is temperature-dependent and is closely associated with **\*capillarity**.

**Surface Water Ocean Topography mission (SWOT)** A mission by **\*NASA** and the Jet Propulsion Laboratory to provide accurate monitoring of local changes in sea level at coasts and changes affecting all surface water bodies, including lakes, reservoirs, and wetlands, with a surface area greater than 250 m<sup>2</sup>. The mission is scheduled to launch in September 2021.



<https://swot.jpl.nasa.gov/mission.htm>

- SWOT mission homepage.

**surface wave** A *\*seismic wave* which propagates along the surface of a medium rather than through it, e.g. *\*Love waves* and *\*Rayleigh waves*. Compare **BODY WAVE**.

**surface wind** The wind close to the Earth's surface, the velocity of which is usually measured at a standard height of 10 m. Surface-wind velocity is reduced by the frictional effect of the underlying surface. The actual wind is a balance of *\*pressure-gradient force*, *\*Coriolis effect*, and frictional effects.

**surf wave** See **SPILLING BREAKER**.

**surge** An expanded, turbulent, dilute flow of gas and *\*pyroclasts*. Three main types are currently recognized. *\*Base surges*, which are cold and wet, are generated during phreatomagmatic or phreatic eruptions; ground surges, which are hot and dry, are generated from the head of pyroclastic flows; and ash-cloud surges, which are also hot and dry, are generated from the overriding gas and *\*ash* cloud above pyroclastic flows.

**Surtseyan eruption** A high-energy volcanic *\*eruption* which occurs when sea or lake water floods into the top of an active open vent, producing an eruption column up to 20 km high and pyroclastic *\*clasts* with extreme fragmentation and moderate dispersal. The term was first used to describe the activity at the new *\*volcano*, Surtsey, which built up on the ocean floor south of Iceland in 1963.

**Surveyor** A series of seven *\*NASA* lunar lander missions that ran from 1966 to 1968. Surveyors 1, 3, 5, 6, and 7 landed on the Moon; Surveyor 2 crashed on the Moon and radio contact with Surveyor 4 was lost.



<http://nssdc.gsfc.nasa.gov/planetary/lunar/surveyor.html>

- A NASA programme to land a series of probes on the Moon.

**survivorship curve** Graphic description of the survival of individuals in a population, from birth to the maximum age attained by any one member. Usually it is plotted as the logarithm of the number of survivors as a

function of age. If a population has a constant mortality rate the graph will be a straight line. The technique may also be used to plot the survivorship of whole populations, *\*species*, genera, or higher *\*taxa*. See also *COHORT*.

**susceptibility, magnetic** See *MAGNETIC SUSCEPTIBILITY*.

**susceptibility meter** An instrument for measuring the *\*magnetic susceptibility* of a sample. It may be low field (less than 10 mT), intermediate (10–100 mT), or high field (more than 100 mT).

**suspect terrane** An area or region that is suspected of being a *\*terrane*, but whose boundary *\*faults* have not been identified.

**suspended load** The part of the total load of a stream that is carried in suspension. It is made up of relatively fine particles that settle at a lower rate than the upward velocity of water eddies. Its highest concentration is in the zone of greatest turbulence, near the bed. It reaches a maximum in shallow streams of high velocity.

**suspension feeder** An aquatic animal that obtains food by straining particles that are suspended in the water, usually by means of tentacles bearing many cilia (see *CILIUM*). Animals feeding near the sea surface feed mainly on *\*plankton*; those dwelling on the sea bed rely more on *\*detritus* drifting down from the more densely populated surface waters.

**sutural angle** See *SUTURE*.

**suture** **1.** A linear belt of highly deformed rocks, including tectonic *\*mélanges*, lenses of *\*ophiolites*, deep-sea *\*sediments*, and usually *\*blueschists*, which is interpreted as the boundary between two collided continents or *\*island arcs*. The location of a suture between collided masses has often led to controversy, and the recognition that *\*collision zones* are in some cases a mosaic of jumbled, sliced, and rotated *\*terranes* has led to the realization that sutures may be diffuse, rather than a narrow belt as was formerly thought. **2.** The line marking the junction between the septa (see *SEPTUM*) and the external wall of a cephalopod (*\*Cephalopoda*) shell that is visible when the shell has been preserved as an internal mould. In some cephalopods the suture lines are simple curves but in ammonoids (*\*Ammonoidea*) the suture becomes crenulate; bends in the suture line that point anteriorly are called ‘saddles’, those pointing posteriorly ‘lobes’. In

gastropods (*\*Gastropoda*), the suture is the line of junction between two *\*whorls* of the shell; the angle the line makes with the horizontal is the 'sutural angle'. **3.** See [CEPHALIC SUTURE](#).

**Svecofennian orogeny** An early to middle *\*Proterozoic* mountain-building episode that affected the Baltic Shield, in what are now Sweden and southern Finland, and that occurred approximately 1900–1800 Ma ago, at approximately the same time as the *\*Hudsonian* and *\*Laxfordian* orogenies.

**Sveconorwegian orogeny** See [DALSLANDIAN OROGENY](#).

**Svedberg unit (S)** The unit of measurement in which *\*sedimentation* coefficients are expressed; it refers to the time it takes for a particle of specified size and shape to settle. It is equal to  $10^{-13}$  seconds, and is written with no space between the number and the symbol, e.g. 64S.

**swale** **1.** A long, narrow depression, approximately parallel to the shoreline, between two ridges on a beach. **2.** A depression in otherwise level ground. **3.** A shallow depression in the undulating surface of a ground *\*moraine*, caused by uneven deposition by the glacier.

**swaley cross-bedding** A form of *\*hummocky* cross-bedding in which there are few or no hummocks, but *\*swales* are preserved.

**Swallow buoy** See [NEUTRALLY BUOYANT FLOAT](#).

**swallow hole** See [DOLINE](#).

**swallowtail twinning** A type of *\*crystal* twinning in which one crystal divides into two along a *\*twin plane*, giving a twinned crystal in the shape of a 'V' or swallowtail. The mineral *\*gypsum* may develop this twinned form.

**Swarm** A three-*\*minisatellite* constellation by the Danish National Space Centre, within the Earth Explorer Opportunity Program of the *\*European Space Agency*, that maps the *\*geomagnetic* field from low Earth orbit. The constellation was launched on 22 November 2013, from the Plesetsk Cosmodrome, Russia, into orbits 3–9 hours apart; Swarms A and C are at an altitude of 460 km or lower, Swarm B at 530 km or lower; Swarms A and C are at an orbital inclination of 87.4°, Swarm B at 88°.

**swash** The turbulent uprush of water that occurs when a wave breaks on a beach. It is an important mechanism for transporting sand and shingle landwards.

**S-wave (secondary wave, shear wave, transverse wave)** An elastic **\*body wave** in which particles oscillate about a fixed point but in a direction perpendicular to the direction of propagation of the wave energy. S-waves cannot travel through a fluid, since a fluid cannot support shear. In an **\*isotropic** and homogeneous medium, S-wave velocity ( $V_s$ ) is given by:  $V_s = \sqrt{(\mu/\rho)}$ , where  $\mu$  is the **\*shear modulus** and  $\rho$  the density of the material. S-waves travel at about half the speed of **\*P-waves** in a given medium, and can be polarized into SV- and SH-waves in which particle motion is restricted to the vertical and horizontal planes respectively. S-waves can also be generated from the non-normal incidence of P-waves on to an **\*interface**, when they are known as 'converted waves' and are mostly SV-waves. S-waves can also be converted into P-waves at an interface.

**Swazian** A S. African **\*stage** of the **\*Archaean**, from about 4000 to 3000 Ma ago, followed by the **\*Randian**, that includes the Onverwacht, Figtree, and Pongola sequences of S. Africa.

**sweepstakes dispersal route** Term coined by G. G. **\*Simpson** in 1940 to describe a possible route of faunal interchange which is unlikely to be used by most animals, but which will, by chance, be used by some. It requires a major barrier that is occasionally crossed. Which groups cross and when they cross are determined virtually at random.

**swell** Long-period waves that have built up sufficient energy to move away from the area where wind stresses created them. The waves assume a uniform pattern and move even through areas where winds are weak or absent. The longer-period waves move faster than shorter-period waves, so the waves spread out as they move away from the storm (dispersion). Swell waves generated south of New Zealand have been recorded arriving on the coast of Alaska.

**swelling coefficient** The amount of swelling pressure generated by the hydration of some **\*clay minerals**, or when water converts to ice. It is measured by an **\*oedometer**.

**swirl** An enigmatic, light-coloured marking on a **\*satellite** or planetary surface. Swirls have no topographic relief. Examples are known from **\*Mercury** and the **\*Moon** (Reiner Gamma). They appear to be antipodal to major impact structures. The Reiner Gamma swirl is associated with strong surface magnetization. It has been suggested that some may be cometary impact sites.

**SwissCube** A 1-unit **\*CubeSat** mission by several Swiss universities to observe airglow phenomena. The CubeSat was launched on 23 September 2009, from Satish Dhawan Space Centre, India, into a Sun-synchronous near-circular orbit at an altitude of 720 km.

**SWOT** See SURFACE WATER OCEAN TOPOGRAPHY MISSION.

**Sycorax (Uranus XVII)** A lesser satellite of **\*Uranus** with a radius of 75 km and a visual **\*albedo** of 0.07.

**syenite** A **\*saturated**, coarse-grained, **\*igneous** rock consisting of **\*essential** **\*alkali** feldspar and **\*ferromagnesian** minerals (**\*biotite**, **\*hornblende**, **\*arfvedsonite**, **\*aegirine-\***augite, and/or aegirine) and **\*accessory** **\*apatite**, **\*zircon**, and iron oxides. The feldspar constitutes more than 65% of the rock. Hypersolvus syenites are characterized by one type of alkali feldspar, usually potassium-rich and displaying perthitic (see **PERTHITE**) texture. Subsolvus syenites are characterized by two types of alkali feldspar, a potassium-rich type displaying perthitic texture and a sodium-rich type displaying antiperthitic texture. Syenites, which are the **\*plutonic** equivalents of **\*trachytes**, are found as ring complexes and as discrete **\*intrusions** on the stable **\*continental crust** and in the cores of some off-axis, ocean-island **\*volcanoes**.

**syenodiorite** See **MONZONITE**.

**syenogabbro** A coarse-grained **\*igneous** rock consisting of **\*essential** **\*alkali** feldspar, calcium-rich **\*plagioclase**, and **\*ferromagnesian** minerals (**\*augite** and **\*biotite**), and **\*accessory** **\*apatite**. The two feldspar types are in equal proportions. The rock is mineralogically half-way between a **\*gabbro** and a **\*syenite**.

**syenoids** **\*Igneous** rocks of **\*syenite** affinity in which **\*feldspathoidal** minerals take the place of **\*alkali** feldspar. **\*Ijolite** is a syenoid.

**sylvite** Mineral, KCl; sp. gr. 2.0; **\*hardness** 2; **\*cubic**; colourless to white, but sometimes shades of blue, yellow, or red; vitreous **\*lustre**; crystals usually **\*cubes**, often in combination with **\*octahedra**; **\*cleavage** perfect cubic; occurs in bedded **\*evaporite** deposits, but is one of the last minerals to precipitate because of its solubility in water; tastes much more bitter than **\*halite**. It is used extensively as a fertilizer.

**symbiosis** A general term describing the situation in which dissimilar organisms live together in close association. As originally defined, the term embraces all types of mutualistic and parasitic relationships. In modern use it is often restricted to mutually beneficial species interactions.

**symmetrical extinction** In optical **\*mineralogy**, the phenomenon which occurs when the **\*vibration** direction of the light bisects the angles between two sets of **\*cleavages** (as seen in basal sections of **\*pyroxenes** and **\*amphiboles**). A special form of symmetrical **\*extinction** may also occur in twinned crystals (see **CRYSTAL TWINNING**) of **\*feldspar**, and this may be used to determine their composition.

**symmetrical fold** A **\*fold** in which the limbs are of equal length.

**symmetrical trend** See **BOW TREND**.

**Symmetrodonta** (infraclass **\*Pantotheria**) Extinct order of **\*mammals** which lived during the late **\*Jurassic** and early **\*Cretaceous**; they may have appeared toward the end of the **\*Triassic**, making them among the earliest mammals known. They had molar teeth with three cusps arranged in symmetrical triangles. They were very small and probably were predators. They are believed to be ancestral to the marsupial and placental mammals. See also **MARSUPIALIA**; **EUTHERIA**.

**symmetry plane** See **CRYSTAL SYMMETRY**.

**sympatric evolution** The development of new **\*taxa** from the ancestral taxon, within the same geographic range; it is geographically possible for interbreeding to occur between the potential new taxa, but for some reason this does not happen. Because of the difficulty of envisaging what the reasons might be, until recently few authorities accepted the reality of sympatric evolution, except for certain special kinds of organism; but recent

studies have shown that **\*chromosomal** mutation can set up a partial barrier to interbreeding, sufficient to permit sympatric speciation.

**sympatry** The occurrence of **\*species** together in the same area. The differences between closely related species usually increase (diverge) when they occur together, in a process called character displacement, which may be morphological or ecological.

**Symphyla** See MYRIAPODA.

**symplectic texture** In **\*petrology**, a **\*texture** characterized by the intergrowth of two **\*minerals** that crystallized simultaneously. A rock with such a texture is called a symplectite.

**symplectite** See SYMPLECTIC TEXTURE.

**symplesiomorphy** The possession of a **\*character** state that is **\*primitive** (**\*plesiomorphic**) and shared between two or more taxa. Shared possession of a symplesiomorph character state is not evidence that the taxa in question are related.

**syn-** From the Greek *sun* meaning ‘with’, a prefix meaning ‘together’, ‘with’, or ‘resembling’.


**synaeresis** The process of subaqueous shrinkage of **\*clays** by the loss of **\*pore** water. The loss of pore waters occurs either because of the change in the volume of some **\*clay minerals** as a result of **\*salinity** changes, or by **\*flocculation** of the clays. The shrinkage results in the production of **\*synaeresis cracks**.

**synaeresis cracks** Irregular, radiating, lenticular-shaped cracks, found on bedding surfaces and often resembling the form of a bird’s foot. These cracks form by subaqueous shrinkage rather than desiccation, and are therefore not an indication of subaerial exposure. Compare **\*DESICCATION CRACKS**.

**synapomorphy** The possession of **\*apomorphic** features by two or more taxa in common (i.e. the features are shared, derived). If the two groups share a **\*character state** that is not the **\*primitive** one, it is plausible that they are related evolutionarily, and only synapomorphic character states can

be used as evidence that taxa are related. **\*Phylogenetic trees** are built by discovering groups united by synapomorphies.

**Synapsida (mammal-like reptiles)** (class **\*Reptilia**) Subclass of reptiles which includes the pelycosaurs (**\*Pelycosauria**) and therapsids (**\*Therapsida**). The pelycosaurs appeared in the Upper **\*Carboniferous** and disappeared in mid-**\*Permian** times, displaced by the therapsids to which they had given rise. The therapsids flourished in the latter part of the Permian and in the **\*Triassic**, but dwindled to **\*extinction** in the early **\*Jurassic**. The therapsids are the ancestors of the **\*mammals**, and share in common with them a synapsid skull (i.e. a skull with one temporal opening).



### Synapsida

**synclinal ridge** An elongated hill underlain by a **\*syncline** whose axis trends parallel with it. Its upstanding nature may result from the relative strength of a compressed downfold compared with the tension-induced weakness of adjacent **\*anticlines**, but other explanations have been proposed. *See also* **INVERTED RELIEF**.

**syncline** A basin or trough-shaped **\*fold** whose upper component **\*strata** are younger than those below.

**synclinorium** A regional synformal structure (*see* **SYNFOLD**) which consists of a series of smaller, higher-order **\*anticlines** and **\*synclines**, some of which may be small enough to be viewed in **\*outcrop**.

**syneresis cracks** See DESICCATION CRACKS.

**synform** A basin or trough-shaped **\*fold** whose younger **\*strata** may be above or below older ones.

**syngenetic ore** An **\*ore** deposit formed simultaneously with the host rock and by similar processes, e.g. bedded **\*ironstone**, **\*magmatic-segregation deposits**.

**synkinematic** See SYNOROGENIC.

**synneusis** The drifting together and mutual attachment of crystals suspended in a **\*magma**. **\*Phenocrysts** may cluster together in this manner to form **\*glomeroporphyritic** aggregates.

**synodic month** The average time between successive new **\*Moons**, which is equal to 29 days, 12 hours, and 44 minutes. This period is curiously correlated with the average length of the human female menstrual cycle (29.5 days), while the average length of human pregnancy (266 days) equals nine synodic months (265.8 days).

**synoptic meteorology** The presentation of the current weather elements of an extensive area at a particular time. Sea-level and upper-level synoptic charts display weather conditions by symbols at selected synoptic stations. 'Synoptic' is from the Greek *sunoptikos*, meaning 'seen together'.

**synorogenic (synkinematic, syntectonic)** Applied to a process or event (e.g. **\*recrystallization** of **\*metamorphic rock**, or the emplacement of **\*plutons**) which occurs at the same time as deformation. Synorogenic **\*sediments** need not show contemporaneous deformation, though the sedimentation will be the result of deformation and uplift in the **\*orogenic belt**. Many authors use 'synkinematic', 'syntectonic', and 'synorogenic' interchangeably; others restrict 'synkinematic' to rocks showing evidence of deformation which happened simultaneously with their formation (e.g. rotated **\*porphyroblasts** in a metamorphic rock). 'Synorogenic' is often restricted to events or processes which happen simultaneously with a major phase in **\*orogeny**.

**synrhabdosome** An association of graptolite (see GRAPTOLITHINA) rhabdosomes (colonies) that are radially arranged. They appear to be life associations.

**synroc** A synthesized substance that has been proposed as a medium for the long-term disposal of *\*radioactive waste*. The waste would be mixed with the synroc ingredients and the mixture would set hard.



<http://www.world-nuclear.org/information-library/nuclear-fuel-cycle/nuclear-wastes/synroc.aspx>

- Synroc Wasteform.

**synrock** An artificially produced substance, based on petroleum, and used for protection, *\*grouting*, etc.

**synsacrum** In birds, a structure in the pelvic girdle formed by the fusion of certain vertebrae.

**syndimentary fault** A *\*fault* that forms by *\*brittle behaviour* during the sedimentary process.

**syndimentary fold** A *\*fold* that forms by *\*ductile behaviour* during the sedimentary process.

**syntaxial growth** A mineral *\*cement* which has grown around a *\*grain* in such a way that the *\*crystal* overgrowth is in *\*optical continuity* with the grain.

**syntectonic** See SYNOROGENIC.

**syntexis** A general term applied to the processes of *\*magma*-rock reactions. *\*Anatexis* and *\*assimilation* can be regarded as end-products of the various process of syntexis. These might include fracturing of *\*country* rock during intrusion, engulfing of country-rock *\*xenoliths* in the magma, and thermal and/or fluid infiltration of xenoliths to initiate melting or solid-state reaction.

**synthem** A major *\*stratigraphic unit* that is *\*unconformity*-bounded.

**synthetic-aperture radar** A *\*real-aperture radar* system used for high altitude and satellite *\*remote sensing*. A long antenna is stimulated resulting in a high resolution in the *\*azimuth* direction by using a *\*Doppler shift* effect to identify *\*backscattered* waves returning from ahead and behind the platform.

**synthetic fault** A *\*fault* whose sense of displacement is the same as that of the main zone of faulting when seen in vertical section. In an extensional regime (see *EXTENSION*), a synthetic fault mimics the displacement of a *\*listric* fault by forming in the active *\*hanging wall*.

**synthetic seismogram** A *\*seismic record* which can be modelled for a layered structure where the individual *\*formation velocities* and thicknesses are known (e.g. from *\*borehole logs*). It can then be compared with an observed *\*seismogram* to test geologic interpretations and geophysical models.

**synthetic thrust** A *\*thrust* whose sense of displacement is in the direction of the major thrust, but which may occur progressively in the *\*footwall* or *\*hanging wall*. A *\*back thrust* in this sense is an antithetic thrust. See *ANTITHETIC FAULT*.

**syntype** All specimens in a type series in which no *\*holotype* was designated.

***Syringopora fischeri*** One of the last representatives of the stony, colonial corals belonging to the subclass *\*Tabulata*; they flourished during the Upper *\*Palaeozoic* and are known from the Upper *\*Permian* of Germany. At the end of the Permian both tabulate and *\*rugose* corals became extinct.

**system** 1. The chronostratigraphic equivalent of the time unit *\*period*. Systems are subdivided into *\*series*, and together several systems constitute an *\*erathem*. When used formally the initial letter of the term is often capitalized, e.g. the *\*Devonian* System. 2. In *\*geomorphology*, a natural arrangement of interrelated objects or variables, the whole possessing properties that make it greater than the sum of the individual parts. It normally possesses stability, expressed by a balance between the input and output of energy and matter. This equilibrium may be upset by internal or external change. If the change is modest, the system quickly regains equilibrium; if it is extreme, a new equilibrium is established. A hillslope, for example, receives *\*precipitation* and exports water, slope debris, and the products of *\*weathering*. The form of the profile represents a balance between input and output. A landslide, perhaps induced by an increase in precipitation, would destroy this equilibrium, and in due course a new balance would be established. Several varieties of system are recognized.

See [CASCADING SYSTEM](#); [CONTROL SYSTEM](#); [MORPHOLOGICAL SYSTEM](#); [PROCESS-RESPONSE SYSTEM](#).

**systematic errors** See [ERRORS](#).

**systematic sampling** See [SAMPLING METHODS](#).

**Système Probatoire d'Observation de la Terre (SPOT)** A series of French observation satellites, the first launched in 1986 by the Centre National d'Études Spatiales (CNES) in collaboration with the Belgian scientific, technical, and cultural services and the Swedish National Space Board (SSTC and SNSB, respectively), which transmits data yielding monochrome, stereoscopic images with a 10 m resolution, superior to those from *\*Landsat*. SPOT 4, launched on 24 March 1998, was in a Sun-synchronous orbit at a height of 822 km; it completed each orbit in 101 minutes and crossed the same surface area every 26 days; the mission ended on 29 June 2013. SPOT 5, launched on 4 May 2002, is in a Sun-synchronous circular orbit identical to those of earlier SPOT satellites at an altitude of 832 km. SPOT-6 was launched on 9 September 2012 and SPOT-7 on 30 June 2014 into Sun-synchronous orbits at 694 km and 660 km, respectively.



<https://earth.esa.int/web/guest/missions/3rd-party-missions/current-missions/spot>

- SPOT.

**systems tract** In *\*sequence* stratigraphy, a three-dimensional depositional unit, defined by its boundaries and internal geometry. Three systems tracts commonly occur within a single cycle of sea-level changes: the *\*highstand* (or highland) systems tract, *\*lowstand (or lowland) systems tract*, and, between them, the *\*transgressive systems tract*. See also [REGRESSIVE SYSTEMS TRACT](#).

**SZ** See [SHENZHOU](#).

# T

**T** See TESLA.

*T* See TAYLOR NUMBER.

**Tabianian** See ZANCLEAN.

**tabula** (*pl.* **tabulae**) A horizontal **\*plate** extending across the interior of a **\*corallite**. See RUGOSA; TABULATA.

**tabulae** See TABULA.

**tabular** Applied to the form of a mineral when it occurs in broad, flat, often rectangular surfaces resembling the top of a table. The form may be due to the presence of a prominent **\*cleavage** direction, as in **\*barite**.

**tabular cross-stratification** See CROSS-STRATIFICATION.

**Tabulata** (subclass **\*Zoantharia**) A **\*Palaeozoic** order of zoantharians that are always colonial, never solitary, and that possess small **\*corallites**. Tabulae (see TABULA) are the major skeletal elements and septa (see SEPTUM) are usually absent. The **\*corallum** is built of individual corallites, not always connected to each other, and may be encrusting, may consist of thin sheets, or may be massive. Other forms may be joined side by side to form fence-like structures. Smaller tabulates appear to have lived in deeper waters, whereas the larger tabulates appear in coral–stromatoporoid (see STROMATOPOROIDEA) associations.

**tabulation** The arrangement of the **\*plates** in the walls of a dinoflagellate (see DINOPHYCEAE) **\*test** that is armoured with a series of plates.

**tachylite (tachylyte)** A black volcanic *\*glass* formed by the chilling of basaltic *\*magma*. The black colour is due to the presence of numerous microscopic crystals within the glass. Tachylite can be found in the chilled rind of *\*pillow lavas*, but is extremely susceptible to alteration by sea water and may be converted to *\*palagonite*.

**tachylyte** See TACHYLITE.

**tachytely** A rate of evolution within a group that is much faster than the average (horotelic) rate. Such accelerated evolution typically occurs when an organism enters a new *\*adaptive zone* and initiates an *\*adaptive radiation* to fill the available niches.

**Taconic orogeny** A middle to late *\*Ordovician* episode of mountain building, named after the Taconic Mountains (a N–S component of the Appalachian system to the east of the Hudson River, New York); it affected the whole of the Appalachians, in an area from what are now New England to eastern Canada, and is known locally in Newfoundland as the Humberian. In the south it was caused by the *\*subduction* of the western margin of the *\*Iapetus Ocean*, leading to the closing of the western basin of the Iapetus and the collision between N. America and the Piedmont microcontinent and the subsidence of the *\*continental shelf* further north. See APPALACHIAN OROGENIC BELT.

**taconite** The name given in the Lake Superior region of N. America to *\*banded iron formations*.

**tadpole plot** A graph of *\*dipmeter* results. The bedding *\*dip* is plotted as a dot on a depth–dip graph, and the dot is given a tail corresponding to the direction of the dip at that point (e.g. the tail points vertically upwards from the dot for a northerly dip direction).

**Tae Weian** See KUNGURIAN.

**tafoni** A hollow, produced by localized *\*weathering* on a steep face. Rock breakdown typically takes place by granular disintegration or by flaking, and the hollow shows a tendency to grow upwards and backward.

**Taghanician** See GIVETIAN.

**tagma** (*pl.* **tagmata**) In arthropods, one of the sections into which the body is divided by differences in size, shape, or function.

**tagmosis** In metamerically segmented animals (see **METAMERIC SEGMENTATION**), functional specialization that leads to differentiation among the segments and the formation of tagmata (see **TAGMA**).

**tailings** Fine waste material from a mineral-processing plant, that is too poor for further treatment. It is often stored in a **\*tailings dam**.

**tailings dam** A dam made from material having sufficient **\*permeability** to allow moisture to drain through over a regulated period. The dam is constructed to retain the water-sodden, fine-grained materials (**\*tailings**) which represent the waste product from a mineral-processing plant.

**Taitai** A **\*series** in the Lower **\*Cretaceous** of New Zealand, underlain by the **\*Oteke** and overlain by the **\*Clarence**. It comprises the Mokoiwian and Korangan, which are roughly contemporaneous with the **\*Neocomian** series and the subsequent **\*Barremian**, **\*Aptian**, and lower **\*Albian** **\*stages**.

**talc** A member of the 2:1 **\*phyllosilicates** (sheet silicates) with composition  $Mg_6[Si_4O_{10}]_2(OH)_4$ ; sp. gr. 2.58–2.83; **\*hardness** 1 (it has the lowest hardness on **\*Mohs's scale of hardness**); **\*monoclinic**; rare crystals are **\*tabular**, often **\*massive**; white to green; **\*cleavage** perfect {001}; massive talc (soapstone or steatite) can be formed during the low-grade **\*metamorphism** of siliceous **\*dolomites**; and as a **\*secondary mineral** during **\*hydrothermal** alteration of **\*ultrabasic** **\*igneous** rocks along shear planes. It is associated with serpentinization with **\*serpentine** changing to talc and **\*magnetite** by addition of CO<sub>2</sub>. It is used extensively as a mineral filler.

**talc schist** A **\*regional metamorphic** rock composed predominantly of **\*talc**, and displaying a **\*schistosity**. The rock forms by the **\*metamorphism** and deformation of **\*ultrabasic** **\*igneous** rocks in regional **\*terranes**.

**taleolae** See **PUNCTATE**.

**talik** See **PERMAFROST**.

**tail cloud** See **CLOUD CLASSIFICATION**.

**talus (scree slope)** A sloping mass of coarse rock fragments accumulated at the foot of a cliff or slope.

**taluvium** A hillslope deposit that consists of a mixture of coarse, rocky rubble and finer particles. The term is a hybrid, derived from ‘\*talus’ and ‘colluvium’ (relatively fine debris that has been moved downhill by slope wash).

**Tame Valley** An \*interglacial in the \*Upton Warren Interstadial, within the Würm (\*Weichselian) glacial.

**Tandem-L** A radar-satellite mission by the German Space Centre (DLR) to monitor dynamic Earth-surface processes with high resolution, including the global measurement of forest biomass, using two \*synthetic-aperture radar instruments. The mission is scheduled for launch in late 2022.



<https://www.dlr.de/dlr/en/Portaldata/1/Resources/documents/2016/Tandem-L-Brochure-2016-06.pdf>

- Tandem-L brochure.

**tangential longitudinal strain** In a \*fold, a buckling (see BUCKLE FOLDING) \*strain distribution in which strain is internal (in contrast to layer-parallel strain), and concentrated in the \*hinge zone, leaving the limbs relatively unstrained. The \*principal strain axes are layer-parallel and there is a neutral surface of zero strain throughout the fold profile.

**TanSat** A Chinese \*minisatellite mission to detect and monitor atmospheric CO<sub>2</sub> (*tan* means carbon in Mandarin). The satellite was launched on 21 December 2016, from Jiuquan Satellite Launch Centre, China, into a Sun-synchronous orbit at an altitude of about 700 km.

**tantalite** See COLUMBITE.

**taphichnia** \*Trace fossils made by animals attempting unsuccessfully to escape.

**taphofacies (taphonomic facies)** A sedimentary rock unit, or association of units, characterized by the combination of preservational features of the fossils contained within it. Compare BIOFACIES.

**taphonomic facies** See TAPHOFACIES.

**taphonomic grade** The degree of preservation with **\*taphofacies**, from which the depositional environment can be inferred.

**taphonomy** The study of the transition of all or part of an organism and its traces from the biosphere into the lithosphere (i.e. **\*fossilization**). The term was coined by J. A. Efremov in 1940.

**taphrogenesis** The formation of **\*rift valleys** and **\*grabens** by vertical movements of the **\*crust**.

**TARANIS** See TOOL FOR THE ANALYSIS OF RADIATIONS FROM LIGHTNINGS AND SPRITES.

**tarsal bone** In **\*tetrapod** vertebrates, one of the distal bones of the hind foot that articulate with the digits (metatarsals).

**tar sand** Oil reservoir where the **\*volatiles** have escaped and the rock has become impregnated with **\*hydrocarbon** residue. Sometimes this occurs in commercial quantities, as at Athabasca, Alberta, Canada; the sands are mined **\*open-cast** and treated with steam and hot water to release the oil.

**tarsometatarsus** A bone formed by the fusion of the **\*metatarsals** with the **\*tarsals** of the appendicular skeleton, and found in birds (**\*Aves**) and some dinosaurs. The beginnings of this fusion can be seen in some **\*Theropoda**, but it is not complete until the early **\*Cretaceous** (**\*Sinornis**). See also TIBIOTARSUS.

**Taupo** See ULTRAPLINIAN ERUPTION.

**taxa** See TAXON.

**taxodont** Applied to a primitive type of **\*hinge \*dentition** present in certain bivalves (**\*Bivalvia**), in which teeth and sockets are small, numerous, and arranged in a row on each side of the **\*shell beak**, on both **\*valves**.

**taxon** (*pl.* **taxa**) Group of organisms of any taxonomic rank, e.g. family, genus, or **\*species**. See CLASSIFICATION.

**taxonomy** (*adj.* **taxonomic, taxonomical**) The formal **\*classification** of organisms, soils, or any other entities, based on degrees of relatedness among the items being considered.

**taxon range zone** (**genus zone**, **species zone**, **total range zone**) The total body of *\*strata* characterized by the presence of a specified *\*taxon*, both in geographical extent and time-stratigraphic range. *See also* RANGE ZONE.

**Taylor, Frank Bursley** (1860–1939) Taylor was a respected American glaciologist, but is better known for his more speculative cosmology, which included the idea that the *\*Moon* was a former comet, captured in the *\*Cretaceous*. He published his ideas on *\*continental drift* three years before *\*Wegener*, but used little evidence to support his theory, and was ignored.

**Taylor number** (*T*) A dimensionless number measuring the influence of rotation on a convecting system. It depends on the scale of the *\*convective cell*, the rate of rotation, and kinematic *\*viscosity*. If *T* is equal to or greater than 1, then rotational effects are significant. *See* CONVECTION.

**TCR** *See* TOTAL CORE RECOVERY.

*t–d* *See* TIME–DISTANCE CURVE.

**T–d curve** A temperature–depth curve.

*T<sub>e</sub>* *See* EFFECTIVE TEMPERATURE.

**tear fault** *See* STRIKE-SLIP FAULT.

**technosols** Soils that occur wherever human activities have sealed the surface, extracted material, or constructed artificial soil. They contain 20% or more of artefacts, or are sealed by hard rock within 5 cm of the surface over at least 95% of the area. Technosols include landfills, sludge, cinders, mine spoils, etc. They are a soil group in the *\*World Reference Base for Soil Resources*.

**tectofacies** A *\*lithofacies* formed by *\*tectonism*.

**tectonic** *See* TECTONISM.

**tectonism** (*adj.* **tectonic**) Deformation within the Earth's *\*crust*, and its consequent structural effects.

**tectonite** A deformed rock with a mineral *\*fabric* which forms a pervasive *\*foliation* or *\*lineation*, or some combination of both. *See* SHAPE FABRIC.

**tectosilicates** *See* SILICATES.

**teeth** 1. See DENTITION. 2. See BONE. 3. See BIVALVIA. 4. See MAMMALIA.

**tegeminal plates** See TEGMEN.

**tegmen** The \*ventral surface of the \*crinoid body between the arms. It is covered with a non-calcareous integument in which small, calcareous plates (tegeminal plates) are usually set.

**Teichichnus** A so-called ‘wall-like’ \*trace fossil in which the horizontal, tube-like trace is underlain by upwardly curving laminae (\*spreiten). It is recorded from the \*Carboniferous of Scotland. The concave, upwardly directed spreiten may be used as ‘way-up’ or \*geopetal structures.

**teilchron** The time unit defined by the local range of a \*species, corresponding to its \*teilzone.

**teilzone (local range zone, topozone)** \*Strata containing part of the total stratigraphic range of a \*fossil \*taxon, occurring in a particular area.

**tektites** Small fragments (usually 2.5–5.0 cm) of \*silica-rich, translucent black \*glass, found scattered over large areas (\*strewn-fields) in particular regions of the Earth. Most tektites exhibit ‘splash’ shapes (tear-drops, dumb-bells, etc.) indicating a rapid cooling and solidification during flight. They are thought to have formed as ejecta from cometary or \*meteorite impacts on to silicate-rich rocks, most likely terrestrial, although this is still a matter of some controversy. The material would have melted on impact, been thrown up into the atmosphere or space, and landed, resolidified, far from the site of origin. The ages of known tektites range from around 0.7 to 35.0 Ma. Microscopic tektites (microtektites) have been recorded from ocean \*sediments off the Ivory Coast, off southern Australia, and from the Indian Ocean.

**teleconnections** Linked atmospheric changes that occur in widely separated parts of the world. For example, the \*southern oscillation links changes in surface atmospheric pressure measured at Darwin, Australia, and Tahiti, in the central South Pacific Ocean; \*El Niño brings dry weather to parts of Brazil, south-western Europe, north-eastern China, and Australia, and wet weather to much of the United States, Israel, and north-western Europe.

**TeLEOS-1** A commercial Earth-observation \*minisatellite mission by ST Electronics, Singapore, to supply up to 1-m-resolution imagery. It was

launched on 16 December 2015, from Satish Dhawan Space Centre, India, into a near-equatorial orbit at an altitude of 550 km.

**Teleostei (teleost, teleostean)** Somewhat loosely defined term (an infraclass or superorder according to some authors) that includes all the living bony fish (*\*Osteichthyes*) with the exception of a few orders of primitive fish. Teleosts arose from holostean (*\*Holostei*) stock in the *\*Jurassic*. A transitional form, *Leptolepis*, from Upper Jurassic marine deposits, was a herring-shaped fish, around 23 cm long, with a *\*homocercal tail* and pelvic fins placed well back on the body. The scales still carried traces of enamel. Teleosts diversified in the *\*Cretaceous* and are now the most abundant *\*vertebrate* group. They exhibit a great variety of form, but are characterized by an internal skeleton entirely of *\*bone*, a reduction in thickness of the scales, and a homocercal tail; typically there is a dorsal swim bladder to control buoyancy, a mobile jaw articulation, and the fin rays may be stiffened into spines.

**teleseism** An *\*earthquake* whose *\*epicentre* is more than 1000 km from, but detectable at, a recording *\*seismometer*. Earthquakes whose epicentres are closer than this are regarded as 'local' events.

**Telesto (Saturn XII)** One of the lesser satellites of *\*Saturn*, discovered in 1980 by *\*Voyager* 1, with a radius measuring 15 × 12.5 × 7.5 km; visual albedo 0.5.

**telethermal** Applied to an *\*ore* deposit far from a source of *\*magma*. It has been produced by *\*hydrothermal activity* at low temperatures and shallow depth with little *\*wall-rock alteration*.

**Television and Infrared Observation Satellite (TIROS)** The first meteorological satellite, launched by the USA on 1 April 1960, and operated by the National Oceanic and Atmospheric Administration (*NOAA*). It monitored strips of territory, missing the poles. Two successors, both called TIROS-N, were launched later. TIROS satellites are now known as NOAA-class satellites.



<http://science.nasa.gov/missions/tiros/>

- A NASA mission launched in 1960 to determine if satellites could be useful in the study of the Earth.

**telinite** See COAL MACERAL.

**telluric anomaly** A perturbation in the flow of a **\*telluric current**. In what is assumed to be an **\*isotropic** homogeneous medium at or near the Earth's surface, telluric current flows uniformly and there are no discernible potential gradients. The presence of a subsurface geologic structure (e.g. a **\*salt dome**) perturbs the line of flow of the telluric current, forming a potential gradient which can be measured (the telluric anomaly) using **\*non-polarizable electrodes** spaced several hundred metres apart.

**telluric current** A naturally occurring electrical current which flows at or near the Earth's surface over very large areas, with a magnitude of about 10 mV/km. At any point from which it is observed it constantly changes direction and magnitude.

**telome theory** A hypothesis suggesting that the fern leaf arose from various rearrangements of branching stem systems.

**telson** The spike-like, terminal segment of the opisthosoma that occurs in certain chelicerates (**\*Chelicerata**), e.g. limulids and eurypterids. The prominent, backward-pointing spine at or near the rear extremity of a trilobite (**\*Trilobita**) is also called a telson, although it may or may not be the equivalent of a true telson.

**Telychian (Ontarian)** A **\*stage** (436–428.2 Ma ago) of the Early **\*Silurian**, underlain by the **\*Aeronian** and overlain by the **\*Sheinwoodian**.

**TEM** See TRANSIENT ELECTROMAGNETIC METHOD.

**Temaikan** A New Zealand **\*stage** (169–161.2 Ma ago) of the Middle **\*Jurassic \*epoch**, preceded by the **\*Uroan** and followed by the Late Jurassic.

**Temispack** A widely used technique for **\*basin modelling**, developed by the Institut Français du Pétrole.

**Tempel–Tuttle** See TEMPLE–TUTTLE.

**temperate climate** General term applied to the characteristics of mid-latitude climates influenced from time to time by both tropical and polar **\*air masses**. Temperature criteria provide subdivisions into warm, cool, or

cold-temperate climates. *See also* CLIMATE CLASSIFICATION; KÖPPEN CLIMATE CLASSIFICATION; STRAHLER CLIMATE CLASSIFICATION.

**temperate glacier** *See* GLACIER.

**temperate zone** In the climate classification devised by Aristotle, the region in either hemisphere between the Arctic or Antarctic Circles and the Tropics of Cancer or Capricorn. In his view, this was the only region habitable for humans. *Compare* FRIGID ZONE; TORRID ZONE.

**temperature–composition diagram** Graphic representation, usually with temperature on the ‘y’ axis, and composition of a *\*phase* or phases (e.g. an *\*isomorphous* series) on the ‘x’ axis, which shows temperature-dependent variations in composition of the solid phase and the associated *\*melt*.

**temperature distribution with depth** *See* GEOTHERMAL GRADIENT; HEAT FLOW.

**temperature inversion** An atmospheric condition in which the typical *\*lapse rate* is reversed and temperature increases vertically through a given layer. In the *\*troposphere* an inversion layer marks conditions of great stability, i.e. a region in which vertical motion is strongly damped, with an absence of turbulence. An inversion acts as a ceiling, preventing further upward convection, and is generally the limit for cloud development. Marked and persistent inversions occur at lower levels, with subsiding air in major anticyclonic cells, such as the *\*Azores high*-pressure zone and cold *\*anticyclones* over continents. *See also* ATMOSPHERIC STRUCTURE; ENVIRONMENTAL LAPSE RATE.

**temperature log** A record of the temperature measured down a *\*borehole*.

**temperature range** Extent of diurnal or seasonal (annual) temperature variation. Generally the highest annual range of temperature is experienced in higher latitudes, especially over 65° N, which reflects the continental influences of N. America and Asia. Tropical land masses tend to have the highest diurnal temperature ranges, which in the equatorial zone are much higher than the annual variations. Annual temperature ranges in general are particularly moderated by maritime influences.

**TEMPEST-D** See TEMPORAL EXPERIMENT FOR STORMS AND TROPICAL SYSTEMS TECHNOLOGY – DEMONSTRATION.

**tempestite (storm deposit)** Material deposited during a single storm, often on a *\*continental shelf* where the *\*tidal range* is small and prevailing winds are strong.

**Templetonian** A *\*stage* (510–508 Ma ago) of the Middle *\*Cambrian* of Australia, underlain by the *\*Ordian* and followed by the *\*Florion*.

**Temple–Tuttle (Tempel–Tuttle)** A *\*comet* with an orbital period of 32.92 years; *\*perihelion* date 27 February 1998; perihelion distance 0.982 AU.

**Temporal Experiment for Storms and Tropical Systems Technology – Demonstration (TEMPEST-D)** A 1-unit *\*CubeSat* mission by Colorado State University to sample convective precipitation events, including cloud formation, ice formation, and cloud dissipation. It was launched on 21 May 2018, from Virginia, into a near-circular orbit at an altitude of about 400 km.

**tenacity** The physical response of a mineral to stress, e.g. vibration, crushing, or bending. Soft minerals which are easily flattened, e.g. *\*copper* and *\*gold*, are described as ‘malleable’. Minerals such as *\*pyrite* and *\*fluorite* crumble easily and are said to be ‘brittle’. Minerals such as *\*mica* are ‘flexible’, meaning they can be bent but will not return to their original shape.

**tennantite** See TETRAHEDRITE.

**tensile strength** The strength of a substance (e.g. rock) under tension. It can be measured by loading cylindrical samples under tension until failure occurs, or by compressing sample discs along a diameter to induce tension failure on a diametrical surface. It can be measured in a *\*point load tester*.

**tensile stress** A *\*normal stress* (negative compressive stress) which pulls apart the material on either side of a plane. Tensile stress greatly weakens rocks, reducing the amount of *\*shear stress* that is needed to produce failure in them.

**tensiometer** A device used to measure the soil-moisture tension in the unsaturated (*\*vadose*) zone. The pressure is negative and cannot be

measured using a conventional open *\*piezometer*. The tensiometer consists of a closed tube with a porous base and a *\*manometer*.

**tension crack** See TENSION FRACTURE.


**tension fracture (tension crack, tension gash)** A discrete, commonly lens-shaped, rock fracture which forms and propagates perpendicularly to the direction of maximum *\*extension*. Tension fractures often occur in *\*en échelon* arrangements.

**tension gash** See TENSION FRACTURE.

**tent rocks (wigwams)** Conical pillars of white rock, often tens of metres tall, formed by the erosion of *\*pyroclastic* rocks, either by the isolation of blocks by intersecting drainage channels, or (more commonly) where resistant blocks of lava or other rocks protect the material beneath them as surrounding rocks are eroded.

**tepee** Fold-like structures developed in *\*calcrete* *\*soil profiles*, tidal areas, and the margins of salt lakes, due to the fluctuations in water levels and changes in chemical *\*precipitation*. Tepee structures are characterized by mineral horizons folded with pointed terminations to the anticlinal closure, resembling the form of a tepee tent.

**tephigram (T $\Phi$ gram)** Diagram showing the vertical variation in properties of the atmosphere, i.e. temperature, potential temperature, and humidity are plotted as a function of pressure, with lines indicating *\*dry* and *\*saturated adiabatic lapse rates*. The changes in temperature and humidity of lifted *\*air* parcels can be compared against the environment (surrounding air) curve, revealing the *\*stability* or *\*instability* of the *\*air mass* and the level at which condensation will occur in the uplifted air. Potential temperature is symbolized by the Greek letter phi ( $\Phi$ ), so the diagram is a T $\Phi$ gram. See also ENVIRONMENTAL LAPSE RATE.



**Tephigram:** Temperature and humidity are plotted against pressure to illustrate the properties of an entire column of air.

**tephra** Collective term applied to all *pyroclastic* particles or fragments ejected from a *volcano*, irrespective of size, shape, or composition. The term is usually applied to air-fall material, rather than pyroclastic flow deposits.


**tephrite** An *undersaturated*, fine-grained, *extrusive*, *igneous* rock consisting of *essential* calcium-*plagioclase*, *nepheline*, and *titanaugite*, with *accessory* *apatite* and iron oxide. The rock is nearly always *porphyritic*. See TRACHYBASALT.

**tephrochronology** A dating method based on the examination of *tephra* (volcanic ejecta); in areas of repeated activity it is often possible to recognize distinctive events within a *pyroclastic* succession, and to use

such markers for local correlation (see **MARKER BED**). The succession, so established, provides useful data on the history of the **\*volcano** as well as a guide to magmatic and geochemical changes operating below ground. In many cases, well removed from the volcanic activity, thin **\*ash** falls may be found within sedimentary successions. Such thin **\*horizons** are regarded as **\*isochronous** and can be used in correlation. The ash is usually altered to form bentonitic clays (see **BENTONITE**), and because these clays still contain traces of volcanic material they can be dated by **\*radiometric** means. This is especially important where the intervening **\*strata** contain **\*fossil** remains (e.g. of early humans in Africa) which can be calibrated by using the dates of the ash falls. Good examples of this application in regional **\*stratigraphy** occur in the Lower **\*Palaeozoic** strata of southern Scandinavia and the mid-**\*Cretaceous** **\*sediments** of the western interior of the USA.

**Teratan** A new Zealand **\*stage** (87–83.7 Ma ago) of the Late **\*Cretaceous** **\*epoch**, preceded by the **\*Mangaotanean** and followed by the **\*Campanian**.

**Terebratulida (terebratulids)** (class **\*Articulata**) Order of **\*Brachiopoda** with **\*punctate** shells, rounded **\*hinge lines**, functional **\*pedicle**, **\*deltoidal plates**, and a lophophore support usually consisting of a pair of **\*crura** and a calcareous loop. The three suborders comprising the order first appeared in the Lower **\*Devonian**; the Centronellidina became extinct at the end of the **\*Permian**, the Terebratulidina and the Terebratellidina are still extant.



**Terebratulida**

**terebratulids** See TEREBRATULIDA.

**terminal node** In a \*phylogenetic tree, the point at the end of a branch representing a progenic taxon.

**terminator** The line on the visible disc of a planet or \*satellite which separates the sunlit and shaded regions, marking sunrise and sunset. Because the Sun angle is very low near the terminator, small details (e.g. lava-flow fronts on the Moon) become clearly visible.

**ternary system** A mineral system in which there are three components, e.g. \*diopside (Di)–\*anorthite (An)–\*albite (Ab). See also COTECTIC CURVE; PHASE; PHASE DIAGRAM.

**Terra** A NASA satellite, 6.8 m long and 3.5 m wide, that was launched on 18 December 1999 into a \*Sun-synchronous orbit at a height of about 700 km, each orbit taking 99 minutes. It became fully operational on 24 February 2000. Terra carried five sensors: advanced spaceborne thermal emission and reflection radiometer (ASTER), clouds and Earth's radiant energy system (CERES), multi-angle imaging spectroradiometer (MISR), moderate-resolution imaging spectroradiometer (MODIS), and measurements of pollution in the troposphere (MOPITT). These collect data on the interactions between the atmosphere, land, oceans, and radiant energy. It was still functioning in November 2018.



[http://www.nasa.gov/mission\\_pages/terra/index.html](http://www.nasa.gov/mission_pages/terra/index.html)

- A NASA mission launched in 1999 to collect data on climate change on Earth.

**terra** (*pl.* **terrae**) The term proposed by Galileo for the white, high-standing regions of the lunar surface, now generally referred to as the Lunar Highlands. These comprise the heavily cratered primary lunar feldspathic crust, forming 83% of the lunar surface. The term is also used for extensive (continent-sized) elevated areas on the surface of any extraterrestrial body (e.g. Ishtar Terra on \*Venus).

**terrace** A nearly flat portion of a landscape, terminated by a steep edge. It may be produced by any one of a range of processes, so the following

varieties are recognized: *\*altiplanation* terrace, *\*kame* terrace, *\*river* terrace, *\*shore platform*, and *\*solifluction* terrace.

**terracette (sheep-walk)** A small-scale land-form consisting of a long, narrow, stepped feature developed in *\*unconsolidated* material mantling a steep hillslope. It may result from slight slippage in material standing at an angle too great for stability.

**terrae** See [TERRA](#).

**terrain** (N. American **terrane**) An area of ground with a particular physical character; an area or region with a characteristic geology, e.g. 'metamorphic terrain'. Many British authors now use the 'terrane' spelling, except for 'terrain correction' (see [TOPOGRAPHIC CORRECTION](#)) and when used in the sense of such expressions as 'rough terrain', i.e. when not meaning 'displaced *\*microcontinent*'. Compare [TERRANE](#).

**terrain component** The influence of the type and features of an area surrounding a proposed mine or engineering site. See also [TERRAIN EVALUATION](#).

**terrain correction** See [TOPOGRAPHIC CORRECTION](#).

**terrain evaluation** The assessment of a *\*terrain* for its suitability as a mining or industrial site. Consideration must be given to whether it can provide: the space required for surface plant; the area needed for disposal, e.g. *\*tailings dams*; adequate areas for stock-piling; parking space for employees; in remote territories, access to an airfield; adequate roads, railways, or access to sea ports; room for future expansion; low fire risk (e.g. the possibility of clearing adjacent woodland to reduce the risk of forest fires); adequate facilities for staff and visitors, e.g. housing, canteens, recreation facilities; space for concentrating plant and a smelting and refining complex. Consideration must also be given to the effect of the development on the environment, e.g. air pollution.

**terrain pattern** The type of landscape surrounding a mine or industrial site. See also [TERRAIN EVALUATION](#).

**terrain unit** A sub-region of an area that is under consideration for use as a mine or industrial site, e.g. whether it has forest, exposed bedrock, *\*talus*,

bog, etc. *See also* TERRAIN EVALUATION.

**terrane** 1. A *\*fault*-bounded area or region which is characterized by a *\*stratigraphy*, structural style, and geologic history distinct from those of adjacent areas, and which is not related to those areas by unconformable contacts or *\*facies* changes. This usage has recently become frequent in studies of *\*orogenic belts* in which terranes can have origins as *\*accretionary wedges*, *\*island arcs*, and *\*microplates*. Initially, ‘displaced’ or ‘allochthonous’ was used to qualify the word ‘terrane’ and to distinguish such terranes from merely ‘*\*suspect terranes*’, but all three qualifiers are falling into disuse. 2. N. American spelling of *\*terrain*, increasingly coming into general usage.

**terra rossa** European soils, red in colour, that developed on iron-oxide-rich residual material over *\*limestone*. They are deep and ancient, some of them being pre-*\*Pleistocene*. They are now classified as *\*inceptisols* or *\*mollisols* in the *\*USDA \*soil taxonomy*.

**terrestrial planet (inner planet)** The term used to classify the four inner, rocky Earth-like planets (*\*Mercury*, *\*Venus*, *\*Earth*, and *\*Mars*) of the *\*solar system*, which, relative to the *\*jovian* (or outer) planets are smaller, have high densities, are composed of metallic (mainly iron) and silicate phases, and have low contents of gaseous elements.

**terrestrial radiation** Long-wave electromagnetic radiation (wavelengths 4–100  $\mu\text{m}$ , with a peak at 10  $\mu\text{m}$ ) from the Earth’s surface and atmosphere. *See also* ATMOSPHERIC ‘WINDOW’; GREENHOUSE EFFECT; RADIATION BUDGET; RADIATION NIGHT.

**terrific horizon** In the *\*World Reference Base for Soil Resources*, a *\*soil horizon* that has been strongly affected by the incorporation of manure.

**terrigenous** The description of a siliciclastic *\*sediment* which has been deposited, or formed, on land.

**Tertiary** A name, now obsolete, for the first sub-era of the *\*Cenozoic \*era*. The Tertiary followed the *\*Mesozoic* and comprised two periods, *\*Palaeocene* and *\*Neogene*. It began 65.5 Ma ago and was followed by the *\*Quaternary*. The name ‘Tertiary’ has been abandoned and been replaced by the *\*Palaeogene* and *\*Neogene periods*.

**tertiary creep** In a material undergoing long-term low stress, the final stage of *\*creep*, characterized by accelerated permanent viscous *\*strain*, leading eventually to failure.

**teschenite** An *\*undersaturated*, medium- to coarse-grained, *\*igneous* rock consisting of *\*essential* calcium-*\*plagioclase*, *\*analcime*, *\*titanaugite*, and *\*barkevikite*, with *\*accessory* iron oxide. The rock is similar to essexite but with analcime taking the place of *\*nepheline*. The teschenite of the Lugar sill, in Ayrshire, Scotland, is a well-known British example.

**tesla (T)** A unit of magnetic field strength.  $1 \text{ T} = 1 \text{ Wb/m}^2 = 10^{-4} \text{ gauss}$ .

**tessera** (*pl.* *tesserae*) On the surface of *\*Venus*, an area of polygonal terrain.

**tesserae** See TESSERA.

**test** Protective shell that covers the cells of some *\*protists* and the soft parts of certain invertebrate animals.

**testate** Having a shell or *\*test*.

**Tethyan realm** The faunal province (*see* FAUNAL REALM) based in the region of the *\*Tethys Sea*. Characteristically *\*Jurassic* to *\*Cretaceous* in age, the Tethyan Realm denotes a warm-water, tropical to subtropical fauna and flora. The name may also be applied to warm-water faunas and floras of *\*Mesozoic* age outside the area of Tethys, especially when used in contrast to the *\*Boreal* (northern) and Austral (southern) realms.

**Tethys (Saturn III)** One of the major satellites of *\*Saturn*, with a radius of 529.9 km; mass  $6.22 \times 10^{20} \text{ kg}$ ; mean density  $1000 \text{ kg/m}^3$ ; visual albedo 0.9. It was discovered in 1684 by G. D. Cassini.


**Tethys Sea (Neotethys)** The sea that more or less separated the two great *\*Mesozoic* *\*supercontinents* of *\*Laurasia* (in the north) and *\*Gondwana* (in the south). That *\*land bridges* between the two supercontinents existed for much of the Mesozoic is attested to by the cosmopolitan character of *\*dinosaur* faunas. See PALAEOTETHYS.

**Tetracorallia** Alternative name for the order *\*Rugosa*.

**tetracorals** See RUGOSA.

**tetrad** See CRYSTAL SYMMETRY; SPORE.

**tetragonal** One of the seven \*crystal systems, with two sets of edges of the same length and a third which is either longer or shorter than the other two. The \*lattice may be referred to three \*crystallographic axes,  $a_1$ ,  $a_2$ , and  $c$  (or  $x$ ,  $y$ , and  $z$ ) where  $a_1$  and  $a_2$  (or  $x$  and  $y$ ) are equal, and  $c$  (or  $z$ ) may be longer or shorter. All three axes are at right angles.



**Tetragonal**

**tetrahedrite** An \*end-member of the tetrahedrite series from tetrahedrite  $(\text{Cu,Fe})_{12}\text{Sb}_4\text{S}_{13}$  to tennantite  $(\text{Cu,Fe})_{12}\text{As}_4\text{S}_{13}$ ; if silver is present in quantity (up to 30%) the mineral is called freibergite; sp. gr. 4.5–5.1; \*hardness 3.0–4.5; greyish-black; \*metallic \*lustre; black \*streak; normally \*massive but \*tetrahedral when crystalline; occurs in \*hydrothermal veins in association with other copper and iron sulphides.


**tetrahedron** A four-faced \*crystal form in which each face is an equilateral triangle. It may be referred to three equal \*crystallographic axes at right angles which join the centres of opposite edges of the crystal. It is a special form within the \*cubic system with the symbol (111).

**Tetrapoda** Vertebrate animals which have four limbs, including the *\*Amphibia*, *\*Reptilia*, *\*Aves*, and *\*Mammalia*.

**Teurian** The basal *\*stage* (65.5–56.5 Ma ago) in the *\*Palaeocene \*epoch* of New Zealand, preceded by the *\*Maastrichtian*, overlain by the *\*Waipawan*, and roughly contemporaneous with the *\*Danian* and lower *\*Thanetian* stages.

**textural maturity** An expression of the *\*sorting*, *\*matrix* content, and *\*grain* angularity in a *\*sediment*. An immature sediment is one with poor sorting, a large proportion of matrix, and angular particles. A texturally mature sediment is matrix poor or matrix free, well sorted, and with well-rounded grains.

**texture** 1. In *\*petrology*, the sizes and shapes of particles in rock and their mutual interrelationships. 2. In pedology, the proportions of *\*sand*, *\*silt*, and *\*clay* in the fine earth of a soil sample, which give a distinctive feel to the soil when handled, and which are defined by classes of soil texture.



## Texture

**Thailand Earth Observation System (THEOS)** An Earth-observation mission that supplies panchromatic and multispectral imagery to assist mapping, land use, agricultural monitoring, forestry management, coastal zone monitoring, and flood risk management. The satellite was launched on 1 October 2008, from Yasný Cosmodrome, Russia, into a Sun-synchronous near-circular orbit at an altitude of 822 km.

**Thalassa (Neptune IV)** A satellite of *\*Neptune*, with a diameter of 80 km; visual albedo 0.06.

**Thalassinoides** An *\*ichnoguild* of branched, horizontal to slightly inclined *\*burrows* or a burrow system, formed by suspension feeders just below water, at the *\*sediment* interface. They usually occur in dense associations. The dimensions of the burrows may reflect the environmental energy level.

*Thalassinoides* burrows are typical of \*clastics and \*detrital \*carbonates, and are abundant in certain \*post-Palaeozoic formations. Remains of Bryozoa have been found within them.

**thalweg** Line joining the lowest points of successive cross-sections, either along a river channel or, more generally, along the valley that it occupies.

**thanatocoenosis** See DEATH ASSEMBLAGE.

**Thanetian** A \*stage of the \*Palaeocene \*epoch, preceded by the Selandian, followed by the \*Ypresian, and dated at 58.7–55.8 Ma ago (Int. Commission on Stratigraphy, 2004). It is roughly contemporaneous with the upper \*Ynezian and \*Bulitian (California), upper \*Teurian and lower \*Waipawan (New Zealand), and the upper \*Wangerripian (Australia). It is named after the Thanet Sands in Kent, England.

**thaw** Onset of melting of ice and snow as the temperature rises above the freezing point.

**Thebe (Jupiter XIV)** The fourth of the known jovian satellites, discovered in 1979, Thebe has a diameter of 100 km ( $\pm 20$  km) (100  $\times$  90 km); mass  $7.77 \times 10^{17}$  kg; mean distance from \*Jupiter 222 000 km.

**theca** (*pl.* thecae) 1. Shell or \*test. 2. The cell wall in some protistan (*see* PROTIST) algae, e.g. \*diatoms and dinoflagellates (*see* DINOPHYCEAE). 3. The dense wall at the margin of a \*corallite in a scleractinian coral (\*Scleractinia). 4. In graptolites (\*Graptolithina), one of the cup-like structures that occur along the \*stipe, each of which houses an individual member of the colony. Thecae vary from simple, straight-sided tubular structures to those with an introverted opening, where the thecal aperture faces downward. There are triangulate thecae with very small apertures and in some species the thecae are long, thin, and separated from one another ('isolate'). In some \*monograptids the thecae vary in structure along the length of an individual stipe as well as from one species to another. Thecal shape also varies through time.

**Thecodontia (thecodonts)** 'Tooth-in-socket' reptiles, and the most primitive order of the \*Archosauria ('ruling reptiles'), ranging from the Upper \*Permian to the Upper \*Triassic. They were ancestral to the \*dinosaurs, \*pterosaurs, and crocodiles. One of the first thecodonts was *Thecodontosaurus browni* which flourished during the Early \*Triassic. It

grew to 2–3 m in length and had a small head and neck. Essentially it was a quadruped, but it could walk on its hind legs.

**thecodonts** See THECODONTIA.

**thematic map** In \*remote sensing, an image which has a \*classification overlaid on to it.

**thematic mapper** An improved \*multispectral scanner that acquires simultaneous images of a scene in different wavelength bands.

**THEMIS** See TIME HISTORY OF EVENTS AND MACROSCALE INTERACTIONS DURING SUBSTORMS.

**theoretical morphology** In evolutionary studies, analysis of the differences between all the forms an organism might have and those which have actually existed. By revealing unfilled regions of \*morphospace the technique may suggest constraints.

**THEOS** See THAILAND EARTH OBSERVATION SYSTEM.

**Therapsida** Order of \*synapsid \*reptiles, ancestral to the \*mammals, ranging from the latter part of the \*Permian to early \*Jurassic times.

**thermal** Used as a noun, the term refers to air that rises in a column over some localized heat source; it rises through the atmospheric environment by convection, as a result of its reduced density that causes cooler, denser air to sink beneath it. Intense heating over land (and particularly over some types of surface) results in \*adiabatic expansion and uplift. When thermals rise to \*condensation level, short-lived clouds (e.g. fair-weather \*cumulus) may form. Continued strong thermals in moist air can result in the growth of the cloud into large cumulus or \*cumulonimbus. When the upward growth reaches a stable layer, the cloud top may spread laterally.

**thermal cleaning** See DEMAGNETIZATION.

**thermal conductivity (K)** A parameter used as a measure of how well a material conducts heat (for example, for copper  $K = 385 \text{ W/m/K}$ , for air  $K = 0.02 \text{ W/m/K}$ ).  $K$  forms the constant of proportionality in the heat-conduction-rate equation:  $dQ/dt = -KA \, d\theta/dl$ , where  $dQ/dt$  is the heat conduction rate (in joules per second),  $A$  is the cross-sectional area of a

uniform sample (in square metres), and  $d\theta/dl$  is the temperature gradient along the sample (in kelvin per metre). *Compare* [THERMAL RESISTIVITY](#).

**thermal emission** The emission of [\\*electromagnetic radiation](#) due to the vibration of molecules as a result of their temperature.

**thermal equator** The zone of highest mean temperature over the Earth, either in the annual or long-term average or at a given moment. On the long-term average, it is located around 5°N latitude. This position north of the geographical equator results from the generally rather higher temperature of the northern hemisphere as compared with the southern hemisphere; this is because the glaciated Antarctic continent maintains colder summers in the southern hemisphere than does the Arctic, with a much smaller land area, in the northern hemisphere. *See also* [COLD POLE](#).

**thermal inertia** A measure of the responsiveness of a material to variations in temperature. In [\\*remote sensing](#) it is measured by diurnal changes in temperature. Materials with a high [\\*heat capacity](#) display high thermal inertia, consequently such materials will show small changes in temperature through the diurnal cycle.

**thermal infrared** [\\*Infrared radiation](#) which has a wavelength between 3.0  $\mu\text{m}$  and 100  $\mu\text{m}$ . At normal environmental temperatures objects emit infrared between these wavelengths; hotter objects, such as fires, emit infrared at wavelengths shorter than thermal infrared. *Compare* [REFLECTED INFRARED](#).

**thermal low** Basically [\\*non-frontal depression](#) associated with strong daytime heating of land surfaces, mainly in summer, occurring for example in Arizona, Spain, and in northern-Indian low-pressure cells. The occurrence of thermal lows is also influenced by imbalances in the main wind-streams in the upper [\\*troposphere](#).

**thermal metamorphism** *See* [CONTACT METAMORPHISM](#).

**thermal resistivity** A measure of how poorly a material conducts heat; the inverse of [\\*thermal conductivity](#).

**thermal wind** A wind that is generated when the air temperature changes by a large amount over a short horizontal distance. When warm and cool air lie adjacent to each other, pressure decreases with height faster in the cool

air than in the warm air, due to its greater density and, therefore, compression. Consequently, a surface of constant pressure slopes upward from the cool to the warm air, and the thickness of each layer (and, therefore, the gradient of this slope) increases with height. The strength of the **\*geostrophic wind** is proportional to the pressure gradient, so the wind strengthens with increasing height. The thermal wind blows with the cool air to its left in the northern hemisphere and to its right in the southern hemisphere. The **\*jet stream** is a thermal wind.

**thermic** See **PERGELIC**.

**thermistor** A semiconductor whose electrical resistance decreases markedly with increasing temperature, and which is used as a sensitive device for measuring temperature.

**thermochronology** The study of the rate at which a rock cooled from a molten state, based on measurements of the amount of **\*daughter** products present in a **\*mineral**. Cooling rocks contain trace amounts of radioactive isotopes, which decay, and as mineral crystal lattices form they retain the daughter products. The amount of daughter products retained in the lattice depends on temperature. Consequently, the amount of a daughter product that has accumulated in a mineral can be used to calculate the time that has elapsed since the mineral was at its **\*blocking temperature**.

**thermoclastic (insolation weathering)** Process of physical **\*weathering** whereby the stresses set up when a rock is alternately heated and cooled become sufficient to cause **\*failure**. It may bring about surface spalling (splintering) because of the low conductivity of rock, and may be particularly effective on polymineralic rocks because of varying coefficients of expansion. Some observational and experimental evidence suggests that the process may be less effective than was believed formerly. See also **EXFOLIATION**.

**thermocline** Generally, a gradient of temperature change, but applied more particularly to the zone of rapid temperature change between the warm surface waters and cooler waters at depth. In the oceans, this zone of rapid temperature change starts at 10–500 m below the surface and can extend down to more than 1500 m. In polar regions the thermocline is generally absent since the ocean surface is covered with ice in winter and solar radiation is small in summer. In thermally stratified lakes in summer the

thermocline separates the warm surface waters (*\*epilimnion*) from the cooler deep waters (*\*hypolimnion*).

**thermograph** Instrument that gives a continuous record of temperature for a day or for a week. The device uses a helical strip of two metals with differing coefficients of expansion. The resulting opening and closing of the coil operates a pen which produces a line over a calibrated chart on a round clock drum.

**thermohaline circulation** Vertical circulation induced by the cooling of surface waters in a large water body. This cooling causes convective overturning and consequent mixing of waters. In the oceans, this circulation usually involves temperature and *\*salinity* variations acting together. See also MERIDIONAL OVERTURNING CIRCULATION.

**thermo-hygrograph** See HYGROTHERMOGRAPH.

**thermokarst** *\*Periglacial* land-form assemblage characterized by enclosed depressions (some with standing water) and so presenting a *\*karst* appearance. It is caused by the selective thaw of ground ice associated with thermal erosion by stream and lake water and may reflect climatic changes or human activity.

**thermoluminescence** A phenomenon whereby certain minerals, when slowly heated to temperatures below their level of incandescence, emit light. Some minerals will do this only when in contact with oxygen (or air) and in this case it is known as oxyluminescence. The light energy emitted is measured using a photomultiplier and recorded as a function of temperature, to produce a 'glow curve'.

**thermoluminescence dating** A technique used to date crystalline material, most commonly ceramics, and sediments. Crystalline materials absorb radiation from the decay of radioactive elements in the soil around them and from *\*cosmic radiation*. When these materials are heated, as in firing ceramics, their absorbed radiation is released as *\*thermoluminescence*. Once cooled, they resume their absorption of ionizing radiation. This releases electrons, a proportion of which become trapped in defects and impurities in the crystal lattice. Crystals in sediments begin absorbing radiation from the time they are buried. Subsequent heating or exposure to light releases thermoluminescence that can be measured and the age of the

sample can be calculated from the amount of radiation emitted. Thermoluminescence dating is used widely in archaeology and has an age range of 1000–500 000 years.



<http://geology.cr.usgs.gov/capabilities/gronemtrac/geochron/thermo/tech.html>

- Explains and describes the thermoluminescence technique.

**thermonuclear reactions** Nuclear fusion reactions occurring between various nuclei of light elements at very high temperatures. These are the source of energy generation in stars. *See also* [NUCLEOSYNTHESIS](#).

**thermophile** An *\*extremophile* (domain *\*Archaea*) that thrives in environments where the temperature is high, typically up to 60 °C. *Compare* [HYPERTHERMOPHILE](#).

**thermopile** Instrument for the measurement of direct and diffuse short-wave radiation, which utilizes a bank of thermocouples covered by a glass or other transparent dome.

**thermoremanent magnetization (TRM)** The *\*remanent magnetization* that heated *\*ferromagnetic* materials acquire as they cool through their *\*Curie temperature* down to room temperature or below.

**thermosphere** The upper zone of the atmosphere, above about 80 km, where solar radiation of the shortest wavelengths is absorbed. In this zone, which includes the *\*ionosphere*, temperature increases with height, but because of the very low atmospheric density there, the heat capacity is minute. *See also* [ATMOSPHERIC STRUCTURE](#).

**Thermosphere Ionosphere Mesosphere Energetics and Dynamics (TIMED)** A joint *\*NASA* and French Centre National d'Études Spatiales (CNES) mission carried on a 587-kg spacecraft that was launched on 7 December 2001 into a circular *\*orbit* at a height of 625 km, inclined 74° to the equator. It began transmitting data on 22 January 2002 and was still functioning in November 2018. TIMED carries four instruments: the global ultraviolet imager (GUVI), solar extreme ultraviolet experiment (SEE), TIMED Doppler interferometer (TIDI), and sounding of the atmosphere using broadband emission radiometry (SABER). These measure the

atmospheric temperatures and energy flows 60–180 km above the surface in the *\*mesosphere* and lower *\*thermosphere*.



<http://www.timed.jhuapl.edu/WWW/index.php>

- A joint CNES and NASA mission to study the influence of the Sun and humans on the mesosphere, ionosphere, and lower thermosphere of Earth's atmosphere.

**Theropoda** Suborder of *\*saurischian* dinosaurs which consists exclusively of bipedal, carnivorous forms. It includes the *\*coelurosaurs* and the *\*carnosaurs*, and ranged from the Upper *\*Triassic* to the *\*Cretaceous*.

**thickness** The difference in height in the atmosphere between particular pressure levels, e.g. between 100 and 50 kPa. Isolines of a given interval on charts are termed 'thickness lines'.

**thin section** A slice of a mineral or rock which is glued to a glass slide with resin of a particular *\*refractive index* (usually 1.540) and ground down to a thickness of 0.03 mm so that light may be transmitted through it when it is examined under a transmitted-light *\*polarizing microscope*. The mineral or rock chip is ground in successive stages, using carborundum or other suitable abrasives mixed with water, and when finished the section is covered with a glass cover slip, again attached with resin. This is the standard method for preparing most of the rock-forming minerals for study.

**thixotropic mud** A type of mud commonly used to cool the drill *\*bit* and remove rock chips during drilling. When the drill motion ceases, the thixotropic properties of the mud prevent rock chips descending and blocking the hole. See *THIXOTROPY*; *SOL*; *QUICK CLAY*.

**thixotropy** The property possessed by some materials of changing from *\*gel* to liquid under *\*shearing stress*, e.g. when shaken, and returning to the original state when at rest. Some muds of the *\*smectite* group have this property and are useful for flushing *\*boreholes*. The change is completely reversible: no change in water content or composition occurs. On a slope the material will flow downwards as long as its velocity is sufficient to maintain minimum shearing stress.

**tholeiite** An extraordinarily abundant, fine-grained, *\*igneous* rock consisting of *\*essential* calcium-*\*plagioclase*, subcalcic *\*augite*, and pigeonite, with *\*interstitial \*glass* or fine *\*quartz-\*feldspar* intergrowths. Tholeiite is a type of *\*basalt*, oversaturated with silica; it occurs as plateau *\*lavas* on the *\*continental crust* and as the main *\*extrusive* component of the ocean floor. *Compare* MID-OCEAN-RIDGE BASALT.

**tholi** *See* THOLUS.

**tholins** A class of molecules that form by the action of ultraviolet radiation or spark discharge on gases of simple compounds, including CH<sub>4</sub>, C<sub>2</sub>H<sub>6</sub>, H<sub>2</sub>O, HCHO, and H<sub>2</sub>S. They are usually brown, sometimes sticky. Tholins (from the Greek *tholos*, meaning muddy) do not form on Earth but are found on the surface of *\*Titan* and other bodies in the outer *\*solar system*. The term was coined by the astronomer Carl Sagan and the chemist Bishun Khare.

**tholoid** *See* DOME.

**tholus** (*pl.* **tholi**) A small, dome-shaped hill on the surface of an extraterrestrial body.

**Thompson, Benjamin (Count Rumford)** (1753–1814) An American-born inventor, Rumford is best known for founding the Royal Institution in Britain in 1799, ‘to promote useful science’. He made studies of heat, leading to investigations of gunpowder, and improvements in lamps, cookers, and fireplaces.

**Thomson, Charles Wyville** (1830–82) An Irish naturalist, Thomson became professor of natural science at Edinburgh University. He was especially interested in life in the deep oceans, which led him to propose an expedition to make detailed studies of the ocean. The result was the *\*Challenger* expedition, at the end of which Thomson began the arrangement of the collections and publications of the results.

**Thomson, William (Lord Kelvin)** (1824–1907) A mathematician and physicist at Glasgow University, Kelvin is best known for his theory of thermodynamics, which he applied to his work on the age of the *\*Earth*. He used studies of the internal heat of the Earth, the age of the *\*Sun*, and *\*tidal friction*, to give an estimate of the Earth’s age of about 20 million years.

This led to conflict with geologists and evolutionists, who saw evidence that the Earth must be much older. See [KELVIN SCALE](#).

**Thoracica** See [CIRRIPEDIA](#).

**thoracic vertebra** See [VERTEBRA](#).

**thorax** The three segments of the body of an arthropod ([\\*Arthropoda](#)) that lie between the head and the abdomen. Each thoracic segment carries a pair of legs. The three thoracic segments are termed the prothorax, mesothorax, and metathorax. In insects ([\\*Insecta](#)), the mesothorax and metathorax may each carry a pair of wings. See [TRILOBITOMORPHA](#).


**thorium–lead dating** A [\\*radiometric](#) dating method based on the [\\*radioactive decay](#) of  $^{232}\text{Th}$ , to yield  $^{208}\text{Pb} + 6\text{He}^4$ , with a half-life (see [DECAY CONSTANT](#)) of 13 900 million years. The minerals used include [\\*sphene](#), [\\*zircon](#), [\\*monazite](#), [\\*apatite](#), and other rare U/Th minerals. The method is not totally reliable and is usually employed in conjunction with other methods. In most cases the results are [\\*discordant](#) as a result of lead loss. The ratio of  $^{208}\text{Pb}:^{232}\text{Th}$  compared with  $^{207}\text{Pb}:^{235}\text{U}$  ratio is particularly useful. The Th–Pb system can also be interpreted by means of [\\*isochron](#) diagrams similar to those used in the [\\*rubidium–strontium](#) method.

**Thornthwaite, Charles Warren** (1889–1963) An American climatologist who devised the system of climate classification that bears his name. Thornthwaite taught on the faculties of the Universities of Oklahoma (1927–43) and Maryland (1940–46), and at Johns Hopkins University (1946–55) before becoming director of the Laboratory of Climatology at Centerton, New Jersey, and professor of climatology at Drexel Institute of Technology, Philadelphia. He was president of the Section of Meteorology of the American Geophysical Union from 1941 to 1944 and in 1951 he was elected president of the Commission for Climatology of the World Meteorological Organization.

**Thornthwaite climate classification** System for describing climates, devised in 1931 and revised in 1948 by the American climatologist [C. W. \\*Thornthwaite](#), that divides climates into groups according to the vegetation characteristic of them, the vegetation being determined by precipitation effectiveness ( $P/E$ , where  $P$  is the total monthly precipitation, and  $E$  is the

total monthly evaporation). The sum of the monthly  $P/E$  values gives the  $P/E$  index, which is used to define five humidity provinces, with associated vegetation. A  $P/E$  index of more than 127 (wet) indicates rain forest; 64–127 (humid) indicates forest; 32–63 (subhumid) indicates grassland; 16–31 (semi-arid) indicates steppe; less than 16 (arid) indicates desert. In 1948 the system was modified to incorporate a moisture index, which relates the water demand by plants to the available precipitation, by means of an index of **\*potential evapotranspiration (PE)**, calculated from measurements of air temperature and day length. In arid regions the moisture index is negative because precipitation is less than the PE. The system also uses an index of thermal efficiency, with accumulated monthly temperatures ranging from 0, giving a frost climate, to more than 127, giving a tropical climate. *See also* [KÖPPEN CLIMATE CLASSIFICATION](#); [STRAHLER CLIMATE CLASSIFICATION](#).

**three-cell model** An attempt to represent the atmospheric circulation systems over a hemisphere by three adjoining vertical cells of meridional surface motion, transferring energy from equatorial to polar regions. This concept of heat transfer by meridional circulation has generally been superseded by more modern ideas of travelling waves. *See also* [HADLEY CELL](#); [ROSSBY WAVES](#).



**Three-cell model**

**three-dimensional seismology (3D seismic)** A technique, used in prospecting for crude oil and natural gas, in which a set of **\*geophones** is located very precisely to detect vibrations from a series of experimentally induced seismic disturbances over a considerable area. In one survey, covering 49 km<sup>2</sup>, 1700 seismic disturbances produced 725 000 geophone traces. The technique has led to the identification of larger reserves than had previously been suspected and increased production.

**threshold** A value beyond which the physical environment adjusts to a change in processes. An example is provided by a **\*glacier**, where a build-up of ice and snow over a number of years reaches a critical level and once that critical level (or threshold) is exceeded there is a sudden change in the process of **\*basal sliding** and the glacier surges forward.

**threshold-slope concept** The mechanism by which valley incision limits the elevation of mountain peaks in a **\*collision zone**. At the same time as the colliding plates push the mountains upward, rivers and glaciers carve deep valleys that increase the slope of mountainsides. When the slopes

exceed a threshold value, landslides (see [MASS-WASTING](#)) reduce the slope to the [\\*angle of repose](#) while removing material from higher levels. A balance is reached at which landslides maintain a constant mountainside slope while also constraining the increase in mountain elevation.

**thrombolites** Structures with a clotted microtexture and no internal laminae, built by [\\*cyanobacteria](#) and found in calcareous, sublittoral [\\*facies](#).

**throughfall** The part of [\\*precipitation](#) which, having been intercepted by vegetation, then falls on to the ground surface. See [INTERCEPTION \(1\)](#).

**throughflow** See [INTERFLOW](#).

**throw** The vertical displacement of a [\\*dip-slip fault](#); its measurement is often complicated by the difficulty of accurately identifying markers on either side of the [\\*fault plane](#).

**thrust** A low-angle (commonly less than 45°) [\\*reverse fault](#), with a significant dip-slip component, in which the [\\*hanging wall](#) overhangs the [\\*footwall](#). [\\*Synthetic thrust](#) sets form [\\*imbricate](#) fan structures which may be thrust-bound, when they form a [\\*duplex](#). Single thrusts typically show a 'staircase' trajectory composed of [\\*ramps](#) and [\\*flats](#).

**thrust-sheet-top basin** See [PIGGYBACK BASIN](#).

**thufur** An earth hummock, some 0.5 m high and 1–2 m wide, which is found in contemporary and past [\\*periglacial](#) environments. Normally it possesses a core of [\\*sediment](#), which suggests that a form of differential heaving by ground ice is responsible for its development.

**thulite** See [ZOISITE](#).

**thundercloud** Popular term for [\\*cumulonimbus](#), which is associated with [\\*thunderstorms](#).

**thunderstorm** A storm of fairly local scale in which strongly developed [\\*cumulonimbus](#) cloud produces thunder and lightning, usually with rain and strong, gusting wind, and often with hail.

**Thuringian** A European stratigraphic term which was originally used as an equivalent to the Zechstein and was later extended to include the whole

Upper *\*Permian*. See also *AUTUNIAN*.

**thuringite** See *CHLORITE*.

**Thurnian** The cold-*\*stage* marine *\*silts* from the middle of the threefold subdivision of the deposits of the Ludham *\*borehole*. They are Lower *\*Pleistocene* in age. See also *ANTIAN*; *BAVENTIAN*; *LUDHAMIAN*; *PASTONIAN*.

**Thvera** A normal *\*polarity subchron* which occurs within the *\*Gilbert* reversed *\*chron*.

**Tibetan Plate** Lying between the Indian and *\*Eurasian Plates*, the Tibetan Plate consists of two continental blocks, northern Tibet and southern Tibet, divided by a *\*suture* with *\*ophiolites*. Most of the *\*Cenozoic* deformation of this part of the Indian–Eurasian collision has been to the south of Tibet, in the Himalayas.

**Tibetan Plateau** An area of Tibet with an elevation of approximately 2 km above sea level, below which the *\*crust* is more than 80 km thick; this has been ascribed to *\*underthrusting* of the crust of the *\*Indo-Australian Plate* (i.e. by *\*A-subduction*) or to horizontal shortening as a result of *\*thrust \*tectonics*. The uplift has been both rapid and recent: one estimate is 2500–3000 m since the *\*Pliocene*.

**tibia** **1.** In a tetrapod vertebrate (*\*Tetrapoda*), the anterior long bone of the lower hind limb (the ‘shin’ bone). **2.** In an insect (*\*Insecta*), the long and often narrow segment of the leg that articulates proximally with the femur (or ‘thigh’) and distally with the tarsus (or ‘foot’).

**tibiotarsus** The bone formed by the fusion of the *\*tarsal bones* and the *\*tibia*, found in birds and some dinosaurs.

**tidal barrage** A dam or barrier built across a tidal channel to allow a power-generating plant to operate. The siting of the barrage requires there to be a large *\*tidal range*, in excess of 5 m, and a large water body open to the sea, with a narrow entrance. One barrage and generating plant in operation since 1960 is on the Rance Estuary in Brittany, France. It generates electricity by the use of two-way turbines, as water flows in and out of the *\*estuary* through the barrage. The Severn Estuary and Morecambe Bay are potential sites for tidal barrages in the British Isles.

**tidal bundle** \*Cross-stratification that comprises the set of sand units and \*mud drapes deposited during the lunar cycle of the \*tides. During \*spring tides the more vigorous water movement results in thick sand units and small mud drapes; the less vigorous water movement during \*neap tides results in thinner sand units and larger mud drapes.

**tidal correction** An adjustment that is made to \*gravimeter data acquired in a \*gravity survey in order to remove the effect due to \*Earth tides, which causes a variation of about 0.3 mgal over 12 hours, depending on latitude and the positions of the Sun and Moon (i.e. the time since the last \*spring tide).

**tidal current** An alternating, horizontal movement of water associated with the rise and fall of the \*tide, these movements being caused by gravitational forces due to the relative motions of Moon, Sun, and Earth. Offshore tidal currents tend to exhibit rotary patterns, while in areas near coasts the currents follow rectilinear paths and reverse periodically (ebb and flow currents). Tidal currents can often reach velocities of 2.5m/sec near shores.

**tidal flat** An intertidal sandflat, \*mudflat, and marsh area developed in some \*lagoons in \*mesotidal areas, and in protected bays and estuarine areas along \*macrotidal coasts. Extensive tidal flats occur in the Wash (eastern England), Wadden Zee (Holland), and along the North-Sea German coast. Tidal flats also occur in warmer climates, as in the Persian Gulf, where \*carbonate and \*evaporite deposits develop. In tropical areas tidal flats tend to be colonized by \*mangrove swamps.

**tidal friction** The friction exerted on the Earth because of the \*phase lag between the \*tides and the gravitational attraction of the Moon, Sun, and planets. It is mostly due to the  $M_2$  ocean tide ( $M_2$  is the principal component of the forces acting in the direction of the Moon).

**tidal heating** The generation of heat due to friction produced by the strong tidal forces exerted by a very massive parent body on a body moving about it in an elliptical orbit. The intensity of tidal heating is proportional to the square of the orbital eccentricity, being zero in a circular orbit and reaching a maximum in a parabolic orbit, and inversely proportional to the size of the orbit.

**tidal inlet** A narrow channel that connects the open sea with a **\*lagoon**. Tidal inlets often occur in **\*barrier island** systems and are typified by small-scale **\*deltas** at each end of the inlets, resulting from the high-velocity, tidal-reversal currents that flow through the channels.

**tidalite** A **\*sediment** deposited under the dominant influence of **\*tidal currents**.


**tidal range** The difference in height between consecutive high and low waters. The tidal range varies from a maximum during **\*spring tides** to a minimum during **\*neap tides**. In tide tables daily high- and low-water heights are given for each geographical locality mentioned.

**tidal rhythmite** A **\*rhythmite** deposited by the action of tides.

**tidal stream** The flow of water in and out of estuaries, bays, and other restricted coastal openings associated with the rise and fall of the **\*tide**. Landward (flood) and seaward (ebb) streams or currents often follow different paths in shallow-water areas, so forming ebb/flood avoidance cells and a braided pattern of sandbanks in these coastal openings.

**tidal theory** Theory of the origin of the **\*solar** system, involving the approach near the Sun of another star. This set up tidal forces, and the instability of the Sun resulted in part of its mass being torn off to form the planets. The theory was proposed by Sir James H. Jeans (1877–1946) and Sir Harold **\*Jeffreys**.

**tide** **1.** The periodic rise and fall of the Earth's oceans, caused by the relative gravitational attraction of the Sun, Moon, and Earth. The effect of the Moon is about twice that of the Sun, giving rise to the **\*spring–\*neap** cycle of tides. Variation in tides is caused by: (a) changes in the relative positions of the Sun, Moon, and Earth; (b) uneven distribution of water on the Earth's surface; and (c) variation in the sea-bed topography. Semi-diurnal tides are those with two high and two low waters (period 12 hours and 25 minutes) during a tidal day (24 hours and 50 minutes). Diurnal tides have one high and one low water during a tidal day. **2.** See **EARTH TIDES**.



## Tide

**tieline** 1. A line on a metamorphic, composition–assemblage diagram (e.g. an *\*ACF* or *\*AFM* diagram), which joins the compositions of *\*minerals* coexisting in equilibrium. 2. A *\*survey* line that crosses others so that all lines can be calibrated to the same value.

**tie point** In *\*remote sensing*, a point on the ground which occurs in two or more images and that can be used to co-register images.

**Tiffanian** A N. American *\*stage* of the *\*Palaeocene* (60.5–56 Ma ago) preceded by the *\*Torrejonian* and followed by the *\*Clarkfordian*.

**‘tiger’s eye’** See RIEBECKITE.

**tight gas** Natural gas that is difficult to extract because it is held in rock or sand (called tight sand) that is relatively impermeable and non-porous.

**tight sand** See TIGHT GAS.

**Tiglian** A late *\*Pliocene* *\*stage* (about 2 Ma ago) which *\*isotope* records from the tropical Atlantic- and Pacific-Ocean cores and other climatic indicators suggest may have been a time of increased cooling.

**tile drain** Short lengths of concrete or ceramic pipe placed end to end to make a drain but also allow water to enter at the joints, laid at any appropriate depth and spacing to remove water from the soil. Nowadays continuous, slotted, plastic piping is often used.

**till** Collective term for the group of *\*sediments* laid down by the direct action of glacial ice without the intervention of water. The sediments may be classified in terms of *\*particle size* (they range between *\*clay*-rich and *\*clast*-dominated types, according to source area and travel distance), or grouped by the basic process of debris release: subglacial melt gives rise to 'lodgement till'; surface *\*ablation* gives 'ablation till', followed by *\*flow till* after further movement; and the general thaw of static ice produces 'melt-out till'. See TILL FABRIC ANALYSIS.

**till fabric analysis** The measurement of the orientation and *\*dip* of the *\*clasts* contained within a *\*till*, and the subsequent plotting and analysis of the data. These may indicate the direction of a former glacial advance: clasts within a lodgement till tend to lie parallel to the direction of ice movement.

**tillite** A lithified deposit of *\*boulder clay* or *\*till* produced by the action of *\*glaciers*. See DIAMICTITE.

**tilloid** General term applied to a chaotic mixture of large blocks set in a *\*clay*-rich *\*matrix*; such mixtures include *\*mudflows*, *\*landslide* deposits, and *\*tillites*.

**till plain** A smooth plain underlain by *\*till*. It may be well preserved, as in mid-western USA, or may be dissected by later *\*erosion*, as in the lowlands of central and eastern England.

**tilt-block tectonics** An extensional tectonic style in which crustal blocks are rotated along *\*normal faults* which tend to become nearly horizontal at depth. It is thought to be important on the *\*passive margin* of a rifted continental mass.

**tiltmeter** An instrument used to measure the change in ground slope. Using laser technology, small changes in the relative levels of three connected but displaced fluid reservoirs are accurately measured to monitor the direction and magnitude of slope changes. Changes of a fraction of a radian can be monitored by this instrument. Tiltmeters are particularly useful on active *\*volcanoes* for measuring the rate of ground inflation or deflation prior to and during an eruptive event. Areas showing large ground inflations are usually correlated with *\*magma* rising into the high-level volcanic plumbing system and may pinpoint the potential site of an eruption.

**time-averaged velocity** See AVERAGE VELOCITY.

**TIMED** See THERMOSPHERE IONOSPHERE MESOSPHERE ENERGETICS AND DYNAMICS.

**time–distance curve** A graph showing the arrival times of \*seismic waves as a function of \*shot–\*geophone \*offset distance. In seismic-\*refraction surveys it is also called a  $t-d$  curve or  $t-x$  curve. In seismic-\*reflection surveys, travel times squared ( $t^2$ ) are plotted against the square of the offset distance ( $x^2$ ) to produce a  $t^2-x^2$  graph. In both sets of graphs, the inverse gradient of straight-line segments provides information about the \*seismic velocity for the appropriate layer.

**time domain** A reference framework in which measurements are related to time, rather than to \*frequency. Compare FREQUENCY DOMAIN.

**Time History of Events and Macroscale Interactions during Substorms (THEMIS)** A \*NASA mission to fly five small spacecraft through explosive geomagnetic disturbances to discover what triggers the northern and southern \*auroras. The five spacecraft were launched on 17 February 2007, from Cape Canaveral, into highly elliptical orbits: probe 1 orbital period 4 days, \*perigee 1.500 Earth radii ( $R_E$ ), \*apogee 31.645  $R_E$ ; probe 2 period 2 days, perigee 1.168  $R_E$ , apogee 19.770  $R_E$ ; probe 3 period 1 day, perigee 1.118  $R_E$ , apogee 12.127  $R_E$ ; probe 4 period 1 day, perigee 1.118  $R_E$ , apogee 12.127  $R_E$ ; probe 5 period 0.8 day, perigee 1.350  $R_E$ , apogee 10.044  $R_E$ .

**time plane** An imaginary surface linking \*interfaces of equal age within a body of \*strata.

**time-rock unit** See CHRONOSTRATIGRAPHIC UNIT.

**timescale** The subdivision of \*geologic time based on one of several different criteria. Timescales related to the rate of sedimentation (see SUBSIDENCE), the rate of \*sea-floor spreading, and \*radiometric dating are used. See APPENDIX B: TIME SCALES.

**time-stratigraphic unit** See CHRONOSTRATIGRAPHIC UNIT.

**tinguaite** An *\*undersaturated*, medium- to coarse-grained, *\*igneous* rock consisting of *\*essential \*alkali* feldspar, *\*nepheline*, and *\*aegirine* (with or without sodic *\*amphibole* or *\*biotite*). Tinguaitite is the *\*hypabyssal* equivalent of *\*phonolite*. The term 'intrusive phonolite' is now preferred.

**tinstone** See CASSITERITE.

**Tioughniogan (Tioughiogan)** See GIVETIAN.

**TIROS** See TELEVISION AND INFRARED OBSERVATION SATELLITE.

**Titan (Saturn VI)** One of the major satellites of *\*Saturn*, with a radius of 2575 km; mass  $1345.5 \times 10^{20}$  kg; mean density 1881 kg/m<sup>3</sup>; visual albedo 0.21. It was discovered in 1655 by C. Huygens. Titan has a dense atmosphere, with a surface pressure of 0.15 MPa. The atmosphere consists primarily of nitrogen, with up to 6% argon and a small amount of methane. The surface temperature is about 94 K. The *\*Cassini* mission reached Titan in 2004 and launched the *\*Huygens* lander in 2005. Huygens photographed a landscape marked by features resembling river channels and the shoreline of a dry lake. It is thought that from time to time liquid methane falls to the surface as rain, forming slushy areas and perhaps filling lakes. Data from Cassini also suggests there is an extensive ocean of liquid water beneath the surface shell.

**titanaugite** A titanium-rich *\*augite*.

**Titania (Uranus III)** One of the major satellites of *\*Uranus*, and the largest. Its radius is 788.9 km; mass  $35.27 \times 10^{20}$  kg; mean density 1710 kg/m<sup>3</sup>; visible albedo 0.18. The surface has many impact craters surrounded by bright ejecta.

**titanite** See SPHENE.

**titanohaematites** See TITANOHEMATITES.

**titanohematites (titanohaematites)** A group of magnetic *\*minerals* belonging to the *\*ilmenite* (FeTiO<sub>3</sub>) to *\*hematite* (Fe<sub>2</sub>O<sub>3</sub>) series that have a *\*Curie temperature* of up to 680°C.

**titanomagnetite** A titanium-rich *\*magnetite*; i.e. a magnetite (Fe<sup>2+</sup>Fe<sup>3+</sup><sub>2</sub>O<sub>4</sub>) in which Ti<sup>2+</sup> replaces some of the Fe<sup>2+</sup>. Titanomagnetites have a *\*Curie*

**temperature** ranging from about 200°C to 580°C.

**Tithonian** A **\*stage** (150.8–145.5 Ma ago) in the Late **\*Jurassic**, preceded by the **\*Kimmeridgian** and followed by the **\*Berriasian**.

**Titius–Bode law (Bode’s law)** An empirical arithmetical relationship between the distances of the planets from the Sun. If the Sun–Earth distance is taken as 10, then the distances of Mercury, Venus, Earth, Mars, Jupiter, and Saturn are approximately satisfied by the sequence: 4, 4 + 3, 4 + 6, 4 + 12, 4 + 48, and 4 + 96. The more accurate version of the ‘law’ is given by the function  $r_n = AB^n$ , where  $r_n$  is the distance of the  $n$ th planet,  $A$  is a constant, and  $B = 1.73$ . This version is due to Mary Blagg (1858–1944). It is not clear that the ‘law’ has any fundamental significance, but may be merely a consequence of gravitational and tidal evolution following planetary and satellite formation. The **\*asteroid** belt occurs about 4 + 24, corresponding to the former ‘missing planet’ in the sequence, that was believed to exist for many years. **\*Uranus**, discovered in 1781, occurs close to the next term, 4 + 192, but the position of **\*Neptune** does not fit the ‘law’. The regular satellites of the giant planets fit modifications of the ‘law’. The ‘law’ was discovered in 1766 by J. D. Titius (1729–96) and was popularized by J. E. **\*Bode** (1747–1826).

**titrimetric analysis** Method of analysis in which a **\*solution** of the substance being determined is treated with a solution of a suitable reagent of exactly known **\*concentration**. The reagent is added to the substance until the amount added is equivalent to the amount of substance to be determined. The point of equivalence (end point) in titrimetric analysis is usually determined by a change in colour of an auxiliary reagent (the indicator), but may also be detected by electrical means (potentiometric, conductometric, titrimetry, etc.).

**tjale (frost table)** A frozen surface at the base of the **\*active layer**, which moves downwards as thaw occurs. The tjale should not be confused with the upper limit of **\*permafrost**, the permafrost table.

**TLE** See TRANSIENT LUMINOUS EVENT.


**toad’s eye tin** See WOOD TIN.

**Toarcian** A *\*stage* in the European Early *\*Jurassic* (183–175.6 Ma ago; Int. Commission on Stratigraphy, 2004), preceded by the *\*Pliensbachian*, followed by the *\*Aalenian*, and roughly contemporaneous with the mid-Uroroan (New Zealand). *See also* LIAS.

**toeset** The asymptotic, basal portion of a *\*Gilbert-type delta*, generally characterized by fine-grained *\*sediment* deposited at the toe of the prograding *\*foreset*. *See also* TOPSET.

**Tolstojan** The second period of the *\*Mercurial geologic timescale*, lasting from 3900–3850 Ma ago, during which the *\*late heavy bombardment* continued and many of the smaller basins on the surface of *\*Mercury* formed.

**tombolo** A *\*spit* that links an island to the mainland or to another island, formed by deposition when waves are refracted round the island.



**Tombolo**

**Tommotian** The basal Russian-Kazakhstan *\*stage* of the *\*Cambrian \*period*, underlain by the Nemakit–Daldynian, overlain by the *\*Atdabanian*, and dated at about 534–530 Ma ago. *See also* CAERFAI EPOCH.

**tomography** Any technique by which details of a subsurface plane can be represented graphically. *See* SEISMIC TOMOGRAPHY.

**tonalite** An oversaturated, coarse-grained, *\*igneous* rock consisting of *\*essential* sodic *\*plagioclase* ( $An_{27-36}$ ), *\*quartz*, *\*hornblende*, and/or *\*biotite*, with *\*accessory \*apatite*, *\*zircon*, and iron oxide.

**Tonawandan** See SCHEINWOODIAN.

**Tonga–Kermadec Trench** The oceanic **\*trench** in the western **\*Pacific Ocean** which forms part of the boundary between the **\*Indo-Australian** and **\*Pacific Plates**. There is a volcanic gap (i.e. a break in the **\*island arc**) to the north-west of the trench, possibly related to the subduction of the Louisville Rise.

**Tongaporutuan** A **\*stage** (10–6 Ma ago) in the **\*Miocene** of New Zealand, underlain by the **\*Waiauan**, overlain by the **\*Kapitean**, and roughly contemporaneous with the upper **\*Tortonian** and **\*Messinian** stages.

**tonguing** See ALBELUVISOLS.

**tonhäutchens** See CUTAN.

**Tonian** The first period of the **\*Neoproterozoic** era, that began 1000 Ma ago and ended 850 Ma ago, both its beginning and end being defined by fixed dates. The Tonian followed the **\*Stenian** period and was followed by the **\*Cryogenian** period.

**tonstein** A compact, **\*kaolinite-rich \*mudstone**, which developed as a kaolinitic **\*palaeosol**, and is frequently found as thin bands within **\*coal** seams or resting directly above the coal. Some tonsteins are laterally extensive and are believed to be the product of weathered volcanoclastic **\*ash**.

**Tool for the Analysis of RAdiations from lightNings and Sprites (TARANIS)** A **\*microsatellite** mission by the French space agency in collaboration with institutions from the US, Denmark, Japan, the Czech Republic, and Poland to study the mechanisms that produce transient luminous emissions and gamma-ray flashes in the atmosphere above areas experiencing thunderstorms (Taranis was the Gallic god of thunder and lightning). The mission is scheduled to launch in 2019.



[https://taranis.cnes.fr/en/TARANIS/GP\\_mission.htm](https://taranis.cnes.fr/en/TARANIS/GP_mission.htm)

- TARANIS mission home page.

**tool mark** The impression made on the surface of a soft bed of **\*sediment** by the impact of an object (tool), or the dragging of an object over the

sediment by a current. Tool marks include bounce, *\*prod*, *\*skip*, *\*groove*, and *\*chevron marks*, which develop by differing interaction of the tools with the sediment.

**topaz** A nesosilicate mineral,  $\text{Al}_2\text{SiO}_4(\text{OH},\text{F})_2$ ; sp. gr. 3.5–3.6; *\*hardness* 8; *\*orthorhombic*; colourless, pale yellow, pale blue, yellowish, or sometimes pink; often transparent; vitreous *\*lustre*; crystals are *\*prismatic* and often *\*bipyramidal* with the vertical faces striated, but it can also be *\*massive* and granular; *\*cleavage* perfect basal {001}; typically occurs in *\*granite* *\*pegmatites*, *\*rhyolite*, and *\*quartz* veins, and extensively as an *\*accessory mineral* in granites, associated with *\*fluorite*, *\*tourmaline*, *\*beryl*, and *\*cassiterite*, also in *\*alluvial* deposits. It is associated with pneumatolytic action (see *PNEUMATOLYSIS*) and is a constituent of greisen. The original cairngorms (see *QUARTZ*) were topaz crystals. It is named after Topazos Island in the Red Sea.

**TOPEX/Poseidon** A joint NASA and French Centre National d'Études Spatiales (CNES) space mission that was launched from French Guiana on 10 August 1992 into a *\*polar orbit* at a height of 1330 km. Its instruments measured sea level, tides, and large-scale ocean features, as well as changes in the heat stored in the oceans. The mission was planned to last three years, but it was not until October 2005 that a malfunction caused the satellite to cease transmitting data and it was turned off on 18 January 2006. See *JASON-1*.



<http://sealevel.jpl.nasa.gov/missions/topex/>

- A joint CNES and NASA mission to measure the ocean topography of Earth.

**toplap** A discordant relationship in which the upper boundary of a *\*depositional sequence* is marked by the termination of initially inclined beds (e.g. those formed in the *\*foresets* of *\*deltas*). Toplap results mainly from nondeposition, possibly with minor *\*erosion*. Coastal toplap occurs during a *\*stillstand* of sea level. Each unit of *\*strata* dips seaward with its upper, terminal edge wedging back towards the land. Successive units of strata build out laterally towards the sea. Compare *BASELAP*.

**topographic correction (terrain correction)** A correction applied to observed geophysical values to remove the effects of topography. In gravity studies, it is the correction applied to each individual determination of gravity to allow for the attraction of rocks occurring as hills above the height of the recording station and as valleys below this level. For seismic corrections see [ELEVATION CORRECTION](#); [STATIC CORRECTION](#).

**topology** In *\*phylogenetics*, the branching pattern of a *\*phylogenetic tree*.

**toposequence** Sequence of *\*soils* in which distinctive soil characteristics are related to the topographic situation.

**topotype** In taxonomy, a specimen found in the type locality of a taxon to which it is thought to belong, but that is not necessarily of that *\*type series*.

**topozone** See [TEILZONE](#).

**topple (debris fall, earth fall)** A type of *\*mass-wasting* in which cohesive blocks of relatively dry earth fall rapidly from steep faces such as river banks.

**topset beds** The upper, near-horizontal layers deposited on a *\*Gilbert-type delta*, generally characterized by the coarsest *\*sediment* found on the prograding (see [PROGRADATION](#)) *\*delta*. See also [FORESET](#); [TOESET](#).

**topsoil** **1.** The superficial layer of *\*soil* that is moved in cultivation. **2.** The A *\*soil horizon* of a *\*soil profile*. **3.** Any surface layer of soil.

**tor** Mass of exposed bedrock, standing abruptly above its surroundings, and typically but not exclusively developed on granitic rocks. It may be formed by selective subsurface *\*weathering* followed by the removal of the weathered debris, by differential frost-shattering, or as an end-product of *\*scarp retreat* under semi-arid conditions.

**torbanite (cannel shale)** A carbonaceous *\*oil shale* that is sapropelic, usually occurs as lenses in *\*coal* seams, and was possibly derived from vegetation-rich mud. See [SAPROPEL](#); [SAPROPELIC COAL](#).

**torbernite** A *\*secondary* mineral, with the formula  $\text{Cu}(\text{UO}_2)_2(\text{PO}_4)_2 \cdot 8\text{--}12\text{H}_2\text{O}$ ; sp. gr. 3.2; *\*hardness* 2.5; *\*tetragonal*; normally bright emerald-green, but occasionally dark green; pale green *\*streak*; vitreous *\*lustre*; crystals *\*tabular*, square, foliated, scaly aggregates; *\*cleavage* perfect basal

{001}; occurs in the oxidized zones of *\*veins* containing uranium and copper minerals, often with *\*autunite*.

**tornado** A relatively small-scale (about 100 m diameter) ‘twisting’ or rotating column of air, like a funnel, with high wind speeds and great destructive force over the narrow path of its movement. Tornadoes can occur anywhere in temperate latitudes, but are especially violent in unstable air conditions in the central parts of the USA. See [APPENDIX C: ENHANCED FUJITA SCALE](#).

**Tornquist Line** See [FENNOSCANDIAN BORDER ZONE](#).

**toroidal field** A magnetic field with no radial components, e.g. where the magnetic lines of force lie on an electrically conducting spherical surface such as the surface of the Earth’s *\*core*. Such a field is not detectable at the Earth’s surface, in contrast to a *\*poloidal field*.

**Torrejonian** A N. American *\*stage* of the *\*Palaeocene* (62.5–60.5 Ma ago), preceded by the *\*Puercan* and followed by the *\*Tiffanian*.

**Torridonian** A *\*stage* of the Upper *\*Proterozoic* of north-western Scotland, from about 1100 to 600 Ma ago, named from Loch Torridon.

**torrid zone** **1.** General climatic term for the region between the northern and southern tropics, broadly the ‘equatorial’ zone. **2.** In the climate classification devised by Aristotle, the region (in his view) uninhabitable for humans in either hemisphere that extends from the equator to the Tropic of Cancer or the Tropic of Capricorn. Compare [FRIGID ZONE](#); [TEMPERATE ZONE](#).

**torsion** **1.** The effect of twisting an object in opposite directions from either end. Torsion tests, using rod-shaped samples, allow stable ductile deformation to be studied up to large *\*strain* states. **2.** In *\*Gastropoda*, the twisting of the body through 180° so that the digestive and nervous systems have a U-shape and the *\*mantle cavity*, anus, gills, and two nephridiopores (excretory openings) occupy an anterior position behind the head.

**torsion balance** A weighing instrument in which the required weight of a sample is set by rotating a lever to move a needle to the appropriate position on a graduated scale, and then the sample is added to the weighing pan, a little at a time, until it counterbalances the pre-set weight precisely and the

needle is brought back to the 'zero' position. This is a useful and rapid technique when large numbers of routine weighings need to be performed in the range of 0.01–5.0 g.

**torta** A low, fairly flat, approximately symmetrical, volcanic dome formed by successive extrusions on level ground, each increment from the vent thinning the previously erupted material as it pushes it outwards. *Torta* is Spanish for 'pie'; volcanic tortas are common in the Andes.

**Tortonian** A Late *\*Miocene* *\*stage*, preceded by the *\*Serravallian*, followed by the *\*Messinian*, and dated at 11.608–7.246 Ma ago (Int. Commission on Stratigraphy, 2004). It is roughly contemporaneous with part of the *\*Mohnian* (California), *\*Waiauian* and lower *\*Tongaporutuan* (New Zealand), and the lower *\*Mitchellian* (Australia). The *\*stratotype* is at Tortona, Italy.

**torus** The shape of a doughnut; a geometrical surface formed by rotating a circle (or other closed curve) about a circle in its plane without intersecting it.

**toscanite** See RHYODACITE.

**total core recovery (TCR)** The total length of *\*core* recovered from a *\*borehole* as a percentage of the length of the borehole.

**total intensity (*F*)** The magnitude of the geomagnetic vector field or of the magnetization of an object.

**total internal reflection** The *\*reflection* of a *\*seismic wave* from the boundary between the layer in which it is travelling and an adjacent layer. A wave in a low-velocity medium which strikes the inner surface of a boundary with a higher-velocity medium at an angle greater than the *\*critical angle*, and therefore is wholly reflected within the layer, is partially converted into another type of wave at the *\*interface*, rather than being refracted out of the layer. See S-WAVE.

**total range zone** See TAXON RANGE ZONE.

**total stress** *In situ* *\*stress* in a rock body, minus the fluid pressure.

**tourmaline** A member of the *\*cyclosilicates* and a borosilicate  $\text{Na}(\text{Mg}, \text{Fe}^{2+}, \text{Mn}, \text{Li}, \text{Al})_3\text{Al}_6 (\text{BO}_3)_3[\text{Si}_6\text{O}_{18}](\text{OH}, \text{F})_4$ . There are three

important members of this family of **\*minerals**: dravite  $(\text{NaMg}_3\text{Al}_6(\text{BO}_3)_3[\text{Si}_6\text{O}_{18}](\text{OH},\text{F})_4)$ ; **\*schorl**  $(\text{Na}(\text{Fe}^{2+},\text{Mn})_3\text{Al}_6(\text{BO}_3)_3[\text{Si}_6\text{O}_{18}](\text{OH},\text{F})_4)$ ; and elbaite  $(\text{Na}(\text{LiAl})_3\text{Al}_6(\text{BO}_3)_3[\text{Si}_6\text{O}_{18}](\text{OH},\text{F})_4)$ ; sp. gr. 2.9–3.2; **\*hardness** 7.0–7.5; **\*trigonal**; black, bluish, pink, or green, never colourless; elongate **\*crystals** common, also **\*acicular** needles and massive or radiating aggregates; **\*cleavage** good {1120} prismatic; occurs in granite **\*pegmatites**, pneumatolytic (see **PNEUMATOLYSIS**) veins, and **\*granites** as elbaite and schorl varieties; it may occur in the rock luxullianite formed by pneumatolytic action after boron has been introduced, where it will occur with **\*topaz**, **\*spodumene**, **\*cassiterite**, **\*fluorite**, and **\*apatite**. Dravite variety occurs in metamorphosed impure **\*limestones** and rarely in some **\*basic \*igneous** rocks; tourmaline is a common detrital ‘heavy’ mineral in sedimentary rocks. Good multicoloured crystals can be used as gemstones.

**tourmalinization** A pneumatolytic (see **PNEUMATOLYSIS**) modification of a pre-existing **\*igneous** rock in which boron-rich, late-stage fluids react with the **\*mineral** assemblage of the primary rock (usually **\*granite**), eventually producing a rock consisting of **\*tourmaline** and **\*quartz**. Intermediate stages in the process, showing **\*alkali feldspar** partly replaced by rosettes of tourmaline, are commonly preserved in granites. Luxullianite, found at Luxulyan, Cornwall, is a fine example.

**Tournaisian** A **\*stage** in the **\*Mississippian \*epoch**, preceded by the **\*Famennian (\*Devonian)**, followed by the **\*Visean**, and dated at 359.2–345.3 Ma ago (Int. Commission on Stratigraphy, 2004). It comprises the Hastarian and Ivorian stages. It is roughly contemporaneous with the lower **\*Carboniferous** Limestone (Britain) and the **\*Kinderhookian** and lower **\*Osagean** (N. America). See also **DINANTIAN**.

**tourquoise** See **TURQUOISE**.

**Toutatis** A **\*solar system** asteroid (No. 4179), measuring  $4.6 \times 2.4 \times 1.9$  km; approximate mass  $10^{13}$  kg; rotational period irregular; orbital period 1.1 years. It is a double object, its two components probably in contact, one estimated to be 2.5 km in diameter and the other 1.5 km. On 29 September 2004 Toutatis reached its closest approach to Earth, at a distance of about 1.5 million km.

**tower karst** A form of karstic morphology (see **KARST**) that is developed mainly in low latitudes, characterized by residual hills of **\*limestone** rising from a flat plain. The hills have near-vertical sides, resembling towers.

**Toyonian** A Russian-Kazakhstanian **\*stage** of the Early **\*Cambrian \*epoch** (518.5–513 Ma ago), preceded by the **\*Botomian** and followed by the **\*Amgan**.

**T-peg** In **\*geomorphology**, a device for measuring the rate of surface **\*creep**. It consists of a metal rod to which a cross-piece is attached. The apparatus is inserted into the **\*regolith** and the cross-piece is levelled. Soil creep causes the cross-piece to tilt, and the degree of inclination is measured with a graduated spirit-level. The relative movement between the ground surface and the depth of insertion can then be inferred.

**trace** A recorded data-set for one channel. On a seismic-**\*refraction** record made for a 12-channel **\*seismograph**, each channel would provide one wave-form which, when viewed with the others, would give a **\*seismic record**. See also **WIGGLE TRACE**.

**trace element 1.** An element that occurs in minute but detectable quantities in minerals and rocks, much less than 1%. All elements except the most common rock-forming elements (O, Si, Al, Fe, Ca, Na, K, Mg, and Ti) generally occur as trace elements, except where locally concentrated in their **\*ores**. **2.** In biology, the occurrence in plant or animal tissue of minor amounts of an element which is essential to its growth.

**trace-element fractionation** The redistribution (fractionation) of elements between solid and liquid **\*phases** caused by heating and partial melting, and which also takes place in crystallization. For example, fractionation of **\*trace elements** apparently occurs in **\*chondritic** meteorites during the formation of **\*chondrules** (small molten droplets produced by heating and melting in outer space which are subsequently quenched).

**trace fossil (ichnofossil)** A **\*biogenic \*sedimentary structure** formed by the behavioural activity of an animal on or within a given substrate. The study of trace fossils is called 'ichnology'. Traces are most frequent at the **\*interface** between different **\*lithologies** (e.g. **\*sandstone** and **\*shale**), and are classified on various criteria including **\*morphology** and preservation. Of these two, the second is preferred as a toponomic classification (i.e.

classification by place of occurrence) and, apart from the processes of preservation, considers the position of the trace within the depositional unit concerned. In 1970, A. Martinsson divided traces into four groups dependent on their relationship to the casting medium: epichnia are surface ridges or grooves; endichnia are tubes or **\*burrows** formed within the casting medium; hypichnia are grooves or ridges preserved on the lower surface of the main body of the casting medium; and exichnia are formed by **\*bioturbation** outside the main body of the casting medium. See FOSSILIZATION.

**tracer** A substance that is used to follow the passage of **\*groundwater** in places where it cannot be observed directly. Typical tracers include fluorescent dyes and salt. The presence of radioactive **\*isotopes**, e.g. tritium and carbon-14, may also be used as tracers in that they allow the age of groundwater to be determined. The presence of small amounts of other substances may also be used to make deductions about the origin and flow path of groundwaters.

**Tracheophyta (vascular plants)** (kingdom **\*Plantae**) Phylum comprising plants that have vascular tissues (xylem and phloem) through which water and nutrients are transported. The division includes the subdivisions **\*Pteridophytina** (spore-bearing vascular plants) and **\*Spermatophyta** (seed-bearing plants).

**trachyandesite** See ANDESITE.

**trachybasalt** The fine-grained, **\*extrusive** equivalent of **\*syenogabbro**. When **\*undersaturated**, **\*feldspathoidal** minerals take the place of **\*alkali** feldspar to generate feldspathoidal trachybasalts known as **\*tephrites** (**\*olivine**-free) and **\*basanites** (olivine present). Trachybasalts are found on the stable **\*continental crust** and on some oceanic islands.

**trachyte** Fine-grained volcanic rock of the **\*alkaline** series of **\*intermediate rocks**. Trachyte is the volcanic equivalent of **\*syenite**.

**trachytic texture** General **\*petrographic** term applied to volcanic rocks that have a fine mat of orientated **\*feldspar** laths, suggestive of flow. See TRACHYTOIDAL.

**trachytoidal** Applied to a **\*foliation** resulting from a parallel alignment of tabular **\*feldspars** in a coarse-grained rock, e.g. **\*syenite**. When the same

alignment occurs in fine-grained rocks, e.g. *\*trachytes*, it is called '*\*trachytic texture*'.

**track** A *\*biogenic \*sedimentary structure* grouped under the Scoyenia assemblage of *\*trace fossils*. The term may refer to a line of vertebrate footprints or to traces left by the limbs of arthropods (*\*Arthropoda*).

**traction carpet** See BED LOAD.

**traction load** See BED LOAD.

**trade-wind inversion** The inversion of temperature *\*lapse rate* with height over a major zone of the trade-wind belt, which is very significant in tropical meteorology. Moist tropical air, extending up to 2–3 km above the surface, is 'boxed in' or 'trapped' by dry, clear, warmer air above, resulting from *\*subsidence* in subtropical *\*anticyclones*. The inversion forms where the subsiding air meets a surface flow of cooler maritime air. The top of a cloud layer marks the base of the inversion.

**trade winds** Old maritime term, much used in meteorology, indicating the steadiness of direction of the prevailing tropical easterly winds, which blow from subtropical high-pressure areas in latitudes 30–40° north and south, generally north-easterly in the northern hemisphere and south-easterly in the southern hemisphere. They are most nearly constant in latitudes centred on 15° north and south. Climatic conditions associated with the belt vary from fine anticyclonic weather in the poleward and eastern margins, caused by *\*subsidence*, to stormier conditions near the equator and western margins, caused by less stable, deeper, moist air.

**Traditional Stratigraphic Scale (TSS)** See STRATIGRAPHIC SCALE.

**trail** 1. An anterior extension of some brachiopod (*\*Brachiopoda*) shells, usually at a large angle to the general plane of the posterior part of the shell.  
2. A *\*biogenic \*sedimentary structure* formed by the movement of snails, clams, or perhaps snakes over the *\*sediment* surface, and classified with *\*tracks* under the Scoyenia assemblage of *\*trace fossils*.

**trailing edge** See PASSIVE MARGIN.

**training area** See CLASSIFICATION.

**tramontana** Local wind in the north-western Mediterranean, which brings dry, cold conditions from the north or north-east across the mountains.

**tranquil flow (sub-critical flow)** See CRITICAL FLOW; FROUDE NUMBER.

**transcurrent fault** See STRIKE-SLIP FAULT.

**transfer fault** A vertical or subvertical \*fault which, via \*dip-slip and \*strike-slip movements, allows the juxtaposition of two \*fault zones which have different displacement characteristics. Lateral \*ramps are the equivalent in \*thrust terrains.


**transfluence** See GLACIAL BREACH.

**transformation twinning** A type of \*crystal twinning caused by a change in the structure of a crystal under different conditions of temperature and pressure (\*polymorphism), e.g. the \*cristobalite–\*tridymite–\*quartz series, where the high-temperature form is \*hexagonal and the low-temperature form is \*trigonal.

**transform fault** A type of \*strike-slip fault in an ocean, occurring at the boundaries of lithospheric \*plates, in which the direction of movement of the crustal blocks is reversed (or ‘transformed’) in comparison with a strike-slip fault on land. For example, at mid-ocean \*ridges the offset between adjacent ridge sections is a transform fault; where the displacement is \*dextral (right lateral) the motion, due to spreading, is left lateral, and vice versa. Generally, transform faults occur at right angles to the ridge itself and indicate the direction of spreading. The active transform fault extends into an inactive fracture zone.

**transgression (marine)** An advance of the sea to cover new land areas, due to a rise in the sea level relative to the land. As a result, shallow-water \*sediments are overlain by those characteristic of deeper water, e.g. shelf mud is deposited on coastal sand. Compare REGRESSION. See ONLAP.

**transgressive systems tract (TST)** In the \*genetic stratigraphic sequence model used in \*sequence stratigraphy, a bounding surface formed by a rapid rise in sea level. Compare HIGHSTAND SYSTEMS TRACT; LOWSTAND SYSTEMS TRACT.



**Transgressive systems tract**

**transient creep** See PRIMARY CREEP.

**transient electromagnetic method (TEM)** An electromagnetic surveying method in which the source signal consists of a train of pulses rather than a continuous wave-form. Quasi-transient methods are used for continuous **\*profiling** (e.g. the **\*INPUT** method), while true TEM methods are most commonly used for depth soundings. It is a **\*time-domain** method.

**transient flow** See STEADY FLOW.


**transient luminous event (TLE)** An optical phenomenon, associated with thunderstorms, that is visible in the upper atmosphere. TLEs include **\*red sprites**, **\*blue jets**, **\*blue starters**, elves, **\*gigantic jets**, **\*gnomes**, **\*pixies**, **\*sprite halos**, **\*trolls**, and **\*upward lightning**.

**transient variation** Any short-term variation. In geomagnetism, aperiodic fluctuations of the **\*geomagnetic field**, usually on scales of microseconds to a few days.

**transit time** The time taken for a seismic **\*P-wave** to travel through one foot (0.3048 m), measured by a **\*sonic** log in units of  $\mu\text{s}/\text{foot}$ .

**translation** See COILING.

**translational slide (planar slide, translational slump, planar slump)** A type of **\*mass-wasting** in which material slides over a plane surface. The material may be wet or dry rock, debris, or soil, and the slide may be slow or, in the case of a rock mass, extremely rapid.



**Translational slide:** Mass-wasting in which material slides across a plane surface.

**translational slump** See TRANSLATIONAL SLIDE.

**translation gliding** See DEFORMATION TWINNING; SLIP.

**translocation** In *\*pedology*, the movement of *\*soil* materials in solution or in suspension from one *\*soil horizon* to another.

**translucent** Applied to a substance (e.g. a mineral) that transmits some light but through which the outlines of objects cannot be seen. This condition is common in varieties of *\*quartz* and *\*fluorite*.

**translucidus** From the Latin *translucidus* meaning ‘transparent’, a variety of cloud, occurring in extensive layers or sheet form, which is *\*translucent*, allowing the Sun or Moon, and occasionally some stars, to be visible. This characteristic can be presented by *\*stratus*, *\*stratocumulus*, *\*altostratus*, and *\*altocumulus*. See also CLOUD CLASSIFICATION.

**transmission coefficient ( $T$ )** The ratio of the amplitude of a transmitted ray ( $A_2$ ) to that of the incident ray ( $A_0$ ), such that  $T = A_2/A_0$ . In the case of normal incidence, and in terms of the *\*acoustic impedances* ( $Z_1$  and  $Z_2$ ) of the two media above and below the boundary,  $T = 2Z_1/(Z_2 + Z_1)$ . Also,  $T = 1 - R$ , where  $R$  is the *\*reflection coefficient*. The transmission coefficient can also be expressed in terms of energy ( $T'$ ), when  $T' = 4Z_1Z_2/(Z_2 + Z_1)^2$ .

**transmission electron microscope** An *\*electron microscope* that is used to explore the internal structure of *\*mineral \*thin sections*. The electron beam passes through the specimen and is focused by lenses on the other side to

produce an image on a phosphor-coated plate. The microscope will resolve images to about 0.3 nm, achieving magnifications of up to 1 000 000 times.

**transmissivity (*T*)** The rate at which \*groundwater is transmitted through a unit width of an \*aquifer under a unit \*hydraulic gradient. It is often expressed as the product of the \*hydraulic conductivity and the full saturated thickness of the aquifer and has units of the form  $\text{m}^3/\text{day}/\text{m}$ .

**transmittance** The ratio of \*electromagnetic radiation passing through a material to that incident upon its surface.

**trans-orogen basin** A continental \*drainage basin in which water moves across the \*strike of a mountain belt. The Columbia River in the N. American Rocky Mountains is an example.

**transparency** Degree to which a substance allows radiation of varying wavelengths to pass through, as opposed to absorbing it. *See also* ABSORPTION.

**transparent** Applied to a substance (e.g. a mineral) which transmits light and through which the outlines of objects can be seen clearly.

**transpiration** The removal of moisture from the \*soil by plant roots, its translocation up the stem to the leaves, and its evaporation through the stomata. The flow of water through the plant is known as the transpiration stream. It reduces leaf temperatures, and is thought to be important for mineral absorption and translocation within plants. The process imposes a number of environmental requirements upon plants; for example, wilting and desiccation result from an insufficient supply of water.

**transpolar drift** The movement of large amounts of \*sea ice across the Arctic Basin, at 1–4 cm/s, from the Siberian coast through the Fram Strait, to the north-east of Greenland, and down the eastern coast of Greenland. The transpolar drift is driven by the Beaufort Gyre, an anticyclonic circulation of water mainly confined to the uppermost 300 m, over the Canadian Basin, off Canada and Alaska. One effect of the transpolar drift is to pile up ice along the northern coasts of Canada and Greenland, where in places it is up to 8 m thick.

**transpression** A tectonic regime which combines both transcurrent \*strike-slip movement with oblique compression. \*Flower structures are commonly

associated with transpressional regimes.

**trans-Saharan seaway** The marine seaway that, during two intervals in the Late *\*Cretaceous*, extended from *\*Tethys* in the north through what are now Libya, Chad, Niger, and Nigeria, to the newly developing S. Atlantic Ocean. On both occasions ammonite (*\*Ammonoidea*) and ostracod (*\*Ostracoda*) faunas are known to have migrated through this seaway. The first event was in the latest *\*Cenomanian* and earliest *\*Turonian*, and was re-established only in the Late *\*Campanian* to Early *\*Maastrichtian* interval. At the southern end (in present-day Nigeria) there was a structural control in the form of the Benue Trough, but the remainder of the sea-way appears to have been controlled only by global sea level change.

**transtension** A tectonic regime combining transcurrent *\*strike-slip* movement with oblique *\*extension*. Such a regime is associated with oceanic spreading *\*ridges* and *\*transform faults*.

**transverse dune** See DUNE.

**transverse-type coast** See ATLANTIC-TYPE COAST.

**transverse wave** See S-WAVE.

**trap** See ANTICLINAL TRAP; DECCAN TRAPS; FAULT TRAP; REEF TRAP; STRATIGRAPHIC TRAP; STRUCTURAL TRAP; UNCONFORMITY TRAP.

**trap-door caldera** A *\*caldera* in which the floor is hinged on one side.

**trapezohedron** A *\*crystal* form where the faces are in the shape of a trapezoid with four sides, none of which are parallel. It is developed in the *\*cubic*, *\*tetragonal*, and *\*hexagonal \*crystal systems*, to give a *\*closed* form similar in shape to a pyramid. It can be referred to the normal axes of symmetry but frequently lacks a horizontal plane of symmetry.

**travel time** The time taken for a wave to travel from a source generator (e.g. a *\*shot* or transmitter), through some media, to a detector (*\*geophone* or receiver), at a known *\*offset*. It is used to compute *\*travel-time curves* and *\*seismic records* in seismic-*\*refraction surveying*. See also ONE-WAY TRAVEL TIME; TWO-WAY TRAVEL TIME. Compare TRANSIT TIME.

**travel-time curve** One of a set of curves which show the *\*travel times* of *\*P-waves* and *\*S-waves* as functions of distance, expressed either in range

kilometres, or in degrees (as epicentral angles). The distance from the recording *\*seismometer* to the *\*epicentre* can be determined by measuring the times that elapse between the arrival of the P-wave and S-wave, and the S-wave and *\*L-wave*.

**traverse** In surveying, a line which connects two points and passes through a series of locations which are to be studied. A traverse may be open-ended and discontinuous, or closed (i.e. it returns to its starting point).

**travertine** Calcium carbonate deposited by *\*precipitation* from carbonate-saturated waters, particularly from hot springs. Travertine deposits are sometimes *\*massive*, but often display a concentric or fibrous internal structure, sometimes building large, concentric, spherical masses. Travertine is also found in cave deposits in the form of *\*stalagmites* and *\*stalactites*. A porous, sponge-textured form of travertine is referred to as '*\*tufa*' or 'calc-sinter'.

**tree-ring analysis** See DENDROCHRONOLOGY.

**trellis drainage pattern** See DRAINAGE PATTERN.

**trema** See ARCHAEOGASTROPODA.

**Tremadocian (Gasconadian)** Oldest (488.3–478.6 Ma ago) of the six *\*stages* that comprise the *\*Ordovician \*period*, preceded by the Late *\*Cambrian \*epoch* and followed by the *\*Arenig* stage. *\*Mudstones* and *\*sandstones* of the Tremadoc *\*series* occur in N. America, Ireland, Wales, England, and Scandinavia. They mark the *\*continental shelf* and slope areas that occur on the southern edge of the northern *\*Iapetus Ocean*.

**tremata** See ARCHAEOGASTROPODA.

**tremolite** An important member of the *\*monoclinic* calcium-rich *\*amphiboles*  $\text{Ca}_2(\text{Mg,Fe}^{2+})_5[\text{Si}_4\text{O}_{11}]_2(\text{OH,F})_2$ , which forms a series with ferroactinolite; sp. gr. 3.02–3.44; *\*hardness* 5.0–6.0; *\*monoclinic*; white to greyish-white; vitreous *\*lustre*; crystals simple, long, *\*prismatic*, and *\*acicular*, often in thin, radial, rod-like, fibrous, and felted masses; *\*cleavage* perfect *\*prismatic* {110}; widespread *\*amphibole* mineral in *\*igneous* rocks, and also in metamorphosed, crystalline *\*limestones*,

**\*dolomites**, and in **\*schists** and **\*hornfels**. It is named after the Tremola Valley near St. Gotthard, Switzerland. *See also* **JADEITE**.

**Trempealeauan** *See* **DOLGELLIAN**.

**trench (oceanic trench)** An elongate depression of the ocean floor which runs parallel to the **\*trend** of adjacent volcanic islands (**\*island arc**) or continent. Oceanic trenches are up to 11 km deep, typically 50–100 km wide, and may be thousands of kilometres long. In cross-section the trench slopes are usually asymmetric, with a steeper slope on the landward side. Most trenches are associated with **\*subduction zones**.

**trend** The **\*azimuth** of a geologic feature, commonly of a **\*fold axis**, and written as a compass bearing.

**Treptichnus pedum** A **\*trace fossil** that is abundant in **\*Ediacaran** sediments. It consists of burrows that turn in ways that produce fan shapes and shapes reminiscent of a twisted rope.

**trevorite** *See* **MAGNETITE**.

**triad** *See* **CRYSTAL SYMMETRY**.

**Triana** *See* **DSCOVER**.

**triangle zone** A triangular area bounded by **\*thrusts**, which has been formed by the truncation of an earlier thrust surface by a later **\*back thrust**.

**triangular facet** *See* **VENTIFACT**.

**Triassic** The earliest (251–199.6 Ma ago) of the three **\*periods** of the **\*Mesozoic era**. As a result of the mass **\*extinctions** of the late **\*Palaeozoic**, Triassic communities contained many new faunal and floral elements. Among these were the ammonites (**\*Ammonoidea**), modern corals, various molluscs (**\*Mollusca**), the **\*dinosaurs**, and certain **\*gymnosperms**.



<https://www.livescience.com/43295-triassic-period.html>

- Triassic Period Facts: Climate, Animals & Plants.


**triaxial cell** An instrument for testing the compressive strength of *\*soils* or *\*rocks*. A load is applied to the sample, which is contained in an impermeable membrane surrounded by a fluid. The loading is increased until failure occurs.

**triaxial compression test** A test for the compressive strength in all directions (*compare* **UNIAXIAL COMPRESSION TEST**) of a *\*rock* or *\*soil* sample, using a *\*triaxial cell*. Tests in which drainage is prevented are called ‘undrained’ tests and the strengths obtained are ‘undrained’ strengths. When *\*pores* are allowed to empty, the tests are called ‘drained’ tests and the strengths obtained are ‘drained’ strengths.

**triaxial ellipsoid** See **PLANE STRAIN**; **SHAPE FABRIC**.

**tributary** A stream that discharges into a larger river.

**triclinic** One of the seven *\*crystal* systems, and the one with the lowest *\*crystal symmetry*, characterized by three *\*crystallographic axes* of unequal length, none of which are at right angles to each other. There is a centre of symmetry about which is oriented a pair of parallel faces. The *\*mineral* *\*plagioclase feldspar* is the best-known example.




Triclinic

**tricolpate sulci** See **POLLEN**.

**Triconodonta** (class *\*Mammalia*, subclass Prototheria) Order that includes the earliest of all mammals, living from the late *\*Triassic* until the early *\*Cretaceous* and distributed over the northern continents. Typically the molar teeth each had a row of three sharp conical cusps, the teeth of the upper and lower jaws forming a shearing device. Premolars and molars were differentiated, probably with some replacement, and probably the young were fed on milk secreted by the mothers. Triconodonta may have been *\*homoiotherms* and nocturnal, and possibly they were arboreal. They are believed to have been true carnivores rather than insectivores. *Triconodon*, one of the larger forms, from the Upper *\*Jurassic*, was the size of a modern cat. The order is believed to have evolved from *\*therapsids* independently of the main line of mammalian evolution and to have left no descendants.

**tridymite** A high-temperature variety of silica (SiO<sub>2</sub>) which is stable between 870 and 1470 °C at normal pressures. It occurs as tabular plates in cavities of volcanic rocks and also in *\*stony meteorites*. See also QUARTZ.


**trigonal (rhombohedral)** One of the seven *\*crystal systems* (closely related to the *\*hexagonal* system) in which the unit cell has the shape of a *\*rhombohedron*, consisting of six *\*crystal* faces all of which have two pairs of parallel sides. The crystals may be referred to four *\*crystallographic axes*: one vertical axis of three-fold *\*crystal symmetry*, and three equal, horizontal axes of two-fold symmetry which are separated by an angle of 120°. The minerals *\*calcite* and *\*quartz* always occur in this system.



**Trigonal**

**Trilete** See SPORE.

**Trilobita (trilobites)** (phylum *\*Arthropoda*) The most primitive arthropod class, known from more than 3900 *\*fossil* species. Inhabitants of *\*Palaeozoic* seas, the trilobites appeared first in the early *\*Cambrian*, had their widest distribution and greatest diversity in the Cambrian and *\*Ordovician* periods, and became extinct in the *\*Permian*. The body was divided into three regions: an anterior *\*cephalon*, comprising at least five, fused segments; a mid-body or *\*thorax*, with a varying number of segments; and a hind region or *\*pygidium*. All three regions were divided by a pair of furrows running the length of the body, giving a trilobite appearance (i.e. a median or axial lobe, flanked on either side by a lateral lobe). The mouth was situated in the middle of the central surface of the cephalon. Paired gill-bearing limbs were attached to the membranaceous, *\*pleural* skeleton. X-ray studies show the eyes to have resembled the compound eyes of living arthropods (see TRILOBITE EYE). Trilobites ranged in size from 0.5 mm long planktonic (see PLANKTON) forms to those nearly 1 m in length; most species were 3–10 cm long. There were nine orders: *\*Redlichiida*; *\*Agnostida*; *\*Naraoiidae*; *\*Corynexochida*; *\*Lichida*; *\*Phacopida*; *\*Ptychopariida*; *Asaphida*; and *\*Proetida*.



**Trilobita**

**trilobite** See TRILOBITA.

**trilobite eye** The eyes of trilobites (*\*Trilobita*) are compound and made up of radially disposed visual units. Most trilobite eyes are 'holochroal' (i.e. many polygonal lenses are in contact with one another and covered by a single cornea). 'Schizochroal' eyes occur in the Phacopina (i.e. the lenses are large, separated from one another, each has its own cornea, and each lens is in two parts); it has been shown that this arrangement produces a sharp focus.

**Trimerophytina** See PSILOPHYTALES.

**trim line** A boundary (or zone) between frost-shattered and glacially scoured bedrock in an upland region. It marks the junction between glacial and *\*periglacial* activity and so indicates the position of a former *\*glacial* limit.

**trimorphism** See POLYMORPHISM.

**Trinculo (Uranus XXI)** A lesser satellite of *\*Uranus* with a radius of 9 km.

**triple core barrel** See CORE BARREL.

**triple junction** The point where three lithospheric *\*plates* meet. This junction may involve three oceanic *\*ridges* (an 'RRR' junction), thought to have developed from a *\*domal uplift*; or involve some other configuration of ridge ('R'), *\*transform fault* ('F'), and oceanic *\*trench* ('T'). Some types of triple junction are stable, while others evolve geologically rapidly into a different configuration. An example of an RRR junction is the point where the *\*African*, *\*South American*, and *\*Antarctic Plates* meet in the S. Atlantic.

**triple-junction method** A mathematical method in which a vector diagram is used to calculate the relative velocity of a third *\*plate*, where the velocities of two of the plates meeting at a *\*triple junction* are known.

**triple point** The point on a *\*phase diagram* at which planes or *\*phases* meet. This is the temperature and pressure at which the three phases (solid, liquid, and gas) of a substance can coexist (*see also* KELVIN SCALE). In a *\*solid-solution* system, the triple point is usually defined by the original pressure and temperature conditions at which the solid, liquid, and gaseous phases of a substance are in equilibrium.

**triserial** Applied to one of the patterns of chambers that occurs in *\*Foraminiferida*, in which there are three chambers to each *\*whorl* in the *\*test*. In some species there can be a change from a single chamber in each whorl ('uniserial'), through two chambers ('biserial'), to triserial.

**tritium clock** Tritium (T) is a naturally occurring radioactive *\*isotope* ( $^3\text{H}$ ) of hydrogen ( $^2\text{H}$ ), having two neutrons and one proton in the nucleus. The isotope decays (half-life = 12.26 years, *see* DECAY CONSTANT) by beta emission (*see* BETA DECAY) to stable helium ( $^3\text{He}$ ) and this can be used to measure the age of water samples back to approximately 30 years. Tritium is produced in the upper atmosphere by the interaction of fast cosmic-ray neutrons with stable  $^{14}\text{N}$ . The tritium combines with hydrogen and oxygen to form HTO, which is then dispersed throughout the *\*hydrosphere*. The natural production rate of tritium is 15–45 atoms/min/cm<sup>2</sup> of the Earth's surface. In some experiments, artificially produced tritium has been introduced into *\*groundwater* and used to trace and time underground

movements. It can also be used to measure and time rates of mixing in oceanic current systems.

**Triton (Neptune I)** A satellite of *\*Neptune*, with a diameter of 2705.2 km; mass  $214.7 \times 10^{20}$  kg; mean density 2054 kg/m<sup>3</sup>; visual albedo 0.7. Its orbit is *\*retrograde*.

**TRMM** See TROPICAL RAINFALL MEASURING MISSION.

**trochiform** Applied to a gastropod (*\*Gastropoda*) shell where the sides of the spire are evenly conical and the base is flat.



**Trochiform**

**trochoid** See SOLITARY CORALS.

**trochospiral** Applied to the growth pattern in *\*Foraminiferida* where material is added to the *\*test* in a helical manner.

**troctolite** A coarse-grained, *\*igneous* rock consisting of *\*essential* magnesium-rich *\*olivine* and calcium-rich *\*plagioclase feldspar* with *\*accessory \*ilmenite*; i.e. it is *\*gabbro* without the *\*pyroxene* component. Troctolites grade into gabbros by a gradual increase in pyroxene content and a gradual decrease in olivine content.

**troilite** An iron sulphide mineral, FeS; sp. gr. 4.8; *\*hardness* 4.0; greyish-brown; *\*metallic \*lustre*; black *\*streak*; *\*massive* or granular; occurs as nodules in *\*iron meteorites*.

**troll** A *\*transient luminous event* that resembles a *\*blue jet* but is red.

**trona** A mineral,  $\text{NaHCO}_3 \cdot \text{Na}_2\text{CO}_3 \cdot 2\text{H}_2\text{O}$ , found in salt lake deposits.

**trondhjemite** See GRANODIORITE.

**tropical air** \*Air masses that form over tropical regions. Broadly, they comprise: (a) maritime tropical air, originating over oceans; and (b) continental tropical air, originating over land masses, and especially over N. Africa and the Middle East. Movements of these air masses poleward, with modification of their original characteristics, can have marked influence on weather in mid-latitudes.

**tropical arid morphoclimatic zone** A \*morphoclimatic region in which the rate of \*mechanical weathering is high, especially \*salt weathering, but there is little \*chemical weathering. Wind action is high. \*Fluvial action is generally low, but occasionally high. There is little \*mass-wasting.

**tropical cyclone (revolving storm)** A generally fairly small but intense, closed low-pressure system which develops over tropical oceans. It is circular in shape, comprising concentric rings of \*cumulus and \*cumulonimbus clouds, separated by bands of relatively clear sky in which air is subsiding. The ‘\*eye’ of the storm is fairly still and clear, with subsiding air that warms by compression; the warm air of the ‘eye’ is diagnostic of a tropical cyclone. Wind speeds of at least 33 m/s (force 12 on the \*Beaufort scale, 64 knots or more) define such storms and distinguish them from less intense systems, e.g. tropical \*depressions (of twice or more than twice the diameter) or tropical storms. The atmospheric pressure gradient in such cyclones commonly ranges from about 95 kPa at the centre to about 100 kPa at the margins and the core pressure defines the intensity of the system. See APPENDIX C: SAFFIR-SIMPSON HURRICANE WIND SCALE.

**Tropical Cyclone Programme (TCP)** A project to improve forecasting and warning systems for tropical cyclones. It forms part of \*World Weather Watch.

**Tropical Rainfall Measuring Mission (TRMM)** A joint space mission between \*NASA and the \*Japan Aerospace Exploration Agency (JAXA), within the Earth Science Enterprise programme of the \*European Space Agency, that was launched on 27 November 1997 from the Tanegashima Space Center, Japan, into a near-circular, low-inclination orbit at a height of

350 km, inclined at 35° to the equator. It carries five instruments: precipitation radar (PR), TRMM microwave imager (TMI), visible and infrared scanner (VIRS), clouds and the Earth's radiant energy systems (CERES), and lightning imaging sensor (LIS). The mission measures diurnal variations in tropical precipitation and evaporation.



<http://global.jaxa.jp/projects/sat/trmm/>

- About Tropical Rainfall Measuring Mission.

**tropical semi-arid morphoclimatic zone** A *morphoclimatic* region in which the rate of *chemical weathering* is low to moderate and that of *mechanical weathering* is high in some places, especially in drier and cooler areas. *Fluvial* action is high, but episodic and wind action is moderate to high. There is a high but sporadic rate of *mass-wasting* locally.

**tropical wet–dry morphoclimatic zone** A *morphoclimatic* region in which the rate of *chemical weathering* is high during the wet season and that of *mechanical weathering* is low to moderate. *Fluvial* action is high during the wet season, with material being transported across the surface as well as along channels. There is a fair amount of *mass-wasting*. Wind action is generally low, but locally moderate during the dry season.

**tropopause** The boundary separating a lower layer of the atmosphere (*troposphere*), in which air temperature generally decreases with height, from the layer above (*stratosphere*), in which temperature remains constant or increases with height. The altitude of the tropopause varies according to sea-surface temperature and season, but also over shorter periods, from an average of 10–12 km over the poles (occasionally descending to 8 km or below) to 17 km over the equator. *See also* ATMOSPHERIC STRUCTURE.

**troposphere** The layer of the atmosphere between the Earth's surface and the *tropopause*, within which the air temperature on average decreases with height at a rate of about 6.5 °C/km, though variations that sometimes occur include inversions (temperature increase with height within some limited layer). Most of the atmospheric turbulence and weather features occur in this layer, which contains almost all the atmospheric water vapour

and most of the **\*aerosols** in suspension in the atmosphere (although there is also an important aerosol layer at about 22 km). *See also* **ATMOSPHERIC STRUCTURE**.

**trough** 1. An extension of low atmospheric pressure from the central regions of a low-pressure system into a zone where generally higher pressure prevails. The term 'trough' is also, and in accordance with this definition, applied to equatorward meanders of the flow of the upper westerly winds over middle latitudes. (The 'equatorial trough', where **\*trade winds** meet, is synonymous with the '**\*intertropical convergence zone**'.) *See also* **LONG WAVE**. 2. The lowest point of a **\*fold** surface.

**trough cross-stratification** *See* **CROSS-STRATIFICATION**.

**trowal** Canadian meteorological term for the line of the upper front of an **\*occlusion** and for the region about it where the warm air, having been lifted off the surface, is still relatively low. It is a **\*trough** or valley of warm air that is still undergoing lifting, and is normally marked by clouds and **\*precipitation**.

**true age** *See* **ABSOLUTE AGE**.

**true dip** The **\*dip** which is measured at right angles to the **\*strike** of a plane and which is the maximum value of dip for that plane. *Compare* **APPARENT DIP**.

**true thickness** The **\*orthogonal thickness** of a structure or bed, measured at right angles to its surface. Where the structure or bed is inclined, a surface exposure or **\*borehole** may give a value greater than the true thickness. In its simplest form true thickness can be calculated from an **\*outcrop** exposure by multiplying the width of the exposed layer by the sine of the angle of **\*dip** of the layer.

**truncated spur** A blunt-ended, sloping ridge which descends the flank of a valley. Its abrupt termination is normally due to **\*erosion** by a **\*glacier** which tends to follow a straighter course than the former river.

**T<sub>s</sub>** Symbol used to indicate the actual average temperature at the surface of the Earth, about 15 °C. *See also* **EFFECTIVE TEMPERATURE**.

**tschermakite** See HORNBLLENDE.

**TSS** Traditional Stratigraphic Scale. See STRATIGRAPHIC SCALE.

**TST** See TRANSGRESSIVE SYSTEMS TRACT.

**tsunami** A seismic sea wave of long period, produced by a submarine \*earthquake, underwater volcanic explosion, or massive \*gravity slide of sea-bed \*sediment. In the open ocean such waves are barely noticeable even though they may be travelling at 700 km/h, but on reaching shallow water they slow, reducing the wavelength while increasing the wave amplitude, building up sometimes to heights of more than 30 m and causing severe damage in coastal areas.

**tsunamites** Distinctive sedimentary rock strata that form on the sea bed as a consequence of the remobilization of sediments by \*tsunamis. Tsunamites preserve a record of past tsunamis.

**t-test** A test to calculate the probability that mean values for a particular measurement are significantly different in two sets of data.

**tuba** From the Latin *tuba* meaning 'trumpet', a supplementary cloud feature of \*cumulonimbus or sometimes \*cumulus, characterized by a column or cone of cloud projecting from the cloud base. See also CLOUD CLASSIFICATION.

**tube feet (podia)** In \*Echinodermata, hollow appendages connected to the water-vascular system, used in some species for locomotion and in others for feeding.

**tubercle** The domed, surface structure in trilobites (\*Trilobita). Some tubercles appear to be domes covering a space, others may have been the sites of sensory organs.

**tubular fenestrae** See FENESTRAE.

**tufa (calc-tufa)** \*Sedimentary rock formed by the deposition or precipitation of calcium carbonate, or more rarely \*silica, as a thin layer around saline springs, or by the encrustations on \*stalactites and \*stalagmites. See also TRAVERTINE.

**tuff** The compacted (lithified) equivalent of a volcanic **\*ash** deposit, which has been generated and emplaced by **\*pyroclastic** processes or was water lain, and in which the grain size of the pyroclasts is less than 2 mm. Where the proportion of **\*lapilli**-sized pyroclasts exceeds 10% the term 'lapilli-tuff' is used.

**tumulus** A small mound or dome-like uplift, up to 20 m or more in diameter, on the crust of a **\*lava** flow. Upwarping of the flow crust is caused by the hydrostatic overpressure of lava within the flow interior, or by excess pressure developed due to the difference in rate of flow between the cooler crust and the more fluid lava below it. Unlike a **\*lava blister**, a tumulus is a solid structure.

**tundra** Treeless plain of the Arctic and Antarctic, characterized by a low, 'grassy' sward. Actually, although grasses are rarely absent, sedges (*Carex* species), rushes (*Juncus* species), and wood rushes (*Luzula* species) are the dominant plants, together with perennial herbs, dwarf woody plants, and various bryophytes (**\*Bryophyta**) and lichens.

**Tundra Soil** One of the Great Soil Groups, within suborder 1 of the order Zonal Soils of the 1949 **\*USDA** system of soil classification, based originally on the work of V. V. **\*Dokuchayev**, but now superseded. Tundra Soils are now classified as **\*inceptisols**. They occur on ground that drains poorly (mainly because of **\*permafrost**), and are acid, are 30–60 cm deep, have a high content of organic matter at the surface, and have a microrelief formed by freezing and thawing; their formation, and the decomposition of organic matter, is inhibited by the low temperature.

**tungstates** A group of non-silicate **\*minerals** in which the  $WO_4$  radical is in combination with a number of metal **\*cations**. Important examples include **\*wolframite** ( $(Fe,Mn)WO_4$ ) and **\*scheelite** ( $CaWO_4$ ). Molybdenum may substitute for tungsten to give related minerals, e.g. the molybdates powellite ( $CaMoO_4$ ) and **\*wulfenite** ( $PbMoO_4$ ).

**tunicates** See UROCHORDATA.

**tunnel trend** See CLEANING-UP TREND.

**tunnel valley** A valley cut by a subglacial stream escaping from beneath an **\*ice sheet**. It is well developed in Denmark (where such valleys are called

‘tunneldale’) and in Germany (‘Rinntaler’). Individual examples may be up to 75 km long and up to 100 m deep, with steep sides and flat floors. The long profile may be irregular, as a result of water under pressure being locally forced uphill, and so the valley may now be occupied by a string of lakes (‘ribbon lakes’).


**TURAM method** A geophysical \***electromagnetic** method in which a very long (up to several hundred metres), insulated cable is used, either grounded at both ends or laid out in a large loop, and energized at low frequencies (less than 1 kHz). A receiver coil is moved perpendicularly across the line of the cable and the two orthogonal components of the secondary field are measured.

**turbidite** Sedimentary deposit laid down by a \***turbidity current**.

**turbidity current** A variety of \***density current** that flows as a result of a density difference created by dispersed \***sediment** within the body of the current. Such currents occur off \***delta fronts**, in lakes, and in oceans, and are initiated by the disturbance of sediments on a slope by strong wave action, \***earthquake** shock, or slumping. Turbidity currents in the oceans are thought to move rapidly (at speeds of up to 7 m/s) down the \***continental slope** or \***submarine** canyons along the sea bed, and to deposit originally shallow-water sediments at the foot of the slope or on the \***abyssal plain**. The ideal sequence of sediments laid down by a waning turbidity current is known as the \***Bouma** sequence.

**turbidity flow** See **GRAIN FLOW**.

**turbinate** Applied to a gastropod (\***Gastropoda**) shell that is shaped like a spinning top, but with a rounded base.



### **Turbinate**

**turbulence** Disturbed flow in a moving stream of air. It is manifested by variations of wind speed and direction (including vertical components), by vertical exchanges of mass, heat, momentum, water vapour, and any pollutants present, caused by eddies.

**turbulent flow** Mode of flow occurring in both air and water, and characterized by the superimposition of transverse movements, notably eddies or vortex behaviour, on the general downstream trend. Local upward movement brings about the lifting and removal of particles from stream beds and from the surface of a sandy area. See [REYNOLDS NUMBER](#).


**turnover rate** Measure of the rate of movement of an element through a [\\*biogeochemical cycle](#). Turnover rate is calculated as the rate of flow into or out of a particular pool, divided by the quantity of the element in that pool. Thus it measures the importance of a particular flux in relation to the pool size. Compare [TURNOVER TIME](#).

**turnover time** Measure of the movement of an element in a [\\*biogeochemical cycle](#); the reciprocal of [\\*turnover rate](#). Turnover time is calculated by dividing the quantity of an element present in a particular pool or reservoir by the flux rate for that element into or out of the pool. Turnover time thus describes the time it takes to fill or empty that particular reservoir.

**Turonian** A Late *Cretaceous* stage, preceded by the *Cenomanian*, followed by the *Coniacian*, and dated at 93.5–89.3 Ma ago (Int. Commission on Stratigraphy, 2004).

**turquoise (tourquoise)** A phosphate  $\text{CuAl}_6(\text{PO}_4)_4(\text{OH})_8 \cdot 4-5\text{H}_2\text{O}$ ; sp. gr. 2.60–2.91; hardness 5.0–6.0; *triclinic*; sky-blue to blue-green; white or greenish *streak*; waxy *lustre*; crystals very rare, normally occurs *massive*, granular to *cryptocrystalline*, or as encrusting masses; *conchoidal fracture*; occurs in *veins* in association with aluminous, *igneous*, or *sedimentary rocks* that have undergone *alteration*. Finer varieties are used for semi-precious stones.

**turreted (turrificate)** Applied to a gastropod (*Gastropoda*) shell that is very high-spined, with a flat or gently rounded base.



**Turreted**

**turrificate** See **TURRETED**.

**tuya** The name given in British Columbia, Canada, to a flat-topped mountain produced by the subglacial eruption of a *central vent volcano*.

**Twenhofel, William Henry** (1875–1957) An American geologist, Twenhofel was a professor at the University of Wisconsin. He made studies

of American *\*Palaeozoic \*sediments*, but is best known for his work on the processes of sedimentation, described in his book *Treatise on Sedimentation* (1926, 1932).

**twilight** Period of half-light caused by scattering and reflection of sunlight in the upper atmosphere at a time when the Sun is some degrees below the horizon.

**twin axis** An axis of a twinned crystal (see *CRYSTAL TWINNING*), normally at right angles to the *\*twin plane* and frequently related to the *\*crystallographic axis* of the crystal. The relation between the different parts of the crystal may be described with reference to a 180° rotation about the twin axis.

**twin gliding** See *DEFORMATION TWINNING*.

**twinkling** In mineral optics, the effect observed in *\*plane-polarized light* when a *\*thin section* of an *\*anisotropic* mineral with widely differing *\*refractive* indices is rotated rapidly on the *\*stage*. The ‘twinkling’ appearance is caused by the rapid changes in relief. *\*Calcite* is a well-known example.

**twin law** The law which describes the fundamental elements along or about which a crystal is twinned, in terms of a *\*twin plane* or *\*twin axis*. The various geometric shapes which result from *\*crystal twinning* may also be described with reference to well-known examples, e.g. *\*orthoclase* feldspar may be twinned on the *\*Carlsbad* law and/or the Baveno law.

**twin plane** The reflection plane which divides a twinned crystal (see *CRYSTAL TWINNING*) such that one half is a mirror image of the other. It is normally parallel to a *\*crystal face*, but in complex multiple twins it may become a highly irregular surface.

**TWINS** See *TWO WIDE-ANGLE IMAGING NEUTRAL-ATOM SPECTROMETERS*.

**two-way travel time** The time taken for a *\*seismic wave* to travel from the *\*shot* down to a *\*reflector* or refractor and back to a *\*geophone* at the surface. For finite *\*offsets*, the two-way travel times are affected by normal *\*moveout*; the normal-incidence two-way travel time is measured at zero offset.

**Two Wide-angle Imaging Neutral-atom Spectrometers (TWINS)** A two-satellite mission by *\*NASA* in collaboration with the Los Alamos National Laboratory and others to study neutral atoms and charged particles, the *\*geomagnetic* field, geomagnetic storms and substorms, *\*auroras*, the 11-year solar cycle, and the effects of space weather on human life on Earth and in space. The satellites were launched on 28 June 2006, from California, into highly elliptical orbits of *\*perigee* about 1100 km and *\*apogee* about 46 000 km (= 7.2 Earth radii), most of the period spent near apogee.

***t*-*x* curve** See TIME-DISTANCE CURVE.

***t*<sup>2</sup>-*x*<sup>2</sup> graph** A time-distance graph, used in *\*reflection* seismology, in which travel times squared (*t*<sup>2</sup>) are plotted as a function of the *\*offset* squared (*x*<sup>2</sup>) to produce straight-line segments each of which represents the reciprocal of the layer *\*root-mean-squared velocity* squared (*v*<sub>rms</sub><sup>2</sup>), and the intercept represents the depth (*z*) to the reflector. (*v*<sub>rms</sub><sup>2</sup>*t*<sup>2</sup> = 4*z*<sup>2</sup> + *x*<sup>2</sup>.) See also T-D CURVE; TIME-DISTANCE CURVE.

**Tycho** Visually, the most prominent lunar crater, 85 km in diameter, formed about 100 Ma ago. See RAYS.

**tympanic bone** In *\*Mammalia*, the bone supporting the tympanic membrane, derived from the angular bone of the lower jaw and, in many mammals, forming the bulla.

**Type I earthquake source** See SINGLE COUPLE.

**Type II earthquake source** See DOUBLE COUPLE.

**type area** The geographical region surrounding the *\*type locality*.

**type locality** The particular site where the type section (*\*stratotype*) for a *\*stratigraphic unit* is located, or where the original type section was first described. Stratigraphic units normally take the geographic component of their name from that of the type locality, or a key physical feature there, e.g. the Kimmeridge Clay from Kimmeridge Bay in Dorset, England. Some early-defined stratigraphic units were named after the *\*type area*, e.g. the *\*Cambrian* system in Wales was named by Adam *\*Sedgwick* (1835) after

*Cambria*, the latinized version of the Welsh *cymry* ('fellow countryman') and *Cymru* (Wales).

**type section** See STRATOTYPE; UNIT-STRATOTYPE; TYPE AREA; TYPE LOCALITY.

**type series** In taxonomy, all the specimens on which the description of a taxon is based.

**type specimen** See HOLOTYPE.

**typhoon** The name given to a \*tropical cyclone that forms over the Pacific, southern Indian Ocean, and China Seas.

**typological method** In stratigraphy, the describing of \*stratigraphic units in terms of the kind of rock or \*fossil they contain. In practice it is the lower-ranking units, e.g. \*zones, that are defined typologically. Compare HIERARCHICAL METHOD.

***Tyrannosaurus rex*** Giant, carnivorous \*dinosaur which lived during the Upper \*Cretaceous in N. America and possibly in Asia. Individuals grew to 12 m in length, 5 m tall, and weighed about 7 tonnes. The apt name means 'king of the tyrant lizards'.

**Tyrrhenian** A European \*stage (0.26–0.01143 Ma ago) of the \*Pleistocene \*epoch, preceded by the \*Sicilian and followed by the \*Holocene epoch.



**Ubendian orogeny** A phase of mountain building whose precise dates are uncertain but which probably occurred about 1800–1700 Ma ago, producing what is now a NW–SE belt in southern Tanzania, northern Zambia, and the eastern Democratic Republic of Congo.

**Udden–Wentworth scale** See PARTICLE SIZE.

**Udocanian** A \*stage of the Lower \*Proterozoic, from about 2600 to 2000 Ma ago (Van Eysinga, 1975), overlain by the \*Ulcanian.

**Ufimian** See ROADIAN.

**Uivakian orogeny** A phase of mountain building that occurred about 3000 Ma ago, producing what is now an approximately N–S belt in north-eastern Labrador, Canada.

**UK-DMC-2** See UNITED KINGDOM — DISASTER MONITORING CONSTELLATION-2.

**Ulatisian** A Californian \*stage (51–48 Ma ago) in the \*Eocene, underlain by the \*Penutian, overlain by the \*Narizian, and roughly contemporaneous with the upper \*Ypresian and lower \*Lutetian stages.

**Ulcanian** A \*stage of the Lower \*Proterozoic, from about 2000 to 1400 Ma ago, underlain by the \*Udocanian and overlain by the \*Burzyan (Burzyanian).

**ulexite** See BORAX.

**ulna** In tetrapods (\*Tetrapoda), the post-axial bone of the fore limb.

**Ulsterian** A \*stage (416–391.8 Ma ago) in the \*Devonian of N. America, preceded by the \*Cayugan, followed by the \*Erian, and containing the

Early Devonian *epoch*.

**ultimate bearing capacity** The *bearing capacity* that is the greatest stress that can be imposed by the weight of a building on a rock or soil without exceeding its strength and causing it to fail. Failure would cause the material to move and the foundations of the building to fail. Because engineers can never be certain of knowing the ultimate bearing capacity ( $q_f$ ) precisely, they divide it by a safety factor ( $F$ ) to produce the *safe bearing capacity* ( $q_s$ ).

**ultimate strength (failure strength)** The maximum *stress* that can be maintained in a body prior to its rupture. It is marked by the highest point on a curve in a *stress–strain diagram*.

**ultisols** Mineral soils, an order in the *USDA soil taxonomy* identified by an *argillic B soil horizon* with a *base saturation* of less than 35%, and red in colour from iron oxide concentration. Ultisols are leached, *acid soils*, associated with humid subtropical environments. *See also RED PODZOLIC SOILS*.

**ultrabasic rock** An *igneous* rock that consists almost entirely of *ferromagnesian minerals* and possesses no free *quartz*, and with less than 45% *silica* ( $\text{SiO}_2$ ). ‘Ultramafic’ is a partial synonym.

**ultramafic** *See ULTRABASIC*.

**ultramylonite** *See DYNAMIC METAMORPHISM*.

**ultraplinian eruption** The most extreme type of *Plinian eruption*, in which the column of ejecta reaches a height of more than 45 km. Such an eruption, with a column up to 50 km high, occurred near Taupo, North Island, New Zealand, in about 186 CE.

**ultraviolet radiation** *Electromagnetic radiation* which has a wavelength between 0.5 nm and 400 nm, located between the visible and X-ray regions of the electromagnetic spectrum. Near ultraviolet occurs at wavelengths between 400 nm and 300 nm, middle ultraviolet between 300 nm and 200 nm, and extreme ultraviolet between 200 nm and 150 nm.

**ultraviolet spectrometer (ultraviolet spectrophotometer; UVS)** A *spectrometer*, used in remote sensing, that measures spectra in the UV

waveband from which information can be derived concerning planetary atmospheres and surfaces.

**ultraviolet spectrophotometer** See ULTRAVIOLET SPECTROMETER.

**ultraviolet–visual spectrophotometry (UV–Vis spectrophotometry)**

Method of spectrophotometric analysis wherein the absorbance (or transmission) of a coloured complex is measured at a specific wavelength in the ultraviolet or visible part of the electromagnetic *\*spectrum*. Any metal or *\*ion* which can be made to form a soluble, coloured complex with appropriate absorption maxima may be quantitatively determined by comparing the absorption of the unknown material with that of standards of appropriate *\*concentration*.

**Ulysses** A *\*NASA* and *\*ESA* probe, launched in 1990, carrying the International Solar Polar Mission to fly over the poles of the Sun, which cannot be seen from Earth, and to study the *\*solar* wind. It travelled first to *\*Jupiter* for a gravity assist. The mission ended on 30 June 2009.



<https://solarsystem.nasa.gov/missions/ulysses/in-depth/>

- Ulysses.

**umber** A *\*mudstone* rich in iron and manganese oxides, with *\*silica*, aluminium oxide, and *\*lime*, that is the source of a pigment of the same name. Compare OCHRE.

**umbilicus** The central cavity of a gastropod (*\*Gastropoda*) shell where the later *\*whorls* do not meet centrally; it is sometimes seen basally as an opening, but is also commonly closed by shell material. In cephalopods (*\*Cephalopoda*), an open space that remains in the axis of coiling when successive whorls do not reach the axis.

**umbo** (*pl.* **umbones**) The first part of a brachiopod (*\*Brachiopoda*) or bivalve (*\*Bivalvia*) shell to be formed. In a brachiopod, the umbo is the posterior part of each *\*valve*; in a bivalve it forms the *\*dorsal* part of the shell.



## Umbo

**umbones** See **UMBO**.

**umbric epipedon** Surface **\*soil horizon** similar to a mollic epipedon but with a **\*base saturation** of less than 50%. It is a **\*diagnostic horizon**.

**umbric horizon** In the **\*World Reference Base for Soil Resources** classification scheme, a **\*soil horizon** that is thick, dark-coloured, with a **\*base saturation** of less than 50% averaged throughout the horizon, and rich in organic matter.

**Umbriel (Uranus II)** One of the major satellites of **\*Uranus**. Its mean radius is 584.7 km; mass  $11.72 \times 10^{20}$  kg; mean density  $1400 \text{ kg/m}^3$ ; visible albedo 0.18. The surface is evenly cratered and darker than the other four major satellites, except for a bright ring 4 km in diameter. The many large craters suggest the surface is old.

**umbrisols** A reference soil group in the **\*World Reference Base for Soil Resources** classification scheme. Umbrisols are soils with an **\*umbric horizon**.

**unavailable water** Water that is present in the soil but that cannot be absorbed by plants because it is held so strongly to the surface of soil


particles.

**uncinus** From the Latin *uncinus* meaning 'hooked', a species of **\*cirrus** cloud that is hooked or tufted at the end of its filaments. *See also* CLOUD CLASSIFICATION.

**unconfined aquifer** *See* AQUIFER.

**unconfined compressive strength (uniaxial compressive strength)** The strength of a **\*rock** or **\*soil** sample when crushed in one direction (uniaxial) without lateral restraint. *See also* UNIAXIAL COMPRESSION TEST.

**unconformity** Surface of contact between two groups of unconformable **\*strata**, which represents a hiatus in the geologic record due to a combination of **\*erosion** and a cessation of sedimentation. *Compare* DIASTEM. *See also* ANGULAR UNCONFORMITY; DISCONFORMITY.



**Unconformity**

**unconformity trap** A *\*stratigraphic trap* formed by *\*folding*, uplift, and *\*erosion* of porous *\*strata*, followed by the deposition of later beds which can act as a seal for oil, gas, or water. Although common structures, these traps contain only 4% of the world's oil, perhaps because of losses that occur during uplift and erosion. See NATURAL GAS; PETROLEUM; POROSITY. Compare ANTICLINAL TRAP; FAULT TRAP; REEF TRAP; STRATIGRAPHIC TRAP; STRUCTURAL TRAP.

**unconformity-type uranium ore** Uranium *\*ore* that is associated with an *\*unconformity*. Uranium is often concentrated in *\*sedimentary rocks* or precipitated from *\*groundwater*, because uranium minerals are readily decomposed under oxidizing conditions. These deposits are all of early *\*Proterozoic* age. In Ontario, uranium is concentrated in the basal bed of the Lorrain Quartzite, which is unconformable on the *\*Archaean* granite, the mineralization being about 1 m thick and covering hundreds of square kilometres.

**unconsolidated** Applied to particles that are loose and not cemented together.

**undae** On the surface of *\*Mars*, *\*Titan*, and *\*Venus*, a field of dunes.

**undaform** The approximately level surface beneath a water body, at a level higher than that of the wave bases so its sediments are stirred by wave action and currents, that gives a seismic reflection.

**undathem** The rock unit produced beneath a water body at a depth higher than the bases of waves, that gives a seismic reflection.

**undepleted mantle** Primary *\*mantle* material from which *\*basalt* has not been extracted. Compare DEPLETED MANTLE.

**undercliff** A stretch of land that lies parallel with and below a major cliff. A good example is the Undercliff of the Isle of Wight, Britain, between Bonchurch and Blackgang, which is overlooked by a conspicuous cliff in Upper Greensand rocks.

**undercooling** The state whereby a liquid must be cooled to well below its *\*solidus* temperature before *\*nucleation* and crystallization are initiated. See SUPERCOOLING.

**underfit stream** See MISFIT STREAM.

**underflow** The flow of \*groundwater in \*alluvial \*sediments, parallel to and beneath a river channel. It forms a significant fraction of the total river flow in coarse gravel alluvium.

**underplating** The addition of material to the underside of a geologic unit, as in the movement of \*crust by \*A-subduction in a \*collision zone so that it underlies the overriding continent. The term has also been used to describe the addition of gabbroic \*plutons beneath \*lavas and \*dykes in a \*constructive margin.

**undersaturated** See SILICA SATURATION.

**underthrusting** The movement of rock such that the lower block moves under a relatively passive upper block, with the plane of contact being gently inclined. Underthrusting is thought to be of major importance in the formation of \*accretionary wedges.

**undertow** The general seaward flow of water beneath individual breaking waves, in contrast to the more localized \*rip-current return flow.

**Undillian** An Australian \*stage (506–504 Ma ago) of the Middle \*Cambrian, preceded by the \*Floridan and followed by the \*Boomerangian.

**undrained test** See TRIAXIAL COMPRESSION TEST.

**undulatus** From the Latin *undulatus* meaning ‘waved’, a variety of cloud whose layers undulate. See also CLOUD CLASSIFICATION.

**undulose extinction** In optical \*mineralogy, the irregular or wavy \*extinction seen in \*thin sections of \*mineral grains (e.g. \*quartz) on rotation of the \*stage. Different parts of the grain go into extinction in different orientations. The phenomenon is thought to result from strain, which may modify crystal orientation slightly.

**ungulate** 1. Any hoofed, grazing mammal, which usually is also adapted for running. Hoofed mammals occur in several mammalian groups (\*Mammalia), and the term ‘ungulate’ no longer has any formal taxonomic use. 2. Hoof-shaped.

**uniaxial compression test** Test of a \*soil or \*rock sample for \*unconfined compressive strength. An undisturbed sample is loaded from above until it

fails. The results depend on the length-to-breadth ratio of the sample and the loading rate.

**uniaxial compressive strength** See UNCONFINED COMPRESSIVE STRENGTH.

**uniaxial interference figure** See INTERFERENCE FIGURE.

**Unibothriocidaris** See BOTHRIOCIDAROIDA.

**unicarinate** Possessing a single carina (*\*keel*). It may occur as a structure along the *\*venter* of a cephalopod (*\*Cephalopoda*), round each *\*whorl* of a gastropod (*\*Gastropoda*), and may also occur in other groups.

**Unified Stratigraphic Timescale (UTS)** A proposed universal timescale that would combine both absolute and relative timescales (see GEOCHRONOLOGY) that have been deduced by all possible means, including *\*biostratigraphy*, *\*chronostratigraphy*, *\*stratigraphy*, *\*magnetostratigraphy*, and *\*radiometric dating*.

**UNIFORM-1** See UNIVERSITY INTERNATIONAL FORMATION MISSION-1.

**uniform flow** **1.** In channels, a flow whose velocity and *\*discharge* do not vary along the channel's length. **2.** In *\*groundwater*, a flow whose velocity and direction are the same at all points in the field of flow. Non-uniform flow in channels results from changes in the cross-sectional shape or discharge along the channel's course.

**uniformitarianism (actualism)** The principle proposed by James *\*Hutton* and paraphrased succinctly as 'the present is the key to the past', i.e. that the formations and structures we see today were produced by processes identical to those operating now. This is a considerable oversimplification, since processes that occurred in historical times may not be occurring now, or may not be observable now, and vice versa.

**unilocular** Applied to the condition in which the *\*test* of a foraminiferid (*\*Foraminiferida*) consists of a single chamber. Where the test consists of two or more chambers it is said to be 'multilocular'.

**uniramous** Of an appendage, unbranched.

**uniserial** **1.** Arranged in a single row. **2.** See TRISERIAL.

**unit cell** See BRAVAIS LATTICE.

**United Kingdom – Disaster Monitoring Constellation-2 (UK-DMC-2)** A second-generation satellite mission working in conjunction with *\*Deimos*, *\*NigeriaSat-2*, and *\*NigeriaSat-X* to provide images of large areas of territory. UK-DMC-2 and Deimos-1 were launched on 29 July 2009, from Baikonur, Kazakhstan, into slightly eccentric, Sun-synchronous orbits, Deimos-1 with *\*perigee* at about 642 km and UK-DMC-2 with perigee at about 632 km.

**United States Department of Agriculture** See *USDA*.

**United States Geological Survey (USGS)** The civilian agency responsible for mapping the United States and for monitoring natural resources. It gathers and analyses data and supplies scientific information and advice to those requiring it. The USGS was established in 1879.



<http://www.usgs.gov/>

- Home page of the USGS.

**unit form** See *PARAMETER*; *PARAMETRAL PLANE*.

**unit hydrograph** See *HYDROGRAPH*.

**unit-stratotype** The standard succession of *\*strata*, selected from the rock successions within a particular locality (the *\*type locality*), that is designated and described as the standard type section for a particular *\*chronostratigraphic* or *\*lithostratigraphic* unit. The unit-stratotype provides the reference against which all other examples of that stratigraphic unit anywhere in the world may be examined and identified. See *STRATOTYPE*.

**unit stress** The amount of force applied to a unit area. The term is most commonly used in engineering geology.

**univariant assemblage** An equilibrium mineral assemblage which, in terms of the *\*phase rule*, has one *\*degree* of freedom; i.e. if equilibrium is to be maintained, a change of any one variable (e.g. pressure) dictates the changes to the values of all the other external variables (temperature and  $P(\text{H}_2\text{O})$ ).

**universal soil loss equation** An equation that is used to predict the extent of *\*soil \*erosion*, principally on farm land:  $\text{soil loss} = R \times K \times LS \times P$ , where  $R$  is an index of rainfall,  $K$  is an index of the susceptibility of the soil to erosion,  $LS$  is an index of slope length ( $L$ ) and steepness ( $S$ ), and  $P$  is an index of conservation measures that will be implemented.

**University International Formation Mission-1 (UNIFORM-1)** A *\*microsatellite* mission by a consortium of universities, funded by the Japanese government, mainly for monitoring forest fires. The microsatellite was launched on 24 May 2014, from the Yoshinobu Launch Complex, Japan, into a Sun-synchronous near-circular orbit at an altitude of 628 km.

**unloading joint** See JOINT.

**unsaturated zone** See VADOSE ZONE; SOIL-WATER ZONE.

**unsaturation index ( $U^{K'}_{37}$ )** An index of di- versus tri-unsaturated  $C_{37}$  *\*alkenones* that is used to estimate past water temperature. The index is calculated as:  $U^{K'}_{37} = C_{37.2}/(C_{37.2} + C_{37.3})$ . Temperature for most marine environments is then given by:  $U^{K'}_{37} = 0.033 T (\text{°C}) + 0.044$ . A different calibration is used for samples from high latitudes and *\*lacustrine* environments.

**unsorted biosparite** See FOLK LIMESTONE CLASSIFICATION.

**unsteady flow** See STEADY FLOW.

**uphole survey** Technique in which seismic sources are energized within a *\*borehole* and arrival times recorded by surface *\*geophones*. It is used particularly to determine weathered-layer velocity.

**uphole time** The time taken for the first direct wave to travel from the *\*shot* fired at the base of a shot-hole to a *\*geophone* at or near the top of the shot-hole.

**upright fold** A *\*fold* whose *\*axial plane* or surface is vertical or nearly so.

**upslope fog** Condensation near the surface in uplifted air on windward slopes.

**upthrow** The relative upward *\*displacement* on one side of a *\*fault*.

**upthrust** Applied to a block that has been moved upwards in a **\*reverse fault**.

**Upton Warren** A warm **\*interstadial** during the Middle **\*Devensian**, between 43 000 and 42 000 years BP, when faunal and floral evidence suggest the landscape was devoid of trees.

**upward continuation** The use of one set of measurements of a potential field (usually gravity or magnetic) over one surface to determine the field at a higher surface. Upward continuation is relatively reliable, in that the field is continued into free space where there are no causative bodies to perturb the field further (unlike **\*downward continuation**). The method effectively attenuates high-**\*wavenumber** anomalies due to near-surface features, thus providing a powerful method for examining deeper structures. Upward continuation is used in gravity surveys to determine the nature of the regional gravity pattern over a large area; in magnetic surveys it is particularly useful in tying together aeromagnetic surveys made from different flying altitudes.

**upward lightning** A **\*transient luminous** event consisting of a lightning flash that rises upward from the top of a thunderstorm cloud, does not flicker, and lasts up to 5 seconds.

**upwelling** In oceans or larger lakes, a water current, or movement of surface water produced by wind and a resulting Ekman spiral, which brings colder water, loaded with nutrient, to the surface from a lower depth. Ocean upwellings occur off Peru, California, W. Africa, and Namibia, and increase the nutrient content of the surface waters, leading to an abundance of marine and bird life. Upwelling also occurs in the open oceans where surface currents diverge, as deep waters rise to the surface to replace the departing waters, and all along the equator as a result of the effects of the NE and SE **\*trade winds**.

**Ur** A postulated **\*supercontinent** that may have existed about 3000 Ma ago, comprising parts of what are now Africa, Australia, and India.

**Uralian orogeny** A phase of mountain building, affecting what is now a N–S belt in Russia, that occurred at the same time as the **\*Hercynian orogeny**, in the late **\*Devonian** and early **\*Carboniferous**.

**uralite** See URALITIZATION.

**uralitization** The alteration of \*primary \*igneous \*pyroxene to a fibrous mass of \*amphibole, usually \*hornblende, during late-stage \*hydrothermal activity, or during low-grade \*metamorphism of the igneous rock. The amphibole was originally thought to be a distinct \*mineral type known as 'uralite'.

**Ural Sea (Obik Sea)** A \*Palaeocene–\*Eocene seaway extending from the present-day Caspian Sea in the south to the Arctic Ocean in the north, covering the area immediately east of the Ural Mountains. The area is now covered by the extensive plain across which flows the River Ob.

**uranian satellites** See ARIEL (URANUS I); BELINDA (URANUS XIV); BIANCA (URANUS VIII); CALIBAN (URANUS XVI); CORDELIA (URANUS VI); CRESSIDA (URANUS IX); CUPID (URANUS II); DESDEMONA (URANUS X); FERDINAND (URANUS XXIV); FRANCISCO (URANUS XXII); JULIET (URANUS XI); MAB (URANUS I); MARGARET (URANUS XXII); MIRANDA (URANUS V); OBERON (URANUS IV); OPHELIA (URANUS VII); PERDITA (URANUS XXV); PORTIA (URANUS XII); PROSPERO (URANUS XVIII); PUCK (URANUS XV); ROSALIND (URANUS XIII); SETEBOS (URANUS XIX); STEPHANO (URANUS XX); SYCORAX (URANUS XVII); TITANIA (URANUS III); TRINCULO (URANUS XXI); UMBRIEL (URANUS II).

**uraninite (pitchblende)** Mineral,  $UO_2$ ; sp. gr. from 8.5 when \*massive to 10 for unaltered crystals; \*hardness 5–6; \*cubic; it is brownish-black or grey-black; brownish-black \*streak; sub-metallic, greasy \*lustre; crystals very rare, the mineral is usually found as \*botryoidal masses; no \*cleavage; radioactive; occurs in \*pegmatites, and in association with \*monazite, \*zircon, and \*tourmaline, in \*hydrothermal veins together with \*cassiterite, \*pyrite, \*chalcopyrite, and \*galena, and as a \*detrital material in \*alluvials. It is a major \*ore mineral for uranium metal.

**uranium deposit** \*Ore deposit containing more than 350 ppm uranium. Six types are recognized: pegmatitic (see PEGMATITE) together with \*disseminated magmatic deposits; hydrothermal veins and \*stockworks (see HYDROTHERMAL ACTIVITY; HYDROTHERMAL MINERAL); vein-unconformity deposits (see VEIN DEPOSIT); \*sandstone deposits; \*placer deposits; and phosphatic \*limestones and black-\*shale deposits.

**uranium–lead dating** All naturally occurring uranium contains  $^{238}\text{U}$  and  $^{235}\text{U}$  (in the ratio 137.7:1). Both **\*isotopes** are the starting points for complex **\*decay series** that eventually produce stable isotopes of lead.  $^{238}\text{U}$  decays to  $^{206}\text{Pb}$  (half-life = 4510 Ma, see **DECAY CONSTANT**) by a process of eight **\*alpha-decay** steps and six **\*beta-decay** steps.  $^{235}\text{U}$  decays to  $^{207}\text{Pb}$  (half-life = 713 Ma) by a similar series of stages that involves seven alpha-decay steps and four beta-decay steps. Also included within this range of methods is that for **\*thorium–lead dating** ( $^{232}\text{Th}$  to  $^{207}\text{Pb}$ ; half-life = 13 900 Ma). Uranium–lead dating was applied initially to uranium minerals, e.g. **\*uraninite** and **\*pitchblende**, but as these are rather restricted in occurrence it is more normal to use the mineral **\*zircon**, even though the uranium is present only in trace amounts. The amount of radiogenic lead from all these methods must be distinguished from naturally occurring lead, and this is calculated by using the ratio with  $^{204}\text{Pb}$ , which is a stable isotope of the element then, after correcting for original lead, if the mineral has remained in a closed system, the  $^{235}\text{U}:^{207}\text{Pb}$  and  $^{238}\text{U}:^{206}\text{Pb}$  ages should agree. If this is the case, they are **\*concordant** and the age determined is most probably the actual age of the specimen. These ratios can be plotted to produce a curve, the Concordia curve (see **CONCORDIA DIAGRAM**). If the ages determined using these two methods do not agree, then they do not fall on this curve and are therefore **\*discordant**. This commonly occurs if the system has been heated or otherwise disturbed, causing a loss of some of the lead daughter atoms. Because  $^{207}\text{Pb}$  and  $^{206}\text{Pb}$  are chemically identical, they are usually lost in the same proportions. The plot of the ratios will then produce a straight line below the Concordia curve. G. W. Wetherill has shown that the two points on the Concordia curve intersected by this straight line will represent the time of initial crystallization and the time of the subsequent lead loss.

**uranium–protactinium dating (protactinium dating)** A method of radiometric dating based on the decay of uranium–235 ( $^{235}\text{U}$ ) to protactinium–231 ( $^{231}\text{Pa}$ ) that is used to date sediments as young as ten years old up to 250 000 years.

**uranium series** See **DECAY SERIES**.

**Uranus** The seventh planet in the *\*solar* system, discovered in 1781 by Sir William Herschel (1738–1822), although he described it as a comet. It was named Uranus by J. E. *\*Bode*. Its equatorial radius is 25 559 km and polar radius 24 973 km; volume 6833 km<sup>3</sup>; mass  $86.83 \times 10^{24}$  kg; mean density 1318 kg/m<sup>3</sup>; visual albedo 0.51; black-body temperature 35.9 K. The inclination of the equator to the plane of the *\*ecliptic* is 97.86°, so the planet is lying on its side (a fact discovered in 1846 by Johann Gottfried Galle (1812–1910)). At its closest approach, Uranus is  $2581.9 \times 10^6$  km from Earth and at its furthest  $3157.3 \times 10^6$  km. Uranus has an atmosphere, with a surface atmospheric pressure well in excess of 10 MPa. The atmosphere is composed of molecular hydrogen (89%) and helium (11%), with *\*aerosols* of methane, ammonia ice, water ice, ammonia hydrosulphide, and possibly methane ice (similar to that of *\*Neptune*). Wind speeds at the surface are 0–200 m/s and the average surface temperature is about 58 K. The total number of known satellites is twenty-seven (see *URANIAN SATELLITES*), and it is likely that more remain to be discovered. Except for Titan, the uranian satellites are denser than those of *\*Jupiter*. Oberon and Titania, the two largest, were discovered in 1787 by Sir William Herschel. Umbriel and Ariel were discovered in 1851 by William Lassell (1799–1880). Miranda was discovered in 1948 by Gerard *\*Kuiper*. Ariel, Oberon, and Titania are probably made of water ice, other ices, and silicates. They are believed to be too cold to have a molten core, but on some there are signs of geological activity. The remaining satellites were revealed in images transmitted to Earth from *\*Voyager 2*.

**urban climate** Modified surface-layer atmospheric conditions caused by the influence of large ‘built-up’ areas. Changes include pollution, reduction in strong wind speeds toward a city centre, *\*turbulence* of air around buildings, warming of air by the heat output from city structures, and increased evaporation and removal (drainage and runoff) of water.

**urban dome** See *URBAN HEAT ISLAND*.

**urban heat island** Air over an urban area that is markedly warmer than air in the surrounding countryside and that is held in an approximately dome-shaped *\*temperature inversion* (urban dome). The inversion tends to trap pollutants. See *URBAN CLIMATE*.

**Urey, Harold Clayton** (1893–1981) An American professor of chemistry from California, Urey is best known as the discoverer of deuterium, but he also did important work on the origin and evolution of the universe and of life. In the field of Earth science he developed the use of **\*oxygen-isotope analysis**, used to indicate ancient climatic temperatures.

**Uriconian** A **\*stage** of the Upper **\*Proterozoic** of Shropshire, England, underlain by the **\*Malvernian** and overlain by the **\*Charnian**.

**Urochordata (tunicates)** (phylum **\*Chordata**) Subphylum comprising the sea-squirts, with reliable **\*fossils** from the **\*Permian**, and possible remains in **\*Silurian \*sediments**. Sea-squirts have tadpole-like larvae with **\*notochords** in their tails, a feature that links them with the chordates.

**Ursidae** See **CARNIVORA**.

**urtite** A coarse-grained, **\*igneous** rock consisting of **\*essential \*nepheline** (about 85% of the rock) and **\*ferromagnesian minerals** (**\*aegirine**, aegirine-**\*augite**, and soda-iron **\*amphibole**). The rock is named from Lujaur-Urt, in the Kola Peninsula, Russia, and is a type of **\*undersaturated \*syenite**.

**Ururoan** A New Zealand **\*stage** (188–169 Ma ago) of the **\*Jurassic**, preceded by the **\*Aratauran** and followed by the **\*Temaikan**.

**Urutawan** A New Zealand **\*stage** (110–104 Ma ago) of the Early Cretaceous epoch, preceded by the **\*Korangian** and followed by the **\*Motuan**.

**USDA (United States Department of Agriculture)** The department of the United States federal government that exists to serve the needs of those engaged in agriculture and rural communities. It comprises a number of agencies, each with its own functions, one of which is the US Soil Survey. In 1960, the USDA published Soil Classification: A Comprehensive System, prepared by the Soil Survey. This method of classifying soils hierarchically was renamed the US Soil Taxonomy in 1970, and was the most widely used taxonomic system prior to the introduction of the international soil classification developed by the **\*FAO**. See **SOIL TAXONOMY**; **WORLD REFERENCE BASE FOR SOIL RESOURCES**.

**UTS** See **UNIFIED STRATIGRAPHIC TIMESCALE**.

**UV** Abbreviation for ultraviolet.

**uvala** An irregularly shaped hollow in a **\*karst** terrain. It is generally 500–1000 m in diameter and may be 100–200 m deep. It is the result of the coalescence of a number of **\*dolines**.

**uvarovite** Member of the **\*garnet** group  $\text{Ca}_3\text{Cr}_2\text{Si}_3\text{O}_{12}$ ; sp. gr. 3.9; **\*hardness** 7.0–7.5; emerald green; occurs as **\*cubic** crystals and **\*massive**; found in **\*serpentinite** in association with **\*chromite**, and in metamorphosed **\*limestones** and skarn ores.

**UVS** *See* ULTRAVIOLET SPECTROMETER.

**UV–Vis spectrophotometry** *See* ULTRAVIOLET–VISUAL SPECTROPHOTOMETRY.

**[UVW]** *See* ZONE SYMBOL.



**V** The *\*vertical* component of a magnetic vector.

**Vaalbara** See PALAEOARCHAEAN.

**Vaalian** A South African *\*stage* (2500–2050 Ma ago), preceded by the *\*Randian* and followed by the *\*Mokolian*.

**vadose zone** See PHREATIC ZONE; SOIL-WATER ZONE.

**Valanginian** A *\*stage* in the European Early *\*Cretaceous* *\*epoch* (140.2–136.4 Ma ago, Int. Commission on Stratigraphy, 2004), preceded by the *\*Berriasian* and followed by the *\*Hauterivian*, for which the *\*type locality* is at Valangin, Switzerland. See also NEOCOMIAN.

**Valdayan/Zyryanka** A *\*Weichselian*, *\*loess-like* *\*silt* from Siberia, in which there is a threefold division with the Zyryanka *\*drift* at the base, then Karginsky *\*interstadial* *\*sediments*, and finally the Sartan drift.

**Valdonnian** A regional *\*stage* (83.5–77 Ma ago) of the Late *\*Cretaceous*, preceded by the *\*Santonian* and followed by the *\*Fuvelian*.

**valency** The bonding potential of an atom, measured by the number of hydrogen *\*ions* (valency 1) that the atom could combine with or replace. In an ionic compound the valency (electrovalency) equals the ionic charge on each ion, e.g. in the compound MgO, Mg<sup>2+</sup> shows a valency of +2, O<sup>2-</sup> a valency of –2. In a covalent compound the valency (covalency) of an atom is equal to the number of bonds it forms, e.g. in CH<sub>4</sub> carbon has a valency of 4, hydrogen a valency of 1.

**valley bog** A peat-forming plant community that develops in badly drained hollows, wet valley bottoms, and valleys that are partly obstructed downstream. Many valley bogs have layers of charcoal beneath the peat,

suggesting that fire may have triggered peat formation by forming an impermeable charcoal layer that caused waterlogging.

**valley bulging** An upward arching of the bedrock along the axis of a valley. It may not be visible at the ground surface due to subsequent *\*erosion*, but it is revealed by the distortion of the geologic structure. It may be due to *\*frost heave*, or to the compressive forces set up when two opposing valley sides approach each other.

**valley glacier** A long, relatively narrow ribbon of ice that is confined between valley walls. The Alpine type is fed by a series of *\*cirque glaciers* that show positive net balances (see *MASS BALANCE*), and is common in the Alps and in the coastal mountains of Alaska, USA. The outlet type is fed by an *\*ice cap* or *\*ice sheet*. The Vatnajökull ice cap in Iceland feeds several outlet glaciers.

**valley-side bench (rock bench)** A terrace-like land-form standing on the flank of a valley, but lacking a veneer of *\*alluvium*. It may have originated as a true *\*river terrace* from which the alluvium has been stripped by subsequent *\*erosion*, or it may simply result from the exposure of a nearly level *\*stratum* of resistant rock.

**valley train** An accumulation of fluvio-glacial deposits laid down in a valley by meltwaters escaping from a decaying *\*glacier*. The surface slopes quite steeply down-valley, and is incised by shifting *\*braided streams*.

**valley wind** An *\*anabatic* wind that blows up-valley during the day in otherwise calm conditions, or a *\*katabatic*, down-valley, night wind. See also *MOUNTAIN WIND*; *RAVINE WIND*.

**vallis** A general term for a valley on a planetary or *\*satellite* surface. The largest example is the Valles Marineris system on *\*Mars*. Mostly the term is confined to smaller, sinuous valleys probably of fluvial origin.

**valve** **1.** One of the two halves of the hinged shell of brachiopods (*\*Brachiopoda*), or molluscs (*\*Mollusca*) of the class *\*Bivalvia*. **2.** One half of the cell wall of a *\*diatom*. **3.** A flap or other constriction that can close to ensure that a fluid flows in only one direction.

**vanadinite** Mineral  $Pb_5(VO_4)_3Cl$  and member of the pyromorphite series with *\*pyromorphite* ( $Pb_5(PO_4)_3Cl$ ) and mimetite ( $Pb_5(AsO_4)_3Cl$ ); sp. gr.

6.88–6.93; **\*hardness** 2.5–3.0; **\*hexagonal**; orange-red, brownish-red, to yellow; white to yellowish **\*streak**; resinous **\*lustre**; crystals sharp, hexagonal **\*prisms**, but occasionally rounded; no **\*cleavage**; found in the oxidized zone of sulphide-ore deposits carrying lead minerals. It can be used as a source of vanadium.

**Van Allen belts** Two belts of high-energy, charged particles trapped by the **\*Earth's** magnetic field within the **\*magnetosphere**. The inner belt starts at an altitude of about 800 km and reaches a maximum intensity at about 2000 km. The outer belt reaches a maximum intensity at between three and four Earth radii (18 000–25 000 km). Although commonly referred to as 'radiation belts', they contain particles, not radiation. Their presence was predicted by James Alfred Van Allen (1914–2006).

**Van Allen Probes** A **\*NASA** mission, formerly known as the Radiation Belt Storm Probes (RBSP) Mission, to study Earth's radiation belts in order to explore the Sun's influence on Earth and near-space. The two identical satellites were launched on 30 August 2012, from Cape Canaveral, into near-equatorial, highly elliptical orbit with a **\*perigee** at about 620 km and **\*apogee** at about 30 500 km (= about 5.8 Earth radii), inclination 10°.

**van der Waals force** The weak attraction of atoms to each other due to the interaction of **\*electrons** and **\*nuclei**. The resultant linkage is called the van der Waals bond (named after the Dutch physicist Johannes van der Waals, 1837–1923).

**vane** A device for indicating the direction of wind. The standard exposure for wind vanes is on a mast at 10 m above unobstructed ground.

**vane test** Test of the shearing strength of **\*soil in situ**. The equipment consists of four thin, rectangular blades which project from a shaft at right angles. This is pressed into the ground and rotated at a uniform rate. The torque required to generate a cylinder of soil determines the shearing strength of the soil.

**van't Hoff, Jacobus Henricus** (1852–1911) A Dutch chemist, and winner of the 1901 Nobel Prize for chemistry for his work on the relationship between osmotic pressure and vapour pressure, van't Hoff was professor of chemistry, mineralogy, and geology at Amsterdam University and later an honorary professor at the Prussian Academy of Sciences. His work on

phase equilibria (see [PHASE DIAGRAM](#)), involving six components of sea water at a range of temperatures, contributed much to the study of sedimentary processes and in particular provided a theoretical basis for understanding the formation of salt deposits.

**vapour-phase crystallization** The crystallization of minerals from hot gases escaping through a volcanic body. Cooling of the escaping gases, which carry elements in solution, promotes the crystallization of minerals in rock cavities or the spaces between [\\*pyroclastic \\*clasts](#) in [\\*ignimbrites](#). Minerals such as [\\*feldspar](#) and [\\*quartz](#) are common vapour-phase crystallization products.

**vapour pressure** The pressure exerted by molecules of a substance in the vapour state, at equilibrium with molecules of the same substance in the liquid state, within a closed container. The magnitude of the vapour pressure exerted depends on the temperature and the identity of the liquid; it does not depend on the amount of liquid in the container. The saturated vapour pressure of water at 0 °C is 610 N m<sup>-2</sup>, rising to 2340 N m<sup>-2</sup> at 20 °C and 7380 N m<sup>-2</sup> at 40 °C. See also [PARTIAL PRESSURE](#).

**vapour-pressure curve** See [DEHYDRATION CURVE](#).

**VAR** See [VARIABLE-AREA DISPLAY](#).

**Varanger** See [VARANGIAN](#).

**Varangian (Varanger)** A [\\*Neoproterozoic \(Cryogenian\) \\*stage](#) from about 650–630 Ma, preceded by the [\\*Sturtian](#) and followed by the [\\*Wonokan](#).

**vardar (vardarac)** A type of [\\*ravine wind](#), which blows in the Moravia–Vardar valley, bringing cold conditions from the north to the Thessaloniki area of Greece.

**variable-area display (VAR)** A way of displaying the wave-form on a [\\*seismic record](#), in which the positive component of the wave is shown blacked in, and the negative-polarity component is absent, unless the VAR display is in the [\\*wiggle-trace](#) mode.



Variable-area display

**variance (mean square)** In statistics, a measure of the spread of the data about the mean. In a set of data, the square of the mean variation about the mean value. It is calculated as the mean of the squared data points minus the mean squared; the standard deviation of a data set is the square root of the variance:  $s^2 = \left( \sum_{i=1}^{i=n} (x_i - \bar{x})^2 \right) / n$  where  $s^2$  is the variance,  $x_i$  is the mean value of the  $i$ th measurement of  $x$ ,  $\bar{x}$  is the mean value of  $x$ , and  $n$  is the number of measurements.

**variolitic** Applied to a spherulitic **texture** consisting of fine, radiating fibres of **plagioclase** or **pyroxene** that is found in the glassy, chilled margin of shallow-level, **basic**, **igneous intrusions** (**dykes** and **sills**) or the glassy **groundmass** of some quenched (see **QUENCHING**) basaltic **lavas**.

**varve** A banded layer of **silt** and **sand** deposited annually in lakes, especially near to **ice** sheets. The coarse, paler material is deposited in summer; the finer, darker material in winter. One varve consists of one light band and one dark band. Varves can be counted to calculate the age of glacial deposits (varve analysis, also called varve chronology or varve

count). Since the pattern of thicknesses of successive varves is often distinctive, correlations can be made between widely separated deposits, using the same principle as that of *\*dendrochronology*.

**varve analysis** See VARVE.

**varve chronology** See VARVE.

**varve count** See VARVE.

**vascular plants** See TRACHEOPHYTA.

**vastitas** (*pl. vastitates*) An extensive plain on the surface of an extraterrestrial body.

**vastitates** See VASTITAS.

**veering** A clockwise shift in the direction of the wind. The reverse change is called backing.

**Vega** Two Soviet missions to *\*Venus* and comet *\*Halley*, both of which were launched in 1984.



<https://stardust.jpl.nasa.gov/comets/vega.html>

- Vega 1 & 2.

**Vegetation and Environment monitoring on a New MicroSatellite (VENμS)** A joint *\*minisatellite* (despite the name) mission between the French and Israeli space agencies to monitor surface vegetation. It was launched on 2 August 2017, from Kourou, French Guiana, into a Sun-synchronous orbit at an altitude of 720 km.



<https://cnes.fr/en/web/CNES-en/3766-joint-french-israeli-vens-mission.php>

- Joint French–Israeli Venus Mission.

**vegetation index** In *\*remote sensing*, a technique to show vegetation in an image. An example is the normalized vegetation index, which calculates how far *\*pixels* in an image plot from the *\*soil line* by dividing the very-

**\*near infrared \*digital** number value by the red digital number value of a multispectral data set.

**vein** A tabular deposit of minerals occupying a fracture, in which particles may grow away from the walls toward the middle.

**vein deposit** An **\*orebody** that is confined within a sheet-like structure, as a result of magmatic activity or deposition from circulating **\*groundwater**. The deposit is usually narrow but persistent. It may pinch and swell and has well-defined walls.

**velocity-depth distribution** The variation that occurs in **\*seismic velocity** with increasing depth, from the surface of the Earth to the **\*core**; this reveals marked discontinuities and boundaries (such as the **\*Mohorovičić** discontinuity, **\*low-velocity zone**, and core–**\*mantle** boundary).

**velocity inversion** See **INVERSION**.

**velocity log** The record of a **\*sonic sonde** while it is being raised through a **\*borehole**.

**velocity profile** The variation of water velocity with vertical distance from the bed of a river, or of wind velocity with distance from the ground. See **VON KARMAN–PRANDTL EQUATION**.

**velocity survey** A series of measurements designed to provide information about the variation and distribution of **\*seismic velocities** throughout a series of media, as a function of depth and horizontal distance. A number of velocity analyses along a **\*seismic section** can be displayed to produce an isovelocity plot in which all points of equal velocity are linked by contours. This reveals the velocity structure and any anomalies in the plane of the section, and provides a useful aid to geologic interpretation.

**VELOX-CI** A **\*minisatellite** mission of Nanyang Technological University of Singapore in collaboration with the Defence Science Organization National Laboratories to measure atmospheric temperature, humidity, and pressure as well as evaluating a navigation system. It was launched on 16 December 2015, from Satish Dhawan Space Centre, India, into a near-equatorial orbit at an altitude of 550 km.

**VELOX-1** A 3-unit **\*CubeSat** mission by Nanyang Technological University of Singapore to acquire high-resolution Earth imagery. It was

launched on 30 June 2014, from Satish Dhawan Space Centre, India, into a Sun-synchronous orbit at an altitude of about 660 km.

**velum** An accessory cloud feature consisting of a widespread veil extending horizontally above other clouds and sometimes connecting the tops of *\*cumulus* clouds. *See also* CLOUD CLASSIFICATION.

**vendavale** A strong, local, south-westerly wind affecting the Straits of Gibraltar, associated with a *\*depression*, and bringing *\*squalls* and heavy rainfall.

**Vendian** The ultimate *\*stage* (650–542 Ma ago) of the *\*Neoproterozoic \*era*, preceded by the *\*Sturtian* and followed by the Early *\*Cambrian \*epoch*. It includes the *\*Ediacaran \*period*.

**Vendobionta** A group of fossils of *\*Ediacaran* age that are believed to comprise a *\*monophyletic* group. Adolf Seilacher proposed in 1984 that they be ranked as a kingdom; in 1994 Leo W. Buss and Seilacher proposed they be ranked as a phylum ancestral to the *\*Cnidaria* and lacking nematocysts (cnidae).

**Venera** A series of sixteen Soviet missions to *\*Venus* that ran from 1967 to 1983.



[http://www.mentallandscape.com/V\\_Venus.htm](http://www.mentallandscape.com/V_Venus.htm)

- An SRI programme to study Venus.

**Venezuelan Remote Sensing Satellite-1 (VRSS-1)** A mission by Venezuela to provide high-resolution images of land and vegetation to assess the country's soil and water resources, assist urban planning, monitor illegal mining and drug activities, and support disaster monitoring. The satellite was launched on 29 September 2012, from the Jiuquan Satellite Launch Centre, China, into a Sun-synchronous near-circular orbit at a mean altitude of 640 km.

**Venice system** A system for the classification of brackish water based on the percentage of chloride contained in the water. *See* HALINITY.

**Vening-Meinesz, Felix Andries** (1887–1966) Professor of *\*geophysics* in the Netherlands, Vening-Meinesz made important contributions to the study

of gravity, and was an early advocate of *\*continental drift*. In 1926 he discovered large negative *\*gravity anomalies* over the *\*Java Trench*, and, using *\*earthquake* data to support his theory, suggested that the trenches were the result of buckling of the *\*crust*, caused by convection currents in the Earth's interior. Vening-Meinesz's use of submarines to improve the precision of his gravity measurements was an imaginative and effective innovation.

**VEN $\mu$ S** See VEGETATION AND ENVIRONMENT MONITORING ON A NEW MICROSATELLITE.

**vent breccia** See BRECCIA. Compare AGGLOMERATE.

**vent conglomerate** Rounded blocks of juvenile and accessory material infilling a volcanic vent. The blocks are derived from the collapse of the vent and surrounding cone walls during and after explosive *\*eruptions*. The *\*clasts* may be rounded by attrition with other blocks in the turbulent vent conditions prevailing during eruptive activity.

**venter** The *\*ventral* side of a cephalopod (*\*Cephalopoda*) where the cephalopod is coiled in a plane spiral. It is the periphery or circumference of the shell.

**Ventersdorp** See RANDIAN.

**ventifact** A pebble that has been faceted by the abrasive action of wind-blown sand and dust. Multiple faceting may reflect pebble movement as underlying sand is disturbed, rather than changes in wind direction. Each facet faces the direction of the abrading wind. A pebble with three facets (triangular facet) is called a 'dreikanter'; one with one facet an 'einkanter'.

**ventral** Towards the lower or underside of an organism (in a vertebrate the side of the animal furthest from the spine); the opposite of *\*dorsal*.

**Venturian** The final *\*stage* (2.2–1.9 Ma ago) in the Late *\*Pliocene* of California, underlain by the *\*Repettian*, overlain by the *\*Wheelerian* (*\*Quaternary*), and roughly contemporaneous with the uppermost *\*Piacenzian* stage.

**Venus** The second planet in the *\*solar system*, orbiting 0.72 AU from the *\*Sun* and  $38.2\text{--}261.0 \times 10^6$  km from Earth. Its radius is 6052 km; mass

$4.869 \times 10^{24}$  kg; mean density 5204 kg/m<sup>3</sup>; surface gravity 8.87 (Earth = 1); visual albedo 0.65; black-body temperature 238.9 K. Venus has a dense atmosphere, with a surface pressure of 9.2 MPa, composed of carbon dioxide (96.5%) and nitrogen (3.5%), with about 150 ppm sulphur dioxide, 70 ppm argon, 20 ppm water, 17 ppm carbon monoxide, 12 ppm helium, and 7 ppm neon. The surface temperature is 737 K and wind speeds range from 0.3 m/s to 1.0 m/s. Its orbit is *\*retrograde*.

**venusian** Of, pertaining to, or characteristic of, the planet *\*Venus*. See also *CYHEREAN*.

**Venus snow** A reflective substance, visually resembling snow, that coats the surface at the top of the mountains of *\*Venus*. It is believed to consist of *\*galena* and *\*bismuthinite* that forms through the condensation of *\*pyrite* minerals that the intense heat vaporizes from the surface at lower elevations.

**veranillo** The short period of dry, hot weather, lasting a few weeks, that occurs during the summer wet season along the western coasts of Central and S. America.

**verano** In parts of tropical Central and S. America, the drought season that occurs from November to April.

**Vereiskian** A Russian *\*stage* (311.7–309.2 Ma ago) in the *\*Pennsylvanian* *\*epoch*, preceded by the *\*Melekesskian* (*\*Bashkirian* epoch) and followed by the *\*Kashirskian*.

**vergence** In an *\*asymmetrical fold*, the apparent direction of movement of the upper limb with respect to the shorter limb; i.e. the *\*facing direction* (*younging*) along the *\*axial plane* at right angles to the axis. *\*Thrust* belts are commonly marked by asymmetric folds with uniform vergence.

**vermiculite** Member of the 2:1 *\*phyllosilicates* (sheet silicates) with composition (Mg,Ca)(Mg, Fe<sup>2+</sup>)<sub>5</sub>(Al,Fe<sup>3+</sup>) [(Si,Al)<sub>8</sub>O<sub>20</sub>](OH)<sub>4</sub>. 8H<sub>2</sub>O and closely related to the *\*smectite* *\*clay minerals*; sp. gr. about 2.3; *\*hardness* about 1.5; *\*monoclinic*; yellow or brown; pearly *\*lustre*; crystals flat and *\*platy*; *\*cleavage* perfect basal {001}; occurs as an *\*alteration* product of *\*biotite*, from the *\*hydrothermal* alteration of biotite and *\*phlogopite*, and

from the alteration of **\*ultrabasic** rocks; expands greatly on heating. It is used extensively for insulation and as a lubricant. *See also* CLAY MINERALS.

**Verrucano** A **\*Permian** formation of **\*sandstones** and **\*conglomerates** with a purplish-red colour.

**vertebra** In the axial skeleton of vertebrates, one of a series of bony segments which replace the notochord, forming the vertebral column (or spinal column or backbone), which encases and so protects the spinal cord. Vertebrae differentiate into five types from anterior to posterior: cervical; thoracic; lumbar; sacral; and caudal. Cervical vertebrae facilitate the mobility of the head. The first two vertebrae of the vertebral column, the atlas and axis, are highly specialized cervical vertebrae, the former articulating with the **\*occipital** region of the **\*cranium**. The thoracic vertebrae articulate with the ribs that fuse with the sternum. Lumbar vertebrae are generally larger, with abbreviated ribs fused to the centrum and supporting the posterior coelomic musculature. Sacral vertebrae fuse with the **\*pelvis**, allowing the transfer of force to the appendicular skeleton. Caudal vertebrae are smaller and less specialized, forming the tail of the organism. Six anatomical features are usually recognizable in vertebrae: the centrum is a solid cylinder which surrounds and often replaces the notochord, forming the central body of the vertebra; the neural arch forms a dorsal ring surrounding the spinal cord; a hemal arch grows ventrally on post-anal vertebrae, enclosing blood vessels; neural and hemal spines are anterior–posterior-oriented blades of bone that project dorsally and ventrally respectively; apophyses are bilaterally paired projections to which musculature is usually attached, including prezygapophyses and postzygapophyses, which occur on the anterior and posterior ends of a vertebra respectively and articulate with zygapophyses of adjacent vertebrae; transverse processes are bilaterally paired lateral projections at each side of the neural arch with which the rib articulates.

**vertebral column** *See* VERTEBRA.

**Vertebrata** *See* CRANIATA.

**vertebrates** *See* CRANIATA.

**vertebratus** A variety of cloud, usually with **\*cirrus**, with cloud elements characterized by a skeletal arrangement, in a form resembling fish

vertebrae. *See also* CLOUD CLASSIFICATION.

**vertical component (V)** The intensity of the \*geomagnetic field normal to the local horizontal plane.

**vertical electrical sounding (VES)** A method for producing information about the subsurface by measuring the \*apparent resistivity at increasing \*electrode spacings. *See* ELECTRICAL SOUNDING.

**vertical fold** *See* NEUTRAL FOLD.

**vertical seismic profile (VSP)** A form of \*seismic record obtained by positioning a detector (\*hydrophone) in a \*borehole at a succession of depths and firing a \*shot from a fixed point at the surface. *See also* WALKAWAY VERTICAL SEISMIC PROFILE.

**vertical stacking** In seismic investigations, the technique in which the signal-to-\*noise ratio is enhanced by summing the signals for a number of \*shots into one set of \*geophones or \*hydrophones at one \*offset. It is used extensively in land seismic-\*refraction surveys, using signal-enhancement \*seismographs which can sum up to 255 shots (e.g. hammer blows, weight-drops, etc.) to produce a good-quality \*seismic record, but is of limited value in marine work, where \*common-depth-point stacks are preferred.

**vertic horizon** In the \*World Reference Base for Soil Resources classification scheme, a \*soil horizon that comprises more than 30% \*clay, that swells significantly when wet and shrinks when dry, and that exhibits \*slickensides.

**vertisols** An order of mineral soils in the \*USDA \*soil taxonomy that contain more than 30% by weight of swelling clay (e.g. \*montmorillonite), and that expand when wet and contract when dry to produce a self-inverting soil and an undulating (\*gilgai) microrelief. Vertisols are associated with seasonally wet and dry environments, and are extensive in the tropics.

**very-long-baseline interferometry (VLBI)** A technique for determining, with precisions of centimetres, the distances between different radiotelescopes, using the \*phase difference between radio signals detected by them that come from very distant radio sources (quasars). Some 135 different sources are routinely used.

**very-low-frequency method (VLF method)** An electromagnetic prospecting method in which permanent, high-power, military transmitters are used as sources of unmodulated carrier waves which induce secondary fields within subsurface conductors that may be many hundreds of kilometres from the source. Various orthogonal components of the secondary field are measured, as a function of the primary field, to obtain information about the nature and position of the subsurface conductors.

**very-near infrared** See NEAR-INFRARED.

**VES** See VERTICAL ELECTRICAL SOUNDING.

**vesicle** (*adj.* **vesicular**) A bubble-shaped cavity in *\*lava*, formed by the expansion of entrapped gases. Such cavities may later become filled with material deposited from solution. Vesicular *\*basalt* (bubbly basalt lava) is basaltic lava containing numerous openings, generally ellipsoidal or cylindrical in shape, formed by the expansion of dissolved gases in the molten rock. See also AMYGDALÉ; SCORIACEOUS.

**vesicular basalt** See VESICLE.

**Vesta** The third largest (after *\*Ceres* and *\*Pallas*) *\*solar* system asteroid (No. 4), diameter 526 km; approximate mass  $3 \times 10^{20}$  kg; rotational period 5.342 hours; orbital period 3.63 years. It was imaged in 1995 by the Hubble Space Telescope and appears to have a basaltic crust overlying an *\*olivine* mantle, indicating that differentiation has occurred. In 2007, the spacecraft *Dawn* took off on a mission to study Vesta and Ceres. The first spacecraft to do so, it entered Vesta's orbit in July 2011 and completed a 14-month survey mission. It went on to Ceres, entering orbit in March 2015.

**Vesuvian eruption (sub-Plinian eruption)** A type of volcanic activity marked by very explosive *\*eruptions* which occur after long periods of dormancy, during which gas pressures in the underlying *\*magma* have built up sufficiently to eject the *\*plug* of solid *\*lava* from the vent. Escaping gases, exsolved in the magma, produce a mobile, frothy lava (*\*pumice*) and clouds of *\*ash* and gases are released into the air. See VOLCANO. Compare HAWAIIAN ERUPTION; PELÉÉAN ERUPTION; PLINIAN ERUPTION; STROMBOLIAN ERUPTION; SURTSEYAN ERUPTION; VULCANIAN ERUPTION.

**vesuvianite** See IDOCRASE.

**VGP** See VIRTUAL GEOMAGNETIC POLE.

**VHN** See VICKERS HARDNESS NUMBER.

**vibration, ground** Ground vibration due to seismic or other activity (e.g. industrial) may be transmitted to structures, the effects on them varying according to the amplitude of the ground motion, the rock or soil through which the vibration travels, and the length of time for which it persists. See VIBROSEIS.

**vibration direction** Light is an electromagnetic vibration, and can be likened to a transfer of energy by vibrating 'particles' along a path from the source to the receiver. In a ray of light of a single wavelength, the wave is generated by the vibration of 'particles' lying along the path of the ray. When, however, a ray of light passes through an *\*anisotropic* mineral plate, the light is split into two unequal components which vibrate in mutually perpendicular directions, and which are related to the *\*refractive* indices of the mineral in those directions.

**Vibroseis** The registered name (trademark) of a device which uses a truck-mounted vibrator plate coupled to the ground to generate a wave train up to seven seconds in duration and comprising a sweep of frequencies. The recorded data from an upswing or downswing (increasing or decreasing frequency respectively) are added together and compared with the source input signals to produce a conventional-looking *\*seismic section*. The device is used increasingly in land surveys instead of explosive sources.

**vicariance** The geographical separation of a species so that two closely related species or a species pair result, one species being the geographical counterpart of the other.

**Vickers hardness number (VHN)** A quantitative measure of the *\*hardness* of *\*ore minerals*, that is used in their identification. A *\*diamond* is indented into a mineral under a predetermined load (usually 100 g), for a given time (usually 15 s), and the cross-sectional area of the indentation is converted into a number, the Vickers hardness number.

**Victoriapithecus** The earliest known Old World monkey; together with a related, more poorly known genus, *Prohylobates*, it is now placed in a family, Victoriapithecidae, separate from advanced Old World monkeys. So far, the family is known only from E. and N. Africa. The *\*bilophodonty*

characteristic of Old World monkeys is only partially developed in Victoriapithecidae and in other respects, too, they are very primitive.

**vidicon** In *\*remote* sensing, an imaging device which uses a transparent material whose electrical conductivity changes with varying incidence of *\*electromagnetic radiation*. A sweeping electron beam is used to measure the varying conductivity which is then translated into an image.

**Vietnam Natural Resources, Environment and Disaster-monitoring Satellite-1A (VNREDSat-1A)** An Earth-observation mission by the Space Technology Institute and Vietnam Academy of Science and Technology that captures images with a resolution 2.5 m for studying the effects of climate change, predicting and taking measures to prevent natural disasters, and managing natural resources. The satellite was launched on 7 May 2013, from Kourou, French Guiana, into a Sun-synchronous orbit at an altitude of 704 km.

**Viking** Two *\*NASA* spacecraft, launched in 1975, that placed vehicles in orbit and landed instruments on the surface of *\*Mars*, Viking 1 landing on 20 July and Viking 2 on 3 September. Both transmitted substantial amounts of data to Earth, as well as the first colour photographs taken from the martian surface.



<http://nssdc.gsfc.nasa.gov/planetary/viking.html>

- A NASA programme of two missions that landed on Mars in 1976.

**Villafranchian** A mammalian *\*age* whose base is dated at approximately 3 Ma ago. It lasted an estimated 2 Ma and therefore transgresses the Late *\*Pliocene*/Early *\*Pleistocene* boundary.

**virga 1. (fall-stripes)** From the Latin *virga* meaning 'rod', a supplementary cloud feature referring to the trails of *\*precipitation* falling from the under surface of cloud but evaporating before reaching the ground. This phenomenon is often seen with *\*cumulus*, *\*cumulonimbus*, *\*altocumulus*, *\*stratocumulus*, *\*cirrocumulus*, *\*nimbostratus*, and *\*altostratus*. *See also* CLOUD CLASSIFICATION. **2. (pl. virgae)** On the surface of *\*Titan*, a stripe of colour.

**virgella** In a graptolite (*\*Graptolithina*), a spine projecting beyond the aperture of the *\*sacula*.

**Virgellina** See GRAPTOLOIDEA.

**Virgilian** The final *\*stage* (305–299 Ma ago) in the *\*Pennsylvanian \*epoch* of N. America, underlain by the *\*Missourian*, overlain by the *\*Wolfcampian (\*Permian)*, and roughly contemporaneous with the upper Charnovnicheskian and Dorogomilovskian stages of the *\*Kasimovian* series plus the *\*Gzhelian* series.

**virtual geomagnetic pole** The point on the Earth's surface at which a magnetic pole would be located if the observed direction of remanence at a particular location was due to a magnetic dipole at the centre of the Earth.

**viscoelastic behaviour (elastoviscous behaviour)** The *\*strain* behaviour of materials which are essentially viscous but which deform elastically under *\*stresses* of short duration. Following the release of a longer-term stress the degree of elastic recovery of viscoelastic materials is somewhat less marked than would be observed in purely elastic materials.

**viscosity** The internal resistance of a substance to flow when a *\*shear* stress is applied. Quantitatively defined, it is the ratio of the shear stress to the *\*strain rate*, in units of pascal seconds (1 Pas = 10 poise). Resistance to flow is caused essentially by molecular or ionic cohesion. In *\*magmas*, molecular cohesion can be very high, especially if the *\*silica* content is high as in *\*rhyolite* magmas, and a yield strength must be overcome before the magma can flow. The presence of solid crystals increases the effective internal cohesion, and dissolved gas reduces it. In general, basaltic magmas have lower viscosities than rhyolite magmas.

**viscous remanent magnetism (VRM)** The magnetization acquired, at room temperature, by a sample lying for a period of time in a constant magnetic field. Usually it is exponentially dependent on the duration of the field. Most rock samples acquire this form of magnetization as they lie in the *\*geomagnetic* field after their formation.

**Visean** A *\*stage* (345.3–326.4 Ma ago) in the *\*Mississippian \*epoch*, the Visean is preceded by the *\*Tournaisian*, followed by the *\*Serpukhovian*, and comprises the Chadian, Arundian, Holkerian, Asbian, and Brigantian *\*substages* (these are also *\*stage* names in western European stratigraphy).

The Visean is roughly contemporaneous with the upper Carboniferous Limestone series (Britain), and the upper *\*Osagean*, *\*Meramecian*, and lower *\*Chesterian* (N. America). *See also* [DINANTIAN](#).

**vishnevite** *See* [CANCRINITE](#).

**visible radiation** *\*Electromagnetic* radiation which has a wavelength between 380 and 780 nm, visible to the human eye.

**visual binary** *See* [STAR PAIR](#).

**vitrage** The resemblance to stained glass that certain minerals exhibit when lit from behind.

**vitrain** *See* [COAL LITHOTYPE](#).

**vitreous** Of a *\*mineral \*lustre*, glassy.

**vitric horizon** In the *\*World Reference Base for Soil Resources* classification, a *\*soil horizon* that contains more than 10% of volcanic glass or other volcanic material.

**vitrinite** *See* [COAL MACERAL](#).

**vitroclastic** Applied to *\*pyroclastic* material that contains fragments of glass.

**vitrophyric** Applied to an *\*igneous \*texture* in which *\*phenocrysts* are embedded in a glassy *\*groundmass*.

**Viverridae** *See* [CARNIVORA](#).

**vivianite** Mineral,  $\text{Fe}_3(\text{PO}_4)_2 \cdot 8\text{H}_2\text{O}$ ; sp. gr. 2.9; *\*hardness* 1.5–2.0; *\*monoclinic*; greyish-blue, occasionally colourless; *\*vitreous \*lustre*; crystals *\*prismatic*, rod-like, *\*acicular*, and stellate masses; *\*cleavage*, one, perfect, *\*pinacoidal*; occurs in phosphorus-rich, sedimentary iron ores and *\*peat* bogs, in association with *\*siderite*, in a reducing environment with organic remains present, and in close association with *\*fossil* bones and shells. It is used as a cheap blue pigment.

**VLBI** *See* [VERY-LONG-BASELINE INTERFEROMETRY](#).

**VLF method** *See* [VERY-LOW-FREQUENCY METHOD](#).

**VNREDSat-1** See VIETNAM NATURAL RESOURCES, ENVIRONMENT AND DISASTER-MONITORING SATELLITE-1A.

**vog** See VOLCANIC HAZE.

**vogesite** A type of \***lamprophyre** characterized by \***essential** \***hornblende** and \***orthoclase** feldspar.

**Vogt, Johan Hermann Lie** (1858–1932) A Norwegian geologist, Vogt was professor of metallurgy at the Technical University of Norway. He did important work on the chemistry of \***silicates**, and on differentiation in cooling \***magmas**. He also made studies of \***ore geology**, especially magmatic ores.

**void** A hole in a rock. If they are interconnected, voids form paths along which water and other fluids may flow. In increasing order of size, the major types of voids are: intercrystalline boundaries; intergranular \***pores** or spaces between the grains of a \***sediment**; microfractures or local cracks, usually extending for only a few tens of centimetres and from a few micrometres to 0.1 mm wide; \***fractures** including \***joints**, small \***faults**, and \***bedding planes**, which are often extensive and may have openings up to a few millimetres wide; fissures formed by solution, \***weathering**, or local gravitational or \***tectonic** displacement, and up to about 10 cm wide; and \***solution channels**, which range up to several metres wide and many hundreds of metres long.

**void ratio** Measurement of the porosity of rocks and soils. The void ratio ( $e$ ) =  $V_v/V_s$ , where  $V_v$  is the volume of air in voids and  $V_s$  is the volume of solid particles.

**Voigt model** See KELVIN–VOIGT MODEL.

**volatile** 1. Applied to a substance with a high \***vapour pressure**, which passes readily into a gaseous \***phase**. 2. (a) A dissolved element in a silicate \***magma** which would be gaseous at that temperature except for the confining pressure and solvent nature of the magma (e.g. Cl, F, and S) and that therefore becomes gaseous when the magma reaches the Earth's surface or a zone of reduced pressure. Common volatiles include water vapour, carbon dioxide, sulphur dioxide, hydrochloric acid, and there are many more. The melting temperatures of late, volatile, saturated \***melts**

may be about 600 °C. They form **\*pegmatites** which have large crystals and contain some metals, e.g. lithium, molybdenum, uranium, and tin. (b) In **\*coal**, a mixture of combustible gases (hydrogen, carbon monoxide, and methane) with other substances, which is given off when coal is heated without air being present. **\*Peat** contains more than 50% volatiles, **\*lignites** about 45%, **\*anthracite** 10%, and **\*graphite** less than 5%.

**volcanic-arc zone** The part of an **\*island arc region** that lies between the **\*fore-arc** and **\*back-arc basins** (or zones). The volcanic-arc zone may consist of an inactive **\*frontal arc** composed of older volcanic rocks on the fore-arc side of the active zone. The frontal and active volcanic arcs are also known as the **\*magmatic arc**.

**volcanic bomb** A lump of ejected **\*lava**, more than 32 mm across, that has acquired one of several characteristic shapes during its trajectory to the ground. For example, **\*bread-crust bombs** have a glassy crust, criss-crossed with cracks; a spindle bomb has a tail formed by the twisting or spinning of the blob as it falls; rope or ribbon bombs are twisted strands of solidified lava; and cannonball bombs are formed from solid lumps of lava that have been rounded by abrasion as they bounced to rest.

**volcanic centres** See ERUPTIVE CENTRES.

**volcanic conduit** The passage in a **\*volcano** through which **\*magma** rises towards the surface.

**volcanic cone** A conical mound of volcanic ejecta accumulated around an eruptive vent. Cones have outer slope angles of about 30° and are topped by a depression or crater over the site of the vent. The type of material which accumulates to form the cone can be used to name the type of cone. For example, alternate layers of **\*lava** with beds of **\*ash** and other **\*pyroclastic** material characterize a **\*stratovolcano** (composite volcano), and **\*spatter** ejected from a vent during a Hawaiian-type eruption would accumulate to form a **\*spatter cone** around the vent. **\*Scoria** ejected from a vent during a Strombolian-type eruption would accumulate to form a **\*scoria cone** around the vent. See VOLCANO.

**Volcanic Constructs** See MARTIAN TERRAIN UNITS.

**volcanic dome** See DOME.

**volcanic dust** Dust, *\*ash*, or other particulate matter commonly suspended in the atmosphere after volcanic *\*eruptions*. After explosive eruptions the dust may be thrown to heights of 20–30 km or more. The fall-out times of dust particles are quite short, a matter of days or weeks, depending on altitude and *\*precipitation*. Volcanogenic *\*aerosols*, usually sulphates, may linger for months, spreading as a long-lived veil in the *\*stratosphere* over much of the Earth.

**volcanic-exhalative processes** Processes associated with penecontemporaneous *\*volcanicity* that produce sulphide *\*ore* deposits, often lenticular in cross-section and commonly located above mineralized *\*stockworks* (the probable channelways to the sea floor). The Kuroko-type copper deposits are typical of continental-arc volcanic-exhalative sulphides.

**Volcanic Explosivity Index (VEI)** A relative measure of the explosive force of volcanic eruptions devised in 1982. See [APPENDIX G](#).

**volcanic haze (vog)** A grey-white haze that forms when sulphur dioxide released by *\*volcanoes* reacts with atmospheric oxygen and water to produce highly reflective sulphate particles.

**volcanicity (volcanism, vulcanicity, vulcanism)** All the processes associated with the transfer of *\*magma* and *\*volatiles* from the interior of the *\*Earth* to its surface. Current volcanicity is confined to regions of the Earth where lithospheric *\*plates* converge, diverge, or pass over possible *\*mantle \*hot spots*.

**volcaniclastic** Applied to a *\*clastic rock* that contains volcanic material. The volcaniclastic particles may be *\*alloglasts*, *\*autoclasts*, *\*epiclasts*, *\*hydroclasts*, or *\*pyroclasts*.

**volcanism** See [VOLCANICITY](#).

**volcanic neck** See [VOLCANIC PLUG](#).

**volcanic pile** See [CENTRAL VENT VOLCANO](#).

**Volcanic Plains** See [MARTIAN TERRAIN UNITS](#).

**volcanic plug (volcanic neck)** The cylindrical filling of an ancient *\*volcano* which, due to its greater resistance, may be preserved after the volcanic edifice has been eroded away. See also [PUY](#).

**Volcanic units** See MARTIAN TERRAIN UNITS.

**volcano** A naturally occurring vent or fissure at the \*Earth's surface through which erupt molten, solid, and gaseous materials. The \*viscosity, gas content, and rate of extrusion of the \*magma probably determine the shape of the mountain built by the \*eruptions. The \*magma may reach the surface either through a single channel (see CENTRAL VENT VOLCANO), or through a series of vertical fractures (see FISSURE VOLCANO). Types of eruptions are named after volcanoes associated with them. See HAWAIIAN ERUPTION; PELÉEAN ERUPTION; PLINIAN ERUPTION; STROMBOLIAN ERUPTION; SURTSEYAN ERUPTION; VESUVIAN ERUPTION; VULCANIAN ERUPTION.

**volcano-tectonic depression** An extremely large, \*caldera- or trough-like, collapse depression, that is usually surrounded by extensive \*ignimbrite sheets. One of the largest such depressions is in the Taupo region of N. Island, New Zealand, and measures 100 × 30 km. Other well-known examples occur in the western USA (e.g. the Yellowstone volcano-tectonic depression) and the Philippines (the Lake Toba volcano-tectonic depression).

**Volgian** A regional \*stage (150.8–142 Ma ago) in the northern European Late \*Jurassic, preceded by the \*Kimmeridgian and followed by the \*Ryzanian, which is distinguished from the \*Tithonian to the south by its Boreal (see BOREAL REALM) ammonite (\*Ammonoidea) fauna. Some authors equate it broadly with the \*Portlandian.

**volume diameter** See PARTICLE SIZE.

**volume diffusion** See DIFFUSION.

**volumenometer** An apparatus used to determine the density of powdered minerals or of substances which react with the \*heavy liquids normally used in the floating equilibrium method. The specimen is placed in a test chamber, graduated with 'upper' and 'lower' marks, and fitted with an airtight lid. The base of the chamber is connected by a flexible U-tube to a mercury reservoir and raising or lowering the reservoir brings mercury into the chamber as far as the 'upper' or 'lower' mark. The volume of compressed air in the chamber is calculated before and after a given weight

of a substance is introduced, so the volume of the substance can be determined. The density can then be calculated from the volume and the atomic weight of the substance.

**volume scattering** In \*radar terminology, the scattering of \*electromagnetic radiation in the interior of a material, such as a vegetation canopy or soil.

**von Karman–Prandtl equation** See KARMAN-PRANDTL EQUATION.

**vorticity** The measure of the amount of rotary or circular motion in a water or \*air mass about a vertical axis. Two types can be distinguished: ‘relative vorticity’ is rotation relative to the Earth’s surface (it is positive if cyclonic, with the rotation in the same direction as the Earth’s direction of rotation, and negative if anticyclonic); ‘absolute vorticity’ is relative vorticity plus the component of the Earth’s rotation about its axis. Absolute vorticity is at a maximum at the poles and zero at the equator.

**Voyager** Two \*NASA spacecraft that were launched in 1977 (Voyager 1 on 5 September and Voyager 2 on 20 August) on a mission to \*Jupiter, \*Saturn, \*Uranus, \*Neptune, and out of the \*solar system. Voyager 1 passed Jupiter in March 1979 and Saturn in November 1980, its camera being used for the last time on 13 February 1990, to take pictures of the entire solar system, except for Mars, Mercury, and Pluto; Voyager 2 reached Jupiter in July 1979 and Saturn in August 1981. They completed their explorations in 1989, but continued on their way. By 2018 Voyager 1 had entered interstellar space and Voyager 2 was in the heliosheath, the outermost layer of the heliosphere, where the pressure from interstellar gas slows the \*solar wind.



<http://voyager.jpl.nasa.gov/>

- A NASA programme of two missions launched in 1977 to study the planetary systems of Jupiter and Saturn and beyond.

**VRM** See VISCOUS REMANENT MAGNETISM.

**VRSS-1** See VENEZUELAN REMOTE SENSING SATELLITE-1.

**VSP** See VERTICAL SEISMIC PROFILE; WALKAWAY VERTICAL SEISMIC PROFILE.

**vug (vugh)** A cavity in a rock, which may contain a lining of crystalline minerals.

**vuggy porosity** A form of *\*secondary* porosity in which the *\*pore* spaces are formed by *\*solution* *\*vugs*. See POROSITY.

**vugh** See VUG.

**Vulcanian eruption** An explosive type of volcanic *\*eruption* that occurs when the pressure of entrapped gases in a relatively viscous *\*magma* becomes sufficient to blow off the overlying crust of solidified *\*lava*. A characteristic feature of Vulcanian eruptions is that the material ejected comes from the older rocks in the volcanic edifice and new magma is not erupted. The activity is often long lasting, with the formation of volcanic gas and *\*ash* clouds and the violent ejection of solid angular fragments of all sizes. See VOLCANO. Compare HAWAIIAN ERUPTION; PELÉÉAN ERUPTION; STROMBOLIAN ERUPTION; SURTSEYAN ERUPTION; VESUVIAN ERUPTION.

**vulcanicity** See VOLCANICITY.

**vulcanism** 1. A theory of the 18th and 19th centuries, based initially on fieldwork in the French Auvergne and associated with Nicolas *\*Desmarest* and James *\*Hall*. They proposed that volcanic rocks were produced from molten material, and that *\*volcanoes* had formerly existed where *\*basalts*, etc. are now present. 2. See VOLCANICITY.



**Waalian** An *interglacial stage* that occurred in northern Europe from about 1.3 to 0.9 Ma ago. It may be equivalent to the *Donau/Günz Interstadial* of the Alpine areas and the *Aftonian* of N. America.

**wacke** A *sandstone* which contains between 15% and 75% mud *matrix*. See DOTT CLASSIFICATION.

**wackestone** A *limestone* defined by the *Dunham* classification as consisting of *carbonate* particles in a mud-*matrix-supported* texture.

**wad** A variety of *psilomelane*, generally with the formula  $\text{BaMn}_8\text{O}_{16}(\text{OH})_4$  but copper and cobalt may substitute for manganese; *hardness* 5.5; dull, greyish-black; amorphous or *earthy aggregates*. It is precipitated from water in poorly drained, boggy ground.

**Wadati-Benioff zone** See BENIOFF ZONE.

**wadi (ouadi)** The Arabic term for an *ephemeral* river channel in a *desert* area. Flow may occur very occasionally.

**Waiauan (Waiaun)** A *stage* (11.5–10 Ma ago) in the *Miocene* of New Zealand, underlain by the *Lillburnian*, overlain by the *Tongaporutuan*, and a substage of the *Tortonian* stage.

**Waiaun** See WAIAUAN.

**Waipawan** A *stage* (56.5–52 Ma ago) in the *Palaeogene* of New Zealand, underlain by the *Teurian*, overlain by the *Mangaorapan*, and roughly contemporaneous with the upper *Thanetian* and lower *Ypresian* stages.

**Waipipian** A *stage* (3.6–2.6 Ma ago) in the *Pliocene* of New Zealand, underlain by the *Opoitian*, overlain by the *Mangapanian*, and a substage

of the [\\*Piacenzian](#) stage.

**Waitakian** A [\\*stage](#) (27–23.03 Ma ago) in the [\\*Miocene](#) of New Zealand, underlain by the [\\*Dunroonian](#), overlain by the [\\*Otaian](#) ([\\*Miocene](#)), and a substage of the [\\*Chattian](#) stage.

**walkaway vertical seismic profile** The [\\*seismic record](#) which is obtained when [\\*hydrophones](#) are positioned down a [\\*borehole](#) to detect [\\*seismic waves](#) originating from successive [\\*shots](#) fired from a surface source with increasing [\\*offset](#). It provides much information about the geophysical characteristics of [\\*reflectors](#) and about the geologic structures in the vicinity of an existing well.

**Walker's steelyard** See [COUNTERPOISED BEAM BALANCE](#).

**Wallace's line** An important zoogeographical division which separates the [\\*Oriental](#) and [\\*Australian](#) faunal realms. Alfred Russel Wallace (1823–1913), a zoogeographer and contemporary of [Charles \\*Darwin](#), first demarcated the boundary, known to this day as 'Wallace's line', between the Oriental faunal realm and the Australian, with its distinctive marsupials ([\\*Marsupialia](#)). This boundary passes east of Java and Bali, northward through the Strait of Makassar (separating Borneo and Sulawesi), then extends eastward, south of Mindanao in the Philippines.

**wall cloud (murus)** See [CLOUD CLASSIFICATION](#).

**wall, law of the** See [LAW OF THE WALL](#).

**wall rock** The rock lining a mineral vein or igneous intrusion.

**wall-rock alteration** A reaction of hydrothermal fluids with enclosing rocks, causing changes in mineralogy that are most marked adjacent to the [\\*vein](#) and become less distinct further away. See [HYDROTHERMAL ACTIVITY](#); [HYDROTHERMAL MINERAL](#).

**Walther's law (law of correlation of facies)** An important statement relating to the manner in which a vertical sedimentary sequence of [\\*facies](#) develops. Walther's law of facies implies that a vertical sequence of facies will be the product of a series of depositional environments which lay laterally adjacent to each other. This law is applicable only to situations where there is no break in the sedimentary sequence.

**Waltonian** The Red Crag (craggs are shelly sands) of Essex and Suffolk, England, which represent part of the Lower **\*Pleistocene** in East Anglia. They mark the first **\*stage** of the British Pleistocene.

**Wangerripian** The basal **\*stage** (61.7–52 Ma ago) in the **\*Palaeogene** of south-eastern Australia, preceded by the **\*Danian** and overlain by the **\*Johannian**.

**waning slope** See **SLOPE PROFILE**.

**Warendian** A **\*stage** (485–475 Ma ago) of the Early **\*Ordovician \*epoch** of Australia, underlain by the **\*Datsonian**, followed by the **\*Bendigonian**, and containing the **\*Lancefieldian**.

**Warepan** A New Zealand **\*stage** (206–203 Ma ago) of the Late **\*Triassic \*epoch**, preceded by the **\*Otamitan** and followed by the **\*Otapirian**.

**warm front** A boundary between advancing warm air that is displacing colder air, e.g. in mid-latitude **\*depressions** where, owing to the convergence of the **\*air masses** and the difference of density between them, the warm air tends to rise over the cold air. Slopes of warm fronts are typically less than 1:100 and the ascent of air is gradual. Stratiform cloud develops in the rising air. High **\*cirrus** cloud followed by lower and thickening **\*altostratus** indicate the approaching front. As the frontal contact with the ground approaches, heavy **\*nimbostratus** and much rain may occur. Passage of the front is marked by a rise of temperature as the cold air ahead of the front gives way to the air in the warm sector behind the front, clearing of precipitation, and (in the northern hemisphere) the wind **\*veering** typically from south or south-easterly to south-westerly.

**warm glacier** See **GLACIER**.

**warm rain** Rain, resulting from the coalescence of droplets, in clouds that are unfrozen (i.e. their upper parts are not at freezing level).

**warm sector** The relatively warm air that occupies the area behind the **\*warm** front in a mid-latitude **\*depression**. Within the warm sector, pressure, wind, and temperature remain fairly steady. Cloud and **\*precipitation** depend on the precise condition of the generally stable air.

**Wasatchian** A N. American **\*stage** of the **\*Eocene** (55.5–50.5 Ma ago), preceded by the **\*Clarkfordian** and followed by the **\*Bridgerian**.

**washboard moraine** See MORAINE.

**washover delta** See WASHOVER FAN.

**washover fan (washover delta)** A fan-shaped body of \*sediment that is transported landward by marine waters flowing through or across a coastal barrier such as a \*barrier bar or island. Such bodies are formed especially during storms when the barriers are likely to be overtopped.

**washplain** A nearly flat surface made up of \*alluvium mantling a thick layer of deeply weathered (see WEATHERING) bedrock and found in a savannah environment. It is washed by seasonally flooding streams which are unable to incise due to the lack of abrasive \*bed load and the large volume of \*sediment.

**water** The water molecule is composed of two hydrogen atoms, each of atomic weight 1.00797, combined with an oxygen atom of atomic weight 15.9994. It can occur as liquid, solid, and gas \*phases. It is a very powerful solvent which is responsible for the transfer of material on and below the Earth's surface.

**water-absorption test** A test to determine the moisture content of \*soil as a percentage of its dry weight (British Standard 1377: 1967). The sample is weighed, dried in an oven, then reweighed under standard conditions. It is calculated as the moisture content, which is equal to: (weight of the container with wet soil minus the weight of the container with dry soil) divided by (weight of the container with dry soil minus the weight of the container); then multiplied by 100 to express it as a percentage.

**water balance (water budget)** A method of assessing the size of future water resources in an \*aquifer, \*catchment area, or geographical region, which involves an evaluation of all the sources of supply or \*recharge in comparison with all known \*discharges or \*abstractions. See MOISTURE BALANCE.

**water budget** See WATER BALANCE.

**water budget, global** The amount of water involved in the \*hydrological cycle each year. Average annual \*precipitation over the whole globe is about 86 cm, of which 77% falls on the oceans and 23% on land. Evaporation and transpiration by plants from the land accounts for 16% of

the total precipitation it receives, and 7% of global precipitation returns to the sea as river and **\*groundwater flows**.

<b>Global reservoirs</b>	
Reservoir	Water content (m <sup>3</sup> )
Oceans	1350 × 10 <sup>15</sup>
Land	33.6 × 10 <sup>15</sup>
Atmosphere	0.013 × 10 <sup>15</sup>

<b>Annual global flux (m<sup>3</sup>yr<sup>-1</sup>)</b>	
Precipitation over oceans	324 × 10 <sup>12</sup>
Precipitation over land	99 × 10 <sup>12</sup>
Runoff/groundwater	17 × 10 <sup>12</sup>
Evaporation/transpiration from land	62 × 10 <sup>12</sup>
Evaporation from oceans	361 × 10 <sup>12</sup>

**water-drive reservoir** An oil reservoir in which the movement of oil is driven by an active aquifer. Water moving into the reservoir from below displaces the oil, which can be recovered until the energy of the aquifer decreases to the point where it ceases to be effective, or the fluid recovered contains too much water to be of value.

**water gun** A marine seismic source which uses compressed air to drive a piston in order to evacuate a chamber flooded with water. The resulting water jet creates a vacuum in the surrounding water which implodes, causing an acoustic pulse. Since no bubbles are formed there is no **\*bubble pulse**, and consequently the short **\*shot** pulse provides higher resolution than can be attained with an **\*airgun**.

**water inventory 1.** An inventory to show how water is consumed, used or otherwise involved in a particular process or place; e.g. within a steel plant or a house. **2.** Globally, approximately 97% of the Earth's water occurs in the oceans. Of the fresh water, 75% is locked up in **\*ice** sheets and **\*glaciers**, almost 25% is **\*groundwater**, and lakes, reservoirs, swamps, river channels, biospheric water, atmospheric water, and soil moisture together

account for the remainder. The amount of water at each stage of the **\*hydrologic cycle** is called the water storage. See **WATER BUDGET, GLOBAL**.

**water potential 1. (osmotic potential, chemical potential)** The difference between the energy of water in the system being considered and of pure, free water at the same temperature. The water potential of pure water is zero, so that of a solution will be negative. If there is a gradient of water potential between two plant cells, water will diffuse down the gradient until equilibrium is reached. **2.** See **CAPILLARY MOISTURE**.

**watershed 1.** See **DIVIDE**. **2.** See **CATCHMENT**.

**waterspout** A **\*tornado**-like vortex that occurs over water and is visible because of the condensation of atmospheric water vapour in the low pressure around the core (not because water is drawn upward from the surface). A waterspout may be a tornado that has moved from over the land, or may form over water, not necessarily beneath a **\*mesocyclone** cloud.

**water storage** See **WATER INVENTORY**.

**water-table** The upper surface of **\*groundwater**, or the level below which an unconfined **\*aquifer** is permanently saturated with water. See also **VADOSE ZONE**.

**water vapour** The gaseous form of water, present in the atmosphere in varying amounts. It is an intermediate stage in the **\*hydrologic cycle**. Water vapour in the atmosphere represents 0.01% of the total **\*water inventory** of the Earth. Water molecules have a short **\*residence** time in the atmosphere, on average about nine days, though this varies over a very wide range, from a few minutes upwards. Water enters the atmosphere by evaporation, transpiration, and **\*sublimation** at the Earth's surface. The concentration of water vapour decreases with increasing altitude up to over 20 km, where some increase may occur. Measures of water-vapour concentration include **\*vapour pressure**, the humidity **\*mixing ratio**, and the **\*absolute**, specific, and **\*relative humidity**. The physical significance of water vapour in the atmosphere is in its condensation to produce cloud and **\*precipitation** (with release of latent heat of condensation, which often accelerates the uplift of rising air, and the transfer of heat from low to high latitudes), its absorption and scattering of radiation, etc. It is the principal cause of the **\*greenhouse effect**.

**water vascular system** See ECHINODERMATA.

**water velocity** The velocity ( $v$  m/s) of sound waves in water is dependent on temperature ( $T$  °C) and \*salinity ( $S$  parts per thousand), and can be calculated for any depth ( $Z$ ) using the formula  $v = 1449 + 4.6T - 0.055T^2 + 0.0003T^3 + (1.39 - 0.012T)(S - 35) + 0.017Z$ .

**water-witching** See DIVINING.

**Waucoban** See EARLY CAMBRIAN EPOCH.

**wave** A periodic disturbance in a solid, liquid, or gas as energy is transmitted through the medium. (Electromagnetic waves e.g. light, can be transmitted through a vacuum.) In water, waves may occur either at the surface or as \*internal waves. The size of the water wave varies from minute \*capillary waves to massive \*tsunami, while \*wave period ranges from a few seconds to several hours.

**wave base** The depth beneath a water mass below which \*wave action ceases to disturb the \*sediments. Wave-base depth is approximately equal to half the wavelength of the surface waves.

**wave-built terrace** A theoretical zone of sedimentation on the outer edge of a \*shore platform or, on the larger scale, of a \*continental shelf. The depth of its surface below sea level was thought to be determined by the wave base (the depth at which wave activity just fails to disturb \*sediment). Modern research has failed to show the existence of such terraces.

**wave clouds** Clouds occurring in an air stream characterized by \*wave motion, usually after passing over a mountain barrier. Lenticular (lens-shaped) clouds and cloudlets appearing stationary in the crests of the waves are a characteristic sight in climates experienced on the lee side of mountain barriers, e.g. in eastern Scotland and Sweden.

**wave-cut bench** See SHORE PLATFORM.

**wave-cut platform** See SHORE PLATFORM.

**wave depression** A low-pressure area developed at the apex of a developing \*wave distortion along a \*front. A series of such systems is typical of mid-latitude \*depression sequences.

**wave diffraction** An effect seen as **\*waves** pass through an opening in a breakwater into protected waters. The waves fan out from the opening into the region beyond, but as they do so their height is diminished.

**wave equation** An equation of **\*wave** mechanics which represents wave displacement ( $\psi$ ) and wave velocity ( $v$ ) as a function of space and time ( $t$ ), where space can be represented by rectangular coordinates ( $x, y, z$ ), such that:  $v^2\psi = (\delta^2\psi/\delta x^2) + \delta^2\psi/\delta y^2 + \delta^2\psi/\delta z^2 = (1/v^2)\delta^2\psi/\delta t^2$ ; or the equation can be represented in spherical coordinates, such that:  $(1/v^2)\delta^2\psi/\delta t^2 = (1/r^2) [\delta(r^2\delta\psi/r)\delta r + (1/\sin \theta)(\delta(\sin \theta \times \delta\psi/\delta\Phi)\delta\theta) + (1/\sin^2 \theta)(\delta^2\psi/\delta\theta^2)]$ , where  $r$  is the radius,  $\theta$  the **\*colatitudes**, and  $\Phi$  the longitude.

**wave-front** The locus of adjacent points with the same **\*phase** in the path of an advancing **\*wave**, e.g. a ripple spreading on the surface of a pond after a stone has been thrown into it.

**wavelength 1.** ( $\lambda$ ) The distance on a **\*wave** between successive points that are in **\*phase**, e.g. for a water wave, the wavelength is the distance from one crest to the next. Wavelength is related to the velocity ( $v$ ) and frequency ( $f$ ) by:  $\lambda = v/f$ . The inverse of wavelength is **\*wavenumber**. **2.** Of a **\*fold** system, the distance between one **\*hinge** or **\*trough** and the next. It is rarely possible, or necessary, to measure the wavelengths of folds precisely.

**wavellite** A **\*secondary mineral**, with the formula  $\text{Al}_3(\text{PO}_4)_2(\text{OH})_3 \cdot 5\text{H}_2\text{O}$ ; sp. gr. 2.3–2.4; **\*hardness** 3.5–4.0; **\*orthorhombic**; normally white, but often greenish, yellow, grey, or brown; vitreous **\*lustre**; normally forms fibrous, radiating aggregates, or hemispherical or globular aggregates; found on **\*joint** surfaces and cavities in rock, especially **\*slates** and **\*shales**, and in association with limonitic **\*orebodies**. It is named after the 18th-century British mineralogist W. Wavell, who discovered it.

**wavenumber 1.** Spatial frequency ( $k$ ); the number of complete **\*wave** cycles per unit distance, and the inverse of **\*wavelength** ( $\lambda$ ), such that  $k = 1/\lambda$ . **2.** Propagation constant; in electromagnetic theory, wavenumber is defined as  $2\pi/\lambda$ , and  $k^2 = \mu\omega(\epsilon\omega + i\sigma)$  for the time factor  $e^{i\omega t}$ , where  $\mu$  is the magnetic **\*permeability**,  $\omega$  the angular frequency in radians per second,  $\epsilon$  the **\*dielectric permittivity**, and  $\sigma$  the **\*electrical conductivity**.

**wave period** The time required for two successive **\*wave** crests to pass a fixed point, or the time for a single wave crest to travel a distance equal to

the length of the wave.

**wave refraction** The process by which the direction of *\*waves* moving in shallow water at an angle to the submarine contours is altered. The part of the wave train travelling in shallower water moves more slowly than that still advancing in deeper water. The lines of the wave crests therefore become more parallel with the submarine contours closer to the coast.

**wave-ripple cross-lamination** The form of cross-lamination (see **CROSS-STRATIFICATION**) produced by the migration of wave-generated *\*ripples*, or combined flow ripples (i.e. ripples formed by a combination of wave action and unidirectional flow). Wave-ripple cross-lamination is characterized by a variety of distinctive features, including: unidirectional cross-laminae, sometimes with drapes (sand laminae) oriented in the opposite direction; lensoid and complexly interwoven cross-sets; irregular, undulatory bases to cross-sets; and laminae which are discordant with the external ripple form.

**wave ripple mark** See **OSCILLATION RIPPLE**.

**wave spectrum** A concept used to describe mathematically the distribution of *\*wave* energy (proportional to the square of the significant wave height) with *\*wave period*. Using this concept, a highly confused pattern of interfering waves can be divided into its constituent wave-forms. The results can be used as an aid to wave forecasting.

**wave velocity** The velocity at which *\*waves* of energy are transmitted through a medium. It depends on the characteristic properties of the medium—in the case of *\*seismic* waves on its elastic properties and density. The wave velocity ( $v$ ) is always related to *\*frequency* ( $f$ ) and *\*wavelength* ( $\lambda$ ) by the expression  $v = f\lambda$ .

**wavy bedding** A form of *\*heterolithic \*sediment* characterized by interbedded rippled *\*sands* and mud layers. Wavy beds are commonly found on storm-dominated shelves (see **SHELF**), but also in lakes, intertidal areas, and other environments where energy levels fluctuate appreciably.

**waxing slope** See **SLOPE PROFILE**.

**waxy** Of a mineral *\*lustre*, smooth in appearance, like wax.

**weakening** In *\*synoptic meteorology*, a decrease in pressure gradient around a pressure system over time. Changes of this kind bring about a weakening of the winds. *See also* INTENSIFICATION.

**Wealden** The name given to a sequence of freshwater deposits in the British pre-*\*Aptian* *\*Cretaceous* system. In Sussex and Kent the Wealden consists of two major sedimentary units: the Hastings Beds Group and the Weald Clay Group. The former is the lower and more sandy unit, whereas the latter is mud-dominated. The name should not be confused with *\*Wealdien*.

**Wealdien** *\*Sand* deposits of the *\*Tithonian* *\*stage* located in France. The name should not be confused with *\*Wealden*.

**wearing course** *See* PAVEMENT.

**weathering** The breakdown of *\*rocks* and *\*minerals* at and below the Earth's surface by the action of physical and chemical processes. Essentially it is the response of Earth materials to the low pressures, low temperatures, and presence of air and water that characterize the near-surface environment, but which were not typical of the environment of formation. There are several varieties of rock breakdown (*see also* MECHANICAL WEATHERING). Simple disintegration may occur, resulting in the production of coarse, angular blocks, of peels or skins (the process of 'desquamation'), of *\*sands*, and of *\*silts*. Minerals may be removed in *\*solution*, and *\*chemical weathering* may form new, often easily eroded substances. *See also* CARBONATION; EROSION; FROST WEDGING; HYDRATION; HYDROLYSIS; THERMOCLASTIC.

**weathering correction** In *\*seismic refraction* and *\*seismic reflection* surveys, a time correction that is made to allow for the *\*low-velocity zone* or *\*weathering layer*. It allows *\*travel times* to be reduced to a common datum by means of the *\*static correction*.

**weathering front** The junction between chemically weathered (*see* CHEMICAL WEATHERING) *\*rock* or *\*regolith* and sound rock. Where the front lies between regolith and unweathered bedrock it may be exposed by subsequent *\*erosion* to form an *\*etchplain*.

**weathering index** A measure of the intensity of **chemical weathering**. It consists of a comparison between a mineral or a chemical compound that is relatively stable with one that is readily removed by weathering. For example, the ratio of **quartz** to **feldspar** is a widely used index. Resistant heavy minerals are also used: the ratio between **zircon** and **tourmaline** (resistant) and the **amphiboles** and **pyroxenes** (less resistant) has been employed.

**weathering layer** A layer at or near the surface, usually marked by a shape transition, in which **seismic velocities** are substantially lower than in the geophysical substrate. The geophysical weathering layer may not correlate exactly with a layer of geologically weathered materials; often it is taken to be the zone above the **water-table**.

**weathering microindices** Indices used in the grade classification of rock material by microscopic examination in the laboratory. The indices are:  $I_{mp}$  (micropetrographic index), the percentage of sound constituents divided by the percentage of unsound constituents; and  $I_{fr}$  (microfracture index), the number of microcracks in a 10 mm traverse of **thin section**.

**weathering pit** See **PANHOLE**.

**weathering profile** A vertical section, from the ground surface to unaltered bedrock, which passes through **weathering zones**. It is usually best developed in the humid tropics, where depths of 100 m have been recorded but where 30 m is more common. The nature of the profile is a complex response to climatic and geologic controls, and to long-term changes in external conditions.

**weathering series** A sequence of common **silicate** minerals, laid out in the order of their susceptibility to **chemical weathering**. A well-known series was suggested by S. S. Goldich in 1938 and runs from **quartz** (most resistant), through **muscovite**, **alkali feldspar**, **biotite**, the **plagioclase feldspars**, and **olivine** (least resistant). This sequence is the reverse of **Bowen's reaction series**, which ranks minerals in the order of their crystallization from a **melt**. An absolute scale of weathering susceptibility is unlikely to be achieved, partly because of variations in environmental conditions.

**weathering zone** A distinctive layer of weathered material that extends roughly parallel to the ground surface. It differs physically, chemically, and mineralogically from the layers above and/or below. A broad distinction may be drawn between the weathering zones in *\*drift*, which are normally distinguished by degrees of oxidation and by carbonate content, and those on bedrock, which are usually separated according to the relative proportions of *\*corestones* and weathered *\*matrix*.

**Weathernews Inc. Satellite-1 (WNISAT-1)** A commercial *\*nanosatellite* mission by Weathernews Inc. of Tokyo, Japan, to monitor the Arctic Ocean. It was launched on 21 November 2013, from Yasnny Cosmodrome, Russia, into a Sun-synchronous near-circular orbit at an altitude of 600 km.

**weather report** A record of observations of meteorological values and the state of the weather, made at a given place and time.

**weather satellite** A satellite that senses the state of the atmosphere, e.g. by photographing cloud distribution, or by using infrared photography to record cloud temperature as an indication of the height of the cloud tops, or by measuring microwave emissions from atmospheric gases, which can be translated into the atmospheric temperature at the height of the emissions. See *MICROWAVE SOUNDING UNIT*.

**wedge** See *RIDGE*.

**wedge-edge trap** A *\*stratigraphic trap* in porous beds, in the form of an inclined wedge, which may trap oil, gas, or water at its upper end. See *NATURAL GAS*; *PETROLEUM*; *POROSITY*.

**wedge tornado** See *CLOUD CLASSIFICATION*.

**Wegener, Alfred** (1880–1930) A German meteorologist and physicist, Wegener is best known for his version of the theory of *\*continental drift*. In *Die Entstehung der Kontinente und Ozeane*, published in 1915 (first English language edition: *The Origin of Continents and Oceans*, 1922) he used a wide range of arguments to support his hypothesis, including palaeontological and palaeoclimatic evidence, the *\*hypsometric curve*, seismic evidence, polar wandering, and the differences between *\*continental* and *\*oceanic crust*. He found little contemporary support, but

has since become regarded as the first to propose the hypothesis in well-argued, scientific terms.

**Weichselian** See DEVENSIAN.

**weight drop** A seismic-energy impulse produced when a heavy weight is allowed to fall and strike the ground. Weights vary from a few kilograms dropped from 2–3 m, for shallow seismic-*\*reflection* surveys, up to several tonnes dropped from 3–4 m, in large-scale, commercial, seismic-*\*reflection* surveys. In an accelerated weight drop, a mass (usually of no more than a few tens of kilograms) is accelerated towards the ground by compressed air, thus increasing the impact force and energy impulse. See also HAMMER; SHOT.

**weighted average** The value assessed from a number of samples, where each sample is given a different value of importance according to its reliability.

**weir** An engineered structure extending across an open water channel. It is possible to calculate the volume of water flowing over a weir from a knowledge of its particular characteristics. Weirs are designed for flow measurement in a variety of shapes but the most common are the suppressed, contracted, V-notch, trapezoid, and broad-crested types. Other weirs may be used to control river water levels or divert flow into channels.

**Weissliegende** See KUNGURIAN; SAXONIAN.

**Weiss zone law** See ADDITION RULE.

**welded ignimbrite** See IGNIMBRITE.

**welded tuff** See IGNIMBRITE.

**well** A completed *\*borehole*. The hole may be ‘dry’ in that it does not produce oil or gas (although containing water), or a producing well.

**well injection method** A method of solution mining in which *\*boreholes* are used to transport a solvent (usually water) to the working face. The mineral is extracted and the resultant solution lifted to the surface either through an adjacent borehole or an annulus formed by equipping the borehole with concentric casings. The pumping mechanism may be located

at the surface or in the borehole. The method is used extensively in *\*halite* mining.

**well logging (wireline logging)** The lowering of various sensors within an instrumental package (a sonde) down a *\*borehole*. The output of these instruments is then measured to produce a series of well logs as the sonde is wound back to the surface. A wide variety of instruments can be combined: *\*caliper*, *\*electrical*, *\*radioactive*, *\*temperature*, etc. The term 'well logging' is also used to refer to the recording of the nature of drill cuttings encountered during the drilling of the well. *See also* BOREHOLE LOGGING; MUD LOG.

**well-point drainage** A method for draining permeable deposits around an excavation that requires small cones of depression. Tubes about 100 mm in diameter, with wire mesh screens, are sunk into the ground and connected by a header pipe to a suction pump at the top. Usually a series of well-points are connected to one header pipe. When these are used in a staged excavation a considerable depth can be drained and the *\*drawdown* restricted, because each well-point acts as a hydrogeologic boundary.

**well screen** A system of mesh screening or holes designed to allow water to enter a *\*well* or *\*borehole* without undue loss of *\*head*, but to exclude *\*sand*, *\*silt*, and other geologic material. Normally, well screens are required only in relatively unconsolidated materials.

**well shooting** Technique in which a series of *\*geophones* are deployed down a *\*borehole* and used to determine *\*seismic velocities*, as a function of depth, by measuring arrival times from surface seismic-energy sources. *See also* UPHOLE SURVEY.

**Weltian** A *\*system* in N. American usage comprising the *\*Archaean* and *\*Priscoan* of Harland et al. (1982), and overlain by the *\*Xenian*.

**Wenlock (Wenlockian)** A *\*sub-epoch* of the Early *\*Silurian \*epoch* (428.2–422.9 Ma ago), underlain by the *\*Llandovery* and overlain by the *\*Ludlow* sub-epochs, throughout Europe. *\*Reef-dwelling* organisms abound on the *\*bedding planes* of the Wenlock *\*limestone* of the Welsh borderlands.

**Wenner electrode array** An *\*electrode* configuration in which four electrodes are deployed in a line, with equal spacing between the two

**\*potential electrodes**, and between each current electrode and its nearest potential electrode. Its **\*geometric factor** ( $K_g$ ) is  $2\pi a$ , where  $a$  is defined for each case. The Wenner array has five variations, three referred to as the tripotential method with  $\alpha$ ,  $\beta$ , and  $\gamma$  configurations, one as the Lee partitioning method (which has a fifth electrode at the **\*array** centre acting as a third potential electrode), and one as the Offset Wenner electrode array, which reduces the effects of lateral inhomogeneities. See **ELECTRODE SEPARATION**.

**Wentworth scale** See **PARTICLE SIZE**.

**Werner, Abraham Gottlob** (1749–1817) Professor of mineral science at the Mining Academy, Freiberg, who developed a system of mineral classification which he called ‘oryctognosy’, based on the external characteristics of minerals. He used the term ‘geognosy’ in preference to geology. The theory of the Earth which he taught became known as **\*neptunism**. He published little but inspired mining students from all over Europe and much of his terminology was adopted even by those who rejected his theory.

**Werrikooian** An Australian **\*stage** (1.806–0.005 Ma ago) of the Early **\*Pleistocene**, preceded by the **\*Pliocene \*epoch**.

**West Australia current** The oceanic current that flows north along the western Australian coast. The flow is strong and steady in summer, but is much reduced during the winter months. Low **\*salinity** (34.5‰) and low temperature (3–7 °C) typify the waters of this current.

**westerlies** Popular term for the prevailing, eastward-moving airstreams in the mid-latitudes of the northern and southern hemispheres.

**Western Boundary Undercurrent** See **CONTOUR CURRENT**.

**western intensification** The tendency of currents along the western margins of all oceans to be particularly strong, swift, and narrow, flowing northwards in the northern hemisphere and southwards in the southern hemisphere. Currents at the eastern margins of all oceans tend to be slower and more diffuse. See **GYRE**.

**Western Pacific Satellite (WESTPAC)** An Australian commercial passive satellite used for communications and **\*geodetic measurement**. It was

launched on 10 July 1998, from Baikonur, Kazakhstan, into a Sun-synchronous circular orbit at an altitude of 835 km.

**West Greenland current** An ocean current that flows northwards parallel to the western coast of Greenland, carrying relatively warm water through the Davis Strait, where the current divides, one section continuing into Baffin Bay and the other joining the *\*Labrador current*. The West Greenland current is a continuation of the *\*North Atlantic Drift* (see *GULF STREAM*).

**West-Kohoutek-Ikemura** A *\*comet* with an orbital period of 6.46 years; *\*perihelion* date 1 June 2000; perihelion distance 1.596 AU.

**WESTPAC** See *WESTERN PACIFIC SATELLITE*.

**Westphalian** A *\*stage* (315–306.5 Ma ago) in the Silesian European epoch (Pennsylvanian), subdivided into Westphalian A, B, C, and D (the original Westphalian E is now included in the *\*Stephanian* series). It is underlain by the *\*Namurian*, overlain by the Stephanian, and is roughly contemporaneous with the Upper *\*Bashkirian* plus most of the *\*Moscovian* series.

**westward drift** The apparent westward motion of features of the *\*geomagnetic field* (mostly the non-dipole component) with time, estimated to be about 0.2° of longitude per year during the last 100 years.

**West Wind Drift** See *ANTARCTIC CIRCUMPOLAR CURRENT*.

**wet-bulb depression** The extent to which the temperature recorded by a ventilated *\*wet-bulb thermometer* falls below the dry-bulb air temperature.

**wet-bulb thermometer** A thermometer, the bulb of which is kept moist by a thin cloth (e.g. muslin) bag connected by a wick to a bath of clean (preferably distilled) water. As long as the air is not saturated, evaporation from the muslin keeps the wet-bulb thermometer at a lower temperature than the dry-bulb thermometer beside it, with which its readings are compared. The depression of the wet-bulb temperature gives a measure of the saturation deficit, from which the *\*relative humidity* and dew-point temperature of the air can be calculated.

**‘wet chemistry’** See *GRAVIMETRIC ANALYSIS*.

**wet mass** The mass of a space satellite including its fuel.

**wet melt** See MELT.

**wetted perimeter** See HYDRAULIC RADIUS.

**W-fold** See M-FOLD.

**Whaingaroan** A *\*stage* (33.9–28 Ma ago) in the *\*Oligocene \*epoch* of New Zealand, underlain by the *\*Priabonian*, overlain by the *\*Duntroonian*, and roughly contemporaneous with the *\*Rupelian* and lower *\*Chattian* stages.

**Wheelerian** The first (1.9–0.01143 Ma ago) of two *\*stages* in the *\*Neogene* of California, underlain by the *\*Venturian (\*Pliocene)*, overlain by the *\*Holocene*, and roughly contemporaneous with the Early Pleistocene *\*series* of southern Europe.

**Whewell, William** (1794–1866) A mineralogist and moral philosopher of Cambridge, who wrote works on mathematics, *\*natural theology*, and the history and philosophy of science. He also studied the theory and causes of *\*tides*, collecting his information with the assistance of members of the British Association for the Advancement of Science. Whewell is responsible for the introduction of such terms as scientist, *\*uniformitarianism*, and *\*catastrophism*.

**whinstone** 1. In the north of England, a quarrymen's name for any fine-grained, dark grey or black to green, *\*igneous* rock, e.g. *\*basalt*, *\*chert*, and *\*dolerite*. 2. In Scotland, the name for basalt.

**whirling psychrometer (sling psychrometer)** A *\*psychrometer* with a handle, which allows rapid rotation of mounted *\*wet-* and dry-bulb thermometers to ensure air flow around the bulbs.

**whirlwind** A spiral wind storm around a low-pressure centre. In arid areas dust may be carried upward several hundred metres.

**white-out** Meteorological condition in which low cloud appears to merge with a snow-covered surface to produce a uniform appearance in which outlines cannot be discerned.

**Whiterockian (White Rockian)** A *\*stage* (471.8–462 Ma ago) of the *\*Ordovician* in the Lower *\*Champlainian \*series* of N. America, preceded

by the **\*Ibexian** and followed by the **\*Mohawkian**.

**‘white smoker’** See **HYDROTHERMAL VENT**.

**whitings** Extensive patches of white, turbid water consisting of a dense suspension of **\*aragonite mud**. These mud suspensions are seen from time to time in **\*carbonate**-dominated shallow seas, and are mainly due to the disturbance of the muddy sea floor by schools of fish, or by turbulence. It was suspected that some whitings were the result of direct precipitation of aragonite from hypersaline, carbonate-saturated waters, but recently this has been shown not to be the case.

**Whitwellian** A regional **\*stage** (426.2–425.4 Ma ago) of the Early **\*Silurian**, underlain by the **\*Sheinwoodian** and overlain by the **\*Gleedonian**.

**whole Earth composition** See **BULK COMPOSITION OF EARTH**.

**whole-mantle convection** The hypothesis that the movement of lithospheric **\*plates** is due to **\*convective** cells that extend throughout the **\*mantle**. **\*Convection** in the outer **\*core** produces a layer of hot material at the core–mantle boundary, which rises through the mantle and is replaced by subsiding cooler, denser material.

**whole-rock dating** **\*Igneous** and **\*metamorphic rocks** that contain no minerals but have particularly high Rb:Sr ratios (see **RUBIDIUM–STRONTIUM DATING**), or rocks that are too fine-grained for mineral separation, can be analysed as whole-rock samples. Whole-rock samples from different parts of the same body generally differ in rubidium content and the  $^{87}\text{Sr}:^{86}\text{Sr}$  ratio of each can be plotted as a function of its  $^{87}\text{Rb}:^{86}\text{Sr}$  ratio in an **\*isochron** diagram. At the time of the initial crystallization different parts of the sample, regardless of rubidium concentration, would have had the same  $^{87}\text{Sr}:^{86}\text{Sr}$  ratio and hence plot as a horizontal line. With the passage of time  $^{87}\text{Rb}$  would be lost and corresponding amounts of radiogenic  $^{87}\text{Sr}$  gained. As the  $^{87}\text{Sr}:^{86}\text{Sr}$  ratio changes in each part of the rock, the slope of the isochron increases progressively, providing a measure of the age of the crystallization. The intercept of the isochron at the ordinate indicates the isotopic composition of common strontium at the beginning of the process.

**whorl** See **COILING**.

**Widmanstätten structure** The pattern revealed on polished and etched **\*iron meteorites**, showing a characteristic **\*texture**. It consists of parallel interlocking bands of kamacite ( $\alpha$ -iron) and taenite ( $\gamma$ -iron) resulting from the slow cooling of the meteorite. Analysis of the pattern indicates the size and structure of the parent body. It is named after Count Alois von Beckh Widmanstätten (1753–1849) of Vienna who described it in 1804.

**Wiechert, Emil** (1861–1928) A German physicist, and founder of the Geophysical Institute at Göttingen (1901), Wiechert made important improvements to the **\*seismograph**, which made it possible to distinguish between different types of **\*seismic waves**. He was thus able to identify and name **\*P-** and **\*S-waves**. He also calculated the diameter and density of the Earth's **\*core** from seismic data.

**Wiener filter** A filter which converts a known input signal into an output signal which, according to a least-squares test, is the one most similar to a desired form of signal output.

**Wien's displacement law** The law which states that the wavelength of **\*electromagnetic radiation** emitted by a material is inversely proportional to the absolute temperature of that material. As the absolute temperature increases, the wavelength of emitted radiation becomes shorter.

**wiggle trace** The classic seismic-**\*trace** output from a galvanometer, which illustrates the wave-form of the recorded data. A modern display of **\*seismic waves** is in the form of a **\*variable-area** display with wiggle.



**Wiggle trace**

**wigwams** See TENT ROCKS.

**Wild 2** A **\*comet** with an orbital period of 6.17 years; **\*perihelion** date 6 May 1997; perihelion distance 1.583 AU.

**wildflysch** Turbiditic, **\*mass-flow** **\*sediments** with numerous ill-sorted **\*exotic** **\*clasts**. In modern terminology wildflysch is better referred to as a form of **\*diamictite**.

**willemite** Mineral,  $Zn_2SiO_4$ ; sp. gr. 4.0; **\*hardness** 5.5; **\*trigonal**; greenish-yellow, but can vary to brown or white; vitreous **\*lustre**; crystals can be **\*prismatic**, but it usually forms granular or **\*massive** aggregates; strongly **\*fluorescent**; occurs mainly in the oxidized zones of zinc deposits. It is named after King William I of the Netherlands.

**Wilson, John Tuzo** (1908–93) Professor of geophysics at the University of Toronto, Tuzo Wilson was best known for his explanation, in 1965, of the **\*transform faults** which offset ocean spreading axes. His ideas were based on studies of linear **\*magnetic** anomalies and **\*seismicity** beneath **\*ocean crust**. He was the first person to use the term **\*plates**, and also invoked the idea of a **\*hot spot** to explain the evolution of the Hawaiian chain of islands. See WILSON CYCLE.

**Wilson cycle** The hypothesis, named after Tuzo **\*Wilson** who proposed it, that an ocean basin has a lifespan with several stages: from opening, through development, to final closing and the destruction of the basin. Six stages, and the **\*plate tectonic** processes involved, have been identified in different parts of the Earth today, and have been postulated for **\*orogenic** belts back to the early **\*Proterozoic**. The earliest (embryonic) stage involves uplift and crustal **\*extension** with the formation of **\*rift** valleys (e.g. the E. African Rifts). The young stage involves further subsidence, plus **\*sea-floor spreading**; the result is a narrow, parallel-sided sea, possibly with **\*evaporites** from intermittent desiccation (e.g. the **\*Red Sea**). At this stage, if the uplift was broadly **\*domal** with a pattern of three radial rifts forming a **\*triple junction**, two of the rifts may widen leaving the third to form an **\*aulacogen** (e.g. the Ethiopian Rift). The next (mature) stage is exemplified by the **\*Atlantic Ocean**, a wide ocean basin flanked by **\*continental shelves** and with the production of new, hot, **\*oceanic** crust along an oceanic **\*ridge**. Eventually this expanding system becomes unstable, and part of the cooled **\*lithosphere**, away from the ridge, sinks into the **\*asthenosphere**, forming an oceanic **\*trench** with an associated **\*island arc**. The shrinking **\*Pacific Ocean** is thought to be at this stage. Further shrinking, with the compression, **\*metamorphism**, and uplift of **\*accretionary wedges** to form young mountain ranges, marks the terminal stage (e.g. the Mediterranean). Finally, all the oceanic crust between the continental masses has subducted and the continents converge in a **\*collision zone**, being joined along a **\*suture**. The suture (e.g. the Indus–Yarlung Zangbo suture in the Himalayas) marks the relic scar between the plates, and the **\*plate margin** finally becomes inactive.

**wilting coefficient** See PERMANENT WILTING PERCENTAGE.

**wilting point** See PERMANENT WILTING PERCENTAGE.

**WIND** A solar-terrestrial mission by **\*NASA**'s Goddard Space Flight Center as part of the Global Geospace Science and International Solar-Terrestrial Physics programmes to provide complete **\*plasma**, energetic particle, and magnetic field data for studies of the **\*magnetosphere** and **\*ionosphere**, determine the magnetospheric output to interplanetary space in the upstream region, investigate basic plasma processes in the near-Earth **\*solar wind**, and provide observations for missions to the inner and outer

heliosphere. The satellite was launched on 1 November 1994, from Cape Canaveral, initially into a double-lunar swingby orbit near the **\*ecliptic plane**, **\*apogee** 80–250  $R_E$ , **\*perigee** 5–10  $R_E$ . In 1996 the mission was extended, and WIND was inserted into a halo orbit about **\*Lagrangian point L1**, at 235–265  $R_E$ .

**Windermere Interstadial (Late-Devensian Interstadial)** A relatively warm period that occurred towards the end of the last (**\*Devensian**) glacial stage in Britain. The event took place about 13 000–11 000 radiocarbon years BP. The Windermere Interstadial includes the **\*Bølling-\*Allerød** and **\*Older Dryas \*chronozones** of Scandinavia.

**wind noise** Electrical **\*noise** generated by **\*geophones** or cables when they are buffeted by the wind; mechanical noise generated when trees and tall structures are moved by the wind and the motion is transmitted to the ground through their roots or foundations.

**window** See APERTURE.

**wind rose** Quantitative diagram (**\*rose diagram**) representing the relative frequencies of different wind directions and wind speeds at a climatic station over a period of time.

**windrow** Streak of foam or row of floating debris, aligned in the prevailing wind direction, formed on the surface of a lake or ocean. Where winds blow across a water surface, vertical circulation cells are set up in near-surface waters. These circulation cells are alternately right- and left-handed vortices, and windrows form along the lines of convergence between adjacent cells at the water surface.

**wind shear** A change of horizontal wind velocity with height, which varies according to the rate of change of temperature with altitude. Vertical wind shear can be a cause of cloud formation in the turbulent mixing taking place in a **\*boundary layer** of air moving at different speeds. The shear between the wind at different levels can be expressed as a vector, measuring the difference in speed and direction.

**windward** ‘Upwind’, i.e. in the direction from which the wind blows.

**wireline** Any system which is used down a **\*borehole** and which can be operated at the end of a cable. The term is used with reference to drilling

techniques and geophysical logging methods.

**wireline logging** See WELL LOGGING.

**Wirtanen** A *comet* with an orbital period of 5.46 years; *perihelion* date 11 June 1997; perihelion distance 0.339 AU.

**Wisconsinian** The last (80 000–10 000 years ago) of four glacial episodes recognized in N. America. As with previous glacials there were several advances of the ice, and these glacial deposits include sequences of *tills* laid down by ice in the western N. American Cordillera. More evidence from areas peripheral to the *ice* sheets suggests that temperatures were perhaps 6 °C cooler than they are now. This glacial is perhaps equivalent to the *Würm* Glaciation or Glacial of the Alpine areas.

**witherite** Member of the *carbonate* group of minerals, with the formula BaCO<sub>3</sub>; sp. gr. 4.3; *hardness* 3.5; greyish-white; *vitreous* *lustre*; crystals *tabular* and twinned (see CRYSTAL TWINNING), but also *massive* and granular; occurs as a *gangue* mineral in hydrothermal-vein deposits (see HYDROTHERMAL MINERAL) associated with *barite* and *galena*.

**within-plate basalt (WPB)** *Basalts* which are generated within a continental or oceanic *plate*, rather than at a *constructive* or *destructive* plate margin. Continental *flood* basalts (e.g. the Columbia River Plateau in the western USA) and *ocean* island basalts are types of within-plate basalts. Although each has its own particular geochemical characteristics, they have been identified as a coherent group using their Ti/100; Zr; Y × 3 ratios (against a standard). This effectively separates the WPB group from the *mid-ocean-ridge basalt (MORB)*, low-potassium *tholeiite*, and *calc-alkaline* basalt groups.

**Witwatersrand** See RANDIAN.

**WMO** See WORLD METEOROLOGICAL ORGANIZATION.

**WNISAT-1** See WEATHERNEWS INC. SATELLITE-1.

**Wolfcampian** A N. American *stage* (299–280 Ma ago) in the Early *Permian* *epoch*, underlain by the *Virgilian* (*Pennsylvanian*), overlain by the *Leonardian*, and roughly contemporaneous with the *Asselian* and lower *Sakmarian* *stages*.

**wolframite** \*Ore mineral for tungsten, with the formula  $(\text{Fe,Mn})\text{WO}_4$ ; sp. gr. 7.0–7.5; \*hardness 5.0–5.5; \*monoclinic; grey-black to brownish-black; brownish-black \*streak; metallic \*lustre; crystals normally \*tabular, \*prismatic, often \*bladed, also occurs granular and \*massive; \*cleavage perfect {010}; occurs in high-temperature hydrothermal veins, \*quartz veins, and \*pegmatites, associated with granitic rocks, with \*cassiterite, \*arsenopyrite, \*tourmaline, \*scheelite, \*galena, \*sphalerite, and quartz.

**wollastonite** Member of the \*nesosilicates (chain silicates)  $\text{CaSiO}_3$  and associated with the \*pyroxenes, although it does not possess a pyroxene atomic lattice and is termed a pyroxenoid (along with pectolite ( $\text{Ca}_2\text{NaH}[\text{SiO}_3]_3$ ) and \*rhodonite); sp. gr. 2.8–3.1; \*hardness 4.5–5.0; \*triclinic; white to grey; vitreous to pearly \*lustre; crystals \*tabular, \*prismatic, also occurs \*massive and cleavable, or fibrous; \*cleavage perfect {100}; occurs in metamorphosed, siliceous \*limestones, and \*alkaline, \*igneous rocks, and associated with \*calcite, \*epidote, and \*tremolite; soluble in hydrochloric acid with the separation of silica. It is used as a source mineral for rock wool of high strength and with long fibres, and is named after the British chemist and mineralogist William Hyde Wollaston (1766–1828).

**Wolstonian** The glaciation which followed the \*Hoxnian \*Interglacial; a pale, chalky \*till (the Gipping Till) occurs in East Anglia, England, and the stone orientation and the included \*erratics suggest a movement from the north. The general \*stratigraphy needs clarification since it is not certain that the till rests on Hoxnian deposits and there are a number of morainic deposits attributed to this glacial which occur in Norfolk. During this ice advance a large lake, Lake Harrison, was ponded up in the Midlands. The Wolstonian perhaps correlates with the \*Saalian. The Wolstonian stage is named after Wolston, Warwickshire, England, where the relevant deposits were first discovered.

**Wonokan** A \*stage of the \*Ediacaran \*period, dated at about 630–570 Ma ago (Int. Commission on Stratigraphy, 2004), preceded by the \*Varangian and followed by the \*Poundian.

**wood tin** A variety of \*cassiterite ( $\text{SnO}_2$ ) with concentric bands and compact and fibrous \*texture. Toad's-eye tin shows similar characteristics

on a smaller scale. It occurs in **\*hydrothermal** veins associated with **\*acid \*igneous** rocks.

**Woodward, John** (1665–1728) Woodward was a physician with a wide knowledge of **\*fossils**, whose large collection formed the basis for the geologic museum at Cambridge, England. He was a diluvialist, believing that fossils and **\*sediments** had settled out of the ocean of Noah's Deluge in the order of their specific gravities. His *Essay towards a Natural History of the Earth* (1695) influenced both English and continental natural philosophers. See **DILUVIALISM**.

**woolsack** See **CORESTONE**.

**Wordian** A **\*stage** (268–265.8 Ma ago) of the **\*Middle \*Permian \*epoch**, preceded by the **\*Roadian** and followed by the **\*Capitanian**.

**work hardening** The effect of a decrease in the ease of deformation of a body undergoing progressive deformation, due to a change in its material configuration, and thought to be caused by the accumulation of defects in the lattice.

**World Climate Programme** A project established in 1979 by the **\*World Meteorological Organization** for the collection and preservation of climatic data.

**World Meteorological Organization (WMO)** The United Nations agency responsible for gathering, analysing, and disseminating meteorological data. Its establishment was agreed in 1947 as a successor to the International Meteorological Organization, it came into being in 1951, and later in 1951 it became a UN agency. It has 185 members (179 member states and 6 dependent territories).

**World Reference Base for Soil Resources (WRB)** A soil classification scheme that was developed between 1961 and 1974 by the Food and Agriculture Organization of the UN (FAO) and the UN Educational, Scientific, and Cultural Organization (UNESCO), supported by the UN Environment Programme (UNEP) and the International Society of Soil Science. It has been revised several times. In 1998 the system was named the World Reference Base for Soil Resources. The WRB divides soils into 32 soil groups and 170 subunits. The soil groups are: **\*acrisols**, **\*albeluvisols**, **\*alisols**, **\*andosols**, **\*anthrosols**, **\*arenosols**, **\*calcisols**,

\*cambisols, \*chernozems, \*cryosols, \*durisols, \*ferralsols, \*fluvisols, \*gleysols, \*gypsisols, \*histosols, \*kastanozems, \*leptosols, \*lixisols, \*luvisols, \*nitisols, \*phaeozems, \*planosols, \*plinthosols, \*podzols, \*regosols, \*solonchaks, \*solonetz, \*stagnosols, \*technosols, \*umbrisols, and \*vertisols.

**WorldView** A series of US commercial imaging satellites by DigitalGlobe Inc. that provide images with a resolution of 0.5 m panchromatic and 2 m multispectral. WorldView-1 was launched on 18 September 2007, from California, into a Sun-synchronous circular orbit at an altitude of 496 km. WorldView-2, with image resolution 0.46 m panchromatic, 1.8 m multispectral, was launched on 8 October 2009, from California, into a Sun-synchronous near-circular orbit at an altitude of 767 km. WorldView-3, image resolution 31 cm panchromatic, 1.24 m multispectral, was launched on 31 August 2014, into a Sun-synchronous orbit at 617 km. WorldView-4, resolution 25 cm panchromatic, 1 m multispectral, was launched on 11 November 2016, from California, into a Sun-synchronous orbit at 617 km.

**World Weather Watch (WWW)** A worldwide system for observing, analysing, and forecasting meteorological conditions, established in 1963 under the auspices of the [\\*World Meteorological Organization](#). It supplies constantly updated weather reports and forecasts to all World Meteorological Organization members, obtaining its data from 4 satellites in [\\*polar orbit](#) and 5 in [\\*geostationary orbit](#), about 10 000 land observation stations, 7000 weather ships, and 300 moored and drifting buoys. The [\\*Tropical Cyclone Programme](#) is one of the programmes forming part of the WWW. *See also* [WORLD CLIMATE PROGRAMME](#).

**WorldWide Standard Seismograph Network (WWSSN)** An international network of [\\*seismometer](#) [\\*arrays](#) designed to detect and locate [\\*earthquakes](#). *See also* [LASA](#).

**WPB** *See* [WITHIN-PLATE BASALT](#).

**WRB** *See* [WORLD REFERENCE BASE FOR SOIL RESOURCES](#).

**wrench fault** *See* [STRIKE-SLIP FAULT](#).

**wrinkle ridge (mare ridge) 1.** A broad, curvilinear swell up to 500 m high and 10 km wide, topped by a narrow, steep-sided ridge up to 200 m high

and 2–4 km wide, extending for tens to hundreds of kilometres concentrically near the edges of lunar *\*mare* basins. They are compressional in origin, possibly due to subsidence of mare lavas. 2. See DORSUM.

**Wuchiapingian (Longtanian)** A *\*stage* (260.4–253.8 Ma ago) of the Late *\*Permian*, preceded by the *\*Capitanian* and followed by the *\*Changhsingian*.


**wulfenite** Mineral,  $\text{PbMoO}_4$ ; sp. gr. 6.5–7.0; *\*hardness* 3; *\*tetragonal*; orange-yellow, olive-green, or brown; white *\*streak*; *\*resinous lustre*; crystals usually square plates or tablets, sometimes occurs *\*massive*, granular, or *\*botryoidal*; *\*cleavage* good {101}; sub-*\*conchoidal* fracture; occurs as a *\*secondary mineral* in the oxidized zones of *\*ore minerals* of lead and molybdenum, and in association with *\*anglesite*, *\*cerussite*, and *\*pyromorphite*.

**Wulff stereographic net** See STEREOGRAM.

**Würm** See DEVENSIAN.

**WWSSN** See WORLDWIDE STANDARD SEISMOGRAPH NETWORK.

**WWW** See WORLD WEATHER WATCH.



**xanthophyllite** See MICA.

**Xenian** A \*system of the Lower \*Proterozoic of N. America, from 2500 to 1600 Ma ago.

**xeno-** Foreign, from the Greek *xenos*, stranger.

**xenoblastic** A textural term applied to \*metamorphic rocks which contain \*anhedral \*porphyroblasts.

**xenocryst** A crystal in an \*igneous rock which has not crystallized from the \*melt but has been introduced into the melt from an external source, e.g. the surrounding \*country rocks or a previously crystallized part of the same igneous body. Xenocrysts, which are usually in disequilibrium with the melt, contrast with \*phenocrysts which have crystallized in equilibrium from the melt.

**xenolith** An inclusion or enclave of a pre-existing rock in an \*igneous rock. Xenoliths are often derived from the \*country rocks that have been invaded by the igneous mass, and they frequently show some evidence of reaction, e.g. rounding of their edges and \*metamorphism.

**xenotime** \*Accessory mineral, YPO and associated with \*monazite (Ce, La, Th) PO<sub>4</sub>; sp. gr. 4.4–5.1; \*hardness 4–5; \*tetragonal; yellowish-brown, greyish-white, or pale yellow; pale brown \*streak; resinous to vitreous \*lustre; crystals \*tetragonal \*prisms, very similar to \*zircon; \*cleavage \*prismatic; occurs in granitic and \*alkaline, \*igneous rocks, as well as in \*pegmatites and \*gneisses.

**xenotopic fabric** The \*fabric of a crystalline \*carbonate rock or \*cement, or of an \*evaporite deposit, in which most of the crystals are \*anhedral.

**XPL** See **CROSSED POLARS**.

**xpols** See **CROSSED POLARS**.

**X-ray diffraction crystallography** A method of analysis in which an X-ray beam of known wavelength is directed at a crystal, and the beam is diffracted by reflections off planes of atoms in the crystal. By recording the angular positions of diffracted beams, the spacing between atomic planes can be determined according to the Bragg equation,  $n\lambda = 2d \sin\theta$ , where  $n$  is any integer,  $\lambda$  is the wavelength of the incident-beam X-ray,  $d$  is the spacing between crystal planes ( $d$  spacing), and  $\theta$  is the angle between the crystal plane and the diffracted beam angle (the Bragg angle) (see **BRAGG'S LAW**). The procedure is repeated for various directions in the crystal and a model of its internal structure established. In geology, the technique is used to identify minerals.

**X-ray fluorescence (XRF)** The secondary X-ray emission that results when **\*electrons** from the outer orbitals of an atom fill vacant inner orbital positions originally created by X- or **\*gamma-ray** excitation. These X-rays are characteristic of excited atoms.

**X-ray fluorescence spectrometry** An analytical method which can be used to determine the concentration of a wide range of chemical elements, using the intensity of their fluorescent X-rays. An X-ray beam is used to excite atoms in a sample; **\*electrons** near the **\*nucleus** emit secondary or fluorescent X-rays on reversion to their original states. Short-wavelength X-rays are sorted by diffraction in a pure analysing crystal of known  $d$ -spacing (see **COVALENT RADIUS**). Since  $n\lambda = 2d \sin\theta$  (see **BRAGG EQUATION**),  $\theta$  can be set to a value and radiation detected for a unique wavelength characteristic of the element being analysed. The intensity of the radiation measured, relative to a standard, is proportional to the concentration of the element. It is an important technique in the geochemical analysis of rocks.

**X-ray photography** A photographic technique in which X-rays are used that may reveal internal detail (e.g. of a **\*fossil**) which is not visible externally. The technique has the advantage that it is non-destructive.

**X-ray powder photograph** A photograph produced by monochromatic X-irradiation of a sample of **\*microcrystalline** powder placed at the centre of a circular camera, e.g. a Debye-Scherrer camera. Diffracted X-rays are

recorded on a strip of film wrapped around the circumference of the camera. The angular position of the diffracted X-rays on the film gives structural information about the sample. *See also* [X-RAY DIFFRACTION CRYSTALLOGRAPHY](#).

**X-ray spectrometer** An instrument for measuring the secondary X-rays emitted when the inner [\\*electrons](#) of an atom (W, Au, etc.) are activated by a primary hard X-ray beam. The content of nearly all elements with an atomic weight greater than that of sodium can be estimated.

**XRF** *See* [X-RAY FLUORESCENCE](#).



**yardang** A streamlined, wind-sculptured hill ranging in length from metres to kilometres and developed in any bedrock that is at least weakly consolidated. Yardangs are restricted to deserts that have high aridity with minimal plant cover and soil development, and that are dominated by strong, unidirectional winds for most of the year.

**Yarmouthian** The second (0.7–0.55 Ma ago) of four *\*interglacial* stages recognized in mid-continental N. America. It follows the *\*Kansan* glacial episode and is equivalent to the upper part of the *\*Günz/Mindel Interglacial* of the Alps. At various times during the Yarmouthian climates were both warmer and cooler than they are now.

**Yatalan** The final *\*stage* (3.4–2 Ma ago) in the *\*Pliocene* in south-eastern Australia, underlain by the *\*Kalimnan*, overlain by the *\*Werrikooian* (*\*Quaternary*), and roughly contemporaneous with the upper *\*Piacenzian* stage.

**yazoo stream** A river tributary that may flow for many kilometres on the *\*floodplain* of a trunk stream before finally joining it. The delayed junction is due to the development of *\*levées* by the main stream. They are named after the Yazoo River, Mississippi, USA, which follows the Mississippi River for a considerable distance before breaking through its natural levée.

**Yeadonian** A regional *\*stage* (315.5–314.5 Ma ago) in the *\*Pennsylvanian* *\*epoch*, preceded by the *\*Marsdenian* and followed by the *\*Cheremshanskian*.

**yedoma permafrost** *\*Loess* *\*permafrost* containing 50–90% ice and with organic matter amounting to about 2% carbon by mass. The permafrost is of *\*Pleistocene age* and covers more than 1 million km<sup>2</sup> in Siberia. Yedoma is the name of a town in NE Siberia.

**yellowcake** Concentrated, precipitated, and dried uranium oxide.

**yield–depression curve** A graph on which the **\*drawdown** is plotted against the yield of a pumped well or **\*borehole**. The resulting plot is invariably curved, and is used to determine the optimum pumping rate for a water supply.

**yield point** See ELASTIC LIMIT.

**yield stress** The **\*stress** at which the yield strength of a material is exceeded and elastic behaviour gives way to viscous behaviour (see VISCOSITY). If continued the stress may lead to the ‘failure stress’, beyond which **\*failure** occurs.

**Ynezian** A **\*stage** (61.5–55.8 Ma ago) in the **\*Palaeocene** of California, underlain by the N. American **\*Danian**, overlain by the **\*Bulitian**, and roughly contemporaneous with the upper Danian and lower **\*Thanetian** stages of Europe.

**Younger Dryas** See DRYAS.

**younging** See FACING DIRECTION.

**young mountain belt crustal type (active orogen)** A type of very unstable **\*crust**, averaging 55 km in thickness, but in some areas up to 70 km, and rising to elevations sometimes greater than 3 km. It occurs in regions where there has been recent and often rapid uplift, and compression resulting in extensive folding (see FOLD) and thrusting (see THRUST) of **\*sedimentary rocks**.

**Young’s modulus ( $E$ )** The ratio of longitudinal **\*stress**  $\sigma$  (force  $F$  divided by area  $A$ , *i.e.*  $\sigma = F/A$ ) to longitudinal **\*strain** (change in length  $\delta L$  divided by original length  $L$ , *i.e.*  $\delta L/L$ ) in the presence of lateral strain:  $E = (F/A)/(\delta L/L)$ . If there were no lateral strain, Young’s modulus would be equal to the **\*axial modulus**.

**Yovian** A **\*system** of the Middle **\*Proterozoic** of N. America, from 1600 to 800 Ma ago.

**Ypeenian** An Australian **\*stage** of the Middle **\*Ordovician** (470–468.1 Ma ago), preceded by the **\*Castlemainian** and followed by the **\*Darriwilian**.

**Ypresian** A *\*stage* (55.8–48.6 Ma ago) in the *\*Eocene \*epoch*, preceded by the *\*Thanetian (\*Palaeocene)* and followed by the *\*Lutetian*, named after the Ypres Clay, Belgium. It is roughly contemporaneous with the *\*Penutian* and lower *\*Ulatisian* (N. America), upper *\*Waipawan*, *\*Mangaorapan* and lower *\*Heretaungan* (New Zealand), and the lower *\*Johannian* (Australia).

**Yurmatian** A *\*substage* of the *\*Mesoproterozoic*, from about 1375 to 1100 Ma ago (Int. Commission on Stratigraphy, 2004), of western Russian origin, preceded by the *\*Burzyan* and followed by the *\*Karatau*.

# Z

**Z** The vertical component of the *\*geomagnetic field*.

**Zanclean (Zanclian)** A *\*stage* (5.33–3.6 Ma ago) in the Early *\*Pliocene*, preceded by the *\*Messinian* (*\*Miocene*) and followed by the *\*Piacenzian* (Late Pliocene). It is roughly contemporaneous with part of the *\*Delmontian* (N. America), upper *\*Kapitean* and *\*Opoitian* (New Zealand), and the upper *\*Mitchellian*, *\*Cheltenhamian*, and lower *\*Kalimnan* (Australia).

**‘zap pits’ (microcraters)** Small craters, ranging in size from less than 1µm to about 1cm, on the exposed surfaces of lunar samples, caused by the hypervelocity impact of micrometeorites and dust particles with masses less than about 10<sup>-3</sup> g; typical craters consist of a glass-lined pit, a ‘halo’ zone of fractured material surrounding and underlying the pit, and a spalled zone concentric to the pit. This *\*spallation* may leave the glass-lined pit standing on a pedestal.

**zastrugi** See *SASTRUGI*.

**Zechstein** The final *\*epoch* of the *\*Permian* period, dated at 260.4–251 Ma ago (Int. Commission on Stratigraphy, 2004), preceded by the *\*Capitanian* and followed by the *\*Induan* *\*stages*.

**Zechstein Sea** Name given to an Upper *\*Permian* shallow gulf sea or depositional sequence of rocks that developed in northern Germany and the North Sea Basin. The depositional sequence laid down in this sea consisted of *\*carbonates* and *\*evaporites*. See also *SALINE GIANT*.

**Zedian** A *\*system* of the Upper *\*Proterozoic* of N. America, from 800 Ma to about 590 Ma ago.

**Zelzate** See DENEKAMP.

**Zemorrian** A \*stage (33.5–22 Ma ago) in the \*Miocene of California, underlain by the \*Refugian, overlain by the \*Saucesian, and roughly contemporaneous with the \*Rupelian and \*Chattian (\*Oligocene) and lower \*Aquitainian (Upper Tertiary) stages.


**zeolite facies** A set of metamorphic mineral assemblages produced by the \*metamorphism of a wide range of starting rock types under the same metamorphic conditions, and typically characterized by the development of the mineral assemblage \*smectite–\*zeolites (in addition to relict \*igneous \*plagioclase and \*pyroxene) in rocks of \*basic igneous composition. Other rocks of contrasting composition, e.g. \*shales, would develop their own specific mineral assemblages, even though they are all metamorphosed under the same conditions. The variation of mineral assemblage with starting rock composition reflects a particular range of pressure, temperature, and  $P(H_2O)$  conditions. Experimental studies of mineral  $P$ – $T$  stability fields indicate that the facies represents a range of conditions involving low pressures (1–4 kb) and low temperatures (300–500 °C), usually developed in (a) thick sedimentary sequences on \*continental margins, (b) rock sequences subjected to hot water-convecting systems, and (c) areas of \*tectonic burial at the leading edge of \*thrust sheets.

**zeolites** Group of hydrated alumina silicates of sodium, potassium, calcium, and barium, which occur in \*geodes, altered \*igneous rocks, \*hydrothermal veins, and some \*sediments. The water molecules are weakly held and the hydration–dehydration reaction has some useful applications, as do the ion-exchange properties of zeolites.

**zephyr** Any gentle breeze, but especially a prevailing light and warming breeze from the west at the time of the summer \*solstice (in the northern hemisphere).

**zero-length spring** A spring system in which the effective length is zero when measured from a fixed point. This type of spring is commonly used in \*gravimeters.

**zeta-form beach (fish-hook beach)** A \*beach that increases in width in the direction of \*beach drift, and ends where it adjoins a headland.



### Zeta-form beach

**zeugen** Mushroom-shaped rock that has been eroded by the abrasive action of wind-blown *\*sand*. The undercutting effect is concentrated near ground level, where sand movement is greatest, and is enhanced in areas of near-horizontal strata when the lowest bed is relatively weak.

**Z-fold** In a *\*parasitic fold*, an asymmetric fold whose profile is Z-shaped, reflecting its location on the respective limb of a major fold. See **MINOR FOLD**.

**Zhangheng 1** See **CHINA SEISMO-ELECTROMAGNETIC SATELLITE**.

**zibar** A low-relief, rounded, coarse-grained, sand dune with no *\*slipfaces*. Regularly spaced zibars produce an undulating surface on otherwise level ground.

**zig-zag fold** See **CHEVRON FOLD**.

**zinc blende** See **SPHALERITE**.

**zincite** See **OXIDES**.

**Zingg diagram** A diagram, introduced by T. Zingg in 1935, that is used to plot the relative dimensions of the long, short, and intermediate axes of a particle, allowing its shape to be classified as bladed (see **BLADE**), *\*oblate*, *\*equant*, or *\*prolate*.

**zinnwaldite** A member of the *\*mica* group of 2:1 *\*phyllosilicates* (sheet silicates) with formula  $K_2(Fe^{2+}_{2-1}, Li_{2-3}, Al_2)[Si_{6-7}Al_{2-1}O_{20}](F, OH)_4$ ; sp. gr. 2.9–3.2; *\*hardness* 2–3; *\*monoclinic*; grey, brown, or even dark green;

\*vitreous to \*pearly \*lustre; forms thick, \*tabular crystals; \*cleavage well marked; occurs in \*granites and \*pegmatites in association with \*scheelite, \*wolframite, \*cassiterite, and \*fluorite. It is pneumatolytic in origin (see PNEUMATOLYSIS). It is named after Zinnwald, Saxony, Germany.

**zircon** Mineral,  $ZrSiO_4$ ; sp. gr. 4.6; \*hardness 7.5; \*tetragonal; most commonly light brown or reddish-brown, but sometimes grey, yellow, or green; vitreous \*lustre; crystals usually square \*prisms with bipyramidal terminations; \*cleavage \*prismatic, indistinct {110} and poor {111}; one of the most widely distributed \*accessory minerals in \*igneous rocks (e.g. \*granite, \*syenite, and \*pegmatites), when the crystals can be quite large, also occurs in \*metamorphic rocks (e.g. \*gneisses and \*schists), and concentrated in \*detrital beach and river \*sands. It is the main source of zirconium metal. Common zircon is used extensively in the foundry industry.

**ZiYuan 1.** The Chinese name for \*CBERS-4. **2. (ZiYuan-1 Number 2 Optical Mission of China)** A high-resolution imaging mission by the China National Space Administration (with no Brazilian participation). ZY-1-02C was launched on 22 December 2011, from the Taiyuan Space Centre, China, into a Sun-synchronous orbit at an altitude of 781 km. **3. ZiYuan-3A (ZY-3A)** A high-resolution, stereoscopic, Earth-mapping mission that will compile a database for the production of 1:50 000 and larger-scale maps as well as data for resource mapping, environmental surveying, disaster monitoring, city planning, and national security requirements. It was launched on 9 January 2012, from the Taiyuan Space Centre, into a Sun-synchronous near-circular orbit at an altitude of 506 km.

**Zoantharia (zoantharian corals, Hexacorallia)** (class \*Anthozoa) Subclass of solitary and colonial anthozoans which are many tentacled and have an enteron (gastrovascular cavity) divided by numerous paired \*mesenteries. The basal tissues secrete a corallum (cup) made of \*aragonite, which has an outer epitheca (skin) and radially arranged septa (see SEPTUM). The mesenteries are placed between the septa. The first group appeared in the \*Ordovician and is divided into four orders: \*Rugosa; \*Scleractinia; \*Tabulata; and \*Heterocorallia. Scleractinian zoantharians (\*Mesozoic to Recent) are responsible at the present time for building coral \*reefs, restricted to tropical areas. They may build wave-

resistant structures, sometimes very thick, and are usually associated with other organisms, e.g. calcareous *\*algae*. Organic build-ups also took place in the *\*Palaeozoic*, and although zoantharians were present they did not always represent the major part of the fauna. Stromatoporoids (*\*Stromatoporoidea*), algae, etc., are associated with zoantharians in these build-ups and it is not always certain whether Palaeozoic reefs were comparable with modern or Mesozoic examples. It is unlikely that Palaeozoic zoantharians were reef-frame builders in the sense that modern scleractinians are, and probable that organisms other than corals formed the wave-resistant structures.

**zodiacal light** A band of light apparent at night in the west, after twilight, and again before morning twilight, caused by atmospheric particles scattering sunlight.

**Zoeppritz's equations** A set of equations that describe the way the energy of seismic *\*P-* and *\*S-waves* is partitioned relative to their *\*angles of incidence* when they meet an interface, typically a different type of rock. The equations were formulated by the German geophysicist Karl Bernhard Zoeppritz (1881–1908), but not published until 1919, long after his death. *See also* **KNOTT'S EQUATIONS**.

**zoisite** Member of the *\*epidote* group of minerals, with the formula  $\text{Ca}_2\text{Al}_3\text{Si}_3\text{O}_{12}\text{OH}$ ; sp. gr. 3.3; *\*hardness* 6.5; greyish-white or greenish; crystals *\*orthorhombic*, *\*prismatic*, but normally *\*massive* or *\*columnar*; occurs in regional-metamorphic (see **REGIONAL METAMORPHISM**) rocks. Thulite is a rose-pink variety.

**zonal** **1.** Applied to winds blowing in a mainly west-to-east or east-to-west direction, particularly in describing the main, broad airstreams of the general or large-scale atmospheric circulation, e.g. the zonal westerly winds of middle latitudes. The zonal (or circulation) index is a conventional measurement indicating the strength of the west-to-east airflow over middle latitudes, e.g. in latitudes 35–55° N. Strong westerlies with the pressure systems also in a west-to-east orientation accompany high values of the zonal index. Cellular or meridional airflow (or sometimes weak and chaotic patterns of circulation) accompany low values of the zonal index. **2.** Applied to features (e.g. soils and vegetation) characteristic of a particular

region that is approximately bounded by lines of latitude (i.e. a region lying parallel to the equator).

**zonal flow** The winds that blow in a mainly west-to-east or east-to-west direction, and particularly to the main, broad airstreams of the general or large-scale atmospheric circulation (e.g. the zonal westerly winds of middle latitudes). The zonal (or circulation) index is a conventional measurement indicating the strength of the west-to-east airflow over middle latitudes.

**zonal index** See ZONAL FLOW.

**zonal scheme** Scheme concerned with the use of numerous \*fossils as stratigraphic indicators for the subdivision of a sequence of rocks.

**zonation** The subdivision of a \*stratigraphic unit or units by means of \*fossils.

**Zond** A series of Soviet lunar missions that ran from 1965 to 1970.



<https://www.lpi.usra.edu/lunar/missions/zond/>

- Zond mission.

**zone 1. (biostratigraphic zone)** A unit of rock characterized by a clearly defined \*fossil content. To avoid confusion with other types of zone the term '\*biozone' (short for 'biostratigraphic zone') is preferred by many authorities, although the term 'biozone' is also used in a different sense. The term is usually qualified to denote the type of zone. See ACME ZONE; ASSEMBLAGE ZONE; CONCURRENT RANGE ZONE; LINEAGE ZONE; OPPEL ZONE; RANGE ZONE; SUBZONE; TAXON RANGE ZONE; TEILZONE; ZONULE; INDEX FOSSIL. **2.** In \*crystallography, a set of \*crystal faces whose intersecting edges are parallel. They are also parallel to, and may be rotated about, a \*zone axis. See CRYSTAL ZONING. **3.** See METAMORPHIC ZONE.

**zone axis** An axis of a crystal, which normally passes through the centre of the crystal and may or may not be parallel to the \*crystallographic axes, but which is parallel to the edges of \*crystal faces that meet in a \*zone, so those edges may be rotated about it.

**zone fossil** See INDEX FOSSIL.

**zone of aeration** See SOIL-WATER ZONE.

**zone of saturation** See PHREATIC ZONE.

**zone refining** 1. In metallurgy, the small-scale production of high-purity metals using the principle that an impure \*melt will deposit pure crystals on solidifying. A rod of the specimen is melted over a very narrow region at one end, and then the molten region is transferred along the rod by means of a moving furnace. Impurities collect in the molten zone and are swept to one end of the metal. 2. In geology, the same mechanism has been invoked whereby \*incompatible \*trace elements are partitioned into an advancing \*partial melt.

**zone symbol ([UVW])** The symbol which defines the position of the \*zone axis of a crystal with respect to the \*crystallographic reference axes of the mineral. It is enclosed in square brackets to indicate that it refers to a line and not a plane. It may be calculated from the \*Miller indices of any two faces in the \*zone, or determined by plotting the faces on a \*stereographic projection.

**zonule** Subdivision of a subzone based essentially on the presence of a given species. Several zonules are recognized, for example, in the Upper \*Cambrian Franconia Sandstone of Minnesota. Here the \*zone was recognized by the presence of the trilobites (\*Trilobita) *Ptychaspis* and *Prosaukia*, the subzone by *Prosaukia*, and zonules by *Prosaukia striata* and *P. granulosa*.

**zoogeographical region** See FAUNAL REALM.

**zoogeomorphology** The study of the effects of animals on \*geomorphology, including their contribution to the \*erosion, transport, and deposition of rock and loose material. These effects arise from animal activities such as trampling, burrowing, mound-building, wallowing, nesting, the removal of plants, and caching food, as well as the direct consumption of \*soil and rock and dam-building by beavers (*Castor* species).

**zoophycus** With \**Chondrites*, an \*ichnoguild of many-branched, radial, \*trace fossils probably made by a worm that moved back and forth through the sediment, each branch of its burrow exploring a new area. They are believed to be \*fodinichnia.

**Zosterophyllum** A Lower \*Devonian psilopsid plant (\*Psilophytales) whose lower parts underwent H-branching, producing a tufted growth habit. Erect branches were smooth and the sporangia (see SPORE) were grouped together on terminal spikes.

**Zosterophyllophytina** See PSILOPHYTALES.

**ZTR index** The relative proportions of \*zircon, \*tourmaline, and \*rutile that are present in a sediment or \*sedimentary rock; the index provides a method for determining the extent of \*weathering the material has undergone and therefore its age. These minerals are used because their high \*specific gravity makes them resistant to weathering.

**Zungxian** A Chinese \*stage (505–501 Ma ago) of the Middle \*Cambrian, preceded by the \*Zuzhuangian and followed by the \*Paibian.

**Zuzhuangian** A Chinese \*stage (509–505 Ma ago) of the Middle \*Cambrian, preceded by the \*Maozhangian and followed by the \*Zungxian.

**zygapophyses** See VERTEBRA.

# Appendix A: Stratigraphic Units as Defined in the North American Stratigraphic Code, 1983

## Material Units

Lithostratigraphic	Lithodemic		Magnetopolarity	Biostratigraphic	Pedostratigraphic	Allostratigraphic
Supergroup	Supersuite	Complex	Polarity superzone			Allogroup
Group	Suite		<i>Polarity zone</i>	<i>Biozone</i> (interval, assemblage, abundance)	<i>Geosol</i>	<i>Alloformation</i>
Formation	<i>Lithodeme</i>		Polarity subzone			Allomember
Member (lens, tongue)			Subbiozone realism			
Bed(s) (flow(s))						

## Temporal and Chronostratigraphic Units

Chronostratigraphic	Geochronologic, geochronometric	Polarity chronostratigraphic	Polarity chronologic	Diachronic	
Eonothem	Eon	Polarity Superchronozone	Polarity Superchron		
Erathem (supersystem)	Era (superperiod)			Diachron	<i>Episode</i>
System (sub-system)	<i>Period</i> (subperiod)	<i>Polarity Chronozone</i>	<i>Polarity Chron</i>		Phase
Series	Epoch				Span
Stage (substage)	Age (subage)	Polarity Subchronozone	Polarity Subchron		Cline
Chronozone	Chron				
(Fundamental units shown in italic)					

# **Appendix B: Timescales**

## Geologic Timescale

Eon/Eonothem	Era/Erathem	Sub-era	Period/System	Epoch/Series	Began Ma
PHANEROZOIC	Cenozoic	Quaternary	Pleistogene	Holocene	0.11
				Pleistocene	1.81
		Neogene	Pliocene	5.3	
			Miocene	23.03	
			Palaeogene	Oligocene Eocene	33.9 55.8
				Palaeocene	65.5
		Mesozoic	Cretaceous	Late	99.6
				Early	145.5
				Jurassic	Late
			Middle		175.6
Early	199.6				
Mesozoic	Triassic		Late	228	
		Middle	245		
		Early	251		
PHANEROZOIC	Palaeozoic	Permian	Late	260.4	
			Middle	270.6	
			Early	299	
		Upper	Carboniferous	Pennsylvanian	318.1
				Mississippian	359.2
		Devonian	Late	385.3	
			Middle	397.5	
			Early	416	
		Lower	Silurian	Late	422.9
				Early	443.7
Late	460.9				

			Ordovician	Middle	471.8
				Early	488.3
	Palaeozoic	Lower		Late	501
			Cambrian	Middle	513
				Early	542
PROTEROZOIC	Neoproterozoic		Ediacaran		600
			Cryogenian		850
			Tonian		1000
	Mesoproterozoic		Stenian		1200
			Ectasian		1400
	Palaeoproterozoic		Calymmian		1600
			Statherian		1800
			Orosirian		2050
			Rhyacian		2300
			Siderian		2500
ARCHAEAN	Neoarchaeon				2800
	Mesoarchaeon				3200
	Palaeoarchaeon				3600
	Eoarchaeon				3800
HADEAN	Swazian				3900
	Basin Groups				4000
	Cryptic				4567.17

(Source: International Union of Geological Sciences, 2004.

Note: Hadean is an informal name. The Hadean, Archaean, and Proterozoic Eons cover the time formerly known as the Precambrian. Tertiary is now an informal name, although it continues to be widely used. Quaternary is equivalent to Pleistogene.)

## Lunar Timescale

Stratigraphic mapping of the lunar surface has established the following Systems, in order of increasing age:

System	Events
Copernican	Young ray craters (younger than about one billion years)
Eratosthenian	Older post-mare craters (about 1–3 billion years)
Imbrian	Main mare basalt flooding, preceded by formation of the Imbrium and Orientale basins (3–3.85 billion years)
Pre-Imbrian:	
Nectarian	Formation of 11 major impact basins preceded by the Nectaris basin (3.85–3.92 billion years)
Pre-Nectarian	Formation of about 30 impact basins preceded by the Procellarumbasin (before 3.92–4.2? billion years)

The crystallization of the anorthositic lunar highland crust has been dated at 4.44 billion years. This event closely followed the formation of the Moon.

Note that the term 'Procellarum System', to encompass the main period of mare lava flooding, has been abandoned.

## Martian (Areological) Timescale

Epoch	Absolute age (Gy) (Hartmann–Tanaka Model)	Absolute age (Gy) (Neukum–Wise Model)
Upper Amazonian	0.25–0.00	0.70–0.00
Middle Amazonian	0.70–0.25	2.50–0.70
Lower Amazonian	1.80–0.70	3.55–2.50
Upper Hesperian	3.10–1.80	3.70–3.55
Lower Hesperian	3.50–3.10	3.80–3.70
Upper Noachian	3.85–3.50	4.30–3.80
Middle Noachian	3.92–3.85	4.50–4.30
Lower Noachian	4.60–3.92	4.60–4.50

(These represent two different models published by the University of Arizona Press)

---

# Appendix C: Wind Strength

## The Beaufort Scale of Wind Strength

**Force 0.** Less than 1 mph (less than 1 km/h). Calm. The air feels still and smoke rises vertically.

**Force 1.** 1–3 mph (1–5.5 km/h). Light air. Wind vanes and flags do not move, but rising smoke drifts.

**Force 2.** 4–7 mph (5.6–11 km/h). Light breeze. Drifting smoke indicates the wind direction.

**Force 3.** 8–12 mph (12–19 km/h). Gentle breeze. Leaves rustle, small twigs move, and flags made from lightweight material stir gently.

**Force 4.** 13–17 mph (20–28 km/h). Moderate breeze. Loose leaves and pieces of paper blow about.

**Force 5.** 18–24 mph (29–38 km/h). Fresh breeze. Small trees that are in full leaf wave in the wind.

**Force 6.** 25–30 mph (39–49 km/h). Strong breeze. It becomes difficult to use an open umbrella.

**Force 7.** 31–38 mph (50–61 km/h). Moderate gale. The wind exerts strong pressure on people walking into it.

**Force 8.** 39–46 mph (62–74 km/h). Fresh gale. Small twigs are torn from trees.

**Force 9.** 47–54 mph (75–88 km/h). Strong gale. Chimneys blown down, slates and tiles torn from roofs.

**Force 10.** 55–63 mph (89–102 km/h). Whole gale. Trees are broken or uprooted.

**Force 11.** 64–73 mph (103–117 km/h). Storm. Trees are uprooted and blown some distance. Cars are overturned.

**Force 12.** 74 mph or more (118 km/h or more). Hurricane. Devastation is widespread. Buildings are destroyed, many trees uprooted. In the original instruction, 'no sail can stand'.

## Saffir–Simpson Hurricane Wind Scale

Category	Sustained wind	Damage
1	74–95 mph (119–153 km/h)	Damage to roofs and gutters; branches may break from large trees; trees with shallow roots may be uprooted; extensive damage to power and telephone lines.
2	96–110 mph (154–177 km/h)	Extensive damage to homes; many shallow-rooted trees broken or uprooted; near-total power loss.
3	111–129 mph (178–208 km/h)	Major damage to homes; many trees snapped or uprooted; loss of power and water supplies.
4	130–156 mph (209–251 km/h)	Catastrophic damage; many homes lose their roofs, some walls collapse; most of the area rendered uninhabitable.
5	157+ mph (252+ km/h)	Damage catastrophic; many homes destroyed; many fallen trees; most power and telephone lines down; area uninhabitable for weeks or months.

## Enhanced Fujita Scale

Rating	Wind speed (km/h; mph)	Damage
EH0	105–137; 65–85	Light
EH1	138–177; 86–110	Moderate
EH2	178–217; 111–135	Considerable
EH3	218–265; 136–165	Severe
EH4	266–322; 166–200	Devastating
EH5	>323; >200	Incredible

---

# Appendix D: SI Units, Conversions, and Multiples

## SI Units (Système International d'Unités)

Quantity	Name of unit	Symbol	Equivalent	Reciprocal
length	metre	m	3.281 feet	1 ft = 0.3048 m
mass	kilogram	kg	2.2 pounds	1 lb = 0.454 kg
time	second	s		
electric current	ampere	A		
thermodynamic temperature	kelvin	K	1 °C (1.8 °F)	1 °C = 1 K
luminous intensity	candela	cd		
amount of substance	mole	mol		

## Supplementary Units

Quantity	Unit	Symbol
plane angle	radian	rad
solid angle	steradian	sr

## Derived SI units

Quantity	Name of unit	Symbol	Equivalent	Reciprocal
frequency	hertz	Hz		
energy	joule	J	0.2388 calories	1 cal = 4.1868 J
force	newton	N	0.225 pounds force	1 lbf = 4.448 N
power	watt	W	0.00134 horse-power	1 hp = 745.7 W
pressure	pascal	Pa	0.00689 lbf/in <sup>2</sup>	1 lbf/in <sup>2</sup> = 145 Pa
electric charge	coulomb	C		
electric potential difference	volt	V		
electric resistance	ohm	Ω		
electric conductance	siemens	S		
electric capacitance	farad	F		
magnetic flux	weber	Wb		
inductance	henry	H		
magnetic flux density	tesla	T		
luminous flux	lumen	lm		
illuminance	lux	lx		
absorbed dose	gray	Gy		
activity	becquerel	Bq		
dose equivalent	sievert	Sv		

## Multiples used with SI units

Name of multiple	Symbol	Value (multiply by)
atto	a	10 <sup>-18</sup>
femto	f	10 <sup>-15</sup>
pico	p	10 <sup>-12</sup>
nano	n	10 <sup>-9</sup>
micro	μ	10 <sup>-6</sup>
milli	m	10 <sup>-3</sup>
centi	c	10 <sup>-2</sup>
deci	d	10 <sup>-1</sup>
deca	da	10
hecto	h	10 <sup>2</sup>
kilo	k	10 <sup>3</sup>
mega	M	10 <sup>6</sup>
giga	G	10 <sup>9</sup>
tera	T	10 <sup>12</sup>
peta	P	10 <sup>15</sup>
exa	E	10 <sup>18</sup>

# Appendix E: Volcanic Explosivity Index (VEI)

VEI	Ejecta (bulk)	Type	Description	Plume	Frequency	Tropospheric injection	Stratospheric injection
0	<10 <sup>4</sup> m <sup>3</sup>	Hawaiian	Effusive	<100 m	Continuous	Negligible	None
1	>10 <sup>4</sup> m <sup>3</sup>	Hawaiian/Strombolian	Gentle	100 m–1 km	Daily	Minor	None
2	>10 <sup>6</sup> m <sup>3</sup>	Strombolian/Vulcanian	Explosive	1–5 km	Fortnightly	Moderate	None
3	10 <sup>7</sup> m <sup>3</sup>	Vulcanian/Peléan/Sub-Plinian	Catastrophic	3–15 km	3 Months	Substantial	Possible
4	>0.1 km <sup>3</sup>	Peléan/Plinian/Sub-Plinian	Cataclysmic	>10 km (Plin. or Sub-Plin.)	18 Months	Substantial	Definite
5	>1 km <sup>3</sup>	Peléan/Plinian	Paroxysmic	>10 km (Plin.)	12 Years	Substantial	Significant
6	>10 km <sup>3</sup>	Plinian/Ultra-Plinian	Colossal	>20 km	50–100 Years	Substantial	Substantial
7	>100 km <sup>3</sup>	Ultra-Plinian	Supercolossal	>20 km	500–1000 Years	Substantial	Substantial
8	>1000 km <sup>3</sup>	Ultra-Plinian	Mega-Colossal	>20 km	>50 000 Years	Vast	Vast

---

# **Appendix F: Torino Impact Hazard Scale**

<b>Category</b>	<b>Number</b>	<b>Implications</b>
No hazard (white zone)	0	Likelihood of collision zero or very low; also applies to small objects that burn up in the atmosphere and falls that cause no damage.
Normal (green zone)	1	Routine discovery; pass near Earth predicted but no unusual level of danger.
Meriting attention from astronomers (yellow zone)	2	Discovery of an object making somewhat close but not unusual pass; no cause for public attention or concern; collision very unlikely.
	3	Close encounter with a 1% or greater chance of collision capable of local destruction; public attention warranted if encounter less than 10 years away.
	4	Close encounter with a 1% or greater chance of collision capable of regional devastation; public attention warranted if encounter less than 10 years away.
Threatening (orange zone)	5	Close encounter posing serious but still uncertain threat of regional devastation; critical attention by astronomers needed to determine whether a collision will occur; governmental contingency planning may be warranted if encounter less than 10 years away.
	6	Close encounter with a large object posing serious but still uncertain threat of global catastrophe; critical attention by astronomers needed to determine whether a collision will occur; governmental contingency planning may be warranted if encounter less than 10 years away.
	7	Very close encounter with a large object which, if occurring this century, poses unprecedented but still uncertain risk of global catastrophe; critical attention by astronomers needed to determine whether a collision will occur; governmental contingency planning may be warranted if encounter less than 30 years away.
Certain collisions (red zone)	8	Collision certain, capable of causing local destruction if on land or tsunami if close offshore; such events occur on average between once per 50 years and once per 1000 years.
	9	Collision certain, capable of causing unprecedented regional devastation if on land or major tsunami if on ocean; such events occur on average between once per 10 000 years and once per 100 000 years.
	10	Collision certain, capable of causing global climatic catastrophe that may threaten future of civilization as we know it, whether impact on land or over ocean; such events occur on average once per 100 000 years or less often.

---

## Appendix G: Avalanche Classes

Class	Damage	Path length
1	Could knock a person down but not bury them.	10 m
2	Could bury, injure, or kill a person.	100 m
3	Could bury and wreck a car, damage a truck, demolish a small building, break trees.	1000 m
4	Could wreck a rail wagon or large truck, demolish several buildings or up to 4 ha of forest.	2000 m
5	Largest known; could destroy a village or up to 40 ha of forest.	3000 m

---

## Appendix H: List of Useful Websites



This is a web-linked dictionary. To access the websites listed below, go to the dictionary's web page at <http://www.oup.com/uk/reference/resources/earthsciences>, click on Web links in the Resources section, and click straight through to the relevant websites.



### Aëtosauria

- Introduction to the Aëtosauria.



### affine

- Affine transformation.



### A'KF diagram

- Describes triangular plots in metamorphic petrology.



### American Geosciences Institute

- Home page of the AGI.



### argon–argon dating

- A tutorial on the  $^{40}\text{Ar}/^{39}\text{Ar}$  step-heating dating technique.



### Austrian Geological Society

- English language home page of the society.



### big bang theory

- An overview of the big bang theory.



### **Brillouin spectroscopy**

- Explains and describes Brillouin spectroscopy.



### **British Geological Survey**

- Home page of the BGS.



### **Bulgarian Geological Society**

- English language home page of the society.



### **Cenozoic**

- The Cenozoic Era.



### **Commission on Ore Mineralogy**

- Forty years of COM.



### **Cretaceous**

- The Cretaceous Period of the Mesozoic Era.



### **Devonian**

- The Devonian Period.



### **diogenites**

- Diogenites.



### **Earth Science Enterprise**

- Earth Science Enterprise.



### **Ediacaran**

- The Ediacaran Period.



### **Eocene**

- The Eocene Epoch.



### **finite rotation**

- Plate Tectonic Modeling: Tools and Methods; Rotation Models.



### **Geological Association of Canada**

- Home page of the GAC.



### **Geological Society of Africa**

- Home page of the society.



### **Geological Society of America**

- Home page of the GSA.



### **Geological Society of Australia**

- Home page of the society.



### **Geological Society of Denmark**

- English-language home page of the society.



### **Geological Society of India**

- English-language home page of the society.



### **Geological Society of Japan**

- English-language home page of the society.



### **Geological Society of London**

- Home page of the Geological Society of London.



### **Geological Society of Malaysia**

- Home page of the society.



### **Geological Society of South Africa**

- Home page of the society.



### **Geological Society of Sweden**

- English-language home page of the society.



### **Geological Society of Trinidad and Tobago**

- Home page of the society.



### **geologic timescale**

- A detailed geological timescale including stages.



### **Geoscience Society of New Zealand**

- Home page of the society.



### **Global Heritage Stone Resource**

- IIGS Subcommission: Heritage Stones.



### **Global Seismographic Network**

- The Global Seismographic Network (GSN).



### **gull**

- Describes two gull caves from the Wiltshire and Avon border.



### **horn**

- Describes glacial horns.



### **Hydrology and Water Resources Programme**

- The World Meteorological Organization's hydrology and water resources programme.



### **International Association for Engineering Geology and the Environment**

- IAEG Homepage.



### **International Federation of Digital Seismic Networks**

- The International Federation of Digital Seismic Networks.



### **International Geophysical Year**

- Explains and describes the IGY.



### **International Polar Year**

- Explains and describes the IPY.



### **Israel Geological Society**

- English-language home page of the society.



### **Itokawa**

- Close-Up on the Asteroid Itokawa.



### **Japanese Meteorological Agency seismic intensity scale**

- Tables explaining the JMA seismic intensity scale.



### **Joint Oceanographic Institutions for Deep Earth Sampling**

- JOIDES Resolution Science Operator.



### **limnology**

- The International Society of Limnology.



### **lithology classification**

- How to classify rocks on site? (Geological or lithological classification).



### **Lost City Hydrothermal Field**

- Geological Evolution of the Lost City Hydrothermal Field.



### **Manning equation**

- An account of Manning's life and work.



### **Mesoproterozoic**

- The Mesoproterozoic Era of the Proterozoic Eon.



### **Mesozoic**

- The Mesozoic Era.



### **metamorphic rock classification**

- Classification of Metamorphic Rocks.



### **Miocene**

- The Miocene Epoch.



### **Neogene**

- Neogene Period.



### **Neoproterozoic**

- Neoproterozoic Era.



### **Nepal Geological Society**

- English-language home page of the society.



### **Nor'easter**

- What is a Nor'easter?



### **Oligocene**

- Oligocene Epoch.



### **Ordovician**

- The Ordovician Period.



### **Palaeocene**

- The Palaeocene Epoch.



### **Palaeogene**

- The Palaeogene Period.



### **palaeontology**

- Describes conditions and life throughout Earth history.



### **Palaeozoic**

- Palaeozoic Era.



### **permanent strain**

- Describes the concept of rheology.



### **Permian**

- The Permian Period.



### **Phanerozoic**

- Phanerozoic.



### **Pleistocene**

- Pleistocene Epoch.



### **Pliocene**

- Pliocene Epoch.



### **Proterozoic**

- Proterozoic Eon.



### **protist**

- What Are Protists?



### **rhenium–osmium dating**

- A review of the physical foundations of the rhenium–osmium method.



### **Saffir–Simpson Hurricane Wind Scale**

- Describes the Saffir–Simpson Hurricane Wind Scale.



### **Silurian**

- The Silurian Period.



### **soil lithology**

- Lithological soil types.



### **Swiss Geological Society**

- English-language home page of the society.



### **synroc**

- Synroc Wasteform.



### **thermoluminescence dating**

- Explains and describes the thermoluminescence technique.



### **Triassic**

- Triassic Period Facts: Climate, Animals & Plants.



### **United States Geological Survey**

- Home page of the USGS.

---

# Appendix I: Satellite Missions

## Abbreviations

CNES	Centre National d'Études Spatiales (France)
CONAE	Comisión Nacional de Actividades Espaciales (Argentina)
DLR	Deutsches Zentrum für Luft und Raumfahrt (Germany)
ESA	European Space Agency
EUMETSAT	European Organisation for the Exploitation of Meteorological Satellites
JAXA	Japan Aerospace Exploration Agency
NASA	National Aeronautics and Space Administration (USA)
NOAA	National Oceanic and Atmospheric Administration (USA)
Roscosmos	Russian Federal Space Agency
RSA	Russian Space Agency
SNSB	Swedish National Space Board
SRI	Space Research Institute (Bulgaria)
SSTC	Belgian Scientific, Technical, and Cultural Services
USDD	United States Department of Defense



The websites below refer to satellite missions described in the main text. Full descriptions of the missions are also available on the companion website.



### Active Cavity Radiometer Irradiance Monitor Satellite

- A NASA mission launched in 1978 to monitor total solar irradiance.



### Advanced Earth Observing Satellite

- A joint CNES, JAXA, NASA, and NASDA mission to study the Earth's environment.



### Apollo

- A NASA programme of missions from 1963 to 1972 to land humans on the Moon and return them to Earth.



### **Aqua**

- A NASA mission launched in 2002 to study the Earth's water.



### **Aquarius**

- A joint CONAE and NASA mission to measure sea surface salinity.



### **Argo**

- Argo Programme home page.



### **A-Train**

- A NASA mission of several satellites placed in polar orbit to study Earth-system science.



### **Aura**

- A NASA mission to monitor complex scientific interactions affecting the globe.



### **AutoNaut**

- AutoNaut Home Page.



### **Cassini**

- A joint ESA and NASA mission that studied Titan, moon of Saturn, and carried the Huygens probe.



### **Ceres**

- Dawn Mission Overview.



### **Clementine**

- A joint NASA and USDD mission launched in 1994 to map the Moon.



### **Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations**

- A joint CNES and NASA mission to study the role that clouds and atmospheric aerosols play in regulating the Earth's weather and climate.



### **CloudSat**

- A NASA mission to further knowledge of atmospheric science.



### **Community Initiative for Continuous Earth Remote Observation**

- Cicero — A Distributed Small Satellite Radio Occultation Pathfinder Mission.



### **Copernicus**

- Copernicus, Observing the Earth.



### **CubeSat Infrared Atmospheric Sounder**

- Mission information.



### **Discovery Program**

- A NASA programme of missions to study asteroids, comets, interstellar dust, Mars, the Moon, and solar wind.



### **Doppler Orbitography and Radiopositioning Integrated by Satellite**

- International DORIS Service.



### **Dynamic Ionosphere CubeSat Experiment**

- ASTRA mission update.



### **Earth Observing System**

- A NASA series of polar-orbiting satellites to study the Earth's land, atmosphere, and oceans.



### **Earth Remote Observation System**

- EROS Satellite Overview.



### **Elektro-L**

- A Roscosmos mission launched in 2011 to enable local and global weather forecasting and measure the Earth's magnetic field.



### **EUMETSAT Advanced Retransmission Service**

- Regional Data Service/EARS.



### **Galileo**

- A NASA mission to study Jupiter and its moons.



### **Genesis**

- A NASA mission to study the origins of the Sun and inner planets.



### **Geostationary Operational Environmental Satellite**

- GOES Satellite Network.



### **Giotto**

- An ESA mission launched in 1985 that studied Halley's Comet.



### **Gravity Recovery and Climate Experiment**

- A joint DLR and NASA mission to accurately map variations in the Earth's gravity field.



### **Hubble Space Telescope**

- About the Hubble Space Telescope.



### **Huygens**

- A joint ESA and NASA probe launched from the Cassini orbiter that landed on Titan, moon of Saturn.



### **Ice, Cloud, and land Elevation Satellite**

- A NASA mission to measure ice sheet mass balance, cloud and aerosol heights.



### **Indian Remote Sensing**

- The Saga of Indian Remote Sensing Satellite System.



### **InSight**

- InSight Mars Lander.



### **International Sun–Earth Explorer 3**

- A joint ESA and NASA mission to study the Earth's magnetosphere.



### Jason-1

- A joint CNES and NASA mission to study global ocean circulation and its link with atmosphere and climate.



### Jason-2

- A joint CNES and NASA mission to study ocean current direction and to measure variations in sea surface height.



### Landsat

- A NASA programme of satellites from 1972 to collect visual information about the Earth.



### Luna

- An RSA mission to collect information about the Moon.



### Lunar Orbiter

- A NASA mission to map the Moon prior to the Apollo manned landing in 1969.



### Magellan

- A NASA mission to study and map the surface of Venus.



### Mariner

- A NASA programme of missions that studied Mars, Venus, and Mercury.



### Mars 96

- An RSA mission launched in 1996 to investigate Mars.



### Mars Global Surveyor

- A NASA mission launched in 1996 to survey and map Mars.



### Mars Pathfinder

- A NASA mission launched in 1996 to study the feasibility of landings on, and explorations of, Mars.



### **Mars Surveyor**

- NASA missions to map and study the climate of Mars.



### **Meteor**

- Meteor Spacecraft Family.



### **Meteor-3M**

- A joint NASA and RSA mission in Sun-synchronous orbit to measure ozone, aerosols, and water vapour in the Earth's atmosphere.



### **Meteosat**

- Meteosat.



### **Monitor-E**

- An RSA mission to study and map the geology and environment of the Earth.



### **Muses-C**

- About Asteroid Explorer 'Hayabusa' (MUSES-C).



### **Near Earth Asteroid Rendezvous**

- NEAR.



### **New Horizons**

- New Horizons: NASA's Mission to Pluto and the Kuiper Belt.



### **New Millennium Deep Space-1**

- A NASA probe designed to test new technology.



### **PARASOL**

- Mission.



### **Phobos**

- An RSA mission to study the Sun, Mars, and Phobos, a moon of Mars.



### Pioneer

- A NASA programme of unmanned space probes launched for planetary exploration.



### QuikSCAT

- Missions: QuikSCAT.



### RADARSAT

- RADARSAT Constellation Mission home page.




### Ranger

- A NASA mission to obtain close-up images of the surface of the Moon.



### Rosetta

- ESA mission website.



### Sakigake

- A Japanese mission to send a test spacecraft close to Halley's Comet, when it last approached the Earth in 1986.



### SeaWiFS

- SeaWiFS Project.



### SELENE

- A JAXA mission to study the Moon's origins and evolution.



### SEOSat/Ingenio

- SEOSAT/INGENIO: Spanish Earth Observation Satellite.



### Solar Radiation and Climate Experiment

- SORCE (Solar Radiation and Climate Experiment).



### **Stardust**

- Stardust.



### **Suisei**

- Suisei.



### **Surface Water Ocean Topography mission**

- SWOT mission homepage.



### **Surveyor**

- A NASA programme to land a series of probes on the Moon.



### **Système Probatoire d'Observation de la Terre**

- SPOT.



### **Tandem-L**

- Tandem-L brochure.




### **Television and Infrared Observation Satellite**

- A NASA mission launched in 1960 to determine if satellites could be useful in the study of the Earth.



### **Terra**

- A NASA mission launched in 1999 to collect data on climate change on Earth.



### **Thermosphere Ionosphere Mesosphere Energetics and Dynamics**

- A joint CNES and NASA mission to study the influence of the Sun and humans on the mesosphere, ionosphere, and lower thermosphere of Earth's atmosphere.



### **Tool for the Analysis of RADIATIONS from lightNings and Sprites**

- TARANIS mission home page.



### **TOPEX/Poseidon**

- A joint CNES and NASA mission to measure the ocean topography of Earth.



### **Tropical Rainfall Measuring Mission**

- About Tropical Rainfall Measuring Mission.



### **Ulysses**

- Ulysses.



### **Vega**

- Vega 1 & 2.



### **Vegetation and Environment monitoring on a New MicroSatellite**

- Joint French–Israeli Venus Mission.



### **Venera**

- An SRI programme to study Venus.



### **Viking**

- A NASA programme of two missions that landed on Mars in 1976.



### **Voyager**

- A NASA programme of two missions launched in 1977 to study the planetary systems of Jupiter and Saturn and beyond.



### **Zond**

- Zond Mission.